

This is a repository copy of *Undertaking studies within a trial to evaluate recruitment and retention strategies for RCTs : lessons learnt from the PROMETHEUS research programme*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/199646/>

Version: Accepted Version

---

**Article:**

Parker, Adwoa [orcid.org/0000-0002-2880-3935](https://orcid.org/0000-0002-2880-3935), Arundel, Catherine Ellen [orcid.org/0000-0003-0512-4339](https://orcid.org/0000-0003-0512-4339), Clark, Laura Kate [orcid.org/0000-0001-9227-5447](https://orcid.org/0000-0001-9227-5447) et al. (16 more authors) (Accepted: 2022) *Undertaking studies within a trial to evaluate recruitment and retention strategies for RCTs : lessons learnt from the PROMETHEUS research programme*. Health Services and Delivery Research. ISSN 2050-4357 (In Press)

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Title page

## Report Title

Undertaking studies within a trial to evaluate recruitment and retention strategies for RCTs: lessons learnt from the PROMETHEUS research programme.

## Keywords

Study within A Trial (SWAT); embedded trials; patient selection\*; research subjects; research design; data accuracy; humans; methodology; Randomized Controlled Trials as Topic; humans; sample size; Research Personnel; Evidence-Based Medicine / standards\*; Practice Guidelines as Topic / standards\*

## Author List

Adwoa Parker<sup>1</sup>, Catherine Arundel<sup>1</sup>, Laura Clark<sup>1</sup>, Elizabeth Coleman<sup>1</sup>, Laura Doherty<sup>1</sup>, Catherine Hewitt<sup>1</sup>, David Beard<sup>2</sup>, Peter Bower<sup>3</sup>, Cindy Cooper<sup>4</sup>, Lucy Culliford<sup>5</sup>, Declan Devane<sup>6,7</sup>, Richard Emsley<sup>8</sup>, Sandra Eldridge<sup>9</sup>, Sandra Galvin<sup>6,7</sup>, Katie Gillies<sup>10</sup>, Alan Montgomery<sup>11</sup>, Chris Sutton<sup>12</sup>, Shaun Treweek<sup>10</sup>, David J Torgerson<sup>1</sup>

<sup>1</sup> York Trials Unit, Department of Health Sciences, University of York, YO10 5DD, UK

<sup>2</sup> Nuffield Department of Orthopaedics, Rheumatology, and Musculoskeletal Science, NIHR Biomedical Research Unit, University of Oxford, Oxford, UK

<sup>3</sup> National Institute for Health Research School for Primary Care Research, Centre for Primary Care and Health Services Research, University of Manchester, Manchester, M13 9PL, UK

<sup>4</sup> School of Health and Related Research, University of Sheffield, 30 Regent Street, Sheffield, S1 4DA, UK

<sup>5</sup> Bristol Trials Centre, Clinical Trials and Evaluation Unit, University of Bristol, Bristol Royal Infirmary, Bristol, UK

<sup>6</sup> School of Nursing and Midwifery, University of Galway, Galway, Republic of Ireland

<sup>7</sup> Health Research Board-Trials Methodology Research Network, Galway, Republic of Ireland

<sup>8</sup> Department of Biostatistics and Health Informatics, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

<sup>9</sup> Institute of Population Health Sciences, Queen Mary University of London, Mile End Road, London, UK

<sup>10</sup> Health Services Research Unit, University of Aberdeen, Health Sciences Building, Forehill, UK  
Aberdeen, Aberdeen, UK AB25 2ZD, UK

<sup>11</sup> University of Nottingham, Nottingham Clinical Trials Unit, University Park Nottingham, Nottinghamshire, NG7 2RD, UK

<sup>12</sup> School of Health Sciences, University of Manchester, Manchester, M13 9PL, UK

## Corresponding Author

David Torgerson, York Trials Unit, Department of Health Sciences, University of York, YO10 5DD; UK  
email - [david.torgerson@york.ac.uk](mailto:david.torgerson@york.ac.uk); Telephone – 01904 321340.

## Disclosure of competing interests

Adwoa Parker holds an NIHR Advanced Fellowship (NIHR302256), received funding from the MRC and NIHR as part of being a co-applicant on this PROMETHEUS programme (MR/R013748/1) (MR/R013748/1), is a co-applicant on an NIHR HTA funded trial (ACTIVE, Award ID: 15/130/84) that was funded by PROMETHEUS, received funding from the Wellcome Trust, through the Centre for Future Health (CFH) at the University of York to developing a training intervention for staff to improve participant recruitment into randomised controlled trials, received funding through an institutional award from an NIHR CTU Infrastructure award (NIHR132547), is Editor for the special SWAT edition for the Journal Research Methods in Medicine & Health Sciences and is a SWAT Collection Advisor for F1000 Research, and is supervising a PhD Studentship (2021-2024) titled ‘The cost effectiveness of SWATs for improving recruitment and retention in RCTs’.

Catherine Arundel received funding from two NIHR HTA grants as a co-applicant (PROFHER-2, award ID 16/73/03; and SWHSI-2, award ID 17/42/94) - both of which received PROMETHEUS funding, received a salary through institutional funding from the MRC and NIHR for the PROMETHEUS programme (MR/R013748/1), received funding through an institutional award for the DISC trial (NIHR HTA award ID 15/102/04)- which received PROMETHEUS funding, and received funding through an institutional award from an NIHR CTU Infrastructure award (NIHR132547).

Laura Clark received a salary through institutional funding from the MRC and NIHR for the PROMETHEUS programme (MR/R013748/1) and received funding through an institutional award from an NIHR CTU Infrastructure award (NIHR132547).

Elizabeth Coleman received a salary through institutional funding from the MRC and NIHR for the PROMETHEUS programme (MR/R013748/1).

Laura Doherty received a salary through institutional funding from the MRC and NIHR for the PROMETHEUS programme (MR/R013748/1) and received funding through an institutional award from an NIHR CTU Infrastructure award (NIHR132547).

Catherine Hewitt received funding from the MRC and NIHR as part of being a co-applicant on this PROMETHEUS programme (MR/R013748/1) (MR/R013748/1), and has also been the recipient of a range of NIHR funding, award IDs: HTA-133865, HTA - 16/111/91, HTA – 132718, HTA - 17/94/36, HTA - 17/76/06, RfPB - PB-PG-0317-20047, HTA – 131483, RfPD – 201176, RfPB - PB-PG-0418-20034, HTA – 15/154/07, HTA – 131784, HTA – 135304, PHR - 16/122/20, HTA – 15/102/04, PHR - 15/05/28, HS&DR - 15/70/26, PGfAR – 201174, HTA - 15/130/84, HTA - 16/167/57, HTA - 16/73/03, HTA – 127739, HTA - 11/36/37, PHR - 14/186/11, HTA – 132808, HTA – 127467, HTA - 15/166/08, HTA – 128625, HTA – 133880, HTA - 17/42/94, HTA - 11/136/52, PGfAR- RP-PG-1214-20017, HS&DR – 129213, HTA – 133784, PGfAR - RP-PG-0216-20002, HTA - 14/49/149, PGfAR – RP-PG-1016-20003, PHR – 128341, HTA – 131805, PGfAR -200607, HTA --133418, HTA -127510, RIGHT -200806, PGfAR - RP-PG-0217-20006, HTA -13/26/01. She is a member of the NIHR HTA Commissioning Committee (2015 to present) and Deputy Chair (2019 to present). Prof Hewitt has also been a member of the HTA Commissioning Sub-Board (2016-2017) and is a current member of the following committees: NIHR CTU Standing Advisory Committee (2020-2024); HTA Post-Funding Committee (2020-2023); HTA Funding Committee Policy Group (2020-2023).

David Beard holds an NIHR Senior Investigator award, has received a range of NIHR funding, award IDs: HS&DR - 131537, HTA - 14/140/63, HTA - 129830, HTA - 127807 RfPB - PB-PG-1215-20041 RfPB - PB-PG-0215-36084, HTA - 134398, HTA - 12/196/08, HTA - 131629, HTA - 15/80/40, HTA - 135260, HTA - 15/103/03, HTA - 15/39/06, HTA – 128043, PGfAR - RP-PG-0615-20002. Prof Beard also received funding from the MRC and NIHR as part of being a co-applicant on this PROMETHEUS programme (MR/R013748/1) (MR/R013748/1) and is a member of the NIHR HTA Commissioning Committee (2020 to present).

Peter Bower holds an NIHR Senior Investigator award, along with a range of other NIHR funding, award IDs: HS&DR – 12/130/33, PRP – 201428, RfPB - PB-PG-0416-20033, RfPB - PB-PG-0816-20035, PGfAR - RP-PG-1016-20010, HS&DR – 17/09/08, HS&DR – 13/54/34, HS&DR – 14/21/21 HTA – 16/84/01 PRP – 200933, HS&DR – 151666, HS&DR – 12/130/15, HS&DR – 134436 HTA – 14/29/01 PDG – 202044, HS&DR – 16/48/07, PGfAR - RP-PG-0216-20003, HS&DR – 14/156/16, PGfAR – 202030, PGfAR - RP-PG-1210-12007, PGfAR - RP-PG-1214-20003, GHRG – 134638. Prof Bower also

received funding from the MRC and NIHR as part of being a co-applicant on this PROMETHEUS programme (MR/R013748/1) (MR/R013748/1).

Cindy Cooper received funding from the MRC and NIHR as part of being a co-applicant on this PROMETHEUS programme (MR/R013748/1) (MR/R013748/1), and has also been the recipient of a range of NIHR funding, award IDs: CTU - 135424, CTU- 132546, CTU- 129761, CTU- 130354, CTU-34 (Infrastructure award), CTU- 129671, CTU- 129743, CTU- 129762, CTU- 135420, CTU- 135458, HTA - 12/21/01, RfPB - PB-PG-0817-20027, HTA - 09/06/01, EME – 133169, HTA - 13/14/01, HTA - 15/08/40, PDG – 202607, HTA - 14/140/80, HS&DR - 14/140/80, Undefined stream – 127963, PDG - RP-DG-0615-10008, HTA - 16/81/01, PHR - 15/49/32, HTA - 15/38/04, HTA - 17/16/04, PGfAR – 201618, HTA - 15/35/03 PGfAR - RP-PG-1016-20006. She is a member of the NIHR CTU Standing Advisory Committee (2016-2023).

Lucy Culliford received funding from the MRC and NIHR as part of being a co-applicant on this PROMETHEUS programme (MR/R013748/1) (MR/R013748/1), and has also been the recipient of a range of NIHR funding, award IDs: EME - 134140 PRP – 203243, RfPB – 201483, RfPB - PB-PG-0418-20029, HTA - 127849, EME - 15/180/55, HTA - 133554, EME - 13/94/15, HTA - 131850, HTA - 127457, HTA - 16/24/09, HTA - 15/101/16, HTA - 16/142/04, EME - 17/145/40, HTA - 16/140/07.

Declan Devane is the Scientific Director of the Health Research Board-Trials Methodology Research Network (HRB-TMRN). The HRB-TMRN is funded by the Health Research Board and its brief includes the funding of SWATs. He is a co-applicant on NIHR funded projects: HS&DR – 150979, HTA – 17/148/07). He is a co-applicant on this PROMETHEUS programme (MR/R013748/1) and participates in a number of Data Safety Monitoring Boards and Advisory Boards.

Richard Emsley has received a range of NIHR funding, award IDs: NIHR300051 NIHR300051, PB-PG 1014-35011 NIHR127755, NIHR130971, 15/187/05, RP-PG-0216-20009, PB-PG-0613-31113, 17/80/09, 15/48/21, 16/111/51, 12/3000/40, NIHR128623, NIHR131175, 13/119/18, NIHR200605, NIHR129926, 16/101/02, NIHR131125, NIHR134702, 15/178/09, 14/29/01, RP-PG-0218-20006, RP-PG-1016-20010, 14/197/65, 15/31/04, NIHR132690, 14/68/08. Prof Emsley is a member of the NIHR HTA Clinical Evaluation and Trials Committee (2018 - 2021), a member of the NIHR Clinical Trials Unit Standing Advisory Committee (2020 - 2023), a member of the NIHR Strategy Board (2022) and is an NIHR Research Professor (2019-2024).

Sandra Eldridge has received a range of NIHR funding, award IDs: CTU-32 (infrastructure award), CTU – 135421, CTU – 129741, PGfAR - RP-PG-0610-10097, HTA - 08/14/41, PGfAR – 202037, PGfAR - RP-PG-0611-20002, PGfAR - RP-PG-0615-20009, PGfAR - RP-PG-1212-20018, PHR - 16/139/01, RIGHT –

200824, PHR - 127793, PGfAR - RP-PG-0609-10181, PGfAR - RP-PG-0617-20004, PGfAR - RP-PG-1016-20008, PGfAR - RP-PG-0612-20001, PGfAR - RP-PG-1209-10038, PGfAR - RP-PG-0218-20001, GHR – 133850. Prof Eldridge is a member of the NIHR CTU Standing Advisory Committee, has been a member of the HTA Clinical Evaluation and Trials Committee (2014-2018).

Sandra Galvin is the National Programme Manager of the Health Research Board - Trials Methodology Research Network (Health Research Board Ireland, employed under funding grant HRB-TMRN-2021-001, which funds SWAT research, is a co-applicant on this PROMETHEUS programme (MR/R013748/1), and is a SWAT Collection Advisor for F1000 Research.

Katie Gillies has been the recipient of a range of NIHR funding, award IDs: HTA – 129248, RfPB - PB-PG-0416-20033, HTA – 132999, HTA - 14/192/71, HTA – 133561, HTA - 17/68/01, HTA – 130310, HTA – 127280, HS&DR – 131537. Dr Gillies is a member of the NIHR HTA Clinical Evaluation and Trials Committee.

Alan Montgomery received funding from the MRC and NIHR as part of being a co-applicant on this PROMETHEUS programme (MR/R013748/1). He has been the recipient of a range of NIHR and MRC funding, award IDs: CTU – 132545, CTU – 135422, CTU-26, RfPB - PB-PG-0613-31083, RfPB - PB-PG-1217-20026,, HTA - 12/190/05 EME - 11/47/03 RfPB - PB-PG-0817-20005, HTA - 15/110/02, HTA - 132711, HTA - 10/57/24, HTA - 129210, HTA - 127393, HTA - 127292, HTA - 131159, HTA - 16/96/09, HTA - 132016 EME - 128240 PGfAR - 203279, HTA - 12/24/02, HTA – 129515, PRP - 015/0309, PDG - RP-DG-0615-10005, HTA - 129917, HTA - 12/127/10, HTA - 17/94/31, HTA – 131440, COVID-19 – 133443, MRC - MR/K025643/1, MRC - MR/V020803/1. Prof Montgomery is an independent member of Trial Steering / Data Monitoring Committees for various NIHR-funded studies. He was a member of the HTA Clinical Evaluation and Trials Committee (2015-2021).

Chris Sutton has received institutional funding from the MRC and NIHR for the PROMETHEUS programme (MR/R013748/1). He has also been the recipient of a range of NIHR funding, award IDs: HS&DR - 130581, RfPB - 201093, RfPB - 203475, HTA - 16/111/31, RfPB - PB-PG-0214-33010, RfPB - PB-PG-1216-20024, RfPB - 203468, RfPB - 203507, HTA – 131483, PDG – 202044, HS&DR - 14/04/16, EME - 132622, HTA – 133418. He is a member of the NIHR HTA Commissioning Committee (2020 to present).

Shaun Treweek received funding from the MRC and NIHR as part of being a co-applicant on this PROMETHEUS programme (MR/R013748/1). He has also been the recipient of NIHR and MRC funding, award IDs: PHR – 129791, HTA - 12/127/12, HTA - 15/130/73, MRC - MR/K025643/1. He is Editor-in-Chief of the journal 'Trials'.

David Torgerson received funding from the MRC and NIHR as part of being the Chief Investigator on this PROMETHEUS programme (MR/R013748/1). He has also been the recipient of a range of NIHR funding, award IDs: HTA - 14/49/149, PHR - 15/05/28, CTU-40 (infrastructure award), HTA - 127510, PHR – 128341, CTU - 132547, RfPB – 203506, PGfAR - RP-PG-0609-10171, HTA - 13/26/01, HTA - 15/166/08, PGfAR - RP-PG-0615-20003, HTA - 15/130/84, HTA - 15/154/07, RfPB - PB-PG-0416-20035, HTA - 16/73/03, PHR - 16/122/20, HTA - 16/167/56, HTA - 16/167/57, HTA - 17/42/94, HTA - 17/94/36, RfPB - PB-PG-0418-20034, HTA - 127739, HTA - 128625, HTA - 132718, PHR - 131745, HTA - 131805, HTA - 132674, RfPB – 202203. Prof Torgerson is an Editorial board member for the Journal Research Methods in Medicine & Health Sciences and is a SWAT Collection Advisor at F1000 Research.

## Total word count

33247

## Abstract

### *Background*

Randomised controlled trials ('trials') are susceptible to poor participant recruitment and retention. Studies Within A Trial (SWATs) are the strongest methods for testing the effectiveness of strategies to improve recruitment and retention. However, relatively few of these have been conducted.

### *Aims*

PROMoting THE USE of Studies Within A Trial (PROMETHEUS) aimed to facilitate at least 25 SWATs evaluating recruitment or retention strategies. We share our experience of delivering the PROMETHEUS programme, and the lessons learnt for undertaking randomised SWATs.

### *Design*

A network of 10 Clinical Trials Units (CTUs) and one primary care research centre committed to conducting randomised controlled SWATs of recruitment and/or retention strategies was established. Promising recruitment and retention strategies were identified from various sources including Cochrane systematic reviews, the SWAT Repository, and existing prioritisation exercises, which were reviewed by patient and public (PPI) members to create an initial priority list of seven recruitment and eight retention interventions.

Host trial teams could apply for funding and receive support from the PROMETHEUS team to undertake SWATs. We also tested the feasibility of undertaking coordinated SWATs, across multiple host trials simultaneously.

### *Setting*

CTU-based trials recruiting or following up participants in any setting in the UK were eligible.

### *Participants*

CTU-based teams undertaking trials in any clinical context in the UK.

### *Interventions*

Funding of up to £5,000 and support from the PROMETHEUS team to design, implement, and report SWATs.

### *Main outcome measures*

Number of host trials funded.



### *Results*

42 SWATs were funded (31 host trials), across 12 CTUs. The mean cost of a SWAT was £3,535. Twelve SWATs tested the same strategy across multiple host trials using a coordinated SWAT design, and four used a factorial design. Two recruitment and five retention strategies were evaluated in more than one host trial. PROMETHEUS will add 18% more SWATs to the Cochrane systematic review of recruitment strategies, and 79% more SWATs to the Cochrane review of retention strategies. For retention, we found that pre-notifying participants by card, letter or email before sending questionnaires was effective, as was the use of pens, and sending personalised text messages to improve questionnaire response. We highlight key lessons learnt to guide others planning SWATs, including involving PPI partners; prioritising and selecting strategies to evaluate and elements to consider when designing a SWAT; obtaining governance approvals; implementing SWATs, including individual and coordinated SWATs, and; reporting SWATs.

### *Limitations*

The COVID-19 pandemic negatively impacted five SWATs, being either delayed (n=2) or prematurely terminated (n=3).

### *Conclusions*

PROMETHEUS significantly increased the evidence base for recruitment and retention strategies. When provided with both funding and practical support, host trial teams successfully implemented SWATs.

### *Future research*

Future research should identify and target gaps in the evidence base, including widening SWAT uptake, undertaking more complex SWATs, and translating SWAT evidence into practice.

### *Study registration*

All SWATs had to be registered with the Northern Ireland Network for Trials Methodology Research SWAT Repository.

### *Funding details*

This project was funded by the Medical Research Council (MR/R013748/1) and National Institute for Health Research as part of the MRC–NIHR Methodology Research programme.

## Table of contents

Report Title	1
Keywords	1
Author List	1
Corresponding Author	2
Disclosure of competing interests	2
Total word count	6
Abstract	7
Background	7
Aims	7
Design	7
Setting	7
Participants	7
Interventions	7
Main outcome measures	7
Results	8
Limitations	8
Conclusions	8
Future research	8
Study registration	8
Funding details	8
Table of contents	9
List of Tables	19
List of Figures	20
Abbreviations	21
Plain English summary	25

Aims of the research	25
Background	25
Methods	25
Results	25
Conclusions	25
Scientific summary	26
Background	26
Aims	26
Methods	26
Results	26
Conclusions	27
Recommendations	27
Recommendations for funders	27
Recommendations for Sponsors	28
Recommendations for involving patient and public involvement partners (PPI) in SWAT research	28
Recommendations for oversight committees	29
Recommendations for journals and reviewers	29
Recommendations for trial teams and methodologists undertaking SWAT research	29
Recommendation for trial conduct and using SWAT evidence	30
Recommendations for future research	30
Study registration	32
Funding details	32
Introduction	33
What is covered in this report?	33
Who should use this report?	33
Chapter 1: Background to the PROMETHEUS programme	35
The importance of randomised controlled trials	35

Recruitment and retention difficulties in trials	35
SWAT methodology	35
Addressing recruitment and retention difficulties using SWATs	36
Aims and objectives	37
Chapter 2: PROMETHEUS methods	38
PROMETHEUS preparatory work	38
Eligibility criteria	39
Host trials	39
SWAT interventions	40
PROMETHEUS programme outcomes	40
Sample size	40
Host trial recruitment	41
Randomisation	41
Blinding	41
Data collection methods	42
Statistical methods	42
Data monitoring	42
Harms	42
Auditing	42
Research ethics approval	43
Consent or assent	43
Confidentiality	43
Access to data	43
Chapter 3: PROMETHEUS results	44
Host trials funded, numbers of recruitment SWATs funded, number of retention SWATs funded	44
Recruitment strategies	46
Retention strategies	46
SWAT costs	47

Individual SWATs	47
Coordinated SWATs	48
Results: findings from the funded SWATs	49
Results of recruitment SWATs	49
Results of retention SWATs	57
Retention strategies demonstrating evidence of effectiveness	77
Reminders and prompts	77
Non-monetary incentives	77
Retention strategies demonstrating a lack of effectiveness, or with the potential to harm retention rates	78
Chapter 4: Discussion of the PROMETHEUS programme	79
Summary of main findings from PROMETHEUS	79
Strengths	80
Limitations	81
Comparison with existing programmes of SWAT research: MRC START and TRECA	82
Patient and public involvement (PPI)	83
Equality, Diversity and Inclusion	83
Conclusions	83
Chapter 5: Lessons learnt for PPI	85
Chapter overview	85
Defining PPI in PROMETHEUS	85
Aims	85
PPI: Methods	85
PPI input prior to funding being awarded	85
PPI involvement in prioritisation of strategies to be funded	86
PPI input into individual SWATs	86
PPI: Results	86
PPI involvement in prioritisation of strategies to be funded	86

PPI input into individual SWATs	87
PPI: Discussion and conclusions	87
PPI: Reflections and critical perspective	87
Lessons learnt for involving PPI partners in SWAT research.	88
Chapter 6: Lessons learnt: SWAT development and funding	89
Chapter overview	89
Selecting which strategy to test in a SWAT: the importance of prioritisation	89
SWATs prioritisation, GRADE and pragmatism	90
Supporting teams to undertake SWATs: Trial Forge Evidence packs and an information repository	91
Design Choices: cluster randomised vs individual randomised	92
Funding for SWATs	94
Lessons learnt for trial teams and methodologists undertaking SWAT research	95
Lessons learnt for funders	95
Chapter 7: Lessons learnt for governance approvals for SWATS	96
Chapter overview	96
Governance approvals	96
Sponsor Approval	96
Oversight Committee Approvals (TMG/TSC/DMC)	97
Ethics/HRA Approval	98
Conclusion	101
Lessons learnt for Sponsors	102
Lessons learnt for oversight committees	102
Lessons learnt for trial teams and methodologists undertaking SWAT research	102
Lessons Learnt for journals and reviewers	102
Chapter 8: Lessons learnt for implementing SWATS	103
Chapter overview	103
Introduction	103

Individual SWATs	103
Advantages of Individual SWATs	104
Disadvantages of individual SWATs	104
Case study of individual SWATs: Pens for retention	105
Background	105
Aims and Objectives	105
Methods	106
<i>Individual SWAT Methods</i>	106
<i>Participants, Interventions and Outcomes</i>	106
<i>Randomisation and Blinding</i>	107
<i>Sample Size</i>	107
<i>Analyses</i>	107
<i>Primary outcome</i>	107
<i>Secondary outcomes</i>	107
<i>SWAT Meta-Analysis Methods</i>	108
Results of PROMETHEUS SWATs testing the use of pens for retention	108
Comparison with Existing Literature	108
Cost-effectiveness	109
Discussion	109
Coordinated SWATs	110
Advantages of coordinated SWATs	110
Disadvantages of coordinated SWATs	110
Case studies of coordinated SWATs	111
Case study - Staff training to improve participant recruitment into surgical randomised controlled trials: a feasibility Study Within A Trial (SWAT) across four host randomised controlled trials simultaneously	111
Background	111
Aims and objectives	112

Methods	112
<i>Host trial and participant recruitment</i>	112
<i>SWAT Intervention</i>	112
<i>SWAT control group</i>	112
<i>Outcomes</i>	112
<i>Randomisation</i>	113
<i>Data collection</i>	113
<i>Statistical analysis</i>	113
Results	114
<i>Feasibility of recruiting host trials</i>	114
<i>Feasibility of recruiting sites, participants and randomisation</i>	115
<i>Feasibility of training course</i>	115
<i>Acceptability of the training course</i>	115
<i>Impact of recruitment training on self-confidence and awareness</i>	115
<i>Impact of training on screening and recruitment rates</i>	115
Discussion	116
Case study - Bah humbug! Association between sending Christmas cards to trial participants and trial retention: randomised study within a trial conducted simultaneously across eight host trials	118
Background	118
Aims and Objectives	118
Methods	118
<i>Design</i>	118
<i>Setting</i>	118
<i>Participants and sample size</i>	118
<i>Intervention and comparator</i>	118
<i>Primary outcome</i>	119
<i>Randomisation and Blinding</i>	119



<i>Statistical analysis</i>	119
Results	119
<i>Host trials and participants</i>	119
<i>Impact of sending Christmas cards on retention rates</i>	121
<i>Costs and carbon footprint</i>	121
Discussion	122
Conclusions	122
Lessons learnt for trial teams and methodologists undertaking SWAT research	123
Lessons learnt for funding bodies	123
Chapter 9: Lessons learnt for statistical considerations	124
Chapter overview	124
Sample size considerations	124
Analysis considerations	125
Meta-analysis	127
Economics and costings	128
Conclusions	128
Lessons learnt for trial teams and methodologists undertaking SWAT research	129
Lessons learnt for funders	129
Chapter 10: Lessons learnt for reporting SWATs	130
Chapter overview	130
The need for transparent reporting of SWATs	130
Reporting SWATs	130
Identifying Peer Reviewers	131
Peer review feedback	132
Conclusions	132
Lessons learnt for trial teams and methodologists undertaking SWAT research	132
Lessons learnt for journals and reviewers	133
Chapter 11: Recommendations for future research practice, direction, and support	134

Chapter overview	134
Recommendations	134
Recommendations for funders	134
Recommendations for Sponsors	135
Recommendations for involving PPI partners in SWAT research	135
Recommendations for oversight committees	135
Recommendations for Journals and Reviewers	136
Recommendations for trial teams and methodologists undertaking SWAT research	136
Recommendation for trial conduct and using SWAT evidence	136
Recommendations for future research	136
Chapter 12: Overall conclusions	139
Acknowledgements	141
Contributions of authors	142
Study registrations	143
Funding details	143
Ethics Statement	143
Data Sharing Statement	145
Additional funding received	145
Publications	146
References	148
Appendix 1	166
List of funded host trials and SWATs	166
Appendix 2	177
PROMETHEUS Christmas Card SWAT Intervention	177
Appendix 3	178
Draft HRA approvals Guidance process for SWATs	178
Guidance for SWAT applicants	178
Appendix 4	186



## List of Tables

Table 1: List of key recruitment and retention questions in priority order .....	39
Table 2: The research area of each of the SWATs funded by the PROMETHEUS programme .....	45
Table 3: Examples of funding for staff time required for individual SWATs .....	48
Table 4: Host trials and results of completed SWATs evaluating recruitment strategies .....	50
Table 5: Ongoing SWATs evaluating recruitment strategies .....	53
Table 6: Host trials and results of completed SWATs evaluating retention strategies .....	58
Table 7: Ongoing SWATs evaluating retention strategies .....	70
Table 8: Prematurely terminated SWATs evaluating retention strategies .....	73
Table 9: Information to include when registering a SWAT in the SWAT Repository .....	92
Table 10: Examples of individual and cluster randomised designs in SWATs .....	93
Table 11: SWAT groups and approvals required .....	101
Table 12: SWAT Participant, Interventions and Outcomes .....	106
Table 13: Details of the trials included in the PROMETHEUS Training SWAT .....	114
Table 14: PROMETHEUS Training SWAT - number of patients approached and agreeing to participate at baseline and follow-up, as reported by recruiters per participant defined typical week in the last month .....	116
Table 15: Descriptions of the host trials involved in PROMETHEUS Christmas Card SWAT .....	120
Table 16: Estimated sample size required for a single SWAT to have 80% or 90% power to detect various differences in proportions .....	125
Table 17: Characteristics of currently funded host randomised controlled trials .....	169
Table 18: Guide to which approvals for should be obtained for individual and overarching SWATs	180
Table 19: Cover letter table .....	183
Table 20: Glossary of terms .....	185
Table 21: The SWAT reporting template. The text in blue italics is text that could be cut and pasted into a document as the starting point for reporting a SWAT evaluation. ....	187

## List of Figures

Figure 1 : Flow chart of SWATs funded by the PROMETHEUS programme.....	44
Figure 2: Summary example in Trial Forge evidence packs, used with kind permission from Trial Forge {Trial Forge, 2021} <sup>119</sup> .....	91
Figure 3: Pooled Odds Ratio for Effectiveness of Pen with a Follow Up Questionnaire (Pre PROMETHEUS) {Sharp 2006, Bell 2016} <sup>140, 141</sup> .....	108
Figure 4: Pooled Effect of Including a Pen with a Follow Up Questionnaire {Sharp 2006, Bell 2016, Mitchell 2020, Cunningham-Burley 2020, James 2020} <sup>140, 141, 92 68 81</sup> .....	109
Figure 5: Christmas Card SWAT Cumulative meta-analysis {Coleman 2021} <sup>49</sup> .....	121
Figure 6: PROMETHEUS Christmas Card SWAT intervention.....	177
Figure 7: SWAT evaluation lead and host trial team roles.....	178

## Abbreviations

ACTIVE - Articular Pilon Fracture Trial

ActWELL - A randomised controlled trial to assess the impact of a lifestyle intervention (ActWELL) in women invited to NHS breast screening

ARTISAN - Acute Rehabilitation following Traumatic anterior shoulder dISlocAtioN

C-Gall - A randomised controlled trial comparing laparoscopic Cholecystectomy with observation/conservative management for preventing recurrent symptoms and complications in adults with uncomplicated symptomatic GALLstones

CHAMP-1 - Community pharmacy: highlighting alcohol use in medication appointments pilot study

CLEAR - A 2x2 factorial randomised open label trial to determine the CLinical and cost-Effectiveness of hypertonic saline (HTS 6%) and carbocisteine for Airway cleaRance versus usual care over 52 weeks in bronchiectasis

CONSORT - Consolidated Standards of Reporting Trials

COVID-19 - Coronavirus Disease 2019

CPIT-III - The Cessation in Pregnancy Incentives Trial (CPIT) III

CRN - Clinical Research Network

CTUs - Clinical Trials Units

DISC - Dupuytren's Interventions: Surgery vs. Collagenase (DISC) Trial

DMC - Data Monitoring Committee

FUTURE - Female Urgency, Trial of Urodynamics as Routine Evaluation

GDPR - General Data Protection Regulation

GRADE - Grading of Recommendations, Assessment, Development and Evaluations

GRASP - Getting it Right: Addressing Shoulder Pain

GRIPP - Guidance for Reporting Involvement of Patients and the Public

HRA - Health Research Authority

HRB-TMRN - Health Research Board – Trials Methodology Research Network

HS&DR - Health and Social Care Delivery Research

HTA - Health Technology Assessment

IBD-BOOST - A Randomised Controlled Trial of supported online self-management for symptoms of fatigue, pain and urgency/incontinence in people with inflammatory bowel disease

IntAct - Intraoperative Fluorescence Angiography to Prevent Anastomotic Leak in Rectal Cancer Surgery

KReBS - Knee Replacement Bandaging Study

L1FE - Lateral Compression Type-1 Fracture Fixation in the Elderly

MAGIC - Melatonin for Anxiety prior to General anaesthesia In Children

MiQuit - The MiQuit study: feasibility trial of a computer-tailored smoking cessation intervention providing individualised written and mobile phone text message support to pregnant smokers

MRC - Medical Research Council

MSS3 - Multiple Symptoms Study 3

NHS - National Health Service

NIHR - National Institute for Health and Care Research

ORRCA - Online Resource for Research in Clinical triAls

OTIS - Occupational Therapist Intervention Study

PEP-TALK - A study investigating whether having group discussions in addition to physiotherapy improves the amount of physical activity following hip and knee replacement

PIL - Patient Information Leaflet

PMG - Project Management Group

POSNOc - POSitive Sentinel NOde: adjuvant therapy alone versus adjuvant therapy plus Clearance or axillary radiotherapy

PPI - Patient and Public Involvement

PRioRiTy - Prioritising Recruitment in Randomised Trials

ProFHER-2 - PROximal Fracture of Humerus: Evaluation by Randomisation-2

PROMETHEUS - PROMoting THE Use of SWATs

PURe - Percutaneous nephrolithotomy, flexible Ureterorenoscopy and Extracorporeal shockwave lithotripsy for lower pole kidney stones

R&D - Research and Development

RCT - Randomised Controlled Trial

REC - Research Ethics Committee

REFLECT - A Randomised controlled trial to Evaluate the effectiveness and cost benefit of prescribing high dose FLuoride toothpaste in preventing and treating dEntal Caries in high-risk older adultS

SAP - Statistical Analysis Plan

SARC - Salbutamol for analgesia in renal colic

SSHeW - Does Slip Resistant Footwear Reduce Slips Among Healthcare Workers? A Randomised Controlled Trial

START - Systematic Techniques for Assisting Recruitment to Trials

START:REACTS - Sub-acromial spacer for Tears Affecting Rotator cuff Tendons: a Randomised, Efficient, Adaptive Clinical Trial in Surgery

SWAP - Study Within a Project

SWAT - Study Within A Trial

SWHSI-2 - Surgical Wounds Healing By Secondary Intention - 2

TOPaZ - Treatment of osteogenesis imperfecta with parathyroid hormone and zoledronic acid



TRECA - TRials Engagement in Children and Adolescents

TMG - Trial Management Group

TMRP - Trial Methodology Research Partnership

TSC - Trial Steering Committee

UK - United Kingdom

UK FROST - United Kingdom FROzen Shoulder Trial

VITA - Lactic acid gel versus metronidazole for recurrent bacterial vaginosis in women aged 16 years and over

WORKWELL - A randomised controlled trial of job retention vocational rehabilitation for employed people with inflammatory arthritis

YTU - York Trials Unit

## Plain English summary

### *Aims of the research*

A Study Within a Trial (SWAT) is a research study nested inside a larger 'host trial'. PROMETHEUS aimed to make doing SWATs routine practice in CTUs by funding and supporting at least 25 SWATs.

### *Background*

The best way to test health and social care treatments is to do a randomised controlled trial ('trial'), where some patients get the treatment being tested and some do not. The results of the different groups are compared to see if the treatment improves care. Recruiting patients and keeping them involved in trials is often very difficult. Research teams often do not know how best to recruit and keep patients engaged as the methods have not been tested to see if they work. The best way to test these methods is by doing a SWAT. We test a programme of SWATs for recruiting and keeping patients engaged in trials.

### *Methods*

Trial teams were able to apply for funding of up to £5000 and receive support from the PROMETHEUS team to do SWATs. We use our experience of doing SWATs to outline lessons learnt for doing SWATs.

### *Results*

We funded 42 SWATs and gave teams advice to do them. We significantly increased the knowledge for both recruitment and retention strategies, and found 'pre-notifying' before sending questionnaires, sending pens and personalised text messages were all effective for increasing responses by participants. We tested SWATs across several different trials at the same time to find out more quickly whether their methods worked. We highlight key lessons learnt to guide others doing SWATs, including involving patient partners; picking the right strategy to test; getting ethical approvals; how to do and report SWATs.

### *Conclusions*

PROMETHEUS was successful and supported more SWATs than planned. We hope our experience will support those doing SWATs in the future.

## Scientific summary

### *Background*

Many delivery aspects of randomised controlled trials have not been subjected to rigorous evaluation. Strategies to enhance recruitment and retention often tend to not be based on evidence. However, there is an increasing interest in embedding RCTs of such strategies. The Medical Research Council (MRC) funded START, which found it feasible to test two strategies for recruitment and retention across a number of trials by performing randomised 'Studies Within A Trial' or SWATs.

### *Aims*

The aim of the PROMoting THE USE of Studies Within A Trial (PROMETHEUS) programme was to build on the START initiative and make embedding SWATs standard practice across Clinical Trials Units (CTUs). We intended to pump prime and facilitate the start of at least 25 SWATs across multiple CTUs within 30 months. The ultimate aim was to make the inclusion of SWATs routine when conducting a trial in a CTU. We share our experience of delivering the PROMETHEUS programme, along with the lessons learnt for undertaking randomised SWATs of recruitment and retention methods.

### *Methods*

A network of 10 CTUs and one primary care research centre committed to conducting randomised controlled studies within trials (SWATs) of recruitment and/or retention interventions was established by the PROMETHEUS team. We identified promising recruitment and retention interventions from a variety of sources including Cochrane systematic reviews, the Northern Ireland Network for Trial Methodology Research SWAT Repository Store, and existing prioritisation exercises. Promising strategies were reviewed by our patient and public (PPI) members to create an initial priority list of seven recruitment and eight retention interventions. The programme allowed host trial teams to apply for funding of up to £5,000 and receive support from the PROMETHEUS team to design, implement, and report SWATs. We also tested the feasibility of undertaking coordinated SWATs, across multiple host trials simultaneously.

### *Results*

The PROMETHEUS programme funded 42 SWATs, embedded within 31 host trials, across 12 CTUs. The mean cost of a funded SWAT was £3,535. Of the 42 SWATs, 12 tested the same SWAT across multiple host trials using a coordinated SWAT design, and four tested more than one strategy in a factorial design. Two recruitment and five retention strategies were evaluated in more than one host trial. In the space of just 4 years, PROMETHEUS will add 18% more SWATs to the Cochrane review of

global recruitment strategies, and 79% more SWATs to the Cochrane review of global retention strategies. The results from the SWATs reported to date found there was no evidence of a significant difference in recruitment for any of the strategies tested. For retention, pre-notifying participants by card prior to sending questionnaires was effective (risk difference 3.3%, 95% CI -3.0% to 9.6%); as was pre-notifying participants by letter or email (risk difference 3.8%, 95% CI -6.1 to 13.6%). Sending personalised text messages was more effective for improving the return of postal questionnaires compared to non-personalised text messages (risk ratio 1.16, 95% CI 1.00 to 1.33); and resulted in fewer completions via telephone compared with a non-personalised text (adjusted OR 0.44, 95% CI 0.22 to 0.87). Including a pen with a questionnaire probably increases retention and response rate (pooled OR 1.21, 95% CI 1.09 to 1.35). We highlight key lessons learnt below (Recommendations).

### *Conclusions*

The PROMETHEUS programme significantly increased the international evidence base for both recruitment and retention strategies within RCTs. The funded SWATs evaluated a wide range of recruitment and retention strategies; however, the COVID-19 pandemic negatively impacted five funded SWATs, with two being delayed and three prematurely terminated. Through this project, we identified that when provided with both funding and practical support, host trial teams successfully implemented SWATs. PROMETHEUS led to an overall increase in the evidence base; however, ongoing 'routine' application of SWATs across RCTs employing the lessons learnt is required to ensure that efficient trial conduct strategies are identified.

### *Recommendations*

#### *Recommendations for funders*

- All trial funders should contribute to the effort to improve the efficiency of trials. Funders should encourage the teams that they fund to undertake SWATs
- Funding streams specifically designed to support SWATs must be made available to trial teams to continue building the trial process evidence base, for recruitment and retention as well as for other stages of the trial design and delivery process. This includes funding streams for undertaking specific SWATs, as well as infrastructure funding to support CTUs and other centres to undertake coordination activities that will support the design, conduct, reporting, and implementation of SWATs and their findings to inform the work of the NIHR, the MRC, and other funders
- PROMETHEUS has demonstrated that coordination of activity remains crucial to the delivery of SWATs. A central, national coordination point that provides hands-on support needs to continue and funding should be allocated for this. Additionally, CTUs should identify a lead

for SWATs, to support SWAT activities and evidence-based trial conduct within the CTU, as well as links with others undertaking SWATs elsewhere to share best practice. The funding for both central and CTU-based support should be ongoing

- SWAT priorities need to be identified and communicated clearly to funders, and funders should use these priorities to inform their funding decisions
- Funders should develop a mechanism to promote SWAT questions that have been identified as a priority during the funding application process
- The mean cost of funding requested for a standalone SWAT within PROMETHEUS was £3,535 (range £500 to £5,000). The coordinated SWATs cost £10,668 (training SWAT), and £1,306.40 (Christmas cards); however, these did not include costs for central coordination, data preparation and sharing by the host trial teams, data cleaning, analysis, and write-up. These costs suggest that the £10,000 being offered by the NIHR for trial teams to include a SWAT should be sufficient for most planned SWATs. However, there may be occasions where trial teams may wish to test strategies that may be more expensive
- When applying for SWAT funding, trial teams should be asked to indicate whether the question they are addressing is a priority SWAT question, and to provide a clear rationale for selecting that particular question
- If teams are unable to undertake a SWAT, funders should ask that recruitment and retention methods are clearly reported to support the evidence base

#### *Recommendations for Sponsors*

- Our experience suggests that there is a need for clear, easily accessible information about the nature of SWATs, as well as the role of the funder in supporting SWATs
- Any future changes proposed by the Health Research Authority to the approvals process need to be communicated clearly and applied consistently to each SWAT

#### *Recommendations for involving patient and public involvement partners (PPI) in SWAT research*

- PPI should be considered when undertaking a SWAT, in the same way PPI is expected to be undertaken in the main trial
- PPI partners should be involved to develop novel and untested recruitment and retention strategies, as well as to adapt existing strategies to the context of their specific host trial and the population being enrolled

### *Recommendations for oversight committees*

- Our experience suggests that Trial Management Groups (TMGs) play a key role in decisions about whether a SWAT is undertaken and continued in the host trial or not. TMG members should encourage the uptake of SWATs in their trials. Whilst the findings of SWATs may not always directly inform their host trial, the findings of SWATs undertaken during the early, or internal pilot, phase of the trial may inform the decisions about which strategies should be used at a later stage, such as in the main trial
- Trial Steering Committees should review the SWAT activity and progress, in the same way that they review sub-studies in a trial
- Data Monitoring Committee review is dependent on the specific host trial and SWAT strategy being evaluated

### *Recommendations for journals and reviewers*

- Journal peer reviewer profiles should be updated to include methodological interests and expertise, to support evidence of suitability to undertake a peer review for a SWAT
- When selecting peer reviewers, the SWAT and methodology interests as registered by reviewers should be used where possible
- SWATs are a niche area and so to increase the pool of reviewers, journals should consider being more flexible when assessing reviewer credentials to review a SWAT, such as allowing relevant experience in place of a PhD
- Reviewers should be advised that in many instances informed consent from participants need not be obtained when undertaking a SWAT, due to the low risk associated with the intervention and in the case of retention SWATs due to existing consent for further research being in place. This may depend on the jurisdiction. In the UK this approach is supported by HRA guidance in relation to grading for SWAT interventions and approvals.
- Robust and transparent reporting is necessary, that is compliant with Consolidated Standards of Reporting Trials (CONSORT)

### *Recommendations for trial teams and methodologists undertaking SWAT research*

- There remains a need for continually updated research priorities to allow researchers to address the questions relevant at that time
- When SWAT priorities are set, methodologists need to provide enough information to enable teams to make informed decisions about evaluating the priorities set
- Further work is needed to help teams identify suitable SWAT strategies for their host trials
- SWAT priorities need to be communicated clearly and consistently to trial teams

### *Recommendation for trial conduct and using SWAT evidence*

- As the evidence base develops for effective and cost-effective recruitment and retention strategies, it will become increasingly important for trial teams to use this evidence base to inform their recruitment and retention activities. Trial teams need to actively engage with the evidence base to inform their practice. Funders will need to actively support the trials they fund to use evidence-informed recruitment and retention strategies

### *Recommendations for future research*

- There remains a substantial need for more high-quality SWAT evidence and so Chief Investigators should be encouraged to consider the embedding of a SWAT at the funding stage. Further work is therefore needed to increase the awareness of the methodological importance of SWAT research with research teams, and to develop engagement strategies to increase SWAT activity
- Future research needs to focus on identifying whether further replications are needed for existing evidence. If so, the gaps in the evidence base should be targeted. More coordination and replication of SWAT evaluations are encouraged
- A 'real-time' and dynamic communication strategy including a clear cost and resource breakdown for each suggested SWAT should be developed. This will alleviate the burden on trials teams to begin costing exercises and enables them to make an informed decision more quickly as to whether they can embed a given SWAT
- There is a need to aid teams to identify and select a suitable SWAT for their host trial populations. Pragmatic decisions on which SWAT may be appropriate and feasible to include should be taken as required. A mechanism to communicate SWAT research priorities is needed, and this information needs to be readily accessible for all trialists to refer to
- Our findings demonstrate that within an individual host trial, there is often a capacity to address more than one SWAT question, either separately, or simultaneously using a factorial design. This suggests that there is a capacity to significantly speed up and strengthen the evidence base through teams undertaking more than one SWAT in their trials where relevant
- For certain strategies, coordinated SWATs should be encouraged. This method could be used to rapidly replicate SWAT evaluations to plug the evidence gap, as well as to evaluate more complex recruitment and retention strategies that may be more challenging to undertake using individual SWATs. Materials should be developed to advise teams on how to

undertake coordinated SWATs, as well as a method of networking to enable teams to promote their coordinated SWAT and collaborate

- As the evidence base develops, it will become increasingly important for trialists to utilise the evidence base in a systematic way to identify both effective and ineffective strategies to inform their practice. Future work should therefore consider issues around the dissemination and implementation of SWATs and develop guidance to enable the wider trials community to undertake, report and adopt the findings of SWATs. Implementation science, the study of methods to promote the uptake of evidence-based practice, could be used to inform any such future work. Funders can also help by questioning strategies proposed by trial teams that are known to be either ineffective or not cost-effective
- Improving the knowledge of the potential 'harms' from implementing interventions that have no evidence of benefit is an important next step to help improve uptake
- Whilst establishing the effectiveness of recruitment or retention strategies is important, the high costs of research waste and limited public finance means that cost considerations around SWATs are just as important. With only one retention strategy having high GRADE certainty of cost-effectiveness, we encourage trial teams to undertake streamlined economic evaluations alongside all their SWATs in the future, for strategies shown to be effective, as well as those that are ineffective. For cost-effectiveness, trial teams should look to report the cost per additional participant recruited or retained (that is, the incremental cost-effectiveness ratio). Value of information analyses can help determine the need for further SWAT evidence where several SWATs already exist
- Many trial teams wish to contribute to developing the evidence base by undertaking non-randomised SWATs. Future work that informs the development of guidance for undertaking non-randomised SWATs would be helpful
- Working with trial teams to develop engagement strategies and training to undertake SWATs would be beneficial. Audience specific guidance should be developed to support SWAT research
- Trial teams have expressed they want to undertake SWATs that are important and necessary to increase the evidence base. Collaboration with funders, working groups involved in priority setting, and trial teams are needed to develop a mechanism to communicate this dynamic and evolving information once priority SWATs have been identified
- Work is needed to identify the barriers that teams have when undertaking a SWAT, and strategies and solutions for addressing these barriers should be identified and implemented



- Continued and proactive collaboration is needed with working groups to enable networking, and collaboration with teams undertaking SWAT research
- Reporting guidance is needed to support teams when writing publications to ensure sufficient information is included to GRADE evaluation can be undertaken

### *Study registration*

All SWATs in the PROMETHEUS programme had to be registered with the Northern Ireland Network for Trials Methodology Research SWAT Repository.

### *Funding details*

This project was funded by the Medical Research Council (MR/R013748/1) and National Institute for Health Research as part of the MRC–NIHR Methodology Research programme.

## Introduction

### *What is covered in this report?*

This report focuses on randomised Studies Within A Trial (SWATs) testing the effectiveness of recruitment and retention strategies. The structure of this report begins by presenting the background to the PROMETHEUS programme (*see Chapter 1*). In Chapter 2, we present the methods used in PROMETHEUS, followed by the results (*see Chapter 3*) and then the discussion of the programme (*see Chapter 4*).

Subsequent Chapters focus on the lessons learnt from PROMETHEUS and how this can inform future work focused on SWATs. This includes lessons learnt for; patient and public involvement (PPI) (*see Chapter 5*), the development of SWATs and its funding (*see Chapter 6*), and ethical and other governance approvals (*see Chapter 7*). In Chapter 8, we focus on different approaches to implementing SWATs, using case studies of standalone SWATs, as well as case studies of coordinated SWATs, undertaken in a coordinated way across multiple host trials simultaneously. Chapter 9 reports lessons learnt for analysing SWATs, and Chapter 10 on lessons learnt for reporting SWATs.

At each point, we provide an outline of key points to consider, using specific examples of work from the PROMETHEUS programme to illustrate our point. Wherever relevant, we highlight what different stakeholder groups such as; trial management teams, PPI partners, trial oversight committees, those overseeing trial governance, statisticians, journals, and peer reviewers can learn from the PROMETHEUS experience.

Finally, we discuss recommendations for future research practice, direction, and support (*see Chapter 11*), and then present conclusions (*see Chapter 12*).

### *Who should use this report?*

This report has been written for anyone with an interest in using randomised controlled trials to inform evidence-based health and social care. Lessons learnt that we detail in this report will be of use to those interested in how the design and delivery of future trials can be improved to benefit patient health and social care. This includes individuals and organisations involved in funding trials; those who design, plan and undertake trials such as Chief Investigators and Trial Managers; trial oversight committees; those providing ethical and other regulatory approvals; patient and public partners in trials; trial infrastructure organisations supporting trial delivery such as the UK Clinical Research Networks, Trials Units and the Research Design Service. The report will be of use to those recruiting and retaining trial participants, as well as those interested in trials methodology.

The report has been written for those with an understanding of trials who are new to SWATs; those with some experience of SWATs who want to learn more; and those with SWAT experience who may wish to use our experience to consider possible methodological innovations of their own. This report is also written for journal editors publishing SWATs, as well as those undertaking peer reviews of SWATs.

## Chapter 1: Background to the PROMETHEUS programme

### *The importance of randomised controlled trials*

Randomised controlled trials ('trials') attempt to improve the health and care of populations and minimise the potentially harmful effects of treatments and interventions. Trials are undertaken on the basis that there are uncertainties about the effects of treatments, and such treatments are tested to adequately reduce these uncertainties. Consequently, trials are usually accepted as the foundation for evidence-based practice,<sup>1</sup> the most complete paradigm for delivering safe and effective health and care for patients and citizens. Globally in 2022, there were more than 400,000 trials being undertaken across 220 countries.<sup>2</sup> The UK National Institute for Health and Social Care Research (NIHR) invested £250 million on research, predominantly in the form of trials, in 2019/2020, and £373 million in 2020/2021.<sup>3,4</sup> To be efficient, trials need to recruit large numbers of participants quickly - often in excess of 1000 participants.<sup>5</sup> To have external validity, a generalisable sample needs to be recruited from all relevant patient and population groups, and to avoid post-randomisation selection bias, retention needs to be kept as high as possible.

### *Recruitment and retention difficulties in trials*

Despite the importance of trials, they are susceptible to both poor participant recruitment and retention. As a consequence of this, many trials fail to recruit to both time and budget, often with lower participant retention than anticipated.<sup>5</sup> The cost of poor participant recruitment can be huge,<sup>6</sup> contributing to significant research waste.<sup>7,8</sup> Furthermore, poor recruitment has a detrimental impact on mortality and morbidity as this increases the time to the implementation of effective care and treatments. An analysis of the RECOVERY trial of COVID-19 treatments, which recruited only 10% of eligible patients, showed that had recruitment been 50% or more of eligible patients, there would have been a significant improvement in lives saved due to earlier reporting.<sup>9</sup> Similarly, poor participant retention, which is often overshadowed by the research focus of poor participant recruitment,<sup>10</sup> can also be detrimental to the success of a study. Poor retention may lead to the underpowering of a study and cause the estimates of an intervention's effect to be biased.<sup>11</sup> Aligned with this, a priority-setting exercise involving 85% of UK Clinical Trials Units (CTUs) placed recruitment and retention as the top two priorities for methodological research,<sup>12</sup> cementing the need to identify and implement tools within trials that help to improve both.

### *SWAT methodology*

To date, there has been a distinct lack of robust evidence to inform trial design and conduct, including for recruitment and retention. However, there is an increasing international movement to improve the efficiency and successful delivery of trials by using robust research methods.<sup>13-15</sup> These

robust studies use methods that are embedded within real ‘host’ trials and are referred to as ‘Studies Within A Trial’ (SWATs). A SWAT has been defined as a: “*self-contained study that has been embedded within a host trial with the aim of evaluating or exploring alternative ways of delivering or organising a particular trial process*”.<sup>15</sup>

Treweek *et al.*, (2018)<sup>15</sup> outline several key features of a SWAT. Firstly, SWATs aim to resolve key uncertainties around trial processes. Secondly, a SWAT is embedded within a host trial, but it should have a formal protocol, in the same way as the host trial. Thirdly, a SWAT must not affect the scientific integrity of the host trial, its rationale, or outcome measures. Fourthly, a SWAT can either be tested in a single trial or embedded across multiple host trials in a coordinated way. Finally, SWATs provide evidence to inform the design and conduct of future trials; however, they may also produce evidence to inform decisions about the host trial.

SWATs can adopt a range of different methods including qualitative,<sup>16, 17</sup> mixed methods,<sup>18</sup> and before and after designs.<sup>19</sup> SWATs can be used to understand and refine implementation processes.<sup>20</sup> SWATs can also be non-randomised<sup>21</sup> or randomised.<sup>22</sup>

#### *Addressing recruitment and retention difficulties using SWATs*

Randomised SWATs are the most rigorous method for evaluating the effectiveness of strategies for improving participant recruitment and retention in trials.<sup>23</sup> Two Cochrane reviews identified 150 SWATs of strategies to increase recruitment and/or retention in trials;<sup>23, 24</sup> however, effective, evidence-based strategies are rare and where evaluations exist, they tend to occur in the context of a single trial, meaning that they usually have limited statistical power and their effects across different trial contexts are unclear.<sup>23</sup> The most recently published Cochrane review on retention interventions concluded that there was no high-certainty evidence for any of the evaluated strategies, as assessed by GRADE.<sup>24</sup>

Prior to the PROMETHEUS programme was the Medical Research Council (MRC) START programme;<sup>25, 26</sup> a feasibility study that successfully developed the conceptual, methodological and logistical framework to improve recruitment through the embedding of two recruitment strategies in twelve host trials in primary care and developed reporting guidelines for embedded trials.<sup>25, 27</sup> In addition, since 2014, the Health Research Board – Trials Methodology Research Network (HRB-TMRN) Ireland has supported and funded Irish researchers to conduct methodological studies to improve the efficient conduct of future trials including SWATs, offering funding of up to €25,000 for evaluations.<sup>28</sup> More recently the NIHR Health Technology Assessment (HTA) programme is encouraging applications to embed at least one high-quality SWAT with funding of up to £10,000 per host trial.<sup>29</sup> The TRials Engagement in Children and Adolescents (TRECA) study, also funded by the

NIHR, aimed to develop multimedia interventions to improve the quality of decision-making about recruitment to trials involving children and young people with long-term conditions and subsequently tested them using SWATs.<sup>30</sup>

### *Aims and objectives*

The PROMETHEUS programme aimed to build on the work of these previous initiatives by making the embedding of recruitment and retention SWATs within host trials standard practice across multiple CTUs, working to at least double the number of SWATs produced by the MRC START project, given a similar level of funding. This was to be achieved through pump-priming and facilitating the start of at least 25 SWATs within 30 months. The ultimate aim was to make the inclusion of SWATs routine practice when conducting a trial in a CTU.

## Chapter 2: PROMETHEUS methods

### *PROMETHEUS preparatory work*

Prior to the initiation of the PROMETHEUS programme, a network of ten CTUs, one primary care research centre in the UK, and the HRB-TMRN in the Republic of Ireland was established, each of whom committed to embedding either a randomised controlled recruitment and/or retention SWAT within at least two host trials.

Promising recruitment and retention strategies (which had some evidence of benefit but with substantial uncertainty) were identified from a variety of sources. These sources included Cochrane systematic reviews,<sup>23, 31</sup> the SWAT Repository Store,<sup>32</sup> and the priorities of recruitment and retention strategies identified by CTUs.<sup>33</sup> In addition, the PRioRiTy list of top ten unanswered questions on trial recruitment, for which there is no current evidence, was also reviewed.<sup>34</sup> A PPI panel was also convened to highlight the top priority strategies for evaluation (*see Chapter 5*).

The identified strategies were then prioritised if they met one or more of the following criteria:

1. Strategies that had already been evaluated with results published in peer-reviewed journal publications
2. Strategies currently under evaluation
3. Easy to implement strategies, i.e., those requiring little additional resource (input or cost) from the host trial, as assessed by the PROMETHEUS investigators. Such strategies might not have a large impact; however, they might still make a useful contribution to the evidence base, based on marginal gains and cost-effectiveness
4. Had the potential to significantly impact participant retention or recruitment (which are often the more challenging, expensive strategies to implement)
5. Strategies identified by host trial teams as suitable for testing in a SWAT within their trial

Prioritisation decisions were made through group discussion and consensus. These priorities formed an initial strategy priority list of seven recruitment and eight retention strategies (Table 1). The priority of these strategies was reassessed and rearranged accordingly throughout the programme, based on emerging SWAT evidence.

Table 1: List of key recruitment and retention questions in priority order

Recruitment strategies
1. What is the effect of adding a pen printed with the trial/university logo to the trial invitation on recruitment rates (SWAT 37)?
2. What is the impact of recruitment sites receiving an extra trial coordinator visit on recruitment rates (SWAT 27)?
3. What is the effectiveness of a brief participant information leaflet (PIL) versus standard length PIL on participant recruitment rates? (SWAT 137)
4. What is the impact of a training workshop for staff recruiting patients into trials on recruitment rates (SWAT 111)?
5. What is the effect of offering financial incentives to potential trial participants on recruitment rates (SWAT 59)?
6. What is the effect of mentioning scarcity of trial places in invitation letters on recruitment of trial participants (SWAT 60)?
7. What is the effectiveness of telephoning people who do not respond to a postal invitation on recruitment to randomised trials (SWAT 61)?
Retention strategies
1. What is the effect of adding a pen printed with the trial/university logo to the trial invitation on retention rates (SWAT 37)?
2. What is the effectiveness of a theoretically informed cover letter on improving response rates to annual postal questionnaires (SWAT 24)?
3. What is the effect of a text message notification versus no text message on questionnaire response rates (SWAT 25/SWAT 31)?
4. What is the effectiveness of a personalised text message versus a standard text message for promoting response to postal questionnaires (SWAT 35)?
5. What is the effect of timing text message prompts to increase trial participant response to postal questionnaires (SWAT 44)?
6. What is the effectiveness of sending pre-notification cards (letters/email) to trial participants 1-month (2 weeks) before outcome measurement to improve retention (SWAT 76 / SWAT 86)?
7. What is the impact of receiving a social incentive strategy cover letter compared with a standard covering letter on response rate to postal questionnaires? (SWAT 144)
8. Do courtesy telephone calls to trial participants following enrolment increase future retention rates? (SWAT 114)

Note: SWAT number correlates to the registration number listed on the Northern Ireland Network for Trials

Methodology Research SWAT Store <sup>32</sup>

### Eligibility criteria

#### Host trials

To be eligible for PROMETHEUS funding, host trials were required to meet the following criteria:



1. Registered or eligible for registration on the UK Clinical Research Network Portfolio<sup>35</sup>
2. In the planning phase; recruiting or following up participants, or be in the process of applying for ethics permission (i.e., any trial stage bar the point of trial closure)
3. Willing to apply for ethics permission or amendment to undertake at least one SWAT of a recruitment or retention strategy
4. Willing to randomise and deliver the recruitment or retention strategy according to a shared protocol
5. Willing to share data with the PROMETHEUS team including patient-level data to allow individual patient-level meta-analysis
6. Willing to use or register their SWAT on the Northern Ireland Network for Trials Methodology Research SWAT Repository, a free-to-use online database of ongoing SWATs if the strategy being evaluated is not already registered<sup>32</sup>
7. Able to provide evidence of funding for the host trial (such as a letter from the funder)

#### *SWAT interventions*

PROMETHEUS prioritised a broad list of recruitment and retention strategies that could be evaluated (Table 1), however host trials could also evaluate their own strategies if they wished to. Support included: assistance in writing SWAT protocols; provision of templates and guidance in achieving Research Ethics Committee (REC) and Health Research Authority (HRA) approvals; guidance on writing and submitting SWATs for publication.

#### *PROMETHEUS programme outcomes*

PROMETHEUS was designed to analyse the effectiveness of recruitment and retention interventions within the context of a single host trial, as well as across a range of different trials through the synthesis of similar SWATs results.

#### *Sample size*

The programme aimed to implement at least 25 SWATs across eligible host trials. Generally, as the sample size of SWATs are driven by the size of the host trial, single SWATs are often underpowered; a known and accepted attribute of most SWATs is that whilst sample size calculations may be made, formal power calculations are not required.<sup>36</sup> Instead, the SWAT's sample size is usually driven by the host trial. Therefore, it was always planned that, where appropriate, SWAT intervention results were to be aggregated. To reduce study heterogeneity, studies were encouraged to follow common SWAT protocols to promote homogeneity in interventions in readiness for pooling data (*see Chapter 9*). This aggregation of SWATs was intended to help with the detection of small differences and

provide evidence as to whether results might generalise to a range of contexts, providing evidence with greater external validity.

#### *Host trial recruitment*

Eligible host trials were largely identified and recruited through a combination of the programme collaborators, advertisement on the University of York Trials Unit (YTU) web page, emails to all registered UK CTUs and conference presentations such as at the International Clinical Trials Methodology Conference 2019. Following their identification, trial teams were provided with a PROMETHEUS Information Sheet, invited to submit an expression of interest form, and apply for funding from the programme for up to £5,000 per SWAT embedded in their host trial. The funding application consisted of submitting a SWAT protocol, including a project timetable and an outline of costs. To support the development of the SWAT protocols, host trial teams were provided with the opportunity to meet with PROMETHEUS team members for support to determine an appropriate recruitment and/or retention strategy.

Two independent members of the PROMETHEUS programme peer-reviewed each host trial application and protocol to ensure methodologically robust replicable research was planned. Reviewers were asked to report their peer review comments and scores using a Peer Review Assessment Form, which was adapted from the peer review form used by the HRB-TMRN.<sup>28</sup> Peer reviewers were asked to comment on the following:

- 1) Eligibility,
- 2) Priority and scientific quality,
- 3) Costings, and
- 4) Overall rating of the application.

#### *Randomisation*

Typically, randomisation was completed by the host trial team and therefore the methods of randomisation, allocation concealment, and implementation were applied at the level of the host trial.

#### *Blinding*

Often it is deemed that SWATs do not require individuals to provide participation consent due to them being generally low risk, rarely imposing additional burden on a participant and due to potential risks of confusing the participant as to what they are consenting to.<sup>23</sup> As such, participants

were not blind to their received intervention (e.g., a thank-you card or SMS), but they were unaware of their participation in the SWAT.

#### *Data collection methods*

SWAT data were collected in line with both the relevant SWAT and host trial protocols. The PROMETHEUS team provided trial teams with support to determine appropriate data collection methods where required. The funded teams were asked to provide the PROMETHEUS team with the data for each SWAT, with the aim of collating these in pooled analyses.

#### *Statistical methods*

To enable pooling of findings, a standardised framework was established, similar to that of MRC START.<sup>25, 27</sup> Template protocols with standard outcomes and a template Statistical Analysis Plan (SAP) were designed. Trial teams were encouraged to analyse the outcomes in terms of absolute differences in recruitment and retention rates, as appropriate and to provide a cost per recruited or retained participant where possible. The primary outcome measure was compared across intervention and control arms using logistic regression. For SWATs of retention, teams also looked at elements such as: time to, and completeness of, and responses to follow-up outcome measures. Where possible, odds ratios from multiple SWATs on the same or similar interventions were combined using a random effects meta-analysis, ideally using a one-stage approach if the individual patient-level data was available.

#### *Data monitoring*

A Project Management Group (PMG) was established to oversee the management of the SWATs work consisting of the project Statistician, Research Fellows, and other co-applicants, and was chaired by the Chief Investigator (author DJT). The role of the PMG was to monitor all aspects of the conduct and progress of the study. The PMG met quarterly by teleconference, with annual face-to-face meetings where feasible, meeting more frequently when there was a need to monitor the programme's progress more closely.

#### *Harms*

Due to the nature of the SWAT strategies, no strategy was likely to be responsible for any health-related adverse events. Therefore, adverse event reporting and review remained with the host trial teams with data not being collected for the PROMETHEUS programme.

#### *Auditing*

The PROMETHEUS team maintained contact with each of the host trials throughout the duration of their SWAT conduct, requesting and routinely monitoring their SWAT progress.

### *Research ethics approval*

Each individual SWAT obtained approval from the host trial's research ethics committee, and institutional governance committees as needed. This approval was sought as part of the initial study application or as an amendment, depending on the status of the study at the time of SWAT implementation and/or the nature of the SWAT. Careful consideration was given, during the application as to whether informed consent was required for SWAT participation. Most SWATs also required approval from NHS site research and development (R&D) departments, in line with HRA procedure, prior to implementation. Given that participants may not have provided informed consent for SWAT participation, patient identifiable information was held by the host trial and not released as part of the SWAT data set.

### *Consent or assent*

Participants were not informed of their involvement as it was not appropriate and may have affected the host trial outcomes by resentful demoralisation.<sup>37</sup> Providing the SWAT did not involve knowingly withholding pertinent information about participant involvement within it, and involvement was non-invasive, it was deemed appropriate that patients were not informed about the SWAT and informed consent for involvement was not obtained.

### *Confidentiality*

All SWAT data transferred to the PROMETHEUS team via trial teams was done anonymously; for instance, by removing identifiers such as date of birth (e.g., simply putting age) and participant identity number (e.g., hospital number) and then randomly sorting the data. This ensured that it would not be possible to re-identify participants in the dataset, in line with General Data Protection Regulation (GDPR) requirements with individual participants being identified by their host trial identification number only, or linked SWAT identification number if the host trial identification were removed. All electronic records were stored on a secure server.

### *Access to data*

PROMETHEUS access to the data for each individual SWAT was obtained via signed data sharing agreements in line with the principles for access to, and use of, MRC funded research data report,<sup>38</sup> negotiated at the outset. Ongoing access to the data depended on agreement with the individual host trial teams, with agreements about authorship and dissemination of results from the individual studies and the combined dataset. Subject to the consent of the host trial teams, primary data was made available in an anonymised format suitable for release in the public domain.

## Chapter 3: PROMETHEUS results

*Host trials funded, numbers of recruitment SWATs funded, number of retention SWATs funded*

In total, the PROMETHEUS programme supported 42 SWATs, which were implemented within 31 host trials, across 13 CTUs and Research Centres (Figure 1).




Figure 1 : Flow chart of SWATs funded by the PROMETHEUS programme

Collectively these 31 host trials spanned 17 research areas (Table 2) (see Appendix 1 for a full list of the funded host trials and their characteristics). It is important to note that not all participants in host trials were available for nor included in the SWATs. Five of the host trials implemented more

than one SWAT, a further four implemented a factorial design SWAT, and 12 were involved in the coordinated evaluation of two SWAT strategies.

Table 2: The research area of each of the SWATs funded by the PROMETHEUS programme

Research area	Host trial (sample size)**	Number of host trials	Number of SWATs
Surgical	ACTIVE (334) DISC <sup>a</sup> (710) L1FE (600) ProFHER-2 <sup>a</sup> (380) START:REACTS (221) UK FROST (503) C-Gall (430) MAGIC (1650) PurE (1044)	9	11
Fall reduction	OTIS <sup>a</sup> (1299) SSHeW (4400)	2	3
Orthopaedic (rehabilitation)	ARTISAN (478) KReBS <sup>a</sup> (2600)	2	3
Respiratory	CLEAR <sup>a</sup> (380)	1	3
Smoking cessation	CPIT-III (940) MiQuit-3 <sup>a</sup> (692)	2	3
Wound care	SWHSI-2 <sup>a</sup> (696)	1	3
Oncology (screening and treatment)	IntAct (880) POSNOC <sup>a</sup> (1900) ActWELL (552)	3	3
Community pharmacy	CHAMP-1 <sup>a</sup> (820)	1	2
Physiotherapy	PEP-TALK (250) GRASP (704)	2	2
Primary care (Signs and symptoms)	MSS3 <sup>a</sup> (376)	1	2
Rheumatology	TOPaZ (380) WORKWELL (240)	2	2
Urology	FUTURE (1096) SARC (118)	2	2
Oral Health	REFLECT (1174)	1	1
Gastrointestinal	IBD-BOOST (680)	1	1
Gynaecology	VITA (1900)	1	1

<sup>a</sup>Host trials which implemented more than 1 SWAT; \*\* Not all of the host trial participants were necessarily included in a SWAT.

In total, 12 of the funded SWATs evaluated recruitment strategies (*See Results: findings from the funded SWATs Tables 4&5*) and 30 SWATs evaluated retention strategies (*See Results: findings from the funded SWATs Tables 6&7*).

#### *Recruitment strategies*

The 12 recruitment strategies have been grouped into the recruitment research domains framework mapped by Online Resource for Research in Clinical triAls (ORRCA).<sup>39</sup> The six ORRCA domains covered recruitment strategies aimed at the following:

- A) Trial design
- B) Pre-trial planning
- C) Trial conduct
- D) Recruitment information needs
- E) Recruiter differences
- F) Incentives

Our classification resulted in the funded recruitment strategies being classified under domain D, 'recruitment information needs' or domain F, incentives (*See Results: findings from the funded SWATs Tables 4&5*). Within domain D, individual recruitment strategies were either classified into subdomains as 'D1. Researcher training needs' (4 SWATs) or 'D2. Participant Information Sheet and Consent Form' (7 SWATs). The only recruitment strategy to fall under domain F was classified into the sub-domain of 'F3. Participant non– monetary incentives'.

#### *Retention strategies*

The 30 retention strategies have been grouped into the retention research domains framework mapped by Online Resource for Research in Clinical triAls (ORRCA2).<sup>40</sup> The five ORRCA domains covered retention strategies aimed at the following:

- A) Data collection
- B) Participants
- C) Sites and site staff
- D) Central Study Management
- E) Study design

Our classification resulted in all the funded retention strategies being classified under domains A 'Data collection' or Domain B 'Participants'. There was only one SWAT classified under domain A, and this SWAT fell under the sub-domain of 'A1. Questionnaire design'. The remaining SWATs in Domain B (Participants) were classified under the sub-domains of 'B1. Reminders and prompts' (13 SWATs); 'B2. Monetary incentives' (1 SWAT); 'B3. Non-monetary incentives' (3 SWATs); and 'B4. Maintaining participant engagement' (12 SWATs).

Results were expected for 36 SWATs (*see SWAT Results*). Six SWATs could not be completed: two SWATs encountered technical issues so had to be abandoned (factorial SWAT);<sup>41</sup> one SWAT was embedded in a host trial that was closed early due to poor recruitment;<sup>42</sup> one SWAT was embedded in a host trial that stopped recruitment early, having answered its question;<sup>43</sup> and three further SWATs could not proceed due to the COVID-19 pandemic, which forced the host trial to change its mode of following-up participants.<sup>42, 44, 45</sup>

We also identified an opportunity to test the methodological feasibility of a coordinated SWAT design, which involved the implementation of a recruitment or retention strategy within multiple pre-identified host trials simultaneously. In this coordinated SWAT, REC approval only needed to be obtained once to allow SWAT implementation within all of the included host trials. The results from each host trial are reported simultaneously within one publication, allowing a more rapid increase of the evidence base. This approach was tested in two SWATs; one evaluated the effect of clinician recruitment training on participant recruitment (SWAT 111),<sup>46</sup> and the other sending of Christmas cards to participants on participant retention (SWAT 82),<sup>47</sup> with each being implemented within four and eight host trials, respectively (*See Chapter 8 for case studies of these SWATs*).<sup>48, 49</sup>

## *SWAT costs*

### *Individual SWATs*

The average cost of funding requested for a SWAT within the PROMETHEUS programme was £2,600 (range £500 to £5000). This figure is calculated from 29 SWATs – each element of factorial SWAT was counted separately, with the costs split between them, the coordinated SWATs (n=12 separate SWATs) were not included due to their costs being unrepresentative of the cost a single SWAT, and one SWAT requested no funding after applying due to such low funds needed (approximately £300). The average cost of consumables (such as pens) was £867 (from 15 SWATs), and £1753 to cover staff time (from 13 SWATs) – it was not always possible to distinguish the separate costs from the funding applications. As many of the PROMETHEUS supported SWATs were performed by York Trials Unit, where the PROMETHEUS programme was based, it is anticipated that the costs were



underestimated, i.e., often did not account for staff time. When excluding YTU SWATs, the average cost rises to £3,535 - with mean staff cost being £2,359 (data from 17 and 9 SWATs respectively).

During the discussions with host trial teams with regards to embedding a SWAT, it was apparent that many teams required additional support and insight regarding how long SWAT-specific tasks may take to enable them to cost activities accurately. To assist with this, example costings were developed for two individual SWATs undertaken as part of the PROMETHEUS programme, as shown in Table 3. These highlight that within different SWATs and different CTUs the associated times required may vary, depending on level of experience and infrastructure, for example, if a Trial Manager was having to manually send text reminders, as opposed to an automated system, the associated time costs would be vastly different.

Table 3: Examples of funding for staff time required for individual SWATs

Type of cost	Recruitment SWAT example	Retention SWAT example
Set up (including amending protocol and documentation; setting up and testing randomisation; providing training)	5 days of a Trial Manager's & 3 days of a Programmer's time	5 days of a Trial Manager's time
Activities involved in the SWAT (e.g., undertaking randomisation, implementing the intervention)	-	15 days of a Trial Manager's time
Data Cleaning & Analysis & Write up	2 days of a Trial Statistician's time	10 - 15 days of a Trial Statistician's time

#### *Coordinated SWATs*

The total cost of the recruitment training SWAT was £10,668. This can be broken down into £2,188.57 for staff time to prepare and deliver the training and £8,479.43 for consumables, including travel costs to deliver and attend the training, venue hire and subsistence, as well as thank you vouchers to participants for completing the follow-up questionnaires. The costs of developing the training package were not included.

The cost calculated for the Christmas card SWAT was £1,306.40 – an average cost of £0.76 per card sent. This included time for staff time to prepare the cards, and for consumables such as the printing, postage, and delivery of the cards.

However, the cost estimates for both SWATs do not include the following, which need to be accounted for when planning to undertake a SWAT of this design:

- Coordination from a central point (i.e., to liaise with host trials about involvement and undertaking the SWAT)
- Data preparation and sharing by the host teams
- Analysis and data cleaning
- Write up

### *Results: findings from the funded SWATs*

#### *Results of recruitment SWATs*

Twelve recruitment SWATs were funded within nine host trials. Six of the SWATs had published findings (Table 4) and six of the SWATs were ongoing at the time of writing this manuscript (Table 5). The published SWATs consisted of only two publications: a 2x2 factorial SWAT,<sup>50</sup> and a coordinated SWAT involving four host trials, which tested the feasibility of staff training to improve participant recruitment to surgical RCTs.<sup>51</sup> There was no evidence of a significant difference in recruitment rates in any of the strategies tested. However, the staff training SWAT found that it was feasible to randomise sites across four surgical trials in a coordinated SWAT design. This SWAT also reported that in the intervention group, there was evidence of increased staff confidence when pre- and post-training scores were compared – but had a limited sample size and so requires further replications.

Table 4: Host trials and results of completed SWATs evaluating recruitment strategies

<b>Number of SWATs</b>	<b>Host trial acronym</b>	<b>SWAT domain</b>	<b>Brief SWAT description</b>	<b>SWAT outcomes</b>	<b>Published SWAT Results</b>
------------------------	---------------------------	--------------------	-------------------------------	----------------------	-------------------------------

4	DISC <sup>52</sup> IntAct <sup>53</sup> ProFHER-2 <sup>54</sup> START:REACTS <sup>55</sup>	D1. Researcher training needs	Staff training to improve participant recruitment into surgical randomised trials (SWAT 111) <sup>46</sup>	<p><i>Primary:</i> The feasibility of recruiting sites across multiple surgical trials in a coordinated way</p> <p><i>Secondary:</i></p> <ol style="list-style-type: none"> <li>1) /Numbers/ types of staff enrolled</li> <li>2) Attendance at training, training acceptability</li> <li>3) Staff confidence in recruiting</li> <li>4) Participant recruitment rates six months later.</li> </ol>	Parker et al., (2022) <sup>51</sup> : Four RCTs (33%) comprising 91 sites participated. Of these, 29 sites agreed to participate (32%) and were randomised to intervention (15 sites, 29 staff) or control (14 sites, 29 staff). Research nurses attended and found the training to be acceptable; no surgeons attended. In the intervention group, there was evidence of increased confidence when pre and post training scores were compared (mean difference in change 1.42; 95% CI 0.56, 2.27; p = 0.002). There was no effect on recruitment rate.
---	---	-------------------------------	--	---	---

2	CLEAR <sup>56</sup>	D2. Participant Information Sheet and Consent Form	<p>2x2 factorial SWAT design:</p> <p>Participant invitation letter with personal wet signature vs generic signature (variant of SWAT 3) <sup>57</sup></p> <p>Participant study invitation including a generic doctor-patient photograph vs including no photograph (SWAT 53) <sup>58</sup></p>	<p><i>Primary:</i> Proportion of invited patients who joined the trial.</p> <p><i>Secondary:</i> Proportion of patients retained in the trial.</p>	<p>Anand et al., (2022) <sup>50</sup>: 368 letters were given to potential participants in the CLEAR trial and 121 (33%) joined. Proportions for each randomised group were generic signature and no photograph: 38% (33/88); generic signature and photograph: 32% (28/88); wet-ink personal signature and no photograph: 29% (26/91); wet-ink personal signature and photograph: 34% (34/101). There was no evidence of a significant difference in recruitment between those receiving the patient invitation letter containing a wet-ink versus generic signature (odds ratio (OR): 0.86, 95% confidence intervals (CI): 0.55 to 1.32, p=0.49) or photograph versus no photograph (OR: 0.99, 95% CI: 0.64 to 1.53, p=0.97). Retention was similar for the wet-ink and generic signature groups (OR: 1.20, 95% CI: 0.35 to 4.16, p=0.77) but significantly better when a photograph was used (OR: 5.40, 95% CI: 1.12 to 26.15, p=0.04, based on 2 withdrawals in the photograph group versus 9 in the no photograph group).</p>
---	---------------------	--	--	--	--

Table 5: Ongoing SWATs evaluating recruitment strategies

<b>Number of SWATs</b>	<b>Host trial acronym</b>	<b>SWAT domain</b>	<b>SWAT question</b>	<b>SWAT outcomes</b>	<b>Expected completion data and details of delay if applicable</b>
------------------------	---------------------------	--------------------	----------------------	----------------------	--

1	IBD-BOOST <sup>59</sup>	D2. Participant Information Sheet and Consent Form	Brief participant information sheet (PIS) provided in addition to a standard-length PIS (SWAT 137) <sup>60</sup>	<p>Primary: Recruitment rate</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Proportion of patients in each group who return an expression of interest form</li> <li>2) Cost-effectiveness of the interventions</li> <li>3) Proportion of patients who return an expression of interest form but are not randomised due to a) ineligibility or b) non-consent</li> <li>4) Time taken to respond to an invitation to participate in the host trial</li> </ol>	Completion date is expected to be December 2022. This SWAT had a delay in site set up due to COVID-19
---	-------------------------	--	--	--	---

2	MSS3 <sup>61</sup>	D2. Participant Information Sheet and Consent Form  F3. Participant non-monetary incentives	2x2 factorial design:  Brief participant information sheet (PIS) provided in addition to a standard length PIS (SWAT 137)  Inclusion or not of a trial logo branded pen (SWAT 137) <sup>60</sup>	Primary: Recruitment rate  Secondary: 1) Proportion of patients who return an expression of interest form 2) Cost-effectiveness of the interventions 3) Proportion of patients who return an expression of interest form but are not randomised due to a) ineligibility or b) non-consent 4) Time taken to respond to an invitation to participate in the host trial.	Recruitment was delayed due to COVID 19, however data collection is now complete. The SWAT is undergoing analysis and is being prepared for publication
1	POSNO <sup>62</sup>	D2. Participant Information Sheet and Consent Form	Addition of a pictorial aid to the patient information leaflet (PIL) (SWAT 102) <sup>63</sup>	Primary: Proportion of patients randomised to the trial.	Data collection is complete



1	SARC <sup>64</sup>	D2. Participant Information Sheet and Consent Form	Optimised PIS vs conventional PIS (SWAT 101)	<p>Primary: Proportion of patients who consent to take part in the interventional trial.</p> <p>Secondary: Qualitative outcomes assessing the impact/value of the PIL in the decision making.</p>	Data collection completed in May 2022; in analysis stage.
1	SWHSI-2 <sup>65</sup>	D2. Participant Information Sheet and Consent Form	Inclusion of an infographic provided in addition to a standard patient information leaflet (SWAT 119) <sup>66</sup>	<p>Primary: Recruitment rate</p> <p>Secondary:</p> <p>1) Proportion of patients who are screened for the study but are not randomised due to a) ineligibility or b) non-consent</p> <p>2) Cost-effectiveness</p>	The SWAT is expected to finish in December 2022

### *Results of retention SWATs*

Thirty SWATs testing retention strategies were funded in 25 host trials. At the time of writing, 21 SWATs had concluded and been published, including two 2x2 factorial SWATs,<sup>67, 68</sup> and the coordinated SWAT consisting of eight trials.<sup>49</sup> (Table 6) Three SWATs were still ongoing (Table 7). Unfortunately, six further SWATs were stopped early (Table 8) and will produce no results: one host trial stopped early prior to reaching its recruitment target,<sup>43</sup> a 2x2 factorial SWAT of text messaging strategies encountered a significant system fault with the text messaging software (two SWATs),<sup>41</sup> and three SWATs could not proceed due to the COVID-19 pandemic.<sup>42, 44, 45</sup>

Table 6: Host trials and results of completed SWATs evaluating retention strategies

Number of SWATs	Host trial acronym	SWAT domain	Brief SWAT description	SWAT outcomes	Published SWAT Results
1	PEP-TALK <sup>69</sup>	A1. Questionnaire design	Printing the primary outcome measure on pink paper versus on white paper (SWAT 110) <sup>70</sup>	<p>Primary: host trial primary outcome measure completion.</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Number of reminders sent</li> <li>2) Proportion of remaining questions completed</li> <li>3) Overall questionnaire returns</li> </ol>	<p>Ooms et al., (2022) <sup>71</sup>: 176 participants were randomised: 88 received pink paper, 88 white paper. Host trial primary outcome measures were returned by 84.1% (74/88 participants) in the pink paper group and by 90.9% (80/88 participants) in the white paper group (risk ratio, 0.92 (95% CI 0.80, 1.06); p = .24). Reminders were sent to 48.9% (43/88 participants) in the pink paper group and to 30.7% (27/88 participants) in the white paper group (risk ratio 1.59 (95% CI 1.09, 2.33); p = .01). No other results were statistically significant.</p>

1	UK FROST <sup>72</sup>	B1. Reminders and prompts	Timing of text message prompts (reminder received prior to questionnaire arrival or 4 days later) (SWAT 44) <sup>73</sup>	<p>Primary: proportion of participants who returned a valid questionnaire.</p> <p>Secondary: A systematic review was undertaken to identify other embedded trials to perform a meta-analysis.</p>	<p>Partha Sarathy et al., (2020)<sup>74</sup>: In the pre-notification arm, 122/135 (90.4%) participants returned a valid questionnaire compared with 119/134 (88.8%) in the post-notification arm (difference of -1.6%; 95% CI of difference: -8.9%, 5.7%). There was no difference in time to response (HR = 1.04; 95% CI: 0.80 to 1.34) or need for additional reminders (OR = 0.71; 95% CI: 0.43 to 1.17). When combined with two RCTs in a meta-analysis, no difference in response rates between groups, in relation to reminders, was observed (OR = 0.78 95% CI: 0.42 to 1.45).</p>
---	------------------------	---------------------------	---	---	---

1	GRASP <sup>75</sup>	B1. Reminders and prompts	Personalised text message versus a standard text message (SWAT 35) <sup>76</sup>	<p>Primary: Questionnaire response rate at six months</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Time to response</li> <li>2) Proportion of participants sent a reminder follow-up questionnaire</li> <li>3) Cost</li> </ol>	<p>Cureton et al., (2021) <sup>77</sup>: 618 participants were randomised to a personalised (n = 309) or standard (n = 309) text message. The overall questionnaire response rate was 87% (n = 537/618); 90% (n = 277/309) of participants responded in the personalised text message group compared to 84% (n = 260/309) in the standard text message group (relative risk (RR) 1.07; 95% CI 1.00 to 1.13). Participants randomised to receive the personalised text message were more likely to return their initial postal questionnaire than those who received the standard text message (n = 185/309; 60% vs. n = 160/309; 52%) (RR 1.16; 95% CI 1.00 to 1.33); this represents an absolute percentage difference between intervention groups of 8%. Post hoc subgroup analysis showed that males under 65 years were the group most likely to return their initial questionnaire if they received a personalised text message.</p>
---	---------------------	---------------------------	---	---	---

2	KReBs <sup>78</sup> .	B1. Reminders and prompts	<p>Two separate SWATs:</p> <p>1) Personalised text message versus a standard text message (SWAT 35)<sup>76</sup></p> <p>2) Pen incentive to enhance retention in a randomised trial (SWAT 92)<sup>79</sup></p>	<p>Primary: the proportion of 12-month questionnaires returned.</p> <p>Secondary:</p> <p>1) The proportion of questionnaires completed</p> <p>2) Time to return the questionnaire</p>	<p>Mitchell et al., 2020<sup>80</sup>: 1465 participants were included in the SWAT. In the personalised group, 644/723 (89.1%) of participants returned a questionnaire, compared to 654/742 (88.1%) in the non-personalised group. The absolute difference in return rate was 0.9% (95% CI: -2.3% to 4.2%; p=0.57). There was no evidence of a difference between the groups in the likelihood of returning a questionnaire (OR 1.09; 95% CI: 0.79 to 1.51; p=0.61), the likelihood of returning a complete questionnaire (OR 1.11; 95% CI: 0.82 to 1.51; p=0.50) nor in time to return (HR 1.05; 95% CI: 0.94 to 1.17; p=0.40).</p> <p>Mitchell et al., 2021<sup>81</sup>: 2305 participants were randomised into the SWAT. In the pen group, 1020/1145 (89.1%) of participants returned a questionnaire, compared to 982/1147 (85.6%) in the no pen group. The absolute difference in questionnaire return rate was 3.5% (95% CI: 0.8% to 6.2%; p=0.01). There were statistically significant differences in questionnaire return rate (OR 1.36; 95% CI: 1.06 to 1.74; p=0.02), questionnaire completion rate (OR 1.40; 95% CI: 1.11 to 1.78; p&lt;0.01) and time to questionnaire</p>
---	-----------------------	---------------------------	--	---	---

					return (HR 1.17; 95% CI: 1.07 to 1.27; p<0.01) favouring the pen group.
--	--	--	--	--	--

2	MiQuit-3 <sup>82</sup>	B1. Reminders and prompts	<p>2x2 factorial SWAT design:</p> <p>Personalised text message versus a standard text message (SWAT 35)<sup>76</sup></p> <p>Timing of text message prompts (SWAT 44)<sup>73</sup></p>	<p>Primary: Completion rate of questionnaire via telephone</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Completion rate via any method</li> <li>2) Time to completion</li> <li>3) Number of reminders required</li> </ol>	<p>Coleman et al., (2021)<sup>67</sup>: 194 participants were randomised into the SWAT; 50 to personalised early text, 47 to personalised late text, 50 to non-personalised early text, and 47 to non-personalised late text. There was no evidence that timing of the text message (early: one week before; or late: one day before) had an effect on any of the outcomes. There was evidence that a personalised text would result in fewer completions via telephone compared with a non-personalised text (adjusted OR 0.44, 95% CI 0.22–0.87, p=0.02). However, there was no evidence to show that personalisation or not was better for any of the secondary outcomes.</p>
---	------------------------	---------------------------	---	--	--



1	ActWELL <sup>83</sup>	B1. Reminders and prompts	Sending pre-notification cards to trial participants before outcome measurement (SWAT 76) <sup>84</sup>	<p>Primary: Number of trial participants who complete the outcome measurement (i.e., are retained).</p> <p>Secondary: Cost per participant retained.</p>	<p>Treweek et al., (2021)<sup>85</sup>: 558 participants were included in the SWAT. Of the 274 women sent a card, 231 attended the primary outcome visit (84.3%) compared to 230/284 (81.0%) for those not receiving a card. Sending a pre-notification card may result in a slight increase in attendance at a face-to-face primary outcome measurement visit at 1-year: risk difference = 3.3% (95% confidence interval = -3.0% to 9.6%). This is GRADE low certainty evidence. The direct cost of producing and sending the cards was £192 GBP (€213 EUR; \$260 USD), or £21.33 (€23.55; \$28.77 per additional retained participant).</p>
---	-----------------------	---------------------------	---	--	---

1	WORKWELL <sup>86</sup>	B1. Reminders and prompts	Pre- notification (SWAT 86) <sup>87</sup> (retention)	<p>Primary: Valid response for primary outcome (yes/no)</p> <p>Secondary:</p> <p>1) Valid response for primary outcome (yes/no) without reminder</p> <p>2) Number of reminders sent</p> <p>3) Time to response (days)</p> <p>4) Costs per participant retained</p>	<p>Sutton et al., (2022)<sup>88</sup>: Two hundred forty-four trial participants took part in the SWAT. Among those sent a pre-reminder, 100/121 (83%) provided a valid response for the WORKWELL primary outcome, compared to 97/123 (79%) of those not sent a pre-reminder. The estimated adjusted odds ratio was 1.28 (95% confidence interval 0.67–2.42), with a risk difference of 3.8% (95% CI -6.1 to 13.6%), favouring the pre-reminder. The estimated intervention cost per additional participant retained was £53.42, and the total cost per additional participant retained was £46.52.</p>
---	------------------------	---------------------------	---	--	---

2	OTIS <sup>89</sup>	B3. Non-monetary incentives	<p>2x2 factorial design: Pen incentive to enhance retention in a randomised trial (SWAT 92)<sup>79</sup></p> <p>Social incentive text cover letter sent with a postal follow-up questionnaire (SWAT 144)<sup>90</sup></p>	<p>Primary: Proportion of who returned the questionnaire.</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Time to return</li> <li>2) Completeness of the questionnaire</li> <li>3) Necessity of a reminder letter</li> <li>4) cost effectiveness</li> </ol>	<p>James et al., (2021)<sup>68</sup>: 12-month questionnaire response rate was 721 out of 755 (95.5%). Neither the pen nor social incentive cover letter had a statistically significant effect on response rate: pen 95.2% vs. no pen 95.8%, adjusted OR 0.90 (95% CI 0.45 to 1.80; p=0.77); social incentive cover letter 95.2% vs. no social incentive cover letter 95.8%, adjusted OR 0.84 (95% CI 0.42 to 1.69, p=0.63). No statistically significant differences were observed between either of the intervention groups on time to response, need for a reminder or completeness. Therefore, neither intervention was cost-effective.</p>
---	--------------------	-----------------------------	---	---	--

1	SSHeW <sup>91</sup>	B3. Non-monetary incentives	Pen incentive to enhance retention in a randomised trial (SWAT 92) <sup>79</sup>	<p>Primary: Proportion of participants who return questionnaire</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Time to response</li> <li>2) Completeness of response</li> <li>3) Reminder notice sent</li> <li>3) Cost effectiveness</li> </ol>	<p>Cunningham-Burley et al., (2020) <sup>92</sup>: 1466 SSHEW trial participants were randomised into the SWAT. In total, 13 withdrew from the host trial before they were due to be sent their follow-up questionnaire, 728 participants received a pen with their questionnaire, and 725 did not receive a pen. A questionnaire was returned from 67.7% of the pen group and 64.7% of the group who did not receive a pen. There was no significant difference in return rates between the two groups (OR 1.15, 95% CI 0.92 to 1.43, p=0.22), nor level of completeness of the questionnaires (AMD -0.01, 95% CI 0.06 to 0.05, p=0.77). There was weak evidence of a reduction in the proportion of participants requiring a reminder and in time to response in the pen group.</p>
---	---------------------	-----------------------------	--	--	---

8	<p>C-GALL <sup>93</sup></p> <p>CPIT-III <sup>94</sup></p> <p>DISC<sup>52</sup></p> <p>FUTURE <sup>95</sup></p> <p>ProFHER-2<sup>54</sup></p> <p>PUR-E-RCT<sup>96</sup></p> <p>REFLECT<sup>97</sup></p> <p>SWHSI-2<sup>65</sup></p>	<p>B4. Maintaining participant engagement</p>	<p>Sending Christmas cards to trial participants (SWAT 82) <sup>47</sup></p>	<p>Primary: Proportion of participants completing their next follow-up</p> <p>Secondary:</p> <p>1) Time to complete the follow-up (defined as number of days between follow-up due and follow-up complete)</p> <p>2) Cost per card sent (including staff time and printing and postage costs)</p> <p>3) Cost per additional participant retained (if applicable)</p>	<p>Coleman et al., (2021) <sup>49</sup>: 8 host trials were recruited. 1469 participants (age 16-94 years; 70% (n=1033) female; 96% (813/847) white ethnicity) across the eight host trials were involved in the analysis (cut short owing to covid-19). No evidence was found of a difference in retention rate between the two arms for any of the host trials when analysed separately or when the results were combined (85.3% (639/749) for cards versus 85.4% (615/720) for no card; odds ratio 0.96, 95% confidence interval 0.71 to 1.29; P=0.77).</p>
---	--	---	--	--	--

1	CLEAR <sup>56</sup>	B4. Maintaining participant engagement	<p>Combined in a three-arm SWAT:</p> <p>Thank you note or card after each study visit (SWAT 54) <sup>98</sup></p> <p>Personalisation (not including patient's name, electronically signed v personalised handwritten name and wet ink signature)</p>	<p>Primary: Proportion of participants who remain in the study.</p> <p>Secondary: Time that participants remain in the study before they withdraw.</p>	<p>Anand (2021) <sup>99</sup> (<i>interim analysis only</i>): 141 participants were randomised into the SWAT (generic thank you card, n=48; personalised thank you card, n=46; no thank you card, n=47).</p> <p>Of these, 67 patients had completed all five visits at the time of the interim analysis. This analysis only uses data for the 11 patients who voluntarily withdrew following randomisation. Proportions withdrawing: personalised thank you card= 8.7%, generic thank you card = 8.3%, No thank you card = 6.4% (Total =7.8%). Generic thank you card versus Personalised thank you card OR 0.95 (95% CI 0.22 to 4.07, P= 0.9498); Generic thank you card versus No thank you card OR 1.33 (95% CI, 0.28 to 6.31, P= 0.7168); Personalised thank you card versus No thank you card, OR 1.40 (95% CI 0.29 to 6.62, P= 0.6737); Card versus No card, OR 1.36 (95% CI- 0.34 to 5.40, P=0.6581). No significant differences were found for any of these four comparisons.</p>
---	---------------------	--	--	--	---

Table 7: Ongoing SWATs evaluating retention strategies

<b>Number of SWATs</b>	<b>Host trial acronym</b>	<b>SWAT domain</b>	<b>SWAT question</b>	<b>SWAT outcomes</b>	<b>Expected completion data and details of delay if applicable</b>
------------------------	---------------------------	--------------------	----------------------	----------------------	--

1	ARTISAN <sup>100</sup>	B4. Maintaining participant engagement	Courtesy telephone calls versus postcards to trial participants following enrolment (SWAT 121) <sup>101</sup>	<p>Primary: Proportion of participants returning questionnaire by post at the 6-months.</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Time to response to the questionnaires at all-time points by post</li> <li>2) Response rates at 6 weeks, and 3 and 12 months</li> <li>3) Response rates at 6 weeks, 3 months, 6 months and 12 months (return of questionnaire data at any point)</li> <li>4) Completeness of responses</li> <li>5) Number of reminder notices required</li> <li>6) Cost of intervention per participant</li> </ol>	<p>In progress</p> <p>Data collection ongoing, expected to be complete in September 2022.</p>
---	------------------------	---	---	--	---



1	MAGIC <sup>102</sup>	B1. Reminders and prompts	Personalised text message versus a standard text message (SWAT 35) <sup>76</sup>	<p>Primary: Questionnaire response rate at 6-months</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Time to response</li> <li>2) Proportion of participants sent a reminder follow-up questionnaire</li> <li>3) Cost.</li> </ol>	<p>In progress</p> <p>Host trial stopped recruitment due to COVID-19, recommencing early 2022. Anticipated to finish recruitment 31st July 2023.</p>
1	SWHSI-2 <sup>65</sup>	B4. Maintaining participant engagement	Thank you card following each study visit (SWAT 119) <sup>103</sup>	<p>Primary: Proportion of participants who complete the 6-month questionnaire</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Completeness of response at 6 months</li> <li>2) Whether a reminder notice is required at 6 months</li> <li>3) Cost of SWAT intervention per participant retained at 6 months</li> </ol>	<p>In progress</p> <p>The SWAT paused for 3 months due to COVID-19 and recommenced July 2020. The SWAT is expected to finish at the end of 2023.</p>

Table 8: Prematurely terminated SWATs evaluating retention strategies

Number of SWATs	Host trial acronym	SWAT domain	SWAT question	SWAT outcomes	Details of reasons the SWAT will not be completed
2	CHAMP-1 <sup>41</sup>	B1. Reminders and prompts	<p>2x2 factorial SWAT design:</p> <p>Personalised text message versus a standard text message (SWAT 35) <sup>76</sup></p> <p>Timing of text message prompts (SWAT 44) <sup>73</sup></p>	<p>Primary: Questionnaire completion rate</p> <p>Secondary:</p> <p>1) Time to response</p> <p>2) Number of attempts to contact (telephone call) required before the questionnaire is recorded as completed or the maximum number of calls</p>	<p>During the course of undertaking this SWAT, there occurred a system error with the text messaging software, resulting in a high proportion of the messages not being sent to participants as planned, so this SWAT could not proceed.</p>

1	ACTIVE <sup>45</sup>	B1. Reminders and prompts	Text message reminder which participants can respond to, compared with a 'no reply' text message on questionnaire response rates (SWAT 109) <sup>104</sup>	Primary: Proportion of questionnaires completed at the 3-month follow-up.  Secondary: 1) Time to questionnaire return 2) Proportion of patients requiring at least one return reminder notice	Due to COVID-19 the SWAT is not continuing
1	TOPaZ <sup>44</sup>	B1. Reminders and prompts	Pre-notification cards to trial participants (SWAT 76) <sup>84</sup> (retention)	Primary: Number of trial participants who complete the outcome measurement  Secondary: Cost per participant retained.	This SWAT was unable to continue due to COVID-19.

1	VITA <sup>43</sup>	B2. Monetary incentives	Conditional financial incentives vs unconditional financial incentives <sup>105</sup>	<p>Primary: The number of participants who complete the primary outcome at two-week follow up.</p> <p>Secondary: The number of reminders sent for each group.</p>	<p>Recruitment in this host trial stopped prior to reaching the target of 1900 participants on recommendation from the Data Monitoring Committee and Trial Steering Committee after a planned review of the results indicated that the research question had been answered. Therefore, the SWAT could not proceed.</p>
---	--------------------	-------------------------	---	---	--

1	L1FE <sup>42</sup>	B4. Maintaining participant engagement	Telephone calls or postcards to trial participants (SWAT 114) <sup>106</sup>	<p>Primary: Proportions of participants who complete and return the questionnaire at 6 weeks, 12 weeks, and 6-month time points.</p> <p>Secondary:</p> <ol style="list-style-type: none"> <li>1) Time to response</li> <li>2) Completeness of response at the 6-month time point</li> <li>3) Whether a reminder notice is required at the 6-month time point</li> <li>4) Cost of SWAT intervention per participant retained.</li> </ol>	The host trial was terminated early by the funder, so the SWAT was abandoned.
---	--------------------	---	---	---	---

## *Retention strategies demonstrating evidence of effectiveness*

The published SWATs reported the following evidence of effectiveness for the strategies evaluated:

### *Reminders and prompts*

For pre-notification of trial participants, SWAT 76 found that sending a pre-notification card may result in a slight increase in attendance at a face-to-face primary outcome measurement visit at 1-year: risk difference = 3.3% (95% confidence interval = -3.0% to 9.6%).<sup>85</sup> SWAT 86 compared sending a pre-notification letter or email before sending a self-report questionnaire, versus no pre-notification on retention rates (valid response for the host trial primary outcome). This SWAT found that of those sent a pre-notification, 100/121 (83%) provided a valid response for the host trial primary outcome, compared to 97/123 (79%) of those not sent a pre-reminder. The estimated adjusted odds ratio was 1.28 (95% confidence interval 0.67–2.42), with a risk difference of 3.8% (95% CI -6.1 to 13.6%), favouring the pre-notification. The estimated intervention cost per additional participant retained was £53.42, and the total cost per additional participant retained was £46.52.<sup>88</sup>

For the text messaging of trial participants, SWAT 35 found that participants randomised to receive a personalised text message were more likely to return their initial postal questionnaire than those who received a standard text message (n = 185/309; 60% vs. n = 160/309; 52%) (RR 1.16; 95% CI 1.00 to 1.33); this represents an absolute percentage difference between intervention groups of 8%. Post hoc subgroup analysis showed that males under 65 years were the group most likely to return their initial questionnaire if they received a personalised text message.<sup>77</sup> Another evaluation of SWAT 35 found that when comparing personalised text messages versus non-personalised text messages, there was evidence that a personalised text would result in fewer completions via telephone compared with a non-personalised text (adjusted OR 0.44, 95% CI 0.22–0.87, p=0.02).<sup>67</sup>

### *Non-monetary incentives*

In the evaluations of the pen strategy, SWAT 92 reported that when participants were randomised to receive a pen compared to no pen with their postal questionnaire, there were statistically significant differences in questionnaire return rate (OR 1.36; 95% CI: 1.06 to 1.74; p=0.02), questionnaire completion rate (OR 1.40; 95% CI: 1.11 to 1.78; p<0.01) and time to questionnaire return (HR 1.17; 95% CI: 1.07 to 1.27; p<0.01) favouring the pen group.<sup>81</sup> However, James et al., also testing SWAT 92, found no statistically significant effect on response rate: pen 95.2% vs. no pen 95.8%, adjusted OR 0.90 (95% CI 0.45 to 1.80; p=0.77).<sup>68</sup> SWAT 37 found no statistically significant difference in return rates when participants were sent a pen with their postal questionnaire

compared to not being sent a pen (OR 1.15, 95% CI 0.92 to 1.43,  $p=0.22$ ), nor level of completeness of the questionnaires (AMD -0.01, 95% CI 0.06 to 0.05,  $p=0.77$ ).<sup>92</sup> However, there was weak evidence of a difference, in favour of the pen group, in both time to return (median time to return 15 vs 18 days; HR 1.12, 95% CI 0.98 to 1.27,  $p=0.09$ ), and in the proportion of participants requiring a reminder (OR 0.83, 95% CI 0.68 to 1.02,  $p=0.08$ ).

In Chapter 8, we present a meta-analysis combining the results of the PROMETHEUS pen SWATs with other published SWATs of this strategy (*see Case study of individual SWATs: Pens for retention*). The pooled effect combining all SWATs indicated that including a trial pen with a questionnaire probably increases retention and response rate (pooled OR 1.21, 95% CI 1.09 to 1.35).<sup>107</sup>

#### *Retention strategies demonstrating a lack of effectiveness, or with the potential to harm retention rates*

There was no evidence of a significant positive difference in retention rates in any of the other 25 strategies tested. This includes the coordinated SWAT testing the effectiveness of sending Christmas cards on retention rates; no evidence was found of a difference in retention rate between the two arms for any of the host trials when analysed separately, nor when the results were combined.<sup>49</sup>

One SWAT identified that there was a potential harmful impact on retention from one of the strategies evaluated. SWAT 110 reported that printing the trial primary outcome on pink paper does not increase data return, when compared to printing the primary outcome on white paper (*risk ratio, 0.92 (95% CI 0.80, 1.06);  $p = .24$* ). This SWAT also showed some evidence that it potentially decreases response and is more burdensome to collect postal data by increasing the necessity for reminders, with reminders sent to 48.9% (43/88 participants) in the pink paper group and in 30.7% (27/88 participants) in the white paper group (*risk ratio 1.59 (95% CI 1.09, 2.33);  $p = .01$* ).<sup>71</sup>

## Chapter 4: Discussion of the PROMETHEUS programme

### *Summary of main findings from PROMETHEUS*

The PROMETHEUS programme successfully embedded 42 SWATs within 31 host RCTs, exceeding the original target of 25 SWATs. This is the biggest single effort to generate SWAT evidence in the world, representing a substantial increase in the global methodological evidence base. The PROMETHEUS-funded SWATs represent an increase of 18% (12/68) more SWATs to the Cochrane systematic review of recruitment strategies,<sup>23</sup> and 79% (30/38) more SWATs to the Cochrane systematic review of retention strategies.<sup>24</sup>

The main findings from the PROMETHEUS SWATs reported to date show that for recruitment, there was no evidence of a significant difference in recruitment rates in any of the strategies tested. For retention, we found that pre-notification of trial participants by sending a card may result in a slight increase in attendance at a face-to-face primary outcome measurement visit; and that sending a pre-notification letter or email before sending a self-report questionnaire by post increases response rates, compared with no pre-notification. We also found that participants randomised to receive a personalised text message were more likely to return their initial postal questionnaire than those who received a non-personalised text message; and that men aged under 65 years were most likely to return their initial questionnaire if they received a personalised text message. Another SWAT comparing personalised text messages versus non-personalised text messages, found evidence that a personalised text would result in fewer telephone follow-ups, which were more resource intensive. When participants were randomised to receive a pen compared to no pen with their postal questionnaire, combined results from PROMETHEUS together with other published SWATs found that including a pen probably increases retention and response rate (pooled OR 1.21, 95% CI 1.09 to 1.35). One SWAT showed evidence of an adverse impact on retention when printing the trial primary outcome on pink paper compared to printing the primary outcome on white paper; this showed a decreased response, and it was also more burdensome to collect postal data as it increased the need for reminders. The other strategies tested showed no evidence of a significant difference in retention rates.

This was possible to fund more SWATs than initially anticipated due to the majority of the SWATs costing less than the proposed funding limit, with the mean SWAT cost being £3,535; less than the proposed £5000. We have shown that typically SWATs can cost less than the £10,000 SWAT funding made available by the HTA. Therefore, PROMETHEUS is currently the largest programme of work to act as a central coordination point to offer both funding and practical support for the embedding of SWATs, independently contributing the largest amount of evidence to the recruitment and retention



strategy evidence base. These SWATs were collectively implemented within a large number of host trials across multiple CTUs, demonstrating the wide reach of the programme. The programme's success confirms the feasibility of implementing methodological research within a vast range of research areas when appropriate resource and infrastructure support is made available. It also confirms the feasibility and acceptability among trial teams of conducting coordinated SWATs, enabling a more rapid evaluation of recruitment and retention strategies.

As host trial teams were often inexperienced in conducting and implementing SWATs, the PROMETHEUS programme acted as an invaluable coordination point, providing teams with the confidence, knowledge, and resources to do so. Further to this, the programme identified that the lack of SWAT funding was often a barrier to implementation; a concept which was reinforced as the number of conducted SWATs increased following the introduction of PROMETHEUS funding. However, despite providing both funding and guidance, the encouragement of external researchers to conduct a SWAT proved difficult; with the majority of the SWATs being conducted by both York Trials Unit (18 SWATs, 43%) and the CTUs which PROMETHEUS co-applicants were associated with (16 SWATs, 38%). It is important to place this seeming lack of poor external engagement in context, given that the original PROMETHEUS focus was to support our collaborating network of CTUs and centres, who had already committed to starting at least two SWATs.

### *Strengths*

Five of the host trials implemented more than one SWAT, and a further four implemented a factorial design SWAT, allowing for the assessment of two SWATs in a coordinated way. This finding demonstrates that, within an individual host trial, there is often a capacity to address more than one SWAT question, either separately (such as testing a strategy aimed at recruitment during the recruitment phase; and then testing a retention strategy during the follow-up phase) or simultaneously, using a factorial design. This suggests that there is capacity to significantly speed up and strengthen the evidence base through teams undertaking more than one SWAT in their trials where relevant. Another pathway to speeding up and strengthening the evidence base is through the coordinated SWAT approach that we pioneered, where the same recruitment or retention strategy is tested across multiple host trials simultaneously. Thus, our work highlights that given finite resources, robust evidence can be generated at speed in two key ways: by doing more SWATs in the same host trial, or by testing the same SWAT question across different host trials in a coordinated way.

Aside from the provision of SWAT funding, the work undertaken within PROMETHEUS also presents added value through working with CTUs, which are the key centres in the UK supporting the delivery

of trials, to support the uptake of SWATs. Uniquely, PROMETHEUS focused on generating evidence for trial recruitment as well as retention, both of which are crucial for the successful delivery of trials, but which until PROMETHEUS had not been addressed together. Despite similar funding as other programmes such as MRC START (which funded 12 SWATs) and TRECA (which funded six), PROMETHEUS has been able to generate more than three and seven times the number of SWATs, respectively, adding substantial, much-needed evidence to support trial process decision-making.

### *Limitations*

Our experience of five funded SWATs failing to complete highlights the reality of SWATs being entirely subject to the fortunes of the host trials; if the host trial stops for any reason or changes its mode of follow-up, it places the SWAT in jeopardy. One study commenced the SWAT but technical errors in the software associated with the intervention (text messaging) resulted in a significant proportion of participants not receiving the intervention and so the SWAT had to be terminated. For two other SWATs recruitment to the host trial was stopped prematurely (one on direction of the associated oversight committees; one by the Funder) and so planned retention SWATs could not proceed. Finally, although we had awarded funding to all of our host trials prior to the COVID-19 pandemic proliferating in the UK in March 2020, the widespread societal changes that this forced resulted in the recruitment pathway of many of the PROMETHEUS host trials either pausing or altering their study pathways. As a result, two SWATs did not proceed, and 10 others remain in progress at the time of writing this report. We will work with the host trial teams of the ongoing SWATs to ensure their findings appropriately contribute to the evidence base.

Further to this, the funded SWATs evaluated a wide range of recruitment and retention strategies. While this led to an overall increase in the evidence base, the evidence generated did not allow for enough replications to have been conducted to conclude the effectiveness of all SWAT strategies evaluated via meta-analyses. This limited our ability to answer most SWAT questions definitively. Despite this, significant progress has been made with some SWATs (e.g., pens for retention) for which only a limited number of further replications in specific populations (male populations, younger adult populations) are now required to reach a definitive evaluation of strategy effectiveness. The development of the Trial Forge SWAT Network should enable additional replications to be completed more rapidly to fully answer these SWAT questions.

We found a greater interest and conduct of SWATs evaluating retention, rather than recruitment strategies. This may be due to retention SWATs being potentially less challenging to undertake than recruitment SWATs, as logistically, there is likely to be more time to introduce a retention strategy when there are multiple follow-up time points. Recruitment SWATs may also present additional

challenges, such as the inaccurate perception that multiple levels of consent are required from individuals being approached to enter a trial (e.g., consent to be invited into the trial, consent to be included in the SWAT, and consent to be enrolled into the trial). Alternatively, there may also be greater time pressure associated with the embedding of a recruitment rather than retention SWAT, due to the additional tasks of site set up. Lessons learnt from the PROMETHEUS programme are also discussed by Clark *et al.*, (2022).<sup>108</sup>

As all the funded SWATs were also conducted within the UK, the SWAT evidence generated through PROMETHEUS is not necessarily applicable to populations within other countries, limiting the reach of the programme.

We included only randomised host trials, which may have limited our ability to identify variation in effectiveness of recruitment and retention strategies across different research contexts. However, we believe that focusing on randomised controlled trials in this particular programme was appropriate, since trials are particularly difficult to recruit to, with randomisation and strong treatment preferences leading to issues around both clinician and participant equipoise, something not experienced in non-randomised studies. A direction for future research is to include non-randomised studies as host studies. We also focused only on randomised SWATs. Non-randomised SWATs such as qualitative studies, observational studies, and surveys have a substantial role to play in improving the evidence base for recruitment and retention. One strong way that non-randomised SWATs can help with the evidence base is through process evaluations, which might include interviews to elicit participants' experiences of being exposed to a recruitment strategy, as well as the experiences of patients who decline the invitation to participate in a trial. This will be crucial to include in future SWATs to ensure both effectiveness in terms of recruitment and retention but also acceptability for participants.

#### *Comparison with existing programmes of SWAT research: MRC START and TRECA*

Building on and improving on the SWAT work initiated by MRC START, which established the feasibility of testing recruitment strategies across multiple host trials,<sup>25, 26</sup> as well as TRECA,<sup>30</sup> which aimed to undertake a programme of SWATs in trials recruiting children and young people, PROMETHEUS has successfully undertaken more SWATs at a faster pace and successfully disseminated findings. However, PROMETHEUS faced challenges that were similar to those in MRC START, such as a large proportion of the SWATs being undertaken by researchers linked with the PROMETHEUS team. There is a range of reasons for the success of PROMETHEUS, which includes prior learning on SWATs gained from MRC START and other work undertaken by its collaborators,

the financial support to undertake SWATs, as well as the support mechanism provided to host trial teams from a well-established registered CTU. This suggests that speed and efficiency for undertaking SWATs can be improved further, but it is important that the momentum and skill set of staff are maintained.

### *Patient and public involvement (PPI)*

PPI was an important part of this programme, both for the overarching PROMETHEUS programme, and for the SWATs undertaken. We actively involved PPI members in identifying, prioritising and implementing the SWATs, and we have dedicated chapter five to outline the PPI undertaken, as well as lessons learnt for the trials community, and PPI partners.

### *Equality, Diversity and Inclusion*

Increasing the diversity of trial participants is fundamental to increase participation rates in trials. An ambition in testing recruitment and retention strategies is not only to increase the numbers of people taking part in trials, but also helping to recruit participants that better reflect those that might benefit from the trial results, or to avoid harm. Our statistical analysis plan reflected this and for each host trial funded, we asked that wherever possible, demographic data on the age, gender, and ethnicity of participants should be recorded. The importance of capturing these data can be illustrated in the GRASP SWAT, which tested personalised vs., non-personalised text messages.<sup>77</sup> In this SWAT, post hoc subgroup analysis showed that males under 65 years were the group most likely to return their initial questionnaire if they received a personalised text message. Capturing such data in all SWATs will also allow subsequent future patient-level meta-analysis to be undertaken, and differences in effect between different populations to be assessed. In time, there may also be a need for testing of population-specific SWATs if there is evidence of differential effects in distinct population groups.

### *Conclusions*

PROMETHEUS originally aimed to fund 25 SWATs and supported 42 SWATs, generating the single biggest body of SWAT activity in the world. When SWAT funding was made available, we found that many teams embedded SWATs into their research. The mean cost of each SWAT was £3,535. In addition to funding SWATs, PROMETHEUS successfully demonstrated the methodological feasibility of undertaking coordinated SWATs, a powerful new tool with the potential to rapidly accelerate the evidence base for recruiting and retaining trial participants. In addition to money, the coordination PROMETHEUS provided was crucial to increasing the recruitment and retention evidence base.

The PROMETHEUS SWATs reported to date found that there was no evidence of a significant difference in recruitment rates in any of the strategies tested. For the retention strategies, we found evidence of effectiveness for sending pre-notification cards, letters, and emails to trial participants; that personalised text messages were more effective than non-personalised text messages, especially for men aged under 65 years; that sending a pen compared to no pen with postal questionnaires was more effective, although this result was not consistent as another SWAT found no difference. One SWAT comparing pink versus white paper for printing the primary outcome showed evidence of a decreased response in the pink paper group, and that it was also more burdensome to collect postal data in this group. We found no evidence of a significant difference in retention rates for any of the other strategies tested. More replications of the recruitment and retention strategies funded by the PROMETHEUS programme are required to provide definitive evidence for these strategies.

In the chapters to follow, we reflect on lessons learnt from undertaking the PROMETHEUS programme.

## Chapter 5: Lessons learnt for PPI

### *Chapter overview*

PPI is important for developing and evaluating interventions aimed at recruiting and retaining patients in RCTs. In this section, we report the PPI in PROMETHEUS using the GRIPP2 Reporting Checklist<sup>109</sup> and reflect on the lessons learnt for involving PPI partners in SWAT research.

### *Defining PPI in PROMETHEUS*

The traditional aim of PPI is to involve end users in the research process. Given the methodological focus of PROMETHEUS, we defined 'patients and the public' broadly, to include those who would potentially be impacted by the recruitment and retention strategies, such as patients, potential patients, carers, and members of the public.

We acknowledge the importance of stakeholders from the trial community who may be involved in undertaking SWATs, recruiting participants, or applying the evidence from SWATs, such as principal investigators, statisticians, trial managers, and staff based within CTUs and other centres undertaking rigorous RCTs. However, we consider the contributions of these stakeholders as separate from PPI.

### *Aims*

The aims of the PPI in PROMETHEUS were to:

1. Inform the overarching prioritisation of the recruitment and retention strategies that host trial teams were funded to undertake
2. Inform how individual host trial teams could select, adopt and/or adapt specific recruitment or retention strategies for their specific patient population and recruitment/retention contexts.

### *PPI: Methods*

#### *PPI input prior to funding being awarded*

A significant amount of PPI underpinned the PROMETHEUS programme. MRC START, which PROMETHEUS builds on, included significant public involvement in developing recruitment interventions and the methodological frameworks around recruitment SWATs.<sup>25</sup> Five members of the PROMETHEUS team (authors Devane, Torgerson, Gillies, Galvin and Treweek) undertook two James Lind Alliance priority setting exercises: one for recruitment<sup>110</sup> and the other for retention strategies<sup>111</sup> with stakeholders including members of the public, clinicians, and researchers; these priority lists were used to inform the prioritisation of the recruitment and retention strategies used

in PROMETHEUS. A number of the existing recruitment and retention strategies registered on the SWAT Repository involved patients and the public in their development;<sup>112, 113</sup> or were led by the host trial's PPI group.<sup>114, 115</sup>

#### *PPI involvement in prioritisation of strategies to be funded*

We hosted a SWAT public involvement panel to ensure the project met the needs of and gained input from patients and the public. We advertised for members through PPI forums and our existing contacts. The panel comprised five men and women with a range of health conditions, and diverse experiences of being involved as PPI members in specific trials, prior involvement in methodological research (such as MRC START) as well as part of research funding panels. Panel members were provided with a 'Lay Summary' of the PROMETHEUS programme, along with a 'Remit and Role Description', which outlined the role and provided practical information such as how expenses and payments would be made, and who to contact for further information. We organised a 'Priority setting meeting', ahead of which we circulated an agenda and a briefing document providing an initial draft outline of a priority list of recruitment and retention strategies that we wanted to fund as part of PROMETHEUS, and the methods that we had used to identify this initial priority list (see Chapter 2). We also specifically asked members of the PPI panel to think of any strategies that they additionally wanted to prioritise to be tested. During this meeting, PPI members commented on the initial priority list and each member ranked the strategies, listing their preferred strategy first, and their next preferred option second. Suggestions made by PPI panel members for additional SWATs not on the initial priority list included thanking participants for taking part in trials, which could include offering 'thank you' gifts, and including a person's name in trial correspondence. We provided a 'PPI Feedback form' for panel members to provide feedback on the meeting and how they felt about their ability to contribute.

#### *PPI input into individual SWATs*

To ensure ongoing PPI, we requested that each host trial consulted with patient and public members involved in their RCTs as part of the implementation process for each SWAT. This gained bespoke advice relevant to the context and population of each host trial. Additionally, where individual host trial teams undertaking funded SWATs lacked access to PPI members, we sought PPI input from the PROMETHEUS PPI panel on their behalf.

#### *PPI: Results*

##### *PPI involvement in prioritisation of strategies to be funded*

We developed a final priority list which the PPI group agreed with (see Chapter 2). These strategies were listed and presented in the order of priority selected by the PPI panel. The PROMETHEUS

priority list was presented on our programme website, included in the information for funding applicants, and highlighted in the funding peer reviewer form to ensure that reviewers prioritised funding applications that were proposed to evaluate the SWATs. Overall, the feedback from PPI panel members was very positive, with members saying that they were encouraged to and felt able to contribute to the meeting discussion, that their ideas were listened to by other members of the group, and that their views were valued.

#### *PPI input into individual SWATs*

PROMETHEUS PPI panel members provided additional input into the implementation of two funded SWATs. They reviewed participant information material in a SWAT which assessed the effectiveness of a brief participant information leaflet (PIL) versus standard length PIL on participant recruitment rates; and a theoretically informed cover letter which aimed to improve response rates to follow-up postal questionnaires. For the Christmas card SWAT, we consulted a PPI group consisting of six members from the Health Services Research Unit PPI Partnership at CHaRT (University of Aberdeen) on the design and content of the card. The group was asked to provide input and comments on the following: the design of the card (two designs were presented); the wording inside the card; and to give feedback on any other aspect. Most of the group members agreed on the card that was ultimately used in all the host trials (*Please refer to Appendix 2 to view the card*) and expressed that the card was acceptable. Other host trials consulted their own PPI groups on the specific SWATs funded as part of PROMETHEUS; however, we did not specifically ask for feedback, so we are unable to report the results of these PPI activities.

#### *PPI: Discussion and conclusions*

There was significant PPI in PROMETHEUS from its earliest stages to prioritisation of the strategies to be funded, to implementation of the funded SWATs. Involving patients and members of the public in methodological research can be challenging, as methodological research is often several steps removed from direct patient care.<sup>116</sup> However, we were able to successfully incorporate PPI in a way that members felt able to contribute to and found rewarding.

#### *PPI: Reflections and critical perspective*

Because of the methodological focus of PROMETHEUS, our PPI panel members were more 'experienced', and had had prior involvement in trials, and some also had prior involvement in methodological research. Whilst this allowed our group to quickly understand and appreciate the aims, objectives, and methods involved in PROMETHEUS, and to contribute in a way that was meaningful and rewarding, this also meant that we may have excluded more 'naïve' PPI members who could potentially have brought novel perspectives to the process. However, on the whole, we



felt that having panel members who had prior experience was more likely to have benefitted rather than hindered our programme.

*Lessons learnt for involving PPI partners in SWAT research.*

- Trial teams should include PPI when planning SWATs: In the same way that PPI is expected for the main trial, PPI should be an important consideration when planning the SWAT and should include those who would potentially be impacted by the recruitment or retention strategy being evaluated
- Depending on the nature of the SWAT, PPI members can include potential or enrolled trial participants
- Helping to identify novel or adapt existing strategies: PPI partners can help trial teams to identify novel or untested strategies to be evaluated. In PROMETHEUS, we asked PPI members to consider potential strategies that they would be interested in getting tested; a strategy suggested in response to this, which was to thank trial participants, led to the development of SWAT 119, which is testing the effectiveness of a thank you card following each study visit. Where a strategy has been tested in a different population, format, or context, PPI partners can be particularly valuable in assisting trial teams to adapt that strategy to be tested within their particular trial
- Accessing PPI partners: For SWAT strategies targeting participants or potential participants, the first port of call should be to approach the PPI members that informed the host trial. For strategies targeting staff, such as a recruitment training course for site staff, the key activity would be to approach staff undertaking the recruitment at sites.

## Chapter 6: Lessons learnt: SWAT development and funding

### *Chapter overview*

In this chapter, we outline how SWAT strategies should be prioritised or selected, as well as key elements to consider when designing a SWAT.

### *Selecting which strategy to test in a SWAT: the importance of prioritisation*

When a decision has been made to include a SWAT within a host trial, the next step is to determine the most suitable SWAT to undertake. To do this the PROMETHEUS group found that there were several areas where research teams wanted support, information, and clarity:

- To have clear SWAT research priorities
- Identifying an intervention suited to a host trial population and trial type
- Communication as to when no further SWAT replications are necessary
- Transparent costs of SWATs
- SWAT interventions for low budgets

This would ensure efficient resource use and maximum research output.

The PROMETHEUS priorities were based on SWATs that could be undertaken with funding of £5000 or less. For further information on how the PROMETHEUS priorities were identified, please refer to Chapter 2 ('PROMETHEUS Preparatory work').

While the PROMETHEUS programme identified a list of priorities for this work (*see Chapter 2*), there remains a need for continually updated research priorities to allow researchers to address the questions relevant at that time. Collaboration is essential when undertaking SWATs. The priority list needs to be developed and finalised in collaboration with working groups that are involved in SWAT and methodology work such as Trial Forge<sup>117</sup> and those involved in the Cochrane reviews.<sup>23, 24</sup> PROMETHEUS has already proactively collaborated with Trial Forge and the MRC NIHR TMRP to begin this work.<sup>118</sup> Together with Trial Forge we have set up the Trial Forge SWAT Network<sup>119</sup> and have utilised membership of the TMRP Recruitment and Retention working groups to promote priorities for SWAT work. In addition to a priority list, to ensure research output is maximised, establishing a real-time method to communicate SWAT priorities to researchers is a recommended future development. The Cochrane reviews on recruitment and retention strategies<sup>23, 24</sup> are updated every few years and so establishing a more rapid method would enable the research gaps to be identified and addressed. Additionally, we identified that there is a need to work with research teams to help establish SWAT priorities based on their experiences as to what is important with their

research areas, settings, and populations. Currently, some of the research priorities that have been identified by both the PRioRITY<sup>110, 111</sup> and PROMETHEUS groups are not suitable for all trial teams and therefore teams have expressed a reluctance to embed them. One such example is the use of SMS for participant retention. Many teams already use SMS and so do not want to undertake a SWAT testing SMS vs no SMS, which the Cochrane review<sup>24</sup> states currently requires further evaluations. This further highlights the importance of working with teams undertaking methodological research to identify SWATs that are acceptable, thus increasing the likelihood of SWATs being undertaken and the findings implemented. There is also a changing landscape that trialists need to be mindful of, for example, SWATs relating to postal follow-up of participants are becoming less relevant with moves toward electronic data collection.

To ensure that SWATs are undertaken in areas of need, it is essential that there is a priority list that demonstrates not only which SWATs are needed but in which trial populations. This will prevent unnecessary SWAT replications from being undertaken and ensure resources are directed to areas that are needed to increase the evidence base. For example, there is high certainty evidence that pens are an effective strategy for older females as a retention strategy but only moderate certainty evidence in a younger population and males.<sup>119</sup> Therefore, it is recommended that if additional replications are undertaken, they are done so within these populations.

There is also a need for more awareness around embedding a SWAT which 'de-adopts' a strategy that has already been planned and budgeted for within a host trial, but further evidence is needed for evidence of its effectiveness, for example, newsletters. These are often sent to sites and participants so these could be tested as part of a low-cost SWAT. Additionally, in-person site initiation visits could be compared to remote site initiation visits and if site initiation visits are 'standard' practice then such a SWAT would save money.<sup>120</sup>

### *SWATs prioritisation, GRADE and pragmatism*

When considering SWAT priorities there is a need for pragmatism; the question needs to be asked as to whether additional replications are actually necessary. It is not possible to complete a SWAT within every population, each replication requires resources, and these resources could be directed to an area where there is still large uncertainty around a strategy.

When assessing the SWAT evidence base GRADE is used to rate the quality of evidence and grade the strength of recommendations.<sup>121</sup> The GRADE domains for rating a judgement down are based on five different criteria: risk of bias, imprecision, inconsistency, indirectness and publication bias, and result in a classification of either very low, low, moderate, or high. Currently, the PROMETHEUS team has collaborated with Trial Forge and assessed SWAT evidence to create Trial Forge evidence

packs<sup>119</sup> that break down recommendations by population according to the GRADE assessments (see the ‘support teams to undertake SWATs: Trial Forge Evidence packs’ section and Figure 2). There is, however, a trade-off between high certainty and directing resources to maximise output and so it is suggested that we may have to accept that we do not have ‘high certainty’ for all populations and a pragmatic consensus will need to be made as to whether further SWAT replications are recommended.

*Supporting teams to undertake SWATs: Trial Forge Evidence packs and an information repository*

Trial Forge<sup>107</sup> have developed a number of Trial Forge evidence packs.<sup>119</sup> These packs provide trialists with information on how many SWAT replications have been undertaken and the evidence base to date, including a summary of GRADE. Figure 2 is an example summary found in a Trial Forge evidence pack. Additionally, they provide teams with practical implementation advice on how to undertake and embed the specific SWAT. These packs enable research teams to make an informed decision - feedback from researchers that have used them is that they are a useful additional resource to support teams.

<b>EVIDENCE SUMMARY</b>	
<b>What is it?</b>	<b>A pen</b> sent with trial questionnaire and/or study materials, which serves as a reminder to return study items. The pen can contain the research institution details and/or the trial name and logo
<b>Does it work?</b>	Sending a trial pen <b>probably increases retention and response rate</b>
<b>How big is the effect?</b>	An increase of <b>1.9%</b> (95% confidence interval = 0.0% to 3.7%).
<b>How certain are we?</b>	GRADE <b>Moderate</b> certainty evidence. GRADE <b>High</b> certainty evidence for an older population.
<b>Recommendation</b>	We <b>recommend that trialists send trial pens to participants</b> to increase retention in trials that use questionnaires.
<b>How can I use this straight away?</b>	See <b>Resource bundle</b> below for support materials.

Figure 2: Summary example in Trial Forge evidence packs, used with kind permission from Trial Forge {Trial Forge, 2021}<sup>119</sup>

To further aid teams in selecting a SWAT, a repository of information could be developed to share necessary information on a freely accessible public platform. This would include not just protocols

which are currently found on the Northern Ireland SWAT store,<sup>32</sup> but additional helpful documents to aid trial teams such as those detailed in Table 9. Currently, steps are being made to progress a repository in collaboration with the Trial Forge SWAT Network.<sup>122</sup> This development will likely take several months to ensure that it is detailed and piloted, and a way to operationalise it presently and in the future is identified. Both the PROMETHEUS and Trial Forge initiatives will lead this and report on the progress via their websites and social media channels.

Table 9: Information to include when registering a SWAT in the SWAT Repository

- |   |
|---|
| <ul style="list-style-type: none"><li>- Details of the SWAT intervention</li><li>- Replications that have been undertaken and in what populations/trial types</li><li>- Replications in progress (including the population/trial types)</li><li>- Evidence to date (meta-analysis)</li><li>- Populations where evidence is needed</li><li>- SWAT protocol</li><li>- Additional supportive information to embed the SWAT</li></ul> |
|---|

#### *Design Choices: cluster randomised vs individual randomised*

In planning a SWAT, an important consideration is whether to randomise at the individual-level (e.g., the participant or potential participant) or at the cluster level (e.g., recruiters or hospitals). If the recruitment or retention strategy is focusing on participants, potential participants, or their proxies, an individual-level randomised design may be the most efficient and appropriate randomisation method. However, for logistical reasons cluster randomisation, perhaps by study site, might be more feasible. On the other hand, if the recruitment or retention strategy affects individual participants or potential participants, but is mainly happening at a different level, such as whether training trial recruiters improve recruitment rates then a cluster design may be more appropriate (*see Chapter 8 for case study*). Cluster randomisation is often used to minimise ‘contamination’ between intervention and control participants.<sup>120</sup> For example, in a SWAT where the intervention is to train Research Nurses recruiting participants on how to improve trial recruitment, it would be difficult to undertake individual patient randomisation within hospital sites (i.e., intervention patients are recruited using the training method; control patients recruited using standard practice) as Nurses who receive the recruitment training might use some of the training received to recruit control patients. Nurses within the same hospital site might also share the recruitment training information (advertently or inadvertently) with colleagues recruiting control patients, who may also change their usual recruitment practice, leading to contamination. This contamination might result in dilution bias

and consequently lead to Type II error (i.e., erroneously concluding there is no effect when the intervention is effective), which cluster randomisation would guard against.

The design of the host trial, as well as the recruitment methods, clinical context, and patient population of the host trial can influence the decision on whether to use individual or cluster randomisation. If the host trial is a cluster randomised trial, the SWAT could use the same clusters as the host trial (although the trial clusters would be randomised separately for the SWAT stratified by the host trial’s cluster status so that half the intervention and half the control clusters would receive the SWAT).

For many SWATs, the decision about whether to randomise at the individual or cluster level is also influenced by the practicalities and ease of implementing the SWAT, which often focuses on minimising the burden on the staff delivering the SWAT. For instance, a retention trial of different letters, or incentives, such as pens, may use a cluster design for practical reasons to avoid centres having to individually randomise follow-up letters to different conditions. The MRC-funded START programme, which tested two recruitment strategies across 12 host trials anticipated that some host trial teams would prefer to use cluster allocation of the START recruitment interventions, to ease the logistical burden on the trial, as well as to reduce the risk of possible contamination.<sup>25</sup> This is also the approach we took in PROMETHEUS, so where individual randomisation was likely to be a challenge or burdensome to those undertaking the SWAT, we suggested to applicants that they consider cluster randomisation. Table 10 outlines examples of individual and cluster randomised designs in SWATs.

Table 10: Examples of individual and cluster randomised designs in SWATs

<b>Types of individual randomisation</b>	<b>Reference</b>
Potential participants are randomised: such as inviting them into a trial, either face-to-face or through written invitation.	See Parker et al., 2018 <sup>49, 123</sup>
Existing trial participants are randomised.	See Chapter 8 Christmas card SWAT, and Coleman <i>et al.</i> , (2021) <sup>49</sup>
Factorial randomisation: Such as a 2x2 factorial SWAT where participants due their follow-up questionnaire are randomised to be sent: a pen; a social incentive text cover letter; both; or neither.	See James <i>et al.</i> , (2021) <sup>68</sup>

Zelen design: Only those randomised to the intervention group are asked for consent.	See Fowell et al., 2006 <sup>124</sup>
<b>Types of cluster randomisation</b>	<b>Reference</b>
Recruiters are randomised: such as individual GPs, surgeons, research nurses or lay advocates.	See Larkey et al., 2002 <sup>122</sup>
Recruiting sites are randomised: such as general practices, hospitals or community based mental health teams.	See Chapter 8 Training SWAT, and Parker et al., 2022 <sup>51</sup>
Time periods are randomised: such as a week or month of recruitment.	See Sheridan et al., 2020 <sup>126</sup>
Matched pairs (typically recruiting sites) are randomised together: such as similar sized hospitals or mental health teams. This can use the same cluster pairs as the host trial.	See Hughes-Morley et al., 2016 <sup>127</sup>

For statistical considerations of SWATs, please see Chapter 9.

### *Funding for SWATs*

There are limited funding streams available for methodological work both in the UK and internationally. PROMETHEUS, funded by the MRC) UK and NIHR as part of the MRC–NIHR Methodology Research programme was able to offer teams funding which can be a limiting factor to researchers undertaking a SWAT.<sup>36</sup> In Chapter 3 we outline the average cost of a SWAT funded through PROMETHEUS. Prior to PROMETHEUS, the UK MRC funded the MRC START programme, which tested the feasibility of undertaking SWATs in a coordinated way.<sup>25</sup>

Some funding agencies have engaged with the need for SWATs to improve trial conduct. These include the Irish Health Research Board which has a programme of funding SWATs up to €25,000 and the UK's NIHR HTA programme that offers UK research teams up to £10,000 to embed a SWAT<sup>4</sup> and up to £30,000 to embed Studies Within A Project (SWAPs) across a number of their funding streams for multiple long-term conditions.<sup>5</sup> Our experience with PROMETHEUS suggests that this level of funding should be suitable for many different SWAT interventions to be embedded within host trials. However, within the PROMETHEUS programme lower cost and easier SWATs were deliberately chosen by the PROMETHEUS Trial Management Group as priorities. Additionally, the NIHR through its Health Services and Delivery Research (HS&DR) and Fellowship streams has also funded SWATs.<sup>30, 127</sup>

### *Lessons learnt for trial teams and methodologists undertaking SWAT research*

- When setting SWAT research priorities, methodologists should provide as much information as possible so that teams are able to make informed decisions. They should consider developing strategies to aid teams to identify and select suitable SWAT interventions for their individual host trials. SWAT priorities need to be clearly communicated to trial teams and funders.
- We would recommend that the priorities are presented alongside estimated costs, resource use (including the time needed to undertake the SWAT) and if possible, a protocol. This was a strategy that we undertook during the PROMETHEUS programme, and feedback received from trial teams was that this was helpful.
- The allocation units of the host trial that can be randomised for the SWAT consist not only trial participants but also sites, recruiting staff, and time periods. The decision on the best allocation unit will be informed by a combination of what is most appropriate methodologically, and what is practical and acceptable to the host trial team, PPI members of the host trial team and staff undertaking recruitment activities.

### *Lessons learnt for funders*

- Trial teams want to know that they were undertaking a SWAT that is necessary and relevant to increase the evidence base. This will ensure efficient resource use and maximum research output. More funding is needed to support the development of these types of resources.
- When applying for funding, trial teams need to indicate whether the question they are addressing is a priority SWAT question, as well as provide a clear rationale for selecting that particular question.



## Chapter 7: Lessons learnt for governance approvals for SWATs

### *Chapter overview*

In this chapter, we discuss our experience of obtaining governance approvals for SWATs, including approvals needed from the host trial Sponsor, trial oversight committees, Research Ethics Committee (REC) and Health Research Authority (HRA) approvals. We outline Key Learning for Sponsors, oversight committees, Trial Management and for journals and peer reviewers. Additionally, we consider the operational ethical issues in relation to embedding SWATs at scale, rather than an empirical analysis of the conceptual ethical issues.

### *Governance approvals*

SWATs are designed to be embedded within a host trial with no implication on its integrity, rationale, or outcomes<sup>15</sup> and, as a result, the obtaining of governance approvals should, in theory, be relatively simple. Despite this, many difficulties including those similar to obtaining approvals for large multi-centre studies continue to prevail.<sup>128-130</sup> These include, but are not limited to, the need for patient consent and/or information regarding the SWAT, study power, and a mis- or lack of understanding with regards to the methodology such as the need for replication. These difficulties pertain at every level of study approval from oversight committees and patient advisory groups, through to Sponsors and governance bodies.

Throughout the PROMETHEUS programme, a range of queries around the approvals process for SWATs were raised. This chapter outlines the queries noted at each level of approval, details some potential ongoing developments in this regard, and outlines suggestions for each group involved in the approvals process for a SWAT.

### *Sponsor Approval*

In the case of individual SWATs, sponsorship usually rests with the host trial Sponsor. Many of the PROMETHEUS SWATs were conducted through York Trials Unit and therefore were sponsored by the University of York; a Sponsor who has for a long time supported SWATs and other methodological work and so has an understanding of the nuances associated with these. As a result, obtaining Sponsor approval of individual SWATs was generally straightforward. SWAT naïve Sponsors however often raised queries with regards to the methodology or processes for adding the SWAT to the host trial approval. For example, Sponsors have asked for the SWAT to be added to the main study protocol thus resulting in a substantial amendment and the addition of additional information on the

SWAT to participant-facing materials which has potential to dilute the SWAT (See Ethics/HRA approvals).

On the converse, the fact that SWATs are usually methodological, rather than clinical, posed difficulties for another sponsorship arrangement. The host trial was sponsored by an NHS Clinical Commissioning Group (CCG), with the SWAT being undertaken by study team members based at the local university. The CCG was keen that the university assume sponsorship for the SWAT, which posed difficulties for amendment submission given two sponsors cannot be detailed in a single application. In the end, the CCG agreed to be Sponsor.

For coordinated SWATs, sponsorship arrangements can be more complicated given the number of host trials involved and that many, if not all, may have different Sponsors. For the two coordinated SWATs undertaken within the PROMETHEUS programme (*see Chapter 8 for case studies*), sponsorship was, ultimately, held by the institution associated with the SWAT coordination (in both cases the University of York). In the instance of the Christmas Card SWAT (*see Chapter 8 for case studies*), the original proposed Sponsor did not understand SWAT methodology and so was unwilling to approve this. The SWAT was only delivered successfully when sponsorship was undertaken by a Sponsor with an understanding of SWATs. As a result, there is undoubtedly potential for difficulties with obtaining Sponsor agreement and approval for coordinated SWATs. It may be difficult to get agreement for sponsorship where the proposed SWAT sponsor is an academic organisation with no existing association with the host trials, or where multiple Sponsors are associated with the host trials but they are unwilling to take sole oversight of the methodological component.

#### *Oversight Committee Approvals (TMG/TSC/DMC)*

Throughout the PROMETHEUS programme, there was little difficulty in obtaining support for SWATs from host trial oversight committees. Trial Management Groups (TMGs) were generally happy to support the inclusion of a SWAT if the Chief Investigator and Trial Manager proposed this. Furthermore, where the trial management team was supportive of undertaking a SWAT then the Trial Steering and/or Data Monitoring Committees (TSC, DMC) were happy to support their inclusion in the host trial. As part of routine study oversight, it is suggested that the TSC particularly should have oversight of SWAT progress given their remit to endorse TMG actions<sup>131</sup> and provide overall study supervision.<sup>132</sup> Given the general low-risk nature of SWATs oversight from the DMC may not be warranted, although this is dependent on the nature of the host trial and SWAT, and DMC agreement.

### *Ethics/HRA Approval*

Many research teams have reported barriers when gaining ethical approval with extensive queries being made during the approvals process. This was particularly frustrating for trialists when the application was for a replication SWAT which had already received ethical approval in another host trial.

Barriers to approval reported by trials participating in the PROMETHEUS programme included ethics committees being concerned that participants would not explicitly be asked to give consent to take part in the SWAT and that participants would not be informed about the SWAT. From a methodological perspective, pursuing consent for a recruitment or retention SWAT may dilute SWAT intervention effects through the Hawthorne effect and/or resentful demoralisation. Recruitment SWATs themselves due to their nature could be very confusing for potential participants as it would not be clear whether they were being asked for consent to take part in the SWAT or the host trial itself, and in the case of recruitment SWATs the participant may not yet have been approached about the host trial. Similarly, for retention SWATs, there is potential for confusion if additional consent is sought. As a result, it may therefore be preferable to not request explicit consent for SWAT participation, although, in accordance with Good Clinical Practice and Research Ethics, this should be considered on an individual basis to ensure participants' interests and rights are protected. It may therefore be appropriate to discuss the need, or not, for consent with the study PPI group or other relevant stakeholders whilst reviewing other elements of the SWAT and to reflect on this discussion, and the agreed consensus, in the submission to the research ethics committee.

Furthermore, existing consent arrangements of the host trial may provide appropriate consent for SWAT inclusion. For example, if the host trial has obtained consent to contact a participant by a range of methods (e.g., postal, email, telephone, SMS) then this would cover SWAT interventions such as newsletters, different cover letters, or SMS reminders. Furthermore, many trials routinely now obtain consent to contact for future research opportunities which would also cover SWAT inclusion for both recruitment and retention interventions.

Where notification and consent of participants regarding the SWAT are not deemed to be required, this can be challenging to explain to a research ethics committee and so the PROMETHEUS team was frequently asked for advice.<sup>133</sup> We suggest that study teams should document explicitly what will happen as part of the SWAT, when this will occur, and which elements comprise host trial procedures and which comprise SWAT processes.<sup>133</sup> It may also be useful to reflect on the lack of current evidence in relation to recruitment or retention, the need to improve research efficiency, and to note any precedent set by similar SWATs which have been approved and conducted.<sup>133</sup>

In addition to the concerns around informed consent, there was also concern that SWATs are underpowered and additional information was requested to explain how SWATs were analysed and the need for replications and evidence synthesis (See Chapter 9).

In many instances, such issues were often resolved when sufficient information and justification was provided, and adding PPI input to these justifications would undoubtedly allay concerns further. Despite this, there was however one instance where the ethical committee refused approval of the SWAT. The ethics committee viewed the question proposed by the SWAT (different information in two cover letters with a follow up questionnaire) as valuable, however, deemed that as this question could be considered for any study that this should be undertaken as a new non-interventional research study and not within the confines of the host trial. Despite the host trial team providing additional information on SWAT methodology, support of their oversight committees, lack of impact on the research sites, and links to the PROMETHEUS programme, it was not possible to change their view on this and so the SWAT did not commence.

Where SWATs are low risk, and supported by PPI input, a risk-based approach may be relevant to ensure streamlined activity. It is worth noting that most, if not all the SWAT strategies and the SWATs themselves undertaken within PROMETHEUS are low risk to host trial participants. In PROMETHEUS we have generally tested strategies that are already being used in trials, but without any real evidence regarding their effectiveness. As a consequence, it is likely that one of the key issues related to risk is the need to blind host trial participants to their participation in the SWAT, in order to avoid (or at least reduce) information (outcome) bias. This would be beneficial for the individual SWAT and for the wider research community; if there were fewer barriers, more SWATs could and perhaps would be undertaken which would increase the evidence base. In the case of recruitment and retention, this would lead to more effective strategies which would result in more efficient research being undertaken and so prevent research waste, which itself becomes an ethical issue when it results in future patients receiving effective treatments more slowly (or prolongs harmful treatment).

During the course of the PROMETHEUS programme, HRA also identified this barrier. As a result, they sought collaboration from both the PROMETHEUS group and Trial Forge<sup>107</sup> to develop a more efficient streamlined approvals process for SWATs. The proposed revised procedure would result in one team holding overall permission for the overarching SWAT evaluation with a specific protocol to be embedded by other host trials. This approach is similar to the approvals process used for the Christmas card SWAT<sup>49</sup> where an overarching approval was granted for the SWAT which covered all

host trials involved. As detailed in Table 11 the proposed HRA process differs slightly as, depending on the type of SWAT, a graded approach to ethical approval is required.

In Group 1, no additional ethical approval would be required, however, HRA study-wide review would still be required. The HRA have deemed this to be appropriate on the basis that these SWATs do not involve or impact directly on participants. Examples of such SWATs include randomising sites to different methods of study training.<sup>46,120</sup>

The HRA propose that Group 2 SWATs would be those which may be participant focussed but are deemed to be very low risk, modest interventions, for example sending a festive greetings card or newsletter to improve retention (Coleman et al, Mitchell et al). Such overarching SWATs would only require ethical approval and HRA study wide review required, should the process targeted be detailed in approved study documentation.

Ethical and HRA review would be required for any Group 3 SWAT due to the impacts this would have on trial procedures or information provided to participants, for example use of alternative information sheets or cover letters.<sup>68,134</sup> These types of strategies are still low risk to participants, but they impact on direct participant-facing materials, which always require ethical approval, so such a review would simply be in line with current review processes.<sup>68, 68,134</sup>

Where required, these approvals may be gained at the start of the trial or as a substantial amendment if the trial has commenced. See Appendix 3 for further details of this.

Table 11: SWAT groups and approvals required

Group	Description	REC Approval Required	UK approvals and amendments
1	The activities targeted by the SWAT are not of a type that needs to be described in detail in a host trial protocol (and so have not required previous approval) but are operational or management activities generally done by the central trial team. For example, randomising the host trial chief investigator to visit some recruiting sites and not others to see if this improves recruitment	The overarching SWAT does not need REC approval, nor are amendments to host trial REC approvals required.	The overarching SWAT would need study wide review.  UK amendment is not required for individual host trials evaluating the SWAT.
2	The SWAT interventions affect participants but affect a process that does not <i>need</i> to be described in detail in a host trial protocol (and so have not <i>required</i> previous approval).  These interventions are provided to the research team via operational or instruction documents, rather than a protocol, which are not reviewed by the REC. However, in some cases these interventions are also described in the protocol. Where this is the case, the intervention remains a Group 2 SWAT due to the low risk and unobtrusive nature of the intervention. For example, randomising whether trial participants are sent a greetings card or not to see if this improves retention.	The overarching SWAT needs REC approval.  Where the host trial protocol does not detail the process targeted by the SWAT, no amendments to host trial REC approval required.  Where the process targeted by the SWAT is described in the host trial protocol, an individual SWAT that changes the process should be submitted to the host trial REC as a substantial amendment.	The overarching SWAT would need study wide review.  A non-substantial amendment to the host trial for UK approval should be submitted.  Where the process targeted by the SWAT is described in the host trial protocol, an individual SWAT that changes the process should be submitted as a substantial amendment to the host trial.
3	These SWAT interventions change the design of the host trial, change procedures undertaken by participants, or the information received by participants. For example, randomising whether potential participants receive a standard information sheet or one designed by PPI members.	The overarching SWAT needs REC approval.  Amendments to host trial REC approvals will be required and will be submitted as a substantial amendment to the REC that approved the host trial.	The overarching SWAT would need study wide review.  A substantial amendment to the host should be submitted.

### Conclusion

The success of obtaining approvals of SWATs undertaken within the PROMETHEUS programme has varied, largely depending on the knowledge and understanding of this methodology. In the main, difficulties have been relatively minor, centring around delays to obtaining approvals, and the associated hassle for busy trialists of responding to ethical queries. There have however been a couple of more substantial issues where sponsorship or ethical approval was not provided. In such instances, this stands to limit the development of the evidence base and so increase or sustain existing levels of research waste.

First and foremost, the key learning for all stakeholders involved in SWAT approvals is developing a better understanding and awareness of the importance of such methodological research, and the

nuances of SWATs versus multicentre trials. The coordinated SWAT approach proposed by the HRA is a step in the right direction and should over time help to improve and simplify the undertaking of SWATs moving forward.

In addition, there are specific learning points for individual stakeholder groups:

#### *Lessons learnt for Sponsors*

- The changes proposed by the HRA to SWAT approvals should be communicated clearly and applied consistently when made publicly available.

#### *Lessons learnt for oversight committees*

- TMG members should encourage and support the undertaking of SWATs within their trials.
- TSCs should routinely review SWAT progress. Whether a DMC review is required is dependent on the host trial and SWAT.

#### *Lessons learnt for trial teams and methodologists undertaking SWAT research*

- The input of PPI members with regards the need for SWAT information and/or consent should be discussed during review of intervention design and content.
- The HRA approach to SWAT approvals should be utilised wherever possible. Where SWATs are prospectively registered with the SWAT repository, consideration should also be given to obtaining approvals to enable building of the evidence base simply and effectively.

#### *Lessons Learnt for journals and reviewers*

- Reviewers should consider the need to query informed consent with authors on the basis that this may not be feasible or appropriate for a SWAT. Transparent reporting (See Lessons Learnt: Reporting) should help to facilitate this.

## Chapter 8: Lessons learnt for implementing SWATs

### *Chapter overview*

In this chapter, we describe our experience of implementing SWATs, and present case studies of different approaches to implementing SWATs using a case study of individual, standalone SWATs testing the effectiveness of pens for retention; and case studies of coordinated SWATs undertaken – one testing a training course for staff recruiting trial participants, and the other testing the effectiveness of sending Christmas cards to participants on retention rates. This chapter also highlights the pros and cons of standalone and coordinated SWATs.

### *Introduction*

Due to the sample size of SWATs often being constrained to the size of the host trial, SWATs should be designed for replication and, ultimately, to be meta-analysed together with other similar SWATs of the same strategies. This provides the statistical power needed to identify small yet meaningful differences and ensure generalisability across different patient groups. However, when undertaking a meta-analysis, careful consideration needs to be given about when it is appropriate to combine data from SWATs in different patient populations, health conditions, and trial contexts.<sup>135</sup>

SWATs can be undertaken in one of two ways; individually and coordinated.

### *Individual SWATs*

Individual SWATs can be undertaken by individual host trial teams on an *ad hoc* basis, based on the interests and needs of the trial team. Individual SWATs independently test the same or similar interventions, within host trials at a time or opportunity that is convenient to the host trial. Multiple evaluations over an extended period may therefore be required in order to reach a definitive answer.

Individual SWATs can also be linked, such as in the MRC START programme, which developed an optimised patient information leaflet intervention and a multimedia information intervention; testing them across multiple host trials in SWATs over several years.<sup>25, 26</sup> The TRECA (Trials Engagement in Children and Adolescents) study similarly aimed to embed the use of multimedia information resources into six host trials in the United Kingdom in a coordinated way.<sup>30</sup> In both MRC START and TRECA, each host trial had a linked but separate SWAT protocol, a separate ethical approval, and analysis, with the aim of undertaking a meta-analysis to determine the effectiveness. A hybrid approach to individual SWATs can be undertaken, with SWATs initially being conducted on an *ad hoc* basis, and then later individual linked SWATs might be conducted to provide a definitive



answer to the research question. The case study of individual SWATs of pens for retention in this chapter is an example of such a hybrid approach.

#### *Advantages of Individual SWATs*

The main advantage of individual SWATs is that these can be easier to implement than coordinated SWATs. There is no reliance on all host trials being able and ready to embed the SWAT at the same time and so trialists can implement a SWAT at a time which best suits their trial, as was the case with the pens for improving retention SWATs.

Individual SWATs also have the benefit of enabling the targeting of specific groups. The majority of SWATs in this case study were undertaken in trials with an older, female population, however, in order to further fill the evidence base, additional, individual replications of this SWAT are required in younger and/or male populations.

They also enable intervention modifications as required or preferred by the host trial. For example, in this case study, SWATs tested the use of a trial or university branded pen, as per the individual trial's choosing. Other options included unbranded pens made from recycled materials or with limited plastic components to reduce carbon footprint, although these were not tested here.

Individual SWATs can take a long time to produce enough evidence to determine whether a strategy is effective or ineffective, and there is a clear economy of effort associated with coordinated SWATs. It is however also possible to derive sufficient data to boost, if not complete the evidence base, with individual SWATs. Three of the five pens for recruitment SWATs were completed individually via the PROMETHEUS programme which demonstrates that targeted effort to support can provide sufficient additional data to boost the evidence base and support more informed trial process decisions.

#### *Disadvantages of individual SWATs*

Due to the sample size of SWATs often being constrained to the size of the host trial, sample size or power calculations are often not undertaken for individual SWATs, which has led to criticisms about the lack of power calculations and also, generally, being underpowered.<sup>36</sup> Even with replication of individual SWATs, it will take a long time to produce enough evidence to determine whether a strategy is effective or ineffective. Of the 137 strategies identified in the Cochrane systematic reviews of recruitment and retention strategies, only three strategies have robust evidence of their effectiveness.<sup>23</sup> Both Cochrane reviews called for more 'depth' to the evidence base, to allow more replication SWATs to develop the evidence more quickly for particular strategies. However, this can take a long time. For the pen SWATs, the first was reported by Sharp in 2006, and it was not until 14 years later in 2020 that there was sufficient data to determine the effectiveness of using pens as a

retention strategy. There is still a need for further evidence in certain populations (younger people, and men), so this case study highlights that generating evidence in an *ad hoc* way using individual SWATs can be slow.

#### *Case study of individual SWATs: Pens for retention*

##### *Background*

Postal questionnaires can be susceptible to low response rates. These are frequently used in trials to collect patient-reported outcomes, and so it is important to use effective strategies to maximise response rates.

The Cochrane review by Brueton *et al.*, (2013)<sup>31</sup> found that non-monetary incentives (e.g., pens, certificates, or other similar tokens) with postal questionnaires had no clear effect in improving response rates or retention. The review however grouped all non-monetary incentives together and so individual strategies within this grouping may subsequently prove to be effective.

Including a pen with a questionnaire, as a non-monetary incentive, has been suggested as a low-cost intervention that may improve response rates. In the context of trial retention, firstly providing a pen with a questionnaire facilitates completion of this. Secondly, the inclusion of a pen acts as an acknowledgement of the participant's involvement in the study, and as a reminder, which may make the recipient more likely to complete the accompanying questionnaire. Thirdly, the theoretical basis of reciprocation, results in participants feeling obliged to respond to this positive behaviour with the same in return.<sup>136-139</sup>

Prior to the PROMETHEUS programme commencing, there was some evidence (two SWATs)<sup>140, 141</sup> to suggest that this may be an effective strategy in improving questionnaire response rates and reducing the number of reminders required. Given the existing evidence, and its match to PRioRiTY II<sup>111</sup> priority six, this strategy was selected as a medium-priority question for the PROMETHEUS programme.

##### *Aims and Objectives*

The individual SWATs were designed to evaluate the effects of providing a pen with a follow-up questionnaire on retention rates.

Results from the series of pen SWATs undertaken prior to or within the PROMETHEUS programme were then combined via meta-analysis by the Trial Forge Initiative in order to determine a definitive answer as to the effectiveness of pens for retention.

## Methods

### Individual SWAT Methods

All pen retention SWATs undertaken in PROMETHEUS followed a core protocol. The only difference was that two SWATs<sup>68, 92</sup> included additional secondary outcomes and associated analyses.

### Participants, Interventions and Outcomes

Table 12 summarises the participants, interventions and outcomes associated with the individual pen SWATs.

Table 12: SWAT Participant, Interventions and Outcomes

<i>Participants</i>	<p>Any participant in the host trial who was due to be sent a follow-up questionnaire at the study's primary outcome time point was included in the SWAT sub-study.</p> <p>Participants who withdrew from host trial follow-up before their follow-up questionnaire was due or those who had received their follow-up questionnaire prior to the start of the pen sub-study were excluded.</p>
<i>Intervention</i>	A pen printed with either the trial or university logo.
<i>Control</i>	Standard practice for the host trial, i.e., no pen.
<i>Outcomes (Primary)</i>	The proportion of participants in each group who return the questionnaire.
<i>Outcomes (Secondary)</i>	<ul style="list-style-type: none"><li>• Time to response: defined as the length of time taken to return the questionnaire</li><li>• Completeness of response: defined as the number of questions completed</li></ul> <p>Cunningham-Burley<sup>92</sup> and James<sup>68</sup> also assessed additional secondary outcomes</p> <ul style="list-style-type: none"><li>• Whether a reminder notice is required (defined as the number of participants requiring a reminder mailing divided by the number of participants who were sent a questionnaire).</li><li>• Cost-effectiveness</li></ul>

### *Randomisation and Blinding*

In each SWAT, participants were allocated using simple randomisation 1:1 to either intervention or control. A statistician who was independent of the questionnaire mailing activity generated the randomisation sequence.

Participants were not informed of and so could not provide informed consent for their participation in the SWATs. Given the nature of the intervention, it was however not possible to blind the participants to receipt, or not, of a pen with the questionnaire nor was it possible to blind research staff implementing the SWAT.

### *Sample Size*

Given their nature, the sample size for each SWAT was determined by the number of participants due to receive their postal questionnaire at the primary outcome time point within the host trial.

### *Analyses*

For each SWAT, data were analysed using the version of Stata current at the time of analysis, on an intention-to-treat basis, using two-sided tests at the 5% significance level.

### *Primary outcome*

- The proportion of participants in each group who returned the questionnaire was compared using logistic regression
- As the SWAT by James et al was factorial (social incentive cover letter, pen) the interaction term was also assessed in the logistic regression model.

### *Secondary outcomes*

- The difference in time to response between the two groups was compared by a Cox proportional hazards model
- Completeness of response between the two groups was compared using a linear regression model<sup>68, 92</sup> or logistic regression<sup>81</sup>
- Requirement for a reminder mailing was compared using logistic regression
- The difference in costs between the intervention and control groups including direct and indirect costs.

Each trial adjusted the analysis models for main trial allocation. In addition, James adjusted for age and gender,<sup>68</sup> and Mitchell for trial site.<sup>81</sup>

### SWAT Meta-Analysis Methods

A meta-analysis of all relevant published SWATs was undertaken by the Trial Forge initiative in 2020 as part of their evidence pack series.<sup>119</sup>

#### Results of PROMETHEUS SWATs testing the use of pens for retention

In total, 4,550 participants were randomised into pen SWATs funded by the PROMETHEUS programme: 2,276 to receive the intervention, and 2,274 to receive the control. Fifty participants either withdrew or were deceased before the SWATs could be implemented leaving 4,500 participants included in the analyses: 2,249 intervention, and 2,251 control. The majority of participants were female (67.1%), and the average age was 64 years.

The results of the individual studies are reported separately.<sup>68, 81, 92</sup>

#### Comparison with Existing Literature

Prior to the PROMETHEUS programme two trials, consisting of 8,512 participants, reported within a 10-year period,<sup>140, 141</sup> suggested that when pooled, as shown in Figure 3, the inclusion of a pen with a follow-up questionnaire may be an effective strategy for improving response rates: Pooled Odds Ratio 1.21, 95% CI 1.05 to 1.40, p=0.01).




Figure 3: Pooled Odds Ratio for Effectiveness of Pen with a Follow Up Questionnaire (Pre PROMETHEUS) {Sharp 2006, Bell 2016}<sup>140, 141</sup>

When the results of Cunningham-Burley,<sup>92</sup> James,<sup>68</sup> and Mitchell<sup>81</sup> were added to the meta-analysis,<sup>138</sup> this contributed a further 4,500 participants (an increase of 52.8%). The pooled effect as shown in Figure 4, indicated that including a trial pen with a questionnaire probably increases retention and response rate (pooled OR 1.21, 95% CI 1.09 to 1.35).<sup>119</sup> The evidence when combined has moderate certainty GRADE evidence overall, but high GRADE evidence for an older population.




Figure 4: Pooled Effect of Including a Pen with a Follow Up Questionnaire {Sharp 2006, Bell 2016, Mitchell 2020, Cunningham-Burley 2020, James 2020}<sup>140, 141, 92 68 81</sup>

### Cost-effectiveness

The cost of retaining one additional participant due to inclusion of a pen with a questionnaire mailing is £25, which is cost-effective if based on the cost of enrolling an average participant in an NIHR funded trial being £2,200.<sup>119</sup>

### Discussion

Over 14 years, five individual SWATs have been undertaken sequentially to ascertain whether the inclusion of a pen with questionnaire mailings is an effective retention strategy.<sup>68, 81, 92, 140, 141</sup> Four of the five SWATs were conducted within a four-year period (2016 – 2020),<sup>68, 81, 92, 140</sup> and three of these were conducted within the PROMETHEUS programme.<sup>68, 81, 92</sup>

The pooled estimate of effect for all five pen SWATs for retention showed an increase in retention of 1.9% (95% CI 0.0 to 3.7%).<sup>142</sup> Depending on population demographics, there is moderate or high certainty evidence in these estimates and so the recommendation is that trialists should include pens to increase retention rates in trials that use postal questionnaires. The bulk of replications undertaken were in female and older populations and so evidence for younger and male populations should be the focus of any further replications.

The costs associated with the inclusion of a pen for retention were estimated, using generic trials information, to be around £25 per additional participant retained.<sup>142</sup> Cunningham-Burley *et al.*, identified a cost of £10.56 per additional participant retained,<sup>92</sup> while others have found lower rates e.g., a cost of \$6.98 (approximately £5.60) for an additional participant to be retained.<sup>142</sup> While these costs are not inconsequential, the intervention is relatively low-cost compared to other

retention interventions and in a large randomised controlled trial, these costs may well be possible to accommodate.

### *Coordinated SWATs*

Simultaneous SWATs are also a type of coordinated SWAT, but they are undertaken across multiple host trials at the same time. Simultaneous SWATs have one protocol for all the host trials, one ethical approval, one analysis, and a write up.

### *Advantages of coordinated SWATs*

A coordinated SWAT usually provides a much larger sample size than individual SWATs. They speed up the process of generating evidence and allow high-quality evidence to be generated at scale for one research question of interest. Due to their scale, it may be possible to embed the SWAT within enough host trials to allow for sufficient evidence to reach a definitive answer in one over-arching evaluation. As the same protocol is standardised across the different host trials and patient populations, data can be pooled in a meta-analysis with greater certainty. They are much faster than undertaking the same number of individual SWATs and then undertaking a meta-analysis. They are also much more efficient, typically requiring just one ethical and regulatory approval for all the SWATs, with one central coordination point which also reduces the burden on host trial teams. They also benefit from an economy of scale, as well as having one central team instead of multiple host trial teams planning, undertaking, analysing, and reporting the SWAT. They can be a useful methodology for testing more complex strategies that may be more challenging and/or costly for individual host trial teams, such as delivering a training course to trial recruiters.

### *Disadvantages of coordinated SWATs*

There are possible disadvantages to undertaking coordinated SWATs. They require coordination from a central point, which includes recruiting host trial teams. This requires a network or infrastructure to exist, to allow host trials to be recruited. In the UK, this has been possible through the infrastructure network established by the NIHR, particularly through the Clinical Research Networks (CRNs), through the speciality groups such as the Surgical Speciality Group, and through the CTU Network.

Due to their novelty and complexity, it may be more difficult to obtain regulatory approvals than just for a single SWAT. As part of PROMETHEUS, we have worked with the HRA in the UK to develop new regulatory approval processes for SWATs, which aim to make it easier for simultaneous SWATs to be approved. Chapter 7 outlines this in detail, and Appendix 3 provides a copy of the draft HRA guidance on SWATs.

There are also logistical challenges. It may be difficult to set up a coordinated SWAT across multiple host trial teams and trials units. For example, all the host trial teams must be ready for the recruitment/retention strategy to be embedded in the SWAT at a similar time. Delivering the SWAT in multiple host trials may also limit flexibility to deliver the SWAT strategy; for example, being restricted to the extent to which the SWAT strategy may be adapted to the host trial, and if the SWAT involves a strategy such as attending a training course, the dates of the training may be limited. Another challenge is that routinely collected data in the participating host trials may differ. Where host trial teams are external to the SWAT coordinating team, it may be difficult to obtain the SWAT data from host trials. Ways to address this are to establish clear timelines for transfer of data; identify during the planning stages how long it is likely to take to clean and securely transfer the data once generated; identify who will clean and transfer the data; and make sufficient funds clearly available to the host trial team to cover the costs of data preparation and sharing.

#### *Case studies of coordinated SWATs*

In PROMETHEUS, we undertook two simultaneous SWATs, which we present here as case studies. The first coordinated SWAT focused on recruitment and was a feasibility study of staff training to improve participant recruitment into surgical trials and was embedded across four host trials.<sup>51</sup> The second was a retention SWAT, which tested the effectiveness of sending Christmas cards to trial participants on retention rates, across eight host trials, which was published as part of the BMJ Christmas edition.<sup>49</sup>

#### *Case study - Staff training to improve participant recruitment into surgical randomised controlled trials: a feasibility Study Within A Trial (SWAT) across four host randomised controlled trials simultaneously*

##### *Background*

The training of trial recruiters has been identified as the top priority topic for recruitment research according to the Directors of UK CTUs,<sup>33</sup> and has been highlighted in a James Lind Alliance priority setting exercise for recruitment research.<sup>110</sup> A systematic review of recruiter training showed that training programmes were well received and increased recruiters' self-confidence.<sup>143</sup> However, only three of the 17 studies included were RCTs. A more recent systematic review looking at the effectiveness of staff training on recruitment concluded that further work on developing a substantial evidence base around the effectiveness of education and training interventions for recruiters to trials is required.<sup>144</sup> As a result, there is a lack of evidence on the effectiveness of staff training to improve recruitment.



### *Aims and objectives*

To test the feasibility of undertaking a coordinated SWAT to train staff who recruit participants into surgical trials, by assessing key uncertainties around recruitment, randomisation, intervention delivery, and data collection.

### *Methods*

#### *Host trial and participant recruitment*

Surgical trials recruiting or likely to be recruiting participants to the host trials in UK hospitals in April 2019 were invited to take part in this SWAT. To be eligible, host trials had to be undertaking face-to-face recruitment. Staff (surgeons and nurses) recruiting participants to the host trials were asked to express interest in attending the training.

#### *SWAT Intervention*

The one-day training course for staff recruiting participants into surgical trials was developed by The University of Bristol's QuinteT team (Qualitative research integrated within Trials).<sup>145</sup> The training aims to share experiences, raise awareness of the hidden challenges of recruitment and equip attendees with strategies to optimise recruitment and informed consent. All staff members involved in recruiting participants at each hospital site were invited to attend a one-day training course relevant to their profession (one date was offered to surgeons, and one to research nurses). Academic researchers within the University of Bristol's QuinteT group delivered the training. The one-day training was supplemented with the GRANULE (Generating surgical recruiters to randomised trials) online e-learning course (<https://learn.nihr.ac.uk/course/view.php?id=385>), which was emailed to all sites in the intervention arm after the course, or in place of if staff were unable to attend.

#### *SWAT control group*

The control group did not receive the recruitment training.

### *Outcomes*

The primary outcome of interest was the ability to recruit multiple surgical trials simultaneously. To be considered feasible, we needed to enrol a minimum of two host trials. Secondary outcomes were:

1. Recruitment: Numbers of recruiting sites and recruiting staff from each site enrolling into the SWAT
2. Randomisation: Number of sites recruited, number of sites randomised and any reasons for sites dropping out after recruitment

3. Intervention delivery: number of intervention training course groups initiated, staff attendance at training in the intervention group, number of participants per group, any reasons for non-attendance or failures in intervention delivery.
4. Acceptability of the training
5. Staff confidence in discussing trial recruitment with potential participants immediately before, immediately after, and at 1–3 months post-training
6. Participant screening and recruitment rate (defined as the proportion of eligible participants who gave their consent and were randomised into the host trial six months following delivery of the course).

#### *Randomisation*

Randomisation was performed separately for each host trial at the cluster (i.e., recruiting hospital site) level. On expressing willingness to participate in the training, recruiting sites within each host trial were randomised to be offered the training workshop (intervention group) or no training (control group) on a 1:1 basis. A computer-generated randomisation schedule was generated using permuted blocks, stratified by recruiting trial.

#### *Data collection*

Online questionnaires were distributed to participants using Qualtrics (QualtricsXM, Provo, UT, USA). Participants were asked to complete questionnaires one month before the workshop (all participants), immediately after the workshop (intervention group only), and 1 to 3 months after the training (all participants).

#### *Statistical analysis*

Statistical analyses were undertaken on an intention-to-treat basis, using a 5% significance level. Baseline data are reported descriptively by SWAT group, using counts and percentages for categorical data, and mean and standard deviation for continuous data. The paired t-test was used to compare pre- and post-training workshop responses, and follow-up are reported as mean and standard deviation by arm. A linear regression model adjusting for baseline score and allocation was run on each of the 11 questions to compare the self-confidence at follow-up between the arms. Recruitment rate and the number of patients screened post-training were compared using linear regression adjusting for host trial and SWAT intervention.

## Results

### Feasibility of recruiting host trials

We identified 12 host trials that were eligible to participate in the SWAT and four trials (33%) comprising 91 sites participated. Recruitment of surgical host trials occurred as planned, and we recruited four host trials in a six-month period.

Details of the host trials can be found in Table 13.

Table 13: Details of the trials included in the PROMETHEUS Training SWAT

Trial	Area	Interventions	Number of sites	Target Sample Size	Primary outcome (time point)
DISC	Dupuytren's Contracture in adults over 18	injection of collagenase or surgery (>=18 years old)	30	710	Patient Evaluation Measure (2years)
IntAct	Rectal cancer in adults over 18	intra-operative fluorescence angiography (IFA), or white light endoscopic surgery	25 (14 UK)	880	Clinical anastomotic leak rate within 90 days post-operation (1 year)
PROFHER-2	Acute 3- and 4-part fractures of the proximal humerus in patients aged over 65 years	reverse shoulder arthroplasty, hemi arthroplasty or non-surgical treatment	40	380	Oxford Shoulder Score (2 years)
START: REACTS	rotator cuff tears that cannot be repaired	arthroscopic debridement or arthroscopic debridement with the InSpace balloon	16	212	Constant-Murley score collected 12 months after surgery.

#### *Feasibility of recruiting sites, participants and randomisation*

Across the four host trials, 29 sites agreed to participate (32%) and were randomised to intervention (15 sites, 29 staff) or control (14 sites, 29 staff). The 58 recruiting staff who enrolled into the SWAT were mostly research nurses (58.6% control and 41.4% intervention) and surgeons (31.0% in both groups). We found it feasible to recruit sites and recruiting staff.

#### *Feasibility of training course*

There were 29 staff in the intervention group who expressed an interest in attending the workshop, and of these, 11 (37.9%) attended. Those who attended the workshop were all female (100%), predominantly research nurses (81.8%) and the modal age group was 40-49 years (45.5%). Only one surgeon agreed to attend the workshop, which resulted in the surgeons' workshop being cancelled. We initiated just one workshop - for Research Nurses, which was relatively well attended, so the training course initiation and attendance was partially feasible.

#### *Acceptability of the training course*

Research Nurses who attended the workshop felt positively about the format of the course with a mean score of 9.3 (0=very poor, 10=excellent; SD 1.0). Participants felt they "learnt a lot" with a mean score of 9.2 (SD 0.9). They also felt that the workshop would make "a lot of difference" to their future recruitment practices, with a mean score of 8.7 (SD 0.9).

#### *Impact of recruitment training on self-confidence and awareness*

When comparing pre-post scores in the intervention group, participants rated their confidence in discussing recruitment higher after the workshop with an increase in mean score of 1.45 (95%CI 0.70 to 2.21,  $p = 0.002$ ). They were also more confident about overcoming recruitment challenges with an increase in the mean score of 2.18 (95%CI 1.19 to 3.17,  $p = 0.001$ ). However, the change in their awareness of the recruitment challenges was not statistically significant with an increase in score of 0.45 (95%CI -1.06 to 1.97,  $p = 0.52$ ).

#### *Impact of training on screening and recruitment rates*

At baseline, the number of patients approached and/or recruited was similar across both groups (Table 14). At 6 months post-training, there was no evidence of a difference in screening between sites randomised to the intervention versus sites allocated to control (coefficient -0.35, 95% CI -7.84 to 7.15,  $p = 0.92$ ). Over the six months post-training, the average eligible to recruited conversion rate across the participating studies was 58%; it was 55% in the intervention group and 63% in the control group. Linear regression of sites (63%) with screening activity post training identified there was no evidence of a difference in recruitment rate between the intervention and control groups (coefficient -0.07, 95% CI -0.43 to 0.29,  $p = 0.66$ ).

Table 14: PROMETHEUS Training SWAT - number of patients approached and agreeing to participate at baseline and follow-up, as reported by recruiters per participant defined typical week in the last month

Summary	Control		Intervention		Total	
	Baseline (n=22)	Follow-up (n=21)	Baseline (n=13)	Follow-up (n=16)	Baseline (n=35)	Follow-up (n=37)
<b>Approached</b>						
0	2 (9.1)	9 (42.9)	0 (0.0)	7 (43.8)	2 (5.7)	16 (43.2)
1-3	11 (50.0)	1 (4.8)	8 (61.5)	0 (0.0)	19 (54.3)	1 (2.7)
4-6	5 (22.7)	5 (23.8)	5 (38.5)	5 (31.2)	19 (54.3)	10 (27.0)
7-10	3 (13.6)	5 (23.8)	0 (0.0)	2 (12.5)	3 (8.6)	7 (18.9)
>10	1 (4.6)	0 (0.0)	0 (0.0)	0 (0.0)	1 (2.9)	0 (0.0)
Missing	0 (0.0)	1 (4.8)	0 (0.0)	2 (12.5)	0 (0.0)	3 (8.1)
<b>Recruited</b>						
0	3 (13.6)	0 (0.0)	1 (7.7)	1 (6.3)	4 (11.4)	1 (2.7)
1-3	12 (54.6)	11 (52.4)	8 (61.5)	9 (56.3)	20 (57.1)	20 (54.1)
4-6	6 (27.3)	6 (28.6)	3 (23.1)	3 (18.8)	9 (25.7)	9 (24.3)
7-10	1 (4.6)	3 (14.3)	0 (0.0)	1 (6.3)	1 (2.9)	4 (10.8)
>10	0 (0.0)	0 (0.0)	1 (7.7)	0 (0.0)	1 (2.9)	0 (0.0)
Missing	0 (0.0)	1 (4.8)	0 (0.0)	2 (12.5)	0 (0.0)	3 (8.1)

### Discussion

This SWAT demonstrated the feasibility of recruiting multiple surgical host trials to undertake a coordinated SWAT to evaluate the effectiveness of a staff training course. We found that there was no evidence of a difference in the number of patients screened between the intervention and control groups. Equally, there was no evidence of a difference in the number of eligible patients recruited and enrolled into the host trials between the two groups.

Undertaking a coordinated evaluation across multiple host trials can significantly speed up generating the evidence base for recruiting participants into trials; and, consequently, improving trial efficiency and reducing research waste. However, this is only the case if key personnel attend training.

We found it challenging to engage surgeons to attend the face-to-face training, with only one surgeon being available to attend the training. As a result, a key cohort of recruiting staff was not represented which may have limited the findings. Delivering the coordinated SWAT in four host trials limited our ability to be flexible with the training dates, which may have made it more difficult for surgeons to have sufficient notice to fit this around their clinical commitments. There was an online component to the course with GRANULE, which was a reduced version of the face-to-face course, to supplement the training. Fully online courses may afford an opportunity to maximise attendance by removing the need to travel. Given the increasing use of remote, video conferencing meetings and training technologies arising from the COVID-19 pandemic, this is now even more pertinent and therefore the medium by which training is delivered should be an important consideration for future studies.

Recruiting host trials for this coordinated SWAT required significant logistical organisation and demanded a central coordination point. The process for recruiting host trial teams typically involved an initial approach to the Chief Investigator (CI). If the CI was interested, then they would discuss the SWAT with their team to obtain buy-in, including from the TMG and Trial Manager. If there was buy-in, then there were further discussions about how the SWAT would align with the host trial, how sites would be approached, and about other logistical and methodological issues. This alignment was not always possible, and one host trial was not recruited because although interested in the training SWAT, they had different ideas about how they wanted the SWAT to be delivered in their trial. For instance, they wanted significantly more training dates for both nurses and surgeons than we were able to offer, and wanted outcomes to be followed up over their entire recruitment phase of 3 years, which we were unable to do for resource reasons.

*Case study - Bah humbug! Association between sending Christmas cards to trial participants and trial retention: randomised study within a trial conducted simultaneously across eight host trials*

#### *Background*

Poor participant retention can adversely impact study validity.<sup>24, 33, 36</sup> Christmas cards are often used to encourage participants to continue participation in a trial. A survey of UK-registered CTUs found that 40% had used Christmas cards as a retention intervention.<sup>33</sup> However, there is no evidence that providing Christmas cards improves retention. For those trials that use resources (both time and money) to send Christmas cards, there is a potential opportunity cost in that these resources might be better invested in other potentially more effective interventions. On the other hand, there could be a case for routine use of this incentive if there was evidence to support it.

One way to evaluate the effectiveness of Christmas cards on retention rates, and allow for rapid collection of evidence, would be to plan to do the same coordinated SWAT in several host trials at the same time.

#### *Aims and Objectives*

To determine the effectiveness of sending Christmas cards to participants in randomised controlled trials to increase retention rate at follow-ups, and to explore the feasibility of doing coordinated SWAT across multiple host trials simultaneously.

#### *Methods*

##### *Design*

Randomised coordinated SWAT conducted across eight host trials.

##### *Setting*

Eight randomised controlled trials researching various areas including surgery and smoking cessation.

##### *Participants and sample size*

3223 trial participants who were still due at least one follow-up from their host randomised controlled trial.

##### *Intervention and comparator*

Participants in the intervention group received a Christmas card sent by post. Participants in the control group did not receive a Christmas card.

### *Primary outcome*

Proportion of participants completing their next follow-up (retention rate) within their host randomised controlled trial.

### *Randomisation and Blinding*

Participants were randomised in a 1:1 ratio (separately by each host trial) to either receive a Christmas card in mid-December 2019 or to not receive a card.

### *Statistical analysis*

Intention-to-treat (ITT) analysis was undertaken, using a 5% significance level. Analysis models adjusted for the host trial allocation and the SWAT allocation and were run for each host trial separately. The results of the individual trials were combined using a random effects meta-analysis. The primary analysis compared the retention rate between the two arms using a logistic regression model. The average cost per card sent was calculated using staff time for preparing the cards, and costs for printing and postage. Assumptions for CO<sub>2</sub> emissions calculation were made, based on previous research were applied which assumes the card weighs 10g, is printed on recycled paper, and posted and recycled in the UK.<sup>146</sup>

### *Results*

#### *Host trials and participants*

Table 15 provides a description of the host trials involved. One thousand, four hundred and sixty-nine (1469) participants (age 16-94 years; 70% (n=1033) female; 96% (813/847) white ethnicity) across the eight host randomised controlled trials were involved in the analysis (cut short owing to COVID-19).



Table 15: Descriptions of the host trials involved in PROMETHEUS Christmas Card SWAT

<b>Acronym</b>	<b>Area</b>	<b>Interventions</b>	<b>Follow-up method</b>	<b>Target Sample Size</b>	<b>Primary outcome</b>
<b>C-Gall</b>	Surgery - gallbladder	Laparoscopic cholecystectomy or conservative management.	Postal	430	Short Form-36 at 18 months
<b>CPIT-3</b>	Smoking in pregnancy	Both groups receive smoking cessation service support and contingent shopping vouchers. Intervention group receives additional shopping vouchers (up to £400).	Telephone	940	Self-reported abstinence from smoking for 8 weeks.
<b>DISC</b>	Surgery/ Drug - hand	Injection of collagenase or surgery.	Clinic / postal	710	Patient evaluation measure at 1 year
<b>FUTURE</b>	Female bladder weakness	Urodynamics plus comprehensive clinical assessment or comprehensive clinical assessment only	Postal	1,096	Patient Global Impression of Improvement at 15 months
<b>ProFHER-2</b>	Surgery - shoulder	Reverse shoulder arthroplasty or hemi-arthroplasty or non-surgical	Clinic / Postal	380	Oxford Shoulder Score at 2 years
<b>PUrE</b>	Surgery - kidney	Extracorporeal shockwave lithotripsy or percutaneous nephrolithotomy or flexible ureterorenoscopy with laser lithotripsy	Postal	1,044	EQ-5D-5L at 12 weeks
<b>REFLECT</b>	Dental	Prescription of 5000ppm fluoride toothpaste or usual care.	Postal	1,174	Proportion of participants receiving dental care due to caries at 36 months
<b>SWHSI-2</b>	Wound healing	Negative pressure wound therapy or usual care (normal dressing)	Postal	696	Time (days) to wound healing.

*Impact of sending Christmas cards on retention rates*

No evidence was found of a difference in retention rate between the two arms for any of the host trials when analysed separately or when the results were combined (85.3% (639/749) for cards versus 85.4% (615/720) for no card; odds ratio 0.96 (95% confidence interval 0.71 to 1.29; P=0.77). A cumulative meta-analysis of the results can be seen in Figure 5.




Figure 5: Christmas Card SWAT Cumulative meta-analysis {Coleman 2021}<sup>49</sup>

*Costs and carbon footprint*

The cost of this intervention was £0.76 (€0.91; \$1.02) per participant, and it will have a carbon footprint of approximately 140 g CO2 equivalent per card. One benefit of this approach was the need to only submit one ethics application.

### *Discussion*

This coordinated SWAT investigated the effectiveness of sending a Christmas card to trial participants on retention rates across eight host trials and showed that sending a Christmas card to trial participants does not increase the retention rate.

This study showed that a coordinated SWAT can be embedded successfully by multiple host trials simultaneously. Not only does this increase the speed at which evidence can be accumulated, it has the additional benefit of allowing only one ethics application to be submitted to cover all the work, which will decrease the burden on individual trial teams.

When this coordinated SWAT was initially planned, it was hoped to involve more than ten trials, and over 10,000 participants to ensure that the question could be answered by this one evaluation. For various reasons, the evaluation was delayed, and some host trials could no longer embed the SWAT which reduced the sample size.

This SWAT was successfully implemented across eight host trials, from two CTUs in the UK, at the same time. This is one of the first instances where a SWAT evaluation has been undertaken in this way, and its success should influence other researchers to consider undertaking simultaneous SWATs in the future, to allow answers to the methodological questions that SWATs pose to be obtained more quickly.

### *Conclusions*

In the PROMETHEUS programme, we have demonstrated that it is feasible to adopt different approaches to undertake SWATs. Host trial teams can undertake individual SWATs focusing on their patient populations and recruitment context to add to the wider evidence base, which in time can be meta-analysed with similar SWATs to provide definitive evidence on the effectiveness of a particular strategy. An alternative is the coordinated SWAT, which potentially offers a much more efficient methodology than individual SWATs, requiring just one regulatory approval for all the SWATs through one central coordination point, providing economies of scale to significantly speed up evidence generation.

Both individual and coordinated SWATs have their advantages and disadvantages, and depending on the situation, one approach may be more appropriate than the other. Indeed, either approach may be suitable in some instances, depending on the preferences of the research team and the resources available to them (such as resources to recruit host trials and coordinate the SWAT), and the methods and interventions being used by the potential host trials. Strategies that are specific to the methods or interventions being used in a host trial, or are highly adapted to the host trial, such as offering a

free yoga class to participants to enhance retention,<sup>147</sup> or a patient and family co-developed participant information,<sup>148</sup> may be more suited to an individual SWAT approach. Strategies that are more widely used, such as sending Christmas cards, text message reminders, and financial incentives all lend themselves well to being tested using a coordinated SWAT approach. A mixture of both approaches may be used to provide a definitive answer to a question. For instance, a simultaneous SWAT was successfully undertaken for the recruitment training course to assess feasibility; however, a conclusion on the effectiveness was not reached. A future evaluation of this training could take a coordinated SWAT approach, or a trial team may wish to undertake an individual evaluation by randomising its recruiting sites to receive the training or not.

#### *Lessons learnt for trial teams and methodologists undertaking SWAT research*

- Embedding a coordinated SWAT in multiple host trials simultaneously, using the same SWAT protocol, has been shown to be a feasible, efficient way to scale up and speed up evidence generation using SWATs. This approach should be used more often to build an evidence base to support the selection of recruitment and retention strategies. This SWAT design could be utilised for other methodological aspects such as protocol/medication compliance strategies, and is suitable within most, if not all, research fields.
- Teams wishing to undertake a coordinated SWAT should plan this carefully. Significant coordination is required to develop a common SWAT protocol that adequately covers all potential host trials, recruit host trials, apply for overarching ethical approvals, prepare and deploy the SWAT intervention, and collect outcome data.
- For most trial teams, individual SWATs remain the most convenient method for testing the effectiveness of strategies for their trial population or trial team; however, where resources allow, a coordinated SWAT approach should be considered as it has the potential to provide definitive answers more rapidly and are more efficient.

#### *Lessons learnt for funding bodies*

- Funding bodies should fund SWATs, particularly coordinated SWATs as they are potentially efficient and are likely to provide robust evidence at speed, which in turn can help to reduce research waste.

## Chapter 9: Lessons learnt for statistical considerations

### *Chapter overview*

Randomised SWATs are like any other trial and as such, the statistical considerations for SWATs are similar to those for a typical trial. Although constrained within the context of the host trial, a SWAT should still follow principles for rigorous analysis, to ensure that the findings are considered reliable. Within this chapter sample size, analyses, and costing considerations when undertaking a SWAT will be detailed, alongside examples of our learnings from the PROMETHEUS programme.

### *Sample size considerations*

It is well known that the sample size for a SWAT is constrained by the host trial. That is, for a recruitment SWAT the maximum number of participants will be the number approached to participate in the host trial, and for a retention SWAT, it is the number who are randomised into the host trial. However, SWATs are not necessarily implemented from the start of a trial, they are often embedded mid-way through recruitment/retention, and as such their sample size is reduced.

Typically, in UK trials funded by NIHR, recruitment of eligible patients is found to be between 50% and 88%, with retention seen to be between 80% and 97%.<sup>5</sup> If a SWAT was powered in a similar way to typical trials, with 80% or 90% power (using a two-sided 5% significance level), the host trial would need to be between 1,438 and 26,256 participants, depending on the initial rates seen to detect a difference between 2% to 4% (see Table 16). With the average target sample size of 1,122 for trials published in the HTA Journal,<sup>149</sup> sufficiently powering these assessments using one trial alone would usually be infeasible. Furthermore, should a SWAT have been undertaken using cluster randomisation, the number of participants required would be larger due to the likely correlation of the participants within the clusters. For example, in MRC START, the minimum sample size for cluster SWATs was appropriately inflated to consider cluster allocations, assuming an intraclass correlation of 0.02 in line with estimates from community studies.<sup>25, 26</sup> Thus multiple replications of a single SWAT evaluation are typically needed, where the results from each are combined to produce a more reliable result.

Table 16: Estimated sample size required for a single SWAT to have 80% or 90% power to detect various differences in proportions

<b>Difference</b>	<b>80% power</b>	<b>90% power</b>
50% to 52%	19,614	26,256
50% to 54%	4,892	6,550
60% to 62%	18,672	24,996
60% to 64%	4,618	6,182
70% to 72%	16,158	21,632
70% to 74%	3,950	5,288
80% to 82%	12,076	16,166
80% to 84%	2,890	3,868
90% to 92%	6,424	8,598
90% to 94%	1,438	1,924

Although it may seem inefficient to need to undertake multiple evaluations of the same strategy, this is actually a beneficial approach. Firstly, this allows for the generalisability of the results – Treweek *et al.*, (2020)<sup>150</sup> suggest that to conclude further SWAT evaluations are no longer a priority, it must have been tested in a wide range of contexts. If it were possible to conduct a large enough SWAT in one host trial for example, the findings would only be directly applicable to that specific trial population, so there is a need for replication in further host trials with a different patient population to ensure the findings are externally valid. This allows for the effectiveness of the strategy as a whole to be determined. As such the PROMETHEUS programme planned to fund several implementations of the same SWAT – for instance, the Christmas Card SWAT – to allow for the results to be combined within a meta-analysis.<sup>47</sup> This allowed us to conclude that Christmas cards are inefficient at increasing retention in an adult population, as it has been evaluated within a range of different host trials, over different research areas, and a range of patient populations, all at once.<sup>49</sup>

### *Analysis considerations*

The approach for the analysis should use ‘intention to treat’. Indeed, it is more usual to use ‘pure intention treat analysis’ in a SWAT than is often the case with the host trials. This is because the primary outcome in a SWAT is recruitment or retention, so there is usually no missing data problem that afflicts most other forms of trials.

As with any analysis undertaken within trials, the analysis of a SWAT should be pre-planned, and detailed with a Statistical Analysis Plan (SAP). This could be within the main SAP for the trial, or as a separate stand-alone document – the choice may depend upon at what point of the trial the SWAT is being undertaken. This SAP should detail the data that is to be collected (i.e., the source of the data), the planned outcomes, and the statistical methods to analyse each of the outcomes – similar to any SAP. Within the PROMETHEUS programme we created a template SAP for the host trial teams to adapt for their SWATs.

Typically, for a recruitment SWAT, the primary outcome will be the proportion of eligible participants consenting to be randomised. Although this is not necessarily always the case as was demonstrated in the ‘training’ SWAT which was a feasibility SWAT (*see Chapter 8, case studies of simultaneous SWATs*), and the primary outcome was measures of recruitment confidence, whilst actual recruitment and randomisation rates were secondary measures. But it is always important to report actual recruitment rates to enable all SWATs to be included in a future meta-analysis.

Similarly, for a retention SWAT, the primary outcome will most likely be the proportion of participants who are retained. The definition of a retained participant may vary; it could be those who have completed a follow-up, attended a clinical visit, or another way, but this should be explicitly stated. Ideally, however, one measure should include the number of participants that provided primary outcome data to, again, allow a meta-analysis. Equally, whether all participants, or only those who were due the follow-up (i.e., excluding those who withdrew prior to the follow-up), are to be included should be detailed.

For both types of SWATs, a logistic regression model is an appropriate analysis model. For retention SWATs this model should be adjusted for the host trial allocation and any stratification variables that were included within the SWAT randomisation. The SWAT design should be taken into account when analysing; for instance, if it was a cluster-randomised SWAT the clustering should be accounted for in the analysis.

Thought should be given as to whether it is appropriate to undertake a sensitivity analysis, where those participants who did not receive the SWAT allocation are excluded (for instance if there is a text message sending failure) - though in most instances this is not required.

Secondary outcomes may include time to events, which would be most appropriately analysed with a Cox proportional hazard model, or count data, such as the number of reminders required, which may require a Poisson or negative binomial regression model. Although for each analysis thought

should be given to whether the relevant assumptions are met, and planned analysis adjusted to be suitable (i.e., using a zero-inflated model).

Once several evaluations of a SWAT have been undertaken, a meta-analysis should be performed to combine the results. Typically, it would be expected that each SWAT would be analysed and reported within its own right, thus a two-stage meta-analysis approach, in which the results from each SWAT are combined, would be used. Should it be possible to obtain the individual data from each trial, a one-stage approach could be considered, though there are typically no additional benefits to this approach apart from considering interactions with pre-specified covariates (e.g., age, gender) and ensuring consistency in model adjustments. A cumulative meta-analysis can be useful in determining if the results appear to have converged, which can be used to help judge if another evaluation is required.<sup>147</sup> When planning a replication of a SWAT, thought should be given to the design and approach taken to ensure that the replications are similar. For instance, using similar wording if evaluating a message-based intervention, or consistent timing, so that the results of the meta-analysis are reliable. The same can be said about the definition of outcomes, is a questionnaire considered complete if it has been returned, if all the questions were answered, or if the primary outcome was completed? These differences can impact the results and should be consistent across SWAT replications wherever possible.

### *Meta-analysis*

A key element of accumulating any evidence for trials is to combine similar trials in a meta-analysis. For SWATs, this is particularly important as, unlike a host trial, they are rarely large enough on their own to demonstrate any difference as being statistically significant. Consequently, for individual SWATs we have a high risk of a Type II error, erroneously concluding there is no important difference when there is. To deal with this important issue we have to rely on combining our SWATs. Typically, this is done using a random effects meta-analysis, as we cannot assume that the effect size will be the same across all SWATs - especially when undertaken in various contexts. In this programme of work, we planned to undertake a one-stage meta-analysis, as the individual patient-level data was to be provided to the team, as part of the funding agreement. However, typically it would not be possible to access this level of data, and thus a two-stage approach, where the estimates of effect from each SWAT are combined is most appropriate. We, therefore, would encourage researchers to undertake a cumulative meta-analysis if there are similar published SWATs available - this recommendation is in line with that provided previously.<sup>147</sup>



### *Economics and costings*

Economic evaluations alongside SWATs aim to assess the costs of the strategies that were being evaluated. Despite the very high costs associated with poor recruitment and retention in trials, there has been very little attention paid to the economic aspects of SWATs. It is not enough to simply identify effective recruitment or retention strategies; we need to identify which strategies are also cost-effective.

Within the PROMETHEUS programme, we encouraged host trials to undertake a cost analysis. To do this, we suggested they collect costs both in terms of resources required (for instance the cost of printing cards), and associated staff time (such as preparing the cards for postage). The importance here is to highlight the additional costs that are being acquired by implementing the strategy, as opposed to the standard treatment. This allows a trial team to determine if, for their trial, the cost would be feasible. For instance, within the Christmas card SWAT we determined that the cost per card sent was on average £0.76 – so if you were to use this strategy in a trial of 1000 participants, it would be a direct cost of £760. However, this analysis can then be taken one step further to allow for an assessment of cost per additional participant recruited or retained – where the total associated cost is divided by the additional participants recruited/retained. However, in the Christmas card example, it was not found to be an effective strategy, and as such no additional participants were retained, so even a relatively small finance cost (such as £760), would be considered wasted resources when assessing retention. In some instances, the information from these evaluations from the PROMETHEUS-funded SWATs have helped inform Trial Forge Evidence packs,<sup>138</sup> to allow trial teams to easily determine if the associated costs are worth the potential benefits.

There is additional work that can be done in this area, for instance for a retention SWAT there may be additional cost benefits if the number of reminders sent also is decreased by the strategy, as this may lead to less paper and postage costs. However, to date, this level of in-depth economic evaluation has not been undertaken within SWAT research, and the area is of growing interest.

### *Conclusions*

SWATs are generally straightforward to analyse. We should use an intention-to-treat approach with a pre-specified SAP. Power calculations are generally not required as the sample sizes are constrained by the size of the host trial. The prospect of each individual SWAT being included in a future meta-analysis should always be at the forefront of the researcher's mind when the analysis is undertaken. Consequently, all the numbers of patients randomised should be reported, and the numbers who respond (i.e., for retention SWAT the number who provided the primary outcome)

and the number who did not, and for recruitment, the numbers who were randomised into the host trial.

#### *Lessons learnt for trial teams and methodologists undertaking SWAT research*

- Generally, the analyses are relatively straightforward and are much less complex than the standard analysis for the host trial.
- A SWAT statistical analysis plan should be produced before the analysis as per the standard practice in trial analysis.
- It is not necessary, generally, to undertake a power calculation because each SWAT is constrained by the size of the existing host trial. The lack of statistical power in most SWATs to show the small, but important, differences in retention and recruitment should be addressed by combining SWATs in a meta-analysis. Consequently, researchers are encouraged when writing up their SWAT to include a cumulative meta-analysis of similar SWATs.

#### *Lessons learnt for funders*

- The analysis of SWATs is generally a much simpler procedure compared with the usual clinical trial analysis. Consequently, the impact on funding for statistical input is relatively light.

## Chapter 10: Lessons learnt for reporting SWATs

### *Chapter overview*

Research teams participating in the PROMETHEUS programme have noted challenges in reporting SWATs. These include accessing funding, identifying a journal that will publish a SWAT, and having resources and dedicated time available to analyse and write up a SWAT publication. This chapter outlines these barriers and identifies progress made by the PROMETHEUS programme in mitigating these.

### *The need for transparent reporting of SWATs*

The need for transparent reporting of clinical trials is well known, and the same applies to SWATs. Given that SWATs are designed to be meta-analysed together with similar evaluations of the same strategies, the need for transparent and timely reporting is arguably more crucial, particularly given the limited evidence base around effective recruitment and retention strategies. Without this, the evidence base will not be increased rapidly, and accurate SWAT priorities cannot be recommended. The findings of a SWAT may be detailed within the host trial publication or in a standalone dedicated publication.

### *Reporting SWATs*

Concise, clear, and structured reporting of SWATs in a timely manner is crucial to building the evidence base. Given these methodological studies utilise a randomised controlled trial approach, many trialists may use and adapt the CONSORT guidance when reporting SWATs. However, there are also guidelines, developed by Madurasinghe *et al.*, (2016),<sup>27</sup> primarily for the reporting of embedded recruitment trials, which offer a useful starting point for the structuring of both recruitment and retention SWAT-related publications. Noting the use of both guidelines in the reporting of SWATs, the PROMETHEUS team, in conjunction with Trial Forge, have developed concise SWAT reporting guidelines (*See Appendix 4*). The guidelines developed by the PROMETHEUS team are designed to be methodologically robust and encourage GRADE adherence, which is crucial to aiding evidence synthesis and so deriving definitive evidence. These have recently been piloted by those writing SWAT publications,<sup>151, 152 80, 81, 92</sup> and once feedback has been received the final iteration of the guidance will be submitted for publication.

SWATs may be published within the host trial publication, which offers benefits in obtaining publication in a potentially high-impact journal. There are, however, disadvantages to this approach, particularly where the SWAT is recruitment focussed and the host trial has a long follow-up period, thus resulting in delays to making the SWAT results publicly available. This, therefore, limits the

prompt inclusion of SWAT findings in evidence synthesis, and this may be further exacerbated by the fact that the host trial publication may not note the SWAT in relevant parameters to allow it to be identified in a search strategy focussing specifically on SWAT methodology. Indeed, publishing the SWAT separately means that 'SWAT' can be included either in the title of the paper or the list of keywords to facilitate identification for future systematic reviews and meta-analyses. Furthermore, where SWATs are submitted as standalone publications, these offer an opportunity for early career researchers to obtain first-author publications. It also allows members of the host trial team to obtain peer-reviewed publications before the host trial is completed. For instance, the SCOOP trial of screening for osteoporosis published three SWATs of retention interventions,<sup>136, 153, 154</sup> which were completed and published before the main SCOOP trial results publication. There is, however, a dearth of methodological journals, which would be an ideal resource for the publication of SWATs, and so population or condition specific journals may often be approached. From the experience of PROMETHEUS, these journals often have limited interest and may reduce the subsequent acceptance of SWAT publications.

Reporting platforms such as F1000 Research<sup>155</sup> can therefore be a simpler, and quicker, alternative to the traditional journal. For example, this platform undertakes pre-publication checks on submission, and once approved the publication is made available on the platform with open peer review and revision following accordingly, which means the publication is available promptly. The costs of platform publication are often calculated on the basis of word count and so may also offer an affordable alternative for open-access publication, although it should be noted that since the PROMETHEUS programme commenced costs for platform publications have increased.

Alternatively, a SWAT might be reported in a preprint server whilst under review by a conventional journal.

### *Identifying Peer Reviewers*

Once a journal has accepted the SWAT for review, an additional barrier is the difficulty in finding suitable peer reviewers to review SWAT publications. This has resulted in substantial delays in the submission of SWATs until the journal finds a peer reviewer. This not only increases the length of publication time, but it also slows down advances in the evidence base and could result in teams continuing to undertake strategies which have no effect or a negative effect on their trials.

The small pool of reviewers is due to several reasons. Firstly, some journals have strict criteria for those who are eligible for being a reviewer (for example, the need to hold a PhD). Secondly, SWATs are a niche area and so many SWAT authors have collaborated that conflict of interests are commonplace, further reducing the pool.

Due to the difficulties observed in PROMETHEUS in identifying potential reviewers, a database of potential peer reviewers has been developed. Currently, this is not publicly available due to data protection and GDPR regulations,<sup>156</sup> although it is hoped that this may be available in the future once permission has been sought from individuals.

Many journals require reviewers to list reviewing preferences or classifications which can be beneficial to journal editors in identifying and allocating publications for review. It is therefore useful for journals to include 'SWATs', 'Trials Methodology', 'Embedded Trials', or other similar fields to reviewer classification fields, and for those with an interest in this methodology to ensure that they have included this in their preferences.

### *Peer review feedback*

As noted previously, SWATs are often constrained to the size of the host trial and so formal sample size calculations are often not completed. Often this leads to peer review feedback, specifically noting the need for a sample size calculation and sample size justifications based on power criteria.

This may be as a result of the difficulties in obtaining appropriate peer review (see above) and so the additional use of methodological classifications may ensure that future reviewers are SWAT experienced rather than perhaps just trials experienced. Where reviewers have limited SWAT experience, it may be beneficial to provide additional guidance on the characteristics of a SWAT and what is expected to be reported methodologically, to assist with mitigating unnecessary queries.

### *Conclusions*

As SWATs are designed to be meta-analysed, the need for transparent and timely reporting is necessary to develop the evidence base, find definitive conclusions on effective (or ineffective) strategies, and identify further SWAT priorities.

During the PROMETHEUS programme progress has been made with improving the reporting of SWATs, including guideline development, identification of publication routes, and improving peer review. In addition, the publication platform F1000 has set up a SWAT Collection,<sup>155</sup> and Research Methods in Medicine & Health Sciences<sup>157</sup> is currently developing a SWAT special edition. From these developments, the following key learning points are suggested for relevant stakeholders:

### *Lessons learnt for trial teams and methodologists undertaking SWAT research*

- SWAT-specific reporting guidelines should be used to ensure that sufficient detail is provided to support GRADE evaluation of SWAT interventions.

- Trialists are encouraged to include SWAT or methodology classifications in their journal reviewer profiles if they have an interest in this methodology and are willing to provide reviews.

*Lessons learnt for journals and reviewers*

- When selecting reviewers, SWAT and methodology classifications should be used where possible.
- Journals should consider potential flexibility to reviewer credentials for niche areas such as SWATs; for example, a PhD can be substituted with relevant experience in the area.

## Chapter 11: Recommendations for future research practice, direction, and support

### *Chapter overview*

Below we outline a range of recommendations for stakeholders such as funders, trial teams who are planning and undertaking recruitment and retention activities and for trial methodologists interested in SWATs.

### *Recommendations*

#### *Recommendations for funders*

- All trial funders should contribute to the effort to improve the efficiency of trials. Funders should encourage the teams that they fund to undertake SWATs
- Funding streams specifically designed to support SWATs must be made available to trial teams to continue building the trial process evidence base, for recruitment and retention as well as for other stages of the trial design and delivery process. This includes funding streams for undertaking specific SWATs, as well as infrastructure funding to support CTUs and other centres to undertake coordination activities that will support the design, conduct, reporting, and implementation of SWATs and their findings to inform the work of the NIHR, the MRC, and other funders
- The PROMETHEUS Programme has demonstrated that coordination of activity remains crucial to the delivery of SWATs. A central, national coordination point that provides hands-on support needs to continue and funding should be allocated for this. Additionally, CTUs should identify a lead for SWATs to support SWAT activities and evidence-based trial conduct within the CTU, as well as a link with others undertaking SWATs elsewhere to share best practice. The funding for both central and CTU based support should be ongoing, in the same way as commissioned NIHR research and developing NICE guidance
- SWAT priorities need to be identified and communicated clearly to funders, and funders should use these priorities to inform their funding decisions
- Funders should develop a mechanism to promote SWAT questions that have been identified as a priority during the application process as a strategy to increase the evidence base for trial conduct using SWATs
- The mean cost of funding requested for a standalone SWAT within PROMETHEUS was £3,535 (range £500 to £5,000). The coordinated SWATs cost £10,668 (training SWAT), and £1,306.40 (Christmas cards); however, these did not include costs for central coordination,

data preparation and sharing by the host trial teams, data cleaning, analysis, and write-up. These costs suggest that the £10,000 being offered by the NIHR for trial teams to include a SWAT should be sufficient for most planned SWATs. However, there may be occasions where trial teams may wish to test strategies that may be more expensive

- When applying for funding, trial teams should be asked to indicate whether the question they are addressing is a priority SWAT question, and to provide a clear rationale for selecting that particular question
- If teams are unable to undertake a SWAT, funders should ask that recruitment and retention methods are clearly reported to support the evidence base

#### *Recommendations for Sponsors*

- Whilst most Sponsors did not raise any issues or difficulties with SWATs being undertaken within their trials, our experience suggests that there is a need for clear, easily accessible information about the nature of SWATs, as well as the role of the funder in supporting SWATs
- Any future changes proposed by the HRA to the approvals process need to be communicated clearly and applied consistently to each SWAT

#### *Recommendations for involving PPI partners in SWAT research*

- PPI should be considered when undertaking a SWAT, in the same way PPI work is expected to be undertaken in the main trial
- PPI partners should be involved to develop novel and untested recruitment and retention strategies, as well as adapting existing strategies to the context of their specific host trial and the population being enrolled
- PPI work can extend to professional stakeholders such as surgeons or research nurses who are intrinsic to site recruitment

#### *Recommendations for oversight committees*

- Our experience suggests that Trial Management Groups (TMGs) play a key role in decisions about whether a SWAT is undertaken and continued in the host trial or not. TMG members should encourage the uptake of SWATs in their trials. Whilst the findings of SWATs may not always directly inform their host trial, the findings of SWATs undertaken during the early phase of the trial (such as during the pilot phase), may inform the decisions about which strategies should be used later, such as in the main trial
- Trial Steering Committees should review the SWAT activity and progress, in the same way that they review sub-studies in a trial



- Data Monitoring Committee review is dependent on the specific host trial and SWAT strategy being evaluated

#### *Recommendations for Journals and Reviewers*

- Journal peer reviewer profiles should be updated to include methodological interests and expertise, to support evidence of suitability to undertake a peer review for a SWAT publication
- When selecting peer reviewers, the SWAT and methodology interests as registered by reviewers should be used where possible
- SWATs are a niche area and so to increase the pool of reviewers, journals should consider being more flexible when assessing reviewer credentials to review a SWAT, such as allowing relevant experience in place of a PhD
- Reviewers should be advised that informed consent from participants is often not obtained when undertaking a SWAT, and that it is not necessary to query this with authors
- Robust and transparent reporting is necessary, that is compliant with Consolidated Standards of Reporting Trials (CONSORT)

#### *Recommendations for trial teams and methodologists undertaking SWAT research*

- There remains a need for continually updated research priorities to allow researchers to address the questions relevant at that time
- When SWAT priorities are set, methodologists need to provide as much information as possible to enable teams to make informed decisions about evaluating the priorities set.
- Further work is needed to help teams identify suitable SWAT strategies for their host trials
- SWAT priorities need to be communicated clearly and consistently to trial teams

#### *Recommendation for trial conduct and using SWAT evidence*

- As the evidence base develops for effective and cost-effective recruitment and retention strategies, it will become increasingly important for trial teams to use this evidence base to inform their recruitment and retention activities. Trial teams need to actively engage with the evidence base to inform their practice, including using evidence from systematic reviews, and from web-based resources such as Trial Forge.<sup>107</sup> Funders will need to actively support the trials they fund to use evidence-informed recruitment and retention strategies

#### *Recommendations for future research*

- There remains a substantial need for more high-quality SWAT evidence and so Chief Investigators should be encouraged to consider the embedding of a SWAT at the funding

stage. Further work is therefore needed to increase the awareness of the methodological importance of SWAT research with research teams and to develop engagement strategies to increase SWAT activity

- Future research needs to focus on identifying whether further replications are needed for existing evidence. If so, the gaps in the evidence base should be targeted. More coordination and replication of SWAT evaluations are encouraged
- A 'real-time' and dynamic communication strategy including a clear cost and resource breakdown for each suggested SWAT should be developed. This will alleviate the burden to trials teams to begin costing exercises and enables them to make an informed decision more quickly as to whether they can embed a given SWAT
- There is a need to aid teams to identify and select a suitable SWAT for their host trial populations. Pragmatic decisions on which SWAT may be appropriate and feasible to include should be taken as required. A mechanism to communicate SWAT research priorities is needed, and this information needs to be readily accessible for all trialists to refer to
- Our findings demonstrate that within an individual host trial, there is often a capacity to address more than one SWAT question, either separately, or simultaneously using a factorial design. This suggests that there is capacity to significantly speed up and strengthen the evidence base through teams undertaking more than one SWAT in their trials where relevant
- For certain strategies, coordinated SWATs should be encouraged. This method could be used to rapidly replicate SWAT evaluations to plug the evidence gap, as well as to evaluate more complex recruitment and retention strategies that may be more challenging to undertake using individual SWATs. Materials should be developed to advise teams on how to undertake coordinated SWATs, as well as a method of networking to enable teams to promote their simultaneous SWAT and collaborate
- As the evidence base develops, it will become increasingly important for trialists to utilise the evidence base in a systematic way to identify both effective and ineffective strategies to inform their practice. Future work should therefore consider issues around the dissemination and implementation of SWATs and develop guidance to enable the wider trials community to undertake, report and adopt the findings of SWATs. Implementation science, the study of methods to promote the uptake of evidence-based practice, could be used to inform any such future work. Funders can also help by questioning strategies proposed by trial teams that are known to be either ineffective or not cost-effective

- Improving the knowledge of the potential 'harms' from implementing interventions that have no evidence of benefit is an important next step to help improve uptake
- Whilst establishing the effectiveness of recruitment or retention strategies is important, the high costs of research waste and limited public finance means that cost considerations around SWATs are just as important. With only one retention strategy having high GRADE certainty of cost-effectiveness, we encourage trial teams to undertake streamlined economic evaluations alongside all their SWATs in the future, for strategies shown to be effective, as well as those that are ineffective. For cost-effectiveness, trial teams should look to report the cost per additional participant recruited or retained (that is, the incremental cost-effectiveness ratio). Value of information analyses can help determine the need for further SWAT evidence where several SWATs already exist
- Many trial teams wish to contribute to developing the evidence base by undertaking non-randomised SWATs. Future work that informs the development of guidance for undertaking non-randomised SWATs would be helpful
- There has been limited assessment currently on the views of participants on individual SWAT interventions. Future work using process evaluations to better understand the acceptability of and impact of SWATs on participants would therefore be useful to better understanding the effectiveness of SWATs
- Working with trial teams to develop engagement strategies and training to undertake SWATs would be beneficial. Audience-specific guidance should be developed to support SWAT research. We suggest building on the successfully executed and attended 'PROMETHEUS hosted webinars' as well as undertaking research with teams to identify what training or support they require
- Trial teams have expressed they want to undertake SWATs that are important and necessary to increase the evidence base. Collaboration with funders, working groups involved in priority setting, and trial teams are needed to develop a mechanism to communicate this dynamic and evolving information once priority SWATs have been identified
- Work is needed to identify the barriers that teams have when undertaking a SWAT, and strategies and solutions for addressing these barriers should be identified and implemented
- Continued and proactive collaboration is needed with working groups to enable networking, and collaboration with teams undertaking SWAT research
- Reporting guidance is needed to support teams when writing publications to ensure sufficient information is included so GRADE evaluation can be undertaken.

## Chapter 12: Overall conclusions

In this manuscript, we have reported the design, methodology, and results of the PROMETHEUS programme. We have used the lessons learnt from undertaking PROMETHEUS to highlight key practical lessons for a range of stakeholders in planning, designing, undertaking, and reporting SWATs. We have used our experience from PROMETHEUS to make a range of recommendations for stakeholders. Given the PROMETHEUS programme was the first of its kind, this enabled a range of observations on the coordination and conduct of SWATs to be identified which have been presented here.

PROMETHEUS originally aimed to fund 25 SWATs and supported 42 SWATs, generating the single biggest body of SWAT activity in the world. When SWAT funding was made available, we found that many teams embedded SWATs into their research. Having a central point of contact that coordinated SWAT activity alongside providing funding has been key in determining the success of PROMETHEUS and will continue to be so to increase the SWAT evidence base. We have used our experience of undertaking PROMETHEUS to provide practical guidance and examples for undertaking SWATs.

The PROMETHEUS SWATs reported to date found that there was no evidence of a significant difference in recruitment rates in any of the strategies tested. For the retention strategies, we found evidence that pre-notification of trial participants by sending a card may result in a slight increase in attendance at a face-to-face primary outcome measurement visit; and that sending a pre-notification letter or email before sending a self-report questionnaire increases response rates, compared with no pre-notification. We also found that participants randomised to receive a personalised text message were more likely to return their initial postal questionnaire than those who received a non-personalised text message, and that men aged under 65 years were the group most likely to return their initial questionnaire if they received a personalised text message. Another SWAT found that when comparing personalised text messages versus non-personalised text messages, there was evidence that a personalised text would result in fewer telephone follow-ups that were more resource intensive. Combined results from PROMETHEUS together with other published SWATs found that including a pen probably increases retention and response rate, compared to not sending a pen with postal questionnaires. One SWAT showed evidence of an adverse impact on retention when printing the trial primary outcome on pink paper compared to printing the primary outcome on white paper; this showed a decreased response and was also more burdensome to collect postal data as it increased the need for reminders. There was no evidence of

a significant difference in retention rates in any of the other strategies tested. More replications of these SWATs are required.

The PROMETHEUS programme has substantially increased the number of high-quality SWATs undertaken and has identified a range of areas where further development is warranted. The conduct of 42 SWATs during this time enabled the identification of key aspects of undertaking SWAT that require further development and coordination. This is essential to ensure that SWATs continue to be undertaken with sufficient replications available to enable meta-analyses to be performed to provide conclusive findings. This will lead to an increase in the evidence base, enabling the identification of effective, ineffective, or harmful recruitment and retention strategies to support trialists to design and undertake efficient research.

There is a need to develop a strategy to aid teams to identify a suitable SWAT for their host trial populations and a mechanism to communicate SWAT research priorities. Ongoing work is needed to increase the awareness of the methodological importance of SWAT research with research teams and develop engagement strategies to increase SWAT activity. Continued collaboration with the HRA is also necessary to refine the SWAT approvals process.

The mean cost of each SWAT was £3,535. In addition to funding SWATs, PROMETHEUS successfully demonstrated the methodological feasibility of undertaking coordinated SWATs, a powerful new tool with the potential to rapidly accelerate the evidence base for recruiting and retaining trial participants. In addition to the financial support, the coordination PROMETHEUS provided was crucial to increasing the recruitment and retention evidence base.

Following the initial success of PROMETHEUS, additional funding was received from the NIHR (award ID: NIHR132547) to extend PROMETHEUS and continue work to investigate how teams can be better supported to implement recruitment and retention SWATs. This allowed us to develop the 'Trial Forge SWAT Network', currently consisting of 30 CTUs and trials centres in the UK, the Republic of Ireland, Iran, and Australia. The Trial Forge SWAT Network will provide ongoing networking and dissemination opportunities. The International Trial Forge SWAT Network has been well received by teams and will offer opportunities to share, disseminate, and increase methodological research. We found webinars to be an effective tool to communicate methodological research and propose that these continue to be utilised in the future.

We have used our experience of undertaking PROMETHEUS to provide practical guidance and examples for undertaking SWATs, as well as make recommendations for all the key stakeholders involved in the conduct and delivery of SWATs. Collectively, these developments should enable and

encourage teams to undertake SWATs, to further develop the evidence base, and ultimately, to prevent research waste by improving trial recruitment and retention.

## Acknowledgements

Firstly, we thank all the PROMETHEUS host trial teams and clinicians that recruited and followed up participants for this programme, as well as the participants in all the host trials for making this research possible.

We thank York Trials Unit (YTU) - an NIHR supported UK Clinical Research Collaboration-registered CTU - for hosting this methodological work. Our particular thanks to Sally Baker and Emma-Louise Rose Brooks at YTU for their support in managing the administrative aspects of the PROMETHEUS programme.

We are grateful to the Health Services Research Unit (HSRU), core funded by the Chief Scientist Office of the Scottish Government Health and Social Care Directorate, for their support, especially with the initiation of the Christmas card coordinated SWAT.

We would also like to thank the Health Research Board – Trials Methodology Research Network (HRB-TMRN) in the Republic of Ireland for kindly sharing materials, such as the ‘peer review assessment form’, which we adapted for PROMETHEUS.

We extend our thanks to The University of Bristol’s team (Qualitative research integrated within Trials) QuinteT, who developed the recruitment training course that we tested as a coordinated SWAT. The training was funded by the MRC Network of Hubs for Trials Methodology Research (MR/L004933/1- R53) and supported by the MRC ConDuCT-II Hub (Collaboration and innovation for Difficult and Complex randomized controlled Trials In Invasive procedures - MR/K025643/1).

We thank Professor Paul Brocklehurst, who was a co-applicant on the original MRC funded application, for his contribution to the PROMETHEUS programme.

We thank the PROMETHEUS PPI members: Christine Allmark, Phillip Bell, Ailsa Donnelly, Judith Hogg, and Kay Walker. Additionally, we would like to thank the HSRU PPI Group who contributed to the Christmas card design for the Christmas card coordinated SWAT, as well as the PPI groups of all the host trials for their feedback and contributions on the SWATs.

### *Contributions of authors*

Adwoa Parker (Assistant Professor), contributed to study conceptualisation, funding acquisition, study methodology and investigation, project administration, drafted the initial version of the manuscript, and read and approved the final version.

Catherine Arundel (Research Fellow) contributed to study conceptualisation, study methodology and investigation, project administration, drafted the initial version of the manuscript, and read and approved the final version.

Laura Clark (Research Fellow) contributed to study investigation, project administration, drafted the initial version of the manuscript, and read and approved the final version.

Elizabeth Coleman (Statistician) contributed to study methodology and investigation, curated the data, carried out analysis and prepared the results for publication, drafted the initial version of the manuscript, and read and approved the final version.

Laura Doherty (Trial Support Officer) contributed to study investigation, project administration, drafted the initial version of the manuscript, and read and approved the final version.

Catherine Hewitt (Professor) contributed to study conceptualisation, funding acquisition, study methodology and investigation, analysis, and preparation of the results, drafting of the manuscript, and read and approved the final version of the manuscript.

David Beard (Professor) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

Peter Bower (Professor) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

Cindy Cooper (Professor) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

Lucy Culliford (Senior Research Fellow) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

Declan Devane (Professor) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

Richard Emsley (Professor) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

Sandra Eldridge (Professor) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

Sandra Galvin (Programme Manager) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

Katie Gillies (Reader) contributed to study methodology and investigation, and read and approved the final version of the manuscript.

Alan Montgomery (Professor) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

Chris Sutton (Senior Lecturer in Clinical Trial Statistics) contributed to study methodology and investigation, and read and approved the final version of the manuscript.

Shaun Treweek (Professor) contributed to study conceptualisation, funding acquisition, study methodology and investigation, and read and approved the final version of the manuscript.

David Torgerson (Professor) conceptualised the study, led on funding acquisition, contributed to study methodology, investigation, drafting the manuscript, provided overall supervision of the programme, and read and approved the final version.

### *Study registrations*

All SWATs were registered with the Northern Ireland Network for Trials Methodology Research SWAT Repository.

### *Funding details*

This project was funded by the Medical Research Council (MR/R013748/1) and National Institute for Health Research as part of the MRC–NIHR Methodology Research programme.

### *Ethics Statement*

Ethical approvals for all of the PROMETHEUS host trials were obtained separately by the host trial teams, and the SWATs undertaken were approved as part of the initial ethical application for the host trial, or through the submission of an amendment to the Research Ethics Committee. The following approvals for the SWATs were obtained:

- ACTIVE: NRES Committee Yorkshire and The Humber – Bradford Leeds (ref: 18/YH/0014)
- ActWELL: East of Scotland Research Ethics Service REC 1 (ref: 17/ES/0073)
- CHAMP-1: South West - Frenchay Research Ethics Committee (ref: 19/SW/0082)
- CLEAR: North East - Tyne & Wear South Research Ethics Committee, (ref: 17/NE/0339)



- GRASP: Berkshire B Research Ethics Committee, substantive amendment to the host trial (Ref. 16/SC/0508)
- IBD-BOOST: London - Surrey Research Ethics Committee (ref: 19/LO/0750)
- KReBs (pen SWAT): North East – Newcastle and North Tyneside on (Ref 16/NE/0400, Amendment 2)
- KReBs (text message SWAT): North East – Newcastle and North Tyneside on (Ref 16/NE/0400), amendment Number 16/NE/0400/AM14)
- L1FE: London - Harrow Research Ethics Committee (ref: 19/LO/0555)
- MiQuit: East Midlands–Nottingham 1 Research Ethics Committee (ref 13/EM/0427 and 17/EM/0327)
- MSS3: North West – Greater Manchester Central Research Ethics Committee (ref: 18/NW/0422)
- OTIS: NHS West of Scotland Research Ethics Committee 3 (ref: 16/WS/0154)
- PEP-TALK: South Central Oxford B – (Ref: 18/SC/0423)
- PONOC: National Research Ethics Service Committee East Midlands—Nottingham 2 (ref: 13/EM/0459)
- SARC: West of Scotland REC 1 (ref: 19/WS/0087)
- SSHeW: Department of Health Sciences Research Ethics Committee at the University of York and the Health Research Authority (Ref: HSRGC/2016/187/A)
- SWHSI-2: Leeds East Research Ethics Committee — reference 19-YH-0054
- TOPaZ: East of Scotland Research Ethics Service (EoSRES) (ref: 16/ES/0110)
- UK-FROST: North East - Newcastle & North Tyneside 2 Ethics Committee, (Ref: 14/NE/1176). Substantial Amendment 1 – REC Favourable Opinion 02/02/2015.
- VITA: London – Harrow Research Ethics Committee on 9 September 2017 (reference 17/LO/1245)
- WORKWELL: West Midlands – Solihull Research Ethics Committee (ref: 18/WM/0327)

The ethical approval for the coordinated recruitment training SWAT across all the host trials was obtained from the University of York’s Department of Health Sciences Research Governance Committee. No further approvals were needed as this was classed as an evaluation of a teaching workshop (Ref 76/76).

The ethical approval for the coordinated Christmas card SWAT across all the host trials was approved by the Yorkshire and the Humber – Leeds West Research Ethics Committee (ref: 19/YH/0349). A single application was submitted to allow the SWAT to be embedded within all the host trials.

### *Data Sharing Statement*

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

### *Additional funding received*

YTU receives core funding from the NIHR CTU Support Funding. Following the initial success of PROMETHEUS, YTU received additional funding to continue some of the work of PROMETHEUS, through an institutional award from the NIHR CTU Infrastructure Support (NIHR132547). The Health Services Research Unit (HSRU) is core funded by the Chief Scientist Office of the Scottish Government Health and Social Care Directorate.

Each host trial received its own funding, as follows:

- Cancer Research UK, the Chief Scientist Office, Scottish Government, Health and Social Care Public Health Agency Northern Ireland, Chest Heart and Stroke Northern Ireland, The Lullaby Trust, Public Health Agency Northern Ireland, and the Scottish Cot Death Trust: CPIT-III
- Medical Research Council and National Institute for Health and Care Research - Efficacy and Mechanism Evaluation: STARTS:REACTS, TOPaZ
- Cancer Research UK and National Institute for Health Research: MiQuit
- National Institute for Health and care Research - Health Technology Assessment Programme: ARTISAN, C-Gall, CLEAR, DISC, FUTURE, GRASP, L1FE, MAGIC, ProFHER-2, PUrE, REFLECT and SWHSI-2, UK FROST, VITA
- National Institute for Health and care Research - Programme Grants for Applied Research: IBD-BOOST
- National Institute for Health Research - Health and Social Care Delivery Research: MSS3
- National Institute for Health Research - Public Health Research Programme: SSHeW
- The Scottish Government: ActWELL
- Versus Arthritis: WORKWELL

Author KG's time was funded by the Chief Scientist Office of the Scottish Government's Health and Social Care Directorate (CZU/3/3). Author RE's time was funded by NIHR Research Professorship (NIHR300051) and the NIHR Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. Author APs time was part-funded by the Wellcome Trust [ref: 204829] through the Centre for Future Health (CFH) at the University of York, which partially supported the delivery of the recruitment training SWAT.

## Publications

1. Coleman E, Arundel C, Clark L, et al. Bah humbug! Association between sending Christmas cards to trial participants and trial retention: randomised study within a trial conducted simultaneously across eight host trials. *BMJ (Clinical research ed)* 2021; 375: e067742. 2021/12/16. <https://doi.org/10.1136/bmj-2021-067742> .
2. Parker A, Mills N, Rooshenas L, et al. Staff training to improve participant recruitment into surgical randomised controlled trials: a feasibility study embedded within four randomised controlled trials. In: *International Clinical Trials Methodology Conference Brighton, 2019*, p.PS8A. BMC.
3. Arundel CE, Clark L, Coleman E, et al. Challenges and Solutions to the Implementation of Studies Within A Trial (SWATs): The Experiences of the PROMETHEUS Programme. *Research in Health Sciences & Medicine*. 2022;0(0). doi:10.1177/26320843221106949
4. Parker A, Arundel C, Mills N, et al. Staff training to improve participant recruitment into surgical randomised controlled trials: a feasibility study embedded within four randomised controlled trials. *Research Methods in Medicine & Health Sciences*. 2022;0(0). <https://doi.org/10.1177/26320843221106950>
5. Doherty L, Parker A, Arundel C, et al. The PROMoting THE USE of SWATs (PROMETHEUS) Programme : lessons learnt and future developments for SWATs. *Research Methods in Medicine & Health Sciences*. Submitted.
6. Clark L, Arundel C, Coleman E, et al. The PROMoting THE USE of SWATs (PROMETHEUS) Programme : lessons learnt and future developments for SWATs. *Research Methods in Medicine & Health Sciences*. 2022;3(4):100-106. <https://doi.org/10.1177/26320843221089632>
7. Treweek S, Gallant S, Anderson AS. SWAT 76 evaluation: randomised evaluation of sending pre-notification cards to trial participants before a face-to-face primary outcome measurement to increase attendance. *F1000Research*. 2021;10. <https://doi.org/10.12688/f1000research.50890.2>
8. Cureton L, Marian IR, Barber VS, Parker A, Torgerson DJ, Hopewell S. Randomised study within a trial (SWAT) to evaluate personalised versus standard text message prompts for increasing trial participant response to postal questionnaires (PROMPTS). *Trials*. 2021 Dec;22(1):1-0. <https://doi.org/10.21203/rs.3.rs-262194/v1>
9. Mitchell AS, Cook L, Dean A, Fairhurst C, Northgraves M, Torgerson DJ, Reed M. Using pens as an incentive for questionnaire return in an orthopaedic trial: an embedded randomised

- controlled retention trial. *F1000Research*. 2021 Sep 7;9:321.  
<https://doi.org/10.12688/f1000research.23018.2>
10. Mitchell AS, Cook L, Dean A, Fairhurst C, Northgraves M, Torgerson DJ, Reed M. An embedded randomised controlled retention trial of personalised text messages compared to non-personalised text messages in an orthopaedic setting. *F1000Research*. 2020;9.  
<https://doi.org/10.12688/f1000research.24244.1>
  11. Coleman E, Whitmore R, Clark LK, Daykin K, Clark M. Pre-notification and personalisation of text-messages to retain participants in a smoking cessation pregnancy RCT: an embedded randomised factorial trial. *F1000Research*. 2021 Jul 22.  
<https://doi.org/10.12688/f1000research.51964.1>
  12. James S, Parker A, Cockayne S, Rodgers S, Fairhurst C, Torgerson DJ, Rhodes S, Cotterill S. Including a pen and/or cover letter, containing social incentive text, had no effect on questionnaire response rate: a factorial randomised controlled study within a trial. *F1000Research*. 2020;9. <https://doi.org/10.12688/f1000research.23767.2>
  13. Cunningham-Burley R, Roche J, Fairhurst C, Cockayne S, Hewitt C, Iles-Smith H, Torgerson DJ, SSHeW Study Team. Enclosing a pen to improve response rate to postal questionnaire: an embedded randomised controlled trial. *F1000Research*. 2020;9.  
<https://doi.org/10.12688/f1000research.23651.1>
  14. Sarathy PP, Kottam L, Parker A, Brealey S, Coleman E, Keding A, Mitchell A, Northgraves M, Torgerson D, Rangan A. Timing of electronic reminders did not improve trial participant questionnaire response: a randomized trial and meta-analyses. *Journal of clinical Epidemiology*. 2020 Jun 1;122:70-7. <https://doi.org/10.1016/j.jclinepi.2020.03.001>

## References

1. Sackett DL, Rosenberg WM, Gray JM, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. In: British Medical Journal Publishing Group; 1996: 71-2.
2. ClinicalTrials.gov. Trends, Charts, and Maps. 2022. URL: <https://clinicaltrials.gov/ct2/resources/trends> (accessed 27/05/2022, 2022).
3. National Institute for Health and Care Research. Annual Report 2019/2020. National Institute for Health and Social Care Research; 2021. URL: [https://www.nihr.ac.uk/documents/about-us/our-contribution-to-research/research-performance/NIHR\\_Annual\\_Report\\_19\\_20.pdf](https://www.nihr.ac.uk/documents/about-us/our-contribution-to-research/research-performance/NIHR_Annual_Report_19_20.pdf) (accessed 27/05/2022, 2022).
4. National Institute for Health and Care Research. Annual Report 2020/2021. National Institute for Health and Social Care Research; 2022. URL: [https://www.nihr.ac.uk/documents/about-us/our-contribution-to-research/research-performance/Final-NIHR-Annual-report%2020\\_21.pdf](https://www.nihr.ac.uk/documents/about-us/our-contribution-to-research/research-performance/Final-NIHR-Annual-report%2020_21.pdf) (accessed 27/05/2022, 2022).
5. Jacques RM, Ahmed R, Harper J, Ranjan A, Saeed I, Simpson RM, et al. Recruitment, consent and retention of participants in randomised controlled trials: a review of trials published in the National Institute for Health Research (NIHR) Journals Library (1997–2020). *BMJ Open* 2022;12:e059230. <https://doi.org/10.1136/bmjopen-2021-059230>
6. Kitterman DR, Cheng SK, Dilts DM, Orwoll ES. The prevalence and economic impact of low-enrolling clinical studies at an academic medical center. *Academic medicine : journal of the Association of American Medical Colleges* 2011;86:1360-6. <https://doi.org/10.1097/ACM.0b013e3182306440>
7. Salman RA-S, Beller E, Kagan J, Hemminki E, Phillips RS, Savulescu J, et al. Increasing value and reducing waste in biomedical research regulation and management. *The Lancet* 2014;383:176-85.
8. Moher D, Glasziou P, Chalmers I, Nasser M, Bossuyt PM, Korevaar DA, et al. Increasing value and reducing waste in biomedical research: who's listening? *Lancet (London, England)* 2016;387:1573-86. [https://doi.org/10.1016/s0140-6736\(15\)00307-4](https://doi.org/10.1016/s0140-6736(15)00307-4)
9. Knowlson C, Torgerson DJ. Effects of rapid recruitment and dissemination on Covid-19 mortality: the RECOVERY trial. *F1000Research* 2020;9.

10. Treweek S, Gillies K. Trial retention: it's time for Cinderella to go to the ball. In: BioMed Central; 2017.
11. Walsh M, Devereaux PJ, Sackett DL. Clinician trialist rounds: 28. When RCT participants are lost to follow-up. Part 1: Why even a few can matter. Clin Trials 2015;12:537-9.  
<https://doi.org/10.1177/1740774515597702>
12. Smith CT, Hickey H, Clarke M, Blazeby J, Williamson P. The trials methodological research agenda: results from a priority setting exercise. Trials 2014;15:1-7.
13. Clarke M, Savage G, Maguire L, McAneney H. The SWAT (study within a trial) programme; embedding trials to improve the methodological design and conduct of future research. Trials 2015;16:1-.
14. Anonymous. Education section – Studies Within A Trial (SWAT). Journal of Evidence-Based Medicine 2012;5:44-5. <https://doi.org/https://doi.org/10.1111/j.1756-5391.2012.01169.x>
15. Treweek S, Bevan S, Bower P, Campbell M, Christie J, Clarke M, et al. Trial forge guidance 1: what is a study within a trial (SWAT)? Trials 2018;19:139.
16. Dwyer CP, Moses A, Rogers FM, Casey D, Joyce R, Hynes SM. A qualitative investigation of reasoning behind decisions to decline participation in a research intervention: A study-within-a-trial. Journal of health psychology 2021; 10.1177/13591053211037736:13591053211037736.  
<https://doi.org/10.1177/13591053211037736>
17. Allan S, Mcleod H, Bradstreet S, Bell I, Whitehill H, Wilson-Kay A, et al. Perspectives of Trial Staff on the Barriers to Recruitment in a Digital Intervention for Psychosis and How to Work Around Them: Qualitative Study Within a Trial. JMIR Human Factors 2021;8:e24055.
18. Racine E, Hurley C, Cheung A, Sinnott C, Matvienko-Sikar K, Baumgartner C, et al. Participants' perspectives and preferences on clinical trial result dissemination: The TRUST Thyroid Trial experience. HRB open research 2018;1:14-. <https://doi.org/10.12688/hrbopenres.12817.2>
19. Clarke M, Maguire L. SWAT 9: Pre-randomisation matching of sites in a cluster randomised trial. In. SWAT Store: Northern Ireland Network for Trials Methodology Research; 2015.
20. Ahmed S, Airlie J, Clegg A, Copey B, Cundill B, Forster A, et al. A new opportunity for enhancing trial efficiency: Can we investigate intervention implementation processes within trials using SWAT (study within a trial) methodology? Research Methods in Medicine & Health Sciences 2022:26320843221080734.

21. Hurley C. SWAT 38: Cost implications of conducting a risk assessment prior to developing a monitoring plan for a multicentre clinical trial. In. 10/01/2016 ed. SWAT Store: Northern Ireland Network for Trials Methodology Research; 2016.
22. Fairhurst C, Roche J, Bissell L, Hewitt C, Hugill-Jones J, Howsam J, et al. A 2x2 randomised factorial SWAT of the use of a pen and small, financial incentive to improve recruitment in a randomised controlled trial of yoga for older adults with multimorbidity [version 2; peer review: 2 approved]. *F1000Research* 2022;10:326. <https://doi.org/10.12688/f1000research.52164.2>
23. Treweek S, Pitkethly M, Cook J, Fraser C, Mitchell E, Sullivan F, et al. Strategies to improve recruitment to randomised trials. *Cochrane Database of Systematic Reviews* 2018;2:MR000013. <https://doi.org/10.1002/14651858.MR000013.pub6>
24. Gillies K, Kearney A, Keenan C, Treweek S, Hudson J, Brueton VC, et al. Strategies to improve retention in randomised trials. *Cochrane Database of Systematic Reviews* 2021;3:MR000032. <https://doi.org/10.1002/14651858.MR000032.pub3>
25. Rick J, Graffy J, Knapp P, Small N, Collier DJ, Eldridge S, et al. Systematic techniques for assisting recruitment to trials (START): study protocol for embedded, randomized controlled trials. *Trials* 2014;15:407. <https://doi.org/10.1186/1745-6215-15-407>
26. Madurasinghe VW, Bower P, Eldridge S, Collier D, Graffy J, Treweek S, et al. Can we achieve better recruitment by providing better information? Meta-analysis of 'studies within a trial' (SWATs) of optimised participant information sheets. *BMC Medicine* 2021;19:218. <https://doi.org/10.1186/s12916-021-02086-2>
27. Madurasinghe VW, Sandra Eldridge on behalf of MRCSG, Gordon Forbes on behalf of the SECG. Guidelines for reporting embedded recruitment trials. *Trials* 2016;17:27. <https://doi.org/10.1186/s13063-015-1126-y>
28. Health Research Board – Trials Methodology Research Network. STUDY WITHIN A TRIAL (SWAT). Health Research Board – Trials Methodology Research Network; 2022. URL: <https://www.hrb-tmrn.ie/research-and-innovation/funding-opportunities/studies-within-a-trial-swats/> (accessed 27/05/2022, 2022).
29. National Institute for Health and Care Research. Studies within a trial (SWAT) and studies within a review (SWAR). 2022. URL: <https://www.nihr.ac.uk/documents/studies-within-a-trial-swat/21512> (accessed 27/05, 2022).

30. Martin-Kerry JM, Bower P, Young B, Graffy J, Sheridan R, Watt IS, et al. Developing and evaluating multimedia information resources to improve engagement of children, adolescents and their parents with trials (TRECA study). Study Protocol for a series of linked randomised controlled trials 2017;18:265. <https://doi.org/10.1186/s13063-017-1962-z>
31. Brueton VC, Tierney J, Stenning S, Harding S, Meredith S, Nazareth I, et al. Strategies to improve retention in randomised trials. Cochrane Database of Systematic Reviews 2013; 10.1002/14651858.MR000032.pub2. <https://doi.org/10.1002/14651858.MR000032.pub2>
32. Northern Ireland Network for Trials Methodology Research. SWAT Store. In. Northern Ireland Network for Trials Methodology Research; 2022.
33. Bower P, Brueton V, Gamble C, Treweek S, Smith CT, Young B, et al. Interventions to improve recruitment and retention in clinical trials: a survey and workshop to assess current practice and future priorities. Trials 2014;15:399-. <https://doi.org/10.1186/1745-6215-15-399>
34. Healy P, Galvin S, Williamson PR, Treweek S, Whiting C, Maeso B, et al. Identifying trial recruitment uncertainties using a James Lind Alliance Priority Setting Partnership – the PRioRiTty (Prioritising Recruitment in Randomised Trials) study. Trials 2018;19:147. <https://doi.org/10.1186/s13063-018-2544-4>
35. National Institute for Health and Care Research. CRN Portfolio. National Institute for Health and Care Research,; 2022. URL: <https://www.nihr.ac.uk/researchers/collaborations-services-and-support-for-your-research/run-your-study/crn-portfolio.htm> (accessed 28/05/2022, 2022).
36. Adamson J, Hewitt CE, Torgerson DJ. Producing better evidence on how to improve randomised controlled trials. BMJ : British Medical Journal 2015;351:h4923. <https://doi.org/10.1136/bmj.h4923>
37. Cook TD, Campbell DT, Day A. Quasi-experimentation: Design & analysis issues for field settings: Houghton Mifflin Boston; 1979.
38. Medical Research Council. MRC Policy and Guidance on Sharing of Research Data from Population and Patient Studies: Medical Research Council; 2017.
39. Kearney A, Harman NL, Rosala-Hallas A, Beecher C, Blazeby JM, Bower P, et al. Development of an online resource for recruitment research in clinical trials to organise and map current literature. Clin Trials 2018;15:533-42. <https://doi.org/10.1177/1740774518796156>



40. Kearney A, Ashford PA, Butlin L, Conway T, Cragg WJ, Devane D, et al. Developing an online, searchable database to systematically map and organise current literature on retention research (ORRCA2). *Clin Trials* 2022;19:71-80. <https://doi.org/10.1177/17407745211053803>
41. Stewart D, van Dongen A, Watson M, Mandefield L, Atkin K, Dhital R, et al. A pilot cluster randomised trial of the Medicines and Alcohol Consultation (MAC): an intervention to discuss alcohol use in community pharmacy medicine review services. *BMC Health Services Research* 2020;20:943. <https://doi.org/10.1186/s12913-020-05797-z>
42. Cook L. Surgical versus conservative treatment of LC1 pelvic fractures in the elderly. In. 08/04/2019 ed: ISRCTN Registry; 2019.
43. Armstrong-Buisseret L, Brittain C, Kai J, David M, Anstey Watkins J, Ozolins M, et al. Lactic acid gel versus metronidazole for recurrent bacterial vaginosis in women aged 16 years and over: the VITA RCT. *Health Technol Assess* 2022;26:1-170. <https://doi.org/10.3310/zzkh4176>
44. Ralston SH. Treatment of osteogenesis imperfecta with parathyroid hormone and zoledronic acid. In. 10/08/2016 ed. ISRCTN Registry; 2016. <https://doi.org/https://doi.org/10.1186/ISRCTN15313991>
45. Flett L, Adamson J, Barron E, Brealey S, Corbacho B, Costa ML, et al. A multicentre, randomized, parallel group, superiority study to compare the clinical effectiveness and cost-effectiveness of external frame versus internal locking plate for complete articular pilon fracture fixation in adults. *Bone & Joint Open* 2021;2:150-63. <https://doi.org/10.1302/2633-1462.23.BJO-2020-0178>
46. Parker A, Mills N, Rooshenas L, Jepson M, Donovan JL, Arundel C, et al. SWAT 111: Staff training to improve participant recruitment into surgical randomised trials In. 2018 ed. SWAT Store: Northern Ireland Network for Trials Methodology Research; 2019.
47. Treweek S, Gillies K, Innes K, MacLennan G. SWAT 82: Sending Christmas cards to trial participants to improve retention. In. 2017 ed. SWAT Store: Northern Ireland Network for Trial Methodology Research 2017.
48. Parker A, Mills N, Rooshenas L, Jepson M, Donovan JL, Arundel C, et al. Staff training to improve participant recruitment into surgical randomised controlled trials: a feasibility study embedded within four randomised controlled trials. *International Clinical Trials Methodology Conference*; Brighton, abstract no. 202, p. PS8A.

49. Coleman E, Arundel C, Clark L, Doherty L, Gillies K, Hewitt C, et al. Bah humbug! Association between sending Christmas cards to trial participants and trial retention: randomised study within a trial conducted simultaneously across eight host trials. *BMJ (Clinical research ed)* 2021;375:e067742. <https://doi.org/10.1136/bmj-2021-067742>
50. Anand R, Bradley J, O'Neill B, McAuley D, Clarke M. A randomised trial of the effects on recruitment and retention of including a wet-ink signature and photograph in the patient invitation letter for a clinical trial: results from a Study Within a Trial (SWAT 3 and SWAT 53) [version 1; peer review: awaiting peer review]. *F1000Research* 2022;11. <https://doi.org/10.12688/f1000research.75339.1>
51. Parker A, Arundel C, Mills N, Rooshenas L, Jepson M, Donovan JL, et al. Staff training to improve participant recruitment into surgical randomised controlled trials: A feasibility study within a trial (SWAT) across four host trials simultaneously. *Research Methods in Medicine & Health Sciences* 2022; 10.1177/26320843221106950. <https://doi.org/10.1177/26320843221106950>
52. Dias J, Arundel C, Tharmanathan P, Keding A, Welch C, Corbacho B, et al. Dupuytren's interventions surgery versus collagenase (DISC) trial: study protocol for a pragmatic, two-arm parallel-group, non-inferiority randomised controlled trial. *Trials* 2021;22:671. <https://doi.org/10.1186/s13063-021-05595-w>
53. Armstrong G, Croft J, Corrigan N, Brown JM, Goh V, Quirke P, et al. IntAct: intra-operative fluorescence angiography to prevent anastomotic leak in rectal cancer surgery: a randomized controlled trial. *Colorectal Dis* 2018;20:O226-o34. <https://doi.org/10.1111/codi.14257>
54. Tharmanathan P, Arundel C. Effectiveness and cost-effectiveness of reverse shoulder arthroplasty versus hemiarthroplasty versus non-surgical care for acute 3 and 4 part fractures of the proximal humerus in patients aged over 65 years – the PROFHER-2 randomised trial. In. 05/04/2018 ed: ISRCTN Registry; 2018.
55. Metcalfe A, Gemperle Mannion E, Parsons H, Brown J, Parsons N, Fox J, et al. Protocol for a randomised controlled trial of Subacromial spacer for Tears Affecting Rotator cuff Tendons: a Randomised, Efficient, Adaptive Clinical Trial in Surgery (START:REACTS). *BMJ Open* 2020;10:e036829. <https://doi.org/10.1136/bmjopen-2020-036829>
56. Bradley JM, Anand R, O'Neill B, Ferguson K, Clarke M, Carroll M, et al. A 2 × 2 factorial, randomised, open-label trial to determine the clinical and cost-effectiveness of hypertonic saline (HTS 6%) and carbocisteine for airway clearance versus usual care over 52 weeks in adults with

bronchiectasis: a protocol for the CLEAR clinical trial. *Trials* 2019;20:747.

<https://doi.org/10.1186/s13063-019-3766-9>

57. Maguire L, Clarke M. SWAT 3: Gender of the person signing an invitation letter for a prospective study. In. 2014 ed. SWAT Store: The Northern Ireland Network for Trials Methodology Research 2014.
58. Anand R, Green N. SWAT 53: Including a photograph on the invitation letter for a prospective study. In. 2017 ed. SWATY Store: Northern Ireland Network for Trial Methodology Research; 2017.
59. Norton C, Syred J, Kerry S, Artom M, Sweeney L, Hart A, et al. Supported online self-management versus care as usual for symptoms of fatigue, pain and urgency/incontinence in adults with inflammatory bowel disease (IBD-BOOST): study protocol for a randomised controlled trial. *Trials* 2021;22:516. <https://doi.org/10.1186/s13063-021-05466-4>
60. White D, Parker A, Torgerson D. SWAT 137: A factorial embedded randomised controlled trial to test the impact on recruitment of a pen incentive and a brief participant information sheet. In. SWAT Store: The Northern Ireland Network for Trials Methodology Research 2018.
61. Mooney C. Multiple Symptoms Study 3. In. 02/10/2018 ed: ISRCTN Registry; 2022. <https://doi.org/https://doi.org/10.1186/ISRCTN57050216>
62. Goyal A, Mann GB, Fallowfield L, Duley L, Reed M, Dodwell D, et al. POSNOC—POSitive Sentinel NOde: adjuvant therapy alone versus adjuvant therapy plus Clearance or axillary radiotherapy: a randomised controlled trial of axillary treatment in women with early-stage breast cancer who have metastases in one or two sentinel nodes. *BMJ Open* 2021;11:e054365. <https://doi.org/10.1136/bmjopen-2021-054365>
63. Montgomery A, Brittain C, Khan S, Lewis M, Tan W, Goyal A, et al. SWAT 102: Addition of a pictorial aid to the patient information leaflet to improve recruitment in a randomised trial. In. 1/07/2019 ed; 2019.
64. Johnson G, Tabner A, Fakis A, Sherman R, Chester V, Bedford E, et al. Salbutamol for analgesia in renal colic: study protocol for a prospective, randomised, placebo-controlled phase II trial (SARC). *Trials* 2022;23:352. <https://doi.org/10.1186/s13063-022-06225-9>
65. Chetter I, Arundel C, Martin BC, Hewitt C, Fairhurst C, Joshi K, et al. Negative pressure wound therapy versus usual care for surgical wounds healing by secondary intention (SWHSI-2 trial): study

protocol for a pragmatic, multicentre, cross surgical specialty, randomised controlled trial. *Trials* 2021;22:739. <https://doi.org/10.1186/s13063-021-05662-2>

66. McCaffery J, Arundel C, Chetter I, Fairhurst C, Joshi K, Mott A, et al. SWAT 116: Impact on recruitment of adding an Infographic to a Patient Information Leaflet. In. 22/04/2019 ed: The Northern Ireland Network for Trials Methodology Research; 2019.

67. Coleman E, Whitemore R, Clark L, Daykin K, Clark M. Pre-notification and personalisation of text-messages to retain participants in a smoking cessation pregnancy RCT: an embedded randomised factorial trial [version 1; peer review: 1 approved, 1 approved with reservations]. *F1000Research* 2021;10. <https://doi.org/10.12688/f1000research.51964.1>

68. James S, Parker A, Cockayne S, Rodgers S, Fairhurst C, Torgerson D, et al. Including a pen and/or cover letter, containing social incentive text, had no effect on questionnaire response rate: a factorial randomised controlled Study within a Trial [version 2; peer review: 2 approved]. *F1000Research* 2021;9:623. <https://doi.org/10.12688/f1000research.23767.2>

69. Smith TO, Parsons S, Fordham B, Ooms A, Dutton S, Hing C, et al. Behaviour change physiotherapy intervention to increase physical activity following hip and knee replacement (PEP-TALK): study protocol for a pragmatic randomised controlled trial. *BMJ Open* 2020;10:e035014. <https://doi.org/10.1136/bmjopen-2019-035014>

70. Ooms A, The PEP-TALK Trial Team. SWAT 110: Printing the primary outcome on Pink Paper versus standard paper to increase participant engagement to postal questionnaires (PEPPER). In. 20/06/2019 ed. SWAT Store: Northern Ireland Network for Trial Methodology Research; 2019.

71. Ooms A, Parsons S, Dutton S, Garrett A, Fordham B, Hing C, et al. SWAT 110: Printing the primary outcome on Pink Paper versus standard paper to increase participant engagement to postal questionnaires (PEPPER). *Research Methods in Medicine & Health Sciences* 2022;3:49-54.

72. Rangan A, Brealey SD, Keding A, Corbacho B, Northgraves M, Kottam L, et al. Management of adults with primary frozen shoulder in secondary care (UK FROST): a multicentre, pragmatic, three-arm, superiority randomised clinical trial. *The Lancet* 2020;396:977-89. [https://doi.org/10.1016/S0140-6736\(20\)31965-6](https://doi.org/10.1016/S0140-6736(20)31965-6)

73. Hughes-Morley A, Brealey S, Keding A, Torgerson D, Hewitt C, Bailey M, et al. SWAT 44: Timing of text message prompts to increase trial participant response to postal questionnaires. In. SWAT Store: The Northern Ireland Network for Trials Methodology Research 2022.

74. Partha Sarathy P, Kottam L, Parker A, Brealey S, Coleman E, Keding A, et al. Timing of electronic reminders did not improve trial participant questionnaire response: a randomized trial and meta-analyses. *Journal of Clinical Epidemiology* 2020;122:70-7.  
<https://doi.org/10.1016/j.jclinepi.2020.03.001>
75. Hopewell S, Keene DJ, Maia Schlüssel M, Dritsaki M, Dutton S, Carr A, et al. Clinical and cost-effectiveness of progressive exercise compared with best practice advice, with or without corticosteroid injection, for the treatment of rotator cuff disorders: protocol for a 2x2 factorial randomised controlled trial (the GRASP trial). *BMJ Open* 2017;7:e018004.  
<https://doi.org/10.1136/bmjopen-2017-018004>
76. Hughes-Morley A, Torgerson D. SWAT 35: Personalised text message versus standard text message prompts for increasing response to postal questionnaires. In: *The Northern Ireland Network for Trials Methodology Research* 2015.
77. Cureton L, Marian IR, Barber VS, Parker A, Torgerson DJ, Hopewell S. Randomised study within a trial (SWAT) to evaluate personalised versus standard text message prompts for increasing trial participant response to postal questionnaires (PROMPTS). *Trials* 2021;22:502.  
<https://doi.org/10.1186/s13063-021-05452-w>
78. Cook L, Northgraves MJ, Fairhurst C, Ronaldson S, Torgerson DJ, Kent J, et al. Knee Replacement Bandaging Study (KReBS) evaluating the effect of a two-layer compression bandage system on knee function following total knee arthroplasty: study protocol for a randomised controlled trial. *Trials* 2019;20:261. <https://doi.org/10.1186/s13063-019-3344-1>
79. Tew G, Tilbrook H, Paul S-A, Howe L, Parker A, Bell K. SWAT 92: Pen incentive to enhance retention in a randomised trial. In. 2019 ed. *SWAT Store: Northern Ireland Network for Trials Methodology Research*; 2019.
80. Mitchell A, Cook L, Dean A, Fairhurst C, Northgraves M, Torgerson D, et al. An embedded randomised controlled retention trial of personalised text messages compared to non-personalised text messages in an orthopaedic setting [version 1; peer review: 1 approved]. *F1000Research* 2020;9. <https://doi.org/10.12688/f1000research.24244.1>
81. Mitchell A, Cook L, Dean A, Fairhurst C, Northgraves M, Torgerson D, et al. Using pens as an incentive for questionnaire return in an orthopaedic trial: an embedded randomised controlled retention trial [version 2; peer review: 1 approved, 1 approved with reservations]. *F1000Research* 2021;9. <https://doi.org/10.12688/f1000research.23018.2>

82. Whitemore R, Leonardi-Bee J, Naughton F, Sutton S, Cooper S, Parrott S, et al. Effectiveness and cost-effectiveness of a tailored text-message programme (MiQuit) for smoking cessation in pregnancy: study protocol for a randomised controlled trial (RCT) and meta-analysis. *Trials* 2019;20:280. <https://doi.org/10.1186/s13063-019-3341-4>
83. Anderson AS, Craigie AM, Gallant S, McAdam C, Macaskill EJ, Mutrie N, et al. Randomised controlled trial to assess the impact of a lifestyle intervention (ActWELL) in women invited to NHS breast screening. *BMJ Open* 2018;8:e024136. <https://doi.org/10.1136/bmjopen-2018-024136>
84. Treweek S, Anderson A, Gallant S. Sending pre-notification cards to trial participants before outcome measurement to improve retention. In: *The Northern Ireland Network for Trials Methodology Research* 2022.
85. Treweek S, Gallant S, Anderson A. SWAT 76 evaluation: randomised evaluation of sending pre-notification cards to trial participants before a face-to-face primary outcome measurement to increase attendance [version 2; peer review: 2 approved]. *F1000Research* 2021;10:84. <https://doi.org/10.12688/f1000research.50890.2>
86. Hammond A, Sutton C, Cotterill S, Woodbridge S, O'Brien R, Radford K, et al. The effect on work presenteeism of job retention vocational rehabilitation compared to a written self-help work advice pack for employed people with inflammatory arthritis: protocol for a multi-centre randomised controlled trial (the WORKWELL trial). *BMC Musculoskeletal Disorders* 2020;21:607. <https://doi.org/10.1186/s12891-020-03619-1>
87. Sutton C, Cotterill S, Forshaw D, Rhodes S, Hammond A. SWAT 86: Advance notification of trial participants before outcome data collection to improve retention. In: 21/08/2018 ed. *SWAT Store: Northern Ireland Network for Trials Methodology Research*; 2018.
88. Sutton CJ, Cotterill S, Forshaw D, Rhodes S, Haig A, Hammond A. Randomised evaluation of pre-notification of trial participants before self-report outcome data collection to improve retention: SWAT86. *Research Methods in Medicine & Health Sciences* 2022:26320843221098427.
89. Cockayne S, Pighills A, Adamson J, Fairhurst C, Drummond A, Hewitt C, et al. Can occupational therapist-led home environmental assessment prevent falls in older people? A modified cohort randomised controlled trial protocol. *BMJ Open* 2018;8:e022488. <https://doi.org/10.1136/bmjopen-2018-022488>
90. McDaid C, Cook L, James S, Ahkter Z, Podmore D. SWAT 144: Social Incentive Retention Cover Letter to Improve Questionnaire Response Rates. In: 14/11/2019 ed; 2019.

91. Cockayne S, Fairhurst C, Frost G, Hewitt C, Liddle M, Zand M, et al. SSheW study protocol: does slip resistant footwear reduce slips among healthcare workers? A randomised controlled trial. *BMJ Open* 2018;8:e026023. <https://doi.org/10.1136/bmjopen-2018-026023>
92. Cunningham-Burley R, Roche J, Fairhurst C, Cockayne S, Hewitt C, Iles-Smith H, et al. Enclosing a pen to improve response rate to postal questionnaire: an embedded randomised controlled trial. *F1000Res* 2020;9:577. <https://doi.org/10.12688/f1000research.23651.1>
93. Ahmed I, Innes K, Brazzelli M, Gillies K, Newlands R, Avenell A, et al. Protocol for a randomised controlled trial comparing laparoscopic cholecystectomy with observation/conservative management for preventing recurrent symptoms and complications in adults with uncomplicated symptomatic gallstones (C-Gall trial). *BMJ Open* 2021;11:e039781. <https://doi.org/10.1136/bmjopen-2020-039781>
94. Sinclair L, McFadden M, Tilbrook H, Mitchell A, Keding A, Watson J, et al. The smoking cessation in pregnancy incentives trial (CPIT): study protocol for a phase III randomised controlled trial. *Trials* 2020;21:183. <https://doi.org/10.1186/s13063-019-4042-8>
95. Abdel-Fattah M, Chapple C, Guerrero K, Dixon S, Cotterill N, Ward K, et al. Female Urgency, Trial of Urodynamics as Routine Evaluation (FUTURE study): a superiority randomised clinical trial to evaluate the effectiveness and cost-effectiveness of invasive urodynamic investigations in management of women with refractory overactive bladder symptoms. *Trials* 2021;22:745. <https://doi.org/10.1186/s13063-021-05661-3>
96. McClinton S, Starr K, Thomas R, MacLennan G, Lam T, Hernandez R, et al. The clinical and cost effectiveness of surgical interventions for stones in the lower pole of the kidney: the percutaneous nephrolithotomy, flexible ureterorenoscopy and extracorporeal shockwave lithotripsy for lower pole kidney stones randomised controlled trial (PUrE RCT) protocol. *Trials* 2020;21:479. <https://doi.org/10.1186/s13063-020-04326-x>
97. Tickle M, Ricketts DJN, Duncan A, O'Malley L, Donaldson PM, Clarkson JE, et al. Protocol for a Randomised controlled trial to Evaluate the effectiveness and cost benefit of prescribing high dose FLuoride toothpaste in preventing and treating dEntal Caries in high-risk older adultTs (reflect trial). *BMC Oral Health* 2019;19:88. <https://doi.org/10.1186/s12903-019-0749-x>
98. Anand R. SWAT 54 Protocol: Giving trial participants a thank you note or card after each study visit. In. 2017 ed. SWAT Store: Northern Ireland Network for Trials Methodology Research; 2017.

99. Anand R. Novel methods within a randomised controlled trial in respiratory disease. Queen's University Belfast.: Queen's University Belfast.; 2021.
100. Kearney RS, Dhanjal G, Parsons N, Ellard D, Parsons H, Haque A, et al. Acute Rehabilitation following Traumatic anterior shoulder dISlocAtioN (ARTISAN): protocol for a multicentre randomised controlled trial. *BMJ Open* 2020;10:e040623. <https://doi.org/10.1136/bmjopen-2020-040623>
101. Dhanjal G. SWAT 121: What are the effects on retention and follow-up of courtesy telephone calls versus postcards to trial participants following enrolment? . In. SWAT Store: The Northern Ireland Network for Trials Methodology Research 2019.
102. Hyslop M. Melatonin for Anxiety prior to General anaesthesia In Children (MAGIC). In. 10/01/2019 ed: ISRCTN Registry; 2019. <https://doi.org/10.1186/ISRCTN18296119>
103. Arundel C, Chetter I, Fairhurst C, Joshi K, McCaffery J, Mott A, et al. SWAT 119: Effects on retention of giving trial participants a thank you card following each study visit. In. 01/11/2019 ed. SWAT Store: The Northern Ireland Network for Trials Methodology Research; 2019.
104. Parker A, Brealey S, Sharma H, Welch C, Keding A, Flett L, et al. SWAT 109: The effectiveness of a text message reminder which participants can respond to, compared with a 'no reply' text message on questionnaire response rates. SWAT Store: The Northern Ireland Network for Trials Methodology Research; 2022. URL: <https://www.qub.ac.uk/sites/TheNorthernIrelandNetworkforTrialsMethodologyResearch/FileStore/Fileupload,943284,en.pdf> (accessed 22nd May, 2022).
105. Bedford L, Young B, Kendrick D, Vedhara K, Gallant S. SWAT 96: Unconditional or conditional incentives for initial and followup postal questionnaires in a clinical trial. In. 01/06/2013 ed: Northern Ireland Network for Trial Methodology Research; 2013.
106. Cook L, Backhouse M, McDaid C, Rex S, Parker A. SWAT 114: Effects of telephone calls or postcards to trial participants following enrolment on retention in a randomised trial. In. 1/JAN/2019 ed. SWAT Store: Northern Ireland Network for Trial Methodology Research; 2019.
107. Trial Forge. Trial Forge. 2022. URL: <https://www.trialforge.org/> (accessed 02/06/2022, 2022).
108. Clark L, Arundel C, Coleman E, Doherty L, Parker A, Hewitt C, et al. The PROMoting THE USE of SWATs (PROMETHEUS) Programme : lessons learnt and future developments for SWATs. *Research Methods in Medicine & Health Sciences* 2022;3:100-6. <https://doi.org/10.1177/26320843221089>



109. Staniszewska S, Brett J, Simera I, Seers K, Mockford C, Goodlad S, et al. GRIPP2 reporting checklists: tools to improve reporting of patient and public involvement in research. *BMJ (Clinical research ed)* 2017;358:j3453. <https://doi.org/10.1136/bmj.j3453>
110. Healy P, Galvin S, Williamson PR, Treweek S, Whiting C, Maeso B, et al. Identifying trial recruitment uncertainties using a James Lind Alliance priority setting partnership—the PRioRiTy (Prioritising recruitment in randomised trials) study. *Trials* 2018;19:1-12.
111. Brunsdon D, Biesty L, Brocklehurst P, Brueton V, Devane D, Elliott J, et al. What are the most important unanswered research questions in trial retention? A James Lind Alliance Priority Setting Partnership: the PRioRiTy II (Prioritising Retention in Randomised Trials) study. *Trials* 2019;20:593. <https://doi.org/10.1186/s13063-019-3687-7>
112. Dwyer C, Hynes S. SWAT 124: Exploring the impact of ineligibility on individuals expressing interest in a trial aimed at improving daily functioning regarding perceptions of self, research and likelihood of future participation: A PPI-infused qualitative SWAT. In. 16/01/2020 ed: Northern Ireland Network for Trials Methodology Research; 2020.
113. Hughes-Morley A. SWAT 26: Improving trial recruitment with a leaflet advertising patient and public involvement. In. 01/03/2014 ed: Northern Ireland Network for Trials Methodology Research; 2014.
114. ROMIO Study PPI Group. SWAT 146: Adding 'Withdrawal Slips' within follow up reminders for questionnaires. In. 29/08/2019 ed: Northern Ireland Network for Trials Methodology Research; 2019.
115. Morrissey E, Casey B, Power R, The D1 Now Young Adult Panel. SWAT 133: Branded gift and letter from PPI group to enhance questionnaire response rate in a randomised trial. In. 01/08/2020 ed: Northern Ireland Network for Trials Methodology Research; 2020.
116. Stewart DC, Bower P, Woolfall K, Gillies K. Actively Involving Patients/Public with Trials Methodology Research (TMR): Report of Workshop: MRC-NIHR Trials Methodology Research Partnership; 2020.
117. Treweek S, Altman DG, Bower P, Campbell M, Chalmers I, Cotton S, et al. Making randomised trials more efficient: report of the first meeting to discuss the Trial Forge platform. *Trials* 2015;16:261. <https://doi.org/10.1186/s13063-015-0776-0>

118. MRC-NIHR. MRC-NIHR Trials Methodology Research Partnership (TMRP). 2021. URL: <https://www.methodologyhubs.mrc.ac.uk/about/tmrp/> (accessed 18.10.2021).
119. Trial Forge. Trial Forge evidence packs 2021. URL: <https://www.trialforge.org/resource-category/evidence-pack/> (accessed 08.10.2021).
120. Jefferson L, Fairhurst C, Brealey S, Coleman E, Cook L, Hewitt C, *et al.* Remote or on-site visits were feasible for the initial setup meetings with hospitals in a multicenter surgical trial: an embedded randomized trial. *Journal of clinical epidemiology* 2018;**100**:13-21.
121. Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, *et al.* GRADE guidelines: 1. Introduction—GRADE evidence profiles and summary of findings tables. *Journal of clinical epidemiology* 2011;**64**:383-94. <https://doi.org/10.1016/j.jclinepi.2010.04.026>
122. Trial Forge. *Trial Forge SWAT network* 2021. URL: [https://www.trialforge.org/2021/06/swat\\_network/](https://www.trialforge.org/2021/06/swat_network/) (accessed September 2021).
123. Parker A, Knapp P, Treweek S, Madhurasinghe V, Littleford R, Gallant S, *et al.* The effect of optimised patient information materials on recruitment in a lung cancer screening trial: an embedded randomised recruitment trial. *Trials* 2018;**19**:1-8.
124. Fowell A, Johnstone R, Finlay I, Russell D, Russell I. Design of trials with dying patients: a feasibility study of cluster randomisation versus randomised consent. *Palliative medicine* 2006;**20**:799-804. <https://doi.org/10.1177/0269216306072554>
125. Larkey LK, Staten LK, Ritenbaugh C, Hall RA, Buller DB, Bassford T, *et al.* Recruitment of Hispanic women to the Women's Health Initiative. the case of Embajadoras in Arizona. *Control Clin Trials* 2002;**23**:289-98.
126. Sheridan R, Knapp P, Bower P, Madurasinghe V, Broadbent DM, Awoyale L, *et al.* Patient recruitment to a diabetic retinopathy screening trial through optimised patient information materials: an embedded study within a trial (SWAT). *F1000Research* 2020;**9**:779.
127. Hughes-Morley A, Hann M, Fraser C, Meade O, Lovell K, Young B, *et al.* The impact of advertising patient and public involvement on trial recruitment: embedded cluster randomised recruitment trial. *Trials* 2016;**17**:1-13. <https://doi.org/10.1186/s13063-016-1718-1>
128. Torgerson DJ, Dumville JC. Ethics review in research. Research governance also delays research 2004;**328**:710. <https://doi.org/10.1136/bmj.328.7441.710-a>
129. Mallick AA, O'Callaghan FJ. Research governance delays for a multicentre non-interventional study. *J R Soc Med* 2009;**102**:195-8. <https://doi.org/10.1258/jrsm.2009.080397>

130. Thompson AGH, France EF. One stop or full stop? The continuing challenges for researchers despite the new streamlined NHS research governance process. *BMC Health Services Research* 2010;10:124. <https://doi.org/10.1186/1472-6963-10-124>
131. Harman NL, Conroy EJ, Lewis SC, Murray G, Norrie J, Sydes MR, et al. Exploring the role and function of trial steering committees: results of an expert panel meeting. *Trials* 2015;16:597. <https://doi.org/10.1186/s13063-015-1125-z>
132. Medical Research Council. MRC guidelines for good clinical practice in clinical trials: Medical Research Council; 1998.
133. Arundel C. Challenges and Solutions to the Implementation of Studies Within A Trial (SWATs): The Experiences of the PROMETHEUS Programme. Submitted to the Research in Health Sciences and Medicine special SWATs edition 2021.
134. Cockayne S, Fairhurst C, Adamson J, Hewitt C, Hull R, Hicks K, et al. An optimised patient information sheet did not significantly increase recruitment or retention in a falls prevention study: an embedded randomised recruitment trial. *Trials* 2017;18:144. <https://doi.org/10.1186/s13063-017-1797-7>
135. Treweek S, Bevan S, Bower P, Campbell M, Christie J, Clarke M, et al. Trial Forge Guidance 1: what is a Study Within A Trial (SWAT)? *Trials* 2018;19:139. <https://doi.org/10.1186/s13063-018-2535-5>
136. Gouldner AW. The Norm of Reciprocity: A Preliminary Statement. *American Sociological Review* 1960;25:161-78. <https://doi.org/10.2307/2092623>
137. Cialdini RB, Vincent JE, Lewis SK, Catalan J, Wheeler D, Darby BL. Reciprocal concessions procedure for inducing compliance: The door-in-the-face technique. *Journal of Personality and Social Psychology* 1975;31:206-15. <https://doi.org/10.1037/h0076284>
138. Regan DT. Effects of a favor and liking on compliance. *Journal of Experimental Social Psychology* 1971;7:627-39. [https://doi.org/https://doi.org/10.1016/0022-1031\(71\)90025-4](https://doi.org/https://doi.org/10.1016/0022-1031(71)90025-4)
139. Falk A, Fischbacher U. A theory of reciprocity. *Games and economic behavior* 2006;54:293-315.
140. Bell K, Clark L, Fairhurst C, Mitchell N, Lenaghan E, Blacklock J, et al. Enclosing a pen reduced time to response to questionnaire mailings. *J Clin Epidemiol* 2016;74:144-50. <https://doi.org/10.1016/j.jclinepi.2015.12.004>

141. Sharp L, Cochran C, Cotton SC, Gray NM, Gallagher ME. Enclosing a pen with a postal questionnaire can significantly increase the response rate. *J Clin Epidemiol* 2006;59:747-54. <https://doi.org/10.1016/j.jclinepi.2005.10.014>
142. Gkekas A, Evans A, Parker A, Ronaldson S, Torgerson D. A systematic review of economic evaluations alongside Studies Within A Trial (SWATs) for improving recruitment and retention in RCTs. *Research Methods in Medicine & Health Sciences* Submitted.
143. Townsend D, Mills N, Savović J, Donovan JL. A systematic review of training programmes for recruiters to randomised controlled trials. *Trials* 2015;16:432-. <https://doi.org/10.1186/s13063-015-0908-6>
144. Delaney H, Devane D, Hunter A, Hennessy M, Parker A, Murphy L, et al. Limited evidence exists on the effectiveness of education and training interventions on trial recruitment; a systematic review. *Journal of clinical epidemiology* 2019;113:75-82. <https://doi.org/10.1016/j.jclinepi.2019.05.013>
145. Mills N, Gaunt D, Blazeby JM, Elliott D, Husbands S, Holding P, et al. Training health professionals to recruit into challenging randomized controlled trials improved confidence: the development of the QuinteT randomized controlled trial recruitment training intervention. *J Clin Epidemiol* 2018;95:34-44. <https://doi.org/10.1016/j.jclinepi.2017.11.015>
146. Schoenfeld D. Partial residuals for the proportional hazards regression model. *Biometrika* 1982;69:239-41.
147. Tew G, Tilbrook H. SWAT 93: Offer of a free yoga class to participants in the control group on completion of the trial follow-up, to enhance retention and reduce contamination. In. 01/04/2018 ed; 2018.
148. O'Neill L, Knapp P, Doyle S, Guinan E, Hussey J. SWAT 100: Patient and family co-developed participant information to improve recruitment rates, retention, and patient understanding of a randomised trial. In. 01/08/2018 ed: Northern Ireland Network for Trials Methodology Research; 2018.
149. Rothwell JC, Julious SA, Cooper CL. A study of target effect sizes in randomised controlled trials published in the Health Technology Assessment journal. *Trials* 2018;19:544. <https://doi.org/10.1186/s13063-018-2886-y>

150. Treweek S, Bevan S, Bower P, Briel M, Campbell M, Christie J, et al. Trial Forge Guidance 2: how to decide if a further Study Within A Trial (SWAT) is needed. *Trials* 2020;21:33. <https://doi.org/10.1186/s13063-019-3980-5>
151. Arundel C, Coleman E, Fairhurst C, Peckham E, Bailey D, Gilbody S. The effectiveness of a contingent financial incentive to improve trial follow up; a randomised study within a trial (SWAT). *F1000Research* 2019;8:1937-. <https://doi.org/10.12688/f1000research.21059.2>
152. Cochrane A, Welch C, Fairhurst C, Cockayne S, Torgerson DJ. An evaluation of a personalised text message reminder compared to a standard text message on postal questionnaire response rates: an embedded randomised controlled trial. *F1000Res* 2020;9:154. <https://doi.org/10.12688/f1000research.22361.1>
153. Mitchell N, Hewitt CE, Lenaghan E, Platt E, Shepstone L, Torgerson DJ. Prior notification of trial participants by newsletter increased response rates: a randomized controlled trial. *Journal of Clinical Epidemiology* 2012;65:1348-52. <https://doi.org/https://doi.org/10.1016/j.jclinepi.2012.05.008>
154. Mitchell N, Hewitt CE, Torgerson DJ. A controlled trial of envelope colour for increasing response rates in older women. *Aging Clin Exp Res* 2011;23:236-40. <https://doi.org/10.1007/bf03337749>
155. F1000Research. Studies Within A Trial (SWAT) Collection. *F1000Research*; 2022. URL: <https://f1000research.com/collections/swat/about-this-collection> (accessed 02/06/2022, 2022).
156. UK Government legislation. Data Protection Act 2018. [legislation.gov.uk](https://www.legislation.gov.uk): The National Archives; 2018. URL: <https://www.legislation.gov.uk/ukpga/2018/12/contents/enacted> (accessed 02/06/2022, 2022).
157. SAGE Journals. *Research Methods in Medicine & Health Sciences*. SAGE Journals; 2022. URL: <https://journals.sagepub.com/home/rmm> (accessed 02/06/2022, 2022).
158. Adamson J, Hewitt CE, Torgerson DJ. Producing better evidence on how to improve randomised controlled trials. *Bmj* 2015;351:h4923.
159. Treweek S, Bevan S, Bower P, Briel M, Campbell M, Christie J, et al. Trial Forge Guidance 2: how to decide if a further Study Within A Trial (SWAT) is needed. *Trials* 2020;21:1-9.
160. ISRCTN. ISRCTN registry In; 2020.

161. ClinicalTrials.gov. ClinicalTrials.gov. 2022. URL: <https://clinicaltrials.gov/ct2/resources/trends> (accessed 27/05/2022, 2022).



## Appendix 1

*List of funded host trials and SWATs*

# PROMETHEUS funded host trials and their characteristics

1. **ACTWELL:** A randomised control trial to assess the impact of a lifestyle intervention (ActWELL) in women invited to NHS breast screening. ISRCTN11057518 (Date registered: 15/02/2017)
2. **ACTIVE:** External frame versus internal locking plate for articular pilon fracture fixation: a multi-centre randomised controlled trial. ISRCTN98152560 (Date registered: 26/02/2018)
3. **ARTISAN:** Acute Rehabilitation following Traumatic anterior shoulder dISlocAtion (ARTISAN) - A Multi Centre RCT. ISRCTN63184243 (Date registered: 03/09/2018)
4. **CHAMP-1:** a multi-centre, cluster randomised controlled, open pilot trial to establish the feasibility of conducting a large scale study comparing an intervention discussing alcohol within routine medication consultations with usual care in community pharmacies. ISRCTN57447996 (Date registered: 19/06/2019)
5. **C-GALL:** A randomised controlled trial comparing laparoscopic cholecystectomy with observation/conservative management for preventing recurrent symptoms and complications in adults with uncomplicated symptomatic gallstones. ISRCTN55215960 (Date registered: 27/05/2016)
6. **CLEAR:** A 2x2 factorial randomised open label trial to determine the clinical and cost-effectiveness of hypertonic saline (HTS 6%) and carbocisteine for airway clearance versus usual care over 52 weeks in bronchiectasis. ISRCTN89040295 (Date Registered: 25/06/2018)
7. **CPIT III:** The smoking cessation in pregnancy incentives trial. ISRCTN15236311 (Date registered: 13/10/2017)
8. **DISC:** Dupuytren's interventions surgery vs collagenase. ISRCTN18254597 (Date registered: 04/04/2017)
9. **FUTURE:** Female Urgency, Trial of Urodynamics as Routine Evaluation; a superiority randomised clinical trial to evaluate the effectiveness and cost effectiveness of invasive

urodynamic investigations in management of women with refractory overactive bladder (OAB) symptoms. ISRCTN63268739 (Date registered: 14/09/2017)

10. GRASP: GRASP: Getting it Right: Addressing Shoulder Pain. ISRCTN16539266 (Date registered: 13/07/2016)

11. IBD BOOST: Living well with inflammatory bowel disease: optimising management of symptoms of fatigue, abdominal pain, and faecal urgency/incontinence via tailored online self-management: the IBD-BOOST programme. ISRCTN71618461 (Date registered: 09/09/2019)

12. IntAct: Intraoperative Fluorescence Angiography to Prevent Anastomotic Leak in Rectal Cancer Surgery. ISRCTN13334746 (Date registered: 02/05/2017)

13. KReBs: A Randomised Controlled Trial of the effect of a Two-layer Compression Bandage System on Knee Function following Total Knee Arthroplasty. ISRCTN87127065 (Date registered: 20/02/2017)

14. LIFE: Surgical versus conservative treatment of LC1 pelvic fractures in the elderly. ISRCTN16478561 (Date registered: 01/04/2019)

15. MAGIC: Melatonin for Anxiety prior to General anaesthesia In Children. ISRCTN18296119 (Date registered: 08/01/2019)

16. MIQUIT 3: RCT and meta-analysis testing effectiveness and cost-effectiveness of a tailored text message programme (MiQuit) for smoking cessation in pregnancy. ClinicalTrials.gov Identifier: NCT03231553 (ate registered: 27/072017)

17. MSS3: Multiple Symptoms Study 3: pragmatic trial of a community based clinic for patients with persistent (medically unexplained) physical symptoms. ISRCTN57050216 (Date registered: 24/09/2018)

18. OTIS: Does Occupational Therapist led environmental assessment and modification reduce falls among high risk older people. ISRCTN22202133 (Date registered: 20/06/2016)

19. PEP-TALK: A behaviour change physiotherapy intervention to increase physical activity following hip and knee replacement: a pragmatic phase III randomised controlled trial. ISRCTN29770908 (Date registered: 23/10/2018)

20. POSNOC: Positive sentinel node: adjuvant therapy alone versus adjuvant therapy plus clearance or axillary radiotherapy. ISRCTN54765244 (Date registered 25/02/2014)

21. PROFHER 2: Effectiveness and cost-effectiveness of reverse shoulder arthroplasty versus hemiarthroplasty versus non-surgical care for acute 3 and 4 part fractures of the proximal humerus in patients aged over 65 years – the PROFHER-2 randomised trial. ISRCTN76296703 (Date registered: 05/04/2018)

22. PUrE RCT: PUrE: percutaneous nephrolithotomy, flexible ureterorenoscopy and extracorporeal shockwave lithotripsy for lower pole kidney stones. ISRCTN98970319 (Date registered: 11/11/2015)



23. REFLECT: A randomised controlled trial to evaluate the cost effectiveness of prescribing high concentration fluoride toothpaste to prevent tooth decay in older adults. ISRCTN11992428 (Date registered: 02/06/20017)
24. SARC: Salbutamol for analgesia in renal colic: A prospective, randomised, placebo-controlled Phase II trial. ISRCTN14552440 (Date registered: 01/07/2019)
25. SSHEW: The SSHeW study - Stopping slips among healthcare workers: A research study about slip resistant footwear in the NHS workplace. ISRCTN33051393 (Date registered: 14/03/2017)
26. START:REACTS: Sub-acromial spacer for Tears Affecting Rotator cuff Tendons: a Randomised, Efficient, Adaptive Clinical Trial in Surgery. ISRCTN17825590 (Date registered: 05/03/2018)
27. SWHSI-2: Surgical Wounds Healing By Secondary Intention – 2. ISRCTN26277546 (Date registered: 15/02/2019)
28. TOPAZ: The TOPaZ Trial - Treatment of Osteogenesis Imperfecta with Parathyroid hormone and Zoledronic acid. ClinicalTrials.gov Identifier: NCT03735537 (Date registered: 8/11/2018)
29. UK FROST: United Kingdom Frozen Shoulder Trial (UK FroST). ISRCTN48804508 (Date registered: 18/07/2014)
30. VITA: Lactic acid gel versus metronidazole for recurrent bacterial vaginosis in women aged 16 years and over: the VITA RCT. ISRCTN14161293 (Date registered: 18/09/2017)
31. WORKWELL: A randomised controlled trial of job retention vocational rehabilitation for employed people with inflammatory arthritis: the WORKWELL trial. ISRCTN61762297 (Date registered: 13/05/2019)

**Table 17: Characteristics of currently funded host randomised controlled trials**

Trial	Aim	Speciality	Condition	Study design	Date of recruitment	Setting	Population	Intervention (s)	Comparator	No. of expected participants	Follow up time points and modality	Primary outcome	Overall trial end date
ACTIVE	Assess the effectiveness of external fixation versus internal fixation for Type C pilon fractures	Trauma and Orthopaedics	Treatment of ankle fractures	2 arm RCT	9/3/2018 to 28/2/2021	Multicentre, Secondary care, UK	Adults with Pilon C fracture	internal fixation	external fixation	334	3, 6, 12 and 24 months by postal questionnaire	Disability Rating Index (DRI) at 12 months post-randomisation	31/08/2022
ACTWELL	Assess the impact of a lifestyle intervention in women invited to NHS breast screening	Not Applicable (NA)	Obesity	2 arm RCT	01/07/2017 to 14/08/2018	Multicentre, UK	Women aged 50-70 years, BMI >25 km/m	Lifestyle intervention	Usual care	552	3 months by telephone 12 months by research nurses	Change in body weight and change in physical activity at 12 months	Unclear
ARTISAN	Assess if a course of physiotherapy is of clear benefit when compared to a single session of advice	Trauma and orthopaedics	Traumatic anterior shoulder dislocation	2 arm RCT	01/11/2018 to 30/09/2020	Multicentre, secondary care, UK	Adults with traumatic shoulder dislocation	A course of physiotherapy	A single advice session	478	6 weeks, 3, 6 and 12 months by postal questionnaire	Oxford Shoulder Instability Score at 6 months	30/11/2021
CHAMP-1	Incorporating discussion of alcohol within medication consultations in community pharmacy	NA	Alcohol misuse	2 arm RCT	24/06/2019 to 07/08/2019	Multicentre, UK	Adults screened positive for unhealthy drinking on a single item alcohol screening question	Incorporate discussion of alcohol	Usual medication consultation	820	2 months by telephone	Alcohol consumption at 2 months	31/12/2022
C-GALL	Conservative management & cholecystectomy in terms of patient QOL and cost-effectiveness	Surgery	Gallstones	2-arm RCT	01/09/2016 to 01/08/2018	Multicentre, Secondary care, UK	Adults aged over 18 with symptomatic uncomplicated gallstone disease	Medical management with analgesia, as required, and dietary advice	Laparoscopic cholecystectomy	430	3, 9 and 18 months questionnaire	Quality of life using area under the curve (AUC) at up to 18 months post-randomisation using the SF-36 bodily pain domain	01/10/2020

Trial	Aim	Speciality	Condition	Study design	Date of recruitment	Setting	Population	Intervention (s)	Comparator	No. of expected participants	Follow up time points and modality	Primary outcome	Overall trial end date
CLEAR	Assess the effectiveness of hypertonic saline and carbocisteine combinations in airway clearance	Respiratory	Bronchiectasis	2x2 factorial RCT	27/06/2018 to 31/12/2020	Multicentre, secondary care, UK	Adults with bronchiectasis	Saline, carbocisteine	Usual Care	Unclear	2, 8, 26, 52 weeks questionnaire at visits	Mean number of exacerbations over 52 weeks	30/06/2022
CPIT III	Assess the effectiveness of financial incentive to encourage pregnant smokers engaging with smoking cessation service and quite smoking	Obstetrics	Smoking cessation	2 arm RCT	01/02/2018 to 31/12/2019	Multicentre, Secondary care, UK	Pregnant women (less than 24 weeks) aged >16	Up to £400 depending on engagement with smoking cessation services and negative CO testing at 3 time-points	Offer of smoking cessation and up to £75 on completion of trial follow-up	940	34-38 weeks gestation and 6 month post-partum for every participant by postal questionnaire, intervention group will be followed up face-to-face at smoking cessation services (SCS) + 4 and 12 weeks post SCS	Self-reported abstinence for at least 8 weeks prior to 34-38 weeks gestation	30/11/2020
DISC	Assess the effectiveness of collagenase injections in Dupuytren's Contractures (DC)	Plastic Surgery	Treatment of DC	2 arm RCT	01/05/2017 to 31/10/2019	Multicentre, Secondary care, UK	Adults >18 with DC	Collagenase injections	limited fasciectomy surgery	710	3, 6, 12 and 24 months at clinics	Patient Evaluation Measure at baseline, 3, 6, 12 and 24 months	31/10/2021
FUTURE	Whether routine Urodynamics investigation & comprehensive clinical assessment significantly improves patient-reported success rates following treatment, vs comprehensive clinical assessment only	Reproductive Health and Childbirth	Overactive bladder (OAB)	2 arm RCT	01/10/2017 to 31/05/2020	Multicentre, Secondary care, UK	Women ≥ 18 years of age	comprehensive clinical assessment only	Urodynamics plus comprehensive clinical assessment	1096	3, 6, and 15 months by postal questionnaire	Patient Global Impression of Improvement - Index (PGI-I)	30/04/2022

Trial	Aim	Speciality	Condition	Study design	Date of recruitment	Setting	Population	Intervention (s)	Comparator	No. of expected participants	Follow up time points and modality	Primary outcome	Overall trial end date
GRASP	Assess cost effectiveness of progressive exercise in rotator cuff disorders	Musculoskeletal	Rotator cuff disorders	2x2 factorial RCT	01/02/2017 to 02/05/2019	Multicentre, secondary care, UK	Adults >18 with rotator cuff disorder	Progressive exercise programme, best practice advice session, injection	best practice advice session	704	2, 6 and 12 month by postal questionnaires	Shoulder pain and disability index	31/08/2021
IBD BOOST	Improve QoL of people with Inflammatory bowel disease (IBD) by reducing symptoms	Gastroenterology	IBD	RCT	Unclear	Secondary care, UK	Patients with IBD	Online training package for nurses, and online self-management programme	Usual care	680	Unclear?	Condition specific quality of life at 6 months	10/2022
IntAct	evaluate whether intraoperative fluorescent angiography (IFA) decreased anastomotic leak (AL) rate	Colorectal surgery	Rectal cancer	2 arm RCT	01/07/2017 to 31/08/2020	Secondary care	Adults >18 with rectal cancer undergoing curative elective surgery	IFA with surgery	Surgery only	880	30 and 90 days by questionnaire and clinical examination	Clinical anastomotic leak rate at 90 days	31/05/2021
KReBs	effect of a Two-layer Compression Bandage System on Knee Function following Total Knee Arthroplasty	Orthopaedics	Total knee replacement	2 arm RCT	01/03/2017 to 31/08/2018	Multicentre, secondary care, UK	Adults having a total knee replacement	2-layer compression bandage	Crepe bandage and synthetic wool layer	2600	10 days, 4 weeks, 6- and 12-month postal questionnaire	Oxford knee score 12 months postoperatively	30/04/2020
L1FE	Assess the effectiveness of surgical fixation with INFIX compared to non-surgical management of LC-1 fragility fractures in older adults	Trauma and Orthopaedics	Treatment of Pelvic fractures	2 arm RCT	09/07/2018 to 31/07/2019	Multicentre, Secondary care, UK	Adults >60 with a LC-1 fracture with severe pain and reduced mobility	Surgical fixation	Non operative management	600	2 weeks by postal questionnaires or telephone interview. 4 weeks, 12 weeks and 6 months follow up in clinic	EQ-5D-5L at 6 months	31/03/2023

Trial	Aim	Speciality	Condition	Study design	Date of recruitment	Setting	Population	Intervention (s)	Comparator	No. of expected participants	Follow up time points and modality	Primary outcome	Overall trial end date
MAGIC	Assess the effectiveness of Melatonin as premedication for surgery in children	Surgery	Dental ophthalmological or ENT surgery	2 arm RCT	10/07/2019 to 15/07/2020	Multicentre, secondary care, UK	Children aged 5-14 years undergoing dental/ophthalmological/ENT surgery	Melatonin	Midazolam	1650	14 days telephone follow-up	Preoperative distress using the modified Yale Preoperative Anxiety Scale	15/07/2021
MIQUIT 3	Determine whether or not MiQuit (text-message support programme) is effective in addition to standard support for smoking cessation in pregnancy	Obstetrics	Smoking cessation	2 arm RCT	unclear	Multicentre, UK	Pregnant women <25 weeks gestation, aged 16 or over	Receive MiQuit text message cessation programme	Usual care	692	Up to 36 weeks gestation, via text messaging	Self-reported smoking abstinence	01/01/2020
MSS3	Determine whether a 50-minute symptoms clinic is cost effective for medically unexplained symptoms (MUS)	General Practice	MUS	2 arm RCT	15/10/2018 to 31/10/2019	Multicentre, Primary care, UK	Adults aged between 18-69 with MUS	50-minute symptoms clinic with 15 minute follow-ups	Usual care	376	12- and 52-weeks postal questionnaire	PHQ-15 at 52 weeks	31/07/2021
OTIS	Assess the effectiveness of an occupational therapist led home environmental assessment and modification for high risk of falling in older people	General Practice	Falls	2 arm RCT	01/18/2016 to 02/08/2018	Multicentre, Primary care, UK	Older people aged 65 and older with risk factors for falls	an occupational therapist led home environmental assessment and a falls prevention leaflet	Usual care	1299	4-, 8- and 12-months postal questionnaire	Number of falls experienced in the 12 months	31/12/2019

Trial	Aim	Speciality	Condition	Study design	Date of recruitment	Setting	Population	Intervention (s)	Comparator	No. of expected participants	Follow up time points and modality	Primary outcome	Overall trial end date
PEP-TALK	Assessing the effectiveness of group discussion in addition to physiotherapy on improving physical activity following hip and knee replacement.	Musculoskeletal diseases	Hip and knee replacement	2 arm RCT	01/01/2019 to 31/12/2019	Multicentre, secondary care, UK	Adults following primary unilateral total hip or knee replacement	Group based exercise sessions in addition to physiotherapy	Physiotherapy only	250	6- and 12-months questionnaire (postal or online)	Physical activity at 6 and 12 months (using the UCLA activity scale)	31/07/2021
POSNOC	Assess the effectiveness of adjuvant therapy alone versus adjuvant therapy plus clearance or axillary radiotherapy in women with early-stage breast cancer.	Cancer	Breast cancer	2 arm RCT	01/01/2014 to 31/08/2021	Multicentre, secondary care, international	Women >18 years with early breast cancer, with macro-metastasis in sentinel node biopsy.	Adjuvant therapy alone	Adjuvant therapy plus axillary treatment (usual care)	1900	3, 6, 12, 24 and 36 months	Axillary recurrence at 5 years	31/12/2023
PROFHER-2	Assess the effectiveness of reverse shoulder arthroplasty, hemiarthroplasty and non-surgical treatment for proximal humerus fractures	Trauma and Orthopaedics	Treatment of proximal humeral fractures	3 arm RCT	01/06/2018 to 31/05/2021	Multicentre, Secondary care, UK	Adults >65 with a 3-4 part humeral fracture	Reverse Shoulder Arthroplasty vs hemiarthroplasty	Conservative management	380	6 months face-to-face follow-up. Postal questionnaire at 1 year and 2 years	Oxford Shoulder Score at 6, 12 and 24 months	31/05/2023
PUR RCT	Assess the effectiveness of three treatment options for lower kidney stones	Urology	Treatment of urolithiasis	2 arm RCT	01/05/2015 to 02/06/2019	Multicentre, Secondary care, UK	Adults >16 with lower pole stones	FURS	Usual care or (PCNL or ESWL)	1044	Weekly up to 12 weeks, 3 months, and 12 months by questionnaires either face-to-face, postal, phone or email	EQ-5D-5L at 1, 2, 4, 8, and 12 weeks	01/12/2020

Trial	Aim	Speciality	Condition	Study design	Date of recruitment	Setting	Population	Intervention (s)	Comparator	No. of expected participants	Follow up time points and modality	Primary outcome	Overall trial end date
REFLECT	To evaluate the effectiveness and cost benefit of General Dental Practitioner (GDP) prescribing of 5000 ppm fluoride toothpaste and usual care compared to usual care alone in individuals 50 years and over with high-risk of caries	Dentistry	Dental Caries	2 arm RCT	10/02/2018 to 30/09/2019	Multicentre, General dental practices, UK	Adults aged 50 and over	Prescription of 5000ppm fluoride toothpaste by dentist	Usual care	1174	12, 24 and 36 months by patient questionnaire; clinical assessments up to 36 months	Number and proportion of individuals requiring restoration or extraction or endodontic treatment due to caries	30/06/2022
SARC	Does salbutamol reduce the pain of kidney stones when used alongside normal pain relief?	Urological and genital diseases	Kidney stones	2 arm RCT	16/09/2019 to 01/08/2021	Secondary care, UK	Adults with a working diagnosis of renal colic	Salbutamol	placebo	118	15-, 30-, 60- and 120-minutes questionnaire during visit	Change in pain score measured using visual analogue scale at 30 minutes	31/12/2021
SSHEW	Whether wearing slip resistant shoes can reduce the number of slips, falls and injuries NHS staff have at work.	Signs and symptoms	Slips, falls and injuries	2 arm RCT	10/03/2017 to 10/01/2019	Multicentre, secondary care, UK	NHS staff	Slip resistant shoes	Usual work footwear	4400	14 weekly text messages	Self-reported slips in the workplace	10/01/2019
START:REACTS	Compare arthroscopic debridement to arthroscopic debridement with InSpace balloon on shoulder function, pain and QOL following shoulder surgery for rotator cuff tears that cannot be repaired.	Musculoskeletal diseases	Rotator cuff tears	2 arm RCT	01/06/2018 to 30/06/2020	Multicentre, secondary care UK	Patients with irreparable rotator cuff tear	Arthroscopic debridement with the InSpace balloon	Usual care	212	3-, 6- and 12-month questionnaire	Shoulder function measured with Constant-Murley score at 12 months	31/12/2021

Trial	Aim	Speciality	Condition	Study design	Date of recruitment	Setting	Population	Intervention (s)	Comparator	No. of expected participants	Follow up time points and modality	Primary outcome	Overall trial end date
TOPAZ	Assess effectiveness of parathyroid hormone & zoledronic acid treatment of osteogenesis imperfecta	Orthopaedics	Osteogenesis imperfecta	2 arm RCT	Unclear	Multicentre, secondary care, UK	Adults with osteogenesis imperfecta	Teriparatide and zoledronic acid	Usual care (bisphosphonate treatment)	380	12- and 24-months visits, telephone calls every 6 months	Incidental fractures (validated by x-ray or other imaging) (approx. 5 years)	01/04/2023
UK FROST	Assess effectiveness of early structured physiotherapy (ESP) versus manipulation under anaesthesia (MUA) versus arthroscopic capsular release (ACR)	Musculoskeletal disease	Frozen shoulder	3 arm RCT	01/01/2015 to 31/12/2017	Multicentre, secondary care, UK	Adults with frozen shoulder	MUA or ACR with MUA	ESP	503	3-, 6- and 12-months questionnaires	Oxford shoulder score at 12 months	30/06/2019
SWHSI-2	Assess the effectiveness of Negative Pressure Wound Therapy (NPWT) on wound healing	General Surgery	Treatment of surgical wounds	2 arm RCT	01/05/2019 to 30/04/2021	Multicentre, Secondary care, UK	Adults >16 with an open surgical wound	NPWT	Usual care	696	weekly phone follow-up until would healing, 3 face to face visits after wound healing, postal questionnaires at 3,6 and 12 months	Time to healing in days from randomisation	30/04/2022
VITA	Lactic acid gel versus metronidazole for recurrent bacterial vaginosis in women aged 16 years and over: the VITA RCT	Gynaecology	Bacterial vaginosis	2 arm RCT	25/09/2017 to 28/06/2019	Multicentre, Secondary care, UK	Women aged 16+ with symptoms of bacterial vaginosis (BV) plus history of one or more episodes of BV in previous 2 years	Intravaginal lactic acid gel	Oral metronidazole	1900	Web-based questionnaires at 2 weeks, 3 months and 6 months	Resolution of bacterial vaginosis based on participant reported resolution of symptoms at 14 days post randomisation	28/06/2020



Trial	Aim	Speciality	Condition	Study design	Date of recruitment	Setting	Population	Intervention (s)	Comparator	No. of expected participants	Follow up time points and modality	Primary outcome	Overall trial end date
WORKWELL	Assessing the effectiveness of work advice for people with arthritis	Musculoskeletal diseases	Inflammatory arthritis	2 arm RCT	18/02/2019 to 30/06/2020	Multicentre, UK	Adults with inflammatory arthritis	Face to face contact with a WORKWELL therapist (identifying work problems and providing advice)	Self-help information pack	240	6- and 12-month questionnaire	Presenteeism (work productivity) measured using Combined Work Activities Limitations Scale at 12 months	31/01/2022

Appendix 2

*PROMETHEUS Christmas Card SWAT Intervention*

Figure 6: PROMETHEUS Christmas Card SWAT intervention



## Appendix 3

### *Draft HRA approvals Guidance process for SWATs*

#### **Study Within A Trial (SWAT)**

##### *Guidance for SWAT applicants*

**Note:** This guidance is primarily intended for coordinated SWATs (see *Glossary*) where there is a coordinated evaluation in several host trials and the individual SWAT evaluations within them.

SWATs planned as a single evaluation within a particular clinical trial should look at the SWAT groups described here and, if approval is required, submit their SWAT as part of their trial submission, or as an amendment, to the ethics committee responsible for the clinical trial approval.

#### **Key principles**

SWATs are run as discreet studies but with the expectation that they will be replicated (in order to boost statistical power and assess the generalisability of the findings). Therefore, it is important to make a distinction between:


1. The overall, overarching SWAT evaluation (see *Glossary*) which will evaluate data from multiple SWATs which are run in multiple trials, and
2. The individual evaluations of a SWAT intervention within individual host trials, each of which will contribute data to the overarching evaluation.

The expectation is that approval will be obtained for the overarching SWAT evaluation (1) and then each individual SWAT (2) will come under the approvals obtained for the host trial; this may be from the start of the trial or as a substantial amendment after the trial has commenced. The SWAT evaluation lead and host trial team roles are shown in figure 7.

Figure 7: SWAT evaluation lead and host trial team roles

The *SWAT evaluation lead* has responsibility for:

- a) applying for and holding required approvals for the overarching SWAT, and sharing these with host trials.
- b) coordinating the overarching SWAT evaluation according to the approved protocol.
- c) collecting data from each of those host trials on the effect of the SWAT intervention in the trial.




Each *host trial team* has approval to:


- a) embed an evaluation of the SWAT intervention into their host trial.
- b) securely transfer data on the effect of the SWAT intervention in the trial to the SWAT evaluation lead.

## SWAT groups

The following table sets out which approvals should be obtained for overarching SWATs and for individual SWATs to be included with a host trial.

Table 18: Guide to which approvals for should be obtained for individual and overarching SWATs

Group	Description
1	<p>The activities targeted by the SWAT are not of a type that needs to be described in detail in a host trial protocol (and so have not required previous approval) but are operational or management activities generally done by the central trial team.</p> <p>Examples of this SWAT group include:</p> <ul style="list-style-type: none"> <li>● Randomising the host trial Chief Investigator to visit some recruiting sites and not others to see whether this improves recruitment.</li> <li>● Randomising monitoring visits to sites being undertaken by either Skype or in person.</li> </ul> <div style="text-align: center; margin-top: 20px;">  <pre> graph TD     A[SWAT Group 1] --&gt; B[REC Review]     A --&gt; C[UK SWR]     A --&gt; D[Host Trial]     B --&gt; E[No]     C --&gt; F[Yes]     D --&gt; G[REC]     D --&gt; H[SWR]     G --&gt; I[No Amendment]     H --&gt; J[No Amendment]             </pre> </div>
2	<p>The SWAT interventions affect participants but affect a process that does not <i>need</i> to be described in detail in a host trial protocol (and so have not <i>required</i> previous approval).</p> <p>These interventions are provided to the research team via operational or instruction documents, rather than a protocol, which are not reviewed by the REC. However, in some cases these interventions are also described in the protocol. Where this is the case, the intervention remains a Group 2 SWAT due to the low risk and unobtrusive nature of the intervention.</p> <p>Examples of this SWAT group include:</p> <ul style="list-style-type: none"> <li>● Variation in the gender of the person who signs a trial invitation letter to assess whether this impacts recruitment</li> <li>● Randomising whether trial participants are sent a Christmas card or not to assess whether this impacts on trial retention</li> </ul>



**REC review**

It is the responsibility of the sponsor to confirm the SWAT group, whether this is the overarching SWAT (SWAT sponsor) or the individual SWATs (host trial sponsor). It is likely that sponsors of overarching SWATs (or a member of the overarching SWAT team working with the sponsor) will want to contact the HRA for advice in most cases. The HRA will provide written advice identifying which group they believe the study to fall into which can then be provided to host trial sponsors

To request this advice, or if a sponsor is unsure about the SWAT group then they should contact [catherine.blewett@hra.nhs.uk](mailto:catherine.blewett@hra.nhs.uk).

## **Overarching SWAT**

Approval applications (SWAT Groups 2 & 3 only) will be booked directly with the SWAT REC [1] (i.e., not via the e-booking system) during the initial phase and should be notified to [catherine.blewett@hra.nhs.uk](mailto:catherine.blewett@hra.nhs.uk). These studies will likely be suitable for Proportionate Review. Overarching SWAT Group 1 applications submitted solely for the purposes of HRA Assessment will be booked directly with [catherine.blewett@hra.nhs.uk](mailto:catherine.blewett@hra.nhs.uk).

When the overarching SWAT sponsor knows when it plans to submit for review, they should contact [catherine.blewett@hra.nhs.uk](mailto:catherine.blewett@hra.nhs.uk) to arrange for the application to be booked. Please note: due to SWAT applications usually being reviewed by one particular REC, it may be necessary to await the next scheduled meeting for that REC.

When an individual SWAT forming part of an overarching SWAT evaluation is submitted as part of a new host trial application, it should be submitted to a single REC as two separate studies (i.e., the overarching SWAT evaluation and the host trial), making clear that the SWAT is part of the overarching SWAT evaluation. These studies may be submitted to the SWAT REC or to any other REC identified by the host trial sponsor. The overarching SWAT sponsor should contact [catherine.blewett@hra.nhs.uk](mailto:catherine.blewett@hra.nhs.uk) to identify the REC that they intend to submit to so that the HRA can make the necessary arrangements for review.

---

[1] Leeds West REC

The covering letter which is submitted with the application should include the following table:

Table 19: Cover letter table

SWAT Group	Choose an item.
SWAT description	
Type of host trial	
Parameters (e.g., disease area, trial population, participant vulnerabilities)	

### Individual SWAT evaluations done in host trials

Host trial teams doing an individual SWAT evaluation as part of an overarching SWAT only need to consider approvals for Group 2 & 3 SWATs. If needed, these may be submitted as a substantial amendment (where the host trial already has a REC favourable opinion) or alongside a new application (where the proposed host trial itself does not already have a REC favourable opinion).

When an individual SWAT is to be submitted as a substantial amendment, it should be submitted to the REC which issued the favourable opinion to the host trial. The Overarching SWAT team should:

- 1) Complete the amendment tool in the first instance and provide this, along with other relevant documents, to the individual SWAT teams to ensure consistency in classification of the amendment and completion of the amendment tool.
- 2) Notify the HRA, at [catherine.blewett@hra.nhs.uk](mailto:catherine.blewett@hra.nhs.uk) which trials will be submitting amendments to incorporate the SWAT intervention, and therefore to which REC(s) the amendment(s) will be submitted.

The individual SWAT teams should then submit the amendment to the relevant REC(s) for review and notify [catherine.blewett@hra.nhs.uk](mailto:catherine.blewett@hra.nhs.uk) when they have done so.



The cover letter to the host trial REC should include the following:

- The table which was submitted and approved for the overarching SWAT\*
- The IRAS ID & REC Reference of the overarching SWAT
- Name of the REC which issued the favourable opinion for the overarching SWAT
- A description of how the individual SWAT complies with the approved scope of the overarching SWAT

\* The lead of the overarching SWAT will be able to provide much of the information in this bullet list. If the table has not been included in the cover letter, you will be contacted and asked to provide this.

An application for an individual SWAT that forms part of an overarching SWAT evaluation should not be submitted unless the overarching SWAT itself has a favourable opinion. If an application is submitted where the overarching SWAT does not yet have a favourable opinion, you will be contacted and advised that the application should be withdrawn and re-submitted after the overarching SWAT has received a favourable opinion.

### **Further help and guidance**

More help and guidance on SWATs is available from the York Trial Forge Centre ([prometheus-group@york.ac.uk](mailto:prometheus-group@york.ac.uk)), or Trial Forge more generally (<http://trialforge.org>; email [info@trialforge.org](mailto:info@trialforge.org)).

## Glossary

Table 20: Glossary of terms

<i>SWAT</i>	Study Within A Trial. These are also called 'embedded trials', 'nested trials' or 'trials in trials'.
<i>Host trial</i>	A clinical trial in which a SWAT is embedded.
<i>SWAT intervention</i>	The trial process intervention that is to be tested. Examples are financial incentives to improve recruitment, Christmas cards to improve retention or the format of a site initiation visit.
<i>Overarching coordinated SWAT</i>	A SWAT evaluation that is planned from the beginning as one that is done in many host trials at the same time. The evaluation will have a lead researcher (the SWAT evaluation lead) who will approach host trial teams to seek their involvement in the SWAT.
<i>SWAT evaluation lead</i>	The SWAT evaluation lead is responsible for developing the SWAT evaluation protocol and for obtaining overarching approval to evaluate the SWAT intervention.
<i>Host trial team</i>	The team that is embedding an evaluation of the SWAT intervention into their own trial under agreement with the evaluation lead.
<i>REC</i>	Research Ethics Committee
<i>UK SWR</i>	UK Study Wide Review

## Appendix 4

*Draft SWAT reporting guidance*

**Trial Forge Guidance [3 or 4]: A template for reporting the results of randomised Studies Within A Trial (SWATs)**

Table 21: The SWAT reporting template. The text in blue italics is text that could be cut and pasted into a document as the starting point for reporting a SWAT evaluation.

CONSORT item to be included in publication		Additional information and example text shown in italics where possible
<b>Title and Abstract</b>		
<b>1a</b>	Term 'SWAT' should be used in the title	SWAT registry number included if available  <i>SWAT [insert number]: [insert title of SWAT]</i>
<b>1b</b>	Structured summary	Structured using these headings: Background, Methods, Results, Conclusion  Details of host trial included in which the SWAT intervention was evaluated
<b>Introduction; Background and objectives</b>		
<b>2a</b>	Scientific background and explanation of rationale for the SWAT	Justify the need for the SWAT; cite systematic review evidence where appropriate  Replication SWAT: Also cite previous SWAT evaluations undertaken as part of the rationale
<b>2b</b>	Specific objectives or hypotheses for the SWAT	State SWAT question as objective  <i>Does [insert SWAT intervention] increase/decrease [outcome] compared to [comparator] in [participants]?</i>
<b>Methods; Trial design</b>		
<b>3a</b>	Description of the SWAT (such as parallel, factorial, cluster) including allocation ratio	Describe trial design and allocation ratio <i>A [insert number of trial arms and trial design] SWAT was undertaken with an allocation ratio of [insert allocation ratio]</i>
<b>3b</b>	State changes (with reasons) to methods of SWAT following commencement	State where the SWAT protocol is registered <i>The SWAT protocol can be found at [insert details of SWAT repository link]</i>  If SWAT protocol is not registered, include as an appendix.  Host trial:

		<p><i>The host trial protocol is available at [insert details]</i></p> <p>Provide a brief description of host trial using PICO format.</p> <p>SWAT: Use a PICO format to guide description – covered within points 4-6.</p> <p>If changes to the SWAT occurred: <i>The following changes occurred once the SWAT started [insert text]</i></p>
<b>Participants</b>		
<b>4a</b>	State eligibility criteria in SWAT, including differences to those from the host trial	State participant eligibility, this can be tabulated.
<b>4b</b>	Include setting(s) and location(s) where SWAT data was collected	<i>SWAT data were collected in the following settings/locations [insert text]</i>
<b>Interventions</b>		
<b>5</b>	Describe SWAT intervention to enable replication, include how and when interventions were administered and recruitment dates.	Briefly describe the SWAT intervention. Reference to the protocol for further details is acceptable as long as the protocol is available to the reader
<b>Outcomes</b>		
<b>6a</b>	State primary and secondary outcome measure for SWAT. Include how and when they are assessed	<p><i>Primary outcome measure: [insert information including how/when assessed].</i></p> <p><i>Secondary outcome measure(s): [insert information including how/when assessed].</i></p> <p>This information can be tabulated.</p> <p>If appropriate:</p>
<b>6b</b>	Include changes (and reasons) to SWAT outcomes after commencement	<i>The following changes occurred once the SWAT started [insert text].</i>
<b>Sample size</b>		
<b>7a</b>	How sample size was determined for the SWAT.	SWATs are often individually underpowered due to the sample size being constrained by the host trial and a reliable estimate of the effect of the SWAT

		intervention might depend on the aggregation of replicated SWAT evaluations. It is not expected that a formal sample size calculation will always be done.
<b>7b</b>	When applicable, explanation of any interim analyses and stopping guidelines for the SWAT	<p><i>The SWAT sample size was dependent on the host trial [insert host trial details], therefore no formal sample size calculation was performed, which is in line with SWAT methodology<sup>15, 160</sup>. [insert any reasoning for a subsample of the host trial being used – e.g., SWAT was included midway through the trial]</i></p> <p>If interim analyses and/or stopping guidelines were planned: <i>The following interim analyses were planned [state analyses here]. The stopping guidelines were [details here]</i></p>
<b>Randomisation: Sequence generation</b>		
<b>8a</b>	Method used to generate the random allocation sequence for the SWAT	<i>Participants were randomised by [insert method with all methodological details].</i>
<b>8b</b>	Type of randomisation; details of any restriction (such as blocking and block size)	
<b>Allocation concealment mechanism</b>		
<b>9</b>	Mechanism used to implement the random allocation sequence (such as sequentially numbered containers), describing any steps taken to conceal the sequence until interventions were assigned for the SWAT	<i>Allocation concealment was achieved by [insert method].</i>
<b>Implementation</b>		
<b>10</b>	Who generated the random allocation sequence, who enrolled participants, and who assigned participants to interventions for the SWAT	<i>Randomisation was performed by [specify centre or personnel], [specify centre or personal] enrolled participants and [specify centre or personal] assigned the participant to the SWAT intervention or comparator.</i>
<b>Blinding</b>		
<b>11a</b>	If done, who was blinded to the SWAT after assignment to interventions (for example, participants, care providers, those assessing outcomes) and how	Explain who was blinded. <i>Blinding was achieved by [insert details].</i>

<b>11b</b>	If relevant, description of the similarity of the SWAT interventions	
<b>Statistical methods</b>		
12a	Statistical methods used to compare groups for primary and secondary outcomes for the SWAT	All analyses for the SWAT should be pre-planned, ideally detailed in a SWAT Statistical Analysis Plan (SAP), which might be a very short component of the SWAT registry entry. Unless detailed thoroughly and extensively in a publicly available SWAT protocol, the analysis for each outcome should be detailed in the methods of the report.  Alternatively, depending on the journal, the SAP could be uploaded as supplementary material. The SAP and subsequent publication details should include the software used, the statistical methods (including significance level for hypothesis testing), and the population used for the analysis (intention-to-treat or per-protocol).
12b	Methods for additional analyses, such as subgroup analyses and adjusted analyses	
<b>Results</b>		
<b>Participant flow</b>		
13a	For each group, the numbers of participants who were randomly assigned, received intended treatment, and were analysed for the primary outcome for the SWAT	Provide a participant flow diagram that includes this data
13b	For each group participating in the SWAT, losses and exclusions after randomisation, together with reasons	
<b>Recruitment</b>		
14a	Dates defining the periods of recruitment and follow-up of the SWAT	<i>Participant recruitment took place between [insert dates].</i>
14b	Why the SWAT ended or was stopped	If the SWAT ended or was stopped early: <i>The SWAT stopped [recruitment/follow up] early due to [insert text].</i>
<b>Baseline data</b>		
<b>15</b>	A table showing baseline demographic and clinical characteristics for each group	The context of the host trial for each SWAT evaluation is likely to be different and contextual information about the host trial should be provided.

		In addition to general information about the host trial (see 'Methods'), we suggest a table of participant baseline characteristics for those allocated to each group of the SWAT evaluation if these details are available.
<b>Numbers analysed</b>		
16	For each group of the SWAT, number of participants (denominator) included in each analysis and whether the analysis was by original assigned groups	
<b>Outcomes and estimation</b>		
17a	For each primary and secondary outcome, results for each group, and the estimated effect size and its precision (such as 95% confidence interval)	Results should be presented in tables as far as possible rather than only being presented in the body of the text. To facilitate meta-analysis, SWATs should report the actual numbers of participants in each group in the SWAT evaluation.
17b	For binary outcomes, presentation of both absolute and relative effect sizes is recommended	A key element of SWAT evidence is the ability for them to be replicated and an important principle for reporting research is that new findings should be placed in the context of existing, relevant evidence. Therefore, we recommend that an updated meta-analysis is included that presents the results of the current SWAT combined with previous evaluations of the SWAT intervention. Presentation as a cumulative meta-analysis is particularly helpful because it would help to inform judgements about the need for further evaluations of a SWAT intervention. <sup>159</sup>
<b>Ancillary analyses</b>		
18	Results of any other analyses performed on the SWAT data, including subgroup analyses and adjusted analyses, distinguishing pre-specified from exploratory	
<b>Harms</b>		
19	All important harms or unintended effects in each group that took part in the SWAT (for specific guidance see CONSORT for harms)	
<b>Discussion</b>		
<b>Limitations</b>		



20	Trial limitations, addressing sources of potential bias, imprecision, and, if relevant, multiplicity of analyses for the SWAT	
<b>Generalisability</b>		
21	Generalisability (external validity, applicability) of the SWAT findings	
<b>Interpretation</b>		
22	Interpretation consistent with results, balancing benefits and harms, and considering other relevant evidence	In addition to a general interpretation, we suggest that the report includes a brief 'Implications for trial practice' and 'Implications for SWAT research'. These could make use of the cumulative meta-analysis and Trial Forge guidance <sup>156</sup> on whether further evaluations of the intervention are warranted.
<b>Other information</b>		
23	Registration Registration number and name of trial registry	Include the information for both the host trial and SWAT.
24	Protocol Where the full trial protocol can be accessed, if available	
25	Funding Sources of funding and other support (such as supply of drugs), role of funders	
<b>Additional</b>		
	Data sharing	Although not part of Madurasinghe et al, 2016 [6] or the CONSORT guidance, we suggest that authors make the data used to generate their results available as a supplementary file, or through data sharing platforms.
	Registration	It is recommended that SWATs are registered on a repository to ensure all SWATs performed can be included in the evidence and base and support future replication. The following repositories are available to register SWATs: the Northern Ireland Methodology Hub's SWAT repository <sup>32</sup> ( <u>this repository is for SWATs only and encourages replications of registered SWATs</u> ), the ISRCTN trial registry, <sup>158</sup> and the <u>Clinical Trials database</u> . <sup>161</sup> SWATs are usually registered in the two latter repositories as part of the host trial.

