

This is a repository copy of *Vertex Representations for Yangians of Kac-Moody algebras*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/200170/>

Version: Other

Preprint:

Guay, Nicolas, Regelskis, Vidas orcid.org/0000-0002-0092-6917 and Wendlandt, Curtis (2018) Vertex Representations for Yangians of Kac-Moody algebras. [Preprint]

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

VERTEX REPRESENTATIONS FOR YANGIANS OF KAC-MOODY ALGEBRAS

NICOLAS GUAY, VIDAS REGELSKIS, AND CURTIS WENDLANDT

ABSTRACT. Using vertex operators, we build representations of the Yangian of a simply laced Kac-Moody algebra and of its double. As a corollary, we prove the PBW property for simply laced affine Yangians.

CONTENTS

1. Introduction	1
2. The Yangian of \mathfrak{g}	3
3. The centrally extended Yangian double of \mathfrak{g}	7
4. The Lie algebras \mathfrak{s} and \mathfrak{t} as central extensions	11
5. Level one vertex representations	13
6. The Poincaré-Birkhoff-Witt Theorem	23
References	26

1. INTRODUCTION

Vertex operators originate from dual resonance models in theoretical physics. They were used by I. Frenkel and V. Kac in their groundbreaking paper [FrKa] to build an explicit realization of the basic representation of a simply-laced affine Lie algebra. Their work was later extended to non-simply laced affine Lie algebras [BTM, GNOS], to quantum affine algebras [FrJi, Be, JiMi, JKM, Ji3, Ji4, ChJi], to twisted quantum affine algebras and more general quantum Kac-Moody algebras [Ji1, Ji2], to toroidal and quantum toroidal algebras [MRY, Sa], and to Lie superalgebra (e.g. [KSU]).

In this paper, we address the problem of developing an analogue of the work of I. Frenkel and V. Kac for Yangians of simply laced Kac-Moody Lie algebras. Yangians form an important family of quantum algebras which originate from physics, but were first properly defined in general by V. Drinfeld in [Dr]. They can be obtained from quantum loop algebras via a limit procedure [Gu] and it turns out that Yangians and quantum loop algebras become isomorphic after passing to certain completions [GTL1]. The first goal of this paper is to construct representations of Yangians, via their centrally extended doubles (see Definition 3.1), using vertex operators which act on a tensor product of a Fock space with a twisted group algebra (see Theorem 5.5). In the case of the Yangian associated to \mathfrak{sl}_n and \mathfrak{gl}_n , this was done in [IoKo, Io, Kh]. It should be noted that our construction is not a direct consequence of the work of I. Frenkel and N. Jing [FrJi, Ji2] on vertex operator representations of quantum affinizations associated to symmetric Kac-Moody algebras. Indeed, our construction differs in at least one essential way from the one in [FrJi, Ji2], namely that we use a different lattice to build the underlying Fock space.

The second goal of this paper is to prove a version of the Poincaré-Birkhoff-Witt Theorem for affine Yangians of simply laced type (Theorem 6.9) using the vertex representations of Theorem 5.5. For Yangians associated to simple Lie algebras, this theorem was proved in general in [Le], and for classical Lie algebras, a version of the PBW Theorem stated in terms of the RTT-presentation of the Yangian can be found in [Mo] and [AMR]; for affine Yangians, only the type $A^{(1)}$ has been considered before [Gu]. A separate proof of the PBW property for simply laced affine Yangians has been announced in [YaZh2]. The argument in *loc. cit.*, which is of independent interest, uses the existence of a morphism from the Yangian of \mathfrak{g} to the reduced Drinfeld double of the spherical subalgebra of a shuffle algebra associated to \mathfrak{g} [YaZh1, Cor. 3.4].

2010 *Mathematics Subject Classification.* Primary 17B37; Secondary 81R10, 17B69.

Our paper is structured as follows. Section 2 presents the definition of the Yangian $Y(\mathfrak{g})$ associated to a symmetrizable Kac-Moody algebra \mathfrak{g} and describes its classical limit as the enveloping algebra of a certain Lie algebra \mathfrak{s} (Proposition 2.7) which coincides with the current algebra $\mathfrak{g}[t]$ when \mathfrak{g} is finite-dimensional. We also recall results of Section 6 from [GNW] about a certain parameter dependent coproduct (Theorem 2.10) which will be needed in Subsection 6.2 in order to build a faithful representation of $Y(\mathfrak{g})$. It is possible to repeatedly apply this coproduct, but since it is not coassociative, one should proceed with care, as explained at the end of Section 2.

In Section 3, we give the definition of the centrally extended Yangian double $DY^c(\mathfrak{g})$ of \mathfrak{g} and study its basic properties. When \mathfrak{g} is a finite-dimensional simple Lie algebra, a definition of $DY^c(\mathfrak{g})$ was given over twenty year ago in [Kh], where it was conjecturally described as the Hopf algebra double of a central extension of $Y(\mathfrak{g})$. Although this interpretation seems to be limited to that setting, a general definition can be obtained by inserting an arbitrary Cartan matrix into the explicit definition of $DY^c(\mathfrak{g})$ provided in [Kh, DiKh]. This procedure leads to Definition 3.1.

After giving the definition of $DY^c(\mathfrak{g})$ (Definition 3.1, Lemma 3.4), we relate its classical limit to the enveloping algebra of a certain Lie algebra \mathfrak{t} (Proposition 3.6), which in the finite-dimensional setting is just the affine Lie algebra $\mathfrak{g}[t^{\pm 1}] \oplus \mathbb{C}K$ associated to \mathfrak{g} . We conclude Section 3 with Proposition 3.8, which makes precise how the Yangian maps into the centrally extended Yangian double.

The aforementioned Lie algebras \mathfrak{s} and \mathfrak{t} can also be described more explicitly when \mathfrak{g} is an untwisted affine Lie algebra: in this case, they are isomorphic to the universal central extensions of two loop algebras. This fact was proved in [MRY] and Section 4 serves to recall this description. In Proposition 4.7, we show that \mathfrak{s} and \mathfrak{t} can be equivalently characterized as the universal central extensions of $\mathfrak{g}'[t]$ and $\mathfrak{g}'[t^{\pm 1}]$, respectively, where $\mathfrak{g}' = [\mathfrak{g}, \mathfrak{g}]$ is the derived subalgebra of \mathfrak{g} . This description of \mathfrak{s} and \mathfrak{t} is also valid when \mathfrak{g} is finite-dimensional. Our PBW Theorem for $Y(\mathfrak{g})$ (namely, Theorem 6.9) is stated as providing an isomorphism between the associated graded ring of $Y(\mathfrak{g})$ (for a certain filtration) and the enveloping algebra of \mathfrak{s} , so the results of Section 4 are relevant for our second main theorem.

The main section of this paper is Section 5. Assuming that \mathfrak{g} is a simply laced Kac-Moody algebra, we construct a representation of the Yangian double $DY^c(\mathfrak{g})$ (and thus of the Yangian $Y(\mathfrak{g})$) which is given by vertex operators and which factors through the Yangian double at level one (see Theorem 5.5 and also Proposition 5.9 and Corollary 5.10 for slightly different versions of that theorem). This representation can be realized in a space built from the tensor product of a Fock space with the twisted group algebra $\mathbb{C}_\varepsilon[Q]$ of the root lattice Q : see Definition 5.3 and (5.3). Its construction generalizes, and has been motivated by, the results of Iohara [Io] for $\mathfrak{g} = \mathfrak{sl}_N$, as well as the results of [Kh] and [IoKo] which were stated for $\mathfrak{g} = \mathfrak{sl}_2$ and $\mathfrak{g} = \mathfrak{gl}_2$, respectively. By considering carefully a certain filtration, our construction leads to a representation of the Lie algebra \mathfrak{t} (Corollary 5.11) which is related, but not always isomorphic, to the representation of \mathfrak{t} obtained from the classical vertex representation construction [FrKa, MRY]: this is made precise in Proposition 5.17.

The last section contains a proof of the PBW Theorem for affine, simply laced Yangians: see Theorem 6.9. We prove that the associated graded ring of the Yangian $Y(\mathfrak{g})$ (for a certain filtration) is isomorphic to the enveloping algebra $U(\mathfrak{s})$ of \mathfrak{s} . As a consequence, we obtain in Theorem 6.10 that the $\mathbb{C}[\hbar]$ -algebra version of the Yangian $Y_\hbar(\mathfrak{g})$ (see Definition 2.1) is a flat deformation of the enveloping algebra $U(\mathfrak{s})$ of \mathfrak{s} . The main point of the proof of Theorem 6.9 is to show the injectivity of the natural epimorphism from $U(\mathfrak{s})$ to the associated graded ring given in Proposition 2.9: this is accomplished by taking tensor products of the vertex representation of $Y(\mathfrak{g})$ constructed in Section 5 (actually, it is necessary to consider a slightly larger Kac-Moody algebra) and, by using a carefully chosen filtration, reducing the proof to the question of the faithfulness of the corresponding vertex representation of \mathfrak{s} , which was addressed previously in [MRY].

Acknowledgements. The authors thank Yaping Yang and Gufang Zhao for sharing a preliminary version of their proof of the Poincaré-Birkhoff-Witt Theorem for simply-laced affine Yangians using the shuffle algebra approach. The first and third named authors gratefully acknowledge the financial support of the Natural Sciences and Engineering Research Council of Canada provided via the Discovery Grant Program and the Alexander Graham Bell Canada Graduate Scholarships - Doctoral Program, respectively. The second named author acknowledges the financial support of the European Social Fund via grant number 09.3.3-LMT-K-712-02-0017.

2. THE YANGIAN OF \mathfrak{g}

In this section we recall the definition of the Yangian and give some of its basic properties. Let \mathfrak{g} be a symmetrizable Kac-Moody Lie algebra associated to an indecomposable Cartan matrix $\mathbf{A} = (a_{ij})_{i,j \in I}$, where I is the set of vertices of the Dynkin diagram of \mathfrak{g} . We assume that \mathbf{A} satisfies the condition

$$(2.1) \quad \min\{|a_{ij}|, |a_{ji}|\} \leq 1 \quad \forall i, j \in I \text{ with } i \neq j.$$

Though this constraint will not play a role until Section 4, the results of [Ji2, Na], together with Lemma 4.2 and Remark 5.15, suggest that the definition of the Yangian (and its centrally extended double) must be modified in order to extend the vertex representation construction of Section 5 beyond the simply-laced case. Let $(,)$ be a fixed non-degenerate invariant symmetric bilinear form on \mathfrak{g} . We denote by $\{\alpha_i\}_{i \in I}$ the set of simple positive roots. Set

$$d_{ij} = \frac{1}{2}(\alpha_i, \alpha_j) \quad \forall i, j \in I.$$

2.1. Definition of the Yangian.

Definition 2.1. *The Yangian $Y_{\hbar}(\mathfrak{g})$ is the unital associative $\mathbb{C}[\hbar]$ -algebra generated by the elements x_{ir}^{\pm} , h_{ir} , for $i \in I$ and $r \in \mathbb{Z}_{\geq 0}$, subject to the relations*

$$(2.2) \quad [h_{ir}, h_{js}] = 0,$$

$$(2.3) \quad [h_{i0}, x_{js}^{\pm}] = \pm 2d_{ij}x_{js}^{\pm},$$

$$(2.4) \quad [x_{ir}^+, x_{js}^-] = \delta_{ij}h_{i,r+s},$$

$$(2.5) \quad [h_{i,r+1}, x_{js}^{\pm}] - [h_{ir}, x_{j,s+1}^{\pm}] = \pm \hbar d_{ij} (h_{ir}x_{js}^{\pm} + x_{js}^{\pm}h_{ir}),$$

$$(2.6) \quad [x_{i,r+1}^{\pm}, x_{js}^{\pm}] - [x_{ir}^{\pm}, x_{j,s+1}^{\pm}] = \pm \hbar d_{ij} (x_{ir}^{\pm}x_{js}^{\pm} + x_{js}^{\pm}x_{ir}^{\pm}),$$

$$(2.7) \quad \sum_{\sigma \in S_m} [x_{ir_{\sigma(1)}}^{\pm}, [x_{ir_{\sigma(2)}}^{\pm}, \dots, [x_{ir_{\sigma(m)}}^{\pm}, x_{js}^{\pm}] \dots]] = 0 \quad \text{for } i \neq j \text{ and } m = 1 - a_{ij}.$$

Remark 2.2. *In the notation of [GNW], the above algebra is equal to $Y_{\hbar}(\mathfrak{g}')$ where \mathfrak{g}' is the derived subalgebra $[\mathfrak{g}, \mathfrak{g}]$. For the definition of the full Yangian, see Definition 2.1 of [GNW]. For \mathbf{A} of finite or affine type, the condition (2.1) only excludes type $A_1^{(1)}$. In this case the appropriate definition of the Yangian is given in [BeTs, §1.2] and [Ko, Def. 5.1].*

Note that $Y_{\hbar}(\mathfrak{g})$ is generated, as a $\mathbb{C}[\hbar]$ -algebra, by x_{ir}^{\pm} , h_{ir} , for $i \in I$ and $0 \leq r \leq 1$: see [GNW, (2.10)]. We also observe that $Y_{\hbar}(\mathfrak{g})$ is equipped with a $\mathbb{Z}_{\geq 0}$ -grading determined by

$$\deg \hbar = 1 \quad \text{and} \quad \deg x_{ir}^{\pm} = \deg h_{ir} = r \quad \forall i \in I, r \geq 0.$$

We now give an equivalent definition of $Y_{\hbar}(\mathfrak{g})$ in terms of generating series which will prove useful in Section 3. The following result is a translation of [GTL2, Prop. 2.3].

Proposition 2.3 (Prop. 2.3 of [GTL2]). *Let $x_i^{\pm}(z) = \sum_{r \geq 0} x_{ir}^{\pm} z^{-r-1}$ and $h_i(z) = \sum_{r \geq 0} h_{ir} z^{-r-1}$ for each $i \in I$. The defining relations of $Y_{\hbar}(\mathfrak{g})$ are equivalent to*

$$(2.8) \quad h_i(z)h_j(w) = h_j(w)h_i(z),$$

$$(2.9) \quad (z - w \mp \hbar d_{ij})h_i(z)x_j^{\pm}(w) = (z - w \pm \hbar d_{ij})x_j^{\pm}(w)h_i(z) \pm 2d_{ij}x_j^{\pm}(w) - [h_i(z), x_{j0}^{\pm}],$$

$$(2.10) \quad (z - w \mp \hbar d_{ij})x_i^{\pm}(z)x_j^{\pm}(w) = (z - w \pm \hbar d_{ij})x_j^{\pm}(w)x_i^{\pm}(z) + [x_{i0}^{\pm}, x_j^{\pm}(w)] - [x_i^{\pm}(z), x_{j0}^{\pm}],$$

$$(2.11) \quad (z - w)[x_i^+(z), x_j^-(w)] = \delta_{ij}(h_i(w) - h_i(z)),$$

$$(2.12) \quad \sum_{\sigma \in S_m} [x_i^{\pm}(z_{\sigma(1)}), [x_i^{\pm}(z_{\sigma(2)}), \dots, [x_i^{\pm}(z_{\sigma(m)}), x_j^{\pm}(w)] \dots]] = 0,$$

where in the last relation $i \neq j$ and $m = 1 - a_{ij}$.

Remark 2.4. *Multiplying (2.9) by z^{-1} and taking the residue at $z = 0$ gives*

$$[h_{i0}, x_j^{\pm}(w)] = \pm 2d_{ij}x_j^{\pm}(w).$$

Multiplying the relation (2.11) by z^{-1} and taking the residue at $z = 0$ yields

$$(2.13) \quad [x_{i0}^+, x_j^-(w)] = \delta_{ij} h_i(w) \quad \forall i, j \in I.$$

Conversely, we have the following:

Proposition 2.5 (Prop. 3.3(3) of [AG]). *The relation (2.11) is a consequence of (2.8), (2.9) and (2.13).*

For each $\zeta \in \mathbb{C}$, let $Y_\zeta(\mathfrak{g})$ be the \mathbb{C} -algebra generated by $\{x_{ir}^\pm, h_{ir}\}_{i \in I, r \geq 0}$ subject to the defining relations of Definition 2.1 with \hbar replaced by ζ . Equivalently,

$$Y_\zeta(\mathfrak{g}) = Y_\hbar(\mathfrak{g}) / (\hbar - \zeta) Y_\hbar(\mathfrak{g}).$$

For the remainder of this paper our focus will primarily be on the Yangian $Y(\mathfrak{g}) = Y_1(\mathfrak{g})$. The emphasis on the single choice $\zeta = 1$ is justified by the fact that the assignment

$$(2.14) \quad x_{ir}^\pm, h_{ir} \in Y(\mathfrak{g}) \mapsto \zeta^{-r} x_{ir}^\pm, \zeta^{-r} h_{ir} \in Y_\zeta(\mathfrak{g})$$

extends to an isomorphism of algebras $Y(\mathfrak{g}) \rightarrow Y_\zeta(\mathfrak{g})$ for each fixed $\zeta \in \mathbb{C}^\times$. Note that $Y(\mathfrak{g})$ is no longer a $\mathbb{Z}_{\geq 0}$ -graded algebra, but rather a filtered algebra with ascending filtration $\{\mathbf{F}_k\}_{k \geq 0}$ determined by assigning filtration degrees r to x_{ir}^\pm and h_{ir} for each $i \in I$ and $r \geq 0$.

2.2. The classical limit.

Definition 2.6. *Let \mathfrak{s} be the Lie algebra generated by $\{X_{ir}^\pm, H_{ir}\}_{i \in I, r \geq 0}$ subject to the defining relations*

$$(2.15) \quad [H_{ir}, H_{js}] = 0,$$

$$(2.16) \quad [H_{ir}, X_{js}^\pm] = \pm 2d_{ij} X_{j, r+s}^\pm,$$

$$(2.17) \quad [X_{ir}^+, X_{js}^-] = \delta_{ij} H_{i, r+s},$$

$$(2.18) \quad [X_{i, r+1}^\pm, X_{js}^\pm] = [X_{ir}^\pm, X_{j, s+1}^\pm],$$

$$(2.19) \quad \text{ad}(X_{i0}^\pm)^{1-a_{ij}}(X_{jr}^\pm) = 0 \quad \text{for } i \neq j.$$

Note that \mathfrak{s} is a $\mathbb{Z}_{\geq 0}$ -graded Lie algebra with $\deg X_{ir}^\pm = \deg H_{ir} = r$ for all $i \in I$ and $r \geq 0$.

In addition, \mathfrak{s} is always an extension of the current algebra $\mathfrak{g}'[t]$. Indeed, if $\{x_i^\pm, h_i\}_{i \in I}$ denote the Chevalley generators of \mathfrak{g}' , normalized so that $(x_i^+, x_i^-) = 1$ and $h_i = [x_i^+, x_i^-]$, then the assignment

$$(2.20) \quad X_{ir}^\pm, H_{ir} \mapsto x_i^\pm \otimes t^r, h_i \otimes t^r \quad \forall i \in I \text{ and } r \geq 0$$

determines a surjective Lie algebra morphism $\mathfrak{s} \twoheadrightarrow \mathfrak{g}'[t]$. This is an isomorphism when \mathfrak{g} is finite-dimensional, but in general this is not the case. We will consider the situation where \mathfrak{g} is of affine type in more detail in Section 4.

The next proposition illustrates that $Y_\hbar(\mathfrak{g})$ is a graded deformation of the enveloping algebra $U(\mathfrak{s})$.

Proposition 2.7. *The assignment*

$$(2.21) \quad X_{ir}^\pm, H_{ir} \mapsto x_{ir}^\pm, h_{ir} \quad \forall i \in I \text{ and } r \geq 0$$

extends to an isomorphism of graded \mathbb{C} -algebras $U(\mathfrak{s}) \rightarrow Y_0(\mathfrak{g})$.

Proof. That (2.21) extends to an epimorphism of algebras $U(\mathfrak{s}) \twoheadrightarrow Y_0(\mathfrak{g})$ is immediate. Since the defining relations of $Y_0(\mathfrak{g})$ are of Lie type, it is isomorphic to $U(\mathfrak{s}')$, where \mathfrak{s}' is the Lie algebra generated by $\{x_{ir}^\pm, h_{ir}\}_{i \in I, r \geq 0}$ subject to the defining relations (2.15), (2.17), (2.18), in addition to the three relations

$$(2.22) \quad [h_{i0}, x_{js}^\pm] = \pm 2d_{ij} x_{js}^\pm, \quad [h_{i, r+1}, x_{js}^\pm] = [h_{ir}, x_{j, s+1}^\pm],$$

$$(2.23) \quad \sum_{\sigma \in S_m} [x_{ir_{\sigma(1)}}^\pm, [x_{ir_{\sigma(2)}}^\pm, \dots, [x_{i, r_{\sigma(m)}}^\pm, x_{js}^\pm] \dots]] = 0 \quad \text{for } i \neq j \text{ and } m = 1 - a_{ij}.$$

To conclude that the assignment $x_{ir}^\pm, h_{ir} \mapsto X_{ir}^\pm, H_{ir}$ extends to $Y_0(\mathfrak{g}) \rightarrow U(\mathfrak{s})$ which is the inverse of the morphism $U(\mathfrak{s}) \rightarrow Y_0(\mathfrak{g})$ defined by (2.21), it suffices to show that the relations of Definition 2.6 imply (2.22) and (2.23).

Since (2.16) implies (2.22), we are left to deduce (2.23) from Definition 2.6. We will prove the stronger result that (2.16), (2.18) and (2.19) imply

$$(2.24) \quad [X_{i, r_1}^\pm, [X_{i, r_2}^\pm, \dots, [X_{i, r_m}^\pm, X_{js}^\pm] \dots]] = 0$$

for all $s, r_1, \dots, r_m \geq 0$ and $i \neq j \in I$.

From (2.18) and induction we obtain

$$(2.25) \quad [X_{i,r+k}^\pm, X_{j,s}^\pm] = [X_{ir}^\pm, X_{j,s+k}^\pm] \quad \forall r, s, k \geq 0 \text{ and } i, j \in I.$$

This implies that for any fixed $n \geq 0$ and $k, s, r_1, \dots, r_n \in \mathbb{Z}_{\geq 0}$, we have

$$\begin{aligned} \text{ad}(X_{i,r_1+k}^\pm) \text{ad}(X_{i,r_2}^\pm) \cdots \text{ad}(X_{i,r_m}^\pm)(X_{j,s}^\pm) \\ = \text{ad}(X_{i,r_1}^\pm) \text{ad}(X_{i,r_2}^\pm) \cdots \text{ad}(X_{i,r_m}^\pm)(X_{j,s+k}^\pm). \end{aligned}$$

This relation implies that if $k_1, \dots, k_n \geq 0$ satisfy $\sum_{a=1}^n k_a = k$, then

$$(2.26) \quad [X_{ik}^\pm, \text{ad}(X_{i,0}^\pm)^{n-1}(X_{js}^\pm)] = \left[X_{i,k_1}^\pm, \left[X_{i,k_2}^\pm, \dots, \left[X_{i,k_n}^\pm, X_{js}^\pm \right] \cdots \right] \right].$$

Now let $s, r_1, \dots, r_m \geq 0$ be arbitrary and set $k = \sum_{a=1}^m r_a$. Applying $\text{ad}(H_{ik})$ to (2.19) and using (2.16) together with (2.26) twice gives (2.24). \square

Remark 2.8. When \mathfrak{g} is finite-dimensional, it is known that $Y_h(\mathfrak{g})$ is a flat deformation of $U(\mathfrak{s})$. We will prove the analogous result for \mathfrak{g} of simply laced affine type in Theorem 6.10.

Recall the filtration $\{\mathbf{F}_k\}_{k \geq 0}$ on $Y(\mathfrak{g})$ defined at the end of Subsection 2.1. Let \bar{x}_{ir}^\pm and \bar{h}_{ir} denote the images of x_{ir}^\pm and h_{ir} in $\mathbf{F}_r/\mathbf{F}_{r-1} \subset \text{gr } Y(\mathfrak{g})$, where $\mathbf{F}_{-1} = \{0\}$. The following result is immediate from the defining relations of $Y(\mathfrak{g})$.

Proposition 2.9. *The assignment*

$$X_{ir}^\pm, H_{ir} \mapsto \bar{x}_{ir}^\pm, \bar{h}_{ir} \quad \forall i \in I \text{ and } r \geq 0$$

extends to an epimorphism of algebras $\phi : U(\mathfrak{s}) \twoheadrightarrow \text{gr } Y(\mathfrak{g})$.

The statement that ϕ is injective is equivalent to the Poincaré-Birkhoff-Witt Theorem for the Yangian. For \mathfrak{g} of finite type this was proven in the early 1990's by Levendorskii [Le], but in the general setting this remains a conjecture. We will prove the injectivity of ϕ for \mathfrak{g} of simply laced affine type in Section 6.

2.3. The coproduct Δ_u . The Yangian of a finite-dimensional simple Lie algebra is well-known to admit the structure of a Hopf algebra. In particular, it is equipped with a coassociative algebra homomorphism $\Delta : Y(\mathfrak{g}) \rightarrow Y(\mathfrak{g}) \otimes Y(\mathfrak{g})$, its coproduct. When the underlying simple Lie algebra is replaced with a more general Kac-Moody algebra, the formulas used to define Δ are no longer well-defined. However, it was shown in [GNW] that, when \mathfrak{g} is affine, there is an algebra homomorphism $\Delta_u : Y(\mathfrak{g}) \rightarrow (Y(\mathfrak{g}) \otimes Y(\mathfrak{g}))((u))$ which, in a strictly formal sense, has limit at $u = 1$ which is in agreement with Δ . The definition of Δ_u is contained in the following theorem. Set $\tilde{h}_{i1} = h_{i1} - \frac{1}{2}h_{i0}^2$ for all $i \in I$ and $\square(a) = a \otimes 1 + 1 \otimes a$ for all $a \in Y(\mathfrak{g})$.

Theorem 2.10 (Thm. 6.2 of [GNW]). *Assume that the Cartan matrix \mathbf{A} of \mathfrak{g} is of affine type, but not of type $A_1^{(1)}$ or $A_2^{(2)}$. Then there is an algebra homomorphism*

$$\Delta_u : Y(\mathfrak{g}) \rightarrow (Y(\mathfrak{g}) \otimes Y(\mathfrak{g}))((u))$$

uniquely determined by

$$(2.27) \quad \begin{aligned} \Delta_u(x_{i0}^\pm) &= x_{i0}^\pm \otimes 1 + 1 \otimes x_{i0}^\pm u^{\pm 1}, \quad \Delta(h_{i0}) = \square(h_{i0}), \\ \Delta_u(\tilde{h}_{i1}) &= \square(\tilde{h}_{i1}) - \sum_{\alpha \in \Delta_+^{\text{re}}} (\alpha, \alpha_i) x_\alpha^- \otimes x_\alpha^+ u^{\text{ht}(\alpha)}, \end{aligned}$$

for all $i \in I$, where Δ_+^{re} is the set of positive real roots, $\text{ht}(\sum_{i \in I} n_i \alpha_i) = \sum_{i \in I} n_i$, and $x_\alpha^\pm \in \mathfrak{g}_{\pm \alpha}$ are such that $(x_\alpha^+, x_\alpha^-) = 1$.

The morphism Δ_u is not coassociative in the traditional sense, but it satisfies the “twisted” coassociativity relation

$$(2.28) \quad (\Delta_u \otimes \text{id}) \circ \Delta_{uv} = (\text{id} \otimes \Delta_v) \circ \Delta_u.$$

By repeated application of Δ_u it is possible to obtain an algebra morphism $\Delta_u^k : Y(\mathfrak{g}) \rightarrow (Y(\mathfrak{g})^{\otimes(k+1)})(u)$ for each $k \geq 0$. However, due to the presence of the parameter u and the twisted coassociativity property (2.28), this must be handled carefully.

Given an associative unital \mathbb{C} -algebra \mathcal{A} and a fixed $k \geq 1$, we denote by $\mathcal{A}((u_k, \dots, u_1))$ the localization of $\mathcal{A}[[u_k, \dots, u_1]]$ at the multiplicative set

$$S = \{u_k^{m_k} \cdots u_1^{m_1} : m_a \geq 0\}.$$

Equivalently, $\mathcal{A}((u_k, \dots, u_1))$ can be realized as the subspace of $\mathcal{A}[[u_k^{\pm 1}, \dots, u_1^{\pm 1}]]$ consisting of elements

$$\sum_{\ell_1, \dots, \ell_k \in \mathbb{Z}} a_{\ell_k, \dots, \ell_1} u_k^{\ell_k} \cdots u_1^{\ell_1}$$

for which there exists $N \geq 0$ such that $a_{\ell_k, \dots, \ell_1} = 0$ whenever $\ell_m < -N$ for any $1 \leq m \leq k$, with product obtained by extending the usual multiplication of formal series in $\mathcal{A}[[u_k, \dots, u_1]]$. The key feature of this algebra we will exploit is that

$$(2.29) \quad \text{ev}_{u,k} : f(u_k, \dots, u_1) \mapsto f(u, \dots, u) \quad \forall f(u_k, \dots, u_1) \in \mathcal{A}((u_k, \dots, u_1)).$$

determines an algebra homomorphism $\text{ev}_{u,k} : \mathcal{A}((u_k, \dots, u_1)) \rightarrow \mathcal{A}((u))$.

To define Δ_u^k we will make use of auxiliary morphisms

$$\Delta_{u_1, \dots, u_k} : Y(\mathfrak{g}) \rightarrow (Y(\mathfrak{g})^{\otimes(k+1)}((u_k))((u_{k-1})) \cdots ((u_1)))$$

which are defined recursively as follows: $\text{id}^{\otimes(k-1)} \otimes \Delta_{u_k}$ extends to a morphism

$$(Y(\mathfrak{g})^{\otimes k})((u_{k-1})) \cdots ((u_1)) \rightarrow (Y(\mathfrak{g})^{\otimes(k+1)}((u_k))((u_{k-1})) \cdots ((u_1))),$$

and the composition of this morphism with $\Delta_{u_1, \dots, u_{k-1}}$ is precisely Δ_{u_1, \dots, u_k} . Inductively, we find that

$$(2.30) \quad \begin{aligned} \Delta_{u_1, \dots, u_k}(h_{i0}) &= \sum_{a=1}^{k+1} (h_{i0})_a, \quad \Delta_{u_1, \dots, u_k}(x_{i0}^{\pm}) = \sum_{a=1}^{k+1} (x_{i0}^{\pm})_a u_1^{\pm 1} \cdots u_{a-1}^{\pm 1}, \\ \Delta_{u_1, \dots, u_k}(\tilde{h}_{i1}) &= \sum_{a=1}^{k+1} (\tilde{h}_{i1})_a - \sum_{a < b} \sum_{\alpha \in \Delta_+^{\text{re}}} (\alpha, \alpha_i) (x_{\alpha}^-)_a (x_{\alpha}^+)_b u_a^{\text{ht}(\alpha)} \cdots u_{b-1}^{\text{ht}(\alpha)}, \end{aligned}$$

where $(X)_a = 1^{\otimes(a-1)} \otimes X \otimes 1^{\otimes(k+1-a)}$, and the product $u_1^{\pm 1} \cdots u_{a-1}^{\pm 1}$ with $a = 1$ is understood to equal 1. Consequently, $\text{Image}(\Delta_{u_1, \dots, u_k}) \subset Y(\mathfrak{g})^{\otimes(k+1)}((u_k, \dots, u_1))$, and we may therefore set

$$(2.31) \quad \Delta_u^k = \text{ev}_{u,k} \circ \Delta_{u_1, \dots, u_k} : Y(\mathfrak{g}) \rightarrow Y(\mathfrak{g})^{\otimes(k+1)}((u)) \quad \forall k \geq 1,$$

where $\text{ev}_{u,k}$ is as in (2.29) with $\mathcal{A} = Y(\mathfrak{g})^{\otimes(k+1)}$.

The explicit formulas (2.30) imply that Δ_u^k is filtered in the sense that

$$(2.32) \quad \Delta_u^k(\mathbf{F}_{\ell}) \subset (\mathbf{F}_{\ell}(Y(\mathfrak{g})^{\otimes(k+1)}((u))))((u)),$$

where $\mathbf{F}_{\ell}(Y(\mathfrak{g})^{\otimes(k+1)}) = \sum_{a_1 + \dots + a_{k+1} = \ell} \mathbf{F}_{a_1} \otimes \cdots \otimes \mathbf{F}_{a_{k+1}}$. By (2.30), the associated graded morphism $\text{gr } \Delta_u^k$ has image contained in $\text{gr } Y(\mathfrak{g})^{\otimes(k+1)}[u^{\pm 1}]$ for each k :

$$(2.33) \quad \text{gr } \Delta_u^k : \text{gr } Y(\mathfrak{g}) \rightarrow \text{gr } Y(\mathfrak{g})^{\otimes(k+1)}[u^{\pm 1}].$$

The family of filtered morphisms $\{\Delta_u^k\}_{k \geq 1}$ will play a decisive role in the proof of the Poincaré-Birkhoff-Witt Theorem in Section 6, as will the analogous morphisms $\{\Delta_{\mathfrak{s},u}^k\}_{k \geq 1}$ for the enveloping algebra $U(\mathfrak{s})$, which we define now.

Let $\Delta_{\mathfrak{s}}$ denote the standard coproduct on $U(\mathfrak{s})$. The assignment $X_{ir}^{\pm} \mapsto u^{\pm 1} X_{ir}^{\pm}$, $H_{ir} \mapsto H_{ir}$ for all $i \in I$ and $r \geq 0$ extends to an algebra morphism $s_u : U(\mathfrak{s}) \rightarrow U(\mathfrak{s})[u^{\pm 1}]$ of $U(\mathfrak{s})$, and we may set

$$\Delta_{\mathfrak{s},u} = (\text{id} \otimes s_u) \circ \Delta_{\mathfrak{s}} : U(\mathfrak{s}) \rightarrow (U(\mathfrak{s}) \otimes U(\mathfrak{s}))[u^{\pm 1}].$$

The morphisms $\Delta_{\mathfrak{s},u}^k : U(\mathfrak{s}) \rightarrow U(\mathfrak{s})^{\otimes(k+1)}[u^{\pm 1}]$ are now constructed in exactly the same way as Δ_u^k (see (2.32)). On generators, we have

$$\Delta_{\mathfrak{s},u}^k(H_{ir}) = \sum_{a=1}^{k+1} (H_{ir})_a, \quad \Delta_{\mathfrak{s},u}^k(X_{ir}^{\pm}) = \sum_{a=1}^{k+1} (X_{ir}^{\pm})_a u^{\pm(a-1)}.$$

In particular, we have the following commutative diagram:

$$(2.34) \quad \begin{array}{ccc} U(\mathfrak{g}) & \xrightarrow{\Delta_{\mathfrak{g},u}^k} & U(\mathfrak{g})^{\otimes(k+1)}[u^{\pm 1}] \\ \phi \downarrow & & \downarrow \phi^{\otimes(k+1)} \\ \mathrm{gr} Y(\mathfrak{g}) & \xrightarrow{\mathrm{gr} \Delta_u^k} & \mathrm{gr} Y(\mathfrak{g})^{\otimes(k+1)}[u^{\pm 1}] \end{array}$$

The map ϕ is the one given in Proposition 2.9.

3. THE CENTRALLY EXTENDED YANGIAN DOUBLE OF \mathfrak{g}

In this section we introduce the centrally extended Yangian double associated to \mathfrak{g} and study its basic algebraic properties.

3.1. Definition of the Yangian double. Let $\delta(w, z) = \sum_{r \in \mathbb{Z}} w^r z^{-r-1} \in \mathbb{C}[[w^{\pm 1}, z^{\pm 1}]]$ denote the formal delta function. Equivalently,

$$(3.1) \quad \delta(w, z) = \frac{z^{-1}}{1 - z^{-1}w} + \frac{w^{-1}}{1 - w^{-1}z}, \quad \text{where} \quad \frac{x^{-1}}{1 - x^{-1}y} = \sum_{k \geq 0} y^k x^{-k-1}.$$

Definition 3.1. The centrally extended Yangian double $DY_{\hbar}^c(\mathfrak{g})$ is the $\mathbb{C}[\hbar]$ -algebra generated by the coefficients $\{h_{ir}, x_{ir}^{\pm}\}_{i \in I, r \in \mathbb{Z}}$ of

$$x_i^{\pm}(z) = \sum_{r \in \mathbb{Z}} x_{ir}^{\pm} z^{-r-1}, \quad h_i^+(z) = 1 + \hbar \sum_{r \in \mathbb{Z}_{\geq 0}} h_{ir} z^{-r-1}, \quad h_i^-(z) = 1 - \hbar \sum_{r \in \mathbb{Z}_{< 0}} h_{ir} z^{-r-1},$$

for all $i \in I$, together with an element \mathfrak{c} , which are subject to the defining relations

$$(3.2) \quad [\mathfrak{c}, h_i^{\pm}(z)] = 0 = [x_i^{\pm}(z), \mathfrak{c}],$$

$$(3.3) \quad h_i^{\pm}(z) h_j^{\pm}(w) = h_j^{\pm}(w) h_i^{\pm}(z),$$

$$(3.4) \quad \frac{1}{\hbar^2} \left(((z-w)^2 - (\mathfrak{c}_{ij}^-)^2) h_i^+(z) h_j^-(w) - ((z-w)^2 - (\mathfrak{c}_{ij}^+)^2) h_j^-(w) h_i^+(z) \right) = 0,$$

$$(3.5) \quad \frac{1}{\hbar} \left((z-w \mp \mathfrak{c}_{ij}^-) h_i^+(z) x_j^{\pm}(w) - (z-w \pm \mathfrak{c}_{ij}^+) x_j^{\pm}(w) h_i^+(z) \right) = 0,$$

$$(3.6) \quad \frac{1}{\hbar} \left((z-w \mp \hbar d_{ij}) h_i^-(z) x_j^{\pm}(w) - (z-w \pm \hbar d_{ij}) x_j^{\pm}(w) h_i^-(z) \right) = 0,$$

$$(3.7) \quad (z-w \mp \hbar d_{ij}) x_i^{\pm}(z) x_j^{\pm}(w) = (z-w \pm \hbar d_{ij}) x_j^{\pm}(w) x_i^{\pm}(z),$$

$$(3.8) \quad [x_i^+(z), x_j^-(w)] = \frac{\delta_{ij}}{\hbar} \left(\delta(w + \hbar \mathfrak{c}, z) h_i^+(w + \frac{\hbar \mathfrak{c}}{2}) - \delta(w, z) h_i^-(z) \right),$$

$$(3.9) \quad \sum_{\sigma \in S_m} [x_i^{\pm}(z_{\sigma(1)}), [x_i^{\pm}(z_{\sigma(2)}), \dots, [x_i^{\pm}(z_{\sigma(m)}), x_j^{\pm}(w)] \dots]] = 0,$$

where $\mathfrak{c}_{ij}^{\pm} = \hbar d_{ij} \pm \frac{\hbar \mathfrak{c}}{2}$ and in the last relation $i \neq j$ and $m = 1 - a_{ij}$.

For each $\kappa \in \mathbb{C}$, we define the Yangian double at level κ to be the $\mathbb{C}[\hbar]$ -algebra

$$DY_{\hbar}^{\kappa}(\mathfrak{g}) = DY_{\hbar}^c(\mathfrak{g}) / (\mathfrak{c} - \kappa) DY_{\hbar}^c(\mathfrak{g}).$$

Remark 3.2. Even though the relations (3.4)-(3.7) and (3.8) involve negative powers of \hbar , this is not the case for the corresponding relations among the generators. (See Lemma 3.4.)

Remark 3.3. The practice of calling $DY_{\hbar}^c(\mathfrak{g})$ the ‘‘Centrally extended Yangian double’’ is explained by the following: when \mathfrak{g} is finite-dimensional, $DY_{\hbar}^0(\mathfrak{g})$ has been conjectured to be equal, after completion, to the Hopf algebra double of $Y_{\hbar}(\mathfrak{g})$ [KhTo], whereas $DY_{\hbar}^c(\mathfrak{g})$ has been conjecturally described, also after completion, as a quotient of the Hopf algebra double of $Y_{\hbar}(\mathfrak{g}) \otimes \mathbb{C}[c]$ by a derivation [Kh]. These conjectures have been proven for $\mathfrak{g} = \mathfrak{sl}_2$: see [KhTo, Prop. 2.1 (ii)] and [Kh, Thm. 3.1].

Although this interpretation of $DY_{\hbar}^c(\mathfrak{g})$ does not extend beyond the finite case, Definition 3.1 is a natural extension of the definitions found in the literature (see in particular [DiKh, §6] and [Io, Cor. 3.4]).

Lemma 3.4. *For each $i \in I$, set $\tilde{h}_i^\pm(z) = \pm \frac{1}{\hbar}(h_i^\pm(z) - 1)$. Then the relations of Definition 3.1 are equivalent to*

$$(3.10) \quad [\mathbf{c}, \tilde{h}_i^\pm(z)] = 0 = [x_i^\pm(z), \mathbf{c}],$$

$$(3.11) \quad \tilde{h}_i^\pm(z) \tilde{h}_j^\pm(w) = \tilde{h}_j^\pm(w) \tilde{h}_i^\pm(z),$$

$$(3.12) \quad ((z-w)^2 - (\mathbf{c}_{ij}^+)^2) \tilde{h}_j^-(w) \tilde{h}_i^+(z) - ((z-w)^2 - (\mathbf{c}_{ij}^-)^2) \tilde{h}_i^+(z) \tilde{h}_j^-(w) \\ = -2d_{ij}\mathbf{c} + 2\hbar d_{ij}(\tilde{h}_j^-(w) - \tilde{h}_i^+(z))\mathbf{c},$$

$$(3.13) \quad (z-w \mp \mathbf{c}_{ij}^-) \tilde{h}_i^+(z) x_j^\pm(w) - (z-w \pm \mathbf{c}_{ij}^+) x_j^\pm(w) \tilde{h}_i^+(z) = \pm 2d_{ij}x_j^\pm(w),$$

$$(3.14) \quad (z-w \pm \hbar d_{ij}) x_j^\pm(w) \tilde{h}_i^-(z) - (z-w \mp \hbar d_{ij}) \tilde{h}_i^-(z) x_j^\pm(w) = \pm 2d_{ij}x_j^\pm(w),$$

$$(3.15) \quad (z-w \mp \hbar d_{ij}) x_i^\pm(z) x_j^\pm(w) = (z-w \pm \hbar d_{ij}) x_j^\pm(w) x_i^\pm(z),$$

$$(3.16) \quad [x_i^+(z), x_j^-(w)] = \delta_{ij} \frac{1}{\hbar} (\delta(w + \hbar \mathbf{c}, z) - \delta(w, z)) \\ + \delta_{ij} \left(\delta(w + \hbar \mathbf{c}, z) \tilde{h}_i^+(w + \frac{\hbar \mathbf{c}}{2}) + \delta(w, z) \tilde{h}_i^-(z) \right),$$

$$(3.17) \quad \sum_{\sigma \in S_m} [x_i^\pm(z_{\sigma(1)}), [x_i^\pm(z_{\sigma(2)}), \dots, [x_i^\pm(z_{\sigma(m)}), x_j^\pm(w)] \dots]] = 0,$$

where in the last relation $i \neq j$ and $m = 1 - a_{ij}$.

It is not difficult to deduce from these relations that $DY_\hbar^c(\mathfrak{g})$ is a \mathbb{Z} -graded algebra with grading determined by

$$\deg \hbar = 1, \quad \deg \mathbf{c} = 0 \quad \text{and} \quad \deg x_{ir}^\pm = \deg h_{ir} = r \quad \forall i \in I, r \in \mathbb{Z}.$$

Next, for each $\zeta \in \mathbb{C}$ we introduce a \mathbb{C} -algebra

$$DY_\zeta^c(\mathfrak{g}) = DY_\hbar^c(\mathfrak{g}) / (\hbar - \zeta) DY_\hbar^c(\mathfrak{g}),$$

and we abbreviate $DY^c(\mathfrak{g}) = DY_1^c(\mathfrak{g})$. Note that, analogously to $Y_\zeta(\mathfrak{g})$, $DY_\zeta^c(\mathfrak{g})$ for $\zeta \in \mathbb{C}^\times$ is precisely the \mathbb{C} -algebra generated by $\{x_{ir}^\pm, h_{ir}\}_{i \in I, r \in \mathbb{Z}}$ and \mathbf{c} subject to the defining relations of Definition 3.1 with \hbar replaced by ζ . For each $\zeta \in \mathbb{C}^\times$, the assignment

$$x_{ir}^\pm, h_{ir}, \mathbf{c} \in DY^c(\mathfrak{g}) \mapsto \zeta^{-r} x_{ir}^\pm, \zeta^{-r} h_{ir}, \mathbf{c} \in DY_\zeta^c(\mathfrak{g}),$$

or equivalently $x_i^\pm(z), \tilde{h}_i^\pm(z), \mathbf{c} \mapsto \zeta x_i^\pm(\zeta z), \zeta \tilde{h}_i^\pm(\zeta z), \mathbf{c}$, extends to an isomorphism of algebras. With this in mind, we will henceforth focus primarily on the $\mathbb{C}[\hbar]$ -algebra $DY_\hbar^c(\mathfrak{g})$ and the \mathbb{C} -algebra $DY^c(\mathfrak{g})$.

The degree assignments $\deg \mathbf{c} = 0$ and $\deg x_{ir}^\pm = \deg h_{ir} = r$ determine a \mathbb{Z} -filtration (but not gradation) on $DY^c(\mathfrak{g})$. For each $k \in \mathbb{Z}$, let \mathbf{F}_k^D denote the subspace of $DY_\zeta^c(\mathfrak{g})$ spanned by monomials of degree $\leq k$, and let

$$\text{gr}_\mathbb{Z} DY_\zeta^c(\mathfrak{g}) = \bigoplus_{k \in \mathbb{Z}} \mathbf{F}_k^D / \mathbf{F}_{k-1}^D$$

denote the corresponding associated graded algebra.

3.2. Classical limit.

Definition 3.5. *Define \mathfrak{t} to be the Lie algebra generated by an element \mathbf{C} together with the coefficients $\{X_{ir}^\pm, H_{ir}\}_{i \in I, r \in \mathbb{Z}}$ of*

$$X_i^\pm(z) = \sum_{k \in \mathbb{Z}} X_{ik}^\pm z^{-k-1} \quad \text{and} \quad H_i(z) = \sum_{k \in \mathbb{Z}} H_{ik} z^{-k-1} \quad \forall i \in I,$$

which are subject to the defining relations

$$(3.18) \quad [\mathbf{C}, H_i(z)] = 0 = [X_i^\pm(z), \mathbf{C}],$$

$$(3.19) \quad [H_i(z), H_j(w)] = -2d_{ij} \delta_z(z, w) \mathbf{C},$$

$$(3.20) \quad [H_i(z), X_j^\pm(w)] = \pm 2d_{ij} \delta(z, w) X_j^\pm(w),$$

$$(3.21) \quad [X_i^+(z), X_j^-(w)] = \delta_{ij} (\delta(z, w) H_i(z) - \delta_z(z, w) \mathbf{C}),$$

$$(3.22) \quad (z-w)[X_i^\pm(z), X_j^\pm(w)] = 0,$$

$$(3.23) \quad \text{ad}(X_{i0}^\pm)^{1-a_{ij}}(X_j^\pm(z)) = 0,$$

where $\delta_z(z, w) = \frac{d}{dz}\delta(z, w) = \sum_{r \in \mathbb{Z}} r z^{r-1} w^{-r-1}$.

Note that the degree assignment

$$\deg \mathbf{C} = 0 \quad \text{and} \quad \deg X_{ir}^\pm = \deg H_{ir} = r \quad \forall i \in I, r \in \mathbb{Z}$$

makes \mathfrak{t} into a \mathbb{Z} -graded Lie algebra. Additionally, just as \mathfrak{s} is an extension of the current algebra $\mathfrak{g}'[t]$, the Lie algebra \mathfrak{t} is an extension of the loop algebra $\mathfrak{g}'[t^{\pm 1}]$. In the notation of (2.20), the assignment

$$X_{ir}^\pm, H_{ir} \mapsto x_i^\pm \otimes t^r, h_i \otimes t^r \quad \forall i \in I \text{ and } r \in \mathbb{Z}$$

defines a surjective Lie algebra morphism $\mathfrak{t} \rightarrow \mathfrak{g}'[t^{\pm 1}]$. We will return to this observation in Section 4.

We now give the analogue of Proposition 2.7 for the Yangian double. Let $\tilde{h}_i^\pm(z)$ be as in Lemma 3.4 (now viewed in $DY_0^c(\mathfrak{g})[[z^{\pm 1}]]$) and set $\tilde{h}_i(z) = \tilde{h}_i^+(z) + \tilde{h}_i^-(z) = \sum_{r \in \mathbb{Z}} h_{ir} z^{-r-1}$.

Proposition 3.6. *The assignment*

$$(3.24) \quad X_i^\pm(z), H_i(z), \mathbf{C} \mapsto x_i^\pm(z), \tilde{h}_i(z), \mathbf{c} \quad \forall i \in I$$

extends to a graded isomorphism of \mathbb{C} -algebras $U(\mathfrak{t}) \rightarrow DY_0^c(\mathfrak{g})$.

Proof. By definition, $DY_0^c(\mathfrak{g})$ is the \mathbb{C} -algebra generated by the coefficients of $x_i^\pm(z)$, $\tilde{h}_i(z)$ and the central element \mathbf{c} , which are subject to the relations of Definition 3.1 with \hbar replaced by 0. Lemma 3.4 implies that, in addition to the centrality of \mathbf{c} , these relations are

$$(3.25) \quad \tilde{h}_i^\pm(z) \tilde{h}_j^\pm(w) = \tilde{h}_j^\pm(w) \tilde{h}_i^\pm(z),$$

$$(3.26) \quad (z-w)^2 [\tilde{h}_j^-(w), \tilde{h}_i^+(z)] = -2d_{ij}\mathbf{c}$$

$$(3.27) \quad (z-w) [\tilde{h}_i^+(z), x_j^\pm(w)] = \pm 2d_{ij} x_j^\pm(w),$$

$$(3.28) \quad (z-w) [x_j^\pm(w), \tilde{h}_i^\pm(z)] = \pm 2d_{ij} x_j^\pm(w),$$

$$(3.29) \quad (z-w) [x_i^\pm(z), x_j^\pm(w)] = 0,$$

$$(3.30) \quad [x_i^+(z), x_j^-(w)] = \delta_{ij} \lim_{\hbar \rightarrow 0} \frac{1}{\hbar} (\delta(w + \hbar \mathbf{c}, z) - \delta(w, z)) + \delta_{ij} \delta(w, z) \tilde{h}_i(w),$$

$$(3.31) \quad \sum_{\sigma \in S_m} [x_i^\pm(z_{\sigma(1)}), [x_i^\pm(z_{\sigma(2)}), \dots, [x_i^\pm(z_{\sigma(m)}), x_j^\pm(w)] \dots]] = 0.$$

It therefore suffices to show that these relations are equivalent to the defining relations of Definition 3.5 (with (H_{ir}, X_{ir}^\pm) replaced by (h_{ir}, x_{ir}^\pm) and \mathbf{C} by \mathbf{c}).

Step 1: ((3.25), (3.26)) \iff (3.19).

Multiplying (3.26) by $\sum_{k \geq 0} (k+1) w^k z^{-k-2} = \left(\frac{z^{-1}}{1-z^{-1}w} \right)^2$ yields the relation

$$[\tilde{h}_j^-(w), \tilde{h}_i^+(z)] = -2d_{ij} \left(\frac{z^{-1}}{1-z^{-1}w} \right)^2 \mathbf{c}.$$

Combining this with (3.25) and using $\tilde{h}_i(z) = \tilde{h}_i^+(z) + \tilde{h}_i^-(z)$, we obtain

$$[\tilde{h}_j(w), \tilde{h}_i(z)] = -2d_{ij} \left(\left(\frac{w^{-1}}{1-w^{-1}z} \right)^2 - \left(\frac{z^{-1}}{1-z^{-1}w} \right)^2 \right) \mathbf{c} = 2d_{ij} \delta_z(z, w) \mathbf{c},$$

where we have used the identity (3.1). Switching $i \leftrightarrow j$ and $z \leftrightarrow w$ yields (3.19).

Conversely, taking the $z^{-r-1} w^{-s-1}$ coefficient of (3.19) gives

$$(3.32) \quad [h_{ir}, h_{js}] = 2rd_{ij} \delta_{r,-s} \mathbf{c}.$$

Multiplying both sides by $z^{-r-1} w^{-s-1}$ and taking the sum separately over $r, s \geq 0$ and $r, s < 0$ gives (3.25).

Switching i and j in (3.32), multiplying both sides by $w^{-r-1} z^{-s-1}$ and taking the sum over $(r, s) \in \mathbb{Z}_{<0} \times \mathbb{Z}_{\geq 0}$ yields

$$[\tilde{h}_j^-(w), \tilde{h}_i^+(z)] = -2d_{ij} \left(\frac{z^{-1}}{1-z^{-1}w} \right)^2 \mathbf{c}.$$

Multiplying both sides by $(z - w)^2$ gives (3.26).

Step 2: ((3.27),(3.28)) \iff (3.20).

Multiplying (3.27) by $\sum_{k \geq 0} w^k z^{-k-1}$ and (3.28) by $\sum_{k \geq 0} z^k w^{-k-1}$ gives

$$[\tilde{h}_i^+(z), x_j^\pm(w)] = \pm \frac{2d_{ij}z^{-1}}{1 - z^{-1}w} x_j^\pm(w) \quad \text{and} \quad [\tilde{h}_i^-(z), x_j^\pm(w)] = \pm \frac{2d_{ij}w^{-1}}{1 - w^{-1}z} x_j^\pm(w).$$

Adding these two relations together gives (3.20). Conversely, taking the $z^{-r-1}w^{-s-1}$ coefficient of (3.20) yields

$$[h_{ir}, x_{js}^\pm] = \pm 2d_{ij} x_{j,r+s}^\pm.$$

Multiplying both sides of this equality by $(z - w)z^{-r-1}w^{-s-1}$ and taking the sum $r \geq 0$ and $s \in \mathbb{Z}$ gives

$$(z - w)[\tilde{h}_i^+(z), x_j^\pm(w)] = \pm 2d_{ij} \sum_{r \geq 0, s \in \mathbb{Z}} x_{j,r+s}^\pm (z^{-r}w^{-s-1} - z^{-r-1}w^{-s}) = \pm 2d_{ij} x_j^\pm(w),$$

which is precisely (3.27). The proof that (3.20) implies (3.28) is similar.

Step 3: (3.30) \iff (3.21), and (3.29) \iff (3.22).

The equivalence of (3.29) with (3.22) is immediate. To prove the (3.30) \iff (3.21), it suffices to show that

$$\lim_{\hbar \rightarrow 0} \frac{1}{\hbar} (\delta(w + \hbar \mathbf{c}, z) - \delta(w, z)) = \delta_w(w, z) \mathbf{c} = -\delta_z(z, w) \mathbf{c},$$

which can be verified directly.

Step 4: (3.31) \implies (3.23), and ((3.20),(3.22),(3.23)) \implies (3.31).

The first implication is obvious. The second implication is proven in the same way as its \mathfrak{s} -analogue in Proposition 2.7. \square

Recall the filtration $\{\mathbf{F}_k^D\}_{k \in \mathbb{Z}}$ defined at the end of Subsection 3.1. Let $\bar{x}_{ir}^\pm, \bar{h}_{ir}$ denote the images of x_{ir}^\pm, h_{ir} in $\mathbf{F}_r^D / \mathbf{F}_{r-1}^D$ and $\bar{\mathbf{c}}$ denote the image of \mathbf{c} in $\mathbf{F}_0^D / \mathbf{F}_{-1}^D$.

Similar verifications to those carried out in the proof of the previous proposition allow us to deduce the following analogue of Proposition 2.9.

Proposition 3.7. *The assignment*

$$X_{ir}^\pm, H_{ir}, \mathbf{C} \mapsto \bar{x}_{ir}^\pm, \bar{h}_{ir}, \bar{\mathbf{c}} \quad \forall i \in I \text{ and } r \in \mathbb{Z}$$

extends to an epimorphism of algebras $\phi_D : U(\mathfrak{t}) \twoheadrightarrow \text{gr}_{\mathbb{Z}} DY^c(\mathfrak{g})$.

Like the epimorphism $\phi : U(\mathfrak{s}) \twoheadrightarrow \text{gr } Y(\mathfrak{g})$ of Proposition 2.9, we expect ϕ_D to be an isomorphism for general \mathfrak{g} . However, the injectivity of ϕ_D will not be considered in this paper.

3.3. From the Yangian to its double. We conclude this section by offering a more precise relation between $Y_{\hbar}(\mathfrak{g})$ and $DY_{\hbar}^c(\mathfrak{g})$. Let $\mathbf{x}_i^\pm(z) = \sum_{r \geq 0} x_{ir}^\pm z^{-r-1} \in DY_{\hbar}^c(\mathfrak{g})[[z^{-1}]]$ for each $i \in I$.

Proposition 3.8. *The assignment*

$$(3.33) \quad \iota_{\hbar} : x_i^\pm(z), h_i(z) \mapsto \mathbf{x}_i^\pm(z \pm \frac{\hbar \mathbf{c}}{2}), \tilde{h}_i^\pm(z) \quad \forall i \in I$$

extends to a morphism of $\mathbb{C}[\hbar]$ -algebras $\iota_{\hbar} : Y_{\hbar}(\mathfrak{g}) \mapsto DY_{\hbar}^c(\mathfrak{g})$. The composition of ι_{\hbar} with the projection $DY_{\hbar}^c(\mathfrak{g}) \twoheadrightarrow DY^c(\mathfrak{g})$ induces a morphism of \mathbb{C} -algebras $\iota : Y(\mathfrak{g}) \rightarrow DY^c(\mathfrak{g})$.

Proof. To distinguish between the generating series of $Y_{\hbar}(\mathfrak{g})$ and $DY_{\hbar}^c(\mathfrak{g})$, we will temporarily denote the series $x_i^\pm(z) \in Y_{\hbar}(\mathfrak{g})[[z^{-1}]]$ from Proposition 2.3 by $\mathcal{X}_i^\pm(z)$. We will prove that ι_{\hbar} preserves the defining relations of $Y_{\hbar}(\mathfrak{g})$ provided by Proposition 2.3.

It is immediate that the relations (3.3) and (3.9) imply that $\tilde{h}_i^\pm(z)$ and $\tilde{\mathbf{x}}_i^\pm(z) = \mathbf{x}_i^\pm(z \pm \frac{\hbar \mathbf{c}}{2})$ satisfy the defining relations (2.8) and (2.12), respectively, of $Y_{\hbar}(\mathfrak{g})$ (with $h_i(z)$ replaced by $\tilde{h}_i^\pm(z)$ and $\mathcal{X}_i^\pm(z)$ replaced by $\tilde{\mathbf{x}}_i^\pm(z)$).

Multiplying (3.5) by z^{-1} and taking the residue at $z = 0$ gives $[h_{i0}, x_j^\pm(w)] = \pm 2d_{ij} x_j^\pm(w)$, and thus

$$(3.34) \quad [h_{i0}, \tilde{\mathbf{x}}_j^\pm(w)] = \pm 2d_{ij} \tilde{\mathbf{x}}_j^\pm(w).$$

Taking instead the $z^{-r-1}w^{-s-1}$ coefficient of (3.5), we obtain

$$[h_{i,r+1}, x_{js}^\pm] - [h_{ir}, x_{j,s+1}^\pm] = \pm (\mathbf{c}_{ij}^- h_{ir} x_{js}^\pm + \mathbf{c}_{ij}^+ x_{js}^\pm h_{ir}).$$

Multiplying both sides by $z^{-r-1}w^{-s-1}$, taking the sum over $r, s \geq 0$, we obtain

$$(z - w \mp c_{ij}^-) \tilde{h}_i^+(z) \mathbf{x}_j^\pm(w) - (z - w \pm c_{ij}^+) \mathbf{x}_j^\pm(w) \tilde{h}_i^+(z) = [h_{i0}, \mathbf{x}_j^\pm(w)] - [\tilde{h}_i^+(z), x_{j0}^\pm].$$

Substituting in the relation (3.34) and applying $w \mapsto w \pm \frac{\hbar c}{2}$ yields (2.9).

The proof that (3.7) implies (2.10) with $\mathcal{X}_i^\pm(z)$ and $\mathcal{X}_j^\pm(w)$ replaced by $\tilde{\mathbf{x}}_i^\pm(z)$ and $\tilde{\mathbf{x}}_j^\pm(w)$, respectively, is similar and will be omitted.

It thus remains to see that the assignment (3.33) preserves the relation (2.11). By Proposition 2.5, it suffices to prove

$$(3.35) \quad [x_{i0}^+, \tilde{\mathbf{x}}_j^-(w)] = \delta_{ij} \tilde{h}_i^+(w) \quad \forall i, j \in I.$$

Taking the residue of (3.16) at $z = 0$ gives

$$(3.36) \quad [x_{i0}^+, x_j^-(w)] = \delta_{ij} \left(\tilde{h}_i^+(w + \frac{\hbar c}{2}) + \tilde{h}_i^-(w) \right),$$

where we have used that $\delta(z, w) \tilde{h}_i^-(z) = \delta(z, w) \tilde{h}_i^-(w)$. The relation (3.35) follows directly from this identity.

The proof is concluded by noting that the second statement of the proposition is an immediate consequence of the first. \square

Observe that ι_{\hbar} (resp. ι) is a graded (resp. filtered) homomorphism. We conjecture that both ι_{\hbar} and ι are injective.

4. THE LIE ALGEBRAS \mathfrak{s} AND \mathfrak{t} AS CENTRAL EXTENSIONS

In Sections 2 and 3 it was noted that the Lie algebras \mathfrak{s} and \mathfrak{t} (see Definitions 2.6 and 3.5) are always extensions of $\mathfrak{g}'[t]$ and $\mathfrak{g}'[t^{\pm 1}]$, respectively. In this section we employ the results of [MRY] to deduce that, when \mathfrak{g} is of untwisted affine type, \mathfrak{s} and \mathfrak{t} are in fact isomorphic to the universal central extensions of $\mathfrak{g}'[t]$ and $\mathfrak{g}'[t^{\pm 1}]$, respectively.

Let \mathfrak{g}_0 be the underlying finite-dimensional, simple Lie algebra of the untwisted affine Lie algebra \mathfrak{g} . We specify the indexing set I to be $\{0, 1, \dots, \ell\}$, the extending vertex of the Dynkin diagram of \mathfrak{g} being labeled by 0. Let \mathcal{A} be a commutative, associative \mathbb{C} -algebra. Then $\mathfrak{g}_0 \otimes_{\mathbb{C}} \mathcal{A}$ is a Lie algebra in a natural way. Denote by $\Omega^1(\mathcal{A})$ the module of Khähler differentials of \mathcal{A} , and let $d\mathcal{A}$ denote the subspace of exact forms (see, for instance, [MRY, §2]).

Theorem 4.1 ([K1], Theorem 3.3). *The Lie algebra $\mathfrak{g}_0 \otimes_{\mathbb{C}} \mathcal{A}$ admits a universal central extension $\mathbf{uce}(\mathfrak{g}_0 \otimes_{\mathbb{C}} \mathcal{A})$ defined by*

$$\mathbf{uce}(\mathfrak{g}_0 \otimes_{\mathbb{C}} \mathcal{A}) = (\mathfrak{g}_0 \otimes_{\mathbb{C}} \mathcal{A}) \oplus \Omega^1(\mathcal{A})/d\mathcal{A}$$

as a vector space, with Lie bracket such that $\Omega^1(\mathcal{A})/d\mathcal{A}$ is central and

$$[X_1 \otimes a, X_2 \otimes b] = [X_1, X_2] \otimes ab + (X_1, X_2) \cdot b(da) \quad \forall X_1, X_2 \in \mathfrak{g}_0 \text{ and } a, b \in \mathcal{A}.$$

We will be interested in the choices $\mathcal{A} = \mathbb{C}[t^{\pm 1}, t_2]$ and $\mathcal{A} = \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}]$. Set

$$\mathfrak{g}_0[t_1^{\pm 1}, t_2] = \mathfrak{g}_0 \otimes_{\mathbb{C}} \mathbb{C}[t_1^{\pm 1}, t_2] \quad \text{and} \quad \mathfrak{g}_0[t_1^{\pm 1}, t_2^{\pm 1}] = \mathfrak{g}_0 \otimes_{\mathbb{C}} \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}].$$

As in [MRY, (3.1)], we let $\mathfrak{t}(\mathbf{A})$ denote the Lie algebra obtained from Definition 3.5 by replacing the defining relation (3.22) with

$$(4.1) \quad [X_i^\pm(z), X_i^\pm(w)] = 0 \quad \forall i \in I.$$

It was proven in [MRY] that, in fact, $\mathfrak{t}(\mathbf{A}) \cong \mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2^{\pm 1}])$. The following lemma asserts that $\mathfrak{t}(\mathbf{A})$ coincides with \mathfrak{t} , and hence that \mathfrak{t} can also be identified with $\mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2^{\pm 1}])$, as will be stated more precisely in Proposition 4.4.

Lemma 4.2. *Assume that \mathfrak{g} is a symmetrizable Kac-Moody Lie algebra with indecomposable Cartan matrix $\mathbf{A} = (a_{ij})_{i,j \in I}$ satisfying the condition (2.1). Then, in the Lie algebra \mathfrak{t} , the relation (3.22) implies the relation (4.1). Conversely, the relations (3.20), (3.21), (3.23) and (4.1) imply that (3.22) holds for all $i, j \in I$. In particular, if \mathfrak{g} is of untwisted affine type (excluding $A_1^{(1)}$), $\mathfrak{t} \cong \mathfrak{t}(\mathbf{A})$.*

Proof. We first prove the implication (3.22) \implies (4.1). The relation (3.22) with $i = j$ implies that there is $A_i(w) \in \mathfrak{t}[[w^\pm]]$ such that

$$[X_i^\pm(z), X_i^\pm(w)] = \delta(z, w)A_i(w).$$

Since the right-hand side is symmetric in w and z and the left-hand side is antisymmetric, both sides must be zero, and hence (4.1) holds.

To prove that ((3.20), (3.21), (3.23), (4.1)) \implies (3.20), we make a few preliminary observations. By taking the residue at $w = 0$ of (3.20) and then also of the relation obtained from (3.20) by exchanging z and w , we arrive at the identity

$$(4.2) \quad [H_i(z), X_{j0}^\pm] = \pm 2d_{ij}X_j^\pm(z) = [H_{i0}, X_j^\pm(z)] \quad \forall i, j \in I.$$

Similarly, from (3.21) we obtain

$$(4.3) \quad [X_{i0}^\mp, X_j^\pm(w)] = \mp \delta_{ij}H_i(w) \quad \forall i, j \in I.$$

Now fix $i, j \in I$ with $i \neq j$. If $a_{ij} = 0$, then (3.23) is the relation $[X_{i0}^\pm, X_j^\pm(w)] = 0$. After applying $\text{ad}(H_i(z))$ to this equation and employing (4.2) and (3.20), it becomes

$$\pm 2d_{ii}[X_i^\pm(z), X_j^\pm(w)] \pm 2d_{ij}\delta(z, w)[X_{i0}^\pm, X_j^\pm(w)] = \pm 2d_{ii}[X_i^\pm(z), X_j^\pm(w)] = 0,$$

which gives (3.22).

If $a_{ij} \neq 0$, then without loss of generality we may assume that $a_{ij} = -1$. The Serre relation (3.23) then reads as $[X_{i0}^\pm, [X_{i0}^\pm, X_j^\pm(w)]] = 0$. Applying $\text{ad}(H_i(z))$ to both sides of this equation, we find that

$$\pm 4d_{ii}[X_{i0}^\pm(z), [X_{i0}^\pm, X_j^\pm(w)]] \pm 2d_{ij}\delta(z, w)[X_{i0}^\pm, [X_{i0}^\pm, X_j^\pm(w)]] = 0,$$

where we have used (3.20), (4.1) and (4.2). Hence, we have

$$[X_i^\pm(z), [X_{i0}^\pm, X_j^\pm(w)]] = 0.$$

Acting on this identity by $\text{ad}(X_{i0}^\mp)$ and employing (3.20) together with (4.2) and (4.3), we deduce that

$$2(d_{ii} + d_{ij})[X_i^\pm(z), X_j^\pm(w)] = -2d_{ij}\delta(z, w)[X_{i0}^\pm, X_j^\pm(w)].$$

By assumption, $-1 = a_{ij} = 2\frac{d_{ij}}{d_{ii}}$, and hence $d_{ii} \neq -d_{ij}$. Multiplying the above equation by $(2d_{ii} + 2d_{ij})^{-1}(z - w)$ therefore produces the relation (3.22). \square

Remark 4.3. The generators X_{ir}^\pm , H_{ir} and \mathbf{C} of \mathfrak{t} are related to the generators $x_r(\pm\alpha_i)$, $\alpha_i^\vee(r)$ and \mathbf{c} of $\mathfrak{t}(\mathbf{A})$ given in [MRY, (3.1)] by

$$X_{ir}^\pm = d_{ii}^{-1/2}x_r(\pm\alpha_i), \quad H_{ir} = d_{ii}^{-1}\alpha_i^\vee(r) \quad \text{and} \quad \mathbf{C} = \mathbf{c}.$$

In order to describe the isomorphism $\mathfrak{t} \cong \mathfrak{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2^{\pm 1}])$ and its \mathfrak{s} -analogue, we will need a more explicit description of $\Omega^1(\mathcal{A})/d\mathcal{A}$ when $\mathcal{A} = \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}]$ or $\mathbb{C}[t_1^{\pm 1}, t_2]$. By [MRY, §2], $\Omega^1(\mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}])/d(\mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}])$ has basis

$$B_t = \{t_1^{-1}dt_1, t_1^k t_2^\ell dt_1, t_1^k t_2^{-1}dt_2 : k \in \mathbb{Z}, \ell \in \mathbb{Z}^\times\}.$$

Similarly, one finds that $\Omega^1(\mathbb{C}[t_1^{\pm 1}, t_2])/d(\mathbb{C}[t_1^{\pm 1}, t_2])$ has basis $B_s \subset B_t$ given by

$$B_s = \{t_1^{-1}dt_1, t_1^k t_2^\ell dt_1 : k \in \mathbb{Z}, \ell \in \mathbb{Z}_{>0}\}.$$

Note that these observations, coupled with Theorem 4.1, imply that $\mathfrak{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2]) \subset \mathfrak{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2^{\pm 1}])$ as a Lie subalgebra. Let $\{X_i^\pm, H_i\}_{i=1}^\ell$ be the Chevalley generators for \mathfrak{g}_0 normalized so that $(X_i^+, X_i^-) = 1$ and $H_i = [X_i^+, X_i^-]$. Let $X_{\pm\theta}$ be root vectors of \mathfrak{g}_0 for the roots $\pm\theta$ normalized so that $(X_\theta, X_{-\theta}) = 1$, where θ is the highest root of \mathfrak{g}_0 . Set $H_\theta = [X_{-\theta}, X_\theta]$.

Proposition 4.4 (Prop. 3.5 of [MRY]). *The assignment $\{X_{ir}^\pm, H_{ir}, \mathbf{C}\}_{i \in I, r \in \mathbb{Z}} \rightarrow \mathfrak{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2^{\pm 1}])$ given by*

$$\begin{aligned} \mathbf{C} &\mapsto t_2^{-1}dt_2, \\ X_{ir}^\pm &\mapsto X_i^\pm \otimes t_2^r, \quad i = 1, \dots, \ell, \\ X_{0r}^\pm &\mapsto X_{\mp\theta} \otimes t_1^{\pm 1}t_2^r, \\ H_{ir} &\mapsto H_i \otimes t_2^r, \quad i = 1, \dots, \ell, \\ H_{0r} &\mapsto H_\theta \otimes t_2^r + t_2^r t_1^{-1}dt_1, \end{aligned}$$

extends to an isomorphism of Lie algebras $\mathfrak{t} \rightarrow \mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2^{\pm 1}])$. Moreover, we have $\mathfrak{s} \cong \mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2])$ with an isomorphism $\mathfrak{s} \rightarrow \mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2])$ given by the above assignment with r taking values in $\mathbb{Z}_{\geq 0}$ and \mathbf{C} omitted.

Remark 4.5. Although the second part of the above proposition (concerning $\mathfrak{s} \cong \mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2])$) was not stated in [MRY, Prop. 3.5], it can be proven in the same way as the first part.

Corollary 4.6. Assume that \mathfrak{g} is of untwisted affine type (excluding $A_1^{(1)}$). Then the natural morphism

$$\mathfrak{s} \rightarrow \mathfrak{t}, \quad X_{ir}^{\pm}, H_{ir} \mapsto X_{ir}^{\pm}, H_{ir} \quad \forall i \in I \text{ and } r \geq 0$$

is an embedding of Lie algebras.

Because of the following proposition, it is also possible to interpret \mathfrak{s} and \mathfrak{t} as universal central extensions of $\mathfrak{g}'[t]$ and $\mathfrak{g}'[t^{\pm 1}]$.

Proposition 4.7. We have isomorphisms of Lie algebras

$$\mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2]) \cong \mathbf{uce}(\mathfrak{g}'[t]) \quad \text{and} \quad \mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2^{\pm 1}]) \cong \mathbf{uce}(\mathfrak{g}'[t^{\pm 1}]).$$

Proof. We begin by noting that, since $\mathfrak{g}'[t]$ and $\mathfrak{g}'[t^{\pm 1}]$ are perfect Lie algebras, the universal central extensions $\mathbf{uce}(\mathfrak{g}'[t])$ and $\mathbf{uce}(\mathfrak{g}'[t^{\pm 1}])$ do in fact exist (see [Ne, Thm. 1.14]).

Since \mathfrak{g} is an untwisted affine Lie algebra, $\mathfrak{g}' \cong \mathfrak{g}_0[t_1^{\pm 1}] \oplus \mathbf{C}K$ with Lie bracket determined by $[K, \mathfrak{g}'] = 0$ and

$$[X_1 \otimes t_1^r, X_2 \otimes t_1^s] = [X_1, X_2] \otimes t_1^{r+s} + r\delta_{r,-s}(X_1, X_2)K$$

for all $X_1, X_2 \in \mathfrak{g}_0$ and $r, s \in \mathbb{Z}$. It follows that $\mathfrak{g}'[t_2] \cong \mathfrak{g}_0[t_1^{\pm 1}, t_2] \oplus \mathbb{C}[t_2]K$ is a central extension of $\mathfrak{g}_0[t_1^{\pm 1}, t_2]$ with natural projection $\pi : \mathfrak{g}'[t] \twoheadrightarrow \mathfrak{g}_0[t_1^{\pm 1}, t_2]$. Let ψ denote the projection $\mathbf{uce}(\mathfrak{g}'[t_2]) \twoheadrightarrow \mathfrak{g}'[t_2]$. Then, by [Ne, Cor. 1.9], $\mathbf{uce}(\mathfrak{g}'[t_2])$ is a universal central extension of $\mathfrak{g}_0[t_1^{\pm 1}, t_2]$ with projection $\pi \circ \psi : \mathbf{uce}(\mathfrak{g}'[t_2]) \twoheadrightarrow \mathfrak{g}_0[t_1^{\pm 1}, t_2]$. This proves that $\mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2]) \cong \mathbf{uce}(\mathfrak{g}'[t_2])$. Replacing t_2 by $t_2^{\pm 1}$, we obtain instead $\mathbf{uce}(\mathfrak{g}_0[t_1^{\pm 1}, t_2^{\pm 1}]) \cong \mathbf{uce}(\mathfrak{g}'[t_2^{\pm 1}])$. \square

5. LEVEL ONE VERTEX REPRESENTATIONS

We now fix \mathfrak{g} to be a simply laced Kac-Moody Lie algebra, and we let $Q = \bigoplus_{i \in I} \mathbb{Z}\alpha_i$ denote the root lattice associated to \mathfrak{g} . In addition, we normalize the invariant form (\cdot, \cdot) so that $(\alpha_i, \alpha_i) = 2$ for all $i \in I$.

In this section, we construct representations of $DY_h^c(\mathfrak{g})$ and $DY^c(\mathfrak{g})$ which are given by vertex operators and which factor through $DY_h^1(\mathfrak{g})$ and $DY^1(\mathfrak{g})$. The main results pertaining to this construction are given in Subsections 5.1 and 5.2.

The vertex operators which define these representations are themselves built from operators arising from the action of a Heisenberg Lie algebra on its Fock space representation. Accordingly, we begin by introducing the appropriate Heisenberg algebra, its Fock space representation, as well as the auxiliary operators which play a central role in our construction.

Definition 5.1. The Heisenberg algebra \mathfrak{H} is the Lie algebra with basis given by the elements $\mathcal{H}_{ir}, \mathcal{C}$ for $i \in I, r \in \mathbb{Z} \setminus \{0\}$ and with the bracket given by

$$[\mathcal{H}_{ir}, \mathcal{C}] = 0, \quad \forall i \in I, \quad \forall r \in \mathbb{Z} \setminus \{0\}, \quad [\mathcal{H}_{ir}, \mathcal{H}_{js}] = r\delta_{rs}\delta_{ij}\mathcal{C}, \quad \forall i, j \in I, \quad \forall r, s \in \mathbb{Z} \setminus \{0\}.$$

Remark 5.2. This is not the usual definition of the Heisenberg algebra associated to Q (see Definition 5.13): rather, it is the Heisenberg algebra associated to the trivial lattice $\mathbb{Z}^{|I|}$.

The polynomial ring $\mathbb{C}[\mathcal{H}_{i,-r}]_{i \in I, r > 0}$ can be equipped with the structure of an \mathfrak{H} -module by defining

$$\mathcal{H}_{j,-s}(f) = \mathcal{H}_{j,-s}f, \quad \mathcal{C}(f) = f, \quad \mathcal{H}_{js}(f) = s \frac{\partial}{\partial \mathcal{H}_{j,-s}}(f) \quad \forall f \in \mathbb{C}[\mathcal{H}_{i,-r}]_{i \in I, r > 0}, \quad j \in I \text{ and } s > 0,$$

yielding the so-called *Fock space* representation of \mathfrak{H} .

Next, fix a bimultiplicative function $\varepsilon : Q \times Q \rightarrow \mathbb{Z}/2\mathbb{Z} = \{\pm 1\}$ satisfying the condition

$$(5.1) \quad \varepsilon(\alpha, \alpha) = (-1)^{\frac{1}{2}(\alpha, \alpha)} \quad \forall \alpha \in Q.$$

The bimultiplicativity of ε implies that $\varepsilon(\alpha, 0) = 1$ for all $\alpha \in Q$, while (5.1) implies that

$$(5.2) \quad \varepsilon(\alpha, \beta) = (-1)^{(\alpha, \beta)} \varepsilon(\beta, \alpha) \quad \forall \alpha, \beta \in Q.$$

The bimultiplicativity of ε also implies that it is a 2-cocycle of Q with values in $\mathbb{Z}/2\mathbb{Z}$, and thus it determines a central extension $\tilde{Q} = \mathbb{Z}/2\mathbb{Z} \times_{\varepsilon} Q$ of Q by $\mathbb{Z}/2\mathbb{Z}$ which is equal to $\mathbb{Z}/2\mathbb{Z} \times Q$ as a set, and has product

$$(\epsilon_a, \alpha)(\epsilon_b, \beta) = (\varepsilon(\alpha, \beta)\epsilon_a\epsilon_b, \alpha + \beta) \quad \forall \alpha, \beta \in Q \text{ and } \epsilon_a, \epsilon_b \in \mathbb{Z}/2\mathbb{Z}.$$

Definition 5.3. Let \mathcal{I} be the two-sided ideal of $\mathbb{C}[\tilde{Q}]$ which is spanned by $e^{(\epsilon_a, \alpha)} - \epsilon_a e^{(1, \alpha)}$ for all $\alpha \in Q$ and $\epsilon_a \in \mathbb{Z}/2\mathbb{Z}$. The twisted group algebra $\mathbb{C}_{\varepsilon}[Q]$ is defined to be the quotient $\mathbb{C}[\tilde{Q}]/\mathcal{I}$.

Since the \mathbb{C} -linear projection $\mathbb{C}[\tilde{Q}] \rightarrow \mathbb{C}[Q]$, $e^{(\epsilon_a, \alpha)} \mapsto \epsilon_a e^{\alpha}$ induces an isomorphism of vector spaces $\mathbb{C}_{\varepsilon}[Q] \rightarrow \mathbb{C}[Q]$, $\mathbb{C}_{\varepsilon}[Q]$ can be equivalently defined as the \mathbb{C} -algebra with basis $\{e^{\alpha}\}_{\alpha \in Q}$ and multiplication given by

$$e^{\alpha} \cdot e^{\beta} = \varepsilon(\alpha, \beta) e^{\alpha + \beta} \quad \forall \alpha, \beta \in Q.$$

Remark 5.4. By [FLM, Proposition 5.2.3], the condition (5.2) determines ε up to equivalence of cocycles, and hence it determines the central extension \tilde{Q} of Q by $\mathbb{Z}/2\mathbb{Z}$ up to isomorphism. In particular, this implies that any two bimultiplicative functions $\varepsilon, \varepsilon'$ satisfying (5.1) will determine the same twisted group algebra up to isomorphism.

The existence of $\varepsilon : Q \times Q \rightarrow \mathbb{Z}/2\mathbb{Z}$ satisfying (5.1) can be established in various ways: see for instance [Ka, §7.8].

Viewing $\mathbb{C}_{\varepsilon}[Q]$ as a left-module over itself, we can form the $U(\mathfrak{H}) \otimes \mathbb{C}_{\varepsilon}[Q]$ -module

$$(5.3) \quad \mathcal{V} = \mathbb{C}[\mathcal{H}_{i, -r}]_{i \in I, r > 0} \otimes \mathbb{C}_{\varepsilon}[Q].$$

We also define an auxiliary family of operators $\{\partial_{\alpha}\}_{\alpha \in Q} \subset \text{End}_{\mathbb{C}} \mathcal{V}$ by

$$\partial_{\alpha}(f \otimes e^{\beta}) = (\alpha, \beta) f \otimes e^{\beta} \quad \forall f \in \mathbb{C}[\mathcal{H}_{i, -r}]_{i \in I, r > 0} \text{ and } \alpha, \beta \in Q.$$

5.1. The $DY_{\hbar}^c(\mathfrak{g})$ -module $\mathcal{V}[[\hbar]]$. We first construct a vertex representation of $DY_{\hbar}^c(\mathfrak{g})$ on the topologically free $\mathbb{C}[[\hbar]]$ -module $\mathcal{V}[[\hbar]]$. The actions of $U(\mathfrak{H}) \otimes \mathbb{C}_{\varepsilon}[Q]$ and of ∂_{α} on \mathcal{V} defined above naturally extend to $\mathcal{V}[[\hbar]]$.

For each $i \in I$, let $N(i)$ denote the set of vertices to which i is connected, i.e. the set of *neighbours* of the vertex i . Define $A_i^{\pm}(z)$ and $B_i^{\pm}(z)$, for each $i \in I$, by

$$A_i^{\pm}(z) = \exp \left(\pm \sum_{r > 0} \frac{\mathcal{H}_{i, -r}}{r} (z^r + (z \mp \hbar)^r) \mp \sum_{r > 0} \sum_{j \in N(i)} \frac{\mathcal{H}_{j, -r}}{r} (z \mp \frac{\hbar}{2})^r \right),$$

$$B_i^{\pm}(z) = \exp \left(\mp \sum_{r > 0} \frac{\mathcal{H}_{ir}}{r} z^{-r} \right).$$

Inspired by [Io], we define the vertex operators $\mathbf{X}_i^{\pm}(z), \mathbf{H}_i^{\pm}(z) \in (\text{End}_{\mathbb{C}[[\hbar]]} \mathcal{V}[[\hbar]])[[z^{\pm 1}]]$, for each $i \in I$, by

$$(5.4) \quad \mathbf{X}_i^{\pm}(z) = \pm A_i^{\pm}(z) B_i^{\pm}(z) e^{\pm \alpha_i} z^{\partial_{\pm \alpha_i}},$$

$$(5.5) \quad \mathbf{H}_i^{+}(z) = B_i^{+}(z + \frac{\hbar}{2}) B_i^{-}(z - \frac{\hbar}{2}) \left(\frac{1 + \frac{\hbar}{2} z^{-1}}{1 - \frac{\hbar}{2} z^{-1}} \right)^{\partial_{\alpha_i}},$$

$$(5.6) \quad \mathbf{H}_i^{-}(z) = A_i^{+}(z) A_i^{-}(z).$$

where, for each $\alpha \in Q$, $z^{\partial_{\alpha}} \in (\text{End}_{\mathbb{C}[[\hbar]]} \mathcal{V}[[\hbar]])[[z^{\pm 1}]]$ is defined on \mathcal{V} by

$$z^{\partial_{\alpha}}(f \otimes e^{\beta}) = z^{(\alpha, \beta)} f \otimes e^{\beta} \quad \forall \beta \in Q \text{ and } f \in \mathbb{C}[\mathcal{H}_{i, -r}]_{i \in I, r > 0}.$$

Equivalently, $z^{\partial_{\alpha}} = \sum_{k \in \mathbb{Z}} P_{\alpha, k} z^k$ with $P_{\alpha, k}(f \otimes e^{\beta}) = \delta_{k, (\alpha, \beta)} f \otimes e^{\beta}$.

Let us explain why $\left(\frac{1 + \frac{\hbar}{2} z^{-1}}{1 - \frac{\hbar}{2} z^{-1}} \right)^{\partial_{\alpha_i}}$, and thus $\mathbf{H}_i^{+}(z)$, belongs to $(\text{End}_{\mathbb{C}[[\hbar]]} \mathcal{V}[[\hbar]])[[z^{-1}]]$. For each invertible series $g(z) \in (\mathbb{C}[[\hbar]])[[z^{-1}]]$, the operator $g(z)^{\partial_{\alpha}}$ (defined on \mathcal{V} by $g(z)^{\partial_{\alpha}}(f \otimes e^{\beta}) = g(z)^{(\alpha, \beta)} f \otimes e^{\beta}$) can be viewed as an element of $(\text{End}_{\mathbb{C}[[\hbar]]} \mathcal{V}[[\hbar]])[[z^{-1}]]$. To see this, first write

$$\sum_{k \in \mathbb{Z}} P_{\alpha, k} g(z)^k = \sum_{k \geq 0} P_{\alpha, k} g(z)^k + \sum_{k < 0} P_{\alpha, -k} (g(z)^{-1})^k.$$

For each $r \geq 0$, the z^{-r} coefficient of $\sum_{k \geq 0} P_{\alpha, k} g(z)^k$ is an infinite sum of the form $\sum_{k \geq 0} a_k(\hbar) P_{\alpha, k}$ with $a_k(\hbar) \in \mathbb{C}[[\hbar]]$. The sum $\sum_{k \geq 0} a_k(\hbar) P_{\alpha, k}$ is a well-defined element of $\text{End}_{\mathbb{C}[[\hbar]]} \mathcal{V}[[\hbar]]$ since, for any fixed $\beta \in Q$,

$P_{\alpha,k}(e^\beta) = 0$ for all but at most one value of k . This implies that $\sum_{k \geq 0} P_{\alpha,k} g(z)^k \in (\text{End}_{\mathbb{C}[[\hbar]]} \mathcal{V}[[\hbar]])[[z^{-1}]]$. The same reasoning can be applied to $\sum_{k > 0} P_{\alpha,-k} (g(z)^{-1})^k$.

With the vertex operators $\{\mathbf{X}_i^\pm(z), \mathbf{H}_i^\pm(z)\}_{i \in I}$ at our disposal, we can now state the main theorem of this section.

Theorem 5.5. *The assignment*

$$(5.7) \quad x_i^\pm(z) \mapsto \mathbf{X}_i^\pm(z), \quad h_i^\pm(z) \mapsto \mathbf{H}_i^\pm(z) \quad \forall i \in I, \quad \mathbf{c} \mapsto 1$$

extends to a homomorphism of $\mathbb{C}[[\hbar]]$ -algebras $\rho_\hbar : DY_\hbar^c(\mathfrak{g}) \rightarrow \text{End}_{\mathbb{C}[[\hbar]]} \mathcal{V}[[\hbar]]$.

The next lemma will be employed to prove this theorem. Let

$$\Gamma_i^\pm(z) = \exp \left(\mp \sum_{r > 0} \frac{\mathcal{H}_{i,\pm r}}{r} z^{\mp r} \right).$$

Lemma 5.6. *Let $\chi_i : I \rightarrow \{0, 1\}$ denote the indicator function of $N(i)$, i.e. $\chi_i(j) = 1$ if $j \in N(i)$ and $\chi_i(j) = 0$ otherwise. Then, for each pair of indices $i, j \in I$, we have*

$$(5.8) \quad \Gamma_i^-(z) \Gamma_i^+(w) = \Gamma_i^+(w) \Gamma_i^-(z) \left(1 - \frac{z}{w}\right)^{-1},$$

$$(5.9) \quad [A_i^\pm(z), A_j^\pm(w)] = [A_i^\pm(z), A_j^\mp(w)] = 0 = [B_i^\pm(z), B_j^\mp(w)] = [B_i^\pm(z), B_j^\pm(w)],$$

$$(5.10) \quad B_i^\pm(z) A_j^\pm(w) = \frac{(1 - z^{-1}w)^{\delta_{ij}} (1 - z^{-1}(w \mp \hbar))^{\delta_{ij}}}{(1 - z^{-1}(w \mp \frac{\hbar}{2}))^{\chi_j(i)}} A_j^\pm(w) B_i^\pm(z),$$

$$(5.11) \quad B_i^\pm(z) A_j^\mp(w) = \frac{(1 - z^{-1}(w \pm \frac{\hbar}{2}))^{\chi_j(i)}}{(1 - z^{-1}w)^{\delta_{ij}} (1 - z^{-1}(w \pm \hbar))^{\delta_{ij}}} A_j^\mp(w) B_i^\pm(z).$$

Proof. Relations of the form (5.8) appear often in the literature: see for instance the proof of Theorem 14.8 in [Ka] and the proof of Proposition 2.9 (a) in [FrKa]. It follows from the fact that

$$\exp(A) \exp(B) = \exp(B) \exp(A) \exp([A, B])$$

for any two operators A and B such that $[A, [A, B]] = 0 = [B, [A, B]]$, together with the relation

$$\left[\sum_{r > 0} \frac{\mathcal{H}_{i,-r}}{r} z^r, \sum_{s > 0} \frac{\mathcal{H}_{j,s}}{s} w^{-s} \right] = \sum_{r,s > 0} \frac{[\mathcal{H}_{i,-r}, \mathcal{H}_{j,s}]}{rs} z^r w^{-s} = -\delta_{ij} \sum_{s > 0} \frac{1}{s} \left(\frac{z}{w} \right)^s = \delta_{ij} \ln \left(1 - \frac{z}{w} \right).$$

The relation (5.9) is immediate from the definition of the operators $A_i^\pm(z)$ and $B_i^\pm(z)$, while (5.10) is a straightforward application of (5.8). The relation (5.11) is a consequence of (5.10) since $B_i^\pm(z) = B_i^\mp(z)^{-1}$. \square

We will also need the following identity, which can be deduced immediately from the definition of $g(z)^{\partial_\alpha}$:

$$g(z)^{\partial_\alpha} e^\beta = e^\beta g(z)^{(\alpha, \beta)} g(z)^{\partial_\alpha} \quad \forall \alpha, \beta \in Q \quad \text{and} \quad g(z) = z \text{ or } g(z) \in ((\mathbb{C}[[\hbar]])[[z^{-1}]])^\times.$$

Proof of Theorem 5.5. The proof is achieved using standard vertex operator calculus. We will prove that the relations of Definition 3.1 are preserved by the assignment (5.7). This is immediate for (3.2), and for (3.3) this is a consequence of the relation (5.9) of Lemma 5.6. The other relations require more elaborate use of Lemma 5.6 and we will treat them independently. Set

$$z_\pm = z \pm \frac{\hbar}{2}.$$

The relation (3.4). Let $c_{ij}^\pm = \frac{\hbar}{2}((\alpha_i, \alpha_j) \pm 1)$ denote the image of \mathbf{c}_{ij}^\pm under (5.7). Then

$$\begin{aligned} \mathbf{H}_i^+(z) \mathbf{H}_j^-(w) &= B_i^+(z_+) B_i^-(z_-) A_j^+(w) A_j^-(w) \left(\frac{z_+}{z_-} \right)^{\partial_{\alpha_i}} \\ &= \frac{(1 - z_-^{-1} w_-)^{\chi_j(i)}}{(1 - z_-^{-1} w)^{\delta_{ij}} (1 - z_-^{-1} (w - \hbar))^{\delta_{ij}}} B_i^+(z_+) A_j^+(w) B_i^-(z_-) A_j^-(w) \left(\frac{z_+}{z_-} \right)^{\partial_{\alpha_i}} \\ &= \frac{(1 - z_-^{-1} w_-)^{\chi_j(i)} (1 - z_+^{-1} w)^{\delta_{ij}} (1 - z_+^{-1} (w - \hbar))^{\delta_{ij}}}{(1 - z_-^{-1} w)^{\delta_{ij}} (1 - z_-^{-1} (w - \hbar))^{\delta_{ij}} (1 - z_+^{-1} w_-)^{\chi_j(i)}} A_j^+(w) B_i^+(z_+) B_i^-(z_-) A_j^-(w) \left(\frac{z_+}{z_-} \right)^{\partial_{\alpha_i}}. \end{aligned}$$

On the other hand, we have

$$\begin{aligned}
\mathbf{H}_j^-(w)\mathbf{H}_i^+(z) &= A_j^+(w)A_j^-(w)B_i^+(z_+)B_i^-(z_-) \left(\frac{z_+}{z_-} \right)^{\partial_{\alpha_i}} \\
&= \frac{(1 - z_+^{-1}w)^{\delta_{ij}}(1 - z_+^{-1}(w + \hbar))^{\delta_{ij}}}{(1 - z_+^{-1}w_+)^{\chi_j(i)}} A_j^+(w)B_i^+(z_+)A_j^-(w)B_i^-(z_-) \left(\frac{z_+}{z_-} \right)^{\partial_{\alpha_i}} \\
&= \frac{(1 - z_+^{-1}w)^{\delta_{ij}}(1 - z_+^{-1}(w + \hbar))^{\delta_{ij}}(1 - z_-^{-1}w_+)^{\chi_j(i)}}{(1 - z_+^{-1}w_+)^{\chi_j(i)}(1 - z_-^{-1}w)^{\delta_{ij}}(1 - z_-^{-1}(w + \hbar))^{\delta_{ij}}} A_j^+(w)B_i^+(z_+)B_i^-(z_-)A_j^-(w) \left(\frac{z_+}{z_-} \right)^{\partial_{\alpha_i}}.
\end{aligned}$$

Therefore, since $\mathcal{V}[\hbar]$ is torsion free, the assignment (5.7) will preserve (3.4) provided

$$\begin{aligned}
&((z - w)^2 - (c_{ij}^-)^2) \frac{(1 - z_-^{-1}w_-)^{\chi_j(i)}(1 - z_+^{-1}w)^{\delta_{ij}}(1 - z_+^{-1}(w - \hbar))^{\delta_{ij}}}{(1 - z_-^{-1}w)^{\delta_{ij}}(1 - z_-^{-1}(w - \hbar))^{\delta_{ij}}(1 - z_+^{-1}w_-)^{\chi_j(i)}} \\
&= ((z - w)^2 - (c_{ij}^+)^2) \frac{(1 - z_+^{-1}w)^{\delta_{ij}}(1 - z_+^{-1}(w + \hbar))^{\delta_{ij}}(1 - z_-^{-1}w_+)^{\chi_j(i)}}{(1 - z_+^{-1}w_+)^{\chi_j(i)}(1 - z_-^{-1}w)^{\delta_{ij}}(1 - z_-^{-1}(w + \hbar))^{\delta_{ij}}}.
\end{aligned}$$

for all $i, j \in I$. This can be checked directly using

$$(5.12) \quad c_{ij}^\pm = \begin{cases} \pm \frac{\hbar}{2} & \text{if } i \neq j, \chi_j(i) = 0, \\ -\frac{\hbar}{2} \pm \frac{\hbar}{2} & \text{if } i \neq j, \chi_j(i) = 1, \\ \hbar \pm \frac{\hbar}{2} & \text{if } i = j. \end{cases}$$

The relation (3.5). Making use of (5.1) together with Lemma 5.6, we deduce that

$$\begin{aligned}
\mathbf{H}_i^+(z)\mathbf{X}_j^\pm(w) &= \pm B_i^+(z_+)B_i^-(z_-)A_j^\pm(w)B_j^\pm(w) \left(\frac{z_+}{z_-} \right)^{\partial_{\alpha_i}} e^{\pm\alpha_j} w^{\partial_{\pm\alpha_j}} \\
&= \pm \frac{(1 - z_-^{-1}w_\mp)^{\pm\chi_j(i)} \left(\frac{z_+}{z_-} \right)^{\pm(\alpha_i, \alpha_j)}}{(1 - z_-^{-1}w)^{\pm\delta_{ij}}(1 - z_-^{-1}(w \mp \hbar))^{\pm\delta_{ij}}} B_i^+(z_+)A_j^\pm(w)B_i^-(z_-)B_j^\pm(w) e^{\pm\alpha_j} w^{\partial_{\pm\alpha_j}} \left(\frac{z_+}{z_-} \right)^{\partial_{\alpha_i}} \\
&= \frac{(1 - z_-^{-1}w_\mp)^{\pm\chi_j(i)}(1 - z_+^{-1}w)^{\pm\delta_{ij}}(1 - z_+^{-1}(w \mp \hbar))^{\pm\delta_{ij}}}{(1 - z_-^{-1}w)^{\pm\delta_{ij}}(1 - z_-^{-1}(w \mp \hbar))^{\pm\delta_{ij}}(1 - z_+^{-1}w_\mp)^{\pm\chi_j(i)}} \left(\frac{z_+}{z_-} \right)^{\pm(\alpha_i, \alpha_j)} \mathbf{X}_j^\pm(w)\mathbf{H}_i^+(z).
\end{aligned}$$

Therefore, the assignment (5.7) will preserve the relation (3.5) if the following identity holds:

$$(z - w \pm c_{ij}^\pm) = (z - w \mp c_{ij}^\mp) \frac{(1 - z_-^{-1}w_\mp)^{\pm\chi_j(i)}(1 - z_+^{-1}w)^{\pm\delta_{ij}}(1 - z_+^{-1}(w \mp \hbar))^{\pm\delta_{ij}}}{(1 - z_-^{-1}w)^{\pm\delta_{ij}}(1 - z_-^{-1}(w \mp \hbar))^{\pm\delta_{ij}}(1 - z_+^{-1}w_\mp)^{\pm\chi_j(i)}} \left(\frac{z_+}{z_-} \right)^{\pm(\alpha_i, \alpha_j)}.$$

This is easily verified using (5.12). If $i \neq j$ and $\chi_j(i) = 0$ then this is clear. If $i \neq j$ and $\chi_j(i) = 1$, then the right-hand side equals

$$(z - w \pm \hbar) \left(\frac{1 - z_-^{-1}w_\mp}{1 - z_+^{-1}w_\mp} \right)^{\pm 1} \left(\frac{z_-}{z_+} \right)^{\pm 1} = (z - w \pm \hbar) \left(\frac{z - w - \frac{\hbar}{2} \pm \frac{\hbar}{2}}{z - w + \frac{\hbar}{2} \pm \frac{\hbar}{2}} \right)^{\pm 1} = z - w,$$

which is the left-hand side. If $i = j$, then $c_{ij}^+ = \frac{3\hbar}{2}$ and $c_{ij}^- = \frac{\hbar}{2}$, and the right-hand side of the equality is

$$(z - w \mp \frac{\hbar}{2}) \left(\frac{(z_+ - w)(z_+ - w \pm \hbar)}{(z_- - w \pm \hbar)(z_- - w)} \right)^{\pm 1} = (z - w \mp \frac{\hbar}{2}) \left(\frac{(z - w + \frac{\hbar}{2})(z - w + \frac{\hbar}{2} \pm \hbar)}{(z - w - \frac{\hbar}{2} \pm \hbar)(z - w - \frac{\hbar}{2})} \right)^{\pm 1} = z - w \pm \frac{3\hbar}{2}.$$

Note that in neglecting the factor of \hbar^{-1} which appears in (3.5), we have made use of the fact that $\mathcal{V}[\hbar]$ is torsion free.

The relation (3.6). Applying again the relations of Lemma 5.6, we obtain

$$\mathbf{H}_i^-(z)\mathbf{X}_j^\pm(w) = \pm A_j^\pm(w)A_i^+(z)A_i^-(z)B_j^\pm(w)e^{\pm\alpha_j}w^{\partial_{\pm\alpha_j}}$$

$$\begin{aligned}
 &= \pm \frac{(1-w^{-1}z)^{\pm\delta_{ij}}(1-w^{-1}(z+\hbar))^{\pm\delta_{ij}}}{(1-w^{-1}z_{\pm})^{\pm\chi_j(i)}} A_j^{\pm}(w) A_i^{\pm}(z) B_j^{\pm}(w) A_i^{\mp}(z) e^{\pm\alpha_j} w^{\partial_{\pm\alpha_j}} \\
 &= \frac{(1-w^{-1}(z+\hbar))^{\pm\delta_{ij}}(1-w^{-1}z_{\pm})^{\pm\chi_j(i)}}{(1-w^{-1}z_{\pm})^{\pm\chi_j(i)}(1-w^{-1}(z-\hbar))^{\pm\delta_{ij}}} \mathbf{X}_j^{\pm}(w) \mathbf{H}_i^{\mp}(z) \\
 &= \left(\frac{1-w^{-1}(z \pm \hbar d_{ij})}{1-w^{-1}(z \mp \hbar d_{ij})} \right) \mathbf{X}_j^{\pm}(w) \mathbf{H}_i^{\mp}(z).
 \end{aligned}$$

(The last equality is obtained by considering the two cases $\chi_j(i) = 1$ and $i = j$.) Multiplying both sides by $(w - z \pm \hbar d_{ij})$ and using the fact that $\mathcal{V}[[\hbar]]$ is torsion free, we find that the relation (3.6) is preserved by the assignment (5.7).

To prove that the remaining three relations of Definition 3.1 are satisfied by $\{\mathbf{X}_i^{\pm}(z), \mathbf{H}_i^{\pm}(z)\}_{i \in I}$, we introduce the following normal ordering: given a finite integer $n \in \mathbb{Z}_{>0}$ together with collections $i_a \in I$ and $\epsilon_a \in \{\pm\}$ for each $1 \leq a \leq n$, set

$$:\mathbf{X}_{i_1}^{\epsilon_1}(z_1) \cdots \mathbf{X}_{i_n}^{\epsilon_n}(z_n): = \left(\prod_{b=1}^n \epsilon_b \right) A_{i_1}^{\epsilon_1}(z_1) \cdots A_{i_n}^{\epsilon_n}(z_n) B_{i_1}^{\epsilon_1}(z_1) \cdots B_{i_n}^{\epsilon_n}(z_n) e^{\epsilon_1 \alpha_{i_1} + \cdots + \epsilon_n \alpha_{i_n}} z_1^{\partial_{\epsilon_1 \alpha_{i_1}}} \cdots z_n^{\partial_{\epsilon_n \alpha_{i_n}}}.$$

Note that with this definition, (5.9) implies that $:\mathbf{X}_{i_1}^{\epsilon_1}(z_1) \cdots \mathbf{X}_{i_n}^{\epsilon_n}(z_n): =: \mathbf{X}_{i_{\sigma(1)}}^{\epsilon_{\sigma(1)}}(z_{\sigma(1)}) \cdots \mathbf{X}_{i_{\sigma(n)}}^{\epsilon_{\sigma(n)}}(z_{\sigma(n)}):$ for each permutation $\sigma \in S_n$.

By (5.1) and Lemma 5.6, we have

$$(5.13) \quad \mathbf{X}_i^{\pm}(z) \mathbf{X}_j^{\mp}(w) = \varepsilon(\alpha_i, \alpha_j) \frac{(1-z^{-1}w_{\pm})^{\chi_j(i)}}{(1-z^{-1}w)^{\delta_{ij}}(1-z^{-1}(w \pm \hbar))^{\delta_{ij} z(\alpha_i, \alpha_j)}} : \mathbf{X}_i^{\pm}(z) \mathbf{X}_j^{\mp}(w) :,$$

$$(5.14) \quad \mathbf{X}_i^{\pm}(z) \mathbf{X}_j^{\pm}(w) = \varepsilon(\alpha_i, \alpha_j) \frac{(1-z^{-1}w)^{\delta_{ij}}(1-z^{-1}(w \mp \hbar))^{\delta_{ij}}}{(1-z^{-1}w_{\mp})^{\chi_j(i)} z^{-(\alpha_i, \alpha_j)}} : \mathbf{X}_i^{\pm}(z) \mathbf{X}_j^{\pm}(w) :,$$

where we have used the fact that $\varepsilon(\pm\alpha, \mp\beta) = \varepsilon(\alpha, \beta) = \varepsilon(\pm\alpha, \pm\beta)$ for all $\alpha, \beta \in Q$.

The relation (3.7). By (5.14), the equality

$$(z - w \mp \hbar d_{ij}) \mathbf{X}_i^{\pm}(z) \mathbf{X}_j^{\pm}(w) = (z - w \pm \hbar d_{ij}) \mathbf{X}_j^{\pm}(w) \mathbf{X}_i^{\pm}(z) \quad \forall i, j \in I$$

will be satisfied provided the following identity holds:

$$(z - w \mp \hbar d_{ij}) \frac{(1-z^{-1}w)^{\delta_{ij}}(1-z^{-1}(w \mp \hbar))^{\delta_{ij}}}{(1-z^{-1}w_{\mp})^{\chi_j(i)} z^{-(\alpha_i, \alpha_j)}} = (-1)^{(\alpha_i, \alpha_j)} (z - w \pm \hbar d_{ij}) \frac{(1-w^{-1}z)^{\delta_{ij}}(1-w^{-1}(z \mp \hbar))^{\delta_{ij}}}{(1-w^{-1}z_{\mp})^{\chi_j(i)} w^{-(\alpha_i, \alpha_j)}}.$$

Using that $(\alpha_i, \alpha_j) = 2\delta_{ij} - \chi_j(i)$, we may rewrite this as

$$(z - w \mp \delta_{ij} \hbar \pm \frac{\hbar \chi_j(i)}{2}) \frac{(z - w)^{\delta_{ij}}(z - w \pm \hbar)^{\delta_{ij}}}{(z - w \pm \frac{\hbar}{2})^{\chi_j(i)}} = (-1)^{\chi_j(i)} (z - w \pm \delta_{ij} \hbar \mp \frac{\hbar \chi_j(i)}{2}) \frac{(w - z)^{\delta_{ij}}(w - z \pm \hbar)^{\delta_{ij}}}{(w - z \pm \frac{\hbar}{2})^{\chi_j(i)}}.$$

If $i \neq j$, both sides are equal to 1, and if $i = j$ both sides are equal to the polynomial

$$(z - w \pm \hbar)(z - w \mp \hbar)(z - w).$$

To prove that the relations (3.8) and (3.9) are preserved by (5.7), we employ the following well-known property of the formal delta function $\delta(z, w)$ which can be found in [Ka, Lemma 7.7] and [LeLi, Proposition 2.1.8 (b)]: given a vector space V and $f(z, w) \in V[[z^{\pm 1}, w^{\pm 1}]]$, we have

$$(5.15) \quad f(z, w) \cdot \delta(z, w) = f(z, z) \cdot \delta(z, w),$$

provided both sides of this equality are well-defined elements of $V[[z^{\pm 1}, w^{\pm 1}]]$.

The relation (3.8). From (5.13) we obtain the equality of operators

$$(5.16) \quad [\mathbf{X}_i^+(z), \mathbf{X}_j^-(w)] = \varepsilon(\alpha_i, \alpha_j) F_{i,j}(z, w) : \mathbf{X}_i^+(z) \mathbf{X}_j^-(w) :,$$

where $F_{i,j}(z, w)$ is given by

$$(5.17) \quad F_{i,j}(z, w) = \left(\frac{(1-z^{-1}w_{+})^{\chi_j(i)} z^{-(\alpha_i, \alpha_j)}}{(1-z^{-1}w)^{\delta_{ij}}(1-z^{-1}(w+\hbar))^{\delta_{ij}}} - (-1)^{(\alpha_i, \alpha_j)} \frac{(1-w^{-1}z_{-})^{\chi_i(j)} w^{-(\alpha_i, \alpha_j)}}{(1-w^{-1}z)^{\delta_{ij}}(1-w^{-1}(z-\hbar))^{\delta_{ij}}} \right).$$

If $i \neq j$ and $\chi_j(i) = 0$, then it is clear that $F_{i,j}(z, w) = 0$. If $i \neq j$ and $\chi_j(i) = 1$, then we again obtain

$$F_{i,j}(z, w) = (1 - z^{-1}w_+)z + (1 - w^{-1}z_-)w = 0.$$

Hence, we have shown that the assignment (5.7) preserves the relation (3.8) when $i \neq j$. If $i = j$, we have

$$\begin{aligned} F_{i,i}(z, w) &= \frac{z^{-2}}{(1 - z^{-1}w)(1 - z^{-1}(w + \hbar))} - \frac{w^{-2}}{(1 - w^{-1}z)(1 - w^{-1}(z - \hbar))} \\ &= \frac{z^{-1}}{1 - z^{-1}w} \left(\frac{z^{-1}}{1 - z^{-1}(w + \hbar)} + \frac{(w + \hbar)^{-1}}{1 - (w + \hbar)^{-1}z} \right) \\ &\quad - \frac{z^{-1}}{1 - z^{-1}w} \frac{(w + \hbar)^{-1}}{1 - (w + \hbar)^{-1}z} - \frac{w^{-2}}{(1 - w^{-1}z)(1 - w^{-1}(z - \hbar))} \\ &= \frac{z^{-1}}{1 - z^{-1}w} \delta(w + \hbar, z) - \frac{w^{-1}}{1 - w^{-1}(z - \hbar)} \delta(w, z), \end{aligned}$$

where we have used the identities (3.1) and $\frac{(w+\hbar)^{-1}}{1-(w+\hbar)^{-1}z} = \frac{w^{-1}}{1-w^{-1}(z-\hbar)}$. Substituting the above expression for $F_{i,i}(z, w)$ into (5.16) and using that $\varepsilon(\alpha_i, \alpha_i) = -1$, we obtain

$$(5.18) \quad [\mathbf{X}_i^+(z), \mathbf{X}_i^-(w)] = -\delta(w + \hbar, z) \frac{z^{-1}}{1 - z^{-1}w} : \mathbf{X}_i^+(z) \mathbf{X}_i^-(w) : + \delta(w, z) \frac{w^{-1}}{1 - w^{-1}(z - \hbar)} : \mathbf{X}_i^+(z) \mathbf{X}_i^-(w) : .$$

By (5.15),

$$\begin{aligned} \delta(w + \hbar, z) \frac{z^{-1}}{1 - z^{-1}w} : \mathbf{X}_i^+(z) \mathbf{X}_i^-(w) : &= \delta(w + \hbar, z) \frac{(w+\hbar)^{-1}}{1 - (w+\hbar)^{-1}w} : \mathbf{X}_i^+(z) \mathbf{X}_i^-(w) : \big|_{z \mapsto w+\hbar} \\ &= -\frac{1}{\hbar} \delta(w + \hbar, z) A_i^+(w + \hbar) A_i^-(w) B_i^+(w + \hbar) B_i^-(w) \left(\frac{w+\hbar}{w} \right)^{\partial_{\alpha_i}} \\ &= -\frac{1}{\hbar} \delta(w + \hbar, z) \mathbf{H}_i^+(w + \frac{\hbar}{2}), \end{aligned}$$

since $A_i^+(w + \hbar) = A_i^-(w)^{-1}$ (see (5.5)). Similarly, (5.15) implies that

$$\begin{aligned} \delta(w, z) \frac{w^{-1}}{1 - w^{-1}(z - \hbar)} : \mathbf{X}_i^+(z) \mathbf{X}_i^-(w) : &= \frac{1}{\hbar} \delta(w, z) : \mathbf{X}_i^+(z) \mathbf{X}_i^-(w) : \big|_{w \rightarrow z} \\ &= -\frac{1}{\hbar} \delta(w, z) A_i^+(z) A_i^-(z) B_i^+(z) B_i^-(z) = -\frac{1}{\hbar} \delta(w, z) \mathbf{H}_i^-(z), \end{aligned}$$

where we have used (5.6) and that $B_i^+(z) = B_i^-(z)^{-1}$. Substituting these identities back into (5.18), we find that

$$[\mathbf{X}_i^+(z), \mathbf{X}_i^-(w)] = \frac{1}{\hbar} (\delta(w + \hbar, z) \mathbf{H}_i^+(w + \frac{\hbar}{2}) - \delta(w, z) \mathbf{H}_i^-(z)),$$

as desired.

The relation (3.9). Observe first that if $(\alpha_i, \alpha_j) = 0$ then $\varepsilon(\alpha_i, \alpha_j) = \varepsilon(\alpha_j, \alpha_i)$ and (5.14) implies

$$[\mathbf{X}_i^\pm(z), \mathbf{X}_j^\pm(w)] = 0.$$

Hence we only need to verify that (3.9) holds when $(\alpha_i, \alpha_j) = -1$. By (5.14), we have

$$(5.19) \quad [\mathbf{X}_i^\pm(z_2), \mathbf{X}_j^\pm(w)] = \varepsilon(\alpha_i, \alpha_j) \left(\frac{z_2^{-1}}{1 - z_2^{-1}w_\mp} + \frac{w^{-1}}{1 - w^{-1}(z_2)_\mp} \right) : \mathbf{X}_i^\pm(z_2) \mathbf{X}_j^\pm(w) :,$$

while repeated application of (5.10) gives

$$\begin{aligned} \mathbf{X}_i^\pm(z_1) : \mathbf{X}_i^\pm(z_2) \mathbf{X}_j^\pm(w) : &= -\varepsilon(\alpha_i, \alpha_j) \frac{z_1(1 - z_1^{-1}z_2)(1 - z_1^{-1}(z_2 \mp \hbar))}{1 - z_1^{-1}w_\mp} : \mathbf{X}_i^\pm(z_1) \mathbf{X}_i^\pm(z_2) \mathbf{X}_j^\pm(w) :, \\ : \mathbf{X}_i^\pm(z_2) \mathbf{X}_j^\pm(w) : \mathbf{X}_i^\pm(z_1) &= \varepsilon(\alpha_i, \alpha_j) \frac{z_2^2 w^{-1}(1 - z_2^{-1}z_1)(1 - z_2^{-1}(z_1 \mp \hbar))}{1 - w^{-1}(z_1)_\mp} : \mathbf{X}_i^\pm(z_1) \mathbf{X}_i^\pm(z_2) \mathbf{X}_j^\pm(w) : . \end{aligned}$$

Combining these last two identities with (5.19) gives

$$[\mathbf{X}_i^\pm(z_1), [\mathbf{X}_i^\pm(z_2), \mathbf{X}_j^\pm(w)]] = -f(z_1, z_2, w) : \mathbf{X}_i^\pm(z_1) \mathbf{X}_i^\pm(z_2) \mathbf{X}_j^\pm(w) :,$$

where

$$f(z_1, z_2, w) = \left(\frac{z_2^{-1}}{1 - z_2^{-1}w_\mp} + \frac{w^{-1}}{1 - w^{-1}(z_2)_\mp} \right) \left(\frac{w^{-1}(z_2 - z_1)(z_2 - z_1 \pm \hbar)}{1 - w^{-1}(z_1)_\mp} + \frac{z_1^{-1}(z_1 - z_2)(z_1 - z_2 \pm \hbar)}{1 - z_1^{-1}w_\mp} \right).$$

Thus, the identity

$$(5.20) \quad f(z_1, z_2, w) + f(z_2, z_1, w) = 0$$

will imply $[\mathbf{X}_i^\pm(z_1), [\mathbf{X}_i^\pm(z_2), \mathbf{X}_j^\pm(w)]] + [\mathbf{X}_i^\pm(z_2), [\mathbf{X}_i^\pm(z_1), \mathbf{X}_j^\pm(w)]] = 0$. Since

$$\begin{aligned} \frac{z_2^{-1}}{1 - z_2^{-1}w_\mp} + \frac{w^{-1}}{1 - w^{-1}(z_2)_\mp} &= \delta(z_2, w_\mp) \mp \frac{\hbar w^{-2}}{(1 - w^{-1}(z_2)_\mp)(1 - w^{-1}(z_2)_\pm)} \quad \text{and} \\ \frac{z_1^{-1}}{1 - z_1^{-1}w_\mp} &= \delta(z_1, w_\mp) - \frac{w^{-1}}{1 - w^{-1}(z_1)_\pm}, \end{aligned}$$

the property (5.15) of the formal delta function implies that

$$\begin{aligned} f(z_1, z_2, w) &= \delta(z_2, w_\mp) \left(\frac{w^{-1}(z_2 - z_1)(z_2 - z_1 \pm \hbar)}{1 - w^{-1}(z_1)_\mp} + \frac{z_1^{-1}(z_1 - z_2)(z_1 - z_2 \pm \hbar)}{1 - z_1^{-1}w_\mp} \right) \\ &\mp \delta(z_1, w_\mp) \frac{\hbar w^{-2}(z_1 - z_2)(z_1 - z_2 \pm \hbar)}{(1 - w^{-1}(z_2)_\mp)(1 - w^{-1}(z_2)_\pm)} \\ &\mp \frac{\hbar w^{-2}}{(1 - w^{-1}(z_2)_\mp)(1 - w^{-1}(z_2)_\pm)} \left(\frac{w^{-1}(z_2 - z_1)(z_2 - z_1 \pm \hbar)}{1 - w^{-1}(z_1)_\mp} - \frac{w^{-1}(z_1 - z_2)(z_1 - z_2 \pm \hbar)}{1 - w^{-1}(z_1)_\pm} \right) \\ &= \pm \hbar \delta(z_2, w_\mp) \mp \hbar \delta(z_1, w_\mp) + \frac{\hbar^2 w^{-4}(z_2 - z_1)(z_2 + z_1 - 2w)}{(1 - w^{-1}(z_2)_\mp)(1 - w^{-1}(z_2)_\pm)(1 - w^{-1}(z_1)_\mp)(1 - w^{-1}(z_1)_\pm)}. \end{aligned}$$

As this expression is antisymmetric in z_1 and z_2 , we may conclude that (5.20) holds, and thus that the vertex operators $\{\mathbf{X}_i^\pm(z)\}_{i \in I}$ satisfy the Serre relations (3.9). \square

Remark 5.7. Taking the coefficient of $z^{-2}w$ in the relation (3.8) with $i = j$ yields

$$[x_{i1}^+, x_{i,-1}^-] = \mathbf{c} + h_{i0}.$$

Combining this with the relation $[x_{i0}^+, x_i^-(w)] = \tilde{h}_i^+(w + \frac{\hbar \mathbf{c}}{2}) + \tilde{h}_i^-(w)$ (see (3.36)), we deduce that $DY_\hbar^c(\mathfrak{g})$ is generated by $\{x_{ir}^\pm\}_{i \in I, r \in \mathbb{Z}}$. Moreover, in the Yangian double $DY_\hbar^\kappa(\mathfrak{g})$ at level $\kappa \in \mathbb{C}^\times$, the series $h_i^\pm(z)$ are uniquely determined by the relations

$$(5.21) \quad \kappa^{-1}(z - w)[x_i^+(z), x_i^-(w)] = \delta(w + \hbar \kappa, z) h_i^+(w + \frac{\hbar \kappa}{2}),$$

$$(5.22) \quad \kappa^{-1}(z - w - \hbar \kappa)[x_i^+(z), x_i^-(w)] = \delta(w, z) h_i^-(z).$$

In particular, the representation ρ_\hbar of Theorem 5.5 is entirely determined by $x_i^\pm(z) \mapsto \mathbf{X}_i^\pm(z)$ for all $i \in I$, and the formulas (5.5) and (5.6) for $\mathbf{H}_i^\pm(z)$ may be deduced from (5.21) and (5.22), as was essentially done below (5.18).

5.2. The $DY^c(\mathfrak{g})$ -module $\tilde{\mathcal{V}}$. As the coefficients of the vertex operators $\mathbf{X}_i^\pm(z)$ and $\mathbf{H}_i^\pm(z)$ are elements of $\text{End}_{\mathbb{C}[[\hbar]]} \mathcal{V}[[\hbar]]$, it is not clear that they can be specialized at $\hbar = \zeta \in \mathbb{C}^\times$ to produce a $DY_\zeta^c(\mathfrak{g})$ representation. In this subsection we exploit the existence of a $(\mathbb{Z} \times Q)$ -grading on \mathcal{V} to show that this can indeed be accomplished after modifying the representation space appropriately.

The $(\mathbb{Z} \times Q)$ -grading on $\mathcal{V} = \mathbb{C}[\mathcal{H}_{i,-r}]_{i \in I, r > 0} \otimes \mathbb{C}_\varepsilon[Q]$ is given by

$$\deg \mathcal{H}_{i,-r} = (-r, 0), \quad \deg e^\alpha = (-\frac{1}{2}(\alpha, \alpha), \alpha) \quad \forall i \in I, r > 0 \text{ and } \alpha \in Q.$$

Note that this choice of grading is different from the more familiar grading on Fock spaces obtained by setting $\deg \mathcal{H}_{i,-r} = (r, 0)$ and $\deg e^\alpha = (\frac{1}{2}(\alpha, \alpha), \alpha)$. Let $\mathcal{V}_{n,\beta}$ denote the subspace of \mathcal{V} spanned by elements of degree (n, β) , so that $\mathcal{V} = \bigoplus_{(n,\beta) \in \mathbb{Z} \times Q} \mathcal{V}_{n,\beta}$. We note the following useful observation:

Lemma 5.8. Setting $\mathcal{P} = \{(n, \beta) \in \mathbb{Z} \times Q : n \leq -\frac{1}{2}(\beta, \beta)\}$, we have

$$\mathcal{V} = \bigoplus_{(n,\beta) \in \mathcal{P}} \mathcal{V}_{n,\beta}.$$

Equivalently, $\mathcal{V}_{n,\beta} = \{0\}$ for all $n > -\frac{1}{2}(\beta, \beta)$.

Next, for each $\beta \in Q$ we set $n(\beta) = -\frac{1}{2}(\beta, \beta)$, so that

$$\mathcal{V}_\beta = \bigoplus_{n \in \mathbb{Z}} \mathcal{V}_{n, \beta} = \bigoplus_{n \leq n(\beta)} \mathcal{V}_{n, \beta} \quad \forall \beta \in Q.$$

Let $\tilde{\mathcal{V}}_\beta = \prod_{n \leq n(\beta)} \mathcal{V}_{n, \beta}$ be the completion of \mathcal{V}_β with respect to this grading, and set

$$\tilde{\mathcal{V}} = \bigoplus_{\beta \in Q} \tilde{\mathcal{V}}_\beta.$$

As $\mathcal{V}_0 = \bigoplus_{n \leq 0} \mathcal{V}_{n, 0}$ is precisely the Fock space $\mathcal{F} = \mathbb{C}[\mathcal{H}_{i, -r}]_{i \in I, r > 0}$, we have the equivalent characterizations $\tilde{\mathcal{V}}_\beta \cong \tilde{\mathcal{F}} \otimes \mathbb{C}e^\beta$ and $\tilde{\mathcal{V}} \cong \tilde{\mathcal{F}} \otimes \mathbb{C}_\varepsilon[Q]$, where $\tilde{\mathcal{F}} = \tilde{\mathcal{V}}_0$.

Now set

$$\mathcal{F}_\hbar = (\mathbb{C}[\hbar])[H_{i, -r}]_{i \in I, r > 0} \cong \mathbb{C}[\hbar] \otimes \mathcal{F}.$$

The $(\mathbb{Z} \times Q)$ -grading on \mathcal{V} extends to a grading on $\mathcal{V}_\hbar = \mathcal{F}_\hbar \otimes \mathbb{C}_\varepsilon[Q]$ after imposing $\deg \hbar = (0, 0)$. We use the same notation as above to denote its graded pieces and \mathbb{Z} -completion:

$$\begin{aligned} \mathcal{V}_\hbar &= \bigoplus_{(n, \beta) \in \mathcal{P}} (\mathcal{V}_\hbar)_{n, \beta} = \bigoplus_{\beta \in Q} (\mathcal{V}_\hbar)_\beta \quad \text{with} \quad (\mathcal{V}_\hbar)_\beta = \bigoplus_{n \leq n(\beta)} (\mathcal{V}_\hbar)_{n, \beta}, \\ \tilde{\mathcal{V}}_\hbar &= \bigoplus_{\beta \in Q} \widetilde{(\mathcal{V}_\hbar)_\beta} \cong \tilde{\mathcal{F}}_\hbar \otimes \mathbb{C}_\varepsilon[Q], \end{aligned}$$

where $\widetilde{(\mathcal{V}_\hbar)_\beta} = \prod_{n \leq n(\beta)} (\mathcal{V}_\hbar)_{n, \beta}$ and $\tilde{\mathcal{F}}_\hbar = \widetilde{(\mathcal{V}_\hbar)_0}$.

Recall that $\mathbf{X}_i^\pm(z) = \sum_{k \in \mathbb{Z}} \mathbf{X}_i^\pm[k]z^{-k-1}$ are the vertex operators which determine the action of $DY_h^c(\mathfrak{g})$ on $\mathcal{V}[[\hbar]]$ (see (5.4)).

Proposition 5.9. *For each $k \in \mathbb{Z}$ and $i \in I$, $\mathbf{X}_i^\pm[k]$ admits an expansion*

$$(5.23) \quad \mathbf{X}_i^\pm[k] = \sum_{a \geq 0} \mathbf{X}_i^\pm[k, a]\hbar^a,$$

with $\mathbf{X}_i^\pm[k, a] \in \text{End}_{\mathbb{C}} \mathcal{V}$ of degree $(k - a, \pm \alpha_i)$. Consequently, $\mathbf{X}_i^\pm(z) \in (\text{End}_{\mathbb{C}[\hbar]} \tilde{\mathcal{V}}_\hbar)[[z^{\pm 1}]]$ and the assignment

$$(5.24) \quad x_i^\pm(z) \mapsto \mathbf{X}_i^\pm(z) \quad \forall i \in I,$$

also determines an algebra morphism $\tilde{\rho}_\hbar : DY_h^c(\mathfrak{g}) \rightarrow \text{End}_{\mathbb{C}[\hbar]} \tilde{\mathcal{V}}_\hbar$.

Proof. The first part of the proposition is proven directly by expanding $\mathbf{X}_i^\pm(z)$ as a formal series in \hbar . To see that $\mathbf{X}_i^\pm[k] \in \text{End}_{\mathbb{C}[\hbar]} \tilde{\mathcal{V}}_\hbar$ for each $k \in \mathbb{Z}$, it suffices to prove that $\mathbf{X}_i^\pm[k] \tilde{\mathcal{V}}_\beta \subset \tilde{\mathcal{V}}_{\beta \pm \alpha_i}$ for each $\beta \in Q$. This is a straightforward consequence of Lemma 5.8 and (5.23).

One may prove the analogous statements for $\mathbf{H}_i^\pm(z)$ in the same way, but as noted in Remark 5.7 the coefficients of $x_i^\pm(z)$ generate $DY_h^c(\mathfrak{g})$ (with $\mathbf{H}_i^\pm(z)$ uniquely determined by (5.21) and (5.22)), and hence this is not necessary and we may conclude that (5.24) determines an algebra morphism $\tilde{\rho}_\hbar : DY_h^c(\mathfrak{g}) \rightarrow \text{End}_{\mathbb{C}[\hbar]} \tilde{\mathcal{V}}_\hbar$. \square

Proposition 5.9 implies that $\mathbf{X}_i^\pm(z)$ can be evaluated at $\hbar = \zeta \in \mathbb{C}$ to produce a well-defined element

$$\mathbf{X}_i^\pm(z, \zeta) = \mathbf{X}_i^\pm(z)_{\hbar \mapsto \zeta} \in (\text{End}_{\mathbb{C}} \tilde{\mathcal{V}})[[z^{\pm 1}]].$$

We will write $\mathbf{X}_i^{\pm, \zeta}[k]$ for the evaluation of $\mathbf{X}_i^\pm[k]$ at $\hbar = \zeta$, so that $\mathbf{X}_i^\pm(z, \zeta) = \sum_{k \in \mathbb{Z}} \mathbf{X}_i^{\pm, \zeta}[k]z^{-k-1}$.

Let $\text{End}_m \mathcal{V}$ denote the subspace of $\text{End}_{\mathbb{C}} \mathcal{V}$ spanned by operators of degree $a \in \mathbb{Z}$. (Here we consider only the \mathbb{Z} -grading on $\mathcal{V} = \bigoplus_{n \in \mathbb{Z}} \mathcal{V}_n$ induced by its $(\mathbb{Z} \times Q)$ -grading.) Consider the direct product $\prod_{m \in \mathbb{Z}} \text{End}_m \mathcal{V}$. The subspace

$$\widetilde{\text{End}}_{\mathbb{C}} \mathcal{V} = \left\{ \sum_{m \in \mathbb{Z}} A_m : A_m = 0 \quad \forall m \gg 0 \right\} \subset \prod_{m \in \mathbb{Z}} \text{End}_m \mathcal{V}$$

is an algebra with multiplication given by the usual product of formal series.

Corollary 5.10. *For each $\zeta \in \mathbb{C}$, $\tilde{\rho}_\hbar$ induces a homomorphism of \mathbb{C} -algebras*

$$\rho_\zeta : DY_\zeta^c(\mathfrak{g}) \rightarrow \text{End}_{\mathbb{C}} \tilde{\mathcal{V}}, \quad x_i^\pm(z) \mapsto \mathbf{X}_i^\pm(z, \zeta) \quad \forall i \in I.$$

Moreover, for each $k \in \mathbb{Z}$ we have

$$(5.25) \quad \mathbf{X}_i^{\pm, \zeta}[k] \in \left(\prod_{m \leq k} \text{End}_m \mathcal{V} \right) \cap \text{End}_{\mathbb{C}} \tilde{\mathcal{V}} \subset \widetilde{\text{End}_{\mathbb{C}} \mathcal{V}},$$

and hence ρ_ζ may be viewed as a morphism $\rho_\zeta : DY_\zeta^c(\mathfrak{g}) \rightarrow \widetilde{\text{End}_{\mathbb{C}} \mathcal{V}}$.

Henceforth, we will adapt the viewpoint that ρ_ζ has codomain $\widetilde{\text{End}_{\mathbb{C}} \mathcal{V}}$, and we will focus almost exclusively on the case where $\zeta = 1$, in which case we shall write $\rho = \rho_1$.

By composing ρ with $\iota : Y(\mathfrak{g}) \rightarrow DY^c(\mathfrak{g})$ from Proposition 3.8, we obtain an algebra morphism

$$(5.26) \quad \varrho = \rho \circ \iota : Y(\mathfrak{g}) \rightarrow \widetilde{\text{End}_{\mathbb{C}} \mathcal{V}}.$$

The algebra $\widetilde{\text{End}_{\mathbb{C}} \mathcal{V}}$ admits a \mathbb{Z} -filtration $\{\mathbf{F}_k(\mathcal{V})\}_{k \in \mathbb{Z}}$ given by

$$\mathbf{F}_k(\mathcal{V}) = \prod_{m \leq k} \text{End}_m \mathcal{V},$$

and we have

$$\text{gr}_{\mathbb{Z}} \widetilde{\text{End}_{\mathbb{C}} \mathcal{V}} = \bigoplus_{m \in \mathbb{Z}} \mathbf{F}_m(\mathcal{V}) / \mathbf{F}_{m-1}(\mathcal{V}) \cong \bigoplus_{m \in \mathbb{Z}} \text{End}_m \mathcal{V} \subset \text{End}_{\mathbb{C}} \mathcal{V}.$$

Set $\mathring{\mathbf{X}}_i^\pm(z) = \sum_{k \in \mathbb{Z}} \mathbf{X}_i^\pm[k, 0] z^{-k-1} \in (\text{End}_{\mathbb{C}} \mathcal{V})[[z^{\pm 1}]]$, where $\mathbf{X}_i^\pm[k, 0]$ is as in (5.23). Explicitly,

$$(5.27) \quad \mathring{\mathbf{X}}_i^\pm(z) = \pm \exp \left(\pm \sum_{r>0} \frac{\tilde{\mathcal{H}}_{i,-r}}{r} z^r \right) \exp \left(\mp \sum_{r>0} \frac{\mathcal{H}_{ir}}{r} z^{-r} \right) e^{\pm \alpha_i} z^{\partial_{\pm \alpha_i}}$$

with $\tilde{\mathcal{H}}_{i,-r} = \sum_{j \in I} (\alpha_i, \alpha_j) \mathcal{H}_{j,-r}$ for each $i \in I$ and $r > 0$. Since ι is a filtered morphism, the relation (5.25) of Corollary 5.10 together with the expansion (5.23) implies the following.

Corollary 5.11. *ρ and ϱ are \mathbb{Z} -filtered morphisms, and the composition of $\text{gr } \rho : \text{gr}_{\mathbb{Z}} DY^c(\mathfrak{g}) \rightarrow \text{End}_{\mathbb{C}} \mathcal{V}$ with the morphism $\phi_D : U(\mathfrak{t}) \rightarrow \text{gr}_{\mathbb{Z}} DY^c(\mathfrak{g})$ of Proposition 3.7 is the representation*

$$(5.28) \quad \rho_0 : U(\mathfrak{t}) \rightarrow \text{End}_{\mathbb{C}} \mathcal{V}, \quad X_i^\pm(z) \mapsto \mathring{\mathbf{X}}_i^\pm(z) \quad \forall i \in I.$$

Remark 5.12. *Here it is understood that the $\mathbb{Z}_{\geq 0}$ -filtration $\{\mathbf{F}_k\}_{k \geq 0}$ on $Y(\mathfrak{g})$ is extended to a \mathbb{Z} -filtration by setting $\mathbf{F}_k = \{0\}$ for all $k < 0$. The representation of \mathfrak{t} given by (5.28) can be obtained directly from ρ_\hbar (see (5.7)) by specializing $\hbar \mapsto 0$, or from ρ_ζ (see Corollary 5.10) by taking $\zeta = 0$. However, the \mathbb{Z} -filtration on $\widetilde{\text{End}_{\mathbb{C}} \mathcal{V}}$ will play a crucial role in Section 6.*

5.3. The \mathfrak{t} -modules \mathcal{V} and $\mathcal{V}_{\mathbf{A}}$. By Corollary 5.11, \mathcal{V} admits the structure of a \mathfrak{t} -module with action encoded by the vertex operators $\mathring{\mathbf{X}}_i^\pm(z)$ defined in (5.27). When the Cartan matrix \mathbf{A} is not invertible, this representation differs from that obtained from the classical construction of vertex representations [FrKa, MRy]. In this subsection we explain the relation between the two constructions.

We begin by recalling the classical setting. By (3.19), the Lie subalgebra of \mathfrak{t} generated by the coefficients of the series $\{H_i(z)\}_{i \in I}$ is a homomorphic image of the following Heisenberg algebra.

Definition 5.13. *The Heisenberg Lie algebra $\mathfrak{H}_{\mathbf{A}}$ associated to the Cartan matrix \mathbf{A} (equivalently, to the root lattice Q) is the Lie algebra over \mathbb{C} with basis $\{H_{ir}\}_{i \in I, r \in \mathbb{Z}} \cup \{\mathbf{C}\}$ subject to the defining Lie bracket relations*

$$[H_{ir}, \mathbf{C}] = 0 \quad \text{and} \quad [H_{ir}, H_{js}] = r(\alpha_i, \alpha_j) \delta_{r,-s} \mathbf{C} \quad \forall i, j \in I \text{ and } r, s \in \mathbb{Z}.$$

For each fixed $\lambda \in Q$, there is a natural action of $\mathfrak{H}_{\mathbf{A}}$ on the polynomial algebra $\mathbb{C}[H_{i,-r}]_{i \in I, r > 0}$ given by

$$H_{j,-s}(f) = H_{j,-s}f, \quad \mathbf{C}(f) = f, \quad H_{j0}(f) = (\alpha_j, \lambda)f, \quad H_{js}(f) = \partial_{js}(f)$$

for all $f \in \mathbb{C}[H_{i,-r}]_{i \in I, r > 0}$, $j \in I$ and $s > 0$, where ∂_{js} is the derivation defined uniquely by

$$\partial_{js}(H_{i,-r}) = s(\alpha_i, \alpha_j) \delta_{s,-r} \quad \forall s > 0 \text{ and } i \in I.$$

We denote $\mathbb{C}[H_{i,-r}]_{i \in I, r > 0}$, equipped with this module structure, by $\mathcal{F}_{\mathbf{A}}^\lambda$.

Now define the vector space $\mathcal{V}_{\mathbf{A}}$ by

$$\mathcal{V}_{\mathbf{A}} = \mathbb{C}[H_{i,-r}]_{i \in I, r > 0} \otimes \mathbb{C}_{\varepsilon}[Q].$$

After identifying $\mathbb{C}[H_{i,-r}]_{i \in I, r > 0} \otimes \mathbb{C}e^{\lambda}$ with $\mathcal{F}_{\mathbf{A}}^{\lambda}$, the space $\mathcal{V}_{\mathbf{A}}$ becomes an \mathfrak{H} -module isomorphic to $\bigoplus_{\lambda \in Q} \mathcal{F}_{\mathbf{A}}^{\lambda}$. To extend this to a \mathfrak{t} -module structure, define for each $\alpha = \sum_{i \in I} n_i \alpha_i \in Q$ operators $\{H_{\alpha,r}\}_{r \in \mathbb{Z}}$ on $\mathcal{V}_{\mathbf{A}}$ by $H_{\alpha,r} = \sum_{i \in I} n_i H_{i,r}$. We then set

$$\Gamma_{\alpha}^{\pm}(z) = \exp \left(\mp \sum_{r > 0} \frac{H_{\alpha, \pm r}}{r} z^{\mp r} \right) \quad \forall \alpha \in Q,$$

and introduce vertex operators $\mathbb{H}_{\alpha}(z), \mathbb{X}_{\alpha}(z) \in (\text{End}_{\mathbb{C}} \mathcal{V}_{\mathbf{A}})[[z^{\pm 1}]]$ by

$$\mathbb{H}_{\alpha}(z) = \sum_{r \in \mathbb{Z}} H_{\alpha,r} z^{-r-1} \quad \text{and} \quad \mathbb{X}_{\alpha}(z) = \Gamma_{\alpha}^{-}(z) \Gamma_{\alpha}^{+}(z) e^{\alpha} z^{\partial_{\alpha}} \quad \forall \alpha \in Q.$$

Proposition 5.14. *Set $\mathbb{X}_i^{\pm}(z) = \pm \mathbb{X}_{\alpha_i}(z)$ and $\mathbb{H}_i(z) = \mathbb{H}_{\alpha_i}(z)$ for all $i \in I$. Then the assignment*

$$(5.29) \quad X_i^{\pm}(z) \mapsto \mathbb{X}_i^{\pm}(z), \quad H_i(z) \mapsto \mathbb{H}_i(z) \quad \forall i \in I, \quad \mathbf{C} \mapsto 1$$

extends to a homomorphism of algebras $\rho_{\mathbf{A}} : U(\mathfrak{t}) \mapsto \text{End}_{\mathbb{C}} \mathcal{V}_{\mathbf{A}}$.

Proof. Although, to the best of our knowledge, the statement of the proposition has only been written down explicitly for \mathbf{A} of finite and of affine type [FrKa, MRY], the argument used to prove the above proposition for \mathfrak{t} associated to the Cartan matrix of an arbitrary simply laced Kac-Moody Lie algebra is the same, and analogous to the proof Theorem 5.5. We refer the reader to [Ka, Thm. 14.8], [MRY, Prop. 4.3] and [LeLi, §6.5] for complete details. The result may also be deduced from [Ji2, Thm. 3.1]. \square

Remark 5.15. *Suppose now that \mathbf{A} is the Cartan matrix of an arbitrary symmetric Kac-Moody Lie algebra (not constrained by the condition (2.1)), and let $\mathfrak{t}_{\mathbf{A}}$ be the Lie algebra defined identically to \mathfrak{t} (see Definition 3.5), but with (3.22) replaced by*

$$(z-w)^{-a_{ij}} [X_i^{\pm}(z), X_j^{\pm}(w)] = 0 \quad \forall i, j \in I.$$

Then the assignment (5.29) determines an algebra homomorphism $U(\mathfrak{t}_{\mathbf{A}}) \rightarrow \text{End}_{\mathbb{C}} \mathcal{V}_{\mathbf{A}}$. The added difficulty in proving this statement is verifying that (5.29) preserves the Serre relation (3.23) when $a_{ij} < -1$. This can again be deduced from [Ji2], although it may also be proven directly using elementary properties of the formal delta function $\delta(z, w)$ and its partial derivatives.

We now turn to relating $\mathcal{V}_{\mathbf{A}}$ with the \mathfrak{t} -module \mathcal{V} from Corollary 5.11. Recall from Definition 5.1 that \mathfrak{H} is the Heisenberg Lie algebra associated to the trivial lattice $\mathbb{Z}^{|I|}$. For each $k \in \mathbb{Z}_{\neq 0}$, set

$$\mathfrak{H}_{\mathbf{A}}^{(k)} = \bigoplus_{i \in I} \mathbb{C} H_{ik} \quad \text{and} \quad \mathfrak{H}^{(k)} = \bigoplus_{i \in I} \mathbb{C} \mathcal{H}_{ik}.$$

Similarly, we set $\mathfrak{H}_{\mathbf{A}}^{(0)} = \bigoplus_{i \in I} \mathbb{C} \mathcal{H}_{i0} \oplus \mathbb{C} \cdot \mathbf{C}$ and $\mathfrak{H}^{(0)} = \mathbb{C} \cdot \mathbf{C}$. Let $\mathfrak{H}'_{\mathbf{A}} = [\mathfrak{H}_{\mathbf{A}}, \mathfrak{H}_{\mathbf{A}}]$ be the derived subalgebra of $\mathfrak{H}_{\mathbf{A}}$:

$$\mathfrak{H}'_{\mathbf{A}} = \bigoplus_{k \neq 0} \mathfrak{H}_{\mathbf{A}}^{(k)} \oplus \mathbb{C} \cdot \mathbf{C} = \bigoplus_{k \in \mathbb{Z}} \mathfrak{H}'_{\mathbf{A}}{}^{(k)}, \quad \text{where} \quad \mathfrak{H}'_{\mathbf{A}}{}^{(k)} = \mathfrak{H}_{\mathbf{A}}^{(k)} \cap \mathfrak{H}'_{\mathbf{A}}.$$

In addition, we denote $\bigoplus_{k \geq 0} \mathfrak{H}'_{\mathbf{A}}{}^{(k)}$ by $\mathfrak{H}_{\mathbf{A}}^{+}$ and $\bigoplus_{k < 0} \mathfrak{H}'_{\mathbf{A}}{}^{(k)}$ by $\mathfrak{H}_{\mathbf{A}}^{-}$, and define \mathfrak{H}^{\pm} analogously.

Lemma 5.16.

(1) *The assignment*

$$\varphi_{\mathbf{A}} : \mathbf{C} \mapsto \mathbf{C}, \quad H_{ir} \mapsto \tilde{H}_{ir} = \begin{cases} \sum_{j \in I} (\alpha_i, \alpha_j) \mathcal{H}_{jr} & \text{if } r < 0, \\ \mathcal{H}_{ir} & \text{if } r > 0, \end{cases}$$

extends to morphism of graded Lie algebras $\varphi_{\mathbf{A}} : \mathfrak{H}'_{\mathbf{A}} \rightarrow \mathfrak{H}$.

(2) *For each $r < 0$, $\varphi_{\mathbf{A}}|_{\mathfrak{H}_{\mathbf{A}}^{(r)}} : \mathfrak{H}_{\mathbf{A}}^{(r)} \rightarrow \mathfrak{H}^{(r)}$ has matrix equal to \mathbf{A} with respect to the bases $\{H_{ir}\}_{i \in I} \subset \mathfrak{H}_{\mathbf{A}}^{(r)}$ and $\{\mathcal{H}_{ir}\}_{i \in I} \subset \mathfrak{H}^{(r)}$. Consequently,*

$$\varphi_{\mathbf{A}}|_{\mathfrak{H}_{\mathbf{A}}^{-}} : \mathfrak{H}_{\mathbf{A}}^{-} \rightarrow \mathfrak{H}^{-}$$

is an isomorphism if and only if \mathbf{A} is invertible, and the same is true for $\varphi_{\mathbf{A}}$.

By the lemma, $\varphi_{\mathbf{A}}|_{\mathfrak{H}_{\mathbf{A}}^-} : \mathfrak{H}_{\mathbf{A}}^- \rightarrow \mathfrak{H}^-$ induces an algebra morphism

$$\Phi_{\mathbf{A}} : U(\mathfrak{H}_{\mathbf{A}}^-) \rightarrow U(\mathfrak{H}^-)$$

which is invertible precisely when \mathbf{A} is. After identifying $U(\mathfrak{H}_{\mathbf{A}}^-)$ and $U(\mathfrak{H}^-)$ with the Fock space representations $\mathbb{C}[H_{i,-r}]_{i \in I, r > 0}$ and $\mathbb{C}[\mathcal{H}_{i,-r}]_{i \in I, r > 0}$ of $\mathfrak{H}'_{\mathbf{A}}$ and \mathfrak{H} , respectively, and equipping $\mathbb{C}[\mathcal{H}_{i,-r}]_{i \in I, r > 0}$ with the structure of a $\mathfrak{H}'_{\mathbf{A}}$ -module via $\varphi_{\mathbf{A}}$, $\Phi_{\mathbf{A}}$ becomes a morphism of $\mathfrak{H}'_{\mathbf{A}}$ -modules. This discussion leads us to the following result.

Proposition 5.17. *The \mathbb{C} -algebra morphism*

$$\Phi_{\mathbf{A}} \otimes \text{id} : \mathbb{C}[H_{i,-r}]_{i \in I, r > 0} \otimes \mathbb{C}_{\varepsilon}[Q] \rightarrow \mathbb{C}[\mathcal{H}_{i,-r}]_{i \in I, r > 0} \otimes \mathbb{C}_{\varepsilon}[Q]$$

is a morphism of \mathfrak{t} -modules $\mathcal{V}_{\mathbf{A}} \rightarrow \mathcal{V}$. It is an isomorphism precisely when \mathbf{A} is invertible.

Proof. Lemma 5.16 and the discussion following it prove that $\Phi_{\mathbf{A}} \otimes \text{id}$ will be invertible exactly when \mathbf{A} is. By comparing the definitions of the vertex operators $\dot{\mathbf{X}}_i^{\pm}(z)$ and $\mathbf{X}_i^{\pm}(z)$ (see (5.27) and Proposition 5.14), we find that $\Phi_{\mathbf{A}} \otimes \text{id}$ will be a morphism of \mathfrak{t} -modules provided $\Phi_{\mathbf{A}}$ is a morphism of $\mathfrak{H}'_{\mathbf{A}}$ -modules in the sense described before the statement of the proposition. As this has already been established, the proposition is proved. \square

6. THE POINCARÉ-BIRKHOFF-WITT THEOREM

We now fix \mathfrak{g} to be a Kac-Moody Lie algebra associated to an indecomposable Cartan matrix \mathbf{A} which is of affine type, and whose associated Dynkin diagram is simply laced with $\ell + 1$ vertices. As in Section 4, we set $I = \{0, 1, \dots, \ell\}$ with $\{1, \dots, \ell\}$ labeling the Dynkin diagram of the underlying finite-dimensional rank ℓ simple Lie algebra \mathfrak{g}_0 .

In this section we will prove that the epimorphism $\phi : U(\mathfrak{s}) \rightarrow \text{gr } Y(\mathfrak{g})$ of Proposition 2.9 is an isomorphism: see Theorem 6.9. By Propositions 4.4 and 4.7, this will imply that $\text{gr } Y(\mathfrak{g}) \cong U(\text{uce}(\mathfrak{g}'[t]))$. As a corollary, we prove in Theorem 6.10 that $Y_{\hbar}(\mathfrak{g})$ is a flat deformation $U(\mathfrak{s}) \cong U(\text{uce}(\mathfrak{g}'[t]))$ (see Remark 2.8).

6.1. A faithful representation of \mathfrak{s} . Our first step in proving the injectivity of ϕ is to use the results of Section 5 to produce a representation of $Y(\mathfrak{g})$ which specializes to a faithful representation of $\mathfrak{s} \cong \text{uce}(\mathfrak{g}'[t])$. To accomplish this, we first enlarge \mathbf{A} to an invertible Cartan matrix.

Set $\dot{I} = I \cup \{-1\}$, and extend \mathbf{A} to a Cartan matrix $\dot{\mathbf{A}} = (a_{ij})_{i,j \in \dot{I}}$ by imposing

$$a_{-1,i} = a_{i,-1} = 2\delta_{-1,i} - \delta_{i,0} \quad \forall i \in \dot{I}.$$

Definition 6.1. *Define $\dot{\mathfrak{g}}$ to be the simply-laced Kac-Moody Lie algebra with Cartan matrix $\dot{\mathbf{A}}$.*

We fix an invariant symmetric non-degenerate bilinear form $\langle \cdot, \cdot \rangle$ on $\dot{\mathfrak{g}}$ extending (\cdot, \cdot) , and assume that it is normalized so that $\langle \alpha_i, \alpha_i \rangle = 2$ for all $-1 \leq i \leq \ell$. In particular $a_{ij} = \langle \alpha_i, \alpha_j \rangle$ for all $-1 \leq i, j \leq \ell$. Let $\dot{Q} = \bigoplus_{-1 \leq i \leq \ell} \mathbb{Z}\alpha_i = \mathbb{Z}\alpha_{-1} \oplus Q$ denote the root lattice of $\dot{\mathfrak{g}}$. The following lemma can be easily deduced.

Lemma 6.2. *The Cartan matrix $\dot{\mathbf{A}}$ is invertible. In particular, $\langle \cdot, \cdot \rangle|_{\dot{Q} \times \dot{Q}}$ is non-degenerate.*

Henceforth, we will use the notation $\dot{\mathcal{V}}$ to denote the space (5.3) corresponding to the above data:

$$\dot{\mathcal{V}} = \mathbb{C}[\mathcal{H}_{i,-r}]_{i \in \dot{I}, r > 0} \otimes \mathbb{C}_{\varepsilon}[\dot{Q}].$$

By Corollaries 5.10 and 5.11, we have a \mathbb{Z} -filtered morphism of \mathbb{C} -algebras

$$\dot{\rho} : DY^c(\dot{\mathfrak{g}}) \rightarrow \widetilde{\text{End}}_{\mathbb{C}} \dot{\mathcal{V}}, \quad x_i^{\pm}(z) \mapsto \mathbf{X}_i^{\pm}(z, 1) \quad \forall i \in \dot{I}.$$

Observe that the assignment

$$\dot{\iota} : x_i^{\pm}(z), h_i^{\pm}(z), c \mapsto x_i^{\pm}(z), h_i^{\pm}(z), c \quad \forall i \in I$$

extends to a filtered algebra homomorphism $\dot{\iota} : DY^c(\mathfrak{g}) \rightarrow DY^c(\dot{\mathfrak{g}})$. We set

$$\dot{\rho} = \dot{\rho} \circ \dot{\iota} : DY^c(\mathfrak{g}) \rightarrow \widetilde{\text{End}}_{\mathbb{C}} \dot{\mathcal{V}}.$$

Define a representation $\dot{\rho}_0$ of \mathfrak{t} on $\dot{\mathcal{V}}$ by setting

$$(6.1) \quad \dot{\rho}_0 = \text{gr}(\dot{\rho} \circ \dot{\iota}) \circ \phi_D : U(\mathfrak{t}) \rightarrow \bigoplus_{m \in \mathbb{Z}} \text{End}_m \dot{\mathcal{V}} \subset \text{End}_{\mathbb{C}} \dot{\mathcal{V}},$$

where ϕ_D is as in Proposition 3.7.

Lemma 6.3. *The \mathfrak{t} -module $\mathring{\mathcal{V}}$, equipped with action given by $\dot{\rho}_0$ above, is a faithful module.*

Proof. Let \mathfrak{t} be the Lie algebra from Definition 3.5 corresponding to $\mathring{\mathbf{A}}$. By (5.17) and Lemma 6.2, the morphism $\Phi_{\mathring{\mathbf{A}}} \otimes \text{id} : \mathcal{V}_{\mathring{\mathbf{A}}} \rightarrow \mathring{\mathcal{V}}$ is an isomorphism of \mathfrak{t} -modules. By pulling back via the natural morphism $\mathfrak{t} \rightarrow \mathfrak{t}$, we obtain an isomorphism of \mathfrak{t} -modules, and the induced \mathfrak{t} -module structure on $\mathring{\mathcal{V}}$ is precisely that given by $\dot{\rho}_0$. That this is a faithful \mathfrak{t} -module now follows from the fact that $\mathcal{V}_{\mathring{\mathbf{A}}}$ is precisely $V(\mathring{Q}, \mathfrak{H}'_{\mathring{\mathbf{A}}})$ from [MRY], and by [MRY, Prop. 4.3], this is a faithful \mathfrak{t} -module. \square

Now set $\dot{\varrho} = \dot{\rho} \circ \iota : Y(\mathfrak{g}) \rightarrow \widetilde{\text{End}}_{\mathbb{C}} \mathring{\mathcal{V}}$, where $\iota : Y(\mathfrak{g}) \rightarrow DY^c(\mathfrak{g})$ is as in Proposition 3.8, and define

$$\dot{\varrho}_0 = \text{gr } \dot{\varrho} \circ \phi : U(\mathfrak{s}) \rightarrow \bigoplus_{m \in \mathbb{Z}} \text{End}_m \mathring{\mathcal{V}} \subset \text{End}_{\mathbb{C}} \mathring{\mathcal{V}},$$

where $\phi : U(\mathfrak{s}) \rightarrow \text{gr } Y(\mathfrak{g})$ is as in Proposition 2.9.

Corollary 6.4. *The \mathfrak{s} -module $\mathring{\mathcal{V}}$, equipped with action given by $\dot{\varrho}_0$ above, is a faithful module.*

Proof. The representation $\dot{\varrho}_0$ is equal to the restriction of $\dot{\rho}_0$ to $U(\mathfrak{s})$ via the embedding of Corollary 4.6, so the result follows immediately. \square

We will use this faithful module, together with the coproduct $\Delta_{\mathfrak{s},u}$ from Subsection 2.3, to construct an embedding of $U(\mathfrak{s})$ into a large algebra built by gluing together endomorphism rings associated to $\mathring{\mathcal{V}}$. We begin with the following general result.

Let \mathfrak{a} be an arbitrary complex Lie algebra and let $\Delta_{\mathfrak{a}}$ and $\varepsilon_{\mathfrak{a}}$ be the coproduct and counit, respectively, of the enveloping algebra $U(\mathfrak{a})$.

Proposition 6.5. *Let V be a faithful representation of \mathfrak{a} with $\rho_{\mathfrak{a}} : U(\mathfrak{a}) \rightarrow \text{End}_{\mathbb{C}} V$ the corresponding morphism. For each $k \geq 0$, set $\rho_{\mathfrak{a},k} = \rho_{\mathfrak{a}}^{\otimes k} \circ \Delta_{\mathfrak{a}}^{(k-1)}$, with $\rho_{\mathfrak{a},0} = \varepsilon_{\mathfrak{a}}$. The universal property of $\prod_{m \geq 0} \text{End}_{\mathbb{C}}(V^{\otimes m})$ dictates that there is a unique morphism*

$$\Phi : U(\mathfrak{a}) \rightarrow \prod_{m \geq 0} \text{End}_{\mathbb{C}}(V^{\otimes m}), \quad \pi_m \circ \Phi = \rho_{\mathfrak{a},m} \quad \forall m \geq 0,$$

where $\pi_m : \prod_{m \geq 0} \text{End}(V^{\otimes m}) \rightarrow \text{End}(V^{\otimes m})$ is the natural projection. Then Φ is injective.

Proof. The result follows from a modification of the argument given in the proof of [AMR, Lemma 3.5]. \square

We would now like to imitate Proposition 6.5 with $\Delta_{\mathfrak{a}}^{(k-1)}$ replaced by $\Delta_{\mathfrak{s},u}^{k-1}$. Let V be a faithful \mathfrak{s} -module with corresponding morphism $\rho_{\mathfrak{s}} : U(\mathfrak{s}) \rightarrow \text{End}_{\mathbb{C}} V$, and for each $k \geq 1$, set

$$\rho_{\mathfrak{s},u}^k = \rho_{\mathfrak{s}}^{\otimes k} \circ \Delta_{\mathfrak{s},u}^{k-1} : U(\mathfrak{s}) \rightarrow \text{End}_{\mathbb{C}}(V^{\otimes k})[u^{\pm 1}].$$

We also set $\rho_{\mathfrak{s},u}^0 = \varepsilon_{\mathfrak{s}} : U(\mathfrak{s}) \rightarrow \mathbb{C} \subset \mathbb{C}[u^{\pm 1}]$. Then there is a unique morphism

$$(6.2) \quad \Phi_u : U(\mathfrak{s}) \rightarrow \prod_{m \geq 0} \text{End}_{\mathbb{C}}(V^{\otimes m})[u^{\pm 1}], \quad \pi_m \circ \Phi_u = \rho_{\mathfrak{s},u}^m \quad \forall m \geq 0,$$

where $\pi_m : \prod_{m \geq 0} \text{End}_{\mathbb{C}}(V^{\otimes m})[u^{\pm 1}] \rightarrow \text{End}_{\mathbb{C}}(V^{\otimes m})[u^{\pm 1}]$ is the natural projection.

Proposition 6.6. *The morphism Φ_u is injective.*

Proof. The evaluation $u \mapsto 1$ induces a morphism

$$\text{ev} : \prod_{m \geq 0} \text{End}_{\mathbb{C}}(V^{\otimes m})[u^{\pm 1}] \rightarrow \prod_{m \geq 0} \text{End}_{\mathbb{C}}(V^{\otimes m}).$$

The composition $\text{ev} \circ \Phi_u : U(\mathfrak{s}) \rightarrow \prod_{m \geq 0} \text{End}_{\mathbb{C}}(V^{\otimes m})$ agrees with the morphism Φ associated to V from Proposition 6.5, and hence is injective. This implies that Φ_u is also injective. \square

Applying Proposition 6.6 with V the faithful \mathfrak{s} -module $\mathring{\mathcal{V}}$ from Corollary 6.4, we obtain the following corollary.

Corollary 6.7. *The morphism of \mathbb{C} -algebras $\Phi_u : U(\mathfrak{s}) \rightarrow \prod_{m \geq 0} \text{End}_{\mathbb{C}}(\mathring{\mathcal{V}}^{\otimes m})[u^{\pm 1}]$, defined by (6.2) with $\rho_{\mathfrak{s}} = \dot{\varrho}_0$, is injective.*

6.2. Statement and proof of the main result. We now construct the Yangian version Ψ_u of the embedding Φ_u from Corollary 6.7, using the morphism $\check{\varrho} : Y(\mathfrak{g}) \rightarrow \widetilde{\text{End}}_{\mathbb{C}} \dot{\mathcal{V}}$. The injectivity of Ψ_u is closely tied to the Poincaré-Birkhoff-Witt Theorem for $Y(\mathfrak{g})$, as we shall explain shortly.

For each $k \geq 1$, $\check{\varrho}^{\otimes k}$ extends to a homomorphism $Y(\mathfrak{g})^{\otimes k}((u)) \rightarrow (\widetilde{\text{End}}_{\mathbb{C}} \dot{\mathcal{V}})^{\otimes k}((u))$. Composing with Δ_u^{k-1} from (2.31), we obtain a morphism

$$\check{\varrho}_u^k : Y(\mathfrak{g}) \rightarrow (\widetilde{\text{End}}_{\mathbb{C}} \dot{\mathcal{V}})^{\otimes k}((u)).$$

As in the $U(\mathfrak{s})$ -case, we set $\check{\varrho}_u^0$ to be the counit.

For each $a \in \mathbb{Z}$, set

$$\text{End}_a(\dot{\mathcal{V}}^{\otimes k}) = \bigoplus_{a_1 + \dots + a_k = a} \left(\text{End}_{a_1} \dot{\mathcal{V}} \otimes \dots \otimes \text{End}_{a_k} \dot{\mathcal{V}} \right) \subset \text{End}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes k}).$$

We let $\widetilde{\text{End}}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes k})$ denote the subspace of $\prod_{a \in \mathbb{Z}} \text{End}_a(\dot{\mathcal{V}}^{\otimes k})$ consisting of summations $\sum_{m \in \mathbb{Z}} A_m$ with $A_m = 0$ for all $m \gg 0$. This is an algebra with multiplication extending that of $\bigoplus_{a \in \mathbb{Z}} \text{End}_a(\dot{\mathcal{V}}^{\otimes k})$. Setting

$$\mathbf{F}_{\ell}(\dot{\mathcal{V}}^{\otimes k}) = \prod_{a \leq \ell} \text{End}_a(\dot{\mathcal{V}}^{\otimes k}) \quad \forall \ell \in \mathbb{Z}$$

equips $\widetilde{\text{End}}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes k})$ with the structure of a \mathbb{Z} -filtered algebra. Recall that $\{\mathbf{F}_{\ell}\}_{\ell \geq 0}$ denotes the $\mathbb{Z}_{\geq 0}$ -filtration on $Y(\mathfrak{g})$ defined above Proposition 2.9, which is extended to a \mathbb{Z} -filtration by setting $\mathbf{F}_{-\ell} = 0$ for $\ell > 0$.

Lemma 6.8. *The image of $\check{\varrho}_u^k$ embeds into $\widetilde{\text{End}}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes k})((u))$. Moreover,*

$$\check{\varrho}_u^k(\mathbf{F}_{\ell}) \subset \mathbf{F}_{\ell}(\dot{\mathcal{V}}^{\otimes k})((u)) \quad \forall \ell \in \mathbb{Z}.$$

Proof. By (2.32), $\Delta_u^{k-1}(\mathbf{F}_{\ell}) \subset \mathbf{F}_{\ell}(Y(\mathfrak{g})^{\otimes k})((u))$, where $\mathbf{F}_{\ell}(Y(\mathfrak{g})^{\otimes k}) = \sum_{a_1 + \dots + a_k = \ell} \mathbf{F}_{a_1} \otimes \dots \otimes \mathbf{F}_{a_k}$. Since $\check{\varrho}$ is also filtered (see Corollary 5.11), we have $\check{\varrho}_u^k(\mathbf{F}_{\ell}) \subset \widetilde{\mathbf{F}}_{\ell}(\dot{\mathcal{V}}^{\otimes k})((u))$, with

$$\widetilde{\mathbf{F}}_{\ell}(\dot{\mathcal{V}}^{\otimes k}) = \sum_{a_1 + \dots + a_k = \ell} \mathbf{F}_{a_1}(\dot{\mathcal{V}}) \otimes \dots \otimes \mathbf{F}_{a_k}(\dot{\mathcal{V}}).$$

As $\mathbf{F}_m(\dot{\mathcal{V}}) = \prod_{a \leq m} \text{End}_a(\dot{\mathcal{V}})$, $\bigotimes_{b=1}^k \mathbf{F}_{a_b}(\dot{\mathcal{V}})$ naturally embeds into the space $\mathbf{F}_{\ell}(\dot{\mathcal{V}}^{\otimes k}) = \prod_{a \leq \ell} \text{End}_a(\dot{\mathcal{V}}^{\otimes k})$, provided $\sum_{b=1}^k a_b = \ell$. This proves the assertion. \square

Consider the algebra

$$\text{End}_u(\dot{\mathcal{V}}^{\otimes k}) = \bigcup_{\ell \in \mathbb{Z}} \left(\mathbf{F}_{\ell}(\dot{\mathcal{V}}^{\otimes k})((u)) \right) \subset \widetilde{\text{End}}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes k})((u)).$$

It is \mathbb{Z} -filtered with $\mathbf{F}_{\ell}(\text{End}_u(\dot{\mathcal{V}}^{\otimes k})) = \mathbf{F}_{\ell}(\dot{\mathcal{V}}^{\otimes k})((u))$ and

$$\text{gr End}_u(\dot{\mathcal{V}}^{\otimes k}) = \bigoplus_{\ell \in \mathbb{Z}} \text{End}_{\ell}(\dot{\mathcal{V}}^{\otimes k})((u)) \subset \text{End}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes k})((u)).$$

Lemma 6.8 implies that $\check{\varrho}_u^k$ can be viewed as a \mathbb{Z} -filtered morphism

$$\check{\varrho}_u^k : Y(\mathfrak{g}) \rightarrow \text{End}_u(\dot{\mathcal{V}}^{\otimes k}).$$

After forming the direct product of algebras $\prod_{m \geq 0} \text{End}_u(\dot{\mathcal{V}}^{\otimes m})$, we obtain an algebra morphism

$$(6.3) \quad \Psi_u : Y(\mathfrak{g}) \rightarrow \prod_{m \geq 0} \text{End}_u(\dot{\mathcal{V}}^{\otimes m}), \quad \pi_m \circ \Psi_u = \check{\varrho}_u^m \quad \forall m \geq 0,$$

where $\pi_m : \prod_{m \geq 0} \text{End}_u(\dot{\mathcal{V}}^{\otimes m}) \rightarrow \text{End}_u(\dot{\mathcal{V}}^{\otimes m})$ is the m -th projection morphism. We are now prepared to state and prove the main result of this section:

Theorem 6.9. *The morphism Ψ_u defined in (6.3) is an embedding of algebras, and the epimorphism*

$$\phi : U(\mathfrak{s}) \cong U(\text{uce}(\mathfrak{g}'[t])) \twoheadrightarrow \text{gr } Y(\mathfrak{g}), \quad X_{ir}^{\pm}, H_{ir} \mapsto \bar{x}_{ir}^{\pm}, \bar{h}_{ir}$$

of Proposition 2.9 is an isomorphism of algebras.

Proof. As, for each $k \geq 0$, $\dot{\varrho}_u^k$ is a filtered morphism $Y(\mathfrak{g}) \rightarrow \text{End}_u(\dot{\mathcal{V}}^{\otimes k})$, we may form the associated graded morphisms

$$\text{gr } \dot{\varrho}_u^k : \text{gr } Y(\mathfrak{g}) \rightarrow \text{End}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes k})(\langle u \rangle).$$

By (2.33), the image of $\text{gr } \dot{\varrho}_u^k$ in fact lies in $\text{End}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes k})[u^{\pm 1}]$. We therefore obtain an algebra morphism

$$\overline{\Psi}_u : \text{gr } Y(\mathfrak{g}) \rightarrow \prod_{m \geq 0} \text{End}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes m})[u^{\pm 1}], \quad \dot{\pi}_m \circ \overline{\Psi}_u = \text{gr } \dot{\varrho}_u^m \quad \forall m \geq 0,$$

where now $\dot{\pi}_m$ is the m -th projection morphism for $\prod_{m \geq 0} \text{End}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes m})[u^{\pm 1}]$.

By definition, $\dot{\varrho}_0 = \text{gr } \dot{\varrho} \circ \phi$ (see (6.1)), and hence the commutativity of the diagram (2.34) implies that

$$(6.4) \quad \overline{\Psi}_u \circ \phi = \Phi_u,$$

where $\Phi_u : U(\mathfrak{s}) \rightarrow \prod_{m \geq 0} \text{End}_{\mathbb{C}}(\dot{\mathcal{V}}^{\otimes k})[u^{\pm 1}]$ is the embedding of Corollary 6.7. This implies that ϕ is also injective, and hence an isomorphism of algebras.

The relation (6.4) also implies that $\overline{\Psi}_u$ is an embedding, from which it follows that Ψ_u is injective using a standard argument. Indeed, given a nonzero element $X \in Y(\mathfrak{g})$, we may take $\ell \geq 0$ minimal such that $X \in \mathbf{F}_{\ell}$. Let $\bar{X} \in \text{gr } Y(\mathfrak{g})$ denote the image of X in $\mathbf{F}_{\ell}/\mathbf{F}_{\ell-1}$, which is nonzero by assumption. If $\Psi_u(X) = 0$, then $\text{gr } \dot{\varrho}_u^m(\bar{X}) = 0$ for all $m \geq 0$ and hence $\overline{\Psi}_u(\bar{X}) = 0$, which is impossible. \square

Recall that if \mathbf{A} is a $\mathbb{Z}_{\geq 0}$ -filtered algebra with ascending filtration $\{\mathbf{F}_k(\mathbf{A})\}_{k \geq 0}$, then the Rees algebra associated to \mathbf{A} is

$$R_{\hbar}(\mathbf{A}) = \bigoplus_{k \geq 0} \hbar^k \mathbf{F}_k(\mathbf{A}) \subset \mathbf{A}[\hbar].$$

The Rees algebra $R_{\hbar}(\mathbf{A})$ always satisfies $R_{\hbar}(\mathbf{A})/(\hbar - 1)R_{\hbar}(\mathbf{A}) \cong \mathbf{A}$ and $R_{\hbar}(\mathbf{A})/\hbar R_{\hbar}(\mathbf{A}) \cong \text{gr } \mathbf{A}$. The next theorem employs the Rees algebra construction to characterize $Y_{\hbar}(\mathfrak{g})$ in terms of $Y(\mathfrak{g})$.

Theorem 6.10. *The assignment $x_{ir}^{\pm}, h_{ir} \mapsto \hbar^r x_{ir}^{\pm}, \hbar^r h_{ir}$ extends to an isomorphism of $\mathbb{C}[\hbar]$ -algebras*

$$\Psi_{\hbar} : Y_{\hbar}(\mathfrak{g}) \rightarrow R_{\hbar}(Y(\mathfrak{g})) \subset Y(\mathfrak{g})[\hbar].$$

Consequently, $Y_{\hbar}(\mathfrak{g})$ is a flat deformation of the algebra $U(\mathfrak{s}) \cong U(\text{ucc}(\mathfrak{g}'[t]))$.

Proof. That the assignment $x_{ir}^{\pm}, h_{ir} \mapsto \hbar^r x_{ir}^{\pm}, \hbar^r h_{ir}$ extends to a homomorphism Ψ_{\hbar} of $\mathbb{C}[\hbar]$ -modules is verified directly (cf. (2.14)). Since $\{\hbar^r x_{ir}^{\pm}, \hbar^r h_{ir}\}_{i \in I, r \geq 0}$ generate $R_{\hbar}(Y(\mathfrak{g}))$ as a $\mathbb{C}[\hbar]$ -algebra, Ψ_{\hbar} is surjective.

The composition π of the isomorphism $R_{\hbar}(Y(\mathfrak{g}))/\hbar R_{\hbar}(Y(\mathfrak{g})) \rightarrow \text{gr } Y(\mathfrak{g})$ with the quotient homomorphism $R_{\hbar}(Y(\mathfrak{g})) \rightarrow R_{\hbar}(Y(\mathfrak{g}))/\hbar R_{\hbar}(Y(\mathfrak{g}))$ satisfies

$$\hbar^k x_{ik}^{\pm}, \hbar^k h_{ir} \mapsto \bar{x}_{ik}^{\pm}, \bar{h}_{ik} \quad \forall i \in I \text{ and } k \geq 0.$$

Moreover, $\pi \circ \Psi_{\hbar}$ sends the ideal $\hbar Y_{\hbar}(\mathfrak{g})$ to zero and thus factors through the quotient $Y_{\hbar}(\mathfrak{g})/\hbar Y_{\hbar}(\mathfrak{g})$ to give $\Psi_0 : Y_0(\mathfrak{g}) \rightarrow \text{gr } Y(\mathfrak{g})$. After using Proposition 2.7 and Theorem 6.9 to identify both the domain and codomain of Ψ_0 with $U(\mathfrak{s})$, Ψ_0 becomes the identity map.

Now suppose that there is a nonzero $X \in \text{Ker } \Psi_{\hbar}$. Let m be the maximal non-negative integer such that $X = \hbar^m Y$ for some $Y \in Y_{\hbar}(\mathfrak{g})$ (that m is finite follows from the fact that $Y_{\hbar}(\mathfrak{g})$ is graded with $\deg \hbar = 1$). Since Ψ_{\hbar} is a $\mathbb{C}[\hbar]$ -algebra morphism and $R_{\hbar}(Y(\mathfrak{g}))$ is torsion free, $Y \in \text{Ker } \Psi_{\hbar}$. By maximality of m , the image \bar{Y} of Y in $Y_0(\mathfrak{g})$ is nonzero. Since $\Psi_0 : Y_0(\mathfrak{g}) \rightarrow \text{gr } Y(\mathfrak{g})$ is an isomorphism, $\Psi_0(\bar{Y}) \neq 0$. This is a contradiction as $\Psi_0(\bar{Y}) = \overline{\Psi_{\hbar}(Y)} = 0$. Therefore Ψ_{\hbar} is injective, and thus an isomorphism. \square

REFERENCES

- [AG] A. Appel, S. Gautam, *An explicit isomorphism between quantum and classical \mathfrak{sl}_n* . [arXiv:1712.03601](#).
- [AMR] D. Arnaudon, A. Molev, E. Ragoucy, *On the R -matrix realization of Yangians and their representations*, Ann. Henri Poincaré **7** (2006), no. 7-8, 1269–1325. [arXiv:math/0511481](#).
- [Be] D. Bernard, *Vertex operator representations of the quantum affine algebra $\mathcal{U}_q(B_r^{(1)})$* Lett. Math. Phys. **17** (1989), no. 3, 239–245.
- [BeTs] M. Bershtein, A. Tsymbaliuk, *Homomorphisms between different quantum toroidal and affine Yangian algebras*, [arXiv:1512.09109](#).
- [BTM] D. Bernard, J. Thierry-Mieg, *Level one representations of the simple affine Kac-Moody algebras in their homogeneous gradations*, Comm. Math. Phys. **111** (1987), no. 2, 181–246.

- [ChJi] V. Chari, N. Jing, *Realization of level one representations of $U_q(\widehat{\mathfrak{g}})$ at a root of unity*, Duke Math. J. **108** (2001), no. 1, 183–197. [arXiv:math/9909118](#).
- [DiKh] J. Ding, S. Khoroshkin, *Weyl group extension of quantized current algebras*, Transform. Groups **5** (2000), no. 1, 35–59. [arXiv:math/9804139](#).
- [Dr] V. Drinfeld, *Degenerate affine Hecke algebras and Yangians*, (Russian) Funktsional. Anal. i Prilozhen. **20** (1986), no. 1, 69–70.
- [FLM] I. Frenkel, J. Lepowsky, A. Meurman, *Vertex operator algebras and the Monster*, Pure and Applied Mathematics, 134. Academic Press, Inc., Boston, MA, 1988. liv+508 pp.
- [FrJi] I. Frenkel, N. Jing, *Vertex representations of quantum affine algebras*, Proc. Nat. Acad. Sci. U.S.A. **85** (1988), no. 24, 9373–9377.
- [FrKa] I. Frenkel, V. Kac, *Basic representations of affine Lie algebras and dual resonance models*, Invent. Math. **62** (1980/81), no. 1, 23–66.
- [GNOS] P. Goddard, W. Nahm, W. Olive, W. Schwimmer, *Vertex operators for non-simply-laced algebras*, Comm. Math. Phys. **107** (1986), no. 2, 179–212.
- [GNW] N. Guay, H. Nakajima, C. Wendlandt, *Coproduct for Yangians of affine Kac-Moody algebras*, submitted for publication, see version available at <https://sites.ualberta.ca/~nguay/publications.html>.
- [GTL1] S. Gautam, V. Toledano Laredo, *Yangians and quantum loop algebras*, Selecta Math. (N.S.) **19** (2013), no. 2, 271336. [arXiv:1012.3687](#).
- [GTL2] S. Gautam, V. Toledano Laredo, *Yangians, quantum loop algebras, and abelian difference equations*, J. Amer. Math. Soc. **29** (2016), no. 3, 775–824. [arXiv:1310.7318](#).
- [Gu] N. Guay, *Affine Yangians and deformed double current algebras in type A*, Adv. Math. **211** (2007), no. 2, 436–484.
- [Io] K. Iohara, *Bosonic representations of Yangian double $DY_h(\mathfrak{g})$ with $\mathfrak{g} = \mathfrak{gl}_N, \mathfrak{sl}_N$* , J. Phys. A **29** (1996), no. 15, 4593–4621. [arXiv:q-alg/9603033](#).
- [IoKo] K. Iohara, M. Kohno, *A central extension of $DY_h(\mathfrak{gl}_2)$ and its vertex representations*, Lett. Math. Phys. **37** (1996), no. 3, 319–328. [arXiv:q-alg/9603032](#).
- [Ji1] N. Jing, *Twisted vertex representations of quantum affine algebras*, Invent. Math. **102** (1990), no. 3, 663–690.
- [Ji2] N. Jing, *Quantum Kac-Moody algebras and vertex representations*, Lett. Math. Phys. **44** (1998), no. 4, 261–271. [arXiv:math/9802036](#).
- [Ji3] N. Jing, *Level one representations of $U_q(C_2^{(1)})$* , Proc. Amer. Math. Soc. **127** (1999), no. 1, 21–27. [arXiv:q-alg/9710023](#).
- [Ji4] N. Jing, *Quantum Z-algebras and representations of quantum affine algebras*, Comm. Algebra **28** (2000), no. 2, 829–844. [arXiv:math/9806043](#).
- [JiMi] N. Jing, K. Misra, *Vertex operators of level-one $U_q(B_n^{(1)})$ -modules*, Lett. Math. Phys. **36** (1996), no. 2, 127–143.
- [JKM] N. Jing, Y. Koyama, K. Misra, *Level one representations of quantum affine algebras $U_q(C_n^{(1)})$* , Selecta Math. (N.S.) **5** (1999), no. 2, 243–255. [arXiv:math/9802123](#).
- [Ka] V. Kac, *Infinite-dimensional Lie algebras*, Cambridge University Press, Cambridge, third edition, 1990.
- [Kh] S. Khoroshkin, *Central extension of the Yangian double*, Algèbre non commutative, groupes quantiques et invariants (Reims, 1995), 119–135, Smin. Congr., **2**, Soc. Math. France, Paris, 1997. [arXiv:q-alg/9602031](#).
- [KhTo] S. Khoroshkin, V. Tolstoy, *Yangian double*, Lett. Math. Phys. **36** (1996), no. 4, 373–402.
- [KI] C. Kassel, *Kähler differentials and coverings of complex simple Lie algebras extended over a commutative algebra*, Proceedings of the Luminy conference on algebraic K-theory (Luminy, 1983), J. Pure Appl. Algebra **34** (1984), no. 2-3, 265–275.
- [Ko] R. Kodera, *Affine Yangian action on the Fock space*. [arXiv:1506.01246](#).
- [KSU] K. Kimura, J. Shiraishi, J. Uchiyama, *A level-one representation of the quantum affine superalgebra $U_q(\widehat{\mathfrak{sl}}(M+1|N+1))$* , Comm. Math. Phys. **188** (1997), no. 2, 367–378. [arXiv:q-alg/9605047](#).
- [Le] S.Z. Levendorski, *On PBW bases for Yangians*, Lett. Math. Phys. **27** (1993), no. 1, 37–42.
- [LeLi] J. Lepowsky, H. Li, *Introduction to vertex operator algebras and their representations*, Progress in Mathematics, **227**. Birkhuser Boston, Inc., Boston, MA, 2004. xiv+318 pp.
- [Mo] A. Molev, *Yangians and Classical Lie Algebra*, Mathematical Surveys and Monographs **143**, American Mathematical Society, Providence, RI, 2007, xviii+400 pp.
- [MRY] R. Moody, S. Rao, T. Yokonuma, *Toroidal Lie algebras and vertex representations*, Geom. Dedicata **35** (1990), no. 1-3, 283–307.
- [Na] H. Nakajima, *Quiver varieties and finite-dimensional representations of quantum affine algebras*, J. Amer. Math. Soc. **14** (2001), no. 1, 145–238. [arXiv:math/9912158](#).
- [Ne] E. Neher, *An introduction to universal central extensions of Lie superalgebras*, Groups, Rings, Lie and Hopf Algebras, Mathematics and Its Applications. Springer, Boston, MA, pp. 141–166.

- [Sa] Y. Saito, *Quantum toroidal algebras and their vertex representations*, Publ. Res. Inst. Math. Sci. **34** (1998), no. 2, 155–177. [arXiv:q-alg/9611030](#).
- [YaZh1] Y. Yang, G. Zhao, *Cohomological Hall algebras and affine quantum groups*, Selecta Math. (N.S.) (2017). [arXiv:1604.01865](#).
- [YaZh2] Y. Yang, G. Zhao, *The PBW Theorem for the affine Yangians*, in preparation.

N.G.: UNIVERSITY OF ALBERTA, DEPARTMENT OF MATHEMATICS, EDMONTON, AB T6G 2G1, CANADA.
E-mail address: nguay@ualberta.ca

V.R.: UNIVERSITY OF YORK, DEPARTMENT OF MATHEMATICS, YORK, YO10 5DD, UK AND VILNIUS UNIVERSITY, INSTITUTE OF THEORETICAL PHYSICS AND ASTRONOMY, SAULËTEKIO AV. 3, VILNIUS 10257, LITHUANIA.
E-mail address: vidas.regelskis@gmail.com

C.W.: UNIVERSITY OF ALBERTA, DEPARTMENT OF MATHEMATICS, EDMONTON, AB T6G 2G1, CANADA.
E-mail address: cwendlan@ualberta.ca