



This is a repository copy of *Induction of silicon defences in wheat landraces is local not systemic and driven by mobilisation of soluble silicon to damaged leaves.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/200434/>

Version: Accepted Version

---

**Article:**

Thorne, S.J. [orcid.org/0000-0003-0476-8466](https://orcid.org/0000-0003-0476-8466), Maathuis, F.J.M. and Hartley, S.E. (2023) Induction of silicon defences in wheat landraces is local not systemic and driven by mobilisation of soluble silicon to damaged leaves. *Journal of Experimental Botany*. erad224. ISSN 0022-0957

<https://doi.org/10.1093/jxb/erad224>

---

© 2023 The Author(s). Except as otherwise noted, this author-accepted version of a journal article published in *Journal of Experimental Botany* is made available via the University of Sheffield Research Publications and Copyright Policy under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Induction of silicon defences in wheat landraces is local not systemic and driven by mobilisation of soluble silicon to damaged leaves

Sarah J. Thorne <sup>1</sup>\* [ID](#), Frans J. M. Maathuis <sup>2</sup>[ID](#), and Susan E. Hartley <sup>1</sup>[ID](#)

<sup>1</sup> Plants, Photosynthesis, and Soil, School of Biosciences, University of Sheffield, Sheffield, S10 2TN, United Kingdom

<sup>2</sup> Department of Biology, University of York, York, YO10 5DD, United Kingdom

\* Correspondence: [s.j.thorne@sheffield.ac.uk](mailto:s.j.thorne@sheffield.ac.uk)

[frans.maathuis@york.ac.uk](mailto:frans.maathuis@york.ac.uk)

[s.hartley@sheffield.ac.uk](mailto:s.hartley@sheffield.ac.uk)

Running title: Mechanism of induction of silicon defences in wheat

Date of submission: 3/5/23

Number of tables: 0

Number of figures: 5

Word count: 4207

Supplementary tables: 3

Supplementary figures: 4

## 1. Highlight

Silicon is an important defence in crops. Here, a new mechanism involving the movement of soluble silicon in the phloem is proposed to explain localised induction of silicon defences.

## 2. Abstract

- In response to herbivory, many grasses, including crops such as wheat, accumulate significant levels of silicon (Si) as an antiherbivore defence. Damage-induced increases in Si can be localised in damaged leaves or more systemic, but the mechanisms leading to these differences in Si distribution remain untested.
- Ten genetically diverse wheat landraces (*Triticum aestivum*) were used to assess genotypic variation in Si induction in response to mechanical damage and how this was affected by exogenous Si supply. Total and soluble Si levels were measured in damaged and undamaged leaves, as were Si levels in the phloem, to test how Si was allocated to different parts of the plant after damage.
- Localised, but not systemic, induction of Si defences occurred, more pronounced when plants had supplemental Si. Damaged plants had significant increases in Si concentration in their damaged leaves, while the Si concentration in undamaged leaves decreased, such that there was no difference in the average Si concentration of damaged and undamaged plants.
- The increased Si in damaged leaves was due to the redirection of soluble Si, present in the phloem, from undamaged to damaged plant parts, potentially a more cost-effective defence mechanism for plants than increased Si uptake.

**Keywords:** damage, genotypic variation, localised response, phloem, reallocation, silicon, wheat (*Triticum aestivum*)

## 3. Introduction

Plants have many different types of defences against herbivores, including both physical and chemical defences (Howe and Jander, 2007). Particularly in grasses, which include many cereal crops, high silicon (Si) accumulation is an effective antiherbivore defence (Massey *et al.*, 2009; Han *et al.*, 2015; Singh *et al.*, 2020; Waterman *et al.*, 2021). Grasses deposit Si in structures such as phytoliths and silicified spines (Hartley *et al.*, 2015). Phytoliths increase leaf abrasiveness making tissues less palatable (Hall *et al.*, 2020), which deters both insect and mammalian herbivores. Furthermore, phytoliths can reduce herbivore digestive efficiencies and hence limit their growth and development (Massey and Hartley, 2006, 2009; Massey *et al.*, 2008, 2009). In addition, Si deposited

33 in the apoplast may act as a physical barrier, potentially preventing the release of insect oral  
34 secretions and oviposition fluids, known as effectors, which are used by herbivores to recognise  
35 compatible host plants (Coskun *et al.*, 2019; Singh *et al.*, 2020).

36 At least in grasses, Si is an inducible defence (Massey *et al.*, 2007): in response to herbivory, overall  
37 Si accumulation increases, and this is correlated with reduced herbivory (Reynolds *et al.*, 2012;  
38 Hartley *et al.*, 2015; Hall *et al.*, 2020). However, the Si response to herbivory varies significantly  
39 between both plant species and genotypes (Hartley and DeGabriel, 2016). For example, different  
40 patterns of Si accumulation and deposition were found in three species of *Festuca* in response to  
41 artificial damage and Si supply (Hartley *et al.*, 2015), and similar genotypic variation in Si uptake and  
42 deposition have been found in wheat (*Triticum aestivum*; Thorne *et al.*, 2021) and rice (Talukdar *et al.*,  
43 2019). Soininen *et al.* (2013) reported both within and between species variation in Si  
44 accumulation among grasses in response to damage.

45 Plants accumulate Si in the form of silicic acid from the soil using a pathway that is relatively well  
46 characterised in rice. Si is transported through the roots by the serial action of two transporters, Lsi1  
47 and Lsi2 (Ma and Yamaji, 2015). Silicic acid is then loaded into the xylem by the action of a third  
48 transporter, Lsi3 (Huang *et al.*, 2022). Subsequently, Si is translocated to the shoots *via* the  
49 transpiration stream (Ma and Yamaji, 2015). A fourth transporter, Lsi6, is required for xylem  
50 unloading of silicic acid (Yamaji *et al.*, 2008). Lsi6, Lsi3, and Lsi2 are also highly expressed in the node  
51 where they are involved in intervascular transfer to direct Si distribution within the plant (Yamaji *et al.*,  
52 2015). High levels of silicic acid result in its autopolymerisation into immobile silica, which is the  
53 predominant form of Si in plants (Yoshida *et al.*, 1962). Thus, plant Si accumulation is affected by  
54 both the transpiration rate and the activity of Si transporters. Genotypic variation in Si accumulation  
55 has been linked to both differences in transpiration (McLarnon *et al.*, 2017) and differences in the  
56 abundance of Si transporters (Ma *et al.*, 2007b). However, a range of biotic and abiotic factors have  
57 been shown to influence genotypic variation in Si accumulation including damage (Soininen *et al.*,  
58 2013), nitrogen availability and plant competition (de Tombeur *et al.*, 2022), and climatic factors  
59 (Johnson *et al.*, 2023).

60 Although damage-induced localised increases in tissue Si have been reported in several grasses  
61 (Hartley *et al.*, 2015; McLarnon *et al.*, 2017) and in cucumber (Islam *et al.*, 2020), the mechanism  
62 underpinning this localised increase in Si remains to be determined. One hypothesis is that the  
63 uptake of Si from the soil increases, and this additional Si is then directed specifically to damaged  
64 tissue. It is unlikely that differences in transpiration rate explain increased Si accumulation in

65 damaged leaves, as a localised damage response can be observed even in plants where transpiration  
66 rates were greatly reduced (McLarnon *et al.*, 2017). Instead, changes in Si transporter abundance  
67 and activity may be responsible, as has been suggested for rice (Ye *et al.*, 2013), though other  
68 studies have not found evidence for this mechanism (McLarnon *et al.*, 2017).

69 A second hypothesis proposes that part or all the “extra” Si gets relocated from undamaged tissue to  
70 damaged tissue. Si accumulation is an active process which involves significant energetic costs (de  
71 Tombeur *et al.*, 2023) and thus the mobilisation of the Si already present in the plant towards the  
72 sites where it is most needed to repel attack may be a more beneficial strategy than increased  
73 uptake from the soil. However, once Si is deposited, it cannot be remobilised (Samuels *et al.*, 1991),  
74 and thus this hypothesis could only pertain to the soluble Si fraction in plants. It has not yet been  
75 experimentally tested, nor has the interaction between genetic variation in patterns of Si  
76 accumulation, damage, and Si supply.

77 Here, the effect of mechanical damage on Si accumulation in a number of genetically diverse  
78 landraces of wheat (*Triticum aestivum*) was investigated and the following questions were  
79 addressed:

- 80 1. Is the induction of silicon-based defences in wheat a localised or systemic response, and  
81 how is the magnitude and pattern of induction affected by Si supply?
- 82 2. If the response is localised, by what mechanism is this achieved? Is the source of the  
83 increased Si in damaged leaves due to *de novo* uptake, or to reallocation of Si, in soluble  
84 form, to damaged leaves from other parts of the plant?

85

## 86 **4. Methods**

### 87 **4.1. Genotype selection, growth conditions, and experimental treatments**

88 To investigate whether there is genotypic variation in the effect of damage on Si accumulation, ten  
89 wheat landraces that differed significantly in their Si accumulation were selected: five high and five  
90 low Si-accumulating landraces (Thorne *et al.*, 2021, Supplementary Table S1). A balanced factorial  
91 experimental design was used with three plants per landrace per damage versus undamaged  
92 treatment. Two seeds were planted in 1 L pots filled with a 2:1 mix of sand and terragreen. One  
93 week after germination, seedlings were thinned to one plant per pot. After thinning, all plants were  
94 fed twice weekly with 200 mL half-strength Hoagland’s solution supplemented with 1.8 mM  
95 dissolved sodium metasilicate ( $\text{Na}_2\text{SiO}_3 \cdot 9\text{H}_2\text{O}$ ). Plants were watered as required.

96 Two of the high and two of the low Si-accumulating landraces were then selected to examine the  
97 effect of Si availability on Si accumulation in response to damage. To control Si supply effectively,  
98 plants were grown hydroponically. Seeds were germinated in sand for 10-11 days, and then  
99 seedlings were transferred to 9 L plastic hydroponics boxes, filled with half-strength Hoagland's  
100 solution. The pH was adjusted to 5.6-6.0 using 1 M HCl or 0.1 M KOH. The nutrient solution was  
101 changed every 3-4 days. The hydroponics solutions were aerated throughout the experiment. A  
102 balanced factorial experimental design was used with plants either damaged or not and  
103 supplemented with Si (+Si) or not (- Si). Plants supplemented with Si received Hoagland's solution  
104 containing 1.8 mM Si. Sodium chloride (NaCl) was used to balance sodium levels for plants not  
105 supplemented with Si. Three plants per landrace, per Si fertilisation level, and per damage treatment  
106 were used.

107 To identify the source of the additional Si leading to the localised induction in damaged leaves, one  
108 landrace (L1) was selected. A balanced factorial experimental design was used with plants either  
109 damaged or not. All plants were grown hydroponically with 1.8 mM Si for the first three weeks.  
110 Immediately prior to damage, half the plants were moved to a medium without Si supplementation  
111 (+/- Si) while the remaining plants continued to be grown with Si supplementation (+/+ Si). Plants  
112 were harvested three days after the first, second, and fourth damage. Three plants per treatment  
113 were used.

114 All plants were grown under controlled glasshouse conditions (16 h daylight; 20 °C /15 °C day/night).  
115 At harvest, roots were washed in deionised water and excess water was removed. Leaf, stem, and  
116 root fresh weight was recorded. Plants were oven-dried at 70 °C until constant mass was achieved,  
117 then dry weight was recorded.

#### 118 **4.2. Damage Treatment**

119 A damage treatment was started four weeks after germination and continued for three weeks. In  
120 the case of the ten landraces grown in soil, plants were damaged three times a week, whereas plants  
121 grown hydroponically were damaged twice weekly. A damage treatment involved removing  
122 approximately half of a newly produced leaf along the midrib. Plants were harvested one day after  
123 the final damage event, seven weeks after germination unless otherwise stated. Plants that were not  
124 mechanically damaged were labelled as undamaged plants. The weight and Si concentration of  
125 damaged and undamaged leaves of damaged plants were analysed separately.

126

127 **4.3. Si Measurements**

128 Portable X-ray fluorescence spectroscopy (P-XRF) was used to measure the leaf Si concentration of  
129 all plants and the root Si concentration of plants grown hydroponically (Reidinger *et al.*, 2012). Dried  
130 leaf material was ball-milled (Retsch MM400 Mixer mill, Haan, Germany) and ground material was  
131 pressed at 10 tons into pellets using a manual hydraulic press with a 13 mm die (Specac, Orpington,  
132 UK). Si analysis (% Si dry weight) was performed using a commercial P-XRF instrument (Nitron  
133 XL3t900 GOLDD analyser: Thermo Scientific Winchester, UK) held in a test stand (SmartStand,  
134 Thermo Scientific, Winchester, UK). The P-XRF machine was calibrated using Si-spiked synthetic  
135 methyl cellulose (Sigma-Aldrich, product no. 274429) and validated using Certified Reference  
136 Materials of NCS DC73349 'Bush branches and leaves' obtained from China National Analysis Center  
137 for Iron and Steel. To avoid signal loss by air absorption, the analyses were performed under a  
138 helium atmosphere (Reidinger *et al.*, 2012). A reading of each side of the pellet was taken,  
139 approximately one hour apart, to account for *u*-drift in the instrument (i.e. variation in readings  
140 between consecutive runs using identical parameters; Johnson, 2014). The two readings were  
141 averaged to obtain the Si concentration (%). The Si concentration of damaged and undamaged  
142 leaves of damaged plants was analysed separately. Si content (mg) was calculated as:

143 
$$Si\ content\ (mg) = \frac{Si\ concentration\ (\%)}{100} \times dry\ weight\ (mg)$$

144 **4.4. Measuring soluble Si**

145 To measure the amount of soluble Si present in the leaves, one landrace (L1) was grown  
146 hydroponically with 1.8 mM Si for seven weeks. Four leaves were damaged simultaneously (~17 % of  
147 the leaves) and plants were harvested five days after the damage. The soluble Si concentration was  
148 measured for ten damaged plants and six undamaged plants. Damaged and undamaged leaves of  
149 damaged plants were analysed separately.

150 Leaf soluble Si was measured using an adapted version of the molybdenum method of measuring  
151 plant available Si in the soil (Sauer *et al.*, 2006). Briefly, samples were oven-dried at 70 °C and ground  
152 in a ball mill. A total of 2 g of sample was added to a 50 mL polyethylene tube with 20 mL 0.01 M  
153 CaCl<sub>2</sub>. Samples were shaken slowly at 30 rpm for 16 h then centrifuged at 2000 rpm for 10 min. The  
154 supernatant was passed through filter paper and the resulting sample was used for Si determination  
155 by the molybdenum assay. For the assay, 1 mL sample, 30 mL 20 % acetic acid, and 10 mL  
156 ammonium molybdate (54 g L<sup>-1</sup>) were mixed in a 50 mL polyethylene tube. Samples were mixed and  
157 then left to stand for 5 minutes. 5 mL 20 % (w/v) tartaric acid and 1 mL of reducing solution were

158 added. The reducing solution comprised 8 g L<sup>-1</sup> Na<sub>2</sub>SO<sub>3</sub>, 1.6 g L<sup>-1</sup> 1-amino-2-naphthol-4-sulfonic acid,  
159 and 100 g L<sup>-1</sup> NaHSO<sub>3</sub> dissolved in deionised water. A final volume of 50 mL was made using 20 %  
160 acetic acid. Samples were left at room temperature for 30 min for colour development then the  
161 absorbance at 810 nm was measured. A standard curve was created using 0, 0.1, 0.5, 1, 5, and 10 µg  
162 Si mL<sup>-1</sup> to determine the Si concentration of the samples. The amount of soluble Si measured using  
163 the molybdenum assay was compared to the total Si concentration measured using XRF to estimate  
164 the total soluble Si present.

#### 165 **4.5. Measuring soluble Si in phloem exudate**

166 Phloem exudate was collected using the EDTA-mediated method as described in Xu *et al.* (2019)  
167 using plants grown hydroponically with 1.8 mM supplementary Si for three weeks. Three leaves per  
168 plant were damaged, then half the plants were moved to medium without supplementary Si.  
169 Phloem exudate was collected 24 h later using three leaves per plant. Phloem exudate was collected  
170 separately for damaged and undamaged leaves of damaged plants, and for undamaged plants. As a  
171 negative control, phloem exudate was collected in water (no EDTA) from undamaged plants. Three  
172 plants per treatment were used. The Si concentration of the phloem exudate was then analysed  
173 using the molybdenum method as described above.

#### 174 **4.6. Measuring expression of Si transporters**

175 To investigate whether damage affected Si transporter gene expression, the expression levels of the  
176 Si transporters *Lsi3* and *Lsi6* were determined using reverse transcriptase quantitative polymerase  
177 chain reaction (RT-qPCR). These transporters were chosen as they are known to be expressed in the  
178 shoots, whereas the other Si transporters, *Lsi1* and *Lsi2*, are expressed predominantly in the roots  
179 (Yamaji *et al.*, 2015). Initial experiments on the leaves of wheat landraces confirmed this with only  
180 very low levels of expression of *Lsi1* and *Lsi2*, but significant expression of *Lsi3* and *Lsi6*, found. Plants  
181 of the landrace L1 were grown hydroponically with 1.8 mM Si for four weeks. Plants were then  
182 moved to – Si medium immediately prior to the first damage event and then subject to either one or  
183 two damage events (with the second damage event occurring 24 hours after the first damage  
184 event). Leaf samples were taken from the middle of the newest expanded leaf 4, 24, and 48 h after  
185 one or two damages. Initially two samples were analysed for each time point and subsequently up to  
186 five biological replicates from a given timepoint were analysed once the most relevant timepoints  
187 had been identified.

188 Primers were designed to match all homoeologs, based on existing wheat sequences where  
189 available, or on homology to the barley sequence (Supplementary Table 2). Leaf tissue was collected



190 and ground under liquid nitrogen using a mortar and pestle and RNA extracted using a RNeasy kit  
191 with DNase treatment (Qiagen), according to the manufacturer's instructions. cDNA synthesis was  
192 performed using Reverse Transcriptase Superscript II M-MLV (Invitrogen) on 500 ng RNA. RT-qPCR  
193 was performed using a QuantiNova SYBR green PCR kit (Qiagen) with 2  $\mu$ L cDNA (diluted 1:10) and  
194 700 nM primer on a Rotor-Gene Q PCR machine (Qiagen). Amplification was performed as follows:  
195 95 °C for 2 min; 40 cycles of 95 °C 5 s and 60 °C for 10 s; determination of melt curve. *Actin* and *TEF1*  
196 were used as reference genes. The qPCR results were analysed using a variation of the  $2^{\text{DDCt}}$  method  
197 as described in Muller *et al.* (2002) and an adapted version of the Q-gene excel software (Simon,  
198 2003).

#### 199 **4.7. Statistical Analysis**

200 All statistical analyses were performed using R software (version 4.2.0, R Core Team, 2022).  
201 Summary statistics were calculated using the Rmisc package (Hope, 2013) and graphs were  
202 produced using the ggplot2 package (Wickham, 2016). Two-way and three-way analyses of variance  
203 (ANOVA) were used to test the effect of Si supply, damage, number of damage events, and landrace  
204 on Si concentration, as relevant to each experiment. Due to the lack of independence between  
205 damaged and undamaged leaves from damaged plants, ANOVAs were performed separately  
206 comparing undamaged plants to either damaged or undamaged leaves of damaged plants.  
207 Additionally, the average leaf Si concentration of damaged plants was calculated by averaging the Si  
208 concentration of damaged and undamaged leaves, accounting for differences in the proportion of  
209 the leaves that were damaged or undamaged. ANOVA was then used to compare the average leaf Si  
210 concentration of damaged plants to undamaged plants.

211 Data normality was checked using Shapiro tests and homogeneity of variance was tested using  
212 Levene's tests. To satisfy the test assumptions, Si concentration was logit transformed and Si content  
213 was log transformed. No transformation was applied to gene expression or soluble Si data. Paired *t*-  
214 tests were used to test for localised induction of Si defences between damaged and undamaged  
215 leaves of damaged plants. A significance level of  $P < 0.05$  was used for all analyses. Significant results  
216 were analysed by performing Tukey's Honest Significance Difference (HSD) *post-hoc* tests using the  
217 emmeans package (Lenth, 2021).

## 218 **5. Results**

### 219 **5.1. Damage results in a localised increase in Si**

220 Averaged across all ten landraces, repeated damage significantly increased Si accumulation in  
221 damaged leaves of damaged plants when compared to undamaged plants (Figure 1;  $F_{1,40} = 67.9$ ,  $P <$

222 0.001). The biggest percentage increase was  $107.6 \pm 28.4$  % in the L3 landrace, compared to an  
223 increase of only  $34.2 \pm 9.3$  % in H5. The localised induction in damaged leaves was significant in all  
224 landraces except landraces H5 and L2, though there was still a trend towards significance even in  
225 these landraces ( $P = 0.055$  and  $0.052$  respectively). Overall, the interaction between landrace and  
226 damage was not significant ( $F_{9,40} = 1.5$ ,  $P = 0.171$ ).

227 Si accumulation was lower in the undamaged leaves of the damaged plants, averaged across all ten  
228 landraces, when compared to the levels in undamaged plants (Figure 1;  $F_{1,40} = 9.2$ ,  $P = 0.004$ ).  
229 However, post-hoc testing revealed that the decrease in undamaged leaves was specific to the low  
230 Si-accumulating landraces L1 ( $P = 0.013$ ) and L5 ( $P = 0.026$ ), with L3 and L4 showing a similar trend  
231 albeit marginally non-significant ( $P = 0.051$  and  $0.054$  respectively). The lack of increase in Si in  
232 undamaged leaves on damaged plants, or even decreases in some landraces, means there was no  
233 overall increase in leaf Si in damaged plants compared to undamaged ones for any landrace  
234 (Supplementary Figure 1). Thus, the induction of Si defences occurred only locally and was not  
235 systemic.

236 On average, high Si-accumulating landraces had higher shoot Si concentrations compared to low Si  
237 accumulating landraces, with an average Si concentration of  $2.97 \pm 0.17$  for undamaged high Si  
238 landraces compared to  $1.89 \pm 0.09$  for low Si landraces. There was no significant correlation between  
239 leaf Si in undamaged plants and the relative increase in leaf Si due to damage ( $r = -0.32$ ,  $P = 0.364$ ).

## 240 **5.2. The effect of damage on Si accumulation depends on Si availability**

241 Repeated damage significantly increased the leaf Si accumulation in the damaged leaves of the four  
242 landraces tested (Figure 2;  $F_{1,47} = 216.0$ ,  $P < 0.001$ ). This was the case in both the +Si (1.8 mM Si  
243 supplied to the hydroponic medium) and -Si (no exogenous Si supply) treatments, although this  
244 increase was significantly greater in the +Si treatment (Figure 2; Si availability x damage interaction:  
245  $F_{1,47} = 23.1$ ,  $P < 0.001$ ).

246

247 As with the ten landraces, damage resulted in only a localised induction of Si defences. Damage  
248 significantly decreased leaf Si accumulation in the undamaged leaves of damaged plants compared  
249 to undamaged plants in the +Si treatment ( $F_{1,24} = 18.2$ ,  $P < 0.001$ ), but this did not occur in the -Si  
250 plants, although leaf Si in this treatment was already an order of magnitude lower ( $F_{1,23} = 0.5$ ,  $P =$   
251  $0.503$ ). There was no significant variation in the Si response to damage among landraces in either  
252 the -Si or +Si treatments. No significant effect of damage on root Si accumulation was found,

253 although, as in the leaf tissue, there was a positive effect of Si supply on Si concentrations  
254 (Supplementary Figure 2).

255

### 256 **5.3. Si is redirected towards damaged leaves in soluble form**

257 To determine whether the observed increase in Si accumulation in damaged leaves was the result of  
258 *de novo* Si uptake or due to reallocation of Si from undamaged leaves, the exogenous Si supply was  
259 removed from plants from the L1 landrace prior to a series of damage events. Increasing the number  
260 of damage events significantly increased Si accumulation in damaged leaves of damaged plants,  
261 although this response was significantly higher in +/+Si (continuous Si supply) plants compared to  
262 +/-Si (Si removed before damage) plants (Figure 3; Si x number of damage events interaction:  $F_{2,24} =$   
263  $21.6, P < 0.001$ ). For +/+Si plants, a greater number of damage events led to a more pronounced Si  
264 increase in damaged leaves. However, in the case of +/-Si plants, the opposite trend occurred: each  
265 damage event was associated with a reduced localised induction, possibly because the total Si  
266 concentration in these plants was “diluted” down by to continuing growth of plant tissue. Thus, after  
267 four damage events, the increase in Si in damaged leaves in +/- Si plants was reduced to 81 % of the  
268 increase occurring after a single damage, whereas for +/+ Si plants, the increase in Si after four  
269 damage events was over three times as great (302 %) of that after a single damage.

270 The distribution of Si in the root, stem, and leaf tissues of damaged and undamaged plants, after  
271 one, two, or four damage treatments, was calculated for both +/+Si and +/-Si plants. The total  
272 amount of Si accumulated in the plants did not differ significantly between damaged and  
273 undamaged plants (Figure 4;  $F_{1,23} = 0.1, P = 0.778$ ), regardless of the number of damage events or  
274 the nature of the Si supply. Si content increased with an increasing number of damage events only  
275 for +/+ Si plants, as would be expected in the case of continuous Si supply. There was no significant  
276 difference in root or stem Si concentration between damaged and undamaged plants, although Si  
277 supply significantly increased both root and stem Si concentration (Supplementary Figure 3). For +/-  
278 Si plants, increasing the number of damage events significantly decreased the stem Si concentration  
279 ( $F_{2,24} = 20.5, P < 0.001$ ).

280 In the absence of external Si supply (+/- Si plants) and given the similarity of total Si content  
281 between damaged and undamaged plants, any increase in Si in damaged leaves most likely  
282 originates from the relocation of Si from undamaged leaves. The soluble Si fraction of damaged and  
283 undamaged leaves was measured in plants of the landrace L1 grown hydroponically for 7 weeks and  
284 damaged once, in order to assess whether it is sufficiently large to account for the extra Si found in  
285 damaged leaves, without the need for additional uptake. Undamaged plants had a soluble Si content

286 of  $2.2 \pm 0.1 \text{ mg g}^{-1}$ , compared to  $4.0 \pm 0.1 \text{ mg g}^{-1}$  for damaged leaves and  $1.8 \pm 0.1 \text{ mg g}^{-1}$  for  
287 undamaged leaves of damaged plants. Based on the difference between undamaged leaves of  
288 damaged plants and undamaged plants, this means that on average there will be  $0.4 \text{ mg g}^{-1}$  soluble  
289 Si available to move from undamaged leaves to damaged leaves of damaged plants. The soluble Si  
290 from undamaged leaves needs to provide the  $1.8 \text{ mg g}^{-1}$  increase in soluble Si observed in damaged  
291 leaves compared to undamaged plants. Thus, as long as the amount of undamaged leaves is 4.5  
292 times as the weight of damaged leaves on a plant, there is sufficient soluble Si to relocate from  
293 undamaged to damaged leaves and account for all the localised induction in Si defences observed.

294 It was hypothesised that the redistribution of Si from undamaged to damaged leaves is the result of  
295 soluble Si being moved *via* the phloem. Supporting this idea, significant concentrations of soluble Si  
296 were measured in the phloem exudate. For +/+ Si plants, soluble Si in the phloem was found to be:  
297  $1.9 \pm 0.6 \text{ mM}$  in undamaged plants,  $1.1 \pm 0.5 \text{ mM}$  in undamaged leaves of damaged plants, and  $1.9 \pm$   
298  $0.4 \text{ mM}$  in damaged leaves. The phloem Si of +/- Si plants was lower at:  $1.1 \pm 0.5 \text{ mM}$  in undamaged  
299 plants,  $0.9 \pm 0.7 \text{ mM}$  in undamaged leaves of damaged plants, and  $1.0 \pm 0.4 \text{ mM}$  in damaged leaves.  
300 The Si in the phloem of damaged leaves was significantly higher in the +/+ Si plants than in the +/- Si  
301 plants ( $F_{1,8} = 9.4$ ,  $P = 0.016$ ) but there was no difference between the phloem Si levels in the  
302 undamaged leaves of damaged plants from the Si two treatments ( $F_{1,8} = 2.0$ ,  $P = 0.196$ ). This suggests  
303 that soluble Si is being directed from undamaged to damaged leaves, particularly in plants with  
304 continuous Si supply, supporting the hypothesis that Si is being mobilised from undamaged to  
305 damaged leaves where Si can be deposited as a physical defence.

306 Soluble Si may be loaded into the phloem by the action of Si transporters. Specifically, the activity of  
307 *Lsi3* and *Lsi6*, which are known to be expressed in the leaves, may be upregulated in damaged  
308 plants. To test this hypothesis, RT-qPCR was applied to determine the expression levels of the *Lsi3*  
309 and *Lsi6* genes. However, no consistent significant differences in *Lsi3* or *Lsi6* gene expression  
310 between damaged and undamaged plants were found, irrespective of the time point (4, 24, or 48 h)  
311 or the number of damage events (one or two; Supplementary Figure 4).

## 312 **6. Discussion**

### 313 **6.1. Is the induction of silicon-based defences a localised or systemic response, and** 314 **how is this affected by Si supply?**

315 Damage resulted in localised, but not systemic, induction of silicon defences. This response was  
316 observed across a range of landraces and at different levels of Si availability, although providing  
317 supplementary Si significantly increased the magnitude of the response. In contrast to the prevailing

318 hypothesis of Si being immediately deposited and thus rendered immobile, this study provides  
319 evidence that soluble Si can be moved from undamaged to damaged leaves to increase Si defences  
320 close to the site of wounding, where they are most needed. This appears to be the result of soluble  
321 Si being transported in the phloem sap to the stem, where it is then loaded into the xylem and  
322 transported to damaged leaves *via* the transpiration stream. However, this study found no evidence  
323 to suggest changes in Si transporter gene expression are involved.

324 In this study, artificial damage was used in place of herbivory to separate the effects of damage to  
325 tissue caused by the herbivore from the effects of molecules in the saliva and other excretions of the  
326 herbivore (Waterman *et al.*, 2019). Damage was found to significantly increase Si accumulation, a  
327 conclusion supported by previous studies which have found that mechanical damage is sufficient to  
328 significantly increase plant Si-based defences (McNaughton *et al.*, 1985; Kim *et al.*, 2014; Ryalls *et*  
329 *al.*, 2018), though it is also clear that actual herbivory can produce greater induction (Massey *et al.*,  
330 2007).

331 This study found significant variation in Si accumulation among wheat landraces. Eight out of ten  
332 landraces responded to damage by significantly increasing Si accumulation in damaged leaves, and  
333 this increase ranged from 34.2 % to 107.6 %. Similar genotypic variation has been found in other  
334 species: Bañuelos and Obeso (2000) reported significant genotypic variation in response to damage  
335 in the grass species *Agrostis tenuis*. Likewise, França *et al.* (2019) reported genotype-specific effects  
336 of Si in rice, such that Si reduced stem damage by stink bugs in only two out of three genotypes  
337 investigated. Using six genotypes for each of four grass species, Soininen *et al.* (2013) found  
338 significant genotypic variation in Si induction in response to damage in only two of the species  
339 examined.

340 This study found only a localised, and not systemic, Si response to damage, with the Si concentration  
341 increasing only in the damaged leaves of damaged plants. Few previous studies have made a  
342 distinction between the Si concentration of damaged and undamaged leaves of damaged plants. In  
343 agreement with the results of this study, insect herbivory significantly increased Si accumulation in  
344 attacked leaves of two genotypes of cucumber (Islam *et al.*, 2020) and damage resulted in localised  
345 induction of Si defences in three grass species (Hartley *et al.*, 2015). However, although damaged  
346 leaves had higher Si compared to undamaged plants for all genotypes, McLarnon *et al.* (2017) found  
347 significantly increased Si in damaged leaves compared to undamaged leaves in only one out of three  
348 tall fescue genotypes investigated.

## 349        **6.2. By what mechanism is localised induction of Si defences achieved?**

350 Overall, the Si concentration of damaged plants was not significantly different to that of undamaged  
351 plants. It was hypothesised that this was a result of the redirection of Si into damaged leaves, with  
352 the Si content of undamaged leaves of damaged plants being significantly lower than that of  
353 undamaged leaves of undamaged plants. To test for this hypothesis, plants were grown with Si and  
354 then moved to –Si medium prior to damage. Increased Si accumulation in damaged leaves was still  
355 observed in these plants, as well as decreased Si accumulation in undamaged leaves, suggesting that  
356 Si redirection had occurred.

357 It has widely been reported in the literature that once deposited as silica, Si cannot be remobilised  
358 (Yoshida *et al.*, 1962; Jones and Handreck, 1967; Samuels *et al.*, 1991; Liu *et al.*, 2019; Frick *et al.*,  
359 2020). However, it is likely that the Si was redirected into damaged leaves prior to deposition as  
360 silica, when it was still in a soluble form. Earlier studies have indicated that there may be relatively  
361 high levels of soluble Si present in the cytoplasm (Gartner *et al.*, 1984; Hodson and Evans, 1995),  
362 which could be used to increase the levels of Si in damaged leaves even after the removal of Si from  
363 the growth medium.

364 To examine whether there was sufficient soluble Si to explain the increased Si localised in damaged  
365 leaves, leaf soluble Si was measured. The soluble Si concentration of undamaged plants was taken as  
366 a baseline for the amount of soluble Si predicted to be present in plants before damage. The  
367 undamaged plants had an average total leaf Si concentration of 1.4 %, of which 15.7 % was soluble  
368 Si. Thus, undamaged plants comprised approximately 0.22 % soluble Si. However, this is likely to be  
369 an underestimate as the formation of silicomolybdate complexes in the molybdenum-based assay  
370 depends on the size of the silicates present. While monomeric and dimeric silicates react quickly,  
371 higher oligomers may not have fully reacted over the course of the 30-minute assay (Coradin *et al.*,  
372 2004). Despite this, it was calculated that there is sufficient soluble Si present in undamaged leaves  
373 to explain the increase in Si in damaged leaves. Thus, it was concluded that the increase in Si in  
374 damaged leaves can be explained by the movement of soluble Si from undamaged leaves into  
375 damaged leaves of damaged plants (Figure 5).

376 It was hypothesised that the increase in Si in damaged leaves would be the result of differences in Si  
377 transporter gene expression. In rice, Si transporters are used to preferentially allocate Si to the  
378 panicle and away from the flag leaf (Yamaji and Ma, 2009; Yamaji *et al.*, 2015) and it is possible that  
379 a similar mechanism results in the preferential allocation of Si to damaged leaves. To test for this,  
380 several time points, after one and two damage events, were used to investigate whether the

381 localised increase in Si in response to damage was the result of changes in Si transporter gene  
382 expression. However, no significant differences in Si transporter gene expression were found,  
383 despite the numerous timepoints tested meaning that it is unlikely any increase in gene expression  
384 was overlooked. Differences in Si transporter gene expression have been found in rice (Ye *et al.*,  
385 2013) but not in other species (McLarnon *et al.*, 2017). It remains possible that post-transcriptional  
386 processes affect the activity of Si transporters as gene expression is not always indicative of protein  
387 activity. Post-translational regulation has been found to be important for aquaporins (Verdoucq *et*  
388 *al.*, 2014). Alternatively, yet to be identified Si transporters may be involved, or the activity of  
389 transcriptional regulators may be involved, as has been found in rice (Wang *et al.*, 2017).

390 To the best of our knowledge, no previous studies have been carried out to directly measure Si in  
391 the phloem, and it is generally suggested that Si may not be phloem-mobile (Raven, 1983). However,  
392 xylem-to-phloem Si transfer has been suggested to occur during grain filling in rice, though this was  
393 not actually tested (Zhou *et al.*, 2021). Likewise, Yang *et al.* (2017) speculated that Si in the phloem  
394 may deter aphid feeding in rice but could not rule out the possibility that their results were due to Si  
395 affecting phloem sap composition. Both boron and arsenic, which share many chemical similarities  
396 to Si, have been observed in the phloem (Carey *et al.*, 2011; Zhao *et al.*, 2012; Umemura and  
397 Takenaka, 2014), although previous studies have not investigated whether this is also the case for Si.  
398 In the case of boron, it must bind to sugars before being transported in the phloem (Umemura and  
399 Takenaka, 2014) and it is possible that a similar mechanism is involved in Si redirection, with *in vitro*  
400 studies supporting the idea that various forms of Si can bind to biologically relevant polymers  
401 (Annenkov *et al.*, 2017). Our study is the first to report significant levels of soluble Si present in the  
402 phloem exudate.

403 The data presented here suggest a model of soluble Si being redirected from undamaged to  
404 damaged leaves *via* the phloem. While it is noted that contamination is often an issue associated  
405 with EDTA-facilitated phloem exudate collection, alternative methods including aphid stylectomy  
406 would not provide sufficient yield for soluble Si analysis (Gaupels *et al.*, 2008). Another issue with  
407 EDTA-facilitated phloem exudate collection is that the amount of exudate released is unknown,  
408 meaning that quantitative comparisons between samples are not possible. Nevertheless, the  
409 presence of Si in the phloem exudate of +/- Si plants which have been grown without exogenous Si  
410 for 24 hr strongly supports the hypothesis that Si is being transported in the phloem as the presence  
411 of soluble Si in such +/- Si plants cannot be explained by *de novo* uptake from the hydroponic  
412 medium. Furthermore, as localised Si induction was observed even nine days after the removal of

413 external Si supply, it is unlikely that there is still enough Si present in the xylem to explain the  
414 observed induction of Si.

415 Due to the low levels of plant-available Si found in many soils, it has recently been suggested that  
416 there are significant costs associated with high Si accumulation and that these are much higher than  
417 previously recognised (de Tombeur *et al.*, 2023). Negative correlations between Si accumulation and  
418 biomass have been reported in several grass species, and in a number of studies (Simpson *et al.*,  
419 2017; Johnson and Hartley, 2018; de Tombeur *et al.*, 2021), suggesting that there is a trade-off  
420 between Si uptake and growth. This could reflect the fact that Si deposition is an active process  
421 involving the use of active efflux transporters (Ma *et al.*, 2007a; Ma and Yamaji, 2015), so there may  
422 be an energetic cost associated with high Si uptake (Simpson *et al.*, 2017). Thus, redirection of  
423 soluble Si from undamaged leaves to sites of damage might be a more energetically favourable way  
424 of increasing defences against herbivores than the more costly mechanism of increasing Si uptake.

### 425 **6.3. Conclusions**

426 By separately measuring Si accumulation in damaged and undamaged leaves, this study has  
427 demonstrated that damage results in only a localised, and not systemic, induction of Si defences.  
428 This localised induction was observed in multiple landraces, although it varied between them, and  
429 the response was stronger when plants were grown at high levels of Si availability. The evidence  
430 presented here suggests that this localised induction is the result of the redirection of soluble Si  
431 from undamaged to damaged leaves *via* the phloem. To the best of our knowledge, this is the first  
432 demonstration of soluble Si being redirected towards the sites of wounding within damaged plants  
433 to increase their defences, a mechanism which has implications for the cost-effectiveness of these  
434 defences, as well as for wider understanding of the fitness benefits of Si accumulation.

## 435 **7. Supplementary data**

436 The following supplementary data are available at JXB online.

437 Table S1. List of landraces used for this study.

438 Table S2. List of primers used for this study.

439 Table S3. ANOVA results for the effect of damage and Si supply on Si accumulation.

440 Fig. S1. Average leaf Si concentration for ten wheat landraces.

441 Fig. S2. Root Si concentration for four landraces grown hydroponically with and without Si  
442 supplementation and subject to damage.

443 Fig. S3. Stem and root Si concentrations for plants grown without Si supplementation after damage.



444 Fig. S4. Si transporter gene expression after damage.

## 445 **8. Acknowledgements**

446 We would like to thank the horticulture team at York for assistance with plant growth. For the  
447 purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to  
448 any Author Accepted Manuscript version arising.

## 449 **9. Author contributions**

450 Conceptualization, S.J.T., S.E.H. and F.J.M.M.; methodology, S.J.T., S.E.H. and F.J.M.M.; formal  
451 analysis, S.J.T.; investigation, S.J.T.; data curation, S.J.T.; writing—original draft preparation, S.J.T.;  
452 writing—review and editing, S.E.H. and F.J.M.M.; funding acquisition, S.E.H. and F.J.M.M. All authors  
453 have read and agreed to the published version of the manuscript.

## 454 **10. Conflict of interest**

455 The authors declare no competing interests.

## 456 **11. Funding statement**

457 This work was supported by the Biotechnology and Biological Sciences Research Council (Award ref  
458 1949569). The University of Sheffield provided additional financial support.

## 459 **12. Data availability**

460 All data generated and analysed in this study are available upon request.

### 13. References

- Annenkov VV, Danilovtseva EN, Pal'shin VA, Verkhovzina ON, Zelinskiy SN, Krishnan UM.** 2017. Silicic acid condensation under the influence of water-soluble polymers: from biology to new materials. *RSC Advances* **7**, 20995–21027.
- Bañuelos MJ, Obeso JR.** 2000. Effect of grazing history, experimental defoliation, and genotype on patterns of silicification in *Agrostis tenuis* Sibth. *Ecoscience* **7**, 45–50.
- Carey AM, Norton GJ, Deacon C, et al.** 2011. Phloem transport of arsenic species from flag leaf to grain during grain filling. *New Phytologist* **192**, 87–98.
- Coradin T, Eglin D, Livage J.** 2004. The silicomolybdic acid spectrophotometric method and its application to silicate/biopolymer interaction studies. *Spectroscopy* **18**, 567–576.
- Coskun D, Deshmukh R, Sonah H, Menzies JG, Reynolds O, Ma JF, Kronzucker HJ, Bélanger RR.** 2019. The controversies of silicon's role in plant biology. *New Phytologist* **221**, 67–85.
- França LL, Dierings CA, Almeida AC de S, Araújo M da S, Heinrichs EA, Silva AR da, Barrigossi JAF, Jesus FG de.** 2019. Resistance in rice to *Tibraca limbativentris* (Hemiptera: Pentatomidae) influenced by plant silicon content. *Florida Entomologist* **101**, 587.
- Frick DA, Remus R, Sommer M, Augustin J, Kaczorek D, Von Blanckenburg F.** 2020. Silicon uptake and isotope fractionation dynamics by crop species. *Biogeosciences* **17**, 6475–6490.
- Gartner S, Roinel N, Paris-Pireyre N.** 1984. Electron probe analysis of sap exudates of two varieties of wheat (*Triticum Aestivum* L.). *Journal de Physique (Paris), Colloque* **45**, 515–518.
- Gaupels F, Knauer T, van Bel AJE.** 2008. A combinatory approach for analysis of protein sets in barley sieve-tube samples using EDTA-facilitated exudation and aphid stylectomy. *Journal of Plant Physiology* **165**, 95–103.
- Hall CR, Dagg V, Waterman JM, Johnson SN.** 2020. Silicon alters leaf surface morphology and suppresses insect herbivory in a model grass species. *Plants* **9**, 643.
- Han Y, Lei W, Wen L, Hou M.** 2015. Silicon-mediated resistance in a susceptible rice variety to the

rice leaf folder, *Cnaphalocrocis medinalis* Guenée (Lepidoptera: Pyralidae). PLoS ONE **10**, 1–13.

**Hartley SE, DeGabriel JL.** 2016. The ecology of herbivore-induced silicon defences in grasses. *Functional Ecology* **30**, 1311–1322.

**Hartley SE, Fitt RN, McLarnon EL, Wade RN.** 2015. Defending the leaf surface: intra- and inter-specific differences in silicon deposition in grasses in response to damage and silicon supply. *Frontiers in Plant Science* **6**, 1–8.

**Hodson MJ, Evans DE.** 1995. Aluminium/silicon interactions in higher plants. *Journal of Experimental Botany* **46**, 161–171.

**Hope RM.** 2013. Rmisc: Ryan Miscellaneous. R package v.1.5. URL <https://CRAN.R-project.org/package=Rmisc>

**Howe GA, Jander G.** 2007. Plant immunity to insect herbivores. *Annual Review of Plant Biology* **59**, 41–66.

**Huang S, Yamaji N, Sakurai G, Mitani-Ueno N, Konishi N, Ma JF.** 2022. A pericycle-localized silicon transporter for efficient xylem loading in rice. *New Phytologist* **234**, 197–208.

**Islam T, Moore BD, Johnson SN.** 2020. Novel evidence for systemic induction of silicon defences in cucumber following attack by a global insect herbivore. *Ecological Entomology* **45**, 1373–1381.

**Johnson J.** 2014. Accurate measurements of low Z elements in sediments and archaeological ceramics using portable X-ray fluorescence (pXRF). *Journal of Archaeological Method and Theory* **21**, 563–588.

**Johnson SN, Hartley SE.** 2018. Elevated carbon dioxide and warming impact silicon and phenolic-based defences differently in native and exotic grasses. *Global Change Biology* **24**, 3886–3896.

**Johnson SN, Vandeger RK, Borevitz JO, Hartley SE, Tissue DT, Hall CR.** 2023. Climatic drivers of silicon accumulation in a model grass operate in low- but not high-silicon soils. *Plants* **12**, 995.

**Jones LHP, Handreck KA.** 1967. Silica in soils, plants, and animals. *Advances in agronomy*. **19**, 107–149.

**Kim YH, Khan AL, Waqas M, Jeong HJ, Kim DH, Shin JS, Kim JG, Yeon MH, Lee IJ.** 2014. Regulation of jasmonic acid biosynthesis by silicon application during physical injury to *Oryza sativa* L. *Journal of Plant Research* **127**, 525–532.

**Lenth R V.** 2021. emmeans: Estimated Marginal Means, aka Least-Squares Means. v. 1.5.4. URL <https://CRAN.R-project.org/package=emmeans>

**Liu WS, Zheng HX, Guo MN, Liu C, Huot H, Morel JL, van der Ent A, Tang YT, Qiu RL.** 2019. Co-deposition of silicon with rare earth elements (REEs) and aluminium in the fern *Dicranopteris linearis* from China. *Plant and Soil* **437**, 427–437.

**Ma JF, Yamaji N.** 2015. A cooperative system of silicon transport in plants. *Trends in Plant Science* **20**, 435–442.

**Ma JF, Yamaji N, Mitani N, Tamai K, Konishi S, Fujiwara T, Katsuhara M, Yano M.** 2007a. An efflux transporter of silicon in rice. *Nature* **448**, 209–212.

**Ma JF, Yamaji N, Tamai K, Mitani N.** 2007b. Genotypic difference in silicon uptake and expression of silicon transporter genes in rice. *Plant Physiology* **145**, 919–924.

**Massey FP, Hartley SE.** 2006. Experimental demonstration of the antiherbivore effects of silica in grasses: impacts on foliage digestibility and vole growth rates. *Proceedings of the Royal Society B: Biological Sciences* **273**, 2299–2304.

**Massey FP, Hartley SE.** 2009. Physical defences wear you down: progressive and irreversible impacts of silica on insect herbivores. *Journal of Animal Ecology* **78**, 281–291.

**Massey FP, Massey K, Roland Ennos A, Hartley SE.** 2009. Impacts of silica-based defences in grasses on the feeding preferences of sheep. *Basic and Applied Ecology* **10**, 622–630.

**Massey FP, Roland Ennos A, Hartley SE.** 2007. Herbivore specific induction of silica-based plant defences. *Oecologia* **152**, 677–683.

**Massey FP, Smith MJ, Lambin X, Hartley SE.** 2008. Are silica defences in grasses driving vole population cycles? *Biology Letters* **4**, 419–422.

**McLarnon E, McQueen-Mason S, Lenk I, Hartley SE.** 2017. Evidence for active uptake and deposition of Si-based defenses in tall fescue. *Frontiers in Plant Science* **8**, 1–11.

**McNaughton SJ, Tarrant JL, McNaughton MM, Davis RD.** 1985. Silica as a defense against herbivory and a growth promotor in African grasses. *Ecology* **66**, 528–535.

**Muller PY, Janovjak H, Miserez AR, Dobbie Z.** 2002. Processing of gene expression data generated by quantitative real-time RT-PCR. *BioTechniques* **32**, 1372–4, 1376, 1378–9.

**R Development Core Team.** 2022. R: A language and environment for statistical computing, v.4.2.0. Vienna, Austria: R Foundation for Statistical Computing. URL <https://www.R-project.org/>.

**Raven JA.** 1983. The transport and function of silicon in plants. *Biological Reviews* **58**, 179–207.

**Reidinger S, Ramsey MH, Hartley SE.** 2012. Rapid and accurate analyses of silicon and phosphorus in plants using a portable X-ray fluorescence spectrometer. *New Phytologist* **195**, 699–706.

**Reynolds JJH, Lambin X, Massey FP, Reidinger S, Sherratt JA, Smith MJ, White A, Hartley SE.** 2012. Delayed induced silica defences in grasses and their potential for destabilising herbivore population dynamics. *Oecologia* **170**, 445–456.

**Ryalls JMW, Moore BD, Johnson SN.** 2018. Silicon uptake by a pasture grass experiencing simulated grazing is greatest under elevated precipitation. *BMC Ecology* **18**, 1–8.

**Samuels AL, Glass ADM, Ehert DL, Menzies JG.** 1991. Mobility and deposition of silicon in cucumber plants. *Plant, Cell & Environment* **14**, 485–492.

**Sauer D, Saccone L, Conley DJ, Herrmann L, Sommer M.** 2006. Review of methodologies for extracting plant-available and amorphous Si from soils and aquatic sediments. *Biogeochemistry* **80**, 89–108.

**Simon P.** 2003. Q-Gene: Processing quantitative real-time RT-PCR data. *Bioinformatics* **19**, 1439–1440.

**Simpson KJ, Wade RN, Rees M, Osborne CP, Hartley SE.** 2017. Still armed after domestication? Impacts of domestication and agronomic selection on silicon defences in cereals. *Functional Ecology*

**31**, 2108–2117.

**Singh A, Kumar A, Hartley S, Singh IK.** 2020. Silicon: Its ameliorative effect on plant defense against herbivory. *Journal of Experimental Botany* **71**, 6730–6743.

**Soininen EM, Bråthen KA, Jusdado JGH, Reidinger S, Hartley SE.** 2013. More than herbivory: Levels of silica-based defences in grasses vary with plant species, genotype and location. *Oikos* **122**, 30–41.

**Talukdar P, Hartley SE, Travis AJ, Price AH, Norton GJ.** 2019. Genotypic differences in shoot silicon concentration and the impact on grain arsenic concentration in rice. *Journal of Plant Nutrition and Soil Science* **182**, 265–276.

**Thorne SJ, Hartley SE, Maathuis FJM.** 2021. The effect of silicon on osmotic and drought stress tolerance in wheat landraces. *Plants* **10**, 814.

**de Tombeur F, Cooke J, Collard L, Cisse D, Saba F, Lefebvre D, Burgeon V, Nacro HB, Cornelis J-T.** 2021. Biochar affects silicification patterns and physical traits of rice leaves cultivated in a desilicated soil (Ferric Lixisol). *Plant and Soil* **460**, 375–390.

**de Tombeur F, Lemoine T, Violle C, Fréville H, Thorne SJ, Hartley SE, Lambers H, Fort F.** 2022. Nitrogen availability and plant–plant interactions drive leaf silicon concentration in wheat genotypes. *Functional Ecology* **36**, 2687–2888.

**de Tombeur F, Raven JA, Toussaint A, et al.** 2023. Why do plants silicify? *Trends in Ecology & Evolution* **38**, 275–288.


**Umemura M, Takenaka C.** 2014. Retranslocation and localization of nutrient elements in various organs of moso bamboo (*Phyllostachys pubescens*). *Science of the Total Environment* **493**, 845–853.

**Verdoucq L, Rodrigues O, Martinière A, Luu DT, Maurel C.** 2014. Plant aquaporins on the move: Reversible phosphorylation, lateral motion and cycling. *Current Opinion in Plant Biology* **22**, 101–107.

**Wang FZ, Chen MX, Yu LJ, et al.** 2017. OsARM1, an R2R3 MYB Transcription factor, is involved in regulation of the response to arsenic stress in rice. *Frontiers in Plant Science* **8**, 1–16.


- Waterman JM, Cazzonelli CI, Hartley SE, Johnson SN.** 2019. Simulated herbivory: The key to disentangling plant defence responses. *Trends in Ecology & Evolution* **34**, 447–458.
- Waterman JM, Cibils-Stewart X, Cazzonelli CI, Hartley SE, Johnson SN.** 2021. Short-term exposure to silicon rapidly enhances plant resistance to herbivory. *Ecology* **102**, 1–8.
- Wickham H.** 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. URL <https://ggplot2.tidyverse.org>
- Xu Q, Ren Y, Liesche J.** 2019. Studying phloem loading with EDTA-facilitated phloem exudate collection and analysis. In: Liesche J. (eds) *Phloem: Methods in Molecular Biology, vol 2014*. Humana, New York, 125–133.
- Yamaji N, Ma JF.** 2009. A transporter at the node responsible for intervascular transfer of silicon in rice. *The Plant Cell* **21**, 2878–2883.
- Yamaji N, Mitatni N, Ma JF.** 2008. A transporter regulating silicon distribution in rice shoots. *The Plant Cell* **20**, 1381–1389.
- Yamaji N, Sakurai G, Mitani-Ueno N, Ma JF.** 2015. Orchestration of three transporters and distinct vascular structures in node for intervascular transfer of silicon in rice. *Proceedings of the National Academy of Sciences* **112**, 11401–11406.
- Yang L, Han Y, Li P, Wen L, Hou M.** 2017. Silicon amendment to rice plants impairs sucking behaviors and population growth in the phloem feeder *Nilaparvata lugens* (Hemiptera: Delphacidae). *Scientific Reports* **7**, 1101.
- Ye M, Song Y, Long J, et al.** 2013. Priming of jasmonate-mediated antiherbivore defense responses in rice by silicon. *Proceedings of the National Academy of Sciences* **110**, E3631–E3639.
- Yoshida S, Ohnishi Y, Kitagishi K.** 1962. Chemical forms, mobility and deposition of silicon in rice plant. *Soil Science and Plant Nutrition* **8**, 15–21.
- Zhao FJ, Stroud JL, Khan AA, McGrath SP.** 2012. Arsenic translocation in rice investigated using radioactive <sup>73</sup>As tracer. *Plant and Soil* **350**, 413–420.

## 14. Figures



**Figure 1: Variation in Si accumulation following damage in ten wheat landraces.** Leaf Si concentration for damaged and undamaged leaves of damaged plants, and undamaged plants. L1-L5 are low Si-accumulating landraces; H1-H5 are high Si-accumulating landraces. Mean values  $\pm$  SE are shown.  $N = 3$ . ANOVA damaged leaves of damaged plants vs undamaged plants: Landrace:  $F_{9,40} = 12.0$ ,  $P < 0.001$ , Damage treatment:  $F_{1,40} = 67.9$ ,  $P < 0.001$ . ANOVA undamaged leaves of damaged plants vs undamaged plants: Landrace  $F_{9,40} = 19.5$ ,  $P < 0.001$ , Damage treatment:  $F_{1,40} = 9.2$ ,  $P = 0.004$ . Statistically significant differences between damaged and undamaged leaves of damaged plants, determined by paired  $t$ -tests, are indicated: \*\*\*  $P < 0.001$ , \*\*  $P < 0.01$ , and \*  $P < 0.05$ .







**Figure 2: Effect of Si supply and damage on Si accumulation.** a) Leaf Si concentration of -Si plants. b) Leaf Si concentration of +Si plants. Note the different scales on the y-axis. Mean values  $\pm$  standard error (SE) are shown.  $N = 3$ . L4 and L5 are low Si-accumulating landrace. H1 and H3 are high Si-accumulating landraces. L indicates a low Si-accumulating landrace. ANOVA for Landrace, Si availability, Landrace x Si availability, and Damage treatment x Si availability at  $P < 0.001$  for both damaged and undamaged leaves of damaged plants vs undamaged plants, see

Supplementary Table 3 for full results. Statistically significant differences between damaged and undamaged leaves of damaged plants, determined by paired  $t$ -tests, are indicated: \*\*\*  $P < 0.001$ , \*\*  $P < 0.01$ , and \*  $P < 0.05$ .

a) Si removed prior to damage (+/- Si)




b) Continuous Si supply (+/+ Si)




**Figure 3: Effect of reducing Si availability on Si accumulation after damage.** Leaf Si concentration for damaged and undamaged leaves of damaged plants, and undamaged plants. a) Plants moved to medium without Si supplementation when damage was started (+/-Si plants). b) Plants grown with continuous Si supplementation (+/+ Si plants). The low Si-accumulating landrace, L1, was used. Mean values  $\pm$  SE are shown. N = 3. ANOVA for Si availability, Damage treatment, and Number of damage events x Si availability at  $P < 0.01$  for both damaged and undamaged leaves of damaged plants vs undamaged plants, see

Supplementary Table 3 for full results. Statistically significant differences between damaged and undamaged leaves of damaged plants, determined by paired *t*-tests, are indicated: \*\*\*  $P < 0.001$ , \*\*  $P < 0.01$ , and \*  $P < 0.05$ .



**Figure 4: Total leaf Si content was unaffected by damage.** Allocation of total Si accumulated to different plant tissues in damaged and undamaged plants after successive damage events. a) +/- Si plants sampled after 1, 2, and 4 damage events. b) ++ Si plants after 1, 2, and 4 damage. Mean values  $\pm$  SE are shown. N = 3. Note different scales of y-axis. Significant ANOVA effects: Number of damage events:  $F_{2,23} = 36.0$ ,  $P < 0.001$ , Si availability  $F_{1,23} = 13.8$ ,  $P = 0.001$ , Number of damage events x Si availability  $F_{2,23} = 8.1$ ,  $P = 0.002$ .



**Figure 5: The localised increase in Si in damaged leaves can be explained by the redirection of soluble Si from undamaged leaves of damaged plants.** An undamaged plant weighing 1 g DW contains a total of 14 mg Si g<sup>-1</sup> leaf on average, of which 2.2 mg g<sup>-1</sup> is soluble. After damage, soluble Si is moved from undamaged leaves to damaged leaves. This movement of soluble Si increases the total Si concentration in damaged leaves to 17 mg Si g<sup>-1</sup>, of which 4 mg g<sup>-1</sup> is soluble. Provided the weight of undamaged leaves is at least 4.5 times more than the weight of damaged leaves, the increase in Si in damaged leaves can be explained by the redirection of soluble Si from undamaged to damaged leaves of damaged plants. This was the case for all the experiments presented here.

Figure created using BioRender (<https://www.biorender.com>).

## Supplementary Information

**Supplementary Table 1: List of landraces used in this study.** H1-H5 are high Si-accumulating landraces while L1-L5 are low Si-accumulating landraces, as characterised in Thorne *et al.* (2021).

Landrace	Collection	Plant ID	Plant Name	Origin
H1	CIMMYT	CWI 2166	K7155.41	Kenya
H2	Watkins	1190195	Gahu (Nepali) or Kyo (Sikkimese)	India
H3	CIMMYT	CWI 3909	OUBAARD	South Africa
H4	Watkins	1190777	Finland 3	Finland
H5	Prague	01C0201531	Orchon	Mongolia
L1	Watkins	1190521	Dandi	India
L2	Watkins	1190568	China 19	China
L3	Watkins	1190605	Karabash	Greece
L4	Watkins	1190662	Samanta 1252	Romania
L5	Watkins	1190751	Armavir	USSR

**Supplementary Table 2: List of primers used for RT-qPCR.**

Target Gene	Forward Primer	Reverse Primer	Product size	Ensembl gene name (all homologues)
<i>Actin</i>	AATGGTCAAGGCTGGTTTCG	ATCACCGACATAGGCATCCTTC	124	TraesCS1A02G020500, TraesCS1B02G024500, TraesCS1D02G020000
<i>Tef1</i>	TTGGTGGCATTGGAAGTGTG	TTGACCTCAGTTGTCAGACCAG	103	TraesCS5B02G418200 TraesCS5D02G423400
<i>Lsi3</i>	TGTTCAAGTACCTCGGCAAC	TTGAGGATGAACTCGGTGAGG	144	TraesCS4A02G412500, TraesCS4B02G312600, TraesCS4D02G310100
<i>Lsi6</i>	TACTCGAACGAGATCCACGAC	TCTCCGATATCACCTTCTTGCC	132	TraesCS7A02G187800, TraesCS7B02G092900, TraesCS7D02G188800


**Supplementary Table 3: ANOVA results for Figures 2 and 3.** Statistically significant results are highlighted in bold. N = 3 for each factor combination.

- a) The effect of landrace, damage, and Si supply on Si concentration in damaged and undamaged leaves of damaged plants compared to undamaged plants.


	Leaf Si (%)						Root Si (%)		
	Undamaged leaves			Damaged leaves			df	F	P
	df	F	P	df	F	P			
Landrace	<b>3</b>	<b>22.39</b>	<b>&lt; 0.001</b>	<b>3</b>	<b>24.81</b>	<b>&lt; 0.001</b>	3	1.71	0.179
Damage	1	0.09	0.764	1	<b>215.99</b>	<b>&lt; 0.001</b>	1	0.79	0.378
Si	<b>1</b>	<b>6188.77</b>	<b>&lt; 0.001</b>	<b>1</b>	<b>9076.47</b>	<b>&lt; 0.001</b>	<b>1</b>	<b>88.54</b>	<b>&lt; 0.001</b>
Landrace x Damage	3	0.20	0.897	3	0.54	0.66	3	0.71	0.550
Landrace x Si	<b>3</b>	<b>23.73</b>	<b>&lt; 0.001</b>	<b>3</b>	<b>27.41</b>	<b>&lt; 0.001</b>	3	1.19	0.326
Damage x Si	<b>1</b>	<b>12.32</b>	<b>&lt; 0.001</b>	<b>1</b>	<b>23.10</b>	<b>&lt; 0.001</b>	1	0.16	0.691
Landrace x Damage x Si	3	0.16	0.923	3	1.39	0.26	3	0.86	0.466

- b) The effect of damage, Si supply, and number of damage events on total plant Si content and leaf Si concentration in damaged and undamaged leaves of damaged plants compared to undamaged plants.


	Leaf Si (%)						Plant Si content (mg)		
	Undamaged leaves			Damaged leaves			df	F	P
	df	F	P	df	F	P			
Number of damage events	<b>2</b>	<b>10.30</b>	<b>&lt; 0.001</b>	2	1.75	0.196	<b>2</b>	<b>36.01</b>	<b>&lt; 0.001</b>
Damage	<b>1</b>	<b>42.81</b>	<b>&lt; 0.001</b>	<b>1</b>	<b>84.23</b>	<b>&lt; 0.001</b>	1	0.08	0.778
Si	<b>1</b>	<b>95.99</b>	<b>&lt; 0.001</b>	<b>1</b>	<b>108.24</b>	<b>&lt; 0.001</b>	<b>1</b>	<b>13.80</b>	<b>0.001</b>
Number x Damage	2	2.62	0.094	2	1.34	0.281	2	0.37	0.698
Number x Si	<b>2</b>	<b>6.79</b>	<b>0.005</b>	<b>2</b>	<b>21.64</b>	<b>&lt; 0.001</b>	<b>2</b>	<b>8.11</b>	<b>0.002</b>
Damage x Si	1	0.15	0.701	1	0.07	0.800	1	0.16	0.694
Number x Damage x Si	2	1.21	0.316	2	1.87	0.176	2	0.35	0.709



**Supplementary Figure 1: Variation in leaf Si accumulation between damaged and undamaged plants for ten wheat landraces.** The average leaf Si concentration for damaged and undamaged leaves of damaged plants was calculated. L1-L5 are low Si-accumulating landraces; H1-H5 are high Si-accumulating landraces. Mean values  $\pm$  SE are shown. N = 3. Statistically significant ANOVA effect: Landrace:  $F_{9,40} = 18.7$ ,  $P < 0.001$ .




**Supplementary Figure 2: Effect of damage and Si supply on root Si accumulation.** (A) Root Si of -Si plants. (B) Root Si of +Si plants. L4 and L5 are low Si-accumulating landraces; H1 and H3 are high Si-accumulating landraces. Mean values  $\pm$  standard error (SE) are shown. N = 3. Statistically significant ANOVA effect: Si:  $F_{1,47} = 88.5$ ,  $P < 0.001$ .



**Supplementary Figure 3: Effect of reducing Si availability on Si accumulation after damage. (A)**

Stem Si concentration for plants moved to medium without Si supplementation when damage was started (+/- Si plants). (B) Stem Si concentration for plants grown continuously with Si supplementation (+/+ Si plants). (C) Root Si concentration for +/- Si plants. (D) Root Si concentration for +/- Si plants. Mean values  $\pm$  SE are shown. N = 3. The low Si accumulating landrace, L1, was used. Significant ANOVA effects for stem Si: Si availability:  $F_{1,24} = 138.5$ ,  $P < 0.001$ ; Number of damage events:  $F_{2,24} = 12.2$ ,  $P < 0.001$ ; Number of damage events x Si availability:  $F_{2,24} = 20.5$ ,  $P < 0.001$ ; for root Si: Si availability:  $F_{1,23} = 12.7$ ,  $P = 0.002$ ; Number of damage events:  $F_{2,23} = 5.2$ ,  $P = 0.013$ .





**Supplementary Figure 4: Si transporter gene expression at different time points.** (A) *Lsi3* expression. (B) *Lsi6* expression. The low Si-accumulating landrace, L1, was used. Mean values  $\pm$  SE are shown. N = 2-5. Dam: number of damage events. No significant differences between damaged and undamaged plants, and between damaged and undamaged leaves of damaged plants, were found.