



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/200608/>

Version: Published Version

---

**Article:**

de Paula, M.A.A., dos Santos Leite, L.C. and Crispino, L.C.B. (2023) Comment on: "Absorption and scattering of massless scalar wave from Regular Black Holes". *General Relativity and Gravitation*, 55 (6). 73. ISSN: 0001-7701

<https://doi.org/10.1007/s10714-023-03122-7>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



## Comment on: “Absorption and scattering of massless scalar wave from Regular Black Holes”

Marco Aurélio Abreu de Paula<sup>1,3</sup> · Luiz Carlos dos Santos Leite<sup>2</sup> ·  
Luís Carlos Bassalo Crispino<sup>1,4</sup>

Received: 10 April 2023 / Accepted: 30 May 2023  
© The Author(s) 2023

Absorption and scattering properties of matter fields in the background of black hole (BH) spacetimes have been investigated in several standard scenarios since the 1960s (see, e.g., Refs. [1–6] and references therein). In the last decade, some attempts to improve the comprehension of these properties in the background of regular BH (RBH) geometries, considering mainly test scalar fields, have been performed (see, e.g., Refs. [7–12]). The authors of the recent work “Absorption and scattering of massless scalar wave from Regular Black Holes” [12] addressed the absorption and scattering cross sections of massless test scalar fields in the background of several well-known RBH solutions. They considered the partial-waves approach and computed the absorption and scattering cross sections through a numerical method for arbitrary values of the frequency  $\omega$  and scattering angle  $\theta$  of the scalar wave.

Nevertheless, the absorption and scattering results obtained by the authors of Ref. [12] using the Numerov numerical method [13] combined with the Pöschl-Teller

---

The authors contributed equally to this work.

✉ Marco Aurélio Abreu de Paula  
marco.paula@icen.ufpa.br  
Luiz Carlos dos Santos Leite  
luiz.leite@ifpa.edu.br  
Luís Carlos Bassalo Crispino  
crispino@ufpa.br

<sup>1</sup> Programa de Pós-Graduação em Física, Universidade Federal do Pará, Belém, Pará 66075-110, Brazil

<sup>2</sup> Campus Altamira, Instituto Federal do Pará, Altamira, Pará 68377-630, Brazil

<sup>3</sup> Consortium for Fundamental Physics, School of Mathematics and Statistics, University of Sheffield, Hicks Building, Hounsfield Road, Sheffield S3 7RH, UK

<sup>4</sup> Departamento de Matemática da Universidade de Aveiro and Centre for Research and Development in Mathematics and Applications (CIDMA), Campus de Santiago, 3810-183 Aveiro, Portugal

potential approximation [14] present some inconsistencies. In this comment, we highlight these inaccuracies, by taking into account some previous results available in the literature [7–10].

Concerning the absorption cross section (ACS), the authors of Ref. [12] obtained that the total ACS decreases as we consider higher BH charge values, and it oscillates around the corresponding geometric cross section, in agreement with some results presented in the literature. However, in the low-frequency regime, the total ACS does not tend to the BH area, as expected from Ref. [15].

Now let us turn our attention to the scattering case. The authors of Ref. [12] obtained that the differential scattering cross section (SCS) of massless test scalar fields in the background of the considered RBH geometries is finite for small scattering angles. This result is incorrect since for scattering potentials with infinite range, the differential SCS diverges in the forward direction [16]. From analytical approximations, for small scattering angles, we also know that the BH charge does not contribute to the dominant term of the differential SCS, in this regime (see, e.g., Refs. [17–20] for Ayón-Beato-García, Bardeen, Bronnikov, and Dymnikova RBHs). Therefore, as  $\theta \rightarrow 0$ , we do not expect to see considerable differences in the differential SCSs as we vary the BH charge. Notwithstanding, the results obtained by the authors show considerable differences in this regime as we increase the BH charge values.

We also emphasize that light rays in the background of RBH geometries associated with nonlinear electrodynamics models follow null geodesics of an effective geometry [21], which is different from the geometry of the spacetime itself. Consequently, the classical results for the absorption and scattering cross section of massless test scalar fields in the background of nonlinear electrodynamics-based RBHs correspond to the trajectory of massless particles with nature other than electromagnetic. Therefore, the geodesic analysis exhibited by the authors of Ref. [12] is not valid for photons, in contrast with what they have stated.

In summary, regarding the absorption and scattering results obtained by M.-Y. Wan and C. Wu using the Numerov numerical method and the Pöschl-Teller potential approximation [12], we conclude that the approach, as applied by them, leads to erroneous results for small values of the frequency and of the scattering angle. Furthermore, the oscillatory pattern of the SCS is unclear. The correct results concerning the Ayón-Beato-García and Bardeen RBHs can be found in Refs. [7–10].

We are grateful to Fundação Amazônia de Amparo a Estudos e Pesquisas (FAPESPA), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)—Finance Code 001, from Brazil, for partial financial support. MP and LC thank the University of Sheffield, in England, and University of Aveiro, in Portugal, respectively, for the kind hospitality during the completion of this work. LL would like to acknowledge IFPA—Campus Altamira for the support. This work has further been supported by the European Union’s Horizon 2020 research and innovation (RISE)

programme H2020-MSCA-RISE-2017 Grant No. FunFiCO-777740 and by the European Horizon Europe staff exchange (SE) programme HORIZON-MSCA-2021-SE-01 Grant No. NewFunFiCO-101086251.

## Declarations

**Conflict of interest** The author declares to have no conflicts of interest.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Futterman, J.A., Handler, F.A., Matzner, R.A.: Scattering from Black Holes. Cambridge University Press, Cambridge (1988)
2. Dolan, S., Doran, C., Lasenby, A.: Fermion scattering by a Schwarzschild black hole. *Phys. Rev. D* **74**, 064005 (2006)
3. Crispino, L.C.B., Dolan, S.R., Oliveira, E.S.: Scattering of massless scalar waves by Reissner–Nordström black holes. *Phys. Rev. D* **79**, 0640022 (2009)
4. Benone, C.L., Crispino, L.C.B.: Superradiance in static black hole spacetimes. *Phys. Rev. D* **93**, 024028 (2016)
5. Leite, L.C.S., Benone, C.L., Crispino, L.C.B.: Scalar absorption by charged rotating black holes. *Phys. Rev. D* **96**, 044043 (2017)
6. Leite, L.C.S., Benone, C.L., Crispino, L.C.B.: On-axis scattering of scalar fields by charged rotating black holes. *Phys. Lett. B* **795**, 6 (2019)
7. Macedo, C.F.B., Crispino, L.C.B.: Absorption of planar massless scalar waves by Bardeen regular black holes. *Phys. Rev. D* **90**, 064001 (2014)
8. Macedo, C.F.B., de Oliveira, E.S., Crispino, L.C.B.: Scattering by regular black holes: planar massless scalar waves impinging upon a Bardeen black hole. *Phys. Rev. D* **92**, 024012 (2015)
9. Paula, M.A.A., Leite, L.C.S., Crispino, L.C.B.: Electrically charged black holes in linear and non-linear electrodynamics: geodesic analysis and scalar absorption. *Phys. Rev. D* **102**, 104033 (2020)
10. de Paula, M.A.A., Leite, L.C.S., Crispino, L.C.B.: Scattering properties of charged black holes in nonlinear and Maxwell's electrodynamics. *Eur. Phys. J. Plus* **137**, 785 (2022)
11. Sanchez, P.A., Bretón, N., Bergliaffa, S.E.P.: Scattering and absorption of massless scalar waves by Born–Infeld black holes. *Ann. Phys. (N.Y.)* **393**, 107 (2018)
12. Wan, M.-Y., Wu, C.: Absorption and scattering of massless scalar wave from Regular Black Holes. *Gen. Relativ. Gravit.* **54**, 148 (2022)
13. Hairer, E., Nørsett, S.P., Wanner, G.: Solving Ordinary Differential Equations I: Nonstiff Problems. Springer, Berlin (1993)
14. Pöschl, G., Teller, E.: Bemerkungen zur quantenmechanik des anharmonischen oszillators. *Z. Phys.* **83**, 143–151 (1933)
15. Higuchi, A.: Low-frequency Scalar absorption cross sections for stationary black holes. *Class. Quantum Gravit.* **18**, L139 (2001); **19**, 599(A) (2002)
16. Newton, R.G.: Scattering Theory of Waves and Particles. Dover Publications, New York (2013)
17. Ghaffarnejad, H., Niad, H.: Weak gravitational lensing from regular Bardeen black holes. *Int. J. Theor. Phys.* **55**, 1492 (2016)
18. Fu, Q.-M., Zhao, L., Liu, Y.-X.: Weak deflection angle by electrically and magnetically charged black holes from nonlinear electrodynamics. *Phys. Rev. D* **104**, 024033 (2021)

19. de Paula, M.A.A., Leite, L.C.S., Crispino, L.C.B.: Massless scalar scattering by a charged regular black hole. *Astron. Nachr.* **344**, e220115 (2023)
20. de Paula, M.A.A., Lima Junior, H.C.D., Cunha, P.V.P., Crispino, L.C.B.: Electrically charged regular black holes in nonlinear electrodynamics: light rings, shadows and gravitational lensing, [arXiv:2305.04776](https://arxiv.org/abs/2305.04776) [gr-qc]
21. Novello, M., De Lorenci, V.A., Salim, J.M., Klippert, R.: Geometrical aspects of light propagation in nonlinear electrodynamics. *Phys. Rev. D* **61**, 045001 (2000)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.