

# Experimental and Theoretical Investigation of the Reaction of NH<sub>2</sub> with NO at Very Low Temperatures

Published as part of *The Journal of Physical Chemistry virtual special issue "Marsha I. Lester Festschrift"*.

Kevin M. Douglas,\* Daniel Lucas, Catherine Walsh, Mark A. Blitz, and Dwayne E. Heard\*



Cite This: *J. Phys. Chem. A* 2023, 127, 7205–7215



Read Online

ACCESS |



Metrics & More




Article Recommendations




Supporting Information

**ABSTRACT:** The first experimental study of the low-temperature kinetics of the gas-phase reaction between NH<sub>2</sub> and NO has been performed. A pulsed laser photolysis-laser-induced fluorescence technique was used to create and monitor the temporal decay of NH<sub>2</sub> in the presence of NO. Measurements were carried out over the temperature range of 24–106 K, with the low temperatures achieved using a pulsed Laval nozzle expansion. The negative temperature dependence of the reaction rate coefficient observed at higher temperatures in the literature continues at these lower temperatures, with the rate coefficient reaching  $3.5 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at  $T = 26 \text{ K}$ . Ab initio calculations of the potential energy surface were combined with rate theory calculations using the MESMER software package in order to calculate and predict rate coefficients and branching ratios over a wide range of temperatures, which are largely consistent with experimentally determined literature values. These theoretical calculations indicate that at the low temperatures investigated for this reaction, only one product channel producing N<sub>2</sub> + H<sub>2</sub>O is important. The rate coefficients determined in this study were used in a gas-phase astrochemical model. Models were run over a range of physical conditions appropriate for cold to warm molecular clouds (10 to 30 K;  $10^4$  to  $10^6 \text{ cm}^{-3}$ ), resulting in only minor changes (<1%) to the abundances of NH<sub>2</sub> and NO at steady state. Hence, despite the observed increase in the rate at low temperatures, this mechanism is not a dominant loss mechanism for either NH<sub>2</sub> or NO under dark cloud conditions.



## 1. INTRODUCTION

The reaction between NH<sub>2</sub> and NO,  $k_1$ , plays an important role in the thermal DeNO<sub>x</sub> process,<sup>1–5</sup> in which NO<sub>x</sub> is thermally removed by NH<sub>3</sub>, and as such has been studied extensively both experimentally<sup>6–25</sup> and theoretically.<sup>10,24–36</sup> The reaction has three main product channels



Product channel R1a, which involves breaking all three chemical bonds in the reactants and forming three new chemical bonds in the products, is chain terminating, converting two radical species (NH<sub>2</sub> + NO) into two closed shell species (N<sub>2</sub> + H<sub>2</sub>O). Conversely, the product channel R1c is chain propagating, as it produces two new radical species (HN<sub>2</sub> + OH). As such, the branching ratio (BR) between channels R1a and R1c,  $\alpha = k_{1c}/k_1$  (where  $k_1$  is the sum of all product channels,  $k_1 = k_{1a} + k_{1b} + k_{1c}$ ), plays an important role in determining the effectiveness of the NH<sub>3</sub> reducing agent in the DeNO<sub>x</sub> cycle. Both experiment and theory suggest channel R1b to be insignificant even at very high temperatures; this is despite the N<sub>2</sub>O + H<sub>2</sub> channel itself being 192 kJ mol<sup>-1</sup> exothermic, and due


to several high activation barriers on the potential energy surface (PES) leading to this channel that lie above the NH<sub>2</sub> + NO entrance channel.<sup>16,29,31,34</sup> After the initial association of NH<sub>2</sub> with NO, the reaction is believed to take place as a single elementary step without collisions, as evidenced by numerous theoretical studies which indicate that the lifetimes of the intermediates are shorter than the time between collisions at temperatures and pressures of interest and by experimental studies which indicate that the rate coefficient is independent of pressure between a few Torr and an atmosphere.<sup>16,20,25</sup> The HN<sub>2</sub> radical formed via channel R1c has been shown both experimentally<sup>37</sup> and theoretically<sup>35,38,39</sup> to have a short lifetime (<0.5 μs) and promptly dissociates to H + N<sub>2</sub> by tunneling through a small potential energy barrier.

Looking at the available experimental data on the reaction, measurements have been made over a wide range of temper-

Received: May 30, 2023

Revised: July 24, 2023

Published: August 17, 2023



atures, from 200 K up to around 2800 K. These measurements can be broadly divided into two groups: the lower temperature range up to around 1000 K, in which experiments are carried out in reactors or flow cells and the reaction is monitored directly,<sup>6,8,10,13,15,16,18,21,24,25</sup> and the higher temperature range around 1000 K and above, in which experiments are carried out in shock tubes or flames and in which the data requires interpretation using high-temperature kinetic modeling.<sup>7,9,11,12,14,17,19,20,22,23</sup> Although there are some discrepancies in the rate coefficients determined in some earlier works, more recent works from both groups are typically in good agreement and show that the reaction has a negative temperature dependence, with the rate coefficient falling from around  $3 \times 10^{-11}$  to  $1 \times 10^{-12}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup> between 200 and 2500 K. The BR between channels R1a and R1c has also been examined in a large number of experimental studies;<sup>6,11–16,18,20–23,25</sup> again despite a number of discrepancies in some of the earlier works, it is now clear from a number of more recent studies that the BR  $\alpha$  increases from around 0.1 at room temperature to around 0.8 at 2500 K. Of the numerous theoretical studies investigating the reaction, only a limited number have attempted to predict this BR.<sup>10,28,31,34</sup> These studies again indicate that the BR  $\alpha$  increases with increasing temperature.

Both NH<sub>2</sub> and NO have been detected in a variety of astrochemical environments, with nitrogen hydrides and nitrogen oxides both thought to play important roles as nitrogen reservoirs and to potentially participate in the formation of more complex species. NH<sub>2</sub> radicals have been detected in diffuse molecular clouds<sup>40</sup> and in high-density star-forming and protostellar regions.<sup>41</sup> Models are often unable to explain both the absolute and relative abundances of the NH<sub>x</sub> hydrides,<sup>41</sup> and gas phase reactions with species such as NO may help to resolve these discrepancies. For example, a gas-phase only model of a diffuse molecular gas cloud underpredicts the observed abundances of both NH and NH<sub>2</sub> by factors of between 10 and 100, while including processes on dust surfaces increases the abundances of these two species, also results in a failure to match the high NH/NH<sub>3</sub> ratio observed.<sup>41</sup> NO, after its first detection toward SgrB2 in 1978,<sup>42</sup> has been detected in dark clouds, star-forming regions, protostellar envelopes, and nuclei of starburst galaxies (e.g.<sup>43–47</sup>). Despite the possible importance of the reaction between NH<sub>2</sub> and NO in affecting the partitioning of nitrogen between different reservoirs (being a reaction that converts a nitrogen hydride and a nitrogen oxide into N<sub>2</sub>), the absence of the reaction in the Kinetic Database for Astrochemistry<sup>48</sup> suggests the reaction is likely missing in many astrochemical models. The reaction is, however, included in the UMIST Database for Astrochemistry (UdFA), with the recommended rate coefficient listed as appropriate at temperatures from 210 K upward.

In this paper, we present measurements of rate coefficients for the reaction between NH<sub>2</sub> and NO in the temperature range 24–106 K, collected using a pulsed laser-photolysis laser-induced fluorescence (PLP-LIF) technique coupled with a Laval nozzle to achieve the low temperatures relevant to the ISM. We also perform a theoretical investigation into the reaction, recreating the ab initio results of an earlier study to produce a PES and combining this with reaction rate theory using the MESMER software package<sup>49</sup> to calculate both rate coefficients and BRs over a wide temperature range. These data were then incorporated into a single-point gas-phase model for cold to warm molecular cloud conditions, and the new results were compared with those predicted by current models.<sup>50</sup>


## 2. METHODOLOGY

**2.1. Experimental Study.** The use of a Laval nozzle expansion coupled with a PLP-LIF technique has been employed by this group to study the kinetics of a range of neutral–neutral reactions at very low temperatures, including reactions of OH with unsaturated hydrocarbons<sup>51</sup> and oxygenated volatile organic compounds,<sup>52–55</sup> <sup>1</sup>CH<sub>2</sub> with atmospheric gases and hydrocarbons,<sup>56,57</sup> both CH<sup>58</sup> and CN<sup>59</sup> with CH<sub>2</sub>O, and NH<sub>2</sub> with CH<sub>2</sub>O.<sup>60</sup> In the current study, we employ the same technique to study the low-temperature kinetics of the reaction of NH<sub>2</sub> with NO. As the experimental apparatus employed in the current and previous studies has been discussed in detail elsewhere,<sup>51–54</sup> only a brief overview is given here.

The reagent and bath gases were combined in a mixing manifold using calibrated mass flow controllers (MFCs; MKS Instruments) prior to being entered into a 2 L gas ballast tank. The NO reagent was introduced as a pure gas, while the NH<sub>3</sub> precursor was introduced as either a pure gas or as a dilute mixture of between 10 and 50% in Ar. Following the gas ballast, the reaction mixture was introduced to a 1 cm<sup>3</sup> stainless-steel reservoir via two pulsed solenoid valves (Parker 9 series) fired at a repetition rate of either 5 or 10 Hz with a pulse duration of around 10 ms. Each pulse of gas underwent a controlled expansion through a convergent-divergent-shaped Laval nozzle into a low-pressure stainless-steel cylindrical chamber (~775 mm length by 240 mm diameter), resulting in a thermalized low-temperature gas flow. A range of nozzles were employed during the experiments to achieve flow temperatures of between 24 and 106 K. The temperature and density profile of the flows were characterized by impact pressure measurements using a Pitot tube, and the temperature of several of the jets was confirmed by rotationally resolved LIF spectroscopy.<sup>56,58</sup> The pressure in the vacuum chamber, as measured by two calibrated capacitance manometers (Leybold Ceravac CTR100N 0–10 Torr and Leybold Ceravac CTR90 0–1000 Torr), was typically in the range of 0.2–1.5 Torr depending on the nozzle and flow conditions employed and controlled by adjusting the pumping speed on the screw pump (Edwards GXS160).

NH<sub>2</sub> radicals were generated from the PLP of NH<sub>3</sub> at 213 nm (Reaction R2) by the fifth harmonic of a Nd/YAG laser (Quantel Q-smart 850), with a typical pulse energy of ~10 mJ. The photolysis laser was introduced colinearly with the axis of the expanded gas flow to produce a uniform radical density. NH<sub>2</sub> radicals were observed by time-resolved LIF spectroscopy, probing the A<sup>2</sup>A<sub>1</sub> (0,9,0) ← X<sup>2</sup>B<sub>1</sub> (0,0,0) transition near 597.6 nm<sup>61,62</sup> using the output of a Nd/YAG-pumped dye laser (a Quantel Q-smart 850 pumping a Sirah Cobra-Stretch). The probe laser was introduced perpendicularly to the photolysis laser beam, crossing the gas flow at the furthest distance from the exit of the nozzle before the flow broke up due to turbulence. This point was determined from impact pressure measurements as the point at which the density of the flow begins to sharply decline and was typically between 10 and 30 cm, depending on the nozzle and bath gas used. The nonresonant fluorescence at ~620 nm was collected via a series of lenses through an optical filter (Semrock Brightline interference filter,  $\lambda_{\text{max}} = 620$  nm, fwhm = 14 nm) and observed by a temporally gated channel photomultiplier (CPM; PerkinElmer C1952P), mounted at 90° to both laser beams. The signal from the CPM was recorded using a digital oscilloscope (LeCroy Waverunner LT264) and sent to a computer using a custom LabView program. The temporal evolution of the LIF signal was recorded by varying the

time delay between photolysis and probe lasers. A typical time-resolved LIF profile (Figure 1) consisted of between 110 and 165 delay steps and resulted from the average of between 6 and 15 individual delay scans.



**Figure 1.**  $\text{NH}_2$  ( $\nu = 0$ ) LIF signal following PLP of  $\text{NH}_3$  at various  $[\text{NO}]$ , given in units of molecules  $\text{cm}^{-3}$ . Solid lines are fits of eq E1 to the data.  $T = 28.2$  K, total Ar density =  $3.20 \times 10^{16}$  molecules  $\text{cm}^{-3}$ .

Photolysis of  $\text{NH}_3$  produces both ground and vibrationally excited  $\text{NH}_2$ ,<sup>62,63</sup> and, as a consequence of this, the profiles of  $\text{NH}_2$  ( $\nu = 0$ ) exhibit a growth resulting from the relaxation of vibrationally excited  $\text{NH}_2$  (see Figure 1). In experiments in which only the  $\text{NH}_3$  precursor and the bath gas were present, this growth was on the order of  $20,000 \text{ s}^{-1}$ , indicating that it takes  $\sim 170 \mu\text{s}$  for 95% of the vibrationally excited  $\text{NH}_2$  to be relaxed down to  $\nu = 0$ . This posed a problem for kinetic measurements of  $\text{NH}_2$  in our system, as the time scale for kinetic experiments was restricted by the length of the uniform supersonic flow, with dynamic times in our system ranging from  $\sim 200$  to  $500 \mu\text{s}$  depending on the nozzle and bath gas used. Therefore, the addition of a species effective in the vibrational relaxation of  $\text{NH}_2$  was required in order to measure accurate rate coefficients for the removal of  $\text{NH}_2$  ( $\nu = 0$ ). Yamasaki et al.<sup>64</sup> have shown that  $\text{CF}_4$  efficiently accelerates vibrational relaxation of  $\text{NH}_2$ . We have found that  $\text{CH}_4$  is also efficient at relaxing vibrationally excited  $\text{NH}_2$ . By adding up to 3%  $\text{CH}_4$  to our flows, we achieved a growth of the  $\text{NH}_2$  ( $\nu = 0$ ) signal on the order of between  $50,000$  and  $100,000 \text{ s}^{-1}$  (depending on the particular flow), allowing us to effectively monitor the removal of  $\text{NH}_2$  ( $\nu = 0$ ) within the time scales of our experiments. To ensure adding up to 3%  $\text{CH}_4$  to our flows did not significantly affect the temperature and density profiles of our low-temperature gas flows, several flows were characterized by impact pressure measurements both with and without 3%  $\text{CH}_4$  present; the temperatures and densities determined for the flows with  $\text{CH}_4$  present were shown to be within 5% of those determined without  $\text{CH}_4$  present.


**2.2. Materials.**  $\text{N}_2$  (99.9995%, BOC), Ar (99.9995%, BOC),  $\text{NH}_3$  (99.98%, BOC), and NO (99.5%, BOC).

**2.3. Theoretical Calculations.** All electronic structure calculations were carried out using the Gaussian 09 suite of programs.<sup>65</sup> Geometric structures of the stationary points [reactants, products, and intermediates, including adducts and transition states (TSs)] given in Diau and Smith<sup>29</sup> were optimized at the B3LYP/6-311G(d,p) level<sup>66–69</sup> to obtain


vibrational frequencies, rotational constants, and electronic and zero point energies (ZPEs). TSs were found to have only one imaginary vibrational frequency, while for the reactants, products, and intermediates, all of the vibrational frequencies were positive. Diau and Smith<sup>29</sup> also demonstrated that the TSs were connected to their designated intermediates by IRC calculations. ZPEs obtained from the harmonic frequencies were corrected with a scaling factor of 0.967 for B3LYP/6-311G(d,p).<sup>70</sup> Rate theory calculations to predict rate coefficients and BRs were performed using the master equation solver for multi-energy well reaction (MESMER) program.<sup>49</sup> Further details of the parameters used in the MESMER program, such as the energy transfer parameters  $\Delta E_{\text{down},\nu}$  are given in the MESMER input file attached as part of the Supporting Information. However, as discussed below, as the reaction between  $\text{NH}_2 + \text{NO}$  is effectively pressure-independent over pressure ranges of interest, the choice of the magnitude of the energy transfer parameters does not play a role in the rate coefficients and BRs calculated.

### 3. RESULTS

Typical  $\text{NH}_2$  LIF profiles produced following the photolysis of  $\text{NH}_3$  can be seen in Figure 1. As can be seen from these profiles, there is an initial growth of the  $\text{NH}_2$  signal with no instant signal observed. As discussed above, this growth in the  $\text{NH}_2$  ( $\nu = 0$ ) signal is due to relaxation of vibrationally excited  $\text{NH}_2$  produced following photolysis of  $\text{NH}_3$ . Relaxation of vibrationally excited  $\text{NH}_2$  occurs via a cascade, such that  $\nu = x \rightarrow \nu = x - 1 \rightarrow \nu = x - 2 \rightarrow \dots \rightarrow \nu = 0$ . As such, the growth of the  $\text{NH}_2$  ( $\nu = 0$ ) signal is not strictly a single exponential growth. However, as higher vibrational levels are relaxed faster than lower levels (for a particular collider),<sup>63</sup> the relaxation of  $\text{NH}_2$  ( $\nu = 1$ ) to  $\text{NH}_2$  ( $\nu = 0$ ) is the rate-limiting step, and as such, the growth of the  $\text{NH}_2$  ( $\nu = 0$ ) signal can be treated as a single-exponential growth (R3)



where X can be  $\text{CH}_4$ , the bath gas,  $\text{NH}_3$ , and the coreagent R under investigation. In practice, relaxation by the bath gas (either Ar or  $\text{N}_2$ ) is slow,<sup>71,72</sup> necessitating the addition of  $\text{CH}_4$  to our flows to promote efficient vibrational relaxation. Reaction R3, together with the reaction for the removal of  $\text{NH}_2$  ( $\nu = 0$ )



and as experiments were carried out under pseudo-first-order conditions (i.e.,  $[\text{NH}_2] \ll [\text{NO}]$  and  $[\text{CH}_4]$ ), the temporal evolution of the  $\text{NH}_2$  ( $\nu = 0$ ) LIF signal is given by


$$[\text{NH}_2 (\nu = 0)]_t = \left( \frac{k'_{\text{rel}}}{k'_{\text{obs}} - k'_{\text{rel}}} \right) [\text{NH}_2 (\nu \geq 1)]_0 (\exp^{-k'_{\text{rel}} t} - \exp^{-k'_{\text{obs}} t}) \quad (\text{E1})$$

and

$$k'_{\text{obs}} = k_1[\text{NO}] + k'_{\text{loss}} \quad (\text{E2})$$

where  $k'_{\text{rel}}$  and  $k'_{\text{obs}}$  are the pseudo-first-order rate coefficients for the reactions producing and removing  $\text{NH}_2$  ( $\nu = 0$ ),  $[\text{NH}_2 (\nu \geq 1)]_0$  is the initial amount of  $\text{NH}_2$  ( $\nu \geq 1$ ) produced following photolysis of  $\text{NH}_3$  that is subsequently relaxed down to  $\text{NH}_2$  ( $\nu = 0$ ), and  $k'_{\text{loss}}$  is the total rate coefficient for other loss processes of  $\text{NH}_2$  ( $\nu = 0$ ), such as diffusion out of the probe laser beam

volume. Equation E1 was fitted to the  $\text{NH}_2$  ( $\nu = 0$ ) LIF profiles, and the parameters  $k_{\text{rel}}'$ ,  $k_{\text{obs}}'$ , and  $[\text{NH}_2$  ( $\nu \geq 1$ )]<sub>0</sub> were extracted. When fitting these  $\text{NH}_2$  profiles, the  $k_{\text{rel}}'$  values were fit globally; that is, for a certain experimental run in which the total pressure, temperature, precursor concentration, and  $[\text{CH}_4]$  were kept constant, all the  $\text{NH}_2$  profiles were fit using a single  $k_{\text{rel}}'$  value that was allowed to float. As can be seen from Figure 1, the  $\text{NH}_2$  traces are satisfactorily fit using a biexponential function, indicating the validity of treating the growth of the  $\text{NH}_2$  ( $\nu = 0$ ) signal as a single process. Plotting  $k_{\text{obs}}'$  vs  $[\text{NO}]$  should then yield a straight line as given by eq E2, with a gradient equal to the bimolecular rate constant,  $k_1$ , and intercept  $k_{\text{loss}}'$ . Examples of such bimolecular plots can be seen in Figure 2. As



**Figure 2.** Bimolecular plots of  $k_{\text{obs}}'$  vs  $[\text{NO}]$  collected at various temperatures, using either (a) Ar or (b)  $\text{N}_2$  as a bath gas. The solid symbols indicate the linear range from which the bimolecular rate coefficient,  $k_1$ , is determined, while the open symbols demonstrate curvature caused by the presence of NO dimers and are excluded from the linear fit. No curvature was observed when  $\text{N}_2$  was used as a bath gas.

can be seen from Figure 2, in some of our bimolecular plots, we observe a curvature in the  $k_{\text{obs}}'$  values at high  $[\text{NO}]$ . We attribute this curvature to the presence of NO dimers, which are formed in our low-temperature flows at high  $[\text{NO}]$ . The  $[\text{NO}]$  at which NO dimers appear in a particular flow will depend on the bath gas used and the temperature and density of the flow, with dimers more likely to form at lower temperatures, higher densities, and with heavier bath gases that act better as third bodies. Indeed, we only observe this curvature in our lowest temperature flows that use Ar as a bath gas, while no curvature was observed in our higher temperature flows using  $\text{N}_2$  as a bath gas. The negative curvature at higher  $[\text{NO}]$  implies that NO dimers do not remove  $\text{NH}_2$  fast enough to counterbalance the loss of NO monomers (i.e., that the NO dimer removes  $\text{NH}_2$  less

than twice as fast as the NO monomer). In bimolecular plots in which curvature was observed, only the linear part of the plot was used to determine  $k_1$  (e.g., solid points in Figure 2), while the other points were excluded (e.g., open points in Figure 2).

An alternative explanation for the curvature that we observe in some of our bimolecular plots is that it is the result of our data fitting procedure rather than due to the presence of NO dimers. At higher  $[\text{NO}]$ , when the kinetic decays become faster, it may be that the  $\text{NH}_2$  ( $\nu \geq 0$ ) relaxation is not complete, resulting in  $\text{NH}_2$  ( $\nu = 0$ ) decays that may appear slower than expected. To determine if this is the case, we have also fit single exponentials to the  $\text{NH}_2$  traces at long times (i.e., from times at which all of the  $\text{NH}_2$  ( $\nu \geq 1$ ) should have been relaxed). Rate coefficients,  $k_1$ , determined fitting single exponential decays to the data were typically within 10% of those determined when using biexponential fits, with many values in much better agreement. In addition, we typically still observe a curving over of the bimolecular plot at higher  $[\text{NO}]$  when using data fitted with a single exponential as well as when fitted with a biexponential. As such, it is likely that this curving over is in fact a result of NO dimerization rather than an artifact of our data fitting procedure. This NO dimerization has not been reported in other low-temperature gas expansion experiments using a Laval nozzle,<sup>73–75</sup> although the characteristics of each low-temperature expansion do differ. NO dimers do, however, form readily in a number of low-temperature molecular beam experiments (e.g.<sup>76,77</sup>), although it should be noted that the temperatures in these experiments are likely significantly lower in our low-temperature flows. The NO dimer itself also has a relatively large binding of  $\sim 14$  kJ mol<sup>-1</sup> for such a small molecule.<sup>78</sup> In either case, whether this curving over is the result of NO dimers or a data analysis artifact, it is the result of a deviation from the true rate coefficient, warranting the exclusion of these points from the bimolecular plot in our determination of  $k_1$ .


The bimolecular rate coefficients for the reaction of  $\text{NH}_2$  with NO ( $k_1$ , Reaction R1) determined in this study are presented in Table 1 and compared with some of the more recent literature data in Figure 3. The errors reported are the 1  $\sigma$  confidence intervals of linear least-squares fits of the bimolecular plots and do not include systematic errors. No change in the rate coefficient was observed as the total pressure of the flows varied by around a factor of 4, which is consistent with previous studies of the system.<sup>10,25</sup>

There have been many previous studies investigating the removal of  $\text{NH}_2$  with NO over temperatures ranging from 200 to 2500 K, employing a range of experimental techniques, with the results presented here being the first low-temperature measurements below 200 K. Figure 3 presents some of the more recent temperature-dependent rate coefficients given in the literature. The highest temperature data above  $\sim 1200$  K are from two shock tube studies carried out by Song et al.,<sup>19,20</sup> in which the  $\text{NH}_2$  is detected using frequency modulation absorption spectroscopy, and from a study by Vandooren et al.,<sup>22</sup> in which either the NO consumption or  $\text{N}_2$  production in a low-pressure ammonia-nitric oxide flame was monitored by mass spectrometry, with the rate coefficients extracted following high-temperature kinetic modeling of the data. The midtemperature range data between  $\sim 200$  and 1100 K come from three studies by Wolf et al.,<sup>25</sup> Diau et al.,<sup>10</sup> and Park and Lin,<sup>16</sup> who, following PLP of  $\text{NH}_3$  in the presence of NO, monitor either the removal of  $\text{NH}_2$  or the growth of  $\text{H}_2\text{O}$  using LIF, cavity ring down spectroscopy, and mass spectrometry, respectively.

Table 1. Rate Coefficients for the Reaction of  $\text{NH}_2 + \text{NO}$  and Relevant Experimental Conditions

$\sim T/\text{K}$	$T^a/\text{K}$	bath gas	$N_{\text{total}}^a/10^{16}$ molecules $\text{cm}^{-3}$	dynamic time/ $\mu\text{s}$	$[\text{CH}_4]/10^{15}$ molecules $\text{cm}^{-3}$	$[\text{NO}]$ range <sup>b</sup> / $10^{13}$ molecules $\text{cm}^{-3}$	$k_1(T)^c/10^{-10}$ $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$
26	$28 \pm 2$	Ar	$3.2 \pm 0.4$	$\sim 500$	0.63	2.8 (4.9)	$3.6 \pm 0.1$
	$26 \pm 2$	Ar	$4.2 \pm 0.5$	$\sim 500$	0.83	2.4 (3.8)	$3.8 \pm 0.2$
	$26 \pm 2$	Ar	$6.2 \pm 0.5$	$\sim 500$	1.3	2.2 (6.2)	$3.4 \pm 0.5$
	$24 \pm 2$	Ar	$7.7 \pm 0.6$	$\sim 500$	1.6	2.6 (8.5)	$3.4 \pm 0.5$
	$41 \pm 3$	Ar	$8.6 \pm 0.8$	$\sim 250$	1.3	3.3 (5.9)	$2.5 \pm 0.2$
	$49 \pm 4$	Ar	$6.3 \pm 0.8$	$\sim 225$	1.2	3.2 (3.7)	$2.9 \pm 0.2$
	$60 \pm 8$	Ar	$7.9 \pm 1.5$	$\sim 270$	1.3	2.9 (4.6)	$2.0 \pm 0.1$
	66	$69 \pm 2$	$\text{N}_2$	$2.2 \pm 0.2$	$\sim 400$	0.43	10.3
$66 \pm 2$		$\text{N}_2$	$3.7 \pm 0.2$	$\sim 400$	0.71	11.0	$2.3 \pm 0.1$
$63 \pm 2$		$\text{N}_2$	$4.9 \pm 0.4$	$\sim 400$	0.97	9.1	$2.4 \pm 0.1$
$64 \pm 2$		$\text{N}_2$	$6.5 \pm 0.4$	$\sim 360$	1.3	11.5	$2.2 \pm 0.1$
$85 \pm 2$		$\text{N}_2$	$7.5 \pm 0.4$	$\sim 180$	1.2	4.5	$1.5 \pm 0.1$
$91 \pm 6$		$\text{N}_2$	$4.9 \pm 0.9$	$\sim 280$	0.90	2.5	$1.4 \pm 0.1$
$106 \pm 9$		$\text{N}_2$	$6.7 \pm 1.5$	$\sim 200$	1.2	3.6	$0.95 \pm 0.05$

<sup>a</sup>Uncertainties in each value of  $T$  and  $N_{\text{total}}$  are  $\pm 1 \sigma$  (the standard deviation) of the measured temperature and density along the axis of the Laval expansion. <sup>b</sup>Gives the  $[\text{NO}]$  range from 0 to the stated value that was used in determining  $k_1$ , while the value in brackets gives the full range of  $[\text{NO}]$  used in the experiment (if different). <sup>c</sup>Uncertainties for each value of  $k(T)$  reported at the  $1 \sigma$  level for the linear least-squares fitting of the pseudo-first-order rate coefficients as a function of  $[\text{NO}]$ .



**Figure 3.** Temperature dependence of the rate coefficient for  $\text{NH}_2 + \text{NO}$  (Reaction R1). Blue squares: this study. Red upward triangles: Wolf, 1997.<sup>25</sup> Yellow downward triangles: Park, 1997.<sup>16</sup> Turquoise left pointing triangles: Diau, 1994.<sup>10</sup> Purple diamonds: Vandooren, 1994.<sup>22</sup> Green circles: Song, 2000<sup>19</sup> and 2001.<sup>20</sup> Black stars: MESMER rate-theory predictions; see Section 4 for details. The inset is zoomed in over the temperature range investigated in this study. Blue and red dotted lines are the parametrized fits to the low- and high-temperature data. The green dashed line is a parametrized fit to all the data (see text). Errors have been left off the main plot for clarity.

As can be seen from Figure 3, our low-temperature data follow the general trend of the higher-temperature data, with the rate of removal of  $\text{NH}_2$  by  $\text{NO}$  increasing with decreasing temperature. Parameterizing the data over the entire temperature range did not give a satisfactory fit to the low-temperature data obtained in this study (see dashed green line in the inset of Figure 3). In particular, an all-temperature parametrization does not take into account the apparent slowdown in the increase in rate coefficient with decreasing temperature, and extrapolating such a fit down to 10 K yields a rate coefficient that is around 4 times faster than that measured at 25 K and that is approaching the theoretical classical capture theory collision limit (estimated to be  $10^{-9} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  at 10 K, by comparison with a similarly sized


system).<sup>59</sup> Instead, both the low-temperature data below 110 K and the high-temperature data above 200 K were parametrized separately (eqs E3 and E4) using a modified Arrhenius expression, giving the blue and red dotted lines shown in Figure 3, respectively. This resulted in a significantly better fit of the low-temperature data provided in this study and takes account of the apparent slowdown in the increase of the rate coefficient. Additionally, extrapolation of the low-temperature parametrized fit yields a rate coefficient at 10 K around twice as fast as that determined at 25 K, and that is well below the theoretical collision limit. All experimental data points were given a 10% error when parametrizing the data to ensure no one data point at a particular temperature was overly weighted. With the exception of the Vandooren et al. data, which report considerably larger rate coefficients than the other studies over a similar temperature range, the majority of the experimental data points lie within 20% of the parametrized rate coefficients, with most deviating by only a few percent. The low- and high-temperature parametrized rate coefficients are given by (errors are  $1 \sigma$ ):

$$k_{(\text{NH}_2+\text{NO})}(10 \leq T/\text{K} \leq 110) = (6.20 \pm 1.02) \times 10^{-11} \times (T/300)^{(-0.704 \pm 0.089)} \quad (\text{E3})$$

$$k_{(\text{NH}_2+\text{NO})}(10 \leq T/\text{K} \leq 2510) = (9.67 \pm 1.63) \times 10^{-12} \times (T/300)^{(-1.08 \pm 0.08)} \times \exp^{[(129 \pm 56)/T]} \quad (\text{E4})$$

## 4. DISCUSSION


**4.1. Theoretical Calculations.** For the reaction between  $\text{NH}_2$  and  $\text{NO}$ , there have been several theoretical investigations, with a number of PESs calculated at various levels of theory.<sup>10,24–36</sup> For this study, we have taken the structures given in Diau and Smith<sup>29</sup> and optimized them at the B3LYP/6-311G(d,p) level in order to obtain vibrational frequencies, rotational constants, and electronic and ZPEs, details of which can be found in the MESMER input file, which is included in the Supporting Information. The calculated energies and vibrational frequencies are in excellent agreement with those reported by Diau and Smith<sup>29</sup> using the same level of theory. Table S1 compares our calculated energies to some of the more recent



**Figure 4.** PES for  $\text{NH}_2 + \text{NO}$ , determined at the B3LYP/6-311G(d,p) level of theory. The  $\text{N}_2\text{O} + \text{H}_2$  product channel (R1b) and the TSs and stationary points leading to it have been excluded (see the text for details).

calculated energies in the literature.<sup>29,31</sup> As can be seen from Table S1, although B3LYP/6-311G(d,p) could be considered a relatively low level of theory by modern computational standards, the energies calculated at this level of theory are in reasonable agreement with those from higher-level calculations. Comparing the energies of the adducts 1–5 and the TSs a–f (see Figure 4), the B3LYP energies are typically within 15  $\text{kJ mol}^{-1}$  of those calculated at the G2M(CC1) level of theory<sup>79</sup> by Diao and Smith<sup>29</sup> and in even better agreement with those calculated at the CCSD(T)/aug-cc-pVTZ level of theory by Fang et al.,<sup>31</sup> being within 9  $\text{kJ mol}^{-1}$ . Comparing the energies of the key three stationary points (TSs c and e, and the  $\text{HN}_2 + \text{OH}$  exothermicity, see discussion below), we typically see even better agreement between the B3LYP and higher-level calculations, and indeed, when fitting these parameters to the experimental data (see below), only minor adjustments to these energies are required. As such, we have chosen not to refine the B3LYP energies by carrying out higher-level electronic energy calculations.

The  $\text{NH}_2 + \text{NO}$  PES is complex, with many deep potential wells and large barriers and three possible product channels




A schematic of the full PES for the reaction can be seen in the Supporting Information (Figure S2). As discussed above, the TSs leading to the  $\text{N}_2\text{O} + \text{H}_2$  product channel (R1b) all lie above the  $\text{NH}_2 + \text{NO}$  entrance channel. As such, both experiment and theory suggest channel R1b to be a very minor channel, even at temperatures as high as 2500 K. Indeed, preliminary MESMER calculations carried out using the full PES indicated that channel R1b accounts for less than 0.2% of the total rate coefficient at 2500 K. As such, all further MESMER calculations were carried out using the reduced surface shown in Figure 4, in which the stationary points leading to channel R1b are excluded. As can be seen from Figure 4, the reaction is initiated by the barrierless addition of  $\text{NH}_2$  to the  $\text{NO}$  to form the adduct  $\text{H}_2\text{NNO}$  (structure 1, Figure 4). This adduct undergoes a H atom shift followed by a series of cis–trans isomerizations, giving rise to four distinct  $\text{HNNOH}$  isomers. All four of these isomers (structures 2–5 in Figure 4) may undergo an endothermic, barrierless bond fission to form the products  $\text{HN}_2 + \text{OH}$ , while

only one of the isomers (no. 5) is configured correctly to form the  $\text{N}_2 + \text{H}_2\text{O}$  products via a four-membered ring TS.

**4.2. Rate Theory Calculations Using MESMER.** Using the program MESMER, we can fit the experimental data by allowing various parameters used in the rate theory calculation or features of the PES itself to be adjusted. In this manner, we are able to improve the agreement between the experimental and calculated rate coefficients and BRs. Rather than fitting all of the available experimental data, in order to reduce the computational time required when carrying out the rate theory calculations, fitting was carried out using only a limited number of rate coefficient and product BR values, as given by the parametrized fits (eqs E3–E5) to the experimental data (see Figures 3 and 5). The temperatures and values used in the fitting, together with the MESMER predicted values, are given in Table 2. The parameters that were adjusted, together with the values obtained from the fitting, are given in Table 3. Initially, each parameter was fitted independently to assess its effect on the predicted rate coefficients/BRs, before a final fitting was carried out in which all the parameters were floated together. When parametrizing the experimental BR data from the literature (see Figure 5), each data point was given a  $\pm 0.1$  error to ensure no one data point at a particular temperature was overly weighted, giving (errors are 1  $\sigma$ )

$$\alpha (300 \leq T/\text{K} \leq 2500) = (0.0811 \pm 0.021) \times (T/300)^{1.12 \pm 0.10} \times \exp^{[(25 \pm 133)/T]} \quad (\text{E5})$$

The key parameter adjusted when fitting the experimental rate coefficients were the inverse Laplace transform (ILT) parameters for the initial association reaction of  $\text{NH}_2$  with  $\text{NO}$ , which take the form of a modified Arrhenius function  $A(T/300)^n$  (with the activation energy being set to zero for the barrierless process). The final fitting returned values of  $A_1 = 2.02 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  and  $n_1 = -0.977$  for the initial association reaction of  $\text{NH}_2$  with  $\text{NO}$ , giving a satisfactory fit to rate coefficients over almost the entire temperature range (60–2500 K, see Figure 3), with the predicted rate coefficients obtained using these  $A$  and  $n$  values typically being within 10% of the rates given by the parametrized fits (eqs E3 and E4; see Table 2). Due to the  $\text{NH}_2 + \text{NO}$  PES containing many deep wells, MESMER was unable to predict or fit rate coefficients and BRs below 60 K.



**Figure 5.** Experimentally determined BR  $\alpha = k_{1c}/k_1$  as a function of temperature. Experimental error bars have not been included for clarity but are quoted below where given. Red upward triangles: Wolf, 1997, error given as  $\pm 0.03$ .<sup>25</sup> Yellow downward triangles: Park, 1997, errors not given.<sup>16</sup> Purple diamonds: Vandooren, 1994, errors not given.<sup>22</sup> Green circles: Song, 2000<sup>19</sup> and 2001, errors not given.<sup>50</sup> Blue hexagons: Halbgewachs 1996, average quoted error  $\pm 0.01$ .<sup>12</sup> Orange stars: Glarborg 1997, average quoted error  $\pm 0.03$ .<sup>11</sup> Light green left facing triangles: Deppe, errors not given, 1997.<sup>9</sup> Brown pentagons: Votsmeier, 1999, estimated average error  $\pm 0.05$ .<sup>23</sup> Black stars: MESMER predictions. The red dotted line is a parametrized fit to literature data (eq E5). The inset shows the temperature dependence of the rate coefficients over the temperature range  $T = 110$ – $2510$  K for; black dotted line: total  $\text{NH}_2 + \text{NO}$  removal rate coefficient ( $k_1$ ). Red dashed line: product channel  $\text{N}_2 + \text{H}_2\text{O}$  ( $k_{1a}$ , eq E6). Blue dot dash line: product channel  $\text{HN}_2 + \text{OH}$  ( $k_{1c}$ , eq E7).

We have identified several parameters that are likely to affect the BR  $\alpha$  for the  $\text{HN}_2 + \text{OH}$  product channel predicted by MESMER. Key among these are the reverse ILT parameters used for the rate coefficients of the barrierless dissociation reactions of the four  $\text{HNNOH}$  isomers (nos. 2–5 in Figure 4). These parameters,  $A_2$  and  $n_2$ , again take the form of a modified Arrhenius function,  $A(T/300)^n$ . Although the  $A$  and  $n$  parameters for the dissociation reaction of each distinct  $\text{HNNOH}$  isomer could be varied individually, we have used the same  $A$  and  $n$  values for all four reactions due to the similarity

in the dissociation reactions, as has been done in previous theoretical investigations.<sup>25,34</sup> Indeed, in the more recent theoretical investigation of this reaction by Fang et al.,<sup>31</sup> in which they do treat the dissociation of the cis and trans isomers separately, they indicate that the dissociation rate coefficients are typically within 30% of one another, indicating that this simplification should not represent too large an error. Fitting of the experimental data returned values of  $A_2 = 1.30 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$  and  $n_2 = -0.079$ , indicating little temperature dependence on the dissociation rate. In addition to the reverse ILT parameters, two other parameters have been shown to be important factors in determining the BR  $\alpha$ ; the  $\text{HN}_2 + \text{OH}$  endothermicity and the heights of the barriers linking the cis–trans isomers (TSe and TSc).<sup>31,34</sup> When adjusting these parameters, we treat the cis–trans isomer barriers as a pair, allowing the energies of both to move up or down in tandem while maintaining the absolute difference in energy between the two TSs, as attempting to float both TSs independently resulted in significantly larger errors in all of the fitted parameters, suggesting that the results were not defined. These two parameters (the  $\text{HN}_2 + \text{OH}$  endothermicity and the TSe/TSc barrier heights) in some ways compensate each other. For example, by lowering the heights of the TSe/TSc pair, we effectively promote the cis–trans isomerization processes, leading to adduct *S*, which promptly dissociates to  $\text{N}_2 + \text{H}_2\text{O}$ , resulting in a decrease in  $\alpha$ . However, this may be countered by decreasing the  $\text{HN}_2 + \text{OH}$  product endothermicity, which in effect promotes this channel, resulting in an increase in  $\alpha$ . How these two parameters are adjusted has also been shown to affect the shape of the BR vs  $T$  plot (Figure 5) by affecting how  $\alpha$  increases with temperature. The best fit to the experimental data required increasing the  $\text{HN}_2 + \text{OH}$  exothermicity by  $2.1 \text{ kJ mol}^{-1}$  (from  $7.5$  to  $9.6 \text{ kJ mol}^{-1}$ ) and decreasing the heights of the TSe/TSc pair by  $4.1 \text{ kJ mol}^{-1}$  (from  $-26.7$  to  $-30.8 \text{ kJ mol}^{-1}$  and from  $-36.2$  to  $-40.3 \text{ kJ mol}^{-1}$  for TSe and TSc, respectively). These are much smaller changes than those made by Fang et al.,<sup>31</sup> in which they move the ZPEs of the  $\text{HN}_2 + \text{OH}$  channel and the TSe/TSc pair, as calculated at the CCSD(T)/aug-cc-pVTZ level of theory, down by  $7$  and  $17 \text{ kJ mol}^{-1}$ , respectively. This in effect moves their values from being around  $5 \text{ kJ mol}^{-1}$  higher than our calculated values to around  $6 \text{ kJ mol}^{-1}$  lower than our values following fitting. It should be

**Table 2.** Comparison of the Rate Coefficient for the  $\text{NH}_2 + \text{NO}$  Reaction and BR  $\alpha = k_{1c}/k_1$  Calculated Using MESMER with the Values Given by the Parametrized Fits to the Experimental Data (eqs E3–E5)

T/K	$k_1/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$			BR $\alpha$ for $\text{HN}_2 + \text{OH}$ product channel (R1c)		
	parameterized fits <sup>a</sup>	MESMER	% difference (%)	parameterized fit	MESMER	difference
60	$1.92 \times 10^{-10}$	$2.26 \times 10^{-10}$	17.7		0.000	
90	$1.45 \times 10^{-10}$	$9.27 \times 10^{-11}$	36.1		0.000	
180	$3.44 \times 10^{-11}$	$3.11 \times 10^{-11}$	9.6		0.000	
300	$1.49 \times 10^{-11}$	$1.53 \times 10^{-11}$	2.7	0.075	0.008	0.067
500	$7.19 \times 10^{-12}$	$7.64 \times 10^{-12}$	6.3	0.137	0.058	0.079
750	$4.26 \times 10^{-12}$	$4.41 \times 10^{-12}$	3.5	0.220	0.175	0.045
1000	$2.99 \times 10^{-12}$	$3.00 \times 10^{-12}$	0.3	0.306	0.308	0.002
1250	$2.28 \times 10^{-12}$	$2.24 \times 10^{-12}$	1.8	0.395	0.432	0.037
1500	$1.84 \times 10^{-12}$	$1.79 \times 10^{-12}$	2.7	0.487	0.536	0.049
1750	$1.54 \times 10^{-12}$	$1.50 \times 10^{-12}$	2.6	0.580	0.620	0.04
2000	$1.32 \times 10^{-12}$	$1.31 \times 10^{-12}$	0.8	0.675	0.685	0.01
2250	$1.15 \times 10^{-12}$	$1.17 \times 10^{-12}$	1.7	0.772	0.735	0.037
2500	$1.02 \times 10^{-12}$	$1.07 \times 10^{-12}$	4.9	0.870	0.773	0.097

<sup>a</sup>Rate coefficients calculated using both the low ( $<110$  K, E3) and high ( $>110$  K, E4) temperature parametrized fits.

**Table 3. Parameters Adjusted during the Fitting to the Experimental Data in the Rate Theory Calculations together with the Values Obtained from the Fitting (Errors are 1  $\sigma$ ). See Text for Details**

parameter	B3LYP <sup>b</sup>	G2M <sup>c</sup>	CCSD(T) <sup>d</sup>	fitting <sup>e</sup>
$A_1/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	$(2.12 \pm 0.11) \times 10^{-11}$	$(1.81 \pm 0.07) \times 10^{-11}$	$(2.95 \pm 0.23) \times 10^{-11}$	$(2.02 \pm 0.19) \times 10^{-11}$
$n_1$	$-0.960 \pm 0.040$	$-1.090 \pm 0.030$	$-0.774 \pm 0.059$	$-0.977 \pm 0.044$
$A_2/\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$	$(8.12 \pm 1.41) \times 10^{-12}$	$(3.97 \pm 0.90) \times 10^{-11}$	$(1.25 \pm 0.26) \times 10^{-11}$	$(1.30 \pm 0.69) \times 10^{-11}$
$n_2$	$-0.101 \pm 0.034$	$-0.075 \pm 0.031$	$-0.198 \pm 0.149$	$-0.079 \pm 0.069$
$\text{HN}_2 + \text{OH ZPE}/\text{kJ mol}^{-1}$	7.5 (fixed)	15.1 (fixed)	10.5 (fixed)	9.6 $\pm$ 3.5
TSe ZPE/ $\text{kJ mol}^{-1}$	-26.7 (fixed)	-33.9 (fixed)	-18.4 (fixed)	-30.8 $\pm$ 8.4
TSc ZPE <sup>a</sup> / $\text{kJ mol}^{-1}$	-36.2 (fixed)	-45.6 (fixed)	-32.6 (fixed)	-40.3 $\pm$ 8.4 (linked <sup>f</sup> )
$\chi^2$ (degrees of freedom) <sup>a</sup>	1.127 (19)	0.983 (19)	1.070 (19)	1.293 (17)

<sup>a</sup>Values given are  $\chi^2/\text{degrees of freedom}$  (experimental points minus parameters, given in brackets). <sup>b</sup>Values obtained using B3LYP/6-311G(d,p) energies. <sup>c</sup>Values obtained using G2M(CC1) energies.<sup>29</sup> <sup>d</sup>Values obtained using CCSD(T)/aug-cc-pVTZ energies.<sup>31</sup> <sup>e</sup>Values obtained using B3LYP/6-311G(d,p) energies and allowing the energies of the key stationary points to float. <sup>f</sup>TSe and c were not floated independently. Instead, TSe was allowed to float while TSc was derived from TSe, meaning both would move up and down in step with each other.

noted that at the level of theory employed in this study, we would expect errors of around 10  $\text{kJ mol}^{-1}$  in our calculated energies.

In addition to floating all of the parameters discussed above, we have also carried out some additional runs in which we only allow the ILT parameters to float while fixing the energies of the stationary points on the PES to either the B3LYP/6-311G(d,p) values determined in this study, the G2M(CC1) values determined by Diau and Smith<sup>29</sup> or the CCSD(T)/aug-cc-pVTZ values determined by Fang et al.<sup>31</sup> in order to compare the surfaces from the three levels of theory. Table 3 gives the values of the fitted parameters together with the goodness of each fit (the  $\chi^2$  values). As can be seen from Table 3, we were able to obtain reasonable fits to the experimental data by fitting the ILT parameters only when using energies from all three levels of theory, with only a slight improvement to the fit when using the higher levels of theory. This is unsurprising considering how similar the energies of the stationary points are at each level of theory (Table S1). What is slightly surprising is that we actually see a small decrease in the quality of the fit when allowing all of the parameters to float when using the B3LYP energies; we put this down to the fact that there are now more degrees of freedom and that the parameters we chose to float (the  $\text{HN}_2 + \text{OH}$  endothermicity and the TSe/TSc barrier heights) only required minor adjustments to achieve a good fit to the experimental data.

BRs predicted by MESMER using the values obtained from the fitting of the experimental data are given in Table 2 and shown in Figure 5. As can be seen from Figure 5, there is reasonable agreement between the experimental and calculated BRs over the entire temperature range of available experimental data (300–2500 K), with the predicted BRs lying within  $\pm 0.1$  of the parametrized experimental values (eq 5; see Table 2). The largest discrepancies between the experimental and theoretical BRs appear at both the high and low ends of the temperature range, with the predicted BRs showing a more pronounced sinusoidal shape than is reflected by the experimental data. Fang et al.<sup>31</sup> also noted the difficulty in calculating a BR that is in good agreement with the literature data at room temperature, suggesting the possibility of experimental error in the room-temperature measurements or other possible shortcomings in the theoretical model, such as understanding the dissociation rates of the different isomers or the treatment of the density of states of some of the TSs, which is used in the calculation of the rate coefficient. However, despite these shortcomings, we are able to predict temperature-dependent BRs that are largely

consistent with experiment. Extending the temperature range over which we predict BRs down to 60 K, we can see that as the temperature drops to 180 K and below, the BR  $\alpha$  has reached zero, indicating that the endothermic  $\text{HN}_2 + \text{OH}$  product channel (R1c) is unviable at these low temperatures; conversely, the BR for the low energy channel producing  $\text{N}_2 + \text{H}_2\text{O}$  (R1a) reaches 1 at 180 K and below. As such, the low-temperature parametrized fit given in Section 3 can be taken as having only one product channel, R1a, producing  $\text{N}_2 + \text{H}_2\text{O}$ .

In order to produce rate coefficient expressions for the two product channels open at higher temperatures, we have applied the BRs predicted by MESMER to the rate coefficients given by the high-temperature parametrized fit in Section 3 (eq E4) to give channel-specific rate coefficients. These channel-specific rate coefficients can be parametrized as (see inset Figure 5; errors are 1  $\sigma$  level of a least-squares fit to the data)

$$k_{(\text{NH}_2+\text{NO}\rightarrow\text{N}_2+\text{H}_2\text{O})}(110 \leq T/\text{K} \leq 2500) = (2.37 \pm 0.40) \times 10^{-11} \times (T/300)^{-2.08 \pm 0.09} \times \exp^{[(-115 \pm 40)/T]} \quad (\text{E6})$$

$$k_{(\text{NH}_2+\text{NO}\rightarrow\text{HN}_2+\text{OH})}(110 \leq T/\text{K} \leq 2500) = (6.23 \pm 0.67) \times 10^{-12} \times (T/300)^{-0.696 \pm 0.056} \times \exp^{[(-1183 \pm 26)/T]} \quad (\text{E7})$$

These rate coefficients and BRs predicted by MESMER are, in effect, the zero pressure rate coefficients and BRs that are applicable to the interstellar medium (ISM). This was confirmed by the fact that rate coefficients and BRs calculated between 60 and 2500 K were shown to be pressure independent over the range  $1 \times 10^{12}$  to  $1 \times 10^{20}$   $\text{molecules cm}^{-3}$ ; only at pressures of  $1 \times 10^{21}$   $\text{molecules cm}^{-3}$  and above was any pressure dependence observed.

**4.3. Astrochemical Implications.** The rate coefficients and product channels for the  $\text{NH}_2 + \text{NO}$  reaction were added to a chemical network used for modeling interstellar and circumstellar material.<sup>80</sup> The gas-phase chemistry in this network is based on the Rate12 release of the UMIST Database for Astrochemistry.<sup>50</sup> A series of single-point gas-phase only chemical models were run, covering the temperature (10–30 K) and density ( $10^4$ – $10^6$   $\text{molecules cm}^{-3}$ ) ranges typical of cold to warm molecular clouds. Fully shielded conditions were assumed (i.e., a high visual extinction), and a cosmic-ray ionization rate of  $1.3 \times 10^{-17} \text{ s}^{-1}$  was used. Initial abundances are taken from Table 3 in McElroy et al.<sup>50</sup> The models were run until a time of  $10^8$  years to steady state. A comparison of the results obtained with and without the updated rate coefficients showed no major changes in the abundances of the reactants



(NH<sub>2</sub> and NO) nor products (H<sub>2</sub>O and N<sub>2</sub>), with maximal differences on the order of ~1%. At a temperature of 10 K, Rate12 adopts a recommended rate coefficient of  $1.49 \times 10^{-12}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>, which is around 450 times lower than the updated value from this work ( $6.70 \times 10^{-10}$  cm<sup>3</sup> molecule<sup>-1</sup> s<sup>-1</sup>). The fact that no species involved in the reaction mechanism exhibits major changes in their abundances using this faster rate coefficient shows that this reaction is not a major loss mechanism for NH<sub>2</sub> nor NO and, correspondingly, is not a major production mechanism for H<sub>2</sub>O nor N<sub>2</sub> under molecular cloud conditions. Inspection of the dominant reactions at 10 K and at steady state showed that loss of NH<sub>2</sub> is dominated by reaction with O to form OH or HNO, whereas loss of NO is dominated by reaction with atomic N to form N<sub>2</sub>.

## 5. CONCLUSIONS

The reaction between NH<sub>2</sub> and NO has been studied experimentally over the temperature range 24–106 K using a pulsed laser-photolysis laser-induced fluorescence (PLP-LIF) technique coupled with a Laval nozzle expansion in order to reach the low temperatures of the ISM. The reaction has been shown to have a negative temperature dependence, in agreement with higher temperature literature data over the temperature range 200–2500 K. Ab initio calculations of the PES of the NH<sub>2</sub> + NO system were carried out, with the structures of the stationary points on the PES given in Diao and Smith<sup>29</sup> optimized in order to obtain vibrational frequencies, rotational constants, and electronic and ZPEs. Using this surface, rate theory calculations were carried out using the MESMER software package. Experimentally determined rate coefficients and BRs have been used to fit the calculated PES. Good agreement between the calculated and experimental rate coefficients was obtained over the entire temperature range by simply adjusting the ILT parameters for the initial association reaction of the NH<sub>2</sub> with NO. Fitting to the experimental BRs from the literature required only minor adjustments to the calculated PES to be made, moving the HN<sub>2</sub> + OH exothermicity down by 2.1 kJ mol<sup>-1</sup>, and the TSe and TSc pair (which control the cis–trans isomerization reactions of the HNNOH adducts) down by 4.1 kJ mol<sup>-1</sup>. In this manner, we were able to predict BRs that are largely consistent with experimentally determined literature (within ±0.1) values over a wide range of temperatures. The calculated BRs show a more pronounced sinusoidal shape than is reflected in the experimental data; as such, the largest discrepancies occur at both the high (2500 K) and low (300 K) temperature ends of the temperature range. Extending the temperature range over which we predict BRs down to 60 K, we see that at 180 K and below, the HN<sub>2</sub> + OH endothermic product channel is no longer open, and the reaction proceeds along the low-temperature channel, forming N<sub>2</sub> + H<sub>2</sub>O. Despite the relatively fast reaction rate measured at low temperatures in this work ( $\sim 10^{-10}$  cm<sup>3</sup> s<sup>-1</sup>), inclusion of the new rates in astrochemical models showed that this reaction is not an important loss mechanism for NH<sub>2</sub> nor NO under molecular cloud conditions; however, the impact of the newly measured rate constant should be tested in other low-temperature astrophysical environments in which radical species are abundant, e.g., photon-dominated regions or the outer winds of AGB stars.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jpca.3c03652>.

Comparison of relative energies of stationary points on the NH<sub>2</sub> + NO PES from this and other recent literature studies; full PES for NH<sub>2</sub> + NO; and energies calculated at the B3LYP/6-311G(d,p) level of theory (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Authors

Kevin M. Douglas – School of Chemistry, University of Leeds, Leeds LS2 9JT, U.K.; [orcid.org/0000-0002-3281-3685](https://orcid.org/0000-0002-3281-3685); Email: [k.m.douglas@leeds.ac.uk](mailto:k.m.douglas@leeds.ac.uk)

Dwayne E. Heard – School of Chemistry, University of Leeds, Leeds LS2 9JT, U.K.; [orcid.org/0000-0002-0357-6238](https://orcid.org/0000-0002-0357-6238); Email: [d.e.heard@leeds.ac.uk](mailto:d.e.heard@leeds.ac.uk)

### Authors

Daniel Lucas – School of Chemistry, University of Leeds, Leeds LS2 9JT, U.K.; [orcid.org/0000-0002-1841-2076](https://orcid.org/0000-0002-1841-2076)

Catherine Walsh – School of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, U.K.; [orcid.org/0000-0001-6078-786X](https://orcid.org/0000-0001-6078-786X)

Mark A. Blitz – School of Chemistry, University of Leeds, Leeds LS2 9JT, U.K.; National Centre for Atmospheric Science (NCAS), University of Leeds, Leeds LS2 9JT, U.K.; [orcid.org/0000-0001-6710-4021](https://orcid.org/0000-0001-6710-4021)

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.jpca.3c03652>

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This study was supported by funding from the UK Science and Technology Facilities Council (grant ST/T000287/1). C.W. acknowledges support from the University of Leeds, STFC, and UKRI (grant no. ST/T000287/1, MR/T040726/1). M.A.B. acknowledges funding from the National Centre for Atmospheric Science, which is funded by the Natural Environmental Research Council.

## ■ REFERENCES

- (1) Glarborg, P.; Damjohansen, K.; Miller, J. A.; Kee, R. J.; Coltrin, M. E. Modeling the thermal DENOx process in flow reactors. Surface effects and Nitrous Oxide formation. *Int. J. Chem. Kinet.* **1994**, *26*, 421–436.
- (2) Lyon, R. K. The NH<sub>3</sub>-NO-O<sub>2</sub> reaction. *Int. J. Chem. Kinet.* **1976**, *8*, 315–318.
- (3) Miller, J. A.; Bowman, C. T. Mechanism and modeling of nitrogen chemistry in combustion. *Prog. Energy Combust. Sci.* **1989**, *15*, 287–338.
- (4) Miller, J. A.; Branch, M. C.; Kee, R. J. A chemical kinetic model for the selective reduction of nitric oxide by ammonia. *Combust. Flame* **1981**, *43*, 81–98.
- (5) Miller, J. A.; Glarborg, P. Modeling the Thermal De-NOx Process: Closing in on a final solution. *Int. J. Chem. Kinet.* **1999**, *31*, 757–765.
- (6) Atakan, B.; Jacobs, A.; Wahl, M.; Weller, R.; Wolfrum, J. Kinetic measurements and product branching ratio for the reaction NH<sub>2</sub>+NO at 294–1027 K. *Chem. Phys. Lett.* **1989**, *155*, 609–613.
- (7) Brown, M. J.; Smith, D. B. Aspects of nitrogen flame chemistry revealed by burning velocity modelling. *Symp. (Int.) Combust., [Proc.]* **1994**, *25*, 1011–1018.

- (8) Bulatov, V. P.; Ioffe, A. A.; Lozovsky, V. A.; Sarkisov, O. M. On the reaction of the  $\text{NH}_2$  radical with NO at 295–620 K. *Chem. Phys. Lett.* **1989**, *161*, 141–146.
- (9) Deppe, J.; Friedrichs, G.; Romming, H. J.; Gg Wagner, H. A kinetic study of the reaction of  $\text{NH}_2$  with NO in the temperature range 1400–2800 K. *Phys. Chem. Chem. Phys.* **1999**, *1*, 427–435.
- (10) Diau, E. W.; Yu, T.; Wagner, M. A. G.; Lin, M. C. Kinetics of the  $\text{NH}_2 + \text{NO}$  Reaction: Effects of Temperature on the Total Rate Constant and the OH/ $\text{H}_2\text{O}$  Branching Ratio. *J. Phys. Chem.* **1994**, *98*, 4034–4042.
- (11) Glarborg, P.; Kristensen, P. G.; DamJohansen, K.; Miller, J. A. Branching fraction of the  $\text{NH}_2 + \text{NO}$  reaction between 1210 and 1370 K. *J. Phys. Chem. A* **1997**, *101*, 3741–3745.
- (12) Halbgewachs, M. J.; Diau, E. W. G.; Mebel, A. M.; Lin, M. C.; Melius, C. F. Thermal reduction of NO by  $\text{NH}_3$ : Kinetic modeling of the  $\text{NH}_2 + \text{NO}$  product branching ratio. *Symp. (Int.) Combust., [Proc.]* **1996**, *26*, 2109–2115.
- (13) Hall, J. L.; Zeitz, D.; Stephens, J. W.; Kasper, J. V. V.; Glass, G. P.; Curl, R. F.; Tittel, F. K. Studies of the  $\text{NH}_2 + \text{NO}$  reaction by infrared kinetic spectroscopy. *J. Phys. Chem.* **1986**, *90*, 2501–2505.
- (14) Kimballin, M. A.; Hanson, R. K. Combustion-driven flow reactor studies of thermal DeNOx reaction kinetics. *Combust. Flame* **1986**, *64*, 337–351.
- (15) Park, J.; Lin, M. C. Direct determination of product branching for the  $\text{NH}_2 + \text{NO}$  reaction at temperatures between 302 and 1060 K. *J. Phys. Chem.* **1996**, *100*, 3317–3319.
- (16) Park, J.; Lin, M. C. Laser-Initiated NO Reduction by  $\text{NH}_3$ : Total Rate Constant and Product Branching Ratio Measurements for the  $\text{NH}_2 + \text{NO}$  Reaction. *J. Phys. Chem. A* **1997**, *101*, 5–13.
- (17) Roose, T. R.; Hanson, R. K.; Kruger, C. H. A shock tube study of the decomposition of no in the presence of  $\text{NH}_3$ . *Symp. (Int.) Combust., [Proc.]* **1981**, *18*, 853–862.
- (18) Silver, J. A.; Kolb, C. E. Kinetic measurements for the reaction of amidogen + nitric oxide over the temperature range 294–1215 K. *J. Phys. Chem.* **1982**, *86*, 3240–3246.
- (19) Song, S.; Hanson, R. K.; Bowman, C. T.; Golden, D. M. Shock tube determination of the overall rate of  $\text{NH}_2 + \text{NO} \rightarrow$  products at high temperatures. *Proc. Combust. Inst.* **2000**, *28*, 2403–2409.
- (20) Song, S.; Hanson, R. K.; Bowman, C. T.; Golden, D. M. Shock tube determination of the overall rate of  $\text{NH}_2 + \text{NO} \rightarrow$  products in the thermal De-NOx temperature window. *Int. J. Chem. Kinet.* **2001**, *33*, 715–721.
- (21) Stephens, J. W.; Morter, C. L.; Farhat, S. K.; Glass, G. P.; Curl, R. F. Branching ratio of the reaction  $\text{NH}_2 + \text{NO}$  at elevated temperatures. *J. Phys. Chem.* **1993**, *97*, 8944–8951.
- (22) Vandooren, J.; Bian, J.; Van Tiggelen, P. J. Comparison of experimental and calculated structures of an ammonia-nitric oxide flame. Importance of the  $\text{NH}_2 + \text{NO}$  reaction. *Combust. Flame* **1994**, *98*, 402–410.
- (23) Votsmeier, M.; Song, S.; Hanson, R. K.; Bowman, C. T. A shock tube study of the product branching ratio for the reaction  $\text{NH}_2 + \text{NO}$  using frequency-modulation detection of  $\text{NH}_2$ . *J. Phys. Chem. A* **1999**, *103*, 1566–1571.
- (24) Wolf, M.; Yang, D. L.; Durant, J. L. Kinetic-studies of  $\text{NH}_x$  radical reactions. *J. Photochem. Photobiol., A* **1994**, *80*, 85–93.
- (25) Wolf, M.; Yang, D. L.; Durant, J. L. A Comprehensive Study of the Reaction  $\text{NH}_2 + \text{NO} \rightarrow$  Products: Reaction Rate Coefficients, Product Branching Fractions, and ab Initio Calculations. *J. Phys. Chem. A* **1997**, *101*, 6243–6251.
- (26) Abou-Rachid, H.; Pouchan, C.; Chaillet, M. Ab initio CI study of the reaction between  $\text{NH}_2$  and NO. *Chem. Phys.* **1984**, *90*, 243–255.
- (27) Casewit, C. J.; Goddard, W. A. Energetics and mechanisms for reactions involving nitrosamide, hydroxydiazenes, and diimide N-oxides. *J. Am. Chem. Soc.* **1982**, *104*, 3280–3287.
- (28) Diau, E. W. G.; Smith, S. C. Temperature dependence of rate coefficients and branching ratios for the  $\text{NH}_2 + \text{NO}$  reaction via microcanonical variational transition state theory. *J. Phys. Chem.* **1996**, *100*, 12349–12354.
- (29) Diau, E. W. G.; Smith, S. C. Theoretical investigation of the potential energy surface for the  $\text{NH}_2 + \text{NO}$  reaction via density functional theory and ab initio molecular electronic structure theory. *J. Chem. Phys.* **1997**, *106*, 9236–9251.
- (30) Duan, X.; Page, M. Theoretical characterization of structures and vibrational frequencies for intermediates and transition states in the reaction of  $\text{NH}_2$  with NO. *J. Mol. Struct.: THEOCHEM* **1995**, *333*, 233–242.
- (31) Fang, D. C.; Harding, L. B.; Klippenstein, S. J.; Miller, J. A. A direct transition state theory based analysis of the branching in  $\text{NH}_2 + \text{NO}$ . *Faraday Discuss.* **2001**, *119*, 207–222.
- (32) Harrison, J. A.; Maclagan, R.; Whyte, A. R. Structures, energies, and vibrational frequencies of intermediates and transition-states in the reaction of  $\text{NH}_2$  and NO. *J. Phys. Chem.* **1987**, *91*, 6683–6686.
- (33) Melius, C. F.; Binkley, J. S. Energetics of the reaction pathways for  $\text{NH}_2 + \text{NO} \rightarrow$  products and  $\text{NH} + \text{NO} \rightarrow$  products. *Symp. (Int.) Combust., [Proc.]* **1985**, *20*, 575–583.
- (34) Miller, J. A.; Klippenstein, S. J. Theoretical Considerations in the  $\text{NH}_2 + \text{NO}$  Reaction. *J. Phys. Chem. A* **2000**, *104*, 2061–2069.
- (35) Phillips, L. F. A priori rate-constant for the reaction  $\text{NH}_2 + \text{NO} \rightarrow \text{N}_2 + \text{H}_2\text{O}$ . *Chem. Phys. Lett.* **1987**, *135*, 269–274.
- (36) Walch, S. P. Theoretical characterization of the reaction  $\text{NH}_2 + \text{NO} \rightarrow$  products. *J. Chem. Phys.* **1993**, *99*, 5295–5300.
- (37) Selgren, S. F.; McLoughlin, P. W.; Gellene, G. I. Observation of dissociative and radiative states of  $\text{N}_2\text{H}$  by neutralized ion-beam techniques. *J. Chem. Phys.* **1989**, *90*, 1624–1629.
- (38) Koizumi, H.; Schatz, G. C.; Walch, S. P. A coupled channel study of  $\text{HN}_2$  unimolecular decay based on a global ab initio potential surface. *J. Chem. Phys.* **1991**, *95*, 4130–4135.
- (39) Walch, S. P.; Duchovic, R. J.; Rohlfing, C. M. Theoretical characterization of the minimum energy path for hydrogen-atom addition to  $\text{N}_2$  - implications for the unimolecular lifetime of  $\text{HN}_2$ . *J. Chem. Phys.* **1989**, *90*, 3230–3240.
- (40) Vandishoeck, E. F.; Jansen, D. J.; Schilke, P.; Phillips, T. G. Detection of the interstellar  $\text{NH}_2$  radical. *Astrophys. J.* **1993**, *416*, L83–L86.
- (41) Persson, C. M.; De Luca, M.; Mookerjee, B.; Olofsson, A. O. H.; Black, J. H.; Gerin, M.; Herbst, E.; Bell, T. A.; Coutens, A.; Godard, B.; et al. Nitrogen hydrides in interstellar gas II. Analysis of Herschel/HIFI observations towards W49N and G10.6–0.4 (W31C). *Astron. Astrophys.* **2012**, *543*, 34.
- (42) Liszt, H. S.; Turner, B. E. Microwave detection of interstellar NO. *Astrophys. J., Lett.* **1978**, *224*, L73–L76.
- (43) Codella, C.; Viti, S.; Lefloch, B.; Holdship, J.; Bachiller, R.; Bianchi, E.; Ceccarelli, C.; Favre, C.; Jiménez-Serra, I.; Podio, L.; et al. Nitrogen oxide in protostellar envelopes and shocks: the ASAI survey. *Mon. Not. Astron. Soc.* **2018**, *474*, 5694–5703.
- (44) Halfen, D. T.; Apponi, A. J.; Ziurys, L. M. Evaluating the N/O Chemical Network: The Distribution of  $\text{N}_2\text{O}$  and NO in the Sagittarius B2 Complex. *Astrophys. J.* **2001**, *561*, 244–253.
- (45) Martin, S.; Mauersberger, R.; Martin-Pintado, J.; Garcia-Burillo, S.; Henkel, C. First detections of extragalactic  $\text{SO}_2$ , NS and NO. *Astron. Astrophys.* **2003**, *411*, L465–L468.
- (46) McGonagle, D.; Irvine, W. M.; Minh, Y. C.; Ziurys, L. M. Detection of Nitric Oxide in the Dark Cloud L134N. *Astrophys. J.* **1990**, *359*, 121.
- (47) Ziurys, L. M.; McGonagle, D.; Minh, Y.; Irvine, W. M. Nitric Oxide in Star-forming Regions: Further Evidence for Interstellar N-O Bonds. *Astrophys. J.* **1991**, *373*, 535.
- (48) Wakelam, V.; Herbst, E.; Loison, J. C.; Smith, I. W. M.; Chandrasekaran, V.; Pavone, B.; Adams, N. G.; Bacchus-Montabonel, M. C.; Bergeat, A.; Beroff, K.; et al. A Kinetic Database for Astrochemistry (KIDA). *Astrophys. J., Suppl. Ser.* **2012**, *199*, 21.
- (49) Glowacki, D. R.; Liang, C.-H.; Morley, C.; Pilling, M. J.; Robertson, S. H. MESMER: An Open-Source Master Equation Solver for Multi-Energy Well Reactions. *J. Phys. Chem. A* **2012**, *116*, 9545–9560.

- (50) McElroy, D.; Walsh, C.; Markwick, A. J.; Cordiner, M. A.; Smith, K.; Millar, T. J. The UMIST database for astrochemistry 2012. *Astron. Astrophys.* **2013**, *550*, A36.
- (51) Taylor, S. E.; Goddard, A.; Blitz, M. A.; Cleary, P. A.; Heard, D. E. Pulsed Laval nozzle study of the kinetics of OH with unsaturated hydrocarbons at very low temperatures. *Phys. Chem. Chem. Phys.* **2008**, *10*, 422–437.
- (52) Caravan, R. L.; Shannon, R. J.; Lewis, T.; Blitz, M. A.; Heard, D. E. Measurements of rate coefficients for reactions of OH with ethanol and propan-2-ol at very low temperatures. *J. Phys. Chem. A* **2015**, *119*, 7130–7137.
- (53) Gomez Martin, J. C.; Caravan, R. L.; Blitz, M. A.; Heard, D. E.; Plane, J. M. C. Low Temperature Kinetics of the CH<sub>3</sub>OH + OH Reaction. *J. Phys. Chem. A* **2014**, *118*, 2693–2701.
- (54) Shannon, R. J.; Blitz, M. A.; Goddard, A.; Heard, D. E. Accelerated chemistry in the reaction between the hydroxyl radical and methanol at interstellar temperatures facilitated by tunnelling. *Nat. Chem.* **2013**, *5*, 745–749.
- (55) Shannon, R. J.; Caravan, R. L.; Blitz, M. A.; Heard, D. E. A combined experimental and theoretical study of reactions between the hydroxyl radical and oxygenated hydrocarbons relevant to astrochemical environments. *Phys. Chem. Chem. Phys.* **2014**, *16*, 3466–3478.
- (56) Douglas, K.; Blitz, M. A.; Feng, W. H.; Heard, D. E.; Plane, J. M. C.; Slater, E.; Willacy, K.; Seakins, P. W. Low temperature studies of the removal reactions of 1CH<sub>2</sub> with particular relevance to the atmosphere of Titan. *Icarus* **2018**, *303*, 10–21.
- (57) Douglas, K. M.; Blitz, M. A.; Feng, W.; Heard, D. E.; Plane, J. M. C.; Rashid, H.; Seakins, P. W. Low temperature studies of the rate coefficients and branching ratios of reactive loss vs quenching for the reactions of 1CH<sub>2</sub> with C<sub>2</sub>H<sub>6</sub>, C<sub>2</sub>H<sub>4</sub>, C<sub>2</sub>H<sub>2</sub>. *Icarus* **2019**, *321*, 752–766.
- (58) West, N. A.; Millar, T. J.; Van de Sande, M.; Rutter, E.; Blitz, M. A.; Decin, L.; Heard, D. E. Measurements of Low Temperature Rate Coefficients for the Reaction of CH with CH<sub>2</sub>O and Application to Dark Cloud and AGB Stellar Wind Models. *Astrophys. J.* **2019**, *885*, 134.
- (59) West, N. A.; Li, L. H. D.; Millar, T. J.; Van de Sande, M.; Rutter, E.; Blitz, M. A.; Lehman, J. H.; Decin, L.; Heard, D. E. Experimental and theoretical study of the low-temperature kinetics of the reaction of CN with CH<sub>2</sub>O and implications for interstellar environments. *Phys. Chem. Chem. Phys.* **2023**, *25*, 7719–7733.
- (60) Douglas, K. M.; Lucas, D. I.; Walsh, C.; West, N. A.; Blitz, M. A.; Heard, D. E. The Gas-phase Reaction of NH<sub>2</sub> with Formaldehyde (CH<sub>2</sub>O) is not a Source of Formamide (NH<sub>2</sub>CHO) in Interstellar Environments. *Astrophys. J., Lett.* **2022**, *937*, L16.
- (61) Copeland, R. A.; Crosley, D. R.; Smith, G. P. In Laser-induced fluorescence spectroscopy of NCO and NH<sub>2</sub> in atmospheric pressure flames. *Symposium (International) on Combustion*; Elsevier, 1985; pp 1195–1203.
- (62) Donnelly, V. M.; Baronavski, A. P.; McDonald, J. R. ArF laser photodissociation of NH<sub>3</sub> at 193 nm: internal energy distributions in NH<sub>2</sub> X<sup>2</sup>B<sub>1</sub> and A<sup>2</sup>A<sub>1</sub>, and two-photon generation of NH A 3Π and b 1Σ<sup>+</sup>. *Chem. Phys.* **1979**, *43*, 271–281.
- (63) Yamasaki, K.; Watanabe, A.; Kakuda, T.; Itakura, A.; Fukushima, H.; Endo, M.; Maruyama, C.; Tokue, I. Vibrational energy distributions of NH<sub>2</sub> (X<sup>2</sup>B<sub>1</sub>) fragments generated in the photolysis of NH<sub>3</sub> at 193 nm: Application of kinetic analysis on vibrational cascade. *J. Phys. Chem. A* **2002**, *106*, 7728–7735.
- (64) Yamasaki, K.; Watanabe, A.; Tanaka, A.; Sato, M.; Tokue, I. Kinetics of the reaction NH<sub>2</sub> (X<sup>2</sup>B<sub>1</sub>, ν<sub>2</sub> = 0 and 1) + NO. *J. Phys. Chem. A* **2002**, *106*, 6563–6569.
- (65) Frisch, M.; Trucks, G.; Schlegel, H.; Scuseria, G.; Robb, M.; Cheeseman, J.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G.; et al. *Gaussian 09*. Revision A.02; Gaussian, Inc.: Wallingford CT, 2016.
- (66) Becke, A. D. Density-functional thermochemistry. II. The effect of the Perdew-Wang generalized-gradient correlation correction. *J. Chem. Phys.* **1992**, *97*, 9173–9177.
- (67) Becke, A. D. Density-functional thermochemistry. I. The effect of the exchange-only gradient correction. *J. Chem. Phys.* **1992**, *96*, 2155–2160.
- (68) Becke, A. D. Density-functional thermochemistry. III. The role of exact exchange. *J. Chem. Phys.* **1993**, *98*, 5648–5652.
- (69) Lee, C. T.; Yang, W. T.; Parr, R. G. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron-density. *Phys. Rev. B: Condens. Matter Mater. Phys.* **1988**, *37*, 785–789.
- (70) NIST Computational Chemistry Comparison and Benchmark Database. <http://cccbdb.nist.gov/>. (accessed April, 2023).
- (71) Gericke, K. H.; Torres, L. M.; Guillory, W. A. State selected removal of vibrationally excited NH<sub>2</sub> [X<sup>2</sup>B<sub>1</sub>(0,ν<sub>2</sub>,0)] radicals. *J. Chem. Phys.* **1984**, *80*, 6134–6140.
- (72) Xiang, T.-X.; Gericke, K.-H.; Torres, L. M.; Guillory, W. A. Vibrational relaxation of NH<sub>2</sub> [X<sup>2</sup>B<sub>1</sub>(0,ν<sub>2</sub>,0)] radicals. *Chem. Phys.* **1986**, *101*, 157–163.
- (73) Bocherel, P.; Herbert, L. B.; Rowe, B. R.; Sims, I. R.; Smith, I. W. M.; Travers, D. Ultralow-Temperature Kinetics of CH(X<sup>2</sup>Π) Reactions: Rate Coefficients for Reactions with O<sub>2</sub> and NO (T = 13–708 K), and with NH<sub>3</sub> (T = 23–295 K). *J. Phys. Chem.* **1996**, *100*, 3063–3069.
- (74) Nunez-Reyes, D.; Hickson, K. M. The reactivity of C(1D) with oxygen bearing molecules NO and O<sub>2</sub> at low temperature. *Chem. Phys. Lett.* **2017**, *687*, 330–335.
- (75) Nunez-Reyes, D.; Hickson, K. M. A low temperature investigation of the gas-phase N(<sup>2</sup>D) + NO reaction. Towards a viable source of N(<sup>2</sup>D) atoms for kinetic studies in astrochemistry. *Phys. Chem. Chem. Phys.* **2018**, *20*, 17442–17447.
- (76) Tsubouchi, M.; de Lange, C. A.; Suzuki, T. Femtosecond time-resolved charged particle imaging studies of the ultraviolet photodissociation of the NO dimer. *J. Chem. Phys.* **2003**, *119*, 11728–11739.
- (77) Wade, E. A.; Cline, J. I.; Lorenz, K. T.; Hayden, C.; Chandler, D. W. Direct measurement of the binding energy of the NO dimer. *J. Chem. Phys.* **2002**, *116*, 4755–4757.
- (78) Ivanic, J.; Schmidt, M. W.; Luke, B. High-level theoretical study of the NO dimer and tetramer: Has the tetramer been observed? *J. Chem. Phys.* **2012**, *137*, 214316.
- (79) Mebel, A. M.; Morokuma, K.; Lin, M. C. Modification of the Gaussian-2 theoretical-model - the use of coupled-cluster energies, density-functional geometries, and frequencies. *J. Chem. Phys.* **1995**, *103*, 7414–7421.
- (80) Walsh, C.; Nomura, H.; van Dishoeck, E. The molecular composition of the planet-forming regions of protoplanetary disks across the luminosity regime. *Astron. Astrophys.* **2015**, *582*, A88.