



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/203466/>

Version: Published Version

Article:

Xu, Qingqing, Tang, Qingyun, Xu, Yang et al. (2023) Biotechnology in Future Food Lipids: Opportunities and Challenges. Annual Review of Food Science and Technology. pp. 225-246. ISSN: 1941-1421

<https://doi.org/10.1146/annurev-food-060721-024353>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Annual Review of Food Science and Technology
**Biotechnology in Future Food
 Lipids: Opportunities and
 Challenges**

Qingqing Xu,^{1,*} Qingyun Tang,^{2,*} Yang Xu,³
 Junjun Wu,⁴ Xiangzhao Mao,⁵ Fuli Li,⁶
 Shian Wang,⁶ and Yonghua Wang^{1,7}

¹School of Food Science and Engineering, South China University of Technology, Guangzhou, Guangdong, China; email: yonghw@scut.edu.cn

²Department of Chemistry, University of York, Heslington, York, United Kingdom

³Department of Molecular and Cellular Biology, University of Guelph, Guelph, Ontario, Canada

⁴College of Food Science and Technology, Nanjing Agricultural University, Nanjing, Jiangsu, China

⁵College of Food Science and Engineering, Ocean University of China, Qingdao, Shandong, China

⁶Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, Qingdao, China

⁷Guangdong Youmei Institute of Intelligent Bio-Manufacturing, Foshan, Guangdong, China

ANNUAL
REVIEWS **CONNECT**

www.annualreviews.org

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

Annu. Rev. Food Sci. Technol. 2023. 14:225–46

First published as a Review in Advance on
 January 25, 2023

The *Annual Review of Food Science and Technology* is
 online at food.annualreviews.org

<https://doi.org/10.1146/annurev-food-060721-024353>

Copyright © 2023 by the author(s). This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See credit lines of images or other third-party material in this article for license information.

*These authors contributed equally to this work.

Keywords

microbial lipids, metabolic engineering, enzymatic modification, structural lipids, phospholipids

Abstract

Lipids are a large group of essential nutrients in daily diets that provide energy and maintain various physiological functions. As the global population is rapidly expanding, there is an urgent need to enhance the production and quality of food lipids. The development of modern biotechnology allows the manipulation of oil production in plants and microorganisms and the improvement of the nutritional value of food lipids. Various metabolic engineering strategies have been exploited to increase oil production and produce value-added oils in traditional oil crops and other novel lipid sources (e.g., plant vegetative tissues, microalgae, and oleaginous microorganisms). Furthermore, natural lipid structures can be modified by lipases to prepare functional lipids, e.g., diacylglycerols, medium–long–medium-type structured triacylglycerols, human milk-fat substitutes, and structural



phospholipids, for specific nutritional demands. In this review, we focus on the recent advances in metabolic engineering of lipid production in plants and microorganisms, and the preparation of functional lipids via biocatalysis.

1. INTRODUCTION

In light of an ever-growing population but increasingly limited natural resources, a major challenge we are facing today is how to feed the world (Hamrick & Chen 2021, Lv et al. 2021). Nearly a billion people all around the world still suffer from hunger or malnourishment (Ghazani & Marangoni 2022), and the demand for food is expected to double over the coming decades because of the ballooning population (Vanhercke et al. 2019b). Lipids are a large group of macronutrients in our diets that provide energy and maintain various physiological functions in our bodies, such as energy storage, cellular signaling, and regulation of the cellular membrane (Ghazani & Marangoni 2022). Insufficient lipid intake can lead to a deficiency in essential fatty acids and fat-soluble vitamins, which is associated with a variety of human pathologies and disorders, including immune dysfunction, cardiovascular disease, vision problems, retinopathy, and inflammation (Shahidi et al. 2021, Wu et al. 2022). Plants and animals are the traditional sources of food lipids, but the current production level barely meets the growing demand of the world. Considering the increasing competition for land, water, and other resources (Godfray et al. 2010), there is a need to develop alternative dietary lipid sources to keep the human food system sustainable while maintaining safety and nutrition (Lv et al. 2021).

Novel lipid sources such as plant vegetative tissues and oleaginous microorganisms (mainly yeast, fungi, microalgae, and bacteria) hold great promise in supporting and complementing the traditional food lipid supply from plants and animals. In plants, more than 60% of the oils are derived from seeds, whereas other tissues, such as plant vegetative tissues, do not normally accumulate oils. Owing to the high biomass in plant vegetative tissues, there is a strong interest in improving oil production from them. Moreover, oleaginous microbes are excellent hosts for lipids and are able to accumulate large quantities of lipids (20–90% of dry cell weight) (Uprety et al. 2022) and assemble the desirable fatty acids into lipid molecules (Kamineni & Shaw 2020). However, the commercially desirable lipids in microbial oils, i.e., those rich in polyunsaturated fatty acids (PUFAs), are often present at low levels and generally mix with lipids of low commercial interest (Uprety et al. 2022). Therefore, different metabolic engineering strategies have been exploited in plants, microalgae, and other microorganisms to improve oil production and produce value-added oils for food purposes (**Figure 1**).

At the other extreme, the world is facing a growing epidemic of obesity and related disorders. High consumption of triacylglycerols (TAGs) with low nutritional value has been a huge trigger for these problems (Ando et al. 2017). Recent findings show that functional lipids such as diacylglycerols (DAGs), structured TAGs (STs), and structured phospholipids (SPLs) can offer better nutritional or health value. Nevertheless, these desirable components are present at low levels in natural lipids and the development of preparation methods is needed. Chemical and enzymatic methods are commonly used for their preparation, but chemical methods are less accepted by consumers when compared with enzymatic methods, along with excessive by-products. Therefore, enzymatic modification of the composition and/or position of fatty acids in lipids represents an effective approach to meeting special nutritional demands (**Figure 1**).

This manuscript reviews advances in the metabolic engineering of plants, microalgae, and microorganisms for improving food lipid production and value and highlights recent progress made in developing functional lipids using enzymatic modification approaches.




Figure 1

Biotechnology in the development of future food lipids. Metabolic engineering of oleaginous plants, microalgae, fungi, yeast, and bacteria can increase lipid production and/or value. Enzymatic modification of lipids can further produce structured lipids such as the human milk-fat substitute for infant formulas, diacylglycerols and medium-long-medium-type structured lipids to control obesity, and PUFA-rich phospholipids for the prevention of cardiovascular diseases. Abbreviations: FA, fatty acids; L, long-chain fatty acids; M, medium-chain fatty acids; O, oleic acid; P, palmitic acid; PG, phosphate group; PUFA, polyunsaturated fatty acids. Figure adapted from images created using BioRender.com and stock.adobe.com.

2. METABOLIC ENGINEERING TO INCREASE LIPID PRODUCTION AND VALUE IN PLANTS

In plant cells, TAG formation is highly compartmentalized and involves the coordinated action of multiple pathways in different subcellular compartments. In brief, TAG biosynthesis starts with *de novo* fatty acid biosynthesis in the plastids, which are then exported to the cytosol and incorporated into TAGs in the endoplasmic reticulum (ER). The assembled TAGs accumulate between the leaflets of the ER membrane and eventually pinch off the ER to deposit lipid droplets in the cytosol. The lipid droplets contain a TAG core surrounded by a monolayer of phospholipids (PLs) with various proteins embedded (Pyc et al. 2017). Those lipid droplet membrane-associated proteins are important in the formation and turnover of lipid droplets (Pyc et al. 2017). Given the fact that TAG formation requires multiple metabolic modules, various metabolic engineering strategies have been explored to enhance TAG production at different metabolic levels, including increasing fatty acid biosynthesis (Push), enhancing TAG assembly (Pull), packaging TAGs into lipid droplets (Package), and preventing TAG turnover (Protect). These metabolic engineering strategies have been exploited individually or in combination to manipulate oil production in oil crops, plant vegetative tissues, and microalgae (Figure 2).

2.1. Engineering Oil Production in Oil Crops

Many genes have been considered key targets for manipulating seed oil production by pushing carbon flux for *de novo* fatty acid biosynthesis, such as *Wrinkled 1 (WRI1)* and *acetyl-CoA*




Figure 2

Metabolic engineering strategies to modify triacylglycerol (TAG) production in plants and microalgae, including increasing fatty acid biosynthesis (Push), enhancing TAG assembly (Pull), packaging TAGs into lipid droplets (Package), and preventing TAG turnover (Protect). Enzymatic reactions are shown by solid black arrows, substrate diffusion/transport is shown by dashed black arrows, transcriptional regulation is shown by dashed red arrows, and acyl modification is shown by solid brown arrows. Abbreviations: ACP, acyl carrier protein; ACCase, acetyl-CoA carboxylase; CoA, coenzyme A; DAG, diacylglycerol; DGAT, diacylglycerol acyltransferase; ER, endoplasmic reticulum; FFA, free fatty acid; LPA, lysophosphatidic acid; PA, phosphatidic acid; PC, phosphatidylcholine; *WRI1*, *Wrinkled 1*.

carboxylase (*ACCase*), and pulling fatty acids into TAGs, such as *DAG acyltransferase* (*DGAT*). *WRI1* is a master transcription factor that regulates the expression of genes involved in fatty acid biosynthesis and glycolysis (Baud et al. 2009, Ruuska et al. 2002). *WRI1* was first identified in the model plant *Arabidopsis thaliana*, in which disruption of *WRI1* led to an 80% reduction in seed oil content (Focks & Benning 1998). Later, *WRI1* orthologs have also been identified in many crop species, which show a similar regulatory role in oil biosynthesis, including in soybean (*Glycine max*) (Chen et al. 2020), canola (*Brassica napus*) (Liu et al. 2010), and maize (*Zea mays*) (Shen et al. 2010). Overexpression of *WRI1* led to increased TAG content in seeds and vegetative tissues in various plant species (Cernac & Benning 2004, Chen et al. 2020, Q. Li et al. 2015, Liu et al. 2010, Sanjaya et al. 2011, Shen et al. 2010). For example, overexpression of maize *WRI1* increased seed oil content by up to 46% in maize without affecting seed germination, grain yield, or seedling growth (Shen et al. 2010). In soybean, soybean *WRI1* overexpressing lines showed up to ~15% increases in seed oil content and improved agronomic traits such as increased seed number per plant, leading to a 33–53% increase in total seed oil production per plant (Guo et al. 2020). These findings highlight *WRI1* as a promising target for improving oil production in crops.

ACCase catalyzes the carboxylation of acetyl-CoA to malonyl-CoA, which is the first committed step of plastid fatty acid synthesis (Salie & Thelen 2016). In most plant plastids, *ACCase* is a heteromeric form consisting of four subunits: biotin carboxylase, biotin carboxyl carrier protein (BCCP), and α - and β -carboxyltransferases (CTs). A homomeric form of *ACCase*, whereby all four enzymatic components are in a single polypeptide, is present in the cytosol of plants for fatty acid elongation. *ACCase* is considered the rate-limiting step in *de novo* fatty acid synthesis and has been subjected to modifications with the aim of improving fatty acid biosynthesis. Enhancing *ACCase* activity by targeting an *Arabidopsis* homomeric *ACCase* to the plastids boosted the oil

content by 5% in canola (Roesler et al. 1997). In *Arabidopsis*, α -CT and β -CT subunits are physically associated, possibly forming heterotetramers ($\alpha\beta_2$), but α -CT is threefold to tenfold less abundant than β -CT and appears to be the limiting subunit in the ACCase complex (Ke et al. 2000, M.M. Wang et al. 2022). Overexpression of pea α -CT resulted in the successful formation of active α/β -CT subunits and increased ACCase activity, eventually leading to an 8–15% increase in oil production in *Arabidopsis* and camelina seeds (M.M. Wang et al. 2022). Recently, three small plastid proteins of the envelope membrane, CT interactors, were found to interact with α -CT to mediate the docking of ACCase to the plastid envelope membrane, thereby attenuating fatty acid biosynthesis. CT interactor knockout lines showed a 22% increase in oil content in *Arabidopsis* leaves (Ye et al. 2020). Biotin/lipoyl attachment domain-containing (BADC) proteins are negative regulators to ACCase, which are BCCP analogs that interact with BCCP but are not biotinylated and thus compete with BCCP to bind with other ACCase subunits (Salie et al. 2016). Downregulation of BADCs increased the seed oil content by up to 32% (Keereetaweep et al. 2018).

DGAT catalyzes the last and committed step of acyl-CoA-dependent TAG assembly by transferring acyl-CoA to the *sn*-3 position of 1,2-DAG to yield TAG, which appears to play a critical role in determining the flux of carbon into oils (Xu et al. 2018). Overexpression of canola *DGAT1* in canola increased the seed oil content by up to ~14% and ~8% (a 3.5% increase in absolute oil content) in greenhouse and field conditions, respectively (Taylor et al. 2009, Weselake et al. 2008). In addition to native DGAT, high-performance DGAT variants represent promising candidates for manipulating oil production. A high-oil DGAT allele was identified in maize, which contains a phenylalanine insertion at position 469 and is responsible for the increased enzyme activity and oil content (Zheng et al. 2008). Overexpression of the high-oil *DGAT* allele increased maize seed oil content by up to 41% (Zheng et al. 2008). Directed evolution has been used to engineer the enzyme performance of canola and soybean *DGAT1* (Roesler et al. 2016, Siloto et al. 2009), and overexpression of performance-enhanced soybean *DGAT1* variants increased soybean seed oil content by up 16% in field trials (Roesler et al. 2016).

Although manipulating single gene expression has significantly enhanced the seed oil content, combinational metabolic engineering by stacking genes involved in different lipid biosynthetic pathways may hold more promise. Gene stacking has proved to be more effective for improving oil production in plant vegetative tissues (see Section 2.2), but only a few reports have focused on seed oils. For example, the combined expression of *WRI1* and *DGAT1* in *Arabidopsis* led to a 16% increase in the seed oil content, which further increased by 20% when combined with the suppression of *Sugar Dependent Lipase 1 (SDPI)*, encoding a lipase for TAG turnover (van Erp et al. 2014). In some cases, however, lipid gene manipulation had only limited or no success. Co-expression of *Arabidopsis WRI1* and *DGAT1* in soybean did not increase the seed oil content; further transcriptomic and metabolomic analyses revealed lipid droplet packaging and fatty acid biosynthesis may be inhibited, whereas TAG turnover and starch/polysaccharide biosynthesis may be enhanced (Arias et al. 2022). These results demonstrate the complexity of lipid metabolism and its interplay with other competing pathways. Indeed, mounting evidence shows that *WRI1*, *DGAT1*, and ACCase are regulated at the post-translational level, and various regulators, interactors, and metabolites can affect their stability, activity, and performance in oil biosynthesis (Caldo et al. 2017, Kong et al. 2020, Ma et al. 2015, Salie et al. 2016).

2.2. Engineering Oil Production in Plant Vegetative Tissues and Microalgae

In addition to engineering oil production of existing oil crops, substantial research has been carried out to develop novel sources of oils such as plant vegetative tissues and microalgae, with the aim to fulfill the potential of oil production and secure a growing supply of vegetable oils.

Vegetative tissues such as leaves and stems make up the majority of the plant biomass and represent a novel platform for oil production (Vanhercke et al. 2019a). Tobacco plants (*Nicotiana tabacum* and *Nicotiana benthamiana*) can produce a high amount of biomass and have been commonly used in engineering oil production. Leaves typically contain no significant levels of oil (<0.5% on a dry weight basis) but metabolic engineering efforts have achieved up to 15–33% (dry weight) of oil accumulation in engineered tobacco leaves so far, highlighting the potential to outperform current oil crops (Vanhercke et al. 2014, 2017). The integrated Push–Pull–Package–Protect metabolic engineering strategy has been found to be more effective than an individual Push, Pull, Package, or Protect strategy at producing high concentrations of oils in vegetative tissues by mimicking metabolic fluxes in oilseeds, where carbon flux is efficiently channeled into TAGs for storage (Vanhercke et al. 2019a). For example, the combined overexpression of *WRI1* (Push), *DGAT1* (Pull), and the lipid droplet protein *OLEOSIN* (Package) yielded up to 15% (dry weight) of oil in engineered tobacco leaf tissues (Vanhercke et al. 2014). Furthermore, silencing *SDPI* (Protect) in this high-oil tobacco background elevated the oil accumulation by up to 33% (Vanhercke et al. 2017). These Push–Pull–Package–Protect strategies have also been used to increase oil content in the vegetative tissues of crops such as sugarcane and potato, where up to 1.9% and 3.3% (dry weight) of oil were achieved in sugarcane leaves and potato tubers, respectively (Hofvander et al. 2016, Liu et al. 2017, Zale et al. 2016). Microalgae hold promise as a sustainable resource for lipid production because of their high lipid accumulation ability and phototrophic growth rate and the fact that they do not need to compete with agricultural crops for arable land (Hu et al. 2008). But microalgal oil production faces many challenges and is still not economically viable (Sun et al. 2019). Thus, similar metabolic engineering approaches have been exploited to manipulate oil production in various microalgal species, such as *Nannochloropsis* spp. and *Pheodactylum tricornerutum* (Hamilton et al. 2014, Liu et al. 2022, Radakovits et al. 2011, Wang et al. 2021, Xu 2022, Xue et al. 2015).

2.3. Producing Value-Added Oils in Plants and Microalgae

The property and value of oils in food, nutrition, or the oleochemical industry are largely determined by the composition of fatty acids, which can be classified based on carbon chain length (e.g., long chain, >C16; short chain, <C8) and saturation degree (saturated, no double bonds; monounsaturated, 1 double bond; polyunsaturated, >2 double bonds). For example, PUFAs such as eicosapentaenoic acid (EPA; 20:5 $\Delta^{5,8,11,14,17}$ *cis*) and docosahexaenoic acid (DHA; 22:6 $\Delta^{4,7,10,13,16,19}$ *cis*) are well recognized for their health-promoting benefits and have been widely used in nutritional supplements (Boyd et al. 2021, Keaney & Rosen 2019); conjugated linolenic acids (18 carbon fatty acids bearing three conjugated double bonds) such as punicic acid (18:3 Δ^{9} *cis,11**trans,13**cis*) have been shown to display various beneficial bioactivities (Holic et al. 2018). Owing to growing demand and scarce supply, metabolic engineering to produce these unusual functional fatty acids in plants and microalgae has attracted substantial attention.

To date, considerable achievements have been made to produce PUFAs such as EPA and DHA in crops by introducing alternate fatty acid desaturases and elongases from the aerobic PUFA metabolic pathways (Napier et al. 2019). By assembling multiple *desaturase* and *elongase* genes, up to 24% EPA and 12% DHA have been achieved in seed oils of transgenic *Camelina sativa*, an emerging oilseed crop (Petrie et al. 2014, Ruiz-Lopez et al. 2014). In field-grown transgenic *C. sativa*, up to 16% EPA and 9% EPA plus 5–10% DHA were accumulated in seed oils, respectively (Han et al. 2020, Usher et al. 2017). The oilseed crop canola has also been engineered to produce considerable levels of EPA and DHA (Petrie et al. 2020). Recently, two different transgenic canola lines, which were developed by BASF and Cargill and Nuseed, CSIRO, and GRDC, that accumulate up to 12% EPA and/or DHA have been approved for commercial growth in

the United States (Napier et al. 2019). Similarly, EPA and DHA production in microalgae *Nannochloropsis oceanica* and *P. tricornutum* have been optimized by modulating the PUFA metabolic pathway and the TAG assembly enzyme DGAT (Hamilton et al. 2014; Haslam et al. 2020; Liu et al. 2022; Xin et al. 2017, 2019; Xu 2022).

As for conjugated fatty acids, modest success has been achieved in crops and plant vegetative tissues in terms of their production using metabolic engineering. The creation of conjugated double bonds stems from the catalytic action of fatty acid conjugases (FADXs), a divergent form of fatty acid desaturase 2 (FAD2) (Holic et al. 2018). By overexpressing a pomegranate FADX in an *Arabidopsis fad3fae1* mutant and canola-type *B. napus* (with *elongase* knocked out, equivalent to the *fae1* mutation in *Arabidopsis*), punicic acid accumulated up to 11.5% and 11% in the seed oils (Mietkiewska et al. 2014, Xu et al. 2020). The punicic acid content in transgenic *Arabidopsis* seeds increased up to 21% and 24.8% when FAD2 and FAD2 plus DGAT2, respectively, from pomegranate were coexpressed with FADX (Mietkiewska et al. 2014, Weselake & Mietkiewska 2014). Similarly, coexpression of tung tree FADX with DGAT2 showed a synergistic effect and resulted in up to 12% of α -eleostearic acid (18:3 $\Delta^{9cis,11trans,13trans}$) production in *Arabidopsis* leaves (Yurchenko et al. 2017). The level of conjugated fatty acids in transgenic plants, however, has been much lower than that of plants that naturally produce them, which is likely due to the inefficient trafficking of conjugated fatty acids from membrane lipids to storage TAG (Holic et al. 2018). Therefore, further identifying unusual fatty acid selective acyltransferases and introducing them together with other important enzymes may be necessary for alleviating the metabolic bottleneck of unusual fatty acid production. Indeed, significant levels of hydroxy fatty acid were achieved in transgenic *Arabidopsis* seed oils by the coexpression of specialized acyltransferases that selectively incorporate hydroxy fatty acid at each stereochemical position of TAG (Lunn et al. 2019).

3. METABOLIC ENGINEERING TO INCREASE LIPID PRODUCTION IN MICROORGANISMS

An overview of anabolism and catabolism in eukaryotic microorganisms is shown in **Figure 3**. Maximizing flux through lipid biosynthesis can be achieved by combining general metabolic engineering strategies: (a) blocking pathways that compete with synthesis of key precursors or consume oleochemical products, (b) pushing flux into lipid biosynthesis by overexpressing rate-limiting enzymes or enzymes that produce intermediates dedicated to lipid biosynthesis, (c) pulling flux through lipid biosynthesis by overexpressing enzymes or regulatory proteins that bypass native regulation, and (d) expressing auxiliary proteins that alleviate stresses and other problems caused by producing specific oleochemicals.

3.1. Increasing the Acetyl-CoA Pool

Acetyl-CoA is the key building block of all oleochemicals and a central metabolite involved in many biochemical systems. The synthesis of acetyl-CoA is highly regulated and competes with the production of several fermentation products such as acetate, ethanol, and lactate. In *Escherichia coli*, deleting such competing pathways is crucial for directing flux via the reversed β -oxidation pathway (Cintolesi et al. 2014; Clomburg et al. 2012, 2015; Dellomonaco et al. 2011; Wu et al. 2020). However, such procedures lead to diminished cell growth rates or substrate uptake rates. For instance, disrupting three pyruvate decarboxylase isozymes eliminates ethanol production in *Saccharomyces cerevisiae* (Dai et al. 2018). Furthermore, these pyruvate decarboxylase mutants exhibit slow growth rates on glucose and accumulate mutations that lead to reduced glucose uptake. To overcome these challenges, a chimeric ATP:citrate lyase–malic enzyme–malate dehydrogenase–citrate transporter Ctp1 was overexpressed and a 20% increase in fatty acid production was observed. Deleting




Figure 3

The native and designed synthetic pathways involved in lipid biosynthesis in eukaryotic oleaginous microorganisms. The solid arrows denote a single step reaction and the dashed arrows denote the process containing more than one biochemical reaction. Abbreviations: 6PG, 6-phosphogluconate; 6PGL, 6-phosphogluconolactone; ACC, acetyl-CoA carboxylase; ACL, ATP citrate lyase; ACP, acyl carrier protein; DHAP, dihydroxyacetone phosphate; E4P, erythrose 4-phosphate; ECR, *trans*-2,3-enoyl-CoA reductase; ER, endoplasmic reticulum; F6P, fructose 6-phosphate; FAS, fatty acid synthase; FBP, fructose-bisphosphatase 1; FFA, free fatty acids; G6P, glucose 6-phosphate; G3P, glyceraldehyde 3-phosphate; HCD, 3-hydroxyacyl-CoA dehydratase; KCR, 3-ketoacyl-CoA reductase; KCS, 3-ketoacyl-CoA synthase; MtFAS, mitochondria fatty acid synthase; MUFA, monounsaturated fatty acid; PalCoA, palmitoyl coenzyme A; PPP, pentose phosphate pathway; PUFA, polyunsaturated fatty acid; R5P, ribose 5-phosphate; Ru5P, ribulose 5-phosphate; SCD, stearoyl-CoA desaturase; Se7P, sedoheptulose 7-phosphate; TAG, triacylglycerol; TCA, tricarboxylic acid cycle; Xu5P, xylulose 5-phosphate. Figure adapted with permission from Xu et al. (2016).

acyl-CoA synthases *FAA1* and *FAA4* and acyl-CoA oxidase *HFD1* and the overexpression of *Rhodospiridium toruloides* *FAS*, *TesA'*, and *ACC1* produced 10.4 g/L free fatty acids (FFAs) in fed-batch fermentation (Zhou et al. 2016). Another study demonstrated that combining adaptive laboratory evolution and metabolic engineering transformed a host strain into a Crabtree-negative yeast (Dai et al. 2018). Further experiments resulted in a strain capable of producing 33.4 g/L FFAs, exhibiting a fourfold improvement over the initial strain in glucose-limited and nitrogen-restricted fed-batch cultivation (Yu et al. 2018). Increasing the acetyl-CoA pool has been shown to be beneficial for increasing lipid production in both nonoleaginous and oleaginous organisms.

3.2. Increasing the Malonyl-CoA Pool

Malonyl-CoA, the elongation unit used in fatty acid biosynthesis, is produced from ACCase, a rate-limiting reaction in fatty acid biosynthesis in several organisms. The production of FFAs was increased via a balanced overexpression of ACCase subunits associated with the coexpression of a thioesterase in *E. coli* (Lennen et al. 2010). In yeast, several strategies were involved to

improve malonyl-CoA production, including (a) replacing the native promoter of ACC1 with a strong constitutive promoter (Qiao et al. 2015), (b) overexpressing a mutated ACC1 (Ser659Ala, Ser1157Ala) that abolishes post-translational phosphorylation inhibition (d’Espaux et al. 2017), (c) bypassing the ACC pathway through the expression of phosphoenolpyruvate carboxylase and methyl malonyl-CoA CT (Shin & Lee 2017), and (d) resolving an allosteric feedback inhibition due to C16–C20 saturated acyl-CoAs via the overexpression of $\Delta 9$ stearoyl-CoA desaturase in *Yarrowia lipolytica* (Qiao et al. 2015) or the overexpression of a $\Delta 9$ -desaturase in *S. cerevisiae* to improve membrane fluidity and the fatty acyl-CoA pool (d’Espaux et al. 2017). Furthermore, it was found that a mutant Mga2p regulator in *Y. lipolytica* resulted in elevated unsaturated fatty acid biosynthesis and lipid accumulation, which might be due to the reduced feedback inhibition of ACC (Liu et al. 2015). Indirect malonyl-CoA biosensors have also been created in *E. coli*, *Pseudomonas putida*, and *Corynebacterium glutamicum* by employing RppA, a type III polyketide synthase that may generate flaviolin, a red-colored pigment, from malonyl-CoA. A genome-wide sRNA knockdown library was also used together with this sensor to identify high malonyl-CoA producers (Yan et al. 2018).

3.3. Leveraging β -Oxidation as a High-Yielding Synthetic Pathway

β -oxidation enzymes may be used in a synthetic direction (r-BOX) to produce medium- and long-chain oleochemicals via acyl-CoAs. The advantage of this pathway is the larger theoretical yields that can be achieved by avoiding ATP consumption required for malonyl-CoA synthesis. The r-BOX is also ideally suited for producing fatty alcohols as this biosynthetic pathway is redox balanced to glucose catabolism. This β -oxidation pathway was first demonstrated by Dellomonaco et al. (2011), and several groups have employed the r-BOX pathway to increase oleochemical biosynthesis in bacteria or yeast. One relevant study found that *E. coli* type II FAS could be employed to make up the pathway (Clomburg et al. 2018). Another study found that the relative expression of essential component enzymes was beneficial for generating medium-chain TAGs (MCTs) (Wu et al. 2017b). The same group further showed that improving NADH availability is important to increasing medium-chain fatty acid production derived from r-BOX in *E. coli* (Wu et al. 2017a). The r-BOX pathway has also been assembled in the cytosol of *S. cerevisiae* to produce butanol, MCTs, and ethyl esters (Mehrer et al. 2018). One main hurdle of assembling r-BOX in yeast is the compartmentalization of r-BOX in the peroxisome away from pyruvate dehydrogenase in the cytosol. The key enzyme in the r-BOX pathway is the reversible thiolase, which elongates the acyl-chains. Specific thiolase variants have different substrate preferences and help determine the ultimate product chain-length. The original description of the r-BOX pathway employs a short-chain acyl-CoA transferase, YqeF, resulting in the accumulation of C4 acyl-CoA and butanol (Clomburg et al. 2012), whereas *Ralstonia eutropha* BktB has been widely employed in r-BOX supporting synthesis of acyl-CoA of up to 10 carbons (Clomburg et al. 2015, Wu et al. 2020).

3.4. Safety of Oleaginous Microbes

The safety of food microorganisms has been strongly highlighted by several global organizations. The United States Pharmacopeia convention develops and publishes standards for food ingredients in the Food Chemicals Codex (Reynolds 1964). The European Food Feed Cultures Association defined food cultures as “safe live bacteria, yeasts or molds used in food production which are in themselves a characteristic food ingredient” (Bourdichon et al. 2018). The Qualified Presumption of Safety (QPS) was developed to provide a generic safety evaluation for biological agents in Europe (Koutsoumanis et al. 2021). Generally recognized as safe (GRAS) rules are used for US Food and Drug Administration regulatory approval (Singh & Gaur 2021). The Chinese

government publishes a microbe list for food usage and allows the usage of those microorganisms in traditional foods (Yao et al. 2022).

Oleaginous microbes are preferred for use in the production of food lipids, especially those in the safe lists published by governments. A couple of oleaginous microorganisms have been characterized as GRAS and QPS status, such as the oleaginous yeasts *Y. lipolytica* and *Xanthophyllomyces dendrorhous* (Koutsoumanis et al. 2021, Nicaud 2012) and the oleaginous fungi *Schizochytrium limacinum* and *Aspergillus oryzae* (Borowitzka 2013, Rousta et al. 2021). The qualification of these microbes is described in the official documents that provide guidance to not only producers but also researchers.

4. BIOCATALYTIC APPROACHES FOR LIPID MODIFICATION

Functional lipids are important food constituents for meeting desired nutritional demands. For example, overconsumption of TAGs, which are the major source of energy within the human diet, has been related to obesity and various health problems (Ando et al. 2017). Replacing TAGs with DAGs can efficiently suppress the accumulation of body fat and help reduce body weight (Prabhavathi Devi et al. 2018). Medium- and long-chain fatty acid TAGs (MLCTs) can reduce calories in lipids and reduce fat accumulation (Nagao & Yanagita 2010). Other STs, for example, *sn*-2 palmitate TAGs, are the main components in human milk-fat substitutes for infant formulas and can meet specific nutritional or functional demands (Wei et al. 2019). Incorporation of PUFAs into STs or SPLs increases their availability and metabolic absorption (Ang et al. 2019).

Lipid modification can be achieved chemically or enzymatically. Enzymatic methods provide superior chemo- and regioselectivity and are more environmentally friendly than chemical methods. Lipases (TAG hydrolases E.C. 3.1.1.3) are the most commonly used enzymes for lipid modification and can catalyze hydrolysis, esterification, and transesterification reactions of lipids. Many of them are commercially available and display regioselectivity to either the *sn*-1,3 or *sn*-2 position of TAGs and have different chain-length preferences. With new developments in protein engineering and bioinformatics, enzymatic functions can also be adjusted to meet different modification purposes.

4.1. Production of Diacylglycerols

DAGs are natural components in edible oils, with levels lower than 10% in natural lipids, and consist mainly of 1,2-DAGs and 1,3-DAGs (3:7). In an era when obesity has become an intractable problem worldwide, DAGs have attracted widespread attention due to their specific physiological functions. The different metabolic pathways of TAGs and 1,3-DAGs (the main component in DAGs) contribute to health-beneficial properties (Prabhavathi Devi et al. 2018). DAGs have, for example, been found to have obviously positive effects in regulating postprandial lipids, relieving type 2 diabetes, and suppressing the accumulation of visceral fat (Li et al. 2019). However, DAGs cannot exert their beneficial effects unless their content is above 27.3% in the edible lipids (Saito et al. 2010). Generally, higher levels of DAGs showed better beneficial effects (Saito et al. 2010); thus, it is necessary to maximize the DAGs content in edible oils by enzymatic methods.

Enzymatic production of DAGs is generally performed in three ways: (a) partial hydrolysis of TAGs with water to produce DAGs, monoacylglycerols (MAGs), and FFAs; (b) glycerolysis reaction between TAGs and glycerol yielding DAGs; and (c) esterification of FFAs with glycerol to form DAGs (Li et al. 2021). Moderate hydrolysis and glycerolysis are commonly used methods for DAG production. However, the obtained DAG level is less than 60% even after purification, and high levels of by-products (MAGs and FFAs) are observed (Lee et al. 2020, Li et al. 2021). Enzymatic esterification has been widely applied in the synthesis of DAGs because it is the most

direct method for DAG production. Furthermore, the usage of lipase with *sn*-1,3-regioselectivity (mainly Lipozyme RM IM and Novozym 435) at suitable ratios between the FFAs and glycerol limits the synthesis of TAGs, which is very hard to separate from DAGs (Lee et al. 2020). However, the DAG content obtained from the reactions catalyzed by these lipases is still limited to the range of 60% to 90% (Li et al. 2016). Certain levels of TAGs are nevertheless generated because of acyl migration during the esterification process. MAG and DAG lipases such as lipase SMG1 from *Malassezia globosa* showed great potential to solve this problem because of their strict specificity for MAGs and DAGs (Li et al. 2016). Wang et al. (2014) adopted SMG1 to produce high-purity DAGs (97% purity after purification). However, preparation of FFAs is a prerequisite step for esterification, which adds to the complexity and cost (Li et al. 2021). Li et al. (2021) used a combination of partial hydrolysis and esterification to solve this problem, and high-purity DAGs (99.3% after purification) were successfully obtained.

It is concluded that high-purity DAGs (generally with DAG levels higher than 80%) could be successfully prepared and commercialized. Moreover, DAGs derived from various sources have been produced to meet individual preferences and needs (Cheong et al. 2007, D.M. Li et al. 2015, Liu et al. 2018, Wang et al. 2019). The challenge encountered is whether higher-purity DAGs (>99%) can be synthesized to facilitate the exploration of their applications in the pharmaceutical field. Also, owing to the high commercial value of DAGs, fast detection methods need to be developed as a means to avoid counterfeit products in the market.

4.2. Production of Structured Triacylglycerols

STs with medium-chain and long-chain fatty acids on the glycerol backbone can be divided into, for example, MLM, MML, LML, and LLM types (M, medium-chain fatty acids; L, long-chain fatty acids). Among them, MLM is gaining more attention (Utama et al. 2019). The medium-chain fatty acids at *sn*-1,3 positions could be hydrolyzed easier in the body than the long-chain fatty acids and thus could be utilized as a quick energy source (Ding et al. 2009). Free medium-chain fatty acids have lower tendencies to be deposited in human adipose tissue (Nagao & Yanagita 2010, Vistisen et al. 2006). Meanwhile, the remaining 2-MAGs with long-chain fatty acids still provide essential fatty acids and can be absorbed well through the intestinal wall. It has been reported that MLM could be used to control obesity, fat malabsorption, and other related disorders (Utama et al. 2019). For example, CyOCy (Cy, caprylic acid; O, oleic acid) was used to treat patients with pancreatic insufficiency as well as to provide people with a rapid energy supply. However, levels of MLM are low in natural oils and fats (Utama et al. 2019); therefore, it is necessary to develop methods for preparation.

There are three commonly used methods for ST production (Utama et al. 2019) (**Figure 4a**): (a) transesterification/exchange of acyl groups between two TAG molecules; (b) direct esterification of FFAs to the glycerol molecules; and (c) acidolysis between TAG molecules and FFAs. The low yield of expected STs (Y.D. Wang et al. 2022) and high cost of the substrate (FFAs) (Yang et al. 2014) limit the application of the esterification method. Transesterification between two natural plant oils could create a complex product with various TAG species, which is not quite suitable for production of STs with certain molecular structures (Y.D. Wang et al. 2022). However, the usage of two single TAG molecules and an *sn*-1,3-specific lipase might be a good choice to solve this problem. Bai et al. (2013) adopted tricaprylin and trilinolenin as substrates and Lipozyme RM IM or Novozym 435 as a catalyst. CyLnCy and CyLnLn (Ln, linolenic acid) were found to be the most dominant TAGs. Acidolysis of long-chain TAGs with medium-chain fatty acids using *sn*-1,3-specific lipases (*Rhizomucor miehei* lipase and *Thermomyces lanuginosus* lipase) is also commonly used for producing MLM (Caballero et al. 2014). Caballero et al. (2014) studied the




Figure 4

Strategies for lipase-catalyzed lipid modification. (a) Production of diacylglycerols (green arrows) through (i) partial hydrolysis of triacylglycerol (TAG), (ii) glycerolysis reaction between TAG and glycerol, or (iii) esterification of free fatty acids (FFAs) with glycerol, catalyzed by *sn*-1,3-specific lipases. Production of structured TAGs (blue arrows) through (i) transesterification between two TAG molecules, (ii) direct esterification of FFA with glycerol, or (iii) acidolysis between TAG and FFAs, catalyzed by *sn*-1,3-specific lipases. PUFA-rich TAGs can also be produced by glycerolysis between glycerol and ethyl ester, catalyzed by nonspecific lipases. Substituted or newly added fatty acids (FAs) are highlighted in red. Abbreviations: DAG, diacylglycerol; PUFA, polyunsaturated fatty acid.

(b) Production of structured phospholipids through acidolysis (green arrows), esterification of lysophospholipids through esterification (orange arrows), and hydrolysis of structured phospholipids to produce lysophospholipids (blue arrows), catalyzed by phospholipase A1 (PLA1), phospholipase A2 (PLA2), *sn*-1,3-, or *sn*-2-specific lipases. Substituted FAs are highlighted in red. X = -H, phosphatidic acid (PA); X = -CH₂CH₂NH₂, phosphatidylethanolamine; X = -CH₂CH₂N+(CH₃)₃, phosphatidylcholine; X = -CH₂CH(OH)CH₂OH, phosphatidylglycerol; X = -CH₂CH(NH₂)COOH, phosphatidylserine.

acidolysis between caprylic acid and avocado oil and found that 29.2% of caprylic acid could be incorporated at the *sn*-1,3 positions after the optimization. To date, it is still challenging to obtain high-purity MLM because of the difficulty of separating the different TAG species. More importantly, the current detection methods are not yet able to easily differentiate the location of fatty acids, which means it is difficult to distinguish between isomers.

Human milk is the best source of energy and nutrition for all healthy infants (Wei et al. 2019). However, breastfeeding is not always possible, and infant formulas are widely considered a good alternative to human milk. Fats are crucial components in human milk and provide energy, essential fatty acids, fat-soluble vitamins, and hormones to improve the growth of infants (Wei et al. 2019). Vegetable oils and milk fat are normally added to infant formula to stimulate the fatty acid composition of human milk fat. However, TAG profiles of human milk fat, plant oils, and bovine milk fat differ substantially, which might influence the health status of infants. Thus, it is necessary to develop a human milk-fat substitute that has similar fatty acid composition and TAG profiles to human milk fat.

The distinguishing feature of human milk fat is the high level of palmitic acid at the *sn*-2 position in TAG molecules (more than 70%) (Wei et al. 2019). Moreover, OPO and OPL (O, oleic acid; P, palmitic acid; L, linoleic acid) are the major TAGs in human milk fat, accounting for 20–40% of total TAGs (Wei et al. 2019). It is thus clear that methods for *sn*-2 palmitate TAGs production must be developed. There are two schemes to prepare the *sn*-2 palmitate TAGs (**Figure 4a**): (a) one-step reactions between TAGs and another TAG/fatty acid ester/FFA and (b) two-step reactions that combine hydrolysis and esterification. Lipases with *sn*-1,3 regiospecificity (from, e.g., *R. miebei*, *Candida lipolytica*, *Candida antarctica*, *Tidestromia lanuginosa*) are important catalysts for modifying the fatty acids at the *sn*-1,3 position in both schemes (Wei et al. 2019). *sn*-2 Palmitate TAGs (65% of palmitic acid at *sn*-2 position) are produced from tripalmitin with oleic acid catalyzed by lipase (*R. miebei*) as the main lipid ingredient for the infant milk formula Betapol®. High-purity (96% of palmitic acid at the *sn*-2 position) *sn*-2 palmitate TAGs can be successfully produced using the two-step route (Ganske & Bornscheuer 2005). For now, it is not a problem to imitate the large number of TAGs in human milk-fat substitutes. There is also a certain level of MLCTs (8–10%, mainly in the form of MLL) and branched-chain fatty acids (Dingess et al. 2017, Wei et al. 2019); however, their importance has yet to be elucidated or examined. This makes it impossible to add these ingredients to current human milk-fat substitutes and simulate breast milk well; thus, it is important to further investigate the importance of the various components in breast milk for the better development of human milk-fat substitutes.

PUFA-rich TAGs are another vital ST for providing important fatty acids to humans. The human body has a limited ability to synthesize PUFAs, but these fatty acids show great benefits with respect to chronic inflammatory diseases, cancer, and sudden death. Thus, PUFAs must be obtained from the diet, for example, via consumption of fish oils. In natural fish oils, PUFAs show low purity, and further processing is necessary. At present, PUFA-rich ethyl esters are the main PUFA source on the market. However, the metabolism of ethyl esters could lead to low bioavailability and the production of ethanol, which tends to cause intoxication in humans with poor alcohol tolerance (Wang et al. 2017). In recent decades, EPA and DHA in the TAG form attracted interest because of their higher stability than that found in FFA forms and higher bioavailability than that found in ethyl ester forms. As a result, a great deal of attention has been given to the synthesis of PUFA-rich TAGs. Current breakthroughs in the preparation of PUFA-rich TAGs are focused on two methods (**Figure 4a**): (a) esterification of free PUFAs to glycerol and (b) glycerolysis between glycerol and ethyl esters. Wang et al. (2017) applied MAS1 from marine *Streptomyces* sp. strain W007 in the esterification of PUFAs and glycerol to produce PUFA-rich TAGs, and a good catalytic reaction effect was found: 92% of PUFAs could be incorporated into TAGs. However, the

free PUFAs are easily oxidized, which limits the application of the esterification method (Wang et al. 2016). Besides, cheap and easily available PUFA-rich ethyl esters are more suitable for the industrial production of PUFA-rich TAGs. Wang et al. (2016) obtained high-purity PUFA-rich TAGs (96.2%) after purification when using PUFA-rich ethyl esters and glycerol as substrates and immobilized MAS1 as a catalyst. It is noted that lipases with no regioselectivity are more suitable catalysts for the glycerolysis of PUFA-rich ethyl esters (Wang et al. 2016).

4.3. Production of Structured Phospholipids

PLs are a class of lipid compounds containing phosphate groups, which are important components of lipids and exhibit unique biological activities (Sun et al. 2018, Zhang et al. 2020). The amphiphilic property of PLs facilitates the bioavailability of fatty acids (Ulven & Holven 2015). It has been confirmed that the ingestion of lipids in the PL form exerts stronger biological effects compared with the TAG form (Ang et al. 2019). Furthermore, PLs with specific fatty acids may change the fatty acid composition of membrane PL in a certain cell type, which might be useful in targeting specific diseases and metabolic conditions (Ang et al. 2019, Kim & Akoh 2015). However, specialized PLs rarely exist in natural lipids and are difficult to obtain via fractionation and extraction. Thus, the synthesis of SPLs was triggered and researched.

SPLs refer to the modification of the structure and position of natural PLs through specific modification technologies (Castejón & Señoráns 2020). There are two general methods used in the modification of PLs (**Figure 4b**): (a) one-step acidolysis (direct transesterification) between PLs and FFAs and (b) a two-step method in combination of hydrolysis and esterification. One-step acidolysis is the direct method for SPL production, but the purity cannot be guaranteed (Adlercreutz et al. 2002). Vikbjerg et al. (2007) adopted an acidolysis reaction for preparing SPLs with caprylic acid and soybean phosphatidylcholine, and the highest incorporation rate of caprylic acid was 36%. The two-step method has the advantage of synthesizing the highly purified SPLs but is more laborious (Vikbjerg et al. 2007) because of the need to separate FFAs in the first step. Currently, the most widely used enzymes [phospholipase A1 (PLA1), phospholipase A2 (PLA2), and lipase] in PL modification vary in their regioselectivity. Both *sn*-1,3-specific lipases (Novozym 435 and Lipozyme TL IM) and PLA1 cleave the *sn*-1 position, whereas *sn*-2-specific lipases (immobilized *C. antarctica* lipase A) and PLA2 act on fatty acids at the *sn*-2 position of PLs (He et al. 2017). The position distribution of fatty acids exerts a great influence on the metabolism and biological effects of SPLs (Hu et al. 2017). For example, higher adsorption and oxidation stability could be obtained when PUFAs are esterified at the *sn*-2 position (Ang et al. 2019). In the future, the design of new SPLs, the analysis of SPL metabolic pathways, and nutritional mechanisms combined with lipidomics technology are expected to advance this field further.

4.4. Enzyme Engineering and Bioinformatics for Developing Catalysts in Lipid Modification

Although many lipases display outstanding catalytic abilities in lipid modification, there are still high demands to obtain lipases with better performance regarding their regioselectivity, fatty acid chain-length selectivity, fatty acid specificity, and higher stability to heat, extreme pH, organic solvents, and processing in industrial applications. Development of modern biotechnology enables engineering of existing lipases and the discovery of novel lipases to obtain desired enzyme functions. Rational design and directed evolution are two main strategies for protein engineering (Kourist et al. 2010). Until now, thousands of 3D lipase structures have been resolved, and their reaction mechanisms, catalytic residues, and enzyme-substrate interactions have been

widely studied (Casas-Godoy et al. 2018, Kazlauskas 1994, Schmid & Verger 1998). With this information, key residues can be predicted and mutated to the desired amino acids, depending on different purposes (Soni 2022, Zorn et al. 2016). Directed evolution is an efficient method to create mutant libraries (i.e., with more than 10^4 – 10^7 variants) that provide a much bigger space for mutagenesis and exploration of enzyme functions (Arnold 2018). It can be a completely random mutagenesis at any residue of the enzyme or limited to certain key residues that are mutated to all 19 other amino acids. Owing to the large quantity of mutants, high-throughput screening methods or selection systems are required, such as colorimetric assays, to detect the hydrolysis products of the product p-nitrophenyl (pNP) from hydrolysis of pNP esters on microtiter plates and the selection of colonies containing active lipases in TAG-agar plates. With the assistance of computational predictive algorithms and machine learning, prediction of protein mutagenesis for desired characteristics has become more precise (Mazurenko et al. 2020, Yang et al. 2019). On the other hand, unlike traditional extraction of lipases from plant/animal tissues or microorganisms, metagenomic approaches enable the discovery and exploration of novel lipases from different resources (Almeida et al. 2020, Shahraki et al. 2022, Verma et al. 2021).

5. CONCLUSIONS AND PERSPECTIVES

Food lipids are essential components in our daily diets, and some lipids (such as EPA and DHA-enriched lipids) have important health-promoting roles and have long been used as nutritional supplements. There is an urgent need to improve oil production to meet the social demand for food and energy. In the past decades, substantial efforts have been devoted to engineering oil production (enhancing oil content and producing value-added oils such as PUFAs) in traditional oil crops and newly developed lipid sources (e.g., plant vegetative tissues, microalgae, and other oleaginous microorganisms), highlighting the possibility of customizing value-added lipids in the future. Although various metabolic engineering strategies have been exploited, only moderate levels of success have been achieved so far, which is largely because of the complexity of lipid metabolism in plants and microorganisms. Therefore, further exploration of the underlying mechanisms of lipid biosynthesis, transport, and storage and their regulation is necessary.

The emergence of functional lipids may help address a substantial problem in the world, namely obesity and related disorders caused by excessive consumption of lipids with low nutritional value. Also, some diseases are caused by underconsumption of certain beneficial lipids, such as how a PUFA deficiency may affect the development of the brain and is detrimental to the prevention of cardiovascular diseases. In recent years, a lot of research has been done on the synthesis of functional lipids, mainly focusing on structural lipids (DAGs, human milk-fat substitutes, MLM, PUFA-rich TAGs, and SPLs). In turn, many functional lipids have been commercialized. However, there is still a problem of insufficient purity, and further research is needed.

Precise nutrition for different groups of the population is considered to be a promising future direction for functional lipid development, requiring customization of the right lipids according to the health status of individuals. Based on the above prospects, there are two main challenges that need to be addressed. First, it is still difficult to figure out the type of fats and oils needed by specific individuals, as current functional lipids can only meet a single need, and it is difficult to meet all the needs of individuals simultaneously. Second, the purity of the functional lipids is still difficult to guarantee. To address the above challenges, efforts need to be made to precisely describe what kind of lipids individuals need. Furthermore, upgrading metabolic engineering strategies and enzyme engineering and separation technology is needed to rationally design structured lipid production in plants and microorganisms and to achieve the preparation and acquisition of high-purity single-structured lipids at an industrial scale.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

The authors acknowledge funding from the National Key R&D Program of China (2019YFD1002403), Distinguished Young Scholars of China (31725022), National Science Fund for Key Program of National Natural Science Foundation of China (31930084), University of Guelph Start-Up Research Grant, and Natural Sciences and Engineering Research Council of Canada Discovery Grant (RGPIN-2022-03459).

LITERATURE CITED

- Adlercreutz D, Budde H, Wehtje E. 2002. Synthesis of phosphatidylcholine with defined fatty acid in the sn-1 position by lipase-catalyzed esterification and transesterification reaction. *Biotechnol. Bioeng.* 78:403–11
- Almeida JM, Alnoch RC, Souza EM, Mitchell DA, Krieger N. 2020. Metagenomics: Is it a powerful tool to obtain lipases for application in biocatalysis? *Biochim. Biophys. Acta* 1868:140320
- Ando Y, Saito S, Miura H, Osaki N, Katsuragi Y. 2017. Consumption of alpha-linolenic acid-enriched diacylglycerol induces increase in dietary fat oxidation compared with alpha-linolenic acid-enriched triacylglycerol: a randomized, double-blind trial. *Nutr. Res.* 48:85–92
- Ang X, Chen H, Xiang J-Q, Wei F, Quek SY. 2019. Preparation and functionality of lipase-catalysed structured phospholipid: a review. *Trends Food Sci. Technol.* 88:373–83
- Arias CL, Quach T, Huynh T, Nguyen H, Moretti A, et al. 2022. Expression of *AtWR11* and *AtDGAT1* during soybean embryo development influences oil and carbohydrate metabolism. *Plant Biotechnol. J.* 20(7):1327–45
- Arnold FH. 2018. Directed evolution: bringing new chemistry to life. *Angew. Chem. Int. Ed.* 57:4143–48
- Bai S, Aziz S, Khodadadi M, Bou Mitri C, St-Louis R, Kermasha S. 2013. Lipase-catalyzed synthesis of medium-long-medium type structured lipids using tricaprylin and trilinolenin as substrate models. *J. Am. Oil Chem. Soc.* 90:377–89
- Baud S, Wuillème S, To A, Rochat C, Lepiniec L. 2009. Role of WRINKLED1 in the transcriptional regulation of glycolytic and fatty acid biosynthetic genes in *Arabidopsis*. *Plant J.* 60:933–47
- Borowitzka MA. 2013. High-value products from microalgae—their development and commercialisation. *J. Appl. Phycol.* 25:743–56
- Bourdichon F, Morelli L, Zuliani V, Laulund S. 2018. *Inventory of microbial food cultures with safety demonstration in fermented food products*. IDF Bull. 455–2012, Int. Dairy Fed., Brussels, Belg.
- Boyd JT, LoCoco PM, Furr AR, Bendele MR, Tram M, et al. 2021. Elevated dietary ω -6 polyunsaturated fatty acids induce reversible peripheral nerve dysfunction that exacerbates comorbid pain conditions. *Nat. Metab.* 3:762–73
- Caballero E, Soto C, Olivares A, Altamirano C. 2014. Potential use of avocado oil on structured lipids MLM-type production catalysed by commercial immobilised lipases. *PLOS ONE* 9:e107749
- Caldo KMP, Acedo JZ, Panigrahi R, Vederas JC, Weselake RJ, Lemieux MJ. 2017. Diacylglycerol acyltransferase 1 is regulated by its N-terminal domain in response to allosteric effectors. *Plant Physiol.* 175:667–80
- Casas-Godoy L, Gasteazoro F, Duquesne S, Bordes F, Marty A, Sandoval G. 2018. Lipases: an overview. In *Lipases and Phospholipases: Methods and Protocols*, ed. G Sandoval, pp. 3–38. New York: Springer
- Castejón N, Señoráns FJ. 2020. Enzymatic modification to produce health-promoting lipids from fish oil, algae and other new omega-3 sources: a review. *New Biotechnol.* 57:45–54
- Cernac A, Benning C. 2004. *WRINKLED1* encodes an AP2/EREB domain protein involved in the control of storage compound biosynthesis in *Arabidopsis*. *Plant J.* 40:575–85

- Chen BB, Zhang GY, Li PH, Yang JH, Guo L, et al. 2020. Multiple GmWRI1s are redundantly involved in seed filling and nodulation by regulating plastidic glycolysis, lipid biosynthesis and hormone signalling in soybean (*Glycine max*). *Plant Biotechnol. J.* 18:155–71
- Cheong L-Z, Tan C-P, Long K, Affandi Yusoff MS, Arifin N, et al. 2007. Production of a diacylglycerol-enriched palm olein using lipase-catalyzed partial hydrolysis: optimization using response surface methodology. *Food Chem.* 105:1614–22
- Cintolesi A, Clomburg JM, Gonzalez R. 2014. In silico assessment of the metabolic capabilities of an engineered functional reversal of the β -oxidation cycle for the synthesis of longer-chain ($C \geq 4$) products. *Metab. Eng.* 23:100–15
- Clomburg JM, Blankschien MD, Vick JE, Chou A, Kim S, Gonzalez R. 2015. Integrated engineering of β -oxidation reversal and ω -oxidation pathways for the synthesis of medium chain ω -functionalized carboxylic acids. *Metab. Eng.* 28:202–12
- Clomburg JM, Contreras SC, Chou A, Siegel JB, Gonzalez R. 2018. Combination of type II fatty acid biosynthesis enzymes and thiolases supports a functional beta-oxidation reversal. *Metab. Eng.* 45:11–19
- Clomburg JM, Vick JE, Blankschien MD, Rodríguez-Moyá M, Gonzalez R. 2012. A synthetic biology approach to engineer a functional reversal of the β -oxidation cycle. *ACS Synth. Biol.* 1:541–54
- Dai ZJ, Huang MT, Chen Y, Siewers V, Nielsen J. 2018. Global rewiring of cellular metabolism renders *Saccharomyces cerevisiae* Crabtree negative. *Nat. Commun.* 9:3059
- Dellomonaco C, Clomburg JM, Miller EN, Gonzalez R. 2011. Engineered reversal of the β -oxidation cycle for the synthesis of fuels and chemicals. *Nature* 476:355–59
- d'Espaux L, Ghosh A, Runguphan W, Wehrs M, Xu F, et al. 2017. Engineering high-level production of fatty alcohols by *Saccharomyces cerevisiae* from lignocellulosic feedstocks. *Metab. Eng.* 42:115–25
- Ding S, Yang JK, Yan YJ. 2009. Optimization of lipase-catalyzed acidolysis of soybean oil to produce structured lipids. *J. Food Biochem.* 33:442–52
- Dingess KA, Valentine CJ, Ollberding NJ, Davidson BS, Woo JG, et al. 2017. Branched-chain fatty acid composition of human milk and the impact of maternal diet: the Global Exploration of Human Milk (GEHM) Study. *Am. J. Clin. Nutr.* 105:177–84
- Focks N, Benning C. 1998. *wrinkled1*: a novel, low-seed-oil mutant of *Arabidopsis* with a deficiency in the seed-specific regulation of carbohydrate metabolism. *Plant Physiol.* 118:91–101
- Ganske F, Bornscheuer UT. 2005. Lipase-catalyzed glucose fatty acid ester synthesis in ionic liquids. *Org. Lett.* 7:3097–98
- Ghazani SM, Marangoni AG. 2022. Microbial lipids for foods. *Trends Food Sci. Technol.* 119:593–607
- Godfray HC, Crute IR, Haddad L, Lawrence D, Muir JF, et al. 2010. The future of the global food system. *Philos. Trans. R. Soc. B* 365:2769–77
- Guo W, Chen LM, Chen HF, Yang HL, You QB, et al. 2020. Overexpression of *GmWRI1b* in soybean stably improves plant architecture and associated yield parameters, and increases total seed oil production under field conditions. *Plant Biotechnol. J.* 18:1639–41
- Hamilton ML, Haslam RP, Napier JA, Sayanova O. 2014. Metabolic engineering of *Phaeodactylum tricorutum* for the enhanced accumulation of omega-3 long chain polyunsaturated fatty acids. *Metab. Eng.* 22:3–9
- Hamrick C, Chen GX. 2021. The challenges of future foods from prevention of nutrient deficiencies to the management of diabetes. *J. Future Foods* 1:47–57
- Han L, Usher S, Sandgrind S, Hassall K, Sayanova O, et al. 2020. High level accumulation of EPA and DHA in field-grown transgenic *Camelina*: a multi-territory evaluation of TAG accumulation and heterogeneity. *Plant Biotechnol. J.* 18:2280–91
- Haslam RP, Hamilton ML, Economou CK, Smith R, Hassall KL, et al. 2020. Overexpression of an endogenous type 2 diacylglycerol acyltransferase in the marine diatom *Phaeodactylum tricorutum* enhances lipid production and omega-3 long-chain polyunsaturated fatty acid content. *Biotechnol. Biofuels* 13:87
- He YJ, Li JB, Kodali S, Chen BL, Guo ZJ. 2017. Rationale behind the near-ideal catalysis of *Candida antarctica* lipase A (CAL-A) for highly concentrating ω -3 polyunsaturated fatty acids into monoacylglycerols. *Food Chem.* 219:230–39
- Hofvander P, Ischebeck T, Turesson H, Kushwaha SK, Feussner I, et al. 2016. Potato tuber expression of *Arabidopsis WRINKLED1* increase triacylglycerol and membrane lipids while affecting central carbohydrate metabolism. *Plant Biotechnol. J.* 14:1883–98

- Holic R, Xu Y, Caldo KMP, Singer SD, Field CJ, et al. 2018. Bioactivity and biotechnological production of punicic acid. *Appl. Microbiol. Biotechnol.* 102:3537–49
- Hu P, Xu XB, Yu LLL. 2017. Interesterified trans-free fats rich in sn-2 nervonic acid prepared using *Acer truncatum* oil, palm stearin and palm kernel oil, and their physicochemical properties. *LWT* 76:156–63
- Hu Q, Sommerfeld M, Jarvis E, Ghirardi M, Posewitz M, et al. 2008. Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances. *Plant J.* 54:621–39
- Kamineni A, Shaw J. 2020. Engineering triacylglycerol production from sugars in oleaginous yeasts. *Curr. Opin. Biotechnol.* 62:239–47
- Kazlauskas RJ. 1994. Elucidating structure-mechanism relationships in lipases: prospects for predicting and engineering catalytic properties. *Trends Biotechnol.* 12:464–72
- Ke J, Wen TN, Nikolau BJ, Wurtele ES. 2000. Coordinate regulation of the nuclear and plastidic genes coding for the subunits of the heteromeric acetyl-coenzyme A carboxylase. *Plant Physiol.* 122:1057–71
- Keaney JFJ, Rosen CJ. 2019. VITAL signs for dietary supplementation to prevent cancer and heart disease. *N. Engl. J. Med.* 380:91–93
- Keereetaweep J, Liu H, Zhai ZY, Shanklin J. 2018. Biotin attachment domain-containing proteins irreversibly inhibit acetyl CoA carboxylase. *Plant Physiol.* 177:208–15
- Kim BH, Akoh CC. 2015. Recent research trends on the enzymatic synthesis of structured lipids. *J. Food Sci.* 80:C1713–24
- Kong Q, Singh SK, Mantyla JJ, Pattanaik S, Guo L, et al. 2020. TEOSINTE BRANCHED1/CYCLOIDEA/PROLIFERATING CELL FACTOR4 interacts with *WRINKLED1* to mediate seed oil biosynthesis. *Plant Physiol.* 184:658–65
- Kourist R, Brundiek H, Bornscheuer UT. 2010. Protein engineering and discovery of lipases. *Eur. J. Lipid Sci. Technol.* 112:64–74
- Koutsoumanis K, Allende A, Alvarez-Ordóñez A, Bolton D, Bover-Cid S, et al. 2021. Update of the list of QPS-recommended biological agents intentionally added to food or feed as notified to EFSA 13: suitability of taxonomic units notified to EFSA until September 2020. *EFSA J.* 19:e06377
- Lee WJ, Zhang Z, Lai OM, Tan CP, Wang Y. 2020. Diacylglycerol in food industry: synthesis methods, functionalities, health benefits, potential risks and drawbacks. *Trends Food Sci. Technol.* 97:114–25
- Lennen RM, Braden DJ, West RM, Dumesic JA, Pfleger BF. 2010. A process for microbial hydrocarbon synthesis: overproduction of fatty acids in *Escherichia coli* and catalytic conversion to alkanes. *Biotechnol. Bioeng.* 106:193–202
- Li DM, Khan FI, Zhao ZX, Wang WF, Yang B, Wang YH. 2016. Diacylglycerol production by genetically modified lipase from *Malassezia globosa*. *J. Mol. Catal. B* 133:S204–12
- Li DM, Qin X, Sun B, Wang W, Wang Y. 2019. A feasible industrialized process for producing high purity diacylglycerols with no contaminants. *Eur. J. Lipid Sci. Technol.* 121:1900039
- Li DM, Qin X, Wang J, Yang B, Wang W, et al. 2015. Hydrolysis of soybean oil to produce diacylglycerol by a lipase from *Rhizopus oryzae*. *J. Mol. Catal. B* 115:43–50
- Li DM, Zhong XR, Faiza M, Wang WF, Lian WS, et al. 2021. Simultaneous preparation of edible quality medium and high purity diacylglycerol by a novel combined approach. *LWT* 150:111949
- Li Q, Shao JH, Tang SH, Shen QW, Wang TH, et al. 2015. Wrinkled1 accelerates flowering and regulates lipid homeostasis between oil accumulation and membrane lipid anabolism in *Brassica napus*. *Front. Plant Sci.* 6:1015
- Liu J, Hua W, Zhan GM, Wei F, Wang XF, et al. 2010. Increasing seed mass and oil content in transgenic *Arabidopsis* by the overexpression of *wri1*-like gene from *Brassica napus*. *Plant Physiol. Biochem.* 48:9–15
- Liu J, Liu MJ, Pan YF, Shi Y, Hu HH. 2022. Metabolic engineering of the oleaginous alga *Nannochloropsis* for enriching eicosapentaenoic acid in triacylglycerol by combined pulling and pushing strategies. *Metab. Eng.* 69:163–74
- Liu LQ, Markham K, Blazek J, Zhou NJ, Leon D, et al. 2015. Surveying the lipogenesis landscape in *Yarrowia lipolytica* through understanding the function of a Mga2p regulatory protein mutant. *Metab. Eng.* 31:102–11
- Liu N, Li DM, Wang WF, Hollmann F, Xu L, et al. 2018. Production and immobilization of lipase PCL and its application in synthesis of α -linolenic acid-rich diacylglycerol. *J. Food Biochem.* 42:e12574

- Liu Q, Guo QG, Akbar S, Zhi Y, El Tahchy A, et al. 2017. Genetic enhancement of oil content in potato tuber (*Solanum tuberosum* L.) through an integrated metabolic engineering strategy. *Plant Biotechnol. J.* 15:56–67
- Lunn D, Wallis JG, Browse J. 2019. Tri-hydroxy-triacylglycerol is efficiently produced by position-specific castor acyltransferases. *Plant Physiol.* 179:1050–63
- Lv XQ, Wu YK, Gong MY, Deng JY, Gu Y, et al. 2021. Synthetic biology for future food: research progress and future directions. *Future Foods* 3:100025
- Ma W, Kong Q, Grix M, Mantyla JJ, Yang Y, et al. 2015. Deletion of a C-terminal intrinsically disordered region of *WRINKLED1* affects its stability and enhances oil accumulation in *Arabidopsis*. *Plant J.* 83:864–74
- Mazurenko S, Prokop Z, Damborsky J. 2020. Machine learning in enzyme engineering. *ACS Catal.* 10:1210–23
- Mehrer CR, Incha MR, Politz MC, Pflieger BF. 2018. Anaerobic production of medium-chain fatty alcohols via a beta-reduction pathway. *Metab. Eng.* 48:63–71
- Mietkiewska E, Miles R, Wickramaratna A, Sahibollah AF, Greer MS, et al. 2014. Combined transgenic expression of *Punica granatum* conjugase (*FADX*) and *FAD2* desaturase in high linoleic acid *Arabidopsis thaliana* mutant leads to increased accumulation of punicic acid. *Planta* 240:575–83
- Nagao K, Yanagita T. 2010. Medium-chain fatty acids: functional lipids for the prevention and treatment of the metabolic syndrome. *Pharmacol. Res.* 61:208–12
- Napier JA, Olsen RE, Tocher DR. 2019. Update on GM canola crops as novel sources of omega-3 fish oils. *Plant Biotechnol. J.* 17:703–5
- Nicaud J-M. 2012. *Yarrowia lipolytica*. *Yeast* 29:409–18
- Petrie JR, Shrestha P, Belide S, Kennedy Y, Lester G, et al. 2014. Metabolic engineering *Camelina sativa* with fish oil-like levels of DHA. *PLOS ONE* 9:e85061
- Petrie JR, Zhou XR, Leonforte A, McAllister J, Shrestha P, et al. 2020. Development of a *Brassica napus* (canola) crop containing fish oil-like levels of DHA in the seed oil. *Front. Plant. Sci.* 11:727
- Prabhavathi Devi BLA, Gangadhar KN, Prasad RBN, Sugasini D, Rao YPC, Lokesh BR. 2018. Nutritionally enriched 1,3-diacylglycerol-rich oil: low calorie fat with hypolipidemic effects in rats. *Food Chem.* 248:210–16
- Pyc M, Cai Y, Greer MS, Yurchenko O, Chapman KD, et al. 2017. Turning over a new leaf in lipid droplet biology. *Trends Plant Sci.* 22:596–609
- Qiao K, Abidi SHI, Liu HJ, Zhang HR, Chakraborty S, et al. 2015. Engineering lipid overproduction in the oleaginous yeast *Yarrowia lipolytica*. *Metab. Eng.* 29:56–65
- Radakovits R, Eduafo PM, Posewitz MC. 2011. Genetic engineering of fatty acid chain length in *Phaeodactylum tricornutum*. *Metab. Eng.* 13:89–95
- Reynolds HL. 1964. Food chemicals codex. *J. Assoc. Off. Agric. Chem.* 47:916–16
- Roesler K, Shen B, Bermudez E, Li C, Hunt J, et al. 2016. An improved variant of soybean type 1 diacylglycerol acyltransferase increases the oil content and decreases the soluble carbohydrate content of soybeans. *Plant Physiol. Biochem.* 171:878–93
- Roesler K, Shintani D, Savage L, Boddupalli S, Ohlrogge J. 1997. Targeting of the *Arabidopsis* homomeric acetyl-coenzyme A carboxylase to plastids of rapeseeds. *Plant Physiol.* 113:75–81
- Rousta N, Hellwig C, Wainaina S, Lukitawesa L, Agnihotri S, et al. 2021. Filamentous fungus *Aspergillus oryzae* for food: from submerged cultivation to fungal burgers and their sensory evaluation: a pilot study. *Foods* 10:2774
- Ruiz-Lopez N, Haslam RP, Napier JA, Sayanova O. 2014. Successful high-level accumulation of fish oil omega-3 long-chain polyunsaturated fatty acids in a transgenic oilseed crop. *Plant J.* 77:198–208
- Ruuska SA, Girke T, Benning C, Ohlrogge JB. 2002. Contrapuntal networks of gene expression during *Arabidopsis* seed filling. *Plant Cell* 14:1191–206
- Saito S, Yamaguchi T, Shoji K, Hibi M, Sugita T, Takase H. 2010. Effect of low concentration of diacylglycerol on mildly postprandial hypertriglyceridemia. *Atherosclerosis* 213:539–44
- Salie MJ, Thelen JJ. 2016. Regulation and structure of the heteromeric acetyl-CoA carboxylase. *Biochim. Biophys. Acta* 1861:1207–13
- Salie MJ, Zhang N, Lancikova V, Xu D, Thelen JJ. 2016. A family of negative regulators targets the committed step of de novo fatty acid biosynthesis. *Plant Cell* 28:2312–25

- Sanjaya Durrett TP, Weise SE, Benning C. 2011. Increasing the energy density of vegetative tissues by diverting carbon from starch to oil biosynthesis in transgenic *Arabidopsis*. *Plant Biotechnol. J.* 9:874–83
- Schmid RD, Verger R. 1998. Lipases: interfacial enzymes with attractive applications. *Angew. Chem. Int. Ed.* 37:1608–33
- Shahidi F, Pinaffi-Langley ACC, Fuentes J, Speisky H, de Camargo AC. 2021. Vitamin E as an essential micronutrient for human health: common, novel, and unexplored dietary sources. *Free Radic. Biol. Med.* 176:312–21
- Shahraki MF, Atanaki FF, Ariaeenejad S, Ghaffari MR, Norouzi-Beirami MH, et al. 2022. A computational learning paradigm to targeted discovery of biocatalysts from metagenomic data: a case study of lipase identification. *Biotechnol. Bioeng.* 119:1115–28
- Shen B, Allen WB, Zheng PZ, Li CJ, Glassman K, et al. 2010. Expression of *ZmLEC1* and *ZmWRI1* increases seed oil production in maize. *Plant Physiol.* 153:980–87
- Shin KS, Lee SK. 2017. Introduction of an acetyl-CoA carboxylation bypass into *Escherichia coli* for enhanced free fatty acid production. *Bioresour. Technol.* 245:1627–33
- Siloto RMP, Truksa M, Brownfield D, Good AG, Weselake RJ. 2009. Directed evolution of acyl-CoA:diacylglycerol acyltransferase: development and characterization of *Brassica napus* DGAT1 mutagenized libraries. *Plant Physiol. Biochem.* 47:456–61
- Singh N, Gaur S. 2021. GRAS fungi: a new horizon in safer food product. In *Fungi in Sustainable Food Production*, ed. X Dai, M Sharma, J Chen, pp. 27–37. Cham, Switz.: Springer
- Soni S. 2022. Trends in lipase engineering for enhanced biocatalysis. *Biotechnol. Appl. Bioc.* 69:265–72
- Sun N, Chen J, Wang D, Lin SY. 2018. Advance in food-derived phospholipids: sources, molecular species and structure as well as their biological activities. *Trends Food Sci. Technol.* 80:199–211
- Sun XM, Ren LJ, Zhao QY, Ji XJ, Huang H. 2019. Enhancement of lipid accumulation in microalgae by metabolic engineering. *Biochim. Biophys. Acta* 1864:552–66
- Taylor DC, Zhang Y, Kumar A, Francis T, Giblin EM, et al. 2009. Molecular modification of triacylglycerol accumulation by over-expression of DGAT1 to produce canola with increased seed oil content under field conditions. *Botany* 87:533–43
- Ulven SM, Holven KB. 2015. Comparison of bioavailability of krill oil versus fish oil and health effect. *Vasc. Health Risk Manag.* 11:511–24
- Uprety BK, Morrison EN, Emery RJN, Farrow SC. 2022. Customizing lipids from oleaginous microbes: leveraging exogenous and endogenous approaches. *Trends Biotechnol.* 40:482–508
- Usher S, Han L, Haslam RP, Michaelson LV, Sturtevant D, et al. 2017. Tailoring seed oil composition in the real world: optimising omega-3 long chain polyunsaturated fatty acid accumulation in transgenic *Camelina sativa*. *Sci. Rep.* 7:6570
- Utama QD, Sitanggang AB, Adawiyah DR, Hariyadi P. 2019. Lipase-catalyzed interesterification for the synthesis of medium-long-medium (MLM) structured lipids: a review. *Food Technol. Biotechnol.* 57:305–18
- van Erp H, Kelly AA, Menard G, Eastmond PJ. 2014. Multigene engineering of triacylglycerol metabolism boosts seed oil content in *Arabidopsis*. *Plant Physiol.* 165:30–36
- Vanhercke T, Belide S, Taylor MC, El Tahchy A, Okada S, et al. 2019a. Up-regulation of lipid biosynthesis increases the oil content in leaves of *Sorghum bicolor*. *Plant Biotechnol. J.* 17:220–32
- Vanhercke T, Divi UK, El Tahchy A, Liu Q, Mitchell M, et al. 2017. Step changes in leaf oil accumulation via iterative metabolic engineering. *Metab. Eng.* 39:237–46
- Vanhercke T, Dyer JM, Mullen RT, Kilaru A, Rahman MM, et al. 2019b. Metabolic engineering for enhanced oil in biomass. *Prog. Lipid Res.* 74:103–29
- Vanhercke T, El Tahchy A, Liu Q, Zhou XR, Shrestha P, et al. 2014. Metabolic engineering of biomass for high energy density: oilseed-like triacylglycerol yields from plant leaves. *Plant Biotechnol. J.* 12:231–39
- Verma S, Meghwanshi GK, Kumar R. 2021. Current perspectives for microbial lipases from extremophiles and metagenomics. *Biochimie* 182:23–36
- Vikbjerg AF, Mu HL, Xu XB. 2007. Synthesis of structured phospholipids by immobilized phospholipase A2 catalyzed acidolysis. *J. Biotechnol.* 128:545–54
- Vistisen B, Mu H, Høy C-E. 2006. Lymphatic recovery of exogenous oleic acid in rats on long chain or specific structured triacylglycerol diets. *Lipids* 41:827–34

- Wang BW, Zhang MG, Ge WH, He K, Cheng FS. 2019. Microencapsulated duck oil diacylglycerol: preparation and application as anti-obesity agent. *LWT* 101:646–52
- Wang MM, Garneau MG, Poudel AN, Lamm D, Koo AJ, et al. 2022. Overexpression of pea α -carboxyltransferase in *Arabidopsis* and *Camelina* increases fatty acid synthesis leading to improved seed oil content. *Plant J.* 110:1035–46
- Wang QT, Feng YB, Lu YD, Xin Y, Shen C, et al. 2021. Manipulating fatty-acid profile at unit chain-length resolution in the model industrial oleaginous microalgae *Nannochloropsis*. *Metab. Eng.* 66:157–66
- Wang WF, Xu Y, Qin XL, Lan DM, Yang B, Wang YH. 2014. Immobilization of lipase SMG1 and its application in synthesis of partial glycerides. *Eur. J. Lipid Sci. Technol.* 116:1063–69
- Wang XM, Li DM, Qu M, Durrani R, Yang B, Wang YH. 2017. Immobilized MAS1 lipase showed high esterification activity in the production of triacylglycerols with n-3 polyunsaturated fatty acids. *Food Chem.* 216:260–67
- Wang XM, Li DM, Wang WF, Yang B, Wang YH. 2016. A highly efficient immobilized MAS1 lipase for the glycerolysis reaction of n-3 PUFA-rich ethyl esters. *J. Mol. Catal. B* 134:25–31
- Wang YD, Zhang T, Liu RJ, Chang M, Wei W, et al. 2022. Reviews of medium- and long-chain triglyceride with respect to nutritional benefits and digestion and absorption behavior. *Food Res. Int.* 155:111058
- Wei W, Jin QZ, Wang XG. 2019. Human milk fat substitutes: past achievements and current trends. *Prog. Lipid Res.* 74:69–86
- Weselake RJ, Mietkiewska E. 2014. *Gene combinations for producing punicic acid in transgenic plants*. US Patent Appl. 14/224,582
- Weselake RJ, Shah S, Tang M, Quant PA, Snyder CL, et al. 2008. Metabolic control analysis is helpful for informed genetic manipulation of oilseed rape (*Brassica napus*) to increase seed oil content. *J. Exp. Bot.* 59:3543–49
- Wu C, Hong B, Jiang SH, Luo X, Lin H, et al. 2022. Recent advances on essential fatty acid biosynthesis and production: clarifying the roles of $\Delta 12/\Delta 15$ fatty acid desaturase. *Biochem. Eng. J.* 178:108306
- Wu JJ, Bao MJ, Duan XG, Zhou P, Chen CW, et al. 2020. Developing a pathway-independent and full-autonomous global resource allocation strategy to dynamically switching phenotypic states. *Nat Commun.* 11:5521
- Wu JJ, Zhang X, Xia XD, Dong MS. 2017a. A systematic optimization of medium chain fatty acid biosynthesis via the reverse beta-oxidation cycle in *Escherichia coli*. *Metab. Eng.* 41:115–24
- Wu JJ, Zhang X, Zhou P, Huang JY, Xia XD, et al. 2017b. Improving metabolic efficiency of the reverse beta-oxidation cycle by balancing redox cofactor requirement. *Metab. Eng.* 44:313–24
- Xin Y, Lu YD, Lee Y-Y, Wei L, Jia J, et al. 2017. Producing designer oils in industrial microalgae by rational modulation of co-evolving type-2 diacylglycerol acyltransferases. *Mol. Plant* 10:1523–39
- Xin Y, Shen C, She YT, Chen H, Wang C, et al. 2019. Biosynthesis of triacylglycerol molecules with a tailored PUFA profile in industrial microalgae. *Mol. Plant* 12:474–88
- Xu P, Qiao KJ, Suk Ahn W, Stephanopoulos G. 2016. Engineering *Yarrowia lipolytica* as a platform for synthesis of drop-in transportation fuels and oleochemicals. *PNAS* 113:10848–53
- Xu Y. 2022. Biochemistry and biotechnology of lipid accumulation in the microalga *Nannochloropsis oceanica*. *J. Agric. Food Chem.* 70(37):11500–9
- Xu Y, Caldo KMP, Pal-Nath D, Ozga J, Lemieux MJ, et al. 2018. Properties and biotechnological applications of Acyl-CoA:diacylglycerol acyltransferase and phospholipid:diacylglycerol acyltransferase from terrestrial plants and microalgae. *Lipids* 53:663–88
- Xu Y, Mietkiewska E, Shah S, Weselake RJ, Chen G. 2020. Punicic acid production in *Brassica napus*. *Metab. Eng.* 62:20–29
- Xue J, Niu YF, Huang T, Yang WD, Liu JS, Li HY. 2015. Genetic improvement of the microalga *Phaeodactylum tricornutum* for boosting neutral lipid accumulation. *Metab. Eng.* 27:1–9
- Yan D, Kim WJ, Yoo SM, Choi JH, Ha SH, et al. 2018. Repurposing type III polyketide synthase as a malonyl-CoA biosensor for metabolic engineering in bacteria. *PNAS* 115:9835–44
- Yang K, Bi Y, Sun S, Yang G, Ma S, Liu W. 2014. Optimisation of Novozym-435-catalysed esterification of fatty acid mixture for the preparation of medium- and long-chain triglycerides (MLCT) in solvent-free medium. *Int. J. Food Sci. Technol.* 49:1001–11

- Yang KK, Wu Z, Arnold FH. 2019. Machine-learning-guided directed evolution for protein engineering. *Nat. Methods* 16:687–94
- Yao S, Wang PH, Bai FR, Cao YH, Zhao T, et al. 2022. Research on the inventory of microbial food cultures in Chinese traditional fermented foods (2nd edition). *Food Ferment. Ind.* 48(1):272–85
- Ye Y, Nikovics K, To A, Lepiniec L, Fedosejevs ET, et al. 2020. Docking of acetyl-CoA carboxylase to the plastid envelope membrane attenuates fatty acid production in plants. *Nat. Commun.* 11:6191
- Yu T, Zhou YJJ, Huang MT, Liu QL, Pereira R, et al. 2018. Reprogramming yeast metabolism from alcoholic fermentation to lipogenesis. *Cell* 174:1549–58.e14
- Yurchenko O, Shockey JM, Gidda SK, Silver MI, Chapman KD, et al. 2017. Engineering the production of conjugated fatty acids in *Arabidopsis thaliana* leaves. *Plant Biotechnol. J.* 15:1010–23
- Zale J, Jung JH, Kim JY, Pathak B, Karan R, et al. 2016. Metabolic engineering of sugarcane to accumulate energy-dense triacylglycerols in vegetative biomass. *Plant Biotechnol. J.* 14:661–69
- Zhang Y, Wu GC, Zhang YJ, Wang XG, Jin QZ, et al. 2020. Advances in exogenous docosahexaenoic acid-containing phospholipids: sources, positional isomerism, biological activities, and advantages. *Compr. Rev. Food Sci. Food Saf.* 19:1420–48
- Zheng P, Allen WB, Roesler K, Williams ME, Zhang S, et al. 2008. A phenylalanine in DGAT is a key determinant of oil content and composition in maize. *Nat. Genet.* 40:367–72
- Zhou YJJ, Buijs NA, Zhu ZW, Qin JF, Siewers V, Nielsen J. 2016. Production of fatty acid-derived oleochemicals and biofuels by synthetic yeast cell factories. *Nat. Commun.* 7:11709
- Zorn K, Oroz-Guinea I, Brundiek H, Bornscheuer UT. 2016. Engineering and application of enzymes for lipid modification, an update. *Prog. Lipid Res.* 63:153–64



Contents

A Comprehensive Review of Nanoparticles for Oral Delivery in Food: Biological Fate, Evaluation Models, and Gut Microbiota Influences <i>Jingyi Xue, Christopher Blesso, and Yangchao Luo</i>	1
Novel Colloidal Food Ingredients: Protein Complexes and Conjugates <i>Fuguo Liu, David Julian McClements, Cuicui Ma, and Xuebo Liu</i>	35
Targeting Interfacial Location of Phenolic Antioxidants in Emulsions: Strategies and Benefits <i>Claire Berton-Carabin and Pierre Villeneuve</i>	63
Molecular Changes of Meat Proteins During Processing and Their Impact on Quality and Nutritional Values <i>Chunbao Li, Anthony Pius Bassey, and Guanghong Zhou</i>	85
A Dual Function of Ferritin (Animal and Plant): Its Holo Form for Iron Supplementation and Apo Form for Delivery Systems <i>Xiaoxi Chang, Chenyan Lv, and Guanghua Zhao</i>	113
Applications of the INFOGEST In Vitro Digestion Model to Foods: A Review <i>Hualu Zhou, Yunbing Tan, and David Julian McClements</i>	135
Predicting Personalized Responses to Dietary Fiber Interventions: Opportunities for Modulation of the Gut Microbiome to Improve Health <i>Car Reen Kok, Devin Rose, and Robert Hutkins</i>	157
Metabolic Signatures from Genebank Collections: An Underexploited Resource for Human Health? <i>Nese Sreenivasulu, Saleh Alseikh, Rhowell N. Tiozon Jr., Andreas Graner, Cathie Martin, and Alisdair R. Fernie</i>	183
Bioinformatic Approaches for Characterizing Molecular Structure and Function of Food Proteins <i>Harrison Helmick, Anika Jain, Genki Terashi, Andrea Liceaga, Arun K. Bhunia, Daisuke Kihara, and Jozef L. Kokini</i>	203

Biotechnology in Future Food Lipids: Opportunities and Challenges <i>Qingqing Xu, Qingyun Tang, Yang Xu, Junjun Wu, Xiangzhaobao Mao, Fuli Li, Shian Wang, and Yonghua Wang</i>	225
Engineering Nutritionally Improved Edible Plant Oils <i>Xue-Rong Zhou, Qing Liu, and Surinder Singh</i>	247
Enzymatic Approaches for Structuring Starch to Improve Functionality <i>Ming Miao and James N. BeMiller</i>	271
Nondigestible Functional Oligosaccharides: Enzymatic Production and Food Applications for Intestinal Health <i>Shaoqing Yang, Chenxuan Wu, Qiaojuan Yan, Xiuting Li, and Zhengqiang Jiang</i>	297
Diet-Derived Antioxidants: The Special Case of Ergothioneine <i>Barry Halliwell, Richard M.Y. Tang, and Irwin K. Cheah</i>	323
Indole-3-Carbinol: Occurrence, Health-Beneficial Properties, and Cellular/Molecular Mechanisms <i>Darshika Amarakoon, Wu-Joo Lee, Gillian Tamia, and Seong-Ho Lee</i>	347
Bacteriophages in the Dairy Industry: A Problem Solved? <i>Guillermo Ortiz Charneco, Paul P. de Waal, Irma M.H. van Rijswijk, Noël N.M.E. van Peij, Douwe van Sinderen, and Jennifer Mabony</i>	367
Bovine Colostrum for Veterinary and Human Health Applications: A Critical Review <i>Kevin Linehan, R. Paul Ross, and Catherine Stanton</i>	387
Addressing Consumer Desires for Sustainable Food Systems: Contentions and Compromises <i>Craig Upright</i>	411
Sensory Analysis and Consumer Preference: Best Practices <i>M.A. Drake, M. E. Watson, and Y. Liu</i>	427
Mechano-Bactericidal Surfaces: Mechanisms, Nanofabrication, and Prospects for Food Applications <i>Yifan Cheng, Xiaojing Ma, Trevor Franklin, Rong Yang, and Carmen I. Moraru</i>	449
Mild Fractionation for More Sustainable Food Ingredients <i>A. Lie-Piang, J. Yang, M.A.I. Schutyser, C.V. Nikiforidis, and R.M. Boom</i>	473
Microbubbles in Food Technology <i>Jiakai Lu, Owen G. Jones, Weixin Yan, and Carlos M. Corvalan</i>	495
How Can AI Help Improve Food Safety? <i>C. Qian, S.I. Murphy, R.H. Orsi, and M. Wiedmann</i>	517

Microgreens for Home, Commercial, and Space Farming:
A Comprehensive Update of the Most Recent Developments
*Zi Teng, Yaguang Luo, Daniel J. Pearlstein, Raymond M. Wheeler,
Christina M. Johnson, Qin Wang, and Jorge M. Fonseca* 539

Errata

An online log of corrections to *Annual Review of Food Science and Technology* articles may be found at <http://www.annualreviews.org/errata/food>