UNIVERSITY of York

This is a repository copy of *Risk factors associated with overweight and obesity in people with severe mental illness in South Asia:Cross-sectional study in Bangladesh, India, and Pakistan.*

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/206252/</u>

Version: Published Version

Article:

Appuhamy, Koralagamage Kavindu, Podmore, Danielle, Mitchell, Alex orcid.org/0000-0001-9311-2092 et al. (15 more authors) (2023) Risk factors associated with overweight and obesity in people with severe mental illness in South Asia:Cross-sectional study in Bangladesh, India, and Pakistan. Journal of nutritional science. e116. ISSN 2048-6790

https://doi.org/10.1017/jns.2023.100

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

JNS Journal of nutritional science



RESEARCH ARTICLE

Risk factors associated with overweight and obesity in people with severe mental illness in South Asia: cross-sectional study in Bangladesh, India, and Pakistan

Koralagamage Kavindu Appuhamy¹* ^(D), Danielle Podmore¹ ^(D), Alex Mitchell¹, Helal Uddin Ahmed², Mark Ashworth³, Jan R. Boehnke^{1,4}, Virtu Chongtham⁵, Asiful Haidar Chowdhury⁶, Olga P. Garcia⁷, Richard I. G. Holt^{8,9}, Rumana Huque⁶, Krishna Prasad Muliyala¹⁰, Eline Klein Onstenk¹¹, Sukanya Rajan¹⁰, David Shiers^{12,13,14}, Najma Siddiqi^{1,15,16}, S. Manjunatha¹⁰ and Gerardo A. Zavala¹ ¹Department of Health Sciences, University of York, York, UK ²National Institute of Mental Health & Hospital, Dhaka, Bangladesh ³School of Lifecourse and Population Sciences, King's College, London, UK ⁴School of Health Sciences, University of Dundee, Dundee, UK ⁵Department of Psychiatry, Government Medical College and Hospital, Chandigarh, India ⁶ARK Foundation, Dhaka, Bangladesh ⁷Facultad de Ciencias Naturales, Universidad Autonoma de Queretaro, Santiago de Querétaro, Mexico ⁸Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK ⁹Southampton National Institute for Health Research Biomedical Research Centre, University Hospital Southampton, Southampton, UK ¹⁰National Institute of Mental Health and Neurosciences, Bangalore, India ¹¹Vrije Universiteit Amsterdam, Amsterdam, the Netherlands ¹²Psychosis Research Unit, Greater Manchester Mental Health NHS Trust, Manchester, UK ¹³Division of Psychology and Mental Health, University of Manchester, Manchester, UK ¹⁴School of Medicine, Keele University, Keele, UK ¹⁵Hull York Medical School, York, UK ¹⁶Bradford District Care NHS Foundation Trust, Bradford, UK (Received 3 August 2023 – Final revision received 13 September 2023 – Accepted 20 October 2023) Journal of Nutritional Science (2023), vol. 12, e116, page 1 of 14 doi:10.1017/jns.2023.100

Abstract

Obesity is one of the major contributors to the excess mortality seen in people with severe mental illness (SMI) and in low- and middle-income countries people with SMI may be at an even greater risk. In this study, we aimed to determine the prevalence of obesity and overweight in people with SMI and investigate the association of obesity and overweight with sociodemographic variables, other physical comorbidities, and health-risk behaviours. This was a multi-country cross-sectional survey study where data were collected from 3989 adults with SMI from three specialist mental health institutions in Bangladesh, India, and Pakistan. The prevalence of overweight and obesity was estimated using Asian BMI thresholds. Multinomial regression models were then used to explore associations between overweight and obesity with various potential determinants. There was a high prevalence of overweight (17·3 %) and obesity (46·2 %). The relative risk of having obesity (compared to normal weight) was double in women (RRR = 2·04) compared with men. Participants who met the WHO recommendations for fruit and vegetable intake had 2·53 (95 % CI: 1·65–3·88) times greater risk of having obesity compared to those not meeting them. Also, the relative risk of having obesity in people with hypertension is 69 % higher than in people without hypertension (RRR = 1·69). In conclusion, obesity is highly prevalent in SMI and associated with chronic disease. The complex relationship between diet and risk of

* Corresponding author: Koralagamage Kavindu Appuhamy, email: kka505@york.ac.uk

© The Author(s), 2023. Published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



obesity was also highlighted. People with SMI and obesity could benefit from screening for non-communicable diseases, better nutritional education, and context-appropriate lifestyle interventions.

Key words: Obesity: Overweight: Severe mental illness: South Asia

Introduction

Severe mental illness (SMI) is broadly defined as a group of mental disorders that are characterised by their persistence and their extensive impact on a person's life.⁽¹⁾ This group includes schizophrenia spectrum disorders, bipolar disorder, and severe depression with psychotic features. People with SMI die on average 10–20 years earlier than the general population. Obesity and its comorbidities are common in people with SMI and are estimated to contribute to one third of the excess mortality.^(2–4)

People with SMI tend to consume a diet low in fruits and vegetables and with more high-calorie convenience foods and sugar-sweetened beverages.^(5–7) This is in part due to the increased hunger caused by antipsychotics acting on various receptors.⁽⁷⁾ People with SMI also tend to be more sedentary due to negative symptoms and lack of motivation associated with their mental condition.⁽⁸⁾

The World Health Organisation (WHO) defines overweight and obesity as 'abnormal or excessive fat accumulation that presents a risk to health'.⁽⁹⁾ Obesity is associated with a range of physical health problems including dyslipidaemia, type 2 diabetes, hypertension, cardiovascular disease, and some cancers.⁽¹⁰⁾ There are also psychosocial sequalae which lead to further disadvantage including lack of self-esteem and motivation, discrimination in several settings including education and employment, and a reduced quality of life.⁽¹¹⁾

The global epidemic of obesity is particularly affecting people from low- and middle-income countries (LMICs) where there have been rapid changes in diet and lifestyle.⁽¹²⁾ There has been a nutrition transition away from traditional diets consisting of non-processed foods and pulses towards more energy-dense foods with added sugars and fats.^(12,13) Furthermore, there has been a decline in physical activity due to more sedentary jobs, and the increasing use of motorised transport.⁽¹²⁾ In addition to the high prevalence of obesity, more than 80 % of people with mental illness live in LMICs.⁽¹⁴⁾

The overlap between obesity and mental illness is therefore likely to be significant in South Asia where mental health service provision continues to be scarce.⁽¹⁴⁾ There is also an increased prevalence of diabetes and other cardiovascular risk factors seen at lower BMI values in the Asian population, as well as higher body fat found at lower BMI values.^(15,16) WHO guidance has, therefore, suggested the use of lower cutoff values for the Asian population based on such risk factors and comorbidities.⁽¹⁷⁾

Despite the increasing prevalence of SMI and obesity in LMICs, there is scarce evidence examining the scale of this comorbidity in the most affected areas; only 20 % of studies related to obesity in people with SMI have been conducted in LMICs.⁽⁴⁾ The disproportionate representation of higher-income settings means that evidence-based strategies may

not take into account the societal and cultural contexts specific to LMICs.

There is an urgent need to understand the prevalence of overweight and obesity in people with SMI in LMICs to guide practice and policy and aid in the development and adaptation of targeted interventions. Determining what association there may be between specific health problems and health-risk behaviours with obesity and overweight will identify those most at risk. This study aims to (1) determine the prevalence of obesity and overweight in adults with SMI in Bangladesh, India, and Pakistan and (2) investigate the association of obesity and overweight with sociodemographic variables, physical health conditions, and health-risk behaviours.

Methods

Study design

This is a study based on a cross-sectional survey that has taken place across mental health institutes in Bangladesh, India, and Pakistan, looking at the physical health of people with SMI, as part of the IMPACT programme.⁽¹⁸⁾

Setting

The cross-sectional survey took place across three national specialist mental health institutions: the National Institute of Mental Health and Hospital (NIMHH) in Dhaka, Bangladesh; the National Institute of Mental Health and Neurosciences (NIMHANS) in Bengaluru, India; the Institute of Psychiatry (IOP) Rawalpindi Medical University, Pakistan.⁽¹⁸⁾ Although they are tertiary care units, the general lack of mental health care provision for SMI at the primary and secondary care level means they serve the general population of people with SMI from across each country.

Participants

Adults (over 18 years old) who were diagnosed with SMI by their local physician and able to provide informed consent were invited to participate in the study. SMI was defined using the International Classification of Disease 10th Revision (ICD-10) as schizophrenia, schizotypal, and delusional disorders (F20-29), bipolar affective disorder (F30, F31), and severe depression with psychotic symptoms (F32·3, F33·3). Diagnosis of SMI was confirmed by trained researchers using the Mini International Neuropsychiatric Interview (MINI) V.6.0.⁽¹⁹⁾

Recruitment and consent

A total of 5801 potential participants were approached during the recruitment period, of these, 3989 adults with SMI from



the three national specialist mental health institutions agreed to participate. Researchers provided written and verbal information about the study to all eligible participants, after which informed consent was sought (or, if incapacitated at the initial point of contact, sought later if capacity was regained during the study period).

We recruited inpatients and outpatients, with approximately 20 % inpatients and 80 % outpatients reflecting the usual proportions seen in practice.⁽¹⁸⁾ At NIMH and NIMHANS, patients were randomly selected. Due to low patient flow at IOP, all patients attending this centre during the study period were invited to participate. The recruitment of participants and data collection took place between June 2019 and January 2022.

Data collection

Face-to-face interviews were carried out to collect information about mental and physical health, risk factors, and health-risk behaviours.⁽²⁰⁾ The survey was translated into the most common local languages spoken in each country (Urdu in Pakistan, Bangla in Bangladesh, Hindi, and Kannada in India).

Dependent variables and measurement

BMI categories. To calculate BMI, we measured the height and weight of all participants in accordance with the WHO guidelines.⁽²¹⁾ Height was measured to a precision of 0.1 cm using a portable height measuring board with participants removing footwear or headgear. Weight was measured in kilograms using a portable weighing scale, with participants in light clothing and no footwear. Both height and weight measurements were taken twice, and the average was used for analysis.

BMI was calculated (weight (kg)/height (m)²) and BMI categories were assigned using both international (normal weight 18.5 kg/m^2 -24.99 kg/m²; overweight 25 kg/m²-29.99 kg/m²; obesity $\geq 30 \text{ kg/m}^2$) and Asian thresholds which have lower cut-offs (normal weight 18.5 kg/m^2 22.99 kg/m²; overweight 23 kg/m²-24.99 kg/m²; obesity $\geq 25 \text{ kg/m}^2$).^(15,17)

Abdominal obesity. To determine the abdominal obesity, waist circumference was measured in duplicate to a precision of 0.1 cm using flexible fibreglass tape at the end of normal expiration, between the lower margin of the last palpable rib and the top of the iliac crest.⁽²¹⁾ Ethnicity-specific cut-off values for waist circumference have been recommended for the Asian population by International Diabetes Federation (IDF): \geq 90 cm for men and \geq 80 cm for women.⁽²²⁾

Independent variables and measurement

Several factors associated with obesity were investigated using the WHO STEPS instrument version 3.2,⁽²⁰⁾ including physical comorbidities, health-risk behaviours, and sociodemographic variables.

Physical comorbidities. Hypertension was defined by blood pressure (BP) exceeding the cut-off (Systolic BP >140 mmHg or Diastolic BP >90 mmHg) when measured during the survey, or those who reported diagnosis from a healthcare professional. BP was measured according to the WHO guidelines,⁽²¹⁾ using an automated BP monitor (OMRON).

Type 2 diabetes was defined by the HbA_{1c} measurement $\geq 6.5\%$ (48 mmol/mol) and those who self-reported. We also defined pre-diabetes according to the American Diabetes Association,⁽²³⁾ as HbA_{1c} between 5.7 to 6.4% (39–47 mmol/mol).

High cholesterol was defined as a low-denisty lipoprotein (LDL) concentration ≥ 1 g/l according to their serum test during the survey and those who self-reported.⁽²⁴⁾ High triglycerides were defined solely based on serum blood tests (≥ 1.8 g/l).⁽²⁴⁾ All blood collection was carried out in accordance with the WHO STEPS surveillance manual.⁽²¹⁾

Health-risk behaviours. Variables were based upon whether the participants followed WHO recommendations for physical activity and fruit and vegetable intake.⁽²⁵⁾ Self-reported current smoking status was also recorded.

Sociodemographic and clinical variables. The variables included SMI diagnosis, SMI duration (years), antipsychotic use, clinical setting (inpatient/outpatient), and sociodemographic variables; age, sex, highest level of education, work status, and income tertiles.

Sample size

A sample size of 865 was originally calculated to be able to estimate the prevalence of diabetes with a precision of 2 % as an example of survey precision. However, this sample size is also sufficient to estimate the prevalence of obesity with a precision <2 %, considering a prevalence estimate of 10 %.⁽¹⁸⁾

Statistical analysis

This study was reported according to STROBE guidelines.⁽²⁶⁾ All statistical analyses were carried out using Stata v.17. Statistical significance was assessed at the 5 % level.

Participant characteristics were summarised descriptively for each country and overall. Continuous variables were reported as means and standard deviations (and/or median, interquartile range, and minimum and maximum as appropriate), and categorical variables were reported as frequencies and percentages.

For each country separately and overall, the prevalence of underweight, normal weight, overweight, and obesity was reported using both the WHO international and the Asian BMI cut-off values. The prevalence of normal weight, overweight and obesity, using Asian cut-off values, was stratified by key characteristics (sex, age, SMI diagnosis and setting), reported by country and overall.

Ŷ

To investigate the associations between BMI categories and other comorbidities, multinomial logistic regression models were fitted with BMI group as the dependent variable. In the interests of accurately quantifying the disease burden of obesity and ensuring it is clinically relevant for the target population, Asian cut-off values were used.

Individuals classified as underweight, according to BMI category, were excluded from this analysis as the examination of the association of being underweight with associated risk factors was not an objective of this study. Physical health comorbidities (diabetes, hypertension, high cholesterol, high triglycerides), health-risk behaviours (smoking status, physical activity, diet), and sociodemographic variables (age group, sex, type of SMI, SMI duration, antipsychotic medication, setting, highest level of education, employment status, income tertile, and country) were included as independent variables. Interactions between the independent variables and country were assessed using a like-lihood ratio test to compare these to the model with no interaction terms. Relative risk ratios (RRR) were reported along with corresponding 95 % confidence intervals and *P*-values. Unadjusted estimates are reported in the appendix.

The associations between abdominal obesity and its determinants were investigated. Logistic regression models were fitted with abdominal obesity as the dichotomous (yes/no) dependent variable. The same variables as in the analysis of BMI categories were included as independent variables. Odds ratios (OR) were reported along with corresponding 95 % confidence intervals and *P*-values. Unadjusted estimates are reported in the appendix.

Analysis models included complete cases only; however, multiple imputation (MI) was performed as a sensitivity analysis, using chained equations to impute missing data. The results were not changed by the MI analysis. Results of the MI analyses are presented in the appendix alongside the complete-case analysis for comparison.

Results

From the 3989 participants included in the study, 3126 (78·4 %) participants were included in the multinomial logistic regression (BMI categories), and 3389 participants ($85 \cdot 0$ %) were included in the binomial logistic regression (abdominal obesity). The excluded participants for each analysis, and the reasons for this, are detailed in the flowchart found in the appendix.

Participant characteristics

The characteristics of study participants are presented in Table 1. A total of 59.1 % of the participants were male. The average age was 35.8 years (sD 11.9; range 18-84 years). Schizophrenia-type disorder was the most common type of SMI with 44.7 % of participants having this diagnosis. The majority of the sample were outpatients (82.5 %), and the lowest income group was the most common (41.4 %).

Prevalence of obesity and overweight

The overall prevalence of obesity across the three countries was $16\cdot0$ % according to the WHO international BMI cut-offs

and 46.2 % according to the WHO Asian cut-offs. The overall prevalence of overweight across the three countries was 30.2 % according to WHO international BMI cut-offs and 17.3 % according to Asian cut-offs.

The overall prevalence of abdominal obesity was 53.8%; however, differences were observed between sexes and countries. Abdominal obesity was less prevalent in men from Bangladesh (27.2%) compared to India (53.1%) and Pakistan (49.4%) despite relatively similar prevalence found amongst the female participants in each country.

Overall and stratified prevalence of BMI categories

According to the Asian cut-offs, the largest proportion of participants (46.2%) were classified as having obesity, compared to the international cut-offs, where the largest proportion (46.0%) were classed as having normal weight (Table 2).

Overall and stratified prevalence for each country

The prevalence of obesity in India and Pakistan was similar (50.7 % [95 % CI: 47.9-53.6] and 52.9 % [95 % CI: 50.2-55.6] respectively), and higher than in Bangladesh (36.8 % [95 % CI: 34.4-39.3]) (Table 3). Overall, the proportion of men with normal weight (31.5 % [95 % CI: 29.6-33.4]) was greater than the proportion of women with normal weight (24.9 % [95 % CI: 22.8-27.0]). Obesity was most prevalent in the 40-54-year age group (56.0 % [95 % CI: 53.0-59.0]) and least prevalent amongst 18-24-year-olds (26.5 % [95 % CI: 23·4-29·8]). Obesity was most prevalent in those with a diagnosis of major depressive disorder with psychotic symptoms in Bangladesh (43.4 % [95 % CI: 32.8-54.7]), and most prevalent in people with bipolar disorder in India (56.7 % [95 % CI: 52.0-61.3]) and Pakistan (57.3 % [95 % CI: 53.1-61.4]). In addition, a greater proportion of outpatients had obesity (48 % [95 % CI: 46.3-49.7]) compared to the inpatients (37.5 % [95 % CI: 33.9-41.2]). The prevalence of underweight was 8.6 % (95 % CI: 7.3-10.2) for Bangladesh, 8.7 % (95 % CI: 7.3-10.3) for Pakistan, and 5.7 % (95 % CI: 4.5-7.2) for India.

Association of predictors of overweight and obesity

The multinomial logistic regression analysis for the association of predictors of overweight and obesity are shown in Table 4a (sociodemographic variables) and Table 4b (comorbidities and health-risk behaviours).

The relative risk of having obesity (compared with normal weight) is double in women compared with men (RRR = 2.04 [95 % CI: 1.56-2.67]). The percentage of participants with overweight was lower in females compared to males, whilst the percentage of participants with obesity was higher in females compared to males. The percentage of patients with overweight or obesity was higher in females than in males (males 1192/1860 (64.1 %)); females 917/1266 (72.4 %)). Compared to 18–24-year-olds, the 40–54-year age group has the greatest relative risk of having obesity (RRR = 2.91 [95 % CI: 2.15–3.92]), and the relative risk of having



Table 1. Participant characteristics summarised overall and by country

Sec. (%) Number with data 150 (100) 175 (100) 1374 (100) 3869 (100) Mate 915 (61.0) 648 (55.1) 796 (100.6) 153 (10.4) 1530 (10.0) Age (yara)		Bangladesh (<i>n</i> 1500)	India (<i>n</i> 1175)	Pakistan (<i>n</i> 1314)	Overall (<i>n</i> 3989)
Number with data 1500 (100) 117, (100) 134 (100) 2359 (100) Farnala 255 (24-9) 517 (28-4) 518 (28-4) 518 (28-4) Agr (varis) 1175 (100) 135 (10-2) 335 (10-2) 335 (10-2) 335 (10-2) n (%) 1500 (100) 175 (100) 138 (10-2) 355 (10-2) 355 (10-2) 355 (10-2) 355 (10-2) Mardian (102h) 130 (100) 175 (100) 134 (100) 386 (100) 175 (100) 134 (100) 386 (100) 16-2-4 years 434 (28-3) 123 (10-5) 159 (12-1) 77 (617) 72 (10-1) 134 (100) 386 (100) 16-2-4 years 235 (17-5) 386 (28-0) 133 (40-0) 1384 (10-1) 386 (100) 25-39 gears 235 (17-5) 326 (28-0) 137 (10-1) 134 (100) 386 (100) 16-2-4 years 235 (17-5) 326 (28-0) 137 (10-1) 1384 (100) 386 (10-1) 25-30 grave 337 (40-3) 375 (10-1) 134 (100) 386 (100) 137 (10-1) 134 (100) 386 (100)	Sex, <i>n</i> (%)				
Male 915 (61-0) 648 (55.1) 796 (60-6) 2359 (64-1) Age (vers)	Number with data	1500 (100)	1175 (100)	1314 (100)	3989 (100)
Fennale 568 (39 0) 527 (44 9) 518 (39 4) (150) (100) n (%) 1500 (100) 1775 (100) 3314 (100) 358 (110) 358 (100) 358	Male	915 (61.0)	648 (55.1)	796 (60.6)	2359 (59.1)
Age (parts) n (%) Moda ((n) Moda ((n) Hore (Moda (140) Hore (Moda (140)	Female	585 (39.0)	527 (44·9)	518 (39.4)	1630 (40.9)
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	Age (years)				
Main (o) 315 (108) 388 (112) 88 (112) 35 (113) Median (IOP) 30 (220-30) 38 (300-46.0) 36 (220-45.0) 35 (250-44.0) Min, Nax 160, 760 150, 810 180, 840 180, 840 180, 840 Min, Nax 150, 1700 1314 (100) 2898 (100) 1775 (100) 1314 (100) 2898 (172) 32-39 years 733 (48, 8) 536 (453) 633 (459) 1873 (47.0) 40-44 years 283 (175) 386 (32.9) 412 (30.6) 1051 (26.3) S5- years 71 (47) 128 (100) 1314 (100) 3989 (100) Schapphenis-type disorder 853 (22.3) 673 (57.3) 176 (13.4) 178 (44.7) Major dynezside biooder with psycholic features 455 (28.1) 259 (25.4) 299 (22.4) 1443 (26.1) Duration of the SML <i>p</i> (%) 1500 (100) 1775 (100) 1314 (100) 3989 (100) s2 years 438 (28.1) 215 (18.3) 239 (22.4) 1433 (26.1) Or ont SML <i>p</i> (%) 1500 (100) 1775 (100) 1314 (100) 398 (100)	n (%)	1500 (100)	1175 (100)	1314 (100)	3989 (100)
	Mean (sd)	31.5 (10.8)	38.8 (11.2)	38.1 (12.3)	35.8 (11.9)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Median (IQR)	30.0 (23.0–38.0)	38.0 (30.0–46.0)	36.0 (28.0–45.0)	35.0 (26.0–44.0)
Age (Bord), n (%) 1175 (100) 1314 (100) 3980 (100) 18-24 years 439 (28 y) 123 (16 5) 159 (12 1) 7716 (17 9) 18-24 years 439 (28 y) 123 (16 5) 159 (12 1) 7716 (17 9) 18-24 years 265 (17 5) 386 (28 9) 402 (20 6) 1051 (26 3) 55 years 265 (17 5) 386 (28 9) 402 (20 6) 1147 (100) 3889 (100) Schizophrenisky of disorder with psycholic features 77 (51 1 63 (57 3) 177 (13 4) 1744 (44 7) Major depressive disorder with psycholic features 77 (51 1 63 (54 4) 601 (45 7) 744 (18 6) Duration of the SM, n (%) The system of the system	Min, Max	18.0, 76.0	18.0, 81.0	18.0, 84.0	18.0, 84.0
Number Instructure Issue (1co) Instructure Instructure <thinstructure< th=""> <thinstructure< th=""> <thi< td=""><td>Age group, n (%)</td><td>1500 (100)</td><td>1175 (100)</td><td>1014 (100)</td><td>2020 (100)</td></thi<></thinstructure<></thinstructure<>	Age group, n (%)	1500 (100)	1175 (100)	1014 (100)	2020 (100)
$\begin{array}{cccccccccccccccccccccccccccccccccccc$		1500 (100)	100 (100)	1314 (100)	3989 (100)
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	18-24 years	434 (28·9) 722 (49 9)	123 (10·5) 529 (45 9)	109 (12·1) 602 (45 0)	/ 10 (17·9) 1972 (47 0)
Ss. mystal Description Description Description Ss. mystal T1 (47) 128 (10.9) 130 (11.4) 349 (87) Number with data 1500 (100) 1175 (100) 134 (100) 389 (100) Solitophronic hype diaorder 935 (82.3) 673 (67.3) 176 (13.4) 1794 (44.7) Major dispressive disorder 935 (82.3) 673 (67.3) 176 (17.4) 389 (100) Duration of the SM, n (%) 1500 (100) 1175 (100) 1314 (100) 3989 (100) 22 years 456 (22.6) 320 (24.4) 1043 (28.6) 320 (24.4) 1043 (28.6) 3-5 years 322 (22.1) 299 (25.4) 299 (25.4) 299 (25.6) 1029 (25.6) Do not know/Dort remember 4 (0.3) 36 (3.1) 7 (0.5) 47 (1.2) Number with data 1500 (100) 1175 (100) 1314 (100) 3898 (100) Inpatient 1187 (75.1) 911 (27.5) 1122 (9.8) 386 (99.9) Number with data 1500 (100) 1174 (99.9) 1312 (99.8) 386 (99.9) No fromal educatio	20-59 years	732 (40·0) 263 (17 5)	386 (32.0)	402 (30 6)	1073 (47.0)
SMI, Tripa The Prior LD (10.5) LD (10.5) LD (10.5) LD (10.5) SDD (10.0) Solit problem with data 5500 (10.0) 175 (10.0) 134 (10.0) 3980 (10.0) Solit problem with data 5500 (10.0) 175 (10.0) 1344 (10.0) 3980 (10.0) Duration of the Solit All solit Al	55+ years	200 (17-0) 71 (4.7)	128 (10.9)	402 (00·0) 150 (11.4)	3/9 (8.7)
Number with data 1500 (100) 1175 (100) 1344 (100) 3989 (100) Schizophreniz-bye disorder 935 (82.3) 673 (67.3) 176 (13.4) 1784 (44.7) Major depressive disorder 935 (82.3) 633 (67.3) 631 (64.7) 741 (18.6) Duration of the SML n (%) 175 (100) 1175 (100) 1344 (100) 3989 (100) Szyaars 436 (22.1) 256 (22.6) 320 (24.4) 1043 (25.1) 3-5 years 328 (22.1) 259 (25.4) 299 (25.4) 939 (30.4) 1022 (25.8) D on know/Dort member 4 (0.3) 36 (3.1) 7 (0.5) 47 (12.2) Setting, n (%) 1177 (70.1) 911 (77.5) 1132 (90.8) 3986 (100) Number with data 1500 (100) 1174 (99.9) 131 (20.8) 3986 (90.9) No formal advatation 151 (10.1) 141 (12.0) 3988 (100) 118 (90.8) 3986 (90.9) No formal advatation 151 (10.1) 141 (12.0) 257 (18.6) 549 (13.8) Primary education 242 (56.1) 401 (34.2) 244 (1.7.8) 1477 (37.1)	SML $n(\%)$	71 (47)	120 (10.3)	100 (11.4)	040 (0.7)
	Number with data	1500 (100)	1175 (100)	1314 (100)	3989 (100)
$\begin{array}{cccc} \mbox{Major depressive disorder with psychotic features} & 77 (5,1) & 63 (5,4) & 601 (45.7) & 741 (16.6) \\ \mbox{Bignolar disorder} & 488 (32.6) & 439 (37.4) & 537 (40.9) & 1464 (36.7) \\ \mbox{Duration of the SMI, n (%) & 1500 (100) & 1175 (100) & 1314 (100) & 3989 (100) \\ \mbox{3.5 years} & 457 (30.5) & 266 (22.6) & 230 (22.4) & 1043 (26.1) \\ \mbox{3.5 years} & 457 (30.5) & 266 (22.6) & 329 (22.8) & 930 (23.3) \\ \mbox{3.5 years} & 457 (30.5) & 266 (22.6) & 329 (22.8) & 930 (23.3) \\ \mbox{3.5 years} & 271 (18.1) & 359 (30.6) & 399 (30.4) & 1029 (25.8) \\ \mbox{3.5 to not know/Don't remember} & 4 (0.3) & 36 (3.1) & 7 (0.5) & 47 (1.2) \\ \mbox{3.5 to not know/Don't remember} & 4 (0.3) & 36 (3.1) & 7 (0.5) & 47 (1.2) \\ \mbox{3.5 to not know/Don't remember} & 4 (0.3) & 313 (20.9) & 224 (22.5) & 122 (9.3) & 699 (17.5) \\ \mbox{3.5 to not know/Don't remember} & 1187 (79.1) & 1177 (100) & 1314 (100) & 3989 (100) \\ \mbox{1.5 through data} & 1500 (100) & 1175 (100) & 1314 (100) & 3989 (100) \\ \mbox{1.5 to not advacation, n (%) \\ \mbox{3.5 to not advacation, n (%) \\ \mbox{3.5 to not advacation} & 151 (10.1) & 141 (12.0) & 257 (19.6) & 349 (13.8) \\ \mbox{3.5 to not advacation} & 152 (10.0) & 1174 (99.9) & 1312 (99.8) & 3986 (99.9) \\ \mbox{3.5 to not advacation} & 234 (15.6) & 228 (19.4) & 273 (20.8) & 735 (16.4) \\ 3.5 tartus (past 12 months), n (%) \\ \mbox{3.5 tartus (past 12 months)$	Schizophrenia-type disorder	935 (62-3)	673 (57.3)	176 (13.4)	1784 (44.7)
	Major depressive disorder with psychotic features	77 (5·1)	63 (5·4)	601 (45.7)	741 (18.6)
$ \begin{array}{c} \text{Duration of the SML}, r(\%) \\ \text{Number with data} \\ 350 (100) \\ 175 (100) \\ 175 (100) \\ 175 (100) \\ 175 (100) \\ 175 (100) \\ 175 (100) \\ 175 (100) \\ 175 (100) \\ 175 (100) \\ 100 \\$	Bipolar disorder	488 (32.5)	439 (37.4)	537 (40.9)	1464 (36.7)
Number with data1500 (100)1175 (100)1134 (100)3989 (100)s2 years436 (26+1)215 (18.3)289 (22-0)940 (22-6)3-5 years457 (30-5)266 (22-6)302 (24-4)1043 (26-1)6-10 years332 (22-1)299 (23-4)299 (23-6)930 (23-3)>10 years271 (18.1)359 (30-6)399 (30-4)1023 (22-8)Do not knowDon't member4 (0-3)36 (3-1)7 (0-5)47 (1-2)Setting, n (%)1175 (100)1314 (100)3989 (100)Inpatient1313 (20-9)224 (22-5)122 (3-3)699 (17.5)Outpatient1187 (79-1)911 (77.5)1182 (30-7)3280 (22-5)Highest level of education, n (%)1174 (199-9)1312 (39-8)396 (99-9)No formal education151 (10.1)141 (12-0)227 (18-5)449 (13-8)Primary education234 (15-6)228 (16-4)273 (20-8)735 (18-4)Higher/more than secondary273 (18-2)404 (34.4)548 (41-8)1225 (30-7)Work status (past 12 months), n (%)1002 (67-8)1307 (99-5)3836 (96-2)Number with data1500 (100)1174 (99-9)1309 (99-6)3983 (99.8)Employed439 (29-3)522 (44-5)507 (38-7)1448 (68.9)Unservind (an)224 (632-1)3052 (861-4)136 (69-5)3836 (96-2)Morthy household income (USD)1477 (99-8)1302 (87-8)1307 (99-5)3836 (96-2)Morthy household income (USD)1497 (99-8)1032 (87-8)<	Duration of the SMI, n (%)				()
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	Number with data	1500 (100)	1175 (100)	1314 (100)	3989 (100)
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	≤2 years	436 (29.1)	215 (18-3)	289 (22.0)	940 (23.6)
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	3–5 years	457 (30.5)	266 (22.6)	320 (24.4)	1043 (26.1)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	6–10 years	332 (22.1)	299 (25.4)	299 (22.8)	930 (23.3)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	>10 years	271 (18.1)	359 (30.6)	399 (30.4)	1029 (25.8)
Satting, n (%)Number with data1500 (100)1175 (100)1314 (100)3989 (100)Inpatient313 (20.9)264 (22.5)122 (9.3)699 (17.5)Outpatient1187 (79.1)911 (77.5)1192 (90.7)3290 (82.5)Highest level of education, n (%)3986 (99.9)3986 (99.9)No formal education151 (10.1)141 (12.0)257 (15.6)549 (13.8)Primary education842 (56.1)401 (34.2)234 (17.8)1477 (37.1)Secondary education234 (15.6)228 (19.4)273 (20.8)733 (18.4)Higher/more than secondary273 (18.2)404 (34.4)548 (41.8)1225 (30.7)Work status (past 12 months), n (%)595 (39.7)227 (19.3)291 (22.2)1113 (27.9)Other, ⁴ 458 (32.1)426 (36.1)425 (36.2)511 (30.0)1422 (35.2)Monthy household income (USD)199.6 (199.8)3386 (96.2)m (%)1497 (99.8)1032 (87.8)1307 (99.5)3836 (96.2)Modian (ICR)176.7 (117.8-235.5)158.9 (66.2-237.8)158.9 (105.9-264.4)Income tertile, n (%)118.1,1777.20.0, 16.551.90.0, 2972.70.0, 16.551.9Income tertile, n (%)118.4 (17.7)1263 (32.9)158.9 (105.9-264.9)Income tertile, n (%)118.4 (17.7)1305 (32.7)126 (32.2)Mindia1497 (99.8)1032 (87.8)1307 (99.5)3836 (96.2)Low with da	Do not know/Don't remember	4 (0.3)	36 (3.1)	7 (0.5)	47 (1·2)
Number with data1500 (100)1175 (100)1314 (100)3989 (100)Inpatient313 (20-9)264 (22-5)122 (9.3)699 (17.5)Outpatient1187 (79-1)911 (77.5)1192 (90-7)3290 (82-5)Highest level of education, n (%)No formal education151 (10-1)141 (12.0)257 (18-6)544 (13.8)Primary education824 (56-1)404 (34-2)224 (17.6)1477 (37.1)Secondary education234 (15.6)228 (19.4)273 (20.8)735 (18.4)Higher/more than secondary273 (18.2)404 (34.4)548 (41.8)1225 (30.7)Work status (past 12 months), n (%)1174 (99.9)1309 (98.6)3983 (99.8)Employed439 (29.3)522 (44.5)507 (38.7)1468 (36.9)Unmber with data1500 (100)1174 (99.9)1309 (98.6)3883 (96.2)Orther466 (31.1)425 (362.2)511 (38.0)1402 (35.2)Monthy household income (USD) n (%)118, 11,777.20.0, 16,551.90.0, 2972.70.0, 16,551.9Income tertile, n (%)118, 11,777.20.0, 16,551.90.0, 2972.70.0, 16,551.9Income tertile, n (%)388 (22.6)220 (21.3)426 (32.6)984 (25.7)Mindial1497 (99.8)1032 (87.8)1307 (99.5)3836 (96.2)Low674 (45.0)349 (33.8)566 (43.3)158.9 (16.9)Number with data1497 (99.8)1032 (87.8)1307 (99.5)3836 (96.2)Low674 (45.0)349 (33.8)566 (43.3) <td>Setting, n (%)</td> <td></td> <td></td> <td></td> <td></td>	Setting, n (%)				
$\begin{array}{llllllllllllllllllllllllllllllllllll$	Number with data	1500 (100)	1175 (100)	1314 (100)	3989 (100)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Inpatient	313 (20.9)	264 (22.5)	122 (9.3)	699 (17.5)
Highest level of education, n (%) Number with data 1500 (100) 1174 (99-9) 1312 (99-8) 3986 (99-9) No formal education 151 (10-1) 141 (12.0) 257 (19-6) 549 (13-8) Primary education 244 (56-1) 401 (34-2) 234 (17-8) 1477 (37-1) Secondary education 234 (15-6) 228 (19-4) 273 (20-8) 735 (18-4) Higher/more than secondary 273 (18-2) 404 (34-4) 548 (41-8) 1225 (30-7) Work status (past 12 months), n (%) Number with data 1500 (100) 1174 (99-9) 1309 (99-6) 3983 (99-8) Employed 439 (29-3) 522 (44-5) 507 (38-7) 1468 (36-9) Unemployed 595 (39-7) 227 (19-3) 291 (22-2) 1113 (27-9) Other ⁴ 466 (31-1) 425 (36-2) 511 (39-0) 1402 (35-2) Monthly household income (USD) n (%) 1497 (99-8) 1032 (87-8) 1307 (99-5) 3836 (96-2) Mean (so) 24-24 (352-1) 305-2 (661-4) 199-8 (199-8) 2374 (513-1) Median (IOR) 176-7 (117-8-235-5) 158-9 (662-264-8) 148-6 (892-237-8) 158-9 (105-9-264-8) Min, Max 11-8, 11.777.2 0-0, 16,551-9 0-0, 2972.7 0-0, 16,551-9 Income terlile, n (%) Munthy data 1497 (99-8) 1032 (87-8) 1307 (99-5) 3836 (96-2) Low 674 (45-0) 349 (33-8) 566 (43-3) 1589 (105-9-264-8) Min, Max 11-8, 11.777.2 0-0, 16,551-9 0-0, 2972.7 0-0, 16,551-9 Income terlile, n (%) Munth data 1497 (99-8) 1032 (87-8) 1307 (99-5) 3838 (96-2) Low 674 (45-0) 349 (33-8) 566 (43-3) 1589 (10-9-264-8) Mindide 445 (32-4) 463 (42-9) 315 (24-1) 1263 (32-9) High 338 (22-6) 220 (21-3) 426 (32-6) 984 (25-7) Marital status, n (%) Number with data 1497 (99-8) 1032 (87-8) 1307 (99-5) 3836 (96-2) Low 674 (45-0) 349 (23-8) 1566 (43-3) 1589 (10-0) Never maried 2539 (35-9) 349 (29-7) 417 (31-7) 1305 (32-7) Currently maried 1500 (100) 1175 (100) 1314 (100) 3989 (100) Never maried 143 (9-5) 115 (9-8) 150 (11-4) 408 (10-2) EMI n (%) 1497 (99-8) 1161 (98-8) 1304 (99-2) 3962 (99-3) Underweight (BMI <18-5) 129 (8-6) 66 (5-7) 113 (8-7) 306 (7-8) Nomal weight (BMI <18-5) 129 (8-6) 66 (5-7) 113 (8-7) 306 (7-8) Nomal weight (BMI <18-5) 129 (8-6) 606 (45-7) 113 (8-7) 306 (7-8) Nomal weight (BMI <18-5) 129 (8-6) 606 (45-7) 113 (8-7) 306 (7-8) Nomal weight (BMI <18-5) 129 (8-6)	Outpatient	1187 (79.1)	911 (77·5)	1192 (90.7)	3290 (82.5)
Number with data1500 (100)1174 (199)1312 (99-8)3986 (99-9)No formal education151 (10-1)141 (12-0)257 (19-6)549 (13.8)Primary education842 (56-1)401 (34-2)234 (17-8)1477 (37-1)Secondary education234 (15-6)228 (19-4)273 (20-8)735 (18-4)Higher/more than secondary273 (18-2)404 (34-4)548 (41-8)1225 (30-7)Work status (past 12 months), n (%)number with data1500 (100)1174 (99-9)1309 (99-6)3983 (99-8)Employed439 (29-3)522 (44-5)507 (38-7)1468 (36-9)Unemployed595 (39-7)227 (19-3)291 (22-2)1113 (27-9)Other*466 (31-1)425 (36-2)511 (39-0)1402 (35-2)Monthy household income (USD)1497 (99-8)1032 (87-8)1307 (99-5)3836 (96-2)Median (ICR)176-7 (117.8-235.5)158 9 (66-2-664.8)148 6 (199.8)237.4 (513-1)Median (ICR)176-7 (117.8-235.5)158 9 (66-2-664.8)148 6 (89-2-237.8)158 9 (105.9-264.8)Income tertile, n (%)1138, 11,777.20.0, 16,551-90.0, 287.270.0, 10, 551-9Income tertile, n (%)388 (22-6)220 (21-3)426 (32-6)984 (42-7)Maridal status, n (%)1134 (100)3989 (100)1497 (99-8)1137 (105)1314 (100)Number with data1500 (100)1175 (100)1314 (100)3989 (100)Never married539 (35-9)349 (29-7)417 (31-7)1305 (32-7)<	Highest level of education, n (%)				
No formal education151 (10-1)141 (12-0)257 (19-6)549 (13-8)Primary education234 (15-6)228 (19-4)273 (20-8)735 (18-4)Higher/more than secondary273 (18-2)404 (34-4)548 (41-8)1225 (30-7)Work staus (past 12 months), n (%)1500 (100)1174 (99-9)1309 (99-6)3983 (99-8)Number with data1500 (100)1174 (99-9)1309 (99-6)3983 (99-8)Unemployed439 (23-3)522 (44-5)507 (38-7)1468 (36-9)Unemployed595 (39-7)227 (19-3)291 (22-2)1113 (27-9)Other*466 (31-1)425 (36-2)511 (39-0)1402 (35-2)Monthy household income (USD)1497 (99-8)1032 (87-8)1307 (99-5)3836 (96-2)Mean (sc)224-6 (352-1)305 2 (861-4)198 6 (199-8)237 4 (513.1)Median (IQR)176 7 (117-8-235-5)158.9 (66-2-264-8)148 6 (89-2-237.8)158.9 (105-9-264-8)Number with data1497 (99-8)1032 (87-8)1307 (99-5)3886 (96-2)Low674 (45-0)349 (33-8)566 (43-3)1589 (41-4)Middle445 (32-4)463 (14-9)315 (24-1)1263 (32-9)High338 (22-6)220 (21-3)426 (32-6)984 (25-7)Marital status, n (%)1175 (100)1314 (100)3989 (100)Number with data1500 (100)1175 (100)1314 (100)3989 (100)Never married539 (35-9)349 (25-7)417 (31-7)1305 (32-7)Low140 (3	Number with data	1500 (100)	1174 (99.9)	1312 (99.8)	3986 (99.9)
Primary Buddation 644 (36-1) 401 (34-2) 224 (17-6) 147 (37-1) Secondary education 234 (15-6) 228 (19-4) 273 (20.8) 735 (18-4) Higher/more than secondary 273 (18-2) 404 (34-4) 548 (41.8) 1225 (30-7) Work stats (past 12 months), n (%) Number with data 1500 (100) 1174 (99-9) 1309 (99-6) 3983 (99-8) Employed 439 (29-3) 522 (44-5) 507 (38-7) 1468 (36-9) Unemployed 595 (39-7) 227 (19-3) 291 (22.2) 1113 (27-9) Other ⁴ 466 (31-1) 425 (36-2) 511 (39-0) 1402 (35-2) Monthly household income (USD) n (%) 1497 (99-8) 1032 (87-8) 1307 (99-5) 3836 (96-2) Mean (so) 224-6 (352-1) 305-2 (86-1) 148-6 (98-2-37-8) 158-9 (16-2-264-8) Min, Max 11-8, 11,777-2 0.0, 16,551-9 0.0, 2972-7 0.0, 16,551-9 Income tertile, n (%) 1138 (10-5 348 (169-2-237-8) 1589 (160-2-264-8) Middle 4495 (32-4) 463 (34-9) 315 (24-1) <td< td=""><td>No formal education</td><td>151 (10-1)</td><td>141 (12·0)</td><td>257 (19.6)</td><td>549 (13.8)</td></td<>	No formal education	151 (10-1)	141 (12·0)	257 (19.6)	549 (13.8)
Secondary education 234 (15-5) 228 (19-4) 273 (20-8) 735 (16-4) Higher/more than secondary 273 (18-2) 404 (34-4) 548 (41-8) 1225 (30-7) Work status (past 12 months), n (%) 1500 (100) 1174 (99-9) 1309 (99-6) 3983 (99-8) Employed 439 (29-3) 552 (44-5) 507 (38-7) 1468 (36-9) Unemployed 595 (39-7) 227 (19-3) 291 (22-2) 1113 (27-9) Other ⁴ 466 (31-1) 425 (36-2) 511 (39-0) 1402 (35-2) Monthly household income (USD) 1497 (99-8) 1032 (87-8) 1307 (99-5) 3836 (96-2) Mean (so) 224 6 (352-1) 305 2 (861-4) 198 6 (199-8) 237 4 (513-1) Mecidan (ICR) 176 7 (117-8-235-5) 158 9 (66-2-264-8) 148 6 (89 2-237-8) 158 9 (105-9-264-8) Min, Max 11-8, 11.777-2 0.0, 16,551-9 0.0, 2972-7 0.0, 16,551-9 Income tertile, n (%) 1497 (99-8) 1032 (87-8) 1307 (99-5) 336 (96-2) Low 674 (45-0) 349 (33-8) 566 (43-3) 1589 (41-4)	Primary education	842 (56-1)	401 (34-2)	234 (17.8)	14/7 (37-1)
IngleminePart (16.2)404 (9.4)346 (41.6)1220 (30.7)Number with data1500 (100)1174 (99-9)1309 (99-6)3983 (99-8)Employed439 (29-3)522 (44-5)507 (38-7)1468 (36-9)Unemployed595 (39.7)227 (19-3)291 (22-2)1113 (27-9)Other ^a 466 (31-1)425 (36-2)511 (39-0)1402 (35-2)Monthly household income (USD)n (%)1497 (99-8)1032 (87-8)1307 (99-5)3836 (96-2)Mean (so)224-6 (352-1)305-2 (861-4)1886 (199-8)237-4 (513-1)Median (IQR)176 7 (117-8-235-5)158-9 (662-264-8)1486 (89-2-237-8)158-9 (105-9-264-8)Min, Max11-8, 11.777-20.0, 16,551-90.0, 2972-70.0, 16,551-9Income tertlie, n (%)Number with data1497 (99-8)1032 (87-8)1307 (99-5)3836 (96-2)Low674 (45-0)349 (33-8)566 (43-3)1589 (41-4)Middle485 (32-4)463 (44-9)315 (24-1)1263 (32-9)High388 (22-6)220 (21-3)426 (32-6)984 (25-7)Marital status, n (%)Number with data1500 (100)1175 (100)1314 (100)3989 (100)Never married ^b 143 (9-5)115 (98-1)150 (11-4)408 (10-2)BMIn (%)140 (95-5)747 (56-8)2276 (57-1)Ever married ^b 143 (9-5)115 (98-1)150 (11-4)408 (10-2)BMI140236 (21-12-26-5)25-3 (21-2-29-3)24-5 (21-2-8-1) <tr< td=""><td>Secondary education</td><td>234 (15·6) 070 (19.0)</td><td>228 (19.4)</td><td>273 (20·8) E49 (41 9)</td><td>735 (18·4) 1005 (20 7)</td></tr<>	Secondary education	234 (15·6) 070 (19.0)	228 (19.4)	273 (20·8) E49 (41 9)	735 (18·4) 1005 (20 7)
Number with data1500 (100)1174 (99-9)1309 (99-6)3983 (99-8)Employed439 (29-3)522 (44-5)507 (38-7)1468 (36-9)Unemployed595 (39-7)227 (19-3)291 (22.2)1113 (27-9)Other*466 (31-1)425 (36-2)511 (39-0)1402 (35.2)Monthy household income (USD)n (%)1497 (99-8)1032 (87-8)1307 (99-5)3836 (96-2)Mean (sc)224-6 (352-1)305-2 (861-4)198-6 (199-8)237-4 (513.1)Median (ICR)176-7 (117-8-235-5)158-9 (662-264-8)148-6 (89-2-237-8)158-9 (105-9-264-8)Nin, Max11-8, 11,777-20-0, 16,551-90-0, 2972-70-0, 16,551-9Income tertile, n (%)111497 (99-8)1032 (87-8)1307 (99-5)3836 (96-2)Number with data1497 (99-8)1032 (87-8)1307 (99-5)3836 (96-2)1000Low674 (45-0)349 (33-8)566 (43-3)1589 (103-9)Mindle485 (32-4)463 (44-9)315 (24-1)1263 (32-9)High338 (22-6)220 (21-3)426 (32-6)984 (25-7)Marital status, n (%)11111Never married510 (11-0)1314 (100)3989 (100)Never married1497 (99-8)1161 (98-8)1304 (99-2)3962 (99-3)Median (ICR)1497 (99-8)1161 (98-8)1304 (99-2)3962 (99-3)Marital status, n (%)Number with data1497 (99-8)<	Mork status (past 12 months) n (%)	273 (10-2)	404 (34.4)	546 (41.0)	1225 (30-7)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Number with data	1500 (100)	1174 (00 0)	1300 (00 6)	3083 (00 8)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Employed	/30 (20.3)	522 (11.5)	507 (38.7)	1/68 (36.9)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Linemployed	595 (39.7)	227 (19.3)	291 (22.2)	1113 (27.9)
Monthly household income (USD)1497 (99.8)1032 (87.8)1307 (99.5)3836 (96.2)Mean (sc)224.6 (352.1)305.2 (861.4)198.6 (199.8)237.4 (513.1)Median (IQR)176.7 (117.8-235.5)158.9 (66.2-264.8)148.6 (89.2-237.8)158.9 (105.9-264.8)Number with data1497 (99.8)1032 (87.8)1307 (99.5)3836 (96.2)Low674 (45.0)349 (33.8)566 (43.3)1589 (41.4)Middle445 (32.4)463 (44.9)315 (24.1)1263 (32.9)High338 (22.6)220 (21.3)426 (32.6)984 (25.7)Marital status, n (%)11500 (100)1175 (100)1314 (100)3989 (100)Never married539 (35.9)349 (29.7)417 (31.7)1305 (32.7)Currently married818 (54.5)711 (60.5)747 (56.8)2276 (57.1)Ever married ¹⁰ 1437 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Mean (sc)24.0 (4.5)25.6 (5.3)25.9 (6.1)25.1 (5.4)Moital status, n (%)1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Median (IQR)23.6 (21.1-26.5)25.0 (22.1-28.5)25.3 (21.8-29.3)24.5 (21.6-28.1)Min, Max1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Median (IQR)23.6 (21.1-26.5)25.0 (22.1-28.5)25.3 (21.8-29.3)24.5 (21.6-28.1)Min, Max1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Underweight (BMI 18-5-4)129 (8-6)506 (43.6)501 (38.4)12	Other ^a	466 (31.1)	425 (36.2)	511 (39.0)	1402 (35.2)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Monthly household income (USD)		120 (00 2)		1102 (00 2)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	n (%)	1497 (99.8)	1032 (87.8)	1307 (99.5)	3836 (96.2)
Median (IQR)176-7 (117.8–235.5)158-9 (66-2–264.8)148-6 (89-2–237.8)158-9 (105.9–264.8)Min, Max11.8, 11,777.20.0, 16,551.90.0, 2972.70.0, 16,551.9Income tertile, n (%)0.001497 (99.8)1032 (87.8)1307 (99.5)3836 (96.2)Low674 (45.0)349 (33.8)566 (43.3)1589 (41.4)Middle485 (32.4)463 (44.9)315 (24.1)1263 (32.9)High338 (22.6)220 (21.3)426 (32.6)984 (25.7)Marital status, n (%)11001175 (100)1314 (100)3989 (100)Number with data1500 (100)1175 (100)1314 (100)3989 (100)Never married539 (35.9)349 (29.7)417 (31.7)1305 (32.7)Currently married818 (54.5)711 (60.5)747 (56.8)2276 (57.1)Ever married ^b 143 (9.5)115 (9.8)150 (11.4)408 (10.2)BMIn (%)1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Mean (sc)24.0 (4.5)25.6 (5.3)25.9 (6.1)25.1 (5.4)Min, Max14.0, 83.813.0, 96.99.7, 65.59.7, 96.9BMI (WHO international classification), n (%)1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Normal weight (BMI 18.5-24.9)817 (54.6)506 (43.6)501 (38.4)1824 (46.0)Overweight (BMI 25.0-29.9)405 (27.1)384 (33.1)406 (31.1)1195 (30.2)Obesity (BMI 25.0-29.9)405 (27.1)384 (33.1)406 (31.1)1195	Mean (sd)	224.6 (352.1)	305.2 (861.4)	198.6 (199.8)	237.4 (513.1)
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	Median (IQR)	176.7 (117.8–235.5)	158.9 (66.2–264.8)	148.6 (89.2–237.8)	158.9 (105.9–264.8)
Income tertile, n (%)Number with data1497 (99-8)1032 (87-8)1307 (99-5)3836 (96-2)Low674 (45-0)349 (33-8)566 (43-3)1589 (41-4)Middle485 (32-4)463 (44-9)315 (24-1)1263 (32-9)High338 (22-6)220 (21-3)426 (32-6)984 (25-7)Marital status, n (%)1175 (100)1314 (100)3989 (100)Never married539 (35-9)349 (29-7)417 (31-7)1305 (32-7)Currently married818 (54-5)711 (60-5)747 (56-8)227 (57-1)Ever married ^b 143 (9-5)115 (9-8)150 (11-4)408 (10-2)BMI1497 (99-8)1161 (98-8)1304 (99-2)3962 (99-3)Mean (sc)24-0 (4-5)25-6 (5-3)25-9 (6-1)25-1 (5-4)Min, Max14-0, 83-813-0, 96-99-7, 65-59-7, 96-9BMI (WHO international classification), n (%)1497 (99-8)1161 (98-8)1304 (99-2)3962 (99-3)Number with data1497 (99-8)1161 (98-8)1304 (99-2)3962 (99-3)Underweight (BMI 18-5)129 (86)66 (5-7)113 (8-7)308 (7-8)Normal weight (BMI 18-5)129 (8-6)66 (5-7)113 (8-7)308 (7-8)Normal weight (BMI 18-5-24-9)817 (54-6)506 (43-6)501 (38-4)1824 (46-0)Overweight (BMI 25-0-29-9)405 (27-1)384 (33-1)406 (31-1)1195 (30-2)Obesity (BMI 230-0)146 (9-8)205 (17-7)284 (21-8) <t< td=""><td>Min, Max</td><td>11.8, 11,777.2</td><td>0.0, 16,551.9</td><td>0.0, 2972.7</td><td>0.0, 16,551.9</td></t<>	Min, Max	11.8, 11,777.2	0.0, 16,551.9	0.0, 2972.7	0.0, 16,551.9
Number with data1497 (99-8)1032 (87-8)1307 (99-5)3836 (96-2)Low674 (45-0)349 (33.8)566 (43.3)1589 (41.4)Middle485 (32.4)463 (44-9)315 (24.1)1263 (32.9)High338 (22-6)220 (21.3)426 (32-6)984 (25-7)Marital status, n (%) </td <td>Income tertile, n (%)</td> <td></td> <td></td> <td></td> <td></td>	Income tertile, n (%)				
Low674 (45·0)349 (33·8)566 (43·3)1589 (41·4)Middle485 (32·4)463 (44·9)315 (24·1)1263 (32·9)High338 (22·6)220 (21·3)426 (32·6)984 (25·7)Marital status, n (%)Number with data1500 (100)1175 (100)1314 (100)3989 (100)Never married539 (35·9)349 (29·7)417 (31·7)1305 (32·7)Currently married818 (54·5)711 (60·5)747 (56·8)2276 (57·1)Ever married ^b 143 (9·5)115 (9·8)150 (11·4)408 (10·2)BMIn (%)1497 (99·8)1161 (98·8)1304 (99·2)3962 (99·3)Median (IQR)23·6 (21·1-26·5)25·0 (22·1-28·5)25·3 (21·8-29·3)24·5 (21·6-28·1)Min, Max14·0, 83·813·0, 96·99·7, 65·59·7, 96·9BMI (WHO international classification), n (%)11497 (99·8)1161 (98·8)1304 (99·2)3962 (99·3)Number with data1497 (99·8)1161 (98·8)1304 (99·2)3962 (99·3)Underweight (BMI <18·5)	Number with data	1497 (99.8)	1032 (87.8)	1307 (99.5)	3836 (96.2)
Middle485 (32.4)463 (44.9)315 (24.1)1263 (32.9)High338 (22.6)220 (21.3)426 (32.6)984 (25.7)Marital status, n (%)Number with data1500 (100)1175 (100)1314 (100)3989 (100)Never married539 (35.9)349 (29.7)417 (31.7)1305 (32.7)Currently married818 (54.5)711 (60.5)747 (56.8)2276 (57.1)Ever married ^b 143 (9.5)115 (9.8)150 (11.4)408 (10.2)BMIn (%)24.0 (4.5)25.6 (5.3)25.9 (6.1)25.1 (5.4)Median (IQR)23.6 (21.1–26.5)25.0 (22.1–28.5)25.3 (21.8–29.3)24.5 (21.6–28.1)Min, Max14.0, 83.813.0, 96.99.7, 65.59.7, 96.9BMI (WHO international classification), n (%)1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Number with data1497 (99.8)1161 (98.8)1304 (99.2)3962 (29.3)Underweight (BMI <18.5)	Low	674 (45·0)	349 (33.8)	566 (43·3)	1589 (41.4)
High338 (22-6)220 (21-3)426 (32-6)984 (25-7)Marital status, n (%)Number with data1500 (100)1175 (100)1314 (100)3989 (100)Never married539 (35-9)349 (29-7)417 (31-7)1305 (32-7)Currently married818 (54-5)711 (60-5)747 (56-8)2276 (57-1)Ever married ^b 143 (9-5)115 (9-8)150 (11-4)408 (10-2)BMI n (%)24-0 (4-5)25-6 (5-3)25-9 (6-1)25-1 (5-4)Mean (sb)23-6 (21-1-26-5)25-0 (22-1-28-5)25-3 (21-8-29-3)24-5 (21-6-28-1)Min, Max1497 (99-8)1161 (98-8)1304 (99-2)3962 (99-3)BMI (WHO international classification), n (%)Number with data1497 (99-8)1161 (98-8)1304 (99-2)3962 (99-3)Underweight (BMI <18-5)	Middle	485 (32.4)	463 (44·9)	315 (24.1)	1263 (32.9)
Marital status, n (%)Number with data1500 (100)1175 (100)1314 (100)3989 (100)Never married539 (35-9)349 (29-7)417 (31-7)1305 (32-7)Currently married818 (54-5)711 (60-5)747 (56-8)2276 (57-1)Ever married ^b 143 (9-5)115 (9-8)150 (11-4)408 (10-2)BMI </td <td>High</td> <td>338 (22.6)</td> <td>220 (21.3)</td> <td>426 (32.6)</td> <td>984 (25.7)</td>	High	338 (22.6)	220 (21.3)	426 (32.6)	984 (25.7)
Number with data1500 (100)1175 (100)1314 (100)3989 (100)Never married539 (35.9)349 (29.7)417 (31.7)1305 (32.7)Currently married818 (54.5)711 (60.5)747 (56.8)2276 (57.1)Ever married ^b 143 (9.5)115 (9.8)150 (11.4)408 (10.2)BMI1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Mean (sD)24.0 (4.5)25.6 (5.3)25.9 (6.1)25.1 (5.4)Median (IQR)23.6 (21.1–26.5)25.0 (22.1–28.5)25.3 (21.8–29.3)24.5 (21.6–28.1)Min, Max14.0, 83.813.0, 96.99.7, 65.59.7, 96.9BMI (WHO international classification), n (%)129 (8.6)66 (5.7)113 (8.7)308 (7.8)Number with data1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Underweight (BMI <18.5)	Marital status, n (%)				
Never married539 (35-9) $349 (29 \cdot I)$ $417 (31 \cdot I)$ $1305 (32 \cdot I)$ Currently married818 (54-5)711 (60-5)747 (56-8)2276 (57-1)Ever married ^b 143 (9-5)115 (9-8)150 (11-4)408 (10-2)BMIn (%)1497 (99-8)1161 (98-8)1304 (99-2)3962 (99-3)Mean (sD)24-0 (4-5)25-6 (5-3)25-9 (6-1)25-1 (5-4)Median (IQR)23-6 (21-1-26-5)25-0 (22-1-28-5)25-3 (21-8-29-3)24-5 (21-6-28-1)Min, Max14-0, 83-813-0, 96-99-7, 65-59-7, 96-9BMI (WHO international classification), n (%)129 (8-6)66 (5-7)113 (8-7)308 (7-8)Number with data1497 (99-8)1161 (98-8)1304 (99-2)3962 (99-3)Underweight (BMI <18-5)	Number with data	1500 (100)	1175 (100)	1314 (100)	3989 (100)
Currently married818 (54-5)/11 (60-5)/47 (56-8)2276 (57-1)Ever married ^b 143 (9-5)115 (9-8)150 (11-4)408 (10-2)BMI </td <td>Never married</td> <td>539 (35.9)</td> <td>349 (29.7)</td> <td>417 (31.7)</td> <td>1305 (32.7)</td>	Never married	539 (35.9)	349 (29.7)	417 (31.7)	1305 (32.7)
Ever married143 (9·5)115 (9·8)150 (11·4)408 (10·2)BMI n (%)1497 (99·8)1161 (98·8)1304 (99·2)3962 (99·3)Mean (sD)24·0 (4·5)25·6 (5·3)25·9 (6·1)25·1 (5·4)Median (IQR)23·6 (21·1–26·5)25·0 (22·1–28·5)25·3 (21·8–29·3)24·5 (21·6–28·1)Min, Max14·0, 83·813·0, 96·99·7, 65·59·7, 96·9BMI (WHO international classification), n (%)Number with data1497 (99·8)1161 (98·8)1304 (99·2)3962 (99·3)Underweight (BMI <18·5)	Currently married	818 (54.5)	/11 (60·5)	747 (56·8)	22/6 (57-1)
DNII1497 (99⋅8)1161 (98⋅8)1304 (99⋅2)3962 (99⋅3)Mean (sD)24⋅0 (4⋅5)25⋅6 (5⋅3)25⋅9 (6⋅1)25⋅1 (5⋅4)Median (IQR)23⋅6 (21⋅1-26⋅5)25⋅0 (22⋅1-28⋅5)25⋅3 (21⋅8-29⋅3)24⋅5 (21⋅6-28⋅1)Min, Max14⋅0, 83⋅813⋅0, 96⋅99⋅7, 65⋅59⋅7, 96⋅9BMI (WHO international classification), n (%)1497 (99⋅8)1161 (98⋅8)1304 (99⋅2)3962 (99⋅3)Number with data1497 (99⋅8)1161 (98⋅8)1304 (99⋅2)3962 (99⋅3)Underweight (BMI <18⋅5)	Ever married ⁻	143 (9.5)	115 (9.8)	150 (11.4)	408 (10-2)
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	BIMI m (9()	1407 (00 0)		1004 (00 0)	
Metal (SD) $24.0 (4.5)$ $25.0 (5.5)$ $25.9 (6.1)$ $25.1 (5.4)$ Median (IQR) $23.6 (21.1-26.5)$ $25.0 (22.1-28.5)$ $25.3 (21.8-29.3)$ $24.5 (21.6-28.1)$ Min, Max14.0, 83.813.0, 96.99.7, 65.59.7, 96.9BMI (WHO international classification), n (%)1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Underweight (BMI <18.5)	// (%) Moan (an)	1497 (99·6) 24 0 (4 5)	1101 (90·0) 25 6 (5 2)	1304 (99·2) 25 0 (6 1)	3902 (99·3) 25 1 (5 4)
Michail (GN)256 (211-205)256 (221-205)256 (216-256)246 (216-256)Min, Max140, 83.813.0, 96.99.7, 65.59.7, 96.9BMI (WHO international classification), n (%)1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Underweight (BMI <18.5)	Modian (IOP)	24.0 (4.3)	25.0 (3.3)	25.3 (0.1)	20.1 (0.4)
Introduct14-0, 60-010-0, 50-5 $5 \cdot 7, 65 \cdot 5$ $9 \cdot 7, 96 \cdot 9$ BMI (WHO international classification), n (%)Number with data1497 (99 \cdot 8)1161 (98 \cdot 8)1304 (99 \cdot 2)3962 (99 \cdot 3)Underweight (BMI <18 \cdot 5)	Min May	14.0 83.8	13.0 06.0	9.7 65.5	9.7 QR.Q
Number with data1497 (99.8)1161 (98.8)1304 (99.2)3962 (99.3)Underweight (BMI <18.5)	BMI (WHO international classification) n (%)	14.0,00.0	10.0, 90.9	3.1,00.0	3.1, 20.2
Underweight (BMI <18.5)129 (8.6)66 (5.7)113 (8.7)308 (7.8)Normal weight (BMI 18.5–24.9)817 (54.6)506 (43.6)501 (38.4)1824 (46.0)Overweight (BMI 25.0–29.9)405 (27.1)384 (33.1)406 (31.1)1195 (30.2)Obesity (BMI ≥30.0)146 (9.8)205 (17.7)284 (21.8)635 (16.0)	Number with data	1497 (99.8)	1161 (98-8)	1304 (99.2)	3962 (99.3)
Normal weight (BMI 18.5–24.9) $817 (54.6)$ $506 (37.6)$ $110 (67.7)$ $506 (7.6)$ Overweight (BMI 25.0–29.9) $405 (27.1)$ $384 (33.1)$ $406 (31.1)$ $1195 (30.2)$ Obesity (BMI $\geq 30.0)$ $146 (9.8)$ $205 (17.7)$ $284 (21.8)$ $635 (16.0)$	Underweight (BMI <18.5)	129 (8.6)	66 (5.7)	113 (8.7)	308 (7.8)
Overweight (BMI 25:0-29:9) $405 (27:1)$ $384 (33:1)$ $406 (31:1)$ $1195 (30:2)$ Obesity (BMI $\geq 30.0)$ 146 (9:8) $205 (17:7)$ 284 (21:8) $635 (16:0)$	Normal weight (BMI 18-5–24-9)	817 (54.6)	506 (43.6)	501 (38.4)	1824 (46.0)
Obesity (BMI ≥30·0) 146 (9·8) 205 (17·7) 284 (21·8) 635 (16·0)	Overweight (BMI 25·0–29·9)	405 (27.1)	384 (33.1)	406 (31.1)	1195 (30.2)
	Obesity (BMI ≥30.0)	146 (9.8)	205 (17.7)	284 (21.8)	635 (16.0)
Continued	- · · ·	. ,	. ,	. /	Continued

cu

5

Table 1. Continued



	Bangladesh (n 1500)	India (<i>n</i> 1175)	Pakistan (<i>n</i> 1314)	Overall (<i>n</i> 3989)
BMI (WHO Asian cut-offs), n (%)				
Number with data	1497 (99.8)	1161 (98.8)	1304 (99.2)	3962 (99.3)
Underweight (BMI <18.5)	129 (8.6)	66 (5.7)	113 (8.7)	308 (7.8)
Normal weight (BMI 18-5-22-9)	523 (34.9)	290 (25.0)	327 (25.1)	1140 (28.8)
Overweight (BMI 23-0-24-9)	294 (19.6)	216 (18.6)	174 (13.3)	684 (17.3)
Obesity (BMI ≥25.0)	551 (36.8)	589 (50.7)	690 (52.9)	1830 (46-2)
Waist circumference (cm)				
Males				
n (% of males)	915 (100)	640 (98.8)	786 (98.7)	2341 (99.2)
Mean (sp)	83.3 (11.2)	90.3 (12.3)	89.9 (16.8)	87.4 (14.0)
Median (IQR)	84.5 (75.8–90.5)	90.0 (82.0-98.0)	89.2 (81.0-99.0)	87.5 (79.2–95.0)
Min, Max	35.5, 163.3	50.6, 140.0	30.3, 191.2	30.3, 191.2
Females				
n (% of females)	582 (99.5)	516 (97·9)	507 (97.9)	1605 (98.5)
Mean (sd)	84.3 (12.2)	87.2 (13.2)	90.6 (22.2)	87.2 (16.5)
Median (IQR)	85.5 (76.5–91.5)	87.5 (79.0–95.3)	92.0 (78.2-105.0)	87.6 (78.0–96.5)
Min, Max	46.5, 174.5	40.0, 130.2	30.0, 193.2	30.0, 193.2
Abdominal obesity, n (%)				
Males				
Number with data	915 (100)	640 (98.8)	786 (98.7)	2341 (99.2)
No abdominal obesity	666 (72.8)	300 (46.9)	398 (50.4)	1364 (58.3)
Has abdominal obesity	249 (27.2)	340 (53.1)	388 (49.4)	977 (41.7)
Females				
Number with data	582 (99.5)	516 (97·9)	507 (97.9)	1605 (98.5)
No abdominal obesity	189 (32.5)	138 (26.7)	133 (26.2)	460 (28.7)
Has abdominal obesity	393 (67.5)	378 (73.3)	374 (73.8)	1145 (71.3)
Males and Females				
Number with data	1497 (99.8)	1156 (98.4)	1293 (98.4)	3946 (98.9)
No abdominal obesity	855 (57.1)	438 (37.9)	531 (41.1)	1824 (46·2)
Has abdominal obesity	642 (42·9)	718 (62.1)	762 (58.9)	2122 (53.8)

^a Other includes: homemaker, student, and retired.

^b Ever married includes: widowed, separated, and divorced. sp, Standard deviation. IQR, Interquartile range.

obesity increased with longer SMI duration; however, the relative risk of having overweight remained consistent with increasing SMI duration. In participants who were taking antipsychotic medication, the relative risk of having obesity was nearly twice that of participants who were not on medication (RRR = 1.93 [95 % CI: 1.17-3.19]). Participants with higher education, and those in the highest income tertile had the greatest relative risk of obesity compared to no formal education (RRR = 1.51 [95 % CI: 1.12-2.05]) and the lowest income tertile (RRR = 1.38 [95 % CI: 1.10-1.72]), respectively.

The relative risk of having obesity in current smokers was lower (RRR = 0.79 [95 % CI: 0.65–0.96]) than in nonsmokers, but the relative risk of being overweight did not differ by smoking status (Table 4b). Participants meeting WHO recommendations for physical activity had 21 % lower risk of having obesity than the less physically active group (RRR = 0.79 [95 % CI: 0.66–0.94]). In contrast, participants who met the WHO recommendations for fruit and vegetable intake had 2.53 (95 % CI: 1.65–3.88) times greater risk of having obesity compared to those not meeting the recommendations.

Participants with pre-diabetes, type 2 diabetes, hypertension, high cholesterol and high triglycerides all had an increased relative risk of having obesity compared to normal weight (Table 4b). The largest relative risk ratio for obesity was seen in participants with hypertriglyceridaemia (RRR = 1.94 [95 % CI: 1.62-2.33]), and this was the only comorbidity

for which the relative risk ratio of being overweight was also significant.

In separate models, interaction terms between country and each variable were included. Only age group (P = 0.002) and high triglycerides (P = 0.048) were identified as having significant interactions with country.

Association of determinants with abdominal obesity

As seen in Table 5 the odds of having abdominal obesity were 3.79 (95 % CI: 2.99-4.80) higher in women compared with men, and the odds increased with increasing age. With longer SMI duration, there were greater odds of having abdominal obesity, and in those on antipsychotic medication, there was 1.64 (95 % CI: 1.05-2.56) times greater odds of having obesity. Participants in the highest income tertile had 1.26 (95 % CI: 1.04–1.54) times greater odds of having abdominal obesity compared to those in the lowest tertile. Both smoking (OR = 0.89 [95 % CI: 0.75 - 1.06]) and physical activity (OR = 1.03[95 % CI: 0.88-1.20]) did not significantly affect the odds of having abdominal obesity in this population; however, participants who ate at least 5 portions of fruit or vegetables per day had 2.35 (95 % CI: 1.65-3.36) times greater odds of abdominal obesity. Participants with type 2 diabetes, hypertension, high cholesterol, and high triglycerides all had higher odds of having abdominal obesity, while no association was found with pre-diabetes.



Table 2. Overall prevalence (using WHO international and WHO Asian cut-offs) and stratified prevalence (using WHO Asian cut-offs) of normal weight, overweight and obesity for all countries (Bangladesh, India, Pakistan)

	All countries (n 3962)					
Frequency prevalence (95 % CI)	Prevalence of normal weight	Prevalence of overweight	Prevalence of obesity			
Overall (WHO Asian cut-offs)	1140/3962	684/3962	1830/3962			
	28.8 (27.4–30.2)	17.3 (16.1–18.5)	46.2 (44.6–47.7)			
Overall (WHO international cut-offs)	1824/3962	1195/3962	635/3962			
	46.0 (44.5–47.6)	30.2 (28.8–31.6)	16.0 (14.9–17.2)			
Stratified prevalence (WHO Asian cut-offs):			· · · ·			
Sex						
Male	739/2349	460/2349	958/2349			
	31.5 (29.6–33.4)	19.6 (18.0–21.2)	40.8 (38.8-42.8)			
Female	401/1613	224/1613	872/1613			
	24.9 (22.8–27.0)	13.9 (12.3–15.7)	54.1 (51.6–56.5)			
Age group	, , , , , , , , , , , , , , , , , , ,					
18–24 years	325/710	113/710	188/710			
	45.8 (42.1–49.5)	15.9 (13.4–18.8)	26.5 (23.4–29.8)			
25–39 years	497/1861	339/1861	891/1861			
	26.7 (24.7–28.8)	18.2 (16.5–20.0)	47.9 (45.6–50.2)			
40–54 vears	224/1044	172/1044	585/1044			
	21.5 (19.1–24.1)	16.5 (14.4–18.9)	56.0 (53.0-59.0)			
55+ vears	94/347	60/347	166/347			
	27.1 (22.7-32.0)	17.3 (13.7–21.6)	47.8 (42.6-53.1)			
SMI						
Bipolar disorder (anv)	387/1455	244/1455	742/1455			
p	26.6 (24.4–28.9)	16.8 (14.9–18.8)	51.0 (48.4–53.6)			
Maior depressive disorder with psychotic features	189/734	103/734	378/734			
· · · · · · · · · · · · · · · · · · ·	25.7 (22.7–29.0)	14.0 (11.7–16.7)	51.5 (47.9-55.1)			
Schizophrenia-type disorder	564/1773	337/1773	710/1773			
	31.8 (29.7–34.0)	19.0 (17.2–20.9)	40.0 (37.8-42.3)			
Setting						
Inpatient	259/686	113/686	257/686			
	37.8 (34.2-41.4)	16.5 (13.9–19.4)	37.5 (33.9-41.2)			
Outpatient	881/3276	571/3276	1573/3276			
	26.9 (25.4–28.4)	17.4 (16.2–18.8)	48.0 (46.3-49.7)			

For participants where BMI was calculable (n = 3962)

Additional models were fitted including an interaction term between country and each variable. When comparing to a model with no interactions, the likelihood ratio test identified variables sex (P < 0.001), age group (P = 0.002), level of education (P = 0.007), work status (P < 0.001), income (P = 0.003), and high triglycerides (P = 0.029) as having a significant interaction effect with country.

Discussion

Obesity is a major public health problem in people with SMI regardless of whether international or Asian-specific thresholds for obesity are used, especially considering its association with other chronic conditions. The prevalence of obesity varied according to SMI and the sociodemographic characteristics of participants.

The prevalence of obesity was considerably lower in Bangladesh than in India and Pakistan, which is mirrored in the general population.⁽²⁷⁾ This may be related to lower income and education in the population in Bangladesh, as explained by other studies.⁽²⁸⁾ This illustrates the complexity of the interplay between socioeconomic and physical determinants of obesity and how other factors like age may be more influential, as the mean age of people with SMI was lowest in Bangladesh.⁽²⁸⁾

Although the psychiatric inpatient setting is considered obesogenic,⁽²⁹⁾ participants in the outpatient setting were more likely to have obesity. Inpatients are more likely to have a refractory degree of SMI which leads to more severe symptoms like catatonia leading to malnutrition.⁽³⁰⁾ This absence of physical activity can lead to a reduction in bone and muscle density which has been associated with underweight.⁽³¹⁾

The higher risk of obesity in women is in line with global trends and is likely driven by socio-cultural factors such as urbanisation where there has been a clear shift in LMICs from agricultural labour to wage labour which is usually more sedentary.⁽³²⁾ The persistent disparity in male and female employment rates, however, shows that women are still more likely to be unemployed and usually in household roles which negatively affects the physical activity of women more than men.^(32,33) Clinical studies show that females on antipsychotic medication gain more weight than males.^(34,35) Furthermore, on a physiological level, women are more susceptible to weight gain due to their fat distribution and their neural responses to food-related stimuli are more positively correlated with BMI.⁽³⁶⁾

Contrary to high income countries (HICs) where the poorest are at higher risk of obesity due to poor diet and unhealthy lifestyle,⁽³⁷⁾ we found the more affluent have an increased risk of obesity. Poorer people in LMICs tend to

Table 3. Overall prevalence (using WHO international and WHO Asian cut-offs) and stratified prevalence (using WHO Asian cut-offs) of normal weight, overweight and obesity in Bangladesh, India, Pakistan^a

	Bangladesh (<i>n</i> 1497)			India (<i>n</i> 1161)			Pakistan (<i>n</i> 1304)		
Frequency prevalence (95 % Cl)	Prevalence of normal weight	Prevalence of overweight	Prevalence of obesity	Prevalence of normal weight	Prevalence of overweight	Prevalence of obesity	Prevalence of normal weight	Prevalence of overweight	Prevalence of obesity
Overall (WHO international cut-offs)	817/1497	405/1497	146/1497	506/1161	384/1161	205/1161	501/1304	406/1304	284/1304
	54·6 (52·0–57·1)	27·1 (24·9–29·4)	9·8 (8·4–11·4)	43·6 (40·8–46·5)	33·1 (30·4–35·8)	17·7 (15·6–20·0)	38·4 (35·8–41·1)	31.1 (28.7–33.7)	21.8 (19.6–24.1)
Overall (WHO Asian cut-offs)	523/1497	294/1497	551/1497	290/1161	216/1161	589/1161	327/1304	174/1304	690/1304
	34·9 (32·6–37·4)	19·6 (17·7–21·7)	36·8 (34·4–39·3)	25·0 (22·6–27·6)	18·6 (16·5–21·0)	50·7 (47·9–53·6)	25·1 (22·8–27·5)	13·3 (11·6–15·3)	52·9 (50·2–55·6)
Stratified prevalence (WHO Asian cut-offs): Sex	, , , , , , , , , , , , , , , , , , ,	, , , , , , , , , , , , , , , , , , ,	, , , , , , , , , , , , , , , , , , ,	, , , , , , , , , , , , , , , , , , ,	,	, , , , , , , , , , , , , , , , , , ,	, , , , , , , , , , , , , , , , , , ,	, , , , , , , , , , , , , , , , , , ,	,
Male	336/915	190/915	300/915	175/643	145/643	280/643	228/791	125/791	378/791
	36·7 (33·7–39·9)	20⋅8 (18⋅3–23⋅5)	32·8 (29·8–35·9)	27·2 (23·9–30·8)	22·6 (19·5–25·9)	43·5 (39·8–47·4)	28·8 (25·8–32·1)	15·8 (13·4–18·5)	47·8 (44·3–51·3)
Female	187/582	104/582	251/582	115/518	71/518	309/518	99/513	49/513	312/513
	32·1 (28·5–36·0)	17·9 (15·0–21·2)	43·1 (39·2–47·2)	22·2 (18·8–26·0)	13·7 (11·0–16·9)	59·7 (55·4–63·8)	19·3 (16·1–22·9)	9·6 (7·3–12·4)	60·8 (56·5–65·0)
Age group	. ,	. ,	. ,		. ,	. ,	. ,	. ,	. ,
18–24 years	196/432	76/432	113/432	53/120	17/120	38/120	76/158	20/158	37/158
	45·4 (40·7–50·1)	17⋅6 (14⋅3–21⋅5)	26·2 (22·2–30·5)	44·2 (35·6–53·1)	14·2 (9·0–21·6)	31.7 (24.0–40.5)	48·1 (40·4–55·9)	12·7 (8·3–18·8)	23·4 (17·5–30·6)
25–39 years	232/731	147/731	296/731	109/531	101/531	294/531	156/599	91/599	301/599
	31.7 (28.5–35.2)	20·1 (17·4–23·2)	40·5 (37·0–44·1)	20·5 (17·3–24·2)	19·0 (15·9–22·6)	55·4 (51·1–59·5)	26·0 (22·7–29·7)	15·2 (12·5–18·3)	50·3 (46·3–54·2)
40–54 years	71/263	56/263	119/263	90/382	72/382	195/382	63/399	44/399	271/399
	27·0 (22·0–32·7)	21·3 (16·8–26·7)	45·2 (39·3–51·3)	23·6 (19·6–28·1)	18·8 (15·2–23·1)	51·0 (46·0–56·0)	15·8 (12·5–19·7)	11.0 (8.3–14.5)	67·9 (63·2–72·3)
55+ years	24/71	15/71	23/71	38/128	26/128	62/128	32/148	19/148	81/148
	33·8 (23·8–45·5)	21·1 (13·2–32·1)	32·4 (22·6–44·1)	29·7 (22·4–38·2)	20·3 (14·2–28·2)	48·4 (39·9–57·1)	21·6 (15·7–29·0)	12·8 (8·3–19·3)	54·7 (46·7–62·6)
SMI									
Bipolar disorder (any)	164/487	101/487	190/487	103/434	70/434	246/434	120/534	73/534	306/534
	33·7 (29·6–38·0)	20·7 (17·4–24·6)	39·0 (34·8–43·4)	23·7 (20·0–28·0)	16∙1 (13∙0–19∙9)	56·7 (52·0–61·3)	22·5 (19·1–26·2)	13·7 (11·0–16·9)	57·3 (53·1–61·4)
Major depressive disorder with psychotic	22/76	17/76	33/76	12/63	17/63	34/63	155/595	69/595	311/595
features	28.9 (19.9–40.1)	22.4 (14.4–33.1)	43.4 (32.8–54.7)	19.0 (11.1–30.6)	27.0 (17.5–39.2)	54.0 (41.7–65.8)	26.1 (22.7–29.7)	11.6 (9.3–14.4)	52.3 (48.2–56.3)
Psychotic disorder	337/934	176/934	328/934	175/664	129/664	309/664	52/175	32/175	73/175
	36·1 (33·1–39·2)	18·8 (16·5–21·5)	35·1 (32·1–38·2)	26·4 (23·1–29·8)	19·4 (16·6–22·6)	46·5 (42·8–50·3)	29·7 (23·4–36·9)	18·3 (13·2–24·7)	41·7 (34·6–49·2)
Setting									
Inpatient	147/311	59/311	80/311	72/255	41/255	119/255	40/120	13/120	58/120
	47·3 (41·8–52·8)	19·0 (15·0–23·7)	25·7 (21·2–30·9)	28·2 (23·0–34·1)	16·1 (12·1–21·1)	46·7 (40·6–52·8)	33·3 (25·5–42·2)	10·8 (6·4–17·8)	48·3 (39·5–57·2)
Outpatient	376/1186	235/1186	471/1186	218/906	175/906	470/906	287/1184	161/1184	632/1184
	31.7 (29.1–34.4)	19·8 (17·6–22·2)	39·7 (37·0–42·5)	24·1 (21·4–27·0)	19·3 (16·9–22·0)	51.9 (48.6–55.1)	24·2 (21·9–26·8)	13·6 (11·8–15·7)	53·4 (50·5–56·2)

^a For participants where BMI was calculable (n = 3962 in total).

Table 4a. Adjusted associations of sociodemographic variables with overweight and obesity, using WHO Asian cut-offs

	Normal weight ^b (<i>n</i> 1017)	Overweight ^b (<i>n</i> 566)	Obesity ^b (<i>n</i> 1543)	Overweight <i>v</i> . normal weight RRR (95 % Cl) <i>P</i> -value	Obese <i>v.</i> normal weight RRR (95 % Cl) <i>P</i> -value
Sex					
Male	668/1860 (35.9)	380/1860 (20.4)	812/1860 (43.7)	Beference	Beference
Female	349/1266 (27.6)	186/1266 (14.7)	731/1266 (57.7)	1.09 (0.78–1.51) <i>P</i> =0.611	2.04 (1.56 - 2.67) P < 0.001
Age group		,			(,
18–24 vears	299/567 (52.7)	100/567 (17.6)	168/567 (29.6)	Reference	Reference
25–39 years	452/1484 (30.5)	284/1484 (19.1)	748/1484 (50.4)	1.73 (1.29–2.32) P<0.001	2.36 (1.84–3.02) P<0.001
40–54 years	188/811 (23.2)	138/811 (17.0)	485/811 (59.8)	2.05 (1.43–2.95) P<0.001	2.91 (2.15–3.92) P<0.001
55+ years	78/264 (29.5)	44/264 (16.7)	142/264 (53.8)	1.59(0.96-2.61)P=0.070	1.63(1.09-2.44)P=0.018
SMI diagnosis		· · ·		· · · ·	, ,
Bipolar disorder	346/1171 (29.5)	201/1171 (17.2)	624/1171 (53.3)	Reference	Reference
Major depressive disorder with psychotic features	180/609 (29.6)	91/609 (14.9)	338/609 (55.5)	0.95 (0.67–1.34) P=0.755	0.81 (0.63–1.06) P=0.130
Schizophrenia-type disorder	491/1346 (36.5)	274/1346 (20.4)	581/1346 (43.2)	0.91 (0.71–1.17) P=0.462	0.69 (0.57–0.85) <i>P</i> < 0.001
SMI duration					
<2 years	297/720 (41.3)	120/720 (16.7)	303/720 (42.1)	Reference	Reference
3–5 years	296/832 (35.6)	170/832 (20.4)	366/8s32 (44·0)	1.32 (0.99–1.77) P=0.062	1.14 (0.90–1.45) <i>P</i> =0.285
6–10 years	232/755 (30.7)	146/755 (19.3)	377/755 (49.9)	1.33 (0.97–1.81) <i>P</i> =0.076	1.30 (1.01–1.68) <i>P</i> =0.044
>10 years	192/819 (23·4)	130/819 (15·9)	497/819 (60.7)	1·36 (0·97–1·91) <i>P</i> =0·077	1·76 (1·35–2·31) <i>P</i> < 0·001
Antipsychotic medication					
No	42/90 (46.7)	15/90 (16.7)	33/90 (36.7)	Reference	Reference
Yes	975/3036 (32.1)	551/3036 (18.1)	1510/3036 (49.7)	1.36 (0.73–2.51) <i>P</i> =0.333	1.93 (1.17–3.19) <i>P</i> =0.010
Setting					
Inpatient	212/486 (43.6)	84/486 (17.3)	190/486 (39.1)	Reference	Reference
Outpatient	805/2640 (30.5)	482/2640 (18.3)	1353/2640 (51.3)	1.42 (1.07–1.90) <i>P</i> =0.017	1·57 (1·24–2·00) <i>P</i> < 0·001
Level of education					
No formal education	132/423 (31.2)	71/423 (16.8)	220/423 (52.0)	Reference	Reference
Primary education	408/1173 (34·8)	239/1173 (20.4)	526/1173 (44·8)	1.15 (0.81 - 1.64) P = 0.437	1.23 (0.92–1.65) <i>P</i> =0.154
Secondary education	183/578 (31.7)	100/578 (17.3)	295/578 (51.0)	1.14 (0.76–1.70) <i>P</i> =0.533	1.44 (1.04–1.98) <i>P</i> =0.027
Higher/more than secondary	294/952 (30.9)	156/952 (16.4)	502/952 (52.7)	1.17 (0.80 - 1.72) P = 0.410	1.51 (1.12–2.05) <i>P</i> =0.007
Work status (past 12 months)					
Employed	357/1166 (30.6)	234/1166 (20.1)	575/1166 (49·3)	Reference	Reference
Unemployed	336/857 (39.2)	171/857 (20.0)	350/857 (40.8)	0.85 (0.65 - 1.11) P = 0.234	0.75 (0.59 - 0.94) P = 0.012
Other ^a	324/1103 (29.4)	161/1103 (14·6)	618/1103 (56.0)	0.82 (0.57–1.17) <i>P</i> =0.270	0.78 (0.58 - 1.04) P = 0.092
Income tertile					
Low	469/1301 (36·0)	236/1301 (18.1)	596/1301 (45.8)	Reference	Reference
Middle	316/1008 (31.3)	198/1008 (19.6)	494/1008 (49.0)	1.22 (0.96–1.56) P=0.105	1.18 (0.96–1.44) P=0.116
High	232/817 (28.4)	132/817 (16·2)	453/817 (55·4)	1.13 (0.85–1.50) <i>P</i> =0.388	1·38 (1·10–1·72) <i>P</i> =0·005
Country					
Bangladesh	492/1297 (37·9)	281/1297 (21.7)	524/1297 (40.4)	Reference	Reference
India	208/708 (29.4)	124/708 (17.5)	376/708 (53.1)	0.91 (0.68–1.23) P=0.537	1.18 (0.93–1.51) <i>P</i> =0.180
Pakistan	317/1121 (28.3)	161/1121 (14.4)	643/1121 (57.4)	0.75 (0.55 - 1.03) P = 0.074	1.25 (0.97–1.61) <i>P</i> =0.083

Parameter estimates from a multinomial logistic regression model are reported (n = 3126).

Note: Adjusted estimates extracted from a multinomial logistic regression model including the independent variables: diabetes, hypertension, high cholesterol, high triglycerides, smoking status, physical activity, diet, age group, sex, type of SMI, SMI duration, antipsychotic medication, setting, highest level of education, employment status, income tertile and country.

^a Other includes: homemaker, student, and retired.

^b Figures in brackets are descriptives which exclude the underweight population and therefore not prevalence data.

RRR, Relative risk ratio; CI, confidence interval.



Table 4b. Adjusted associations of comorbidities and health-risk behaviours with overweight and obesity, using WHO Asian cut-offs

	Normal weight ^a (<i>n</i> 1017)	Overweight ^a (<i>n</i> 566)	Obesity ^a (<i>n</i> 1543)	Overweight <i>v.</i> normal weight RRR 95 % Cl) <i>P</i> -value	Obese <i>v.</i> normal weight RRR (95 % Cl) <i>P</i> -value
Current s	moker				
No	616/1991 (30.9)	335/1991 (16.8)	1040/1991 (52.2)	Reference	Reference
Yes	401/1135 (35.3)	231/1135 (20.4)	503/1135 (44.3)	0.95 (0.75–1.20) <i>P</i> =0.650	0.79 (0.65 - 0.96) P = 0.018
Adequate	physical activity				
No	488/1613 (30.3)	260/1613 (16.1)	865/1613 (53.6)	Reference	Reference
Yes	529/1513 (35.0)	306/1513 (20.2)	678/1513 (44.8)	1.03 (0.83 - 1.28) P = 0.786	0.79 (0.66 - 0.94) P = 0.009
Meet WH	O recommendations for f	ruit/veg per day			
No	983/2961 (33·2)	535/2961 (18.1)	1443/2961 (48.7)	Reference	Reference
Yes	34/165 (20.6)	31/165 (18.8)	100/165 (60.6)	1.76 (1.05–2.93) P=0.031	2.53 (1.65–3.88) P<0.001
Pre-diabe	tes (HbA1c 5.7-6.4%)			, , , , , , , , , , , , , , , , , , ,	Ϋ́Υ,
No	808/2411 (33.5)	446/2411 (18.5)	1157/2411 (48.0)	Reference	Reference
Yes	209/715 (29.2)	120/715 (16.8)	386/715 (54.0)	0.94 (0.72–1.22) P=0.642	1.31 (1.06–1.62) P=0.012
Type 2 Di	abetes (HbA1c ≥6.5%)				
No	947/2797 (33.9)	515/2797 (18.4)	1335/2797 (47.7)	Reference	Reference
Yes	70/329 (21.3)	51/329 (15.5)	208/329 (63.2)	1.06 (0.71–1.58) P=0.791	1.55 (1.13–2.13) P=0.007
Hypertens	sion (BP ≥140/90 mmHg))			
No	911/2618 (34.8)	491/2618 (18·8)	1216/2618 (46.4)	Reference	Reference
Yes	106/508 (20.9)	75/508 (14.8)	327/508 (64.4)	1.20 (0.86–1.68) P=0.291	1.69 (1.30–2.19) P<0.001
High chol	esterol (LDL ≥1 g/l)				
No	572/1527 (37.5)	281/1527 (18.4)	674/1527 (44.1)	Reference	Reference
Yes	445/1599 (27.8)	285/1599 (17.8)	869/1599 (54.3)	1.20 (0.97–1.48) P=0.093	1.32 (1.11–1.56) P=0.002
High trigly	/cerides (≥1.8 g/l)				
No	706/1896 (37.2)	335/1896 (17.7)	855/1896 (45.1)	Reference	Reference
Yes	311/1230 (25.3)	231/1230 (18.8)	688/1230 (55.9)	1.56 (1.25–1.94) <i>P</i> < 0.001	1.94 (1.62–2.33) <i>P</i> < 0.001

Parameter estimates from a multinomial logistic regression model are reported (n = 3126).

Note: Adjusted estimates extracted from a multinomial logistic regression model including the independent variables: diabetes, hypertension, high cholesterol, high triglycerides, smoking status, physical activity, diet, age group, sex, type of SMI, SMI duration, antipsychotic medication, setting, highest level of education, employment status, income tertile and country.

^a Figures in brackets are descriptives which exclude the underweight population and therefore not prevalence data.

RRR, relative risk ratio; CI, confidence interval.

be engaged in more manual and physically demanding labour which causes increased energy expenditure.⁽³⁸⁾ Also, our study found that those with a higher level of education are more at risk of being obese compared to no formal education, which may show that food literacy does not correlate with educational attainment. Research has shown that in more developed countries, education can offset the obesogenic effects of increased wealth; however, in LMICs, no interaction was seen between these factors, and both were in fact independent and positively correlated with BMI.⁽³⁹⁾ This shows that in HICs the more affluent are more likely to purchase better quality and healthier foods, whereas in LMICs the wealthiest have more access to all foods and may gain more weight.⁽⁴⁰⁾ This is supported by the finding that participants who met WHO recommendations of fruit and vegetable intake had more than double the risk of obesity compared to those not following this guidance. It is likely that those that can afford to buy and eat more than five fruits or vegetables a day are also the people that can afford more food which increases the relative risk of obesity.⁽²⁸⁾ The three countries are highly dependent on cereal diets, in the public distribution systems cereals are available at below the market price which in turn increases the consumption of carbohydrates, especially in people from low socioeconomic status.⁽⁴¹⁾ So, although greater vegetable consumption is generally associated with better health outcomes, this should be taken in the context of the whole diet rather than its individual components. Further research should investigate all dietary components.

The results show that people with bipolar disorder are at greatest risk of obesity, which is mirrored in global literature, where the SMI subgroup with the highest prevalence of obesity is bipolar disorder.⁽⁴⁾ This is possibly because people with bipolar disorder can experience periods of severe depression which are associated with weight gain, similar to those with major depressive disorder, however they are also likely to experience obesogenic side effects from antipsychotic medication and mood stabilisers.⁽⁴²⁾

Similar to the general population, obesity was associated with higher relative risk of diabetes, hypertension, high cholesterol, and hypertriglyceridaemia. These are all considered the key features of metabolic syndrome which is associated with three times greater risk of cardiovascular disease and five times greater risk of developing type 2 diabetes,⁽²²⁾ explaining why people with SMI have a 53 % higher risk of developing cardiovascular disease.^(22,43) Obesity was associated with hyertriglyceridaemia, which is considered the hallmark of dyslipidaemia and possibly the major cause of all other lipid abnormalities seen in this BMI range.⁽⁴⁴⁾ Better screening of lipid abnormalities in people with SMI is required to identify those at risk of dyslipidaemia in this population. Also, obesity was associated with pre-diabetes, which supports the theory of obesity being a strong determinant of pre-diabetes due to the vital role of adipose tissue in systemic insulin resistance.⁽⁴⁵⁾

As in the general population, smoking is likely to decrease the risk of obesity due to the appetite suppressing effects of nicotine.^(46,47) However, chronic smoking is still considered an important modifiable risk factor with regards to the excess



Table 5. Adjusted associations of sociodemographic variables, comorbidities, and health-risk behaviours with abdominal obesity

	Has abdominal obesity	No abdominal obesity	OB (95 % CI)	<i>P</i> -value
	(// // 0/)			
Sex		1000/0000 (00.0)	D (
Male	807/2029 (39.8)	1222/2029 (60-2)		R < 0.001
	960/1360 (70-8)	400/1360 (29:4)	3.79 (2.99-4.00)	P < 0.001
18–24 years	216/642 (33.6)	426/642 (66.4)	Reference	
25–39 years	809/1598 (50.6)	789/1598 (49.4)	1.87 (1.50-2.33)	<i>P</i> < 0.001
40–54 vears	539/861 (62.6)	322/861 (37.4)	2.42 (1.86–3.16)	P<0.001
55+ vears	203/288 (70.5)	85/288 (29.5)	3.14 (2.18–4.54)	P<0.001
SMI diagnosis			- (/	
Bipolar disorder	681/1238 (55.0)	557/1238 (45.0)	Reference	
Major depressive disorder with psychotic features	390/669 (58.3)	279/669 (41.7)	0.76 (0.60-0.96)	P = 0.023
Schizophrenia-type disorder	696/1482 (47.0)	786/1482 (53.0)	0.76 (0.63-0.91)	P = 0.003
SMI duration				
<2 years	355/799 (44.4)	444/799 (55.6)	Reference	
3–5 years	440/906 (48.6)	466/906 (51.4)	1.25 (1.00–1.55)	P = 0.045
6–10 years	436/810 (53.8)	374/810 (46·2)	1.34 (1.06–1.68)	P = 0.012
>10 years	536/874 (61.3)	338/874 (38.7)	1.46 (1.15–1.85)	P = 0.002
Antipsychotic medication			- <i>i</i>	
No	43/104 (41.3)	61/104 (58.7)	Reference	
Yes	1724/3285 (52.5)	1561/3285 (47.5)	1.64 (1.05–2.56)	P = 0.030
Setting	000/507 (44.0)	005/507 (50.0)	Deference	
Inpatient	232/527 (44-0)	295/527 (56.0)		R 0.000
Oulpatient	1535/2862 (53.6)	1327/2862 (46-4)	1.27 (1.02–1.58)	P = 0.029
No formal education	286/472 (60.6)	186/472 (30 4)	Poforonco	
Primany education	610/1265 (48.2)	655/1265 (51.8)		P-0.620
Secondary education	313/624 (50.2)	311/624 (49.8)	1.12 (0.84–1.49)	P = 0.436
Higher/more than Secondary	558/1028 (54-3)	470/1028 (45.7)	1.31 (1.00–1.71)	P = 0.051
Work status (past 12 months)	000,1020 (010)	110,1020 (107)		, = 0 001
Employed	577/1243 (46.4)	666/1243 (53.6)	Reference	
Unemployed	383/968 (39.6)	585/968 (60.4)	0.83 (0.68–1.01)	P = 0.066
Other ^a	807/1178 (68.5)	371/1178 (31.5)	1.12 (0.86-1.45)	P = 0.394
Income tertile	× ,	× ,	· · · · ·	
Low	687/1439 (47.7)	752/1439 (52.3)	Reference	
Middle	578/1074 (53.8)	496/1074 (46·2)	1.20 (1.00–1.43)	P = 0.050
High	502/876 (57.3)	374/876 (42.7)	1.26 (1.04–1.54)	P=0.019
Country				
Bangladesh	604/1421 (42.5)	817/1421 (57.5)	Reference	
India	449/746 (60·2)	297/746 (39.8)	1.61 (1.30–2.00)	<i>P</i> < 0.001
Pakistan	714/1222 (58·4)	508/1222 (41.6)	1.48 (1.19–1.85)	<i>P</i> < 0.001
Current smoker			D /	
No	1212/2146 (56.5)	934/2146 (43.5)	Reference	
	555/1243 (44-7)	688/1243 (55·3)	0.89 (0.75–1.06)	P = 0.192
Adequate physical activity	004/1750 (54.0)		Deference	
NO	904/1759 (54-6)	795/1759 (45·2) 907/1620 (50 7)		D 0 745
Tes Moot WHO recommondations for fruit/yea per day	803/1630 (49-3)	827/1830 (50-7)	1.03 (0.00-1.20)	P=0.745
No	1651/3214 (51.4)	1563/3214 (48.6)	Reference	
Yes	116/175 (66.3)	59/175 (33.7)	2.35 (1.65-3.36)	P<0.001
Pre-diabetes (HbA1c 5.7–6.4%)		00,110 (00 1)	2 00 (1 00 0 00)	,
No	1364/2632 (51.8)	1268/2632 (48-2)	Reference	
Yes	403/757 (53.2)	354/757 (46.8)	1.06 (0.88-1.28)	P = 0.539
Type 2 Diabetes (HbA1c ≥6.5%)				
No	1532/3045 (50.3)	1513/3045 (49.7)	Reference	
Yes	235/344 (68.3)	109/344 (31.7)	1.33 (1.01–1.76)	P = 0.041
Hypertension (BP ≥140/90 mmHg)				
No	1404/2863 (49.0)	1459/2863 (51.0)	Reference	
Yes	363/526 (69.0)	163/526 (31.0)	1.69 (1.35–2.11)	<i>P</i> < 0.001
High cholesterol (LDL ≥1 g/l)				
No	807/1683 (48.0)	876/1683 (52.0)	Reference	_
Yes	960/1706 (56·3)	746/1706 (43.7)	1.17 (1.00–1.36)	P = 0.045
High triglycerides (≥1.8 g/l)			Defen	
NO	1016/2097 (48.5)	1081/2097 (51.5)		
res	/51/1292 (58-1)	541/1292 (41.9)	1.69 (1.44–1.98)	P<0.001

Parameter estimates from a logistic regression model are reported (n = 3389).

Note: Adjusted estimates extracted from a logistic regression model including the independent variables: diabetes, hypertension, high cholesterol, high triglycerides, smoking status, physical activity, diet, age group, sex, type of SMI, SMI duration, antipsychotic medication, setting, highest level of education, employment status, income tertile and country. ^a Other includes: homemaker, student, and retired.

OR, odds ratio. CI, confidence interval.

https://doi.org/10.1017/jns.2023.100 Published online by Cambridge University Press

mortality of people with SMI due to its impact on the cardiovascular system through atherosclerosis and the increased risk of lung cancer and chronic obstructive pulmonary disease.^(6,48) Smoking cessation interventions used in HICs have been less successful in LMICs hence the urgent need for culturally relevant interventions to be developed.⁽⁴⁹⁾

Strengths and limitations

There are limitations that deserve further attention. First, due to the cross-sectional nature of the study, it is not possible to determine the causality of the associations. Second, we found there was a lack of standardisation in the HbA1c laboratory analysis across the sites; although we used a laboratory in each country which was used in routine clinical practice. Further research is needed to investigate the complexities of HbA1c measurement and show how it affects the prevalence of diabetes across these countries. Third, there was considerable variability in the classification of SMI diagnosis across the different countries, and we found that far more people were diagnosed with major depression with psychotic features using the MINI v6.0. We cross-analysed this with the selfreported diagnoses and found that the vast majority were matching, which suggests that assessor error was unlikely. Fourth, the sample was exclusively from a tertiary centre cohort rather than from the community which may have implications for the interpretation of the results. However, a community survey would be prohibitively resource intensive; moreover, patients at these centres are likely to be similar to those in community, primary or secondary care, as they serve as 'walk in' and first point of access services, in the absence of any community mental healthcare.

Despite these limitations, the cross-sectional survey spanned three countries and recruited nearly 4000 participants providing good levels of precision for a population that is often neglected in this area of research. The study also included participants with all forms of SMI, providing evidence about the prevalence of obesity in each type of severe mental disorder. By including analyses using the Asian cut-offs for BMI, our results are easily comparable to other literature in South Asia.

Conclusion

There is a high prevalence of obesity in the SMI population in Bangladesh, India, and Pakistan. Obesity was associated with chronic disease in this population and, contrary to HICs, people with higher income and higher levels of educational attainment were at greater risk. Food literacy may not correlate with healthier dietary choices and so better dietary education should be prioritised for people with SMI from all levels of educational attainment. People with SMI and obesity could benefit from screening programmes for non-communicable diseases and context appropriate lifestyle interventions to prevent and treat obesity. We have identified the population at higher risk of obesity which provides useful information for intervention development; however, more research is required to identify key barriers for a healthy lifestyle in this population.



Supplementary material

The supplementary material for this article can be found at https://doi.org/10.1017/jns.2023.100.

Acknowledgments

The authors would like to acknowledge Abu Musa Robin, Anjuman Jum Tithi, Tanvir Arafat, Lipon Saha, Tasmia Rahman, Dr. Khaleda Islam, Dr. Mamun, Dr. Bappi, from the team in Bangladesh, Archith Krishna, Sathish Kumar, Neeta P.S, Venkatalakshmi, Bhuvneshwari L, Manjunatha S, Sobin George, Krishna Jayanthi from the team in India and Rubab Ayesha, Nida Afsheen, Najma Hayat, Zaheen Amin, and Aniqa Maryam from the team in Pakistan for conducting all the interviews and physical measurements, providing technical and managerial support. The authors also would like to thank all the participants who consented and provided their time to complete the interview.

N. S. and J. B. conceived the study. K. A., D. P., A. M., and G. Z. designed the study, K. A., D. P., A. M., V. C., A. C., R. H., K. P. M., and S. R. were involved in data curation. D. P., A. M., V. C., A. C., R. H., K. P. M., S. R., and G. Z. were all involved in data analysis. K. A., D. P., A. M., and G. Z. contributed to the original draft and all authors revised and approved the final manuscript.

This research was funded by the National Institute for Health Research (NIHR) (Grant: GHRG 17/63/130:) using UK aid from the UK Government to support global health research. The views expressed in this publication are those of the author(s) and not necessarily those of the NIHR or the UK Department of Health and Social Care.

D. S. is an expert advisor to the NICE Centre for Guidelines; the views expressed are the authors' and not those of NICE.

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the ethics committees of the Department of Health Sciences, University of York, UK (HSRGC-3/17); the Centre for Injury Prevention and Research, Bangladesh (CIPRB/ERC/ 2OI 8/003); the Institute Ethics Committee, National Institute of Mental Health and Neurosciences, India (BEH.SC.DIV 20/19); the Health Ministry Screening Committee, India (HMSC12/18); and the National Bioethics Committee, Pakistan (4-18/NBC-413/19). Written informed consent was obtained from all patients.

References

- Johnson DL. Overview of severe mental illness. Clin Psychol Rev. 1997;17(3):247–257.
- Bradshaw T & Mairs H. Obesity and serious mental III health: a critical review of the literature. *Healthc (Basel)*. 2014;2(2):166–182, https://doi.org/10.3390/healthcare2020166.
- DE Hert M, Schreurs V, Vancampfort D & Van Winkel R. Metabolic syndrome in people with schizophrenia: a review. World J. Psychiatry. 2009;8(1):15–22.
- 4. Holt R, Afzal M, Ahmad B, et al. Prevalence of overweight and obesity in people with severe mental illness: systematic review

and meta-analysis. *Front Endocrinol.* 2021;**12**:769309, https://doi. org/10.3389/fendo.2021.769309.

- DE Hert M, Correll CU, Bobes J, et al. Physical illness in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in health care. World Psychiatry. 2013;10 (1):52–77.
- Gilbody S, Peckham E, Bailey D, *et al.* Smoking cessation for people with severe mental illness (SCIMITAR+): a pragmatic randomised controlled trial. *Lancet Psychiat.* 2019;6(5):379–390.
- Teasdale SB, Ward PB, Samaras K, et al. Dietary intake of people with severe mental illness: systematic review and meta-analysis. Br J Psychiatry. 2019;214(5):251–259.
- Vancampfort D, Firth J, Schuch FB, *et al.* Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. *World Psychiatry.* 2017;**16**(3):308–315.
- 9. WHO. Obesity [Internet]. 2023. https://www.who.int/health-topics/obesity#tab=tab_1
- Powell-Wiley TM, Poirier P, Burke LE, et al. Obesity and cardiovascular disease: a scientific statement from the American Heart Association. *Circ.* 2021;143(21):e984–1010.
- Sarwer DB & Polonsky HM. The psychosocial burden of obesity. *Endocrinol Metab Clin North Am.* 2016;45(3):677–688.
- Ford ND, Patel SA & Narayan KMV. Obesity in Low- and middle-income countries: burden, drivers, and emerging challenges. *Annu Rev Public Health.* 2017;20(38):145–164.
- Popkin BM, Adair LS & Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev.* 2012;70 (1):3–21.
- Rathod S, Pinninti N, Irfan M, et al. Mental health service provision in Low- and middle-income countries. *Health Serv Insights.* 2017;10:1178632917694350, https://doi.org/10.1177/ 1178632917694350.
- WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet.* 2004;**363**(9403):157–163.
- Misra A. Ethnic-Specific criteria for classification of body mass index: a perspective for Asian Indians and American diabetes association position statement. *Diabetes Technol Ther.* 2015;**17**(9):667–671.
- Asia-Pacific Steering Committee. The Asia-Pacific Perspective: Redefining Obesity and Its Treatment. Sydney: Western-Pacific Region: Health Communications Australia; 2000.
- Zavala GA, Prasad-Muliyala K, Aslam F, *et al.* Prevalence of physical health conditions and health risk behaviours in people with severe mental illness in South Asia: protocol for a cross-sectional study (IMPACT SMI survey). *BMJ Open.* 2020;10(10):e037869, https://doi.org/10.1136/bmjopen-2020-037869.
- Lecrubier Y, Sheehan DV, Weiller E, *et al.* The Mini International Neuropsychiatric Interview (MINI). a short diagnostic structured interview: reliability and validity according to the CIDI. *Eur Psychiatry.* 1997;**12**:224–231.
- Bonita R, Winkelmann R, Douglas KA & de Courten M. The WHO stepwise approach to surveillance (steps) of noncommunicable disease risk factors. In: McQueen DV, and Puska P, eds. *Global Behavioral Risk Factor Surveillance*. Boston, MA: Springer US; 2003:9–22.
- World Health Organization. WHO STEPS Surveillance Manual.: The WHO STEPwise Approach to Chronic Disease Risk Factor Surveillance. Geneva: World Health Organization; 2005.
- Alberti KGMM, Zimmet P & Shaw J. Metabolic syndrome a new world-wide definition. a consensus statement from the International Diabetes Federation. *Diabet Med.* 2006;23(5):469–480.
- Basit A, Fawwad A, Qureshi H, Shera AS & NDSP Members. Prevalence of diabetes, pre-diabetes and associated risk factors: second national diabetes survey of Pakistan (NDSP), 2016–2017. *BMJ Open.* 2018;8(8):e020961, https://doi.org/10.1136/bmjopen-2017-020961.
- 24. Mendis S. Global Status Report on Noncommunicable Diseases 2014. Geneva: World Health Organization; 2015.

- World Health Organization. Global Recommendations on Physical Activity for Health. Switzerland: World Health Organization; 2010.
- von Elm E, Altman DG, Egger M, *et al.* Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Br Med J.* 2007;335 (7624):806–808.
- Zaman MM, Rahman MM, Rahman MR, Bhuiyan MR, Karim MN & Chowdhury MAJ. Prevalence of risk factors for noncommunicable diseases in Bangladesh: results from STEPS survey 2010. *Indian J Public Health*. 2016;60(1):17–25.
- Biswas T, Garnett SP, Pervin S & Rawal LB. The prevalence of underweight, overweight and obesity in Bangladeshi adults: data from a national survey. *PLoS One*. 2017;**12**(5):e0177395, https:// doi.org/10.1371/journal.pone.0177395.
- Faulkner GEJ, Gorczynski PF & Cohn TA. Psychiatric illness and obesity: recognizing the "obesogenic" nature of an inpatient psychiatric setting. *Psychiatr Serv.* 2009;60(4):538–541.
- Bhati MT, Datto CJ & O'Reardon JP. Clinical manifestations, diagnosis, and empirical treatments for catatonia. *Psychiatry*. 2007;4(3): 46–52.
- Sugai T, Suzuki Y, Yamazaki M, *et al.* High prevalence of underweight and undernutrition in Japanese inpatients with schizophrenia: a nationwide survey. *BMJ Open.* 2015;5(12):e008720, https://doi.org/10.1136/bmjopen-2015-008720.
- Kanter R & Caballero B. Global gender disparities in obesity: a review. Adv Nutr. 2012;3(4):491–498.
- Sundari S. Structural changes and quality of women's labour in India. *Indian J Labour Econ.* 2020;63(3):689–717.
- Dayabandara M, Hanwella R, Ratnatunga S, Seneviratne S, Suraweera C & de Silva VA. Antipsychotic-associated weight gain: management strategies and impact on treatment adherence. *Neuropsychiatr Dis Treat.* 2017;13:2231–2241, https://doi.org/10. 2147/NDT.S113099.
- Covell NH, Weissman EM & Essock SM. Weight gain with clozapine compared to first generation antipsychotic medications. *Schizophr Bull.* 2004;30(2):229–240.
- Cooper AJ, Gupta SR, Moustafa AF & Chao AM. Sex/gender differences in obesity prevalence, comorbidities, and treatment. *Curr Obes Rep.* 2021;10(4):458–466.
- Begley A, Paynter E, Butcher LM & Dhaliwal SS. Examining the association between food literacy and food insecurity. *Nutrients*. 2019;**11**(2):445, https://doi.org/10.3390/nu11020445.
- Dinsa GD, Goryakin Y, Fumagalli E & Suhrcke M. Obesity and socioeconomic status in developing countries: a systematic review. *Obes Rev.* 2012;13(11):1067–1079.
- Aitsi-Selmi A, Bell R, Shipley MJ & Marmot MG. Education modifies the association of wealth with obesity in women in middle-income but not low-income countries: an interaction study using seven national datasets, 2005–2010. *PLoS One.* 2014;9 (3):e90403, https://doi.org/10.1371/journal.pone.0090403.
- 40. Templin T, Cravo Oliveira Hashiguchi T, Thomson B, Dieleman J & Bendavid E. The overweight and obesity transition from the wealthy to the poor in low- and middle-income countries: a survey of household data from 103 countries. *PLoS Med.* 2019;**16**(11): e1002968.
- Mughal M & Fontan Sers C. Cereal production, undernourishment, and food insecurity in south Asia. *Rev Dev Econ.* 2020;24 (2):524–545.
- McElroy SL & Keck PE Jr. Obesity in bipolar disorder: an overview. *Curr Psychiatry Rep.* 2012;14(6):650–658.
- Correll CU, Solmi M, Veronese N, Bortolato B, Rosson S, Santonastaso P, *et al.* Prevalence, incidence and mortality from cardiovascular disease in patients with pooled and specific severe mental illness: a large-scale meta-analysis of 3,211,768 patients and 113,383,368 controls. *World Psychiatry.* 2017;16(2): 163–180.
- Klop B, Elte JWF & Cabezas MC. Dyslipidemia in obesity: mechanisms and potential targets. *Nutrients*. 2013;5(4):1218–1240, https://doi.org/10.3390/nu5041218.



- Miao Z, Alvarez M, Ko A, *et al.* The causal effect of obesity on prediabetes and insulin resistance reveals the important role of adipose tissue in insulin resistance. *PLoS Genet.* 2020;**16**(9): e1009018.
- Dare S, Mackay DF & Pell JP. Relationship between smoking and obesity: a cross-sectional study of 499,504 middle-aged adults in the UK general population. *PLoS One*. 2015;**10**(4):e0123579, https:// doi.org/10.1371/journal.pone.0123579.
- Jo YH, Talmage DA & Role LW. Nicotinic receptor-mediated effects on appetite and food intake. J Neurobiol. 2002;53(4):618–632.
- 48. Centers for Disease Control and Prevention (US), National Center for Chronic Disease Prevention and Health Promotion (US), Office on Smoking and Health (US). *How Tobacco Smoke Causes Disease: The Biology and Bebavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General.* Atlanta (GA): Centers for Disease Control and Prevention (US); 2011.
- Iqbal S, Barolia R, Ladak L & Petrucka P. Smoking cessation interventions in South Asian countries: Protocol for scoping review. BMJ Open. 2021;11(2):e038818, https://doi.org/10.1136/% 20bmjopen-2020-038818.