



This is a repository copy of *Cu-catalyzed coupling of aliphatic amines with alkylboronic esters*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/207053/>

Version: Published Version

Article:

Dennis, F., Arenas, A.R., Rodgers, G. et al. (4 more authors) (2024) Cu-catalyzed coupling of aliphatic amines with alkylboronic esters. *Chemistry - A European Journal*, 30 (19). e202303636. ISSN 0947-6539

<https://doi.org/10.1002/chem.202303636>

© 2024 The Authors. Except as otherwise noted, this author-accepted version of a journal article published in *Chemistry - A European Journal* is made available via the University of Sheffield Research Publications and Copyright Policy under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Cu-Catalyzed Coupling of Aliphatic Amines with Alkylboronic Esters

Francesca M. Dennis,^[a] Antonio Romero Arenas,^[a] George Rodgers,^[a] Muralidharan Shanmugam,^[b] Jonathan A. Andrews,^[a] Samantha L. Peralta-Arriaga,^[a] and Benjamin M. Partridge^{*[a]}


We report a Cu-catalyzed oxidative coupling of aliphatic amines with benzylic and aliphatic boronic esters to give high value alkyl amines, products found widely in applications from medicinal chemistry to materials science. This operationally simple reaction, which can be performed on gram scale, runs under mild conditions and exhibits broad functional group tolerance. The terminal oxidant of the reaction is O₂ from the

air, avoiding the need for additional chemical oxidants. Investigation into the reaction mechanism suggests that the boronic ester is activated by an aminyl radical, formed through oxidation of the amine by the Cu catalyst, to give a key alkyl radical intermediate. To demonstrate its utility and potential for late-stage functionalization, we showcase the method as the final step in the total synthesis of a TRPV1 antagonist.


Methods to form C–N bonds are of high synthetic value due to the wide prevalence of N-based functional groups in biologically active compounds, functional materials, and catalysts. Advances in transition metal catalysis have made methods to form of Csp²–N bonds become synthetically reliable and routinely used.^[1–9] However, perhaps the preparation of alkyl amines is now a greater synthetic challenge.^[10] Traditional methods to make alkyl amines, such as alkylation reactions and reductive amination^[11,12] often suffer from poor selectivity and the need for protecting group strategies. Other methods to form alkyl amines, such as hydrogenation^[13] and biocatalysis^[14–16] strategies are excellent, particularly for process chemistry applications, but are less easily applied in discovery chemistry settings. A practical and reliable cross coupling method for the formation of alkyl amines is therefore desirable, comparable to versatility of the Buchwald-Hartwig reaction for the formation of aryl amines.^[17]

Our approach to this challenge has been to develop an alkyl variant of the Chan-Lam amination,^[4–7] the oxidative coupling of an organoboron reagent with an N-based nucleophile. We and others have made some progress in this field (scheme 1),^[18–26] but there is room for improvement particularly in the scope of reaction of both amine and boron coupling partners. Herein, we


a) Previous Chan-Lam amination with alkylboronic esters



b) Previous alkyl Chan-Lam Coupling of aliphatic amines (ref 23):



c) This work:



Scheme 1. Previous examples of Chan-Lam amination with alkylboron reagents.

describe conditions for the Cu-catalyzed amination of alkylboronic esters with aliphatic amines, coupling partners that have previously been found challenging in alkyl Chan-Lam reactions.^[23] Importantly, our method uses O₂ from air (at atmospheric pressure) as the terminal oxidant, rather than a peroxide-based oxidant (which may pose an additional safety risk), making the reaction easy to perform.^[27] This is despite the propensity of alkylboronic esters to react through competing oxidative pathways under Cu-catalysis in the presence of air.^[28]

Discussion


We initially found that by heating boronic ester **1** with CuBr₂ in neat morpholine, amine **2** was formed in quantitative yield (Table 1, entry 1). While this provided proof of concept, it was clear that we needed to reduce the loading of amine and identify a suitable solvent to generate a more general procedure. After investigating a range of Cu-salts and ancillary ligands, we identified that conducting the reaction under air

[a] F. M. Dennis, A. Romero Arenas, G. Rodgers, J. A. Andrews, S. L. Peralta-Arriaga, B. M. Partridge
Department of Chemistry, University of Sheffield, Dainton Building, Sheffield, S3 7HF, United Kingdom
E-mail: b.m.partridge@sheffield.ac.uk


[b] M. Shanmugam
Department of Chemistry and Photon Science Institute, The University of Manchester, Manchester, M13 9PL, United Kingdom

Supporting information for this article is available on the WWW under <https://doi.org/10.1002/chem.202303636>


© 2024 The Authors. Chemistry - A European Journal published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Table 1. Evaluation of reaction conditions on the yield of amine **2**.^a



Entry	CuBr ₂ mol%	L (mol%)	Amine equiv.	solvent	Yield 1	2	3	4
1 ^b	100	–	40	–	< 5 %	> 95 %	< 5 %	–
2 ^c	25	L1 (25)	4	toluene	< 5 %	81 %	9 %	10 %
3 ^d	25	L2 (25)	40	–	< 5 %	95 % ^e	–	–
4 ^d	25	L3 (25)	40	–	< 5 %	67 % ^e	–	–
5	25	L1 (25)	4	toluene/IPA (1:1)	< 5 %	95 %	< 5 %	2 %
6	10	L1 (10)	4	toluene/IPA (1:1)	< 5 %	95 %	3 %	–
7	10	L1 (10)	3	toluene/IPA (1:1)	< 5 %	79 %	6 %	–
8	10	–	3.5	toluene/IPA (1:1)	< 5 %	95 %	0 %	–
9	0	–	3.5	toluene/IPA (1:1)	< 5 %	< 5 %	4 %	58 %
10 ^b	10	–	3.5	toluene/IPA (1:1)	93 %	3 %	0 %	0 %
11 ^f	10	–	1	toluene/IPA (1:1)	20 % ^g	46 %	35 % ^g	12 % ^g



L1



L2




L3

a) Reactions performed using 0.5 mmol of boronic ester **1** unless otherwise stated. Yields determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard. b) reaction carried out under an inert atmosphere. c) reaction carried out at 60 °C. d) reaction carried out at 40 °C. e) formed as a racemate. f) reaction using morpholine as the limiting reagent and 3.5 equivalents of boronic ester **1**. g) yield based on amount of boronic ester **1** instead of the limiting reagent. IPA = isopropyl alcohol.

using a combination of CuBr₂ and N,O-ligand **L1** in toluene gave amine **2** in high yield (entry 2).^[29] While ketone **3** and alcohol **4** side products were also observed, unlike our previous findings^[18,28] these were only formed in moderate amounts. A survey of ligands, including chiral ligands **L2** and **L3**, gave little improvement (entry 3–4). Investigation of the reaction solvent led to improved results. While the use of many solvents led to formation of amine **2** in good yield (see SI for more details), amination was not sufficiently selective, with ketone **3** and alcohol **4** side products formed in significant levels. The presence of these side products can make product isolation for some substrates challenging, and so more selective conditions are desirable. A 1:1 mixture of toluene/isopropanol was found to give a high yield of amine **2** with only traces of oxidation side products observed (entry 5). Under these conditions, the loading of CuBr₂ could be reduced to 10 mol% without loss of yield. However, a compromise is needed between loading of amine and yield of **2** (entries 6–7). A control experiment with CuBr₂ but without ligand suggested that the presence of **L1** had minimal effect on the reaction (entry 8). We therefore believe the likely effect of ligands, such as **L1**–**L3**, is to aid solubilisation of CuBr₂, rather than generate discrete Cu

complexes that act as a catalyst. This perhaps explains why all reactions we have conducted using chiral ligands have generated **2** as a racemate. Other control experiments showed that CuBr₂ is required for amination to occur (entry 9). Under an inert atmosphere only trace amounts of amine **2** was formed (entry 10), suggesting that O₂ is the terminal oxidant in the reaction. By switching the limiting reagent to morpholine (entry 11), and using an excess of boronic ester **1**, amine **2** was still formed albeit in moderate yield. Finally, our best conditions (entry 8) were successfully performed on gram scale, with amine **2** isolated in excellent yield (scheme 2).

We next looked to explore the scope and limitations of our reaction conditions (scheme 2). Pleasingly, a broad range of secondary amines could be coupled in good to excellent yield with boronic ester **1**. This includes simple cyclic amines such as pyrrolidine, piperidine and azepane (**5a–5c**). Substituted piperidines (**5d–5e**) and mono-protected piperazines (**5h–5i**) were also tolerated well. The coupling of amines of relevance to medicinal chemistry including difluoropiperidine (**5d**)^[30] and bicyclic amine (**5g**)^[31] was successful. Acyclic secondary amines could also be reacted successfully (**5j–5n**), though the yield is typically a slightly lower. Pleasingly, the reaction conditions are




Scheme 2. Scope of amines in the amination of boronic ester 1. Reactions were conducted on a 0.5 mmol scale. Yields reported are of isolated material of the corresponding product unless otherwise stated. a) Yield determined by ¹H NMR analysis using 1,3,5-trimethoxybenzene as an internal standard.

also successful when coupling primary aliphatic amines (5o–5r), the first time these have been reported in an alkyl Chan-Lam reaction with boronic esters. The yield of reactions with these amines are generally lower, which correlates with the lower nucleophilicity of primary versus secondary amines. Anilines can also be successfully coupled (5s–5u), although the yield is typically lower than with our previously reported conditions.^[18] Unfortunately, the coupling of benzylamine, and α-amino acids and esters was not successful. In the case of amino acids (and the corresponding ester), presumably coordination of both the carbonyl and amine to Cu leads to an inactive complex. Overall, the process shows excellent functional group tolerance, including groups such as acetals, alcohols, nitriles, sulfonamides and tertiary amines.

We were keen to test the stereoselectivity of the reaction. However, attempts to use enantiomerically enriched boronic ester (S)-1 only resulted in the formation of racemic product. To investigate this further, we stopped the reaction early allowing the isolation of unreacted boronic ester 1. Chiral HPLC analysis confirmed that the boronic ester had not undergone racemisa-

tion under the reaction conditions. Furthermore, we explored whether the reaction is diastereoselective when coupling chiral amines (5v–5y). In each case, essentially a 1:1 d.r. of the respective product was observed. Attempts to use amines with larger substituents at the 2-position were unsuccessful, presumably due reaching the reaction's limit of steric congestion.

The stereoablative nature of the amination reaction is consistent with the formation of an alkyl radical intermediate. However, we cannot rule out at this stage the formation of an alkyl copper intermediate, which have been reported to be configurationally unstable above –50 °C.^[32] To probe the possibility of a radical intermediate, we performed the amination of cyclopropane 6 in a radical clock experiment (eq 1). This resulted in a mixture of two amines, amine 7 with the cyclopropane intact and amine 8 in which the cyclopropane ring has opened. This suggests a radical intermediate is formed, with recombination occurring on a similar timescale to cyclopropane ring opening.^[33]



To gather further evidence, we performed experiments in which we added TEMPO, known to form adducts with alkyl radicals,^[34] to our reaction. Adduct **9** was observed in each case, suggesting that the boronic ester acts as an alkyl radical precursor. When 1 equivalent of TEMPO was used, the major product from the reaction was adduct **9**. However, when 10 mol% of both TEMPO and CuBr₂ were added, amine **2** still was formed in high yield along with 10% of adduct **9**. This suggests that TEMPO does not irreversibly inhibit the copper catalyst.

While formation of alkyl radicals from alkylboronic esters has been previously reported, this typically requires prior formation of a boron 'ate' complex through coordination of an alkoxide, pyridine or aryl lithium.^[35–39] However, such an 'ate' complex was not observed by ¹H or ¹¹B NMR spectroscopy when analysing mixtures of boronic ester **1** with morpholine in the presence of isopropanol.^[29]

To investigate this further, we conducted a series of cyclic voltammetry experiments. Boronic ester **1** showed a high oxidation potential ($E_{on} > 1.44$ V) in MeCN (figure 1a).^[40] The voltammogram of CuBr₂ does not change in the presence of the boronic ester, suggesting there is not a direct interaction between **1** and the Cu salt.^[41,42] However, cyclic voltammetry of a mixture of morpholine and **1** does show the formation of a new peak (oxidation potential of $E_{pc} = 1.25$ V; figure 1b). Given that this is greater than the oxidation potential of morpholine ($E_{pc} = 0.81$ V), we suggest this could arise from a new species generated upon interaction of **1** and a morpholine radical, which can be oxidised at a lower potential than the boronic ester without activation. Aminyl radicals have been reported to react with trialkylboranes^[43–45] and alkylboronic esters^[46,47] leading to the formation of alkyl radicals.

To further understand the nature of the radical intermediates formed during the amination reaction, EPR studies were performed using 5,5-dimethyl-1-pyrroline N-oxide (DMPO) as a spin trap. Analysis of a solution of CuBr₂ in toluene:IPA (1:1) showed an intense, broad EPR signal at around 3150 G. This signal did not change upon addition of boronic ester **1** suggesting that the electronic structure of Cu does not change in the presence of the boronic ester. This is consistent with the cyclic voltammetry data above (Figure 1a). When a mixture of morpholine and CuBr₂ was analysed, a different EPR signal was observed, presumably in part due to ligation of morpholine to




Figure 1. Cyclic voltammetry experiments, conducted in MeCN under N₂ using nBu₄NPF₆ as the electrolyte, of a) boronic ester **1** and CuBr₂, and b) boronic ester **1** and morpholine.

Cu (Figure S1). When CuBr₂ with morpholine, or CuBr₂ with boronic ester **1**, samples were measured in the presence of DMPO, the Cu(II) EPR signal disappears, presumably due to reduction of Cu(II) to Cu(I) (Figure S2). Reduction and regeneration of the Cu(II) signals are observed in the reaction mixtures of CuBr₂ with morpholine, or CuBr₂ with boronic ester **1** and morpholine, even in the absence of DMPO (Figure S3). However, this process is slow (Figure S3). This implies that Cu(II) reduction and regeneration is part of the catalytic cycle and this process is enhanced in the presence of DMPO (Figure S2). No DMPO-adduct signals were detected for CuBr₂ with boronic ester **1** (Figure 2A, red trace). However, for CuBr₂ with morpholine a new signal is observed (Figure 2A, black trace) which increases steadily with time (Figure 2B). This spectrum is consistent with literature data for aminyl-radical adducts of DMPO.^[48] When a mixture of CuBr₂ with morpholine and boronic ester **1** was measured with DMPO, complex EPR spectra were observed due to multiple DMPO-adducts (Figure 2C).^[49,50] We could model the spectra as a mixture of three components (Figure S4): (1) a DMPO-morpholine radical adduct, with the same parameters as the species found in the absence **1**, and which is the dominant species (2) a carbon-centred DMPO-R adduct, and (3) an unknown radical with triplet-of-doublets spectrum that increases for more than 12 hours. The spectrum of this unknown radical is possibly due to decomposition of the primary DMPO-





Figure 2. Data from EPR experiments, performed in solutions of toluene:IPA (1:1) at 80 °C. a) EPR spectra for CuBr_2 + morpholine + DMPO (black trace) and CuBr_2 + boronic ester **1** + DMPO (red trace); b) Plot of EPR signal intensity at 3362 G versus time for a solution of CuBr_2 + morpholine + DMPO; c) EPR spectra of a mixture of CuBr_2 + morpholine + boronic ester **1** + DMPO collected ~100 minutes after mixing (black trace) and of DMPO (red trace) as a negative control.


adducts, but is not consistent with the oxidation product DMPOX which is often seen in DMPO spin trapping experiments.^[51] While the exact identity of the DMPO–R species cannot be uniquely determined from this data, a benzylic radical would be consistent with the TEMPO-trapping experiment above (eq 2).

Based on the experimental data above, we tentatively propose a reaction mechanism outlined in Scheme 3. This involves oxidation of the amine **18** by a Cu(II) complex **20**,^[52] to generate an aminyl radical **19** and Cu(I) complex **21**. Aminyl radical **19** can then combine with alkylboronic ester **17** to give intermediate **16**, which can undergo homolytic bond cleavage to generate alkyl radical **14** and boronate **15**. The alkyl radical **14** can then combine with an amino Cu(II) complex **10**, leading to a Cu(III) complex **12** that can undergo reductive elimination to give the alkyl amine product **12** and a Cu(I) complex **13**. Re-oxidation of Cu(I) complexes **13** and **21** by O_2 from air regenerates Cu(II) complexes **10** and **20** respectively. The fact that two equivalents of amine are consumed in the proposed mechanism presumably explains why the yield of amination is reduced with lower amine loadings or when the amine is the limiting reagent (see Table 1). Further investigations into elucidating this complex reaction mechanism are ongoing.

We have also explored the scope of reaction with respect to the boronic ester (scheme 4). A range of secondary benzylic boronic esters were successfully reacted with morpholine in good to excellent yield. Varying the electronics of the substitutes on the aryl ring of the boronic ester did not lead to great variation in yield. However, the reaction of boronic esters with more electron donating MeO- (**23a**) or electron withdrawing CF_3- (**23g**) substituents showed a small reduction in yield, in part to a small increase in corresponding alcohol and ketone side products observed in the crude reaction mixture. Methyl ester-containing boronic ester was also successfully reacted (**23f**), though a change of reaction solvent to PrOAc was made to avoid formation of the corresponding isopropyl ester trans-esterification side product, which was observed when using the standard IPA/toluene solvent system. However, the yield of amination is reduced compared with our standard conditions, in part due to larger quantities of alcohol and ketone side products observed when using the alternative reaction solvent. Importantly, aryl bromides and chlorides (**23e**, **23i**, **23j**, **23v**) were tolerated, with no products from Ullmann-



Scheme 3. Proposed mechanism for the Cu-catalysed amination reaction.



Goldberg amination observed. In addition, functional groups including azides, ethers and indole were also tolerated. Pleasingly, an alkyl chloride was also tolerated, with no products of amination through nucleophilic substitution observed in the reaction mixture (**23o**). A primary benzylic boronic ester was also successfully coupled in high yield (**23r**).

The amination reaction was also successful with primary and secondary aliphatic boronic esters, despite the radical intermediates formed from these unactivated boronic esters being less stable than those from benzylic boronic esters. Primary alkylboronic esters could be coupled (**23s–23x**) in good yield, though with slightly reduced efficiency compared with benzylic boronic esters. While less reactive still, we could also form the amination product **23y** from a secondary aliphatic boronic ester.

Subjecting tertiary boronic esters **24a** and **24b** to our reaction conditions did lead to formation of a C-tertiary amines **25a** and **25b**, albeit in lower yield compared to our previous results when coupling anilines.^[18] In reaction of **24a**, the corresponding alkene **25** was formed as the major product from the mixture. Unfortunately, tert-butyl boronic ester was found to be unreactive under these reaction conditions.

Finally, we wanted to apply our amination reaction as the final step in the synthesis of a complex molecule, to simulate how the method could be used in a discovery chemistry scenario. We targeted amine **28**, a TRVP1 inhibitor which is a potential treatment for chronic pain.^[53] Due to the poor solubility of **27** in most organic solvents, we required the use of DMSO as reaction solvent in which **27** was partially soluble. Despite this, and without further optimisation, amine **28** could be isolated successfully, and in a yield that would allow the material to be tested for its biological activity. We therefore believe our method could find application in settings such as medicinal chemistry, and allow chemists to develop structure-activity relationships through the coupling of a common advanced amine or boronic ester intermediates.

Conclusions

In conclusion, we have developed conditions for the Cu-catalyzed amination of alkylboronic esters with aliphatic amines. This method expands the scope of the alkyl Chan-Lam reaction, enabling, the coupling of both primary and secondary alkyl amines with aliphatic boronic esters for the first time. Of note, the method uses oxygen from air as the terminal oxidant for the reaction, without need for additional chemical oxidants. The reaction conditions tolerate a broad range of functional groups, and have been applied successfully as the final step in the synthesis of a TRVP1 inhibitor. Mechanistic studies suggest that the reaction proceeds via an alkyl radical intermediate, generated from the boronic ester upon activation by an aminyl radical. The investigation of related Cu-catalyzed oxidative couplings of alkylboron reagents is ongoing, and will be reported in due course.

Acknowledgements

We thank the EPSRC (Grant EP/T009292/1), the Royal Society (Grant RSG\R1\180065), and the University of Sheffield for financial support. We are grateful to the EPR National Facility at the University of Manchester (funder by EPSRC grants EP/V0353231/1 and EP/S033181/1) for conducting EPR experiments and valuable discussion.

Conflict of Interests

The authors declare no conflict of interest.

Data Availability Statement

The data that support the findings of this study are available in the supplementary material of this article.

Keywords: amination · amines · boron · Chan-Lam · copper

- [1] M. M. Heravi, Z. Kheilkordi, V. Zadsirjan, M. Heydari, M. Malmir, *J. Organomet. Chem.* **2018**, *861*, 17–104.
- [2] P. Ruiz-Castillo, S. L. Buchwald, *Chem. Rev.* **2016**, *116*, 12564–12649.
- [3] I. P. Beletskaya, A. V. Cheprakov, *Organometallics* **2012**, *31*, 7753–7808.
- [4] J. X. Qiao, P. Y. Lam, *Synthesis* **2011**, *6*, 829–856.
- [5] F. Monnier, M. Taillefer, in *Topics in Organometallic Chemistry*, (Eds. M. Taillefer, D. Ma); Springer Berlin Heidelberg: Berlin, Heidelberg, **2013**; pp 173–204.
- [6] I. Munir, A. F. Zahoor, N. Rasool, S. A. R. Naqvi, K. M. Zia, R. Ahmad, *Mol. Divers.* **2019**, *23*, 215–259.
- [7] M. J. West, J. W. B. Fyfe, J. C. Vantourout, A. J. B. Watson, *Chem. Rev.* **2019**, *119*, 12491–12523.
- [8] C. Sambriago, S. P. Marsden, A. J. Blacker, P. C. McGowan, *Chem. Soc. Rev.* **2014**, *43*, 3525–3550.
- [9] F. Monnier, M. Taillefer, *Angew. Chemie Int. Ed.* **2009**, *48*, 6954–6971.
- [10] A. Trowbridge, S. M. Walton, M. J. Gaunt, *Chem. Rev.* **2020**, *120*, 2613–2692.
- [11] A. F. Abdel-Magid, S. J. Mehrman, *Org. Process Res. Dev.* **2006**, *10*, 971–1031.
- [12] O. I. Afanasyev, E. Kuchuk, D. L. Usanov, D. Chusov, *Chem. Rev.* **2019**, *119*, 11857–11911.
- [13] A. Cabré, X. Verdager, A. Riera, *Chem. Rev.* **2022**, *122*, 269–339.
- [14] M. D. Patil, G. Grogan, A. Bommaris, H. Yun, *ACS Catal.* **2018**, *8*, 10985–11015.
- [15] S. Simić, E. Zukić, L. Schermund, K. Faber, C. K. Winkler, W. Kroutil, *Chem. Rev.* **2022**, *122*, 1052–1126.
- [16] J. J. Sangster, J. R. Marshall, N. J. Turner, J. Mangas-Sanchez, *ChemBioChem* **2022**, *23*, e202100464.
- [17] K. R. Campos, P. J. Coleman, J. C. Alvarez, S. D. Dreher, R. M. Garbaccio, N. K. Terrett, R. D. Tillyer, M. D. Truppo, E. R. Parmee, *Science* **2019**, *363*, eaat0805.
- [18] J. D. Grayson, F. M. Dennis, C. C. Robertson, B. M. Partridge, *J. Org. Chem.* **2021**, *86*, 9883–9897.
- [19] L. M. Mori-Quiroz, K. W. Shimkin, S. Rezazadeh, R. A. Kozlowski, D. A. Watson, *Chem. Eur. J.* **2016**, *22*, 15654–15658.
- [20] E. Racine, F. Monnier, J.-P. Vors, M. Taillefer, *Chem. Commun.* **2013**, *49*, 7412–7414.
- [21] S. Sueki, Y. Kuninobu, *Org. Lett.* **2013**, *15*, 1544–1547.
- [22] D. S. Kim, H. G. Lee, *J. Org. Chem.* **2021**, *86*, 17380–17394.
- [23] S. Bénard, L. Neuville, J. Zhu, *Chem. Commun.* **2010**, *46*, 3393–3395.
- [24] K. Singh, M. Kumar, E. Pavada, K. Naran, D. F. Warner, P. G. Ruminski, K. Chibale, *Bioorg. Med. Chem. Lett.* **2014**, *24*, 2985–2990.
- [25] M. Larrosa, C. Guerrero, R. Rodríguez, J. Cruces, *Synlett* **2010**, *2010*, 2101–2105.


- [26] I. González, J. Mosquera, C. Guerrero, R. Rodríguez, J. Cruces, *Org. Lett.* **2009**, *11*, 1677–1680.
- [27] A video showing our reaction set up and tips to help solve problems can be viewed at: https://digitalmedia.sheffield.ac.uk/id/1_isl6hrng.
- [28] J. D. Grayson, B. M. Partridge, *ACS Catal.* **2019**, *9*, 4296–4301.
- [29] For more information, see the supporting information.
- [30] N. A. Meanwell, *J. Med. Chem.* **2018**, *61*, 5822–5880.
- [31] A. Zask, J. Kaplan, J. C. Verheijen, D. J. Richard, K. Curran, N. Brooijmans, E. M. Bennett, L. Toral-Barza, I. Hollander, S. Ayril-Kaloustian, K. Yu, *J. Med. Chem.* **2009**, *52*, 7942–7945.
- [32] J. Skotnitzki, V. Morozova, P. Knochel, *Org. Lett.* **2018**, *20*, 2365–2368.
- [33] T. A. Halgren, J. D. Roberts, J. H. Horner, F. N. Martinez, C. Tronche, M. Newcomb, *J. Am. Chem. Soc.* **2000**, *122*, 2988–2994.
- [34] E. G. Bagryanskaya, S. R. A. Marque, *Chem. Rev.* **2014**, *114*, 5011–5056.
- [35] F. Lima, M. A. Kabeshov, D. N. Tran, C. Battilocchio, J. Sedelmeier, G. Sedelmeier, B. Schenkel, S. V. Ley, *Angew. Chemie Int. Ed.* **2016**, *55*, 14085–14089.
- [36] A. J. J. Lennox, J. E. Nutting, S. S. Stahl, *Chem. Sci.* **2018**, *9*, 356–361.
- [37] V. Corcé, C. Ollivier, L. Fensterbank, *Chem. Soc. Rev.* **2022**, *51*, 1470–1510.
- [38] C. Shu, A. Noble, V. K. Aggarwal, *Angew. Chemie Int. Ed.* **2019**, *58*, 3870–3874.
- [39] S. Pillitteri, P. Ranjan, E. V. Van der Eycken, U. K. Sharma, *Adv. Synth. Catal.* **2022**, *364*, 1643–1665.
- [40] MeCN was used as a solvent to enable solubility of the $n\text{Bu}_4\text{NPF}_6$ electrolyte, which was insoluble in toluene/IPA (1:1). When the standard reaction was performed using MeCN as solvent, amination was successful with amine **2** formed in 78% yield. See the supporting information for details.
- [41] G. Sorin, R. Martinez Mallorquin, Y. Contie, A. Baralle, M. Malacria, J.-P. Goddard, L. Fensterbank, *Angew. Chemie Int. Ed.* **2010**, *49*, 8721–8723.
- [42] C. F. Lane, *J. Organomet. Chem.* **1971**, *31*, 421–431.
- [43] A. G. Davies, S. C. W. Hook, B. P. Roberts, *J. Organomet. Chem.* **1970**, *22*, c37–c38.
- [44] A. G. Davies, S. C. W. Hook, B. P. Roberts, *J. Organomet. Chem.* **1970**, *23*, C11–C13.
- [45] J. G. Sharefkin, H. D. Banks, *J. Org. Chem.* **1965**, *30*, 4313–4314.
- [46] E. Speckmeier, T. C. Maier, *J. Am. Chem. Soc.* **2022**, *144*, 9997–10005.
- [47] Z. Wang, N. Wierich, J. Zhang, C. G. Daniliuc, A. Studer, *J. Am. Chem. Soc.* **2023**, *145*, 8770–8775.
- [48] C. Chen, R. P. Veregin, J. R. Harbour, M. L. Hair, S. L. Issler, J. Tromp, *J. Phys. Chem.* **1989**, *93*, 2607–2609.
- [49] G. R. Buettner, *Free Radic. Biol. Med.* **1987**, *3*, 259–303.
- [50] Z. Huang, M. Shanmugam, Z. Liu, A. Brookfield, E. L. Bennett, R. Guan, D. E. Vega Herrera, J. A. Lopez-Sanchez, A. G. Slater, E. J. L. McInnes, X. Qi, J. Xiao, *J. Am. Chem. Soc.* **2022**, *144*, 6532–6542.
- [51] P. Bilski, K. Reszka, M. Bilska, C. F. Chignell, *J. Am. Chem. Soc.* **1996**, *118*, 1330–1338.
- [52] T. Kauffmann, J. Albrecht, D. Berger, J. Legler, *Angew. Chemie Int. Ed.* **1967**, *6*, 633–634.
- [53] H.-L. Wang, J. Katon, C. Balan, A. W. Bannon, C. Bernard, E. M. Doherty, C. Dominguez, N. R. Gavva, V. Gore, V. Ma, N. Nishimura, S. Surapaneni, P. Tang, R. Tamir, O. Thiel, J. J. S. Treanor, M. H. Norman, *J. Med. Chem.* **2007**, *50*, 3528–3539.

Manuscript received: November 2, 2023

Accepted manuscript online: January 3, 2024

Version of record online: ■■■

RESEARCH ARTICLE



>40 examples, 28-87% yield

- **C(sp³)-N coupling**
- **Oxidant is O₂ from air**
- **Operationally simple**

We report the Cu-catalysed coupling of alkylboronic esters and aliphatic amines. This mild method is functional group tolerant, and uses O₂ from air as the terminal oxidant. Investigation

into the mechanism suggest that the boronic ester reacts with an aminyl radical, formed in situ, to generate an alkyl radical intermediate.

F. M. Dennis, A. Romero Arenas, G. Rodgers, M. Shanmugam, J. A. Andrews, S. L. Peralta-Arriaga, B. M. Partridge*

1 – 9

Cu-Catalyzed Coupling of Aliphatic Amines with Alkylboronic Esters

