



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/207735/>

Version: Accepted Version

Article:

Abrishami, T., Chudnovsky, M., Dibek, C. et al. (2022) Graphs with polynomially many minimal separators. *Journal of Combinatorial Theory, Series B*, 152. pp. 248-280. ISSN: 0095-8956

<https://doi.org/10.1016/j.jctb.2021.10.003>

© 2021, Elsevier. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Graphs with polynomially many minimal separators

Tara Abrishami¹ Maria Chudnovsky^{1,*} Cemil Dibek^{1,*}

Stéphan Thomassé^{2,†} Nicolas Trotignon^{2,†} Kristina Vušković^{3,‡}

¹Princeton University, Princeton, NJ 08544

²Univ Lyon, EnsL, UCBL, CNRS, LIP, F-69342, LYON Cedex 07, France

³School of Computing, University of Leeds, UK

September 22, 2021

Abstract

We show that graphs that do not contain a theta, pyramid, prism, or turtle as an induced subgraph have polynomially many minimal separators. This result is the best possible in the sense that there are graphs with exponentially many minimal separators if only three of the four induced subgraphs are excluded. As a consequence, there is a polynomial time algorithm to solve the maximum weight independent set problem for the class of (theta, pyramid, prism, turtle)-free graphs. Since every prism, theta, and turtle contains an even hole, this also implies a polynomial time algorithm to solve the maximum weight independent set problem for the class of (pyramid, even hole)-free graphs.

1 Introduction

All graphs in this paper are finite and simple. Let $G = (V, E)$ be a graph. A set $C \subseteq V(G)$ is a *minimal separator* of G if there are two distinct connected components L, R of $G \setminus C$ such that $N(L) = N(R) = C$. A class \mathcal{G} of graphs is said to have the *polynomial separator property* if there exists a constant c such that every graph $G \in \mathcal{G}$ has at most $|V(G)|^c$ minimal separators.

The polynomial separator property has proven to be a desirable property due to its connection with potential maximal cliques and the maximum weight treewidth k induced subgraph problem. Given a graph G , a nonnegative weight function on $V(G)$, and an integer k , the MAXIMUM WEIGHT TREewidth k INDUCED SUBGRAPH problem (MWT k ISG) asks for a maximum-weight induced subgraph of G of treewidth less than k . The MAXIMUM WEIGHT INDEPENDENT SET problem (MWIS), which asks for an independent set of G with maximum weight, and the FEEDBACK VERTEX SET problem (FVS), which asks for a minimum-size set $X \subseteq V(G)$ such that $G \setminus X$ is a forest, are special cases of MWT k ISG when $k = 1$ and $k = 2$, respectively. Recently, significant progress

*Supported by NSF Grant DMS-1763817. This material is based upon work supported by, or in part by, the U.S. Army Research Laboratory and the U. S. Army Research Office under grant number W911NF-16-1-0404.

†Partially supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program Investissements d’Avenir (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR), and by the French National Research Agency under research grant ANR DIGRAPHS ANR-19-CE48-0013-01.

‡Partially supported by EPSRC grant EP/N0196660/1.

was made regarding the complexity of MWIS in various graph classes using potential maximal cliques, originally developed by Bouchitté and Todinca [5,6]. A milestone result with this approach was obtained in 2014 by Lokshtanov, Vatschelle, and Villanger [11], who designed a polynomial-time algorithm for MWIS in P_5 -free graphs. Later, using the same framework, Grzesik et al. [10] provided a polynomial-time algorithm for MWIS in P_6 -free graphs. More recently, Abrishami et al. [1] extended the framework of potential maximal cliques to MWT k ISG, and gave a polynomial-time algorithm for MWIS in graphs with no induced cycle of length five or greater, and for FVS in P_5 -free graphs.

Given an integer k , it is known that MWT k ISG can be solved in polynomial time for graphs that have polynomially many potential maximal cliques. Minimal separators are closely related to potential maximal cliques: it was shown in [6] that a graph has polynomially many potential maximal cliques if and only if it has polynomially many minimal separators. Consequently, MWT k ISG is polynomial-time solvable in any class of graphs that has the polynomial separator property. It is therefore interesting to find classes of graphs where the number of minimal separators is bounded by a polynomial. We now define four graphs of interest to us (see also Figure 1):

- A *theta* is a graph G consisting of two nonadjacent vertices a, b and three paths P_1, P_2, P_3 , each from a to b , and otherwise vertex-disjoint, such that for $1 \leq i < j \leq 3$, $V(P_i) \cup V(P_j)$ induces a hole in G . In particular, each of P_1, P_2, P_3 has at least two edges. We say that G is a theta between a and b .
- A *pyramid* is a graph G consisting of a vertex a and a triangle $\{b_1, b_2, b_3\}$, and three paths P_1, P_2, P_3 , such that: P_i is between a and b_i for $i = 1, 2, 3$; for $1 \leq i < j \leq 3$, P_i, P_j are vertex-disjoint except for a and $V(P_i) \cup V(P_j)$ induces a hole in G ; and in particular at most one of P_1, P_2, P_3 has only one edge. We say that G is a pyramid from a to $b_1 b_2 b_3$.
- A *prism* is a graph G consisting of two vertex-disjoint triangles $\{a_1, a_2, a_3\}, \{b_1, b_2, b_3\}$, and three paths P_1, P_2, P_3 , pairwise vertex-disjoint, where each P_i has ends a_i, b_i , and for $1 \leq i < j \leq 3$, $V(P_i) \cup V(P_j)$ induces a hole in G . In particular, each of P_1, P_2, P_3 has at least one edge. We say G is a prism between $a_1 a_2 a_3$ and $b_1 b_2 b_3$.
- A *turtle* is a graph G consisting of two vertex-disjoint paths P_1, P_2 and two adjacent vertices $x, y \in V(G) \setminus (V(P_1) \cup V(P_2))$ such that for $i = 1, 2$, P_i is from a_i to b_i , a_1 is adjacent to a_2 , b_1 is adjacent to b_2 , $V(P_1) \cup V(P_2)$ induces a hole in G , x has at least three neighbors in P_1 and no neighbors in P_2 , and y has at least three neighbors in P_2 and no neighbors in P_1 . We say that G is an xy -*turtle* where we call x and y the *centers* of G .




Figure 1: Theta, pyramid, prism, and turtle

Thetas, pyramids, prisms, and turtles are interesting because they provide examples of graphs with exponentially many minimal separators. Specifically, we have the following examples of graphs with exponentially many minimal separators (see also Figure 2).

- A k -*theta* is a graph G with vertex set $V(G) = \{a, a_1, \dots, a_k, b, b_1, \dots, b_k\}$, and its set of edges consists of the pairs of the following form: aa_i, bb_i , and $a_i b_i$ for $1 \leq i \leq k$.

- A k -*pyramid* is a graph G with vertex set $V(G) = \{a, a_1, \dots, a_k, b_1, \dots, b_k\}$, and its set of edges consists of the pairs of the following form: aa_i and $a_i b_i$ for $1 \leq i \leq k$, and $b_i b_j$ for $1 \leq i < j \leq k$.
- A k -*prism* is a graph G consisting of two cliques of size k and a k -edge matching between them. More precisely, $V(G) = \{a_1, \dots, a_k, b_1, \dots, b_k\}$, each of the sets $\{a_1, \dots, a_k\}$ and $\{b_1, \dots, b_k\}$ is a clique, and $a_i b_i \in E(G)$ for $1 \leq i \leq k$, and there are no other edges in G .
- A k -*turtle* is a graph G with two non-adjacent vertices $a, b \in V(G)$, two paths P_1 and P_2 from a to b , vertex-disjoint except for a and b , such that $V(P_1) \cup V(P_2)$ induces a hole H in G . Also, for $1 \leq i \leq k$, $x_i, y_i \in V(G) \setminus V(H)$ such that $x_i y_i \in E(G)$, and x_i has at least three neighbors in P_1 and no neighbors in P_2 , and y_i has at least three neighbors in P_2 and no neighbors in P_1 . Furthermore, the neighbors of x_i 's in P_1 and the neighbors of y_i 's in P_2 are nested along P_1 and P_2 as shown in Figure 2. Specifically, the neighbors of x_i in P_1 are between a and the neighbors of x_j in P_1 for all $1 \leq i < j \leq k$ and the neighbors of y_i in P_2 are between a and the neighbors of y_j in P_2 for all $1 \leq i < j \leq k$.




Figure 2: k -theta, k -pyramid, k -prism, k -turtle, and k -ladder

There are other examples of graphs with exponentially many minimal separators, such as the k -ladder shown in Figure 2. The k -ladder also contains a pyramid. In view of these examples, it is natural to ask whether excluding theta, pyramid, prism, and turtle in a graph is enough to obtain a polynomial number of minimal separators. This was conjectured in [7]:

Conjecture 1.1 ([7]). *There is a polynomial P such that every graph G that contains no theta, pyramid, prism, or turtle has at most $P(|V(G)|)$ minimal separators.*

Here we prove Conjecture 1.1. Let \mathcal{C} be the class of (theta, pyramid, prism, turtle)-free graphs. We prove that the graphs in \mathcal{C} have polynomially many minimal separators. Note that in view of the results in [3], listing the minimal separators of a graph can be done in polynomial time in the size of the graph and the number of its minimal separators. Our proof that the graphs in \mathcal{C} have polynomially many minimal separators is algorithmic in nature, so we include a polynomial-time algorithm here to construct minimal separators of graphs in \mathcal{C} for completeness.

Theorem 1.2. *Let $G \in \mathcal{C}$. One can construct a set \mathcal{S} of size at most $|V(G)|^{18}$ in polynomial time such that \mathcal{S} is the set of all minimal separators of G .*

Since the graphs in Figure 2 have exponentially many minimal separators, Theorem 1.2 is in a sense the best possible. Moreover, as explained above, given an integer k , Theorem 1.2 implies that MWTkISG can be solved in polynomial time for graphs in \mathcal{C} . To be more precise, let n, m, p, s denote, respectively, the number of vertices, the number of edges, the number of potential maximal

cliques, and the number of minimal separators of a graph G . It is proved in [3] that computing the minimal separators of G can be done in time $\mathcal{O}(n^3s)$. In [6], it is proved that $p \leq \mathcal{O}(ns^2 + ns + 1)$ and that the potential maximal cliques of G can be listed in time $\mathcal{O}(n^2ms^2)$. In [1], it is proved that given the list of potential maximal cliques of G and an integer k , if p is polynomial in n , then MWT k ISG can be solved in time $n^{\mathcal{O}(k)}$. By Theorem 1.2, for a graph $G \in \mathcal{C}$, we have $s \leq \mathcal{O}(n^{18})$, and so $p \leq \mathcal{O}(n^{37})$. Therefore, MWT k ISG can be solved in time $n^{\mathcal{O}(k)}$ in \mathcal{C} . Using results from [11], a better complexity for MWIS can be achieved. In [11], based on [9], it is proved that, given the list of potential maximal cliques, MWIS can be solved in time $\mathcal{O}(n^5mp)$ in any graph. Therefore, MWIS can be solved in time $\mathcal{O}(n^{44})$ in \mathcal{C} .

It is easy to observe that every prism, theta, and turtle contains an even hole. Therefore, the following is an immediate corollary of Theorem 1.2.

Corollary 1.3. *The class of (pyramid, even hole)-free graphs has the polynomial separator property.*

In [7], a better bound than the one given in Theorem 1.2 is achieved for (pyramid, even hole)-free graphs. In particular, Corollary 1.3 implies that MWT k ISG and MWIS can be solved in (pyramid, even hole)-free graphs in polynomial time. A *cap* is a cycle of length at least five with exactly one chord and that chord creates a triangle with the cycle. Since every pyramid contains a cap, Corollary 1.3 generalizes a result of [8] where it is shown that MWIS can be solved in (cap, even hole)-free graphs in polynomial time.

We conjecture a stronger version of Theorem 1.2. For an integer $k \geq 3$, a graph G is called a *k-creature* if it is given as follows: $V(G) = A \cup B \cup \{x_1, \dots, x_k\} \cup \{y_1, \dots, y_k\}$ such that

- (i) $G[A]$ and $G[B]$ are connected, and A is anticomplete to B ,
- (ii) for $i = 1, \dots, k$, $x_i y_i \in E(G)$, x_i has a neighbor in A and is anticomplete to B , y_i has a neighbor in B and is anticomplete to A , and
- (iii) for $1 \leq i, j \leq k$ with $i \neq j$, $x_i y_j \notin E(G)$.

We observe that if G is a k -creature, then G contains a theta, pyramid, prism, or turtle; see Lemma 4.1 for details. We conjecture the following:

Conjecture 1.4. *There exists $f : \mathbb{N} \rightarrow \mathbb{N}$ such that if no induced subgraph of G is a k -creature, then G has at most $|V(G)|^{f(k)}$ minimal separators.*

Observe that even if Conjecture 1.4 is true, it does not provide a full characterization of classes with the polynomial separator property. For example, for every integer $k \geq 1$, let T_k be a k -turtle such that the two paths P_1 and P_2 both have length 2^{2^k} . Let \mathcal{D} be the class of graphs formed by all induced subgraphs of the graphs T_k , $k \geq 1$. Observe that T_k has 2^k minimal separators, which is polynomial in $|V(T_k)|$ since $|V(T_k)| \geq 2^{2^k}$, and so \mathcal{D} has the polynomial separator property. However, \mathcal{D} contains k -creatures with k arbitrarily large.

We prove a weaker version of Conjecture 1.4. The proof can also be found in [7]. We say that a graph G is an *immature k-creature* if $V(G)$ can be partitioned into two sets $X = \{x_1, \dots, x_k\}$ and $Y = \{y_1, \dots, y_k\}$ such that the only edges between X and Y are the edges $x_i y_i$ for $i = 1, \dots, k$. The edges among vertices of X and vertices of Y are unrestricted.

Theorem 1.5. *Let $k \geq 1$ be an integer and let G be a graph on n vertices such that no induced subgraph of G is an immature k -creature. Then, G has at most $\mathcal{O}(n^{2^{k-2}})$ minimal separators that can be enumerated in time $\mathcal{O}(n^{2^k})$.*

Proof. Let a and b be two non-adjacent vertices in G . Let C be a minimal separator that separates a and b . Let A and B be the components of $G \setminus C$ that contain a and b , respectively. By the minimality

of C , every vertex in C has a neighbor in A . It is therefore well-defined to consider an inclusion-wise minimal subset X_A of A such that $C \subseteq N(X_A)$. For every $x \in X_A$, there exists a vertex $c \in C$ such that $xc \in E(G)$ and no other vertex of X_A is adjacent to c , for otherwise, $X_A \setminus \{x\}$ would contradict the minimality of X_A . It follows that $G[X_A \cup C]$ contains an immature $|X_A|$ -creature, and so $|X_A| < k$. We define a similar set $X_B \subseteq B$, and we observe that $C = N(X_A) \cap N(X_B)$.

Now, the following algorithm enumerates all minimal separators of G : for every pair of sets X_A, X_B with $|X_A|, |X_B| < k$, compute $C = N(X_A) \cap N(X_B)$ and check whether C is a minimal separator. Since $\binom{n}{i} \leq n^i$, we have $\binom{n}{0} + \dots + \binom{n}{k-1} \leq kn^{k-1}$. Therefore, the algorithm enumerates at most $\mathcal{O}(n^{2k-2})$ minimal separators in time $\mathcal{O}(n^{2k})$. \square

We note that there exist graphs in \mathcal{C} of arbitrarily large cliquewidth. In [2], examples of even-hole-free graphs of arbitrarily large cliquewidth are presented. Those graphs are also diamond-free and they have no clique separators. (A *diamond* is the graph with vertex set $\{a, b, c, d\}$ with all possible edges except ab .) However, they are not in \mathcal{C} because they contain pyramids. In [7], a procedure to obtain graphs in \mathcal{C} with unbounded cliquewidth by modifying graphs defined in [2] is explained in detail. Moreover, those graphs contain arbitrarily large immature k -creatures, and so the main result of the current paper is not a corollary of Theorem 1.5.

The rest of the paper is devoted to the proof of Theorem 1.2. In Section 2, we prove a useful theorem about star cutsets of graphs in \mathcal{C} . In Section 3, we describe the structure of proper separators of graphs in \mathcal{C} . In Section 4, we construct a list of all minimal separators of graphs in \mathcal{C} and prove Theorem 1.2.

Definitions

Let $G = (V, E)$ be a graph. For $X \subseteq V(G)$, $G[X]$ denotes the induced subgraph of G with vertex set X and $G \setminus X$ denotes the induced subgraph of G with vertex set $V(G) \setminus X$. We use induced subgraphs and their vertex sets interchangeably throughout the paper. We say that G *contains* a graph H if G has an induced subgraph isomorphic to H . A graph G is *H -free* if it does not contain H . When \mathcal{H} is a set of graphs, we say that G is \mathcal{H} -free if G is H -free for every $H \in \mathcal{H}$. For a graph H , we say that a set $X \subseteq V(G)$ is an H in G if $G[X]$ is isomorphic to H .

Let $X \subseteq V(G)$. The *neighborhood* of X in G , denoted by $N(X)$, is the set of all vertices in $V(G) \setminus X$ with a neighbor in X . The *closed neighborhood* of X in G , denoted $N[X]$, is given by $N[X] = N(X) \cup X$. For $u \in V(G)$, $N(u) = N(\{u\})$ and $N[u] = N[\{u\}]$. For $u \in V(G) \setminus X$, $N_X(u) = N(u) \cap X$. Let $Y \subseteq V(G)$ be disjoint from X . We say X is *complete* to Y if every vertex in X is adjacent to every vertex in Y , and X is *anticomplete* to Y if every vertex in X is non-adjacent to every vertex in Y . Note that the empty set is complete and anticomplete to every $X \subseteq V(G)$. We say that a vertex v is *complete* (*anticomplete*) to $X \subseteq V(G)$ if $\{v\}$ is complete (anticomplete) to X , and an edge $e = uv$ is *complete* (*anticomplete*) to X if $\{u, v\}$ is complete (anticomplete) to X .

A *clique* in G is a set of pairwise adjacent vertices, and an *independent set* is a set of pairwise non-adjacent vertices. A *triangle* is a clique of size three. A *path* in G is an induced subgraph isomorphic to a graph P with $k + 1$ vertices p_0, p_1, \dots, p_k and with $E(P) = \{p_i p_{i+1} : i \in \{0, \dots, k-1\}\}$. We write $P = p_0 p_1 \dots p_k$ to denote a path with vertices p_0, p_1, \dots, p_k in order. We say that P is a path from p_0 to p_k . For a set $Y \subseteq V(G)$, if $P \setminus \{p_0, p_k\} \subseteq Y$, we say that P is a path from p_0 to p_k through Y . The *length* of a path P is the number of edges in P . A path is *odd* if its length is odd, and *even* otherwise. If $a, b \in P$, we denote by aPb the subpath of P from a to b . For a path P with ends a, b , the *interior* of P , denoted P^* , is the set $V(P) \setminus \{a, b\}$. For an integer $k \geq 4$, a

hole of length k in G is an induced subgraph isomorphic to the k -vertex cycle C_k . A hole is *odd* if its length is odd, and *even* if its length is even.

If $X, Y, Z \subseteq V(G)$ are such that $X \cap Z = \emptyset$, we say that the path $P = p_0 \dots p_k$ is a path from X to Z through Y if $p_0 \in X$, $V(P) \setminus \{p_0\} \subseteq Y$, $V(P) \setminus \{p_k\}$ is anticomplete to Z , and p_k has a neighbor in Z . When $X = \{p_0\}$, we say that P is a path from p_0 to Z through Y . Note that P is disjoint from Z , and possibly $P = p_0$ (when p_0 is a vertex from X with neighbors in Z). A path from X to Z through Y , when it exists, can be computed in time $\mathcal{O}(|V(G)|^2)$ as follows. In $G[X \cup Y \cup Z]$, compute a shortest path $q_0 \dots q_k$ from X to Z by a breadth-first search (that also terminates when no such path exists). Then, the path $q_0 \dots q_{k-1}$ is from X to Z through Y . Observe that this algorithm detects when no path from X to Z through Y exists (when no connected component of $G[X \cup Y \cup Z]$ contains vertices of both X and Z).

2 Star cutsets in (theta, pyramid, prism, turtle)-free graphs

Let G be a graph and H a hole in G . A *minor vertex* for H is a vertex $u \in G \setminus H$ such that u has neighbors in H and $N_H(u)$ is contained in a three-vertex path. A vertex $v \in G \setminus H$ is a *major vertex* for H if v has neighbors in H and v is not minor. A set $X \subseteq V(G)$ is a *star cutset* of G if $G \setminus X$ is not connected, and there exists $x \in X$ such that x is complete to $X \setminus x$. We call x the *center* of the star cutset X . The goal of this section is to prove that major vertices for holes of graphs $G \in \mathcal{C}$ are the centers of star cutsets of G . The following two lemmas describe major and minor vertices.

Lemma 2.1. *Let $G \in \mathcal{C}$. Every major vertex u for a hole H of G has at least four neighbors in H or has exactly three neighbors in H that are pairwise non-adjacent.*

Proof. Let u be a major vertex for a hole H . Suppose u has exactly two neighbors in H . The neighbors of u are non-adjacent, because they are not contained in a three-vertex path. Thus, $H \cup \{u\}$ forms a theta between the neighbors of u in H , a contradiction. If u has exactly three neighbors in H , they are pairwise non-adjacent, otherwise $H \cup \{u\}$ forms a pyramid. \square

Lemma 2.2. *Let $G \in \mathcal{C}$. Then, every minor vertex u for a hole H of G satisfies one of the following:*

- u has a unique neighbor in H (we say that u is a pendant of H)
- u has two adjacent neighbors in H (we say that u is a cap of H)
- u has three neighbors in H which induce a path xyz (we say that u is a clone of y in H)

Proof. Since u is minor, $N_H(u)$ is contained in a three-vertex path. Therefore, the only possibility not listed above is that u has two non-adjacent neighbors in H , in which case $H \cup \{u\}$ is a theta. \square

Suppose H is a hole of a graph G and u is a clone of y in H . We denote by $H_{u \setminus y}$ the hole of G induced by $(V(H) \setminus \{y\}) \cup \{u\}$. Note that y is a clone of u in $H_{u \setminus y}$.

Lemma 2.3. *Let H be a hole in a graph $G \in \mathcal{C}$, let v be a major vertex for H , let u be a clone of y in H , and suppose $uv \in E(G)$. If $yv \notin E(G)$, then u and v have a common neighbor in H .*

Proof. Let $N_H(u) = \{x, y, z\}$. Suppose $uv \in E(G)$ and $yv \notin E(G)$. Because v is major, by Lemma 2.1, v has at least three neighbors in H . If v is anticomplete to $\{x, z\}$, then $H \cup \{u, v\}$ is a uv -turtle in G , a contradiction. Therefore, v is adjacent to at least one of x, z , and so u and v have a common neighbor in H . \square

Let $u \in G \setminus H$ be a vertex with at least two neighbors in H . A u -*sector* is a path $P = u' \dots u''$ such that $P \subseteq H$, u' and u'' are neighbors of u , and P^* is anticomplete to u .

Lemma 2.4. *Let u and v be two non-adjacent major vertices for a hole H of a graph $G \in \mathcal{C}$. Let $P = u' \dots u''$ be a u -sector of H . Then one of the following holds:*

- (i) P contains at most one neighbor of v , and if it has one, it is either u' or u'' ,
- (ii) $u'u'' \in E(G)$ and v is adjacent to both u' and u'' ,
- (iii) P contains at least 3 neighbors of v ,
- (iv) $H \cup \{u, v\}$ is a cube (see Figure 3).




Figure 3: The cube graph

Proof. Let $R = H \setminus P$, let x be the neighbor of u' in R , and let y be the neighbor of u'' in R . First, suppose that P contains exactly two neighbors of v . We may assume that $u'u'' \notin E(G)$, otherwise outcome (ii) holds. Let the neighbors of v in P be given by v', v'' . Then, G contains a theta between v' and v'' in $P \cup \{u, v\}$ unless $v'v'' \in E(G)$, so $v'v'' \in E(G)$. Since either $v' \neq u', u''$ or $v'' \neq u', u''$, we may assume that $v' \neq u', u''$ and v' is between u' and v'' in P (possibly $v'' = u''$). If both u and v have neighbors in R^* , then G contains a pyramid from u to $vv'v''$ via paths through u', u'' , and R^* , so at least one of u, v has no neighbors in R^* . By Lemma 2.1, either $N_H(u) = \{x, u', u'', y\}$ or $N_H(v) = \{x, v', v'', y\}$. Since v is major, v has a neighbor in $R \setminus y$. Let v''' be the neighbor of v in $R \setminus y$ closest to x . If $ux \in E(G)$, then G contains a prism from uxu' to $v''v'v''$ through the paths $u-u''-P-v''$, $u'-P-v'$, and $x-R-v'''-v$, so $ux \notin E(G)$ and thus $N_H(v) = \{x, v', v'', y\}$. Now, G contains a pyramid from u' to $vv'v''$ via the paths $u'-x-v$, $u'-u-u''-P-v''$, and $u'-P-v'$, a contradiction.

Now, suppose P contains exactly one neighbor v' of v . We may assume that $v' \neq u', u''$, otherwise outcome (i) holds. If u and v both have neighbors in R^* , then G contains a theta between u and v' through u', u'' , and $R^* \cup \{v\}$, so at least one of u, v has no neighbors in R^* . By Lemma 2.1, either $N_H(u) = \{x, u', u'', y\}$ or $N_H(v) = \{x, v', y\}$. Since v is major, v has a neighbor in $R \setminus y$. Let v'' be the neighbor of v in R closest to x . If $ux \in E(G)$, then G contains a pyramid from v' to $u'ux$ through the paths $v'-P-u'$, $v'-v-v''-R-x$, and $v'-P-u''-u$, so $ux \notin E(G)$. By symmetry, $uy \notin E(G)$. It follows that $N_H(v) = \{x, v', y\}$. Because u is major, u has a neighbor in R^* . Let u''' be the neighbor of u in R^* closest to y . Then G contains a theta between u and y through the paths $u-u''-y$, $u-u'-x-v-y$, and $u-u'''-R-y$, unless $xu''' \in E(G)$. Similarly, G contains a theta between u and x unless $yu''' \in E(G)$. If $u'v' \notin E(G)$, then G contains a theta between u' and v' through the paths $u'-P-v'$, $u'-x-v-v'$, and $u'-u-u''-P-v'$, so $v'u' \in E(G)$. Similarly, $v'u'' \in E(G)$. Now, $H \cup \{u, v\}$ is a cube, and so outcome (iv) holds. This completes the proof. \square

Let H be a hole in a graph G and let $u, v \in V(G) \setminus V(H)$. We say that u and v are *nested* with respect to H if there exist distinct $a, b \in V(H)$ such that one ab -path of H contains all the neighbors of u and the other ab -path of H contains all the neighbors of v . Note that pendants, caps, and vertices of $G \setminus H$ with no neighbor in H are nested with all other vertices of $G \setminus H$. The vertices u and v are *strictly nested* with respect to H if u and v are nested with respect to H and $N_H(u) \cap N_H(v) = \emptyset$.

Lemma 2.5. *Let H be a hole in a graph $G \in \mathcal{C}$, let u and v be major or clones for H such that u and v are nested, and suppose $uv \in E(G)$. Then, u and v have a common neighbor in H .*

Proof. Since u and v are major or clones, u and v have at least three neighbors in H . If u and v have no common neighbors in H , then $H \cup \{u, v\}$ is a uv -turtle in G , a contradiction. \square

Two vertices u and v not in H that are not nested with respect to H are said to *cross*. The following lemmas characterize the behavior of vertices that cross.

Lemma 2.6. *Let H be a hole in a graph $G \in \mathcal{C}$ and let u and v be two vertices not in H that cross. Then, one of the following holds:*

- (i) H contains four distinct vertices u', v', u'', v'' that appear in this order along H such that $u', u'' \in N_H(u)$ and $v', v'' \in N_H(v)$,
- (ii) $N_H(u) = N_H(v)$, $N_H(u)$ is an independent set, $|N_H(u)| = 3$, and $uv \in E(G)$,
- (iii) $N_H(u) = N_H(v)$, both u and v are clones in H , and $uv \in E(G)$.

Proof. Since u and v are not nested, u and v are major or clones for H . Suppose u and v are both clones. Since u and v cross, it follows that either u and v are clones of adjacent vertices, so outcome (i) holds, or u and v are clones of the same vertex. Let u and v both be clones of y and let $N_H(u) = N_H(v) = \{x, y, z\}$. Then, G contains a theta between x and z in $H_{u \setminus y} \cup \{v\}$ unless $uv \in E(G)$, so outcome (iii) holds. Now, suppose u is a clone of y and v is major. Because u and v cross, $vy \in E(G)$, and because v is major, v has at least one neighbor in $V(H) \setminus N_H(u)$. Therefore, outcome (i) holds. Finally, suppose u and v are both major. Assume that u has a neighbor x in H such that $vx \notin E(G)$. Then, x is contained in a v -sector $P = v' \dots v''$ of H . Because u and v cross, u has a neighbor in $H \setminus P$, so outcome (i) holds. Therefore, we may assume that $N_H(u) = N_H(v)$. If $|N_H(u)| > 3$, outcome (i) holds, so $|N_H(u)| = 3$. Because u is major and $|N_H(u)| = 3$, it follows from Lemma 2.1 that $N_H(u)$ is an independent set. Let $N_H(u) = \{x, y, z\}$. Then, G contains a theta between x and y in $(H \setminus \{z\}) \cup \{u, v\}$ unless $uv \in E(G)$, so outcome (ii) holds. \square

Let $H = h_1-h_2-\dots-h_k-h_1$ be a hole in a graph $G \in \mathcal{C}$ and let $u, v \in V(G) \setminus V(H)$ be two non-adjacent major vertices for H . The following lemma shows that if u and v cross, then $H \cup \{u, v\}$ is a *major non-adjacent cross (MNC) configuration*. We describe MNC configurations as follows (see also Figure 4).

- MNC configuration (1): $k = 4$, and $\{u, v\}$ are complete to H .
- MNC configuration (2): $k = 5$, and $\{u, v\}$ are complete to H .
- MNC configuration (3): $k = 6$, $N_H(u) = \{h_1, h_3, h_5\}$, and $N_H(v) = \{h_2, h_4, h_6\}$.
- MNC configuration (4): $\{u, v\}$ is complete to $\{h_1, h_2, h_3, h_4\}$, v has no other neighbors in H , and u has at least one other neighbor in H .

Let $3 < i < k - 1$. Let H_1 be the path from h_1 to h_{i+1} in $H \setminus \{h_2\}$ and let H_2 be the path from h_2 to h_i in $H \setminus \{h_1\}$.

- MNC configuration (5): $\{u, v\}$ is complete to $\{h_1, h_2, h_i, h_{i+1}\}$, u and v both have other neighbors in H , $N_H(u) \subseteq H_1 \cup \{h_2, h_i\}$, and $N_H(v) \subseteq H_2 \cup \{h_1, h_{i+1}\}$.
- MNC configuration (6): $\{u, v\}$ is complete to $\{h_1, h_2, h_i, h_{i+1}\}$, v has no other neighbors in H , and u has neighbors both in H_1^* and H_2^* .
- MNC configuration (7): $\{u, v\}$ is complete to $\{h_1, h_2, h_i\}$, u and v both have other neighbors in H , $N_H(u) \subseteq H_1 \cup \{h_2, h_i\}$, and $N_H(v) \subseteq H_2 \cup \{h_1\}$.
- MNC configuration (8): $\{u, v\}$ is complete to $\{h_1, h_2\}$, u and v both have other neighbors in H , $N_H(u) \subseteq H_1 \cup \{h_2\}$, and $N_H(v) \subseteq H_2 \cup \{h_1\}$.




Figure 4: MNC configurations (dashed lines represent possible edges)

Lemma 2.7. *Let H be a hole in a graph $G \in \mathcal{C}$ and suppose that u and v are two major vertices for H . If $uv \in E(G)$, then either u and v cross or u and v have a common neighbor in H . If u and v cross, then either $uv \in E(G)$ or G contains an MNC configuration.*

Proof. The first statement follows from Lemma 2.5. It remains to prove the second statement. Assume that u and v cross and $uv \notin E(G)$. Let $H = h_1 - \dots - h_k - h_1$.

(1) *No three common neighbors of u and v in H form an independent set.*

If u and v have three common neighbors $x, y, z \in V(H)$ such that $\{x, y, z\}$ is an independent set, then $\{u, v, x, y, z\}$ is a theta in G between u and v , a contradiction. This proves (1).

(2) *Suppose there exist $i, j \in \{1, \dots, k\}$ with $j > i$ such that $h_i h_j \notin E(G)$, $\{u, v\}$ is complete to $\{h_i, h_j\}$, and there are no common neighbors of u and v in $P = h_{j+1} - h_{j+2} - \dots - h_{i-1}$. Then, u and v do not both have neighbors in P .*

If both u and v have neighbors in P^* , then G contains a theta between u and v , through h_i, h_j , and P^* . Therefore, we may assume that h_{j+1} is adjacent to u . If v has a neighbor in P^* , let v' be the neighbor of v in P^* closest to h_{j+2} . Then, G contains a pyramid from v to $uh_j h_{j+1}$ through the paths $v-h_j$, $v-h_i-u$, and $v-v'-P-h_{j+1}$, a contradiction. Therefore, we may assume that $N_P(v) = \{h_{i-1}\}$. Then, G contains a prism from $uh_j h_{j+1}$ to $h_i v h_{i-1}$ through the paths $u-h_i$, h_j-v , and $h_{j+1}-P-h_{i-1}$, a contradiction. This proves (2).

Let $N = N_H(u) \cap N_H(v)$.

(3) *We may assume that $R = H[N]$ is a subpath of H of length at most one.*

If $N = V(H)$, then by (1), $H \cup \{u, v\}$ is MNC configuration (1) or (2). So, we may assume that $N \neq V(H)$. Suppose first that R is a subpath of H of length greater than one. By (1), R is of length at most three. If it is of length two, then by (2), one of u, v is a clone, a contradiction. Suppose R is of length three, say $R = h_i - h_{i+1} - h_{i+2} - h_{i+3}$. Then, by (2), at most one of u, v has a neighbor in $H \setminus R$. If one of u, v has a neighbor in $H \setminus R$, then $H \cup \{u, v\}$ is MNC configuration (4), otherwise G contains a theta between h_i and h_{i+3} through u, v , and $H \setminus R$.

Next, suppose that R is not a subpath of H . By (1), it follows that R is the disjoint union of two subpaths Q_1, Q_2 of H of length at most one. Let $Q_1 = h_1 - \dots - h_i$ and $Q_2 = h_s - \dots - h_t$, where $k > t \geq s \geq i + 2$, $i \leq 2$, and $t \leq s + 1$. By (2), we may assume that v has no neighbors in $h_{t+1} - h_{t+2} - \dots - h_k$, and that not both u and v have neighbors in $h_{i+1} - \dots - h_{s-1}$. Since u and v cross, at least one of Q_1 and Q_2 has length one, so we may assume $i = 2$.

Note that at least one of u and v has a neighbor in $h_{t+1} \dots h_k$, otherwise G contains a theta between h_1 and h_t through $H \setminus \{h_2, h_s\}$, u , and v . Similarly, at least one of u and v has a neighbor in $h_{i+1} \dots h_{s-1}$. It follows that if $t = s + 1$, then G contains MNC configuration (5) or (6), and if $t = s$, then G contains MNC configuration (7). This proves (3).

(4) *We may assume that $N = \emptyset$.*

We may assume that $H \cup \{u, v\}$ is not a cube (MNC configuration (3)) or MNC configuration (8). By (3), we may assume that $R = h_1$ or $R = h_1 h_2$. Let $i = 1$ if $R = h_1$, and $i = 2$ if $R = h_1 h_2$. We claim that we may assume that $H \setminus R$ contains three distinct vertices u', v', u'' such that h_i, u', v', u'', h_1 appear in this order and $u', u'' \in N_H(u)$, $v' \in N_H(v)$. If $i = 2$, then this follows from the assumption that G does not contain MNC configuration (8), and if $i = 1$, it follows from Lemma 2.6. Let u', v', u'' be chosen such that the path from u' to u'' in $H \setminus R$ is a u -sector. Let H_a be the path from h_i to v' in $H \setminus \{u''\}$, let H_b be the path from h_1 to v' in $H \setminus \{u'\}$, and let H_u be the path from u' to u'' in $H \setminus R$. Since H_a contains a v -sector that contains u' , it follows from Lemma 2.4 that there are at least three neighbors of u in H_a , and so there at least two neighbors of u in H_a^* . Since H_u is a u -sector, there is a neighbor of u in H_a^* between h_i and u' . Similarly, there are at least three neighbors of u in H_b , and so at least two neighbors of u in H_b^* . Since H_u is a u -sector, there is a neighbor of u in H_b^* between h_1 and u'' . Finally, since H_u is a u -sector that contains v' , there are at least three neighbors of v in H_u^* . Let v_1 be the neighbor of v in H_u^* closest to u' , and let v_2 be the neighbor of v in H_u^* closest to u'' . Then, G contains a theta from u to v , through $u-h_1-v$, $u-u'-H_u-v_1-v$, and $u-u''-H_u-v_2-v$. This proves (4).

It follows from Lemma 2.6 and (4) that H contains four distinct vertices u', v', u'', v'' that appear in that order along H , such that $u', u'' \in N_H(u)$ and $v', v'' \in N_H(v)$. By (4), u and v have no common neighbors in H , and so every neighbor of u in H is in the interior of a v -sector and every neighbor of v in H is in the interior of a u -sector. Let u', v', u'', v'' be chosen so that $P = u' \dots u''$ is a u -sector and $Q = v' \dots v''$ is a v -sector. We may assume that $H \cup \{u, v\}$ is not a cube (i.e. MNC configuration (3)). Because v' is in P^* , it follows from Lemma 2.4 that there are at least three neighbors of v in P^* . Since there are no neighbors of v in Q^* , we may assume that v_1, v_2 are neighbors of v in P^* between u' and v' in that order. Similarly, there are at least three neighbors of u in Q^* . Since there are no neighbors of u in P^* , we may assume that u_1, u_2 are neighbors of u in Q^* between u'' and v'' in that order. Finally, because v'' is in the interior of a u -sector and $Q = v' \dots v''$ is a v -sector, there is another neighbor of v between v'' and u' . Then, G contains a theta between u and v through the paths $u-u_2-Q-v''-v$, $u-u''-Q-v'-v$, and $u-u'-P-v_1-v$, a contradiction. \square

Lemma 2.8. *Let H be a hole of length greater than six in a graph $G \in \mathcal{C}$ and suppose u and v are non-adjacent vertices of $G \setminus H$ that cross. Then, H contains a $\{u, v\}$ -complete edge.*

Proof. Since u and v cross, u and v are major or clones. If u and v are both major, it follows from Lemma 2.7 that $H \cup \{u, v\}$ is MNC configuration (4), (5), (6), (7), or (8), so H contains a $\{u, v\}$ -complete edge. Now, suppose u is a clone of y in H and $N_H(u) = \{x, y, z\}$. Because u and v cross, it follows that $vy \in E(G)$. We may assume that $xv, zv \notin E(G)$ since otherwise H contains a $\{u, v\}$ -complete edge. Note that $x-u-z$ is a subpath of $H_{u \setminus y}$ that contains all the neighbors of y in $H_{u \setminus y}$ and no neighbors of v . If v has at least three neighbors in $H_{u \setminus y}$, then G contains a vy -turtle. So v has two neighbors in $H_{u \setminus y}$, say v_1 and v_2 , and hence three neighbors in H . By Lemma 2.1 applied to H and v , $v_1 v_2 \notin E(G)$. But then $H_{u \setminus y}$ and v form a theta, a contradiction. \square

The following lemma describes the behavior of paths whose endpoints are nested with respect to H and whose internal vertices are anticomplete to H .

Lemma 2.9. *Let H be a hole in a graph $G \in \mathcal{C}$ and let $P = u \dots v$ be a path of length at least 1, vertex-disjoint from H , such that u and v have neighbors in H and are nested with respect to H , and no internal vertex of P has a neighbor in H . Then, u and v have a common neighbor in H , or u and v are both pendants of H with adjacent neighbors in H . In particular, if u and v are strictly nested with respect to H , then u and v are both pendants of H with adjacent neighbors in H .*

Proof. We may assume that u and v do not have a common neighbor in H . If u (resp. v) has at least two neighbors in H , then let u' and u'' (resp. v' and v'') be the endpoints of the u -sector (resp. v -sector) that contains all neighbors of v (resp. u) in H , and otherwise let $u' = u''$ (resp. $v' = v''$) be its unique neighbor in H . Without loss of generality, u', v', v'', u'' appear in this order along H . If u and v both have two non-adjacent neighbors in H (and so by Lemma 2.1 and Lemma 2.2, u and v each has at least three neighbors in H) and $uv \in E(G)$, then $H \cup \{u, v\}$ is a uv -turtle, a contradiction. If u and v both have two non-adjacent neighbors in H and $uv \notin E(G)$, then G contains a theta between u and v through the paths $u-u'-H \setminus \{u'', v''\}-v'-v$, $u-u''-H \setminus \{u', v'\}-v''-v$, and $u-P-v$, a contradiction. If u has two non-adjacent neighbors in H and v is a cap, then G contains a pyramid from u to $vv'v''$ through $u-P-v$, $u-u'-H \setminus \{u''\}-v'$, and $u-u''-H \setminus \{u'\}-v''$, a contradiction. If u has two non-adjacent neighbors in H and v is a pendant, then G contains a theta between u and v' through $u-u'-H \setminus \{u''\}-v$, $u-P-v-v'$, and $u-u''-H \setminus \{u'\}-v'$, a contradiction. Thus, neither u nor v has two non-adjacent neighbors in H .

If u and v are both caps, then $H \cup P$ is a prism between $uu'u''$ and $vv'v''$, a contradiction. If u is a cap and v is a pendant, then $H \cup P$ is a pyramid from v' to $uu'u''$. If u and v are both pendants, then $H \cup P$ is a theta between u' and v' , unless $u'v'$ is an edge. \square

Let $G \in \mathcal{C}$, let H be a hole in G , and let w be a major vertex for H . A path $N \subseteq H$ is an *extended neighborhood* of w in H if there exists a w -sector $Q = x \dots y$ such that $N = Q \cup (\{x', y'\} \cap N_H(w))$, where x' and y' are the neighbors of x and y in $H \setminus Q$, respectively. Two vertices $a, b \in H$ are *distant in H with respect to w* if a, b are not contained in an extended neighborhood of w in H . Note that if a vertex $v \in H$ is not adjacent to w , then v is in exactly one extended neighborhood of w in H .

Suppose H is a hole in a graph $G \in \mathcal{C}$ and u is a major vertex for H . The vertex u is called a *hub* if $N_H(u) = \{x, y, z, w\}$ where the vertices x, y, z, w appear in that order in H , $xy, zw \in E(G)$, and $xw, zy \notin E(G)$.

Lemma 2.10. *Let $G \in \mathcal{C}$, let H be a hole in G of length greater than six, and let w be a major vertex for H . Let $p \in V(G) \setminus V(H)$ be such that $pw \notin E(G)$. Then, either $N_H(p)$ is contained in an extended neighborhood of w in H or $H \cup \{p, w\}$ is MNC configuration (6).*

Proof. If p and w are nested, then $N_H(p)$ is contained in an extended neighborhood of w in H , so we may assume that p and w cross. It follows that p is either a clone or a major vertex for H . If p is a clone, then it follows from Lemma 2.8 that $N_H(p)$ is contained in an extended neighborhood of w . Now, suppose p is major. By Lemma 2.7, it follows that $H \cup \{w, p\}$ is MNC configuration (4), (5), (6), (7), or (8), and therefore either $N_H(p)$ is contained in an extended neighborhood of w in H , or $H \cup \{w, p\}$ is MNC configuration (6). \square

Let $G \in \mathcal{C}$, let H be a hole in G , and let w be a major vertex for H . We say that a path $P = p_1 \dots p_k$ is (H, w) -*significant* if there exist $a, b \in V(H)$ such that $a \in N_H(p_1) \setminus N_H(w)$, $b \in N_H(p_k)$, and a and b are distant in H with respect to w .

Lemma 2.11. *Let $G \in \mathcal{C}$, let H be a hole in G of length greater than six, and let w be a major vertex for H such that either w is not a hub or every major vertex for H is a hub. Let $P = p_1 \dots p_k$ be (H, w) -significant with $a, b \in V(H)$ as in the definition of a significant path. Let $Q = x \dots y$*

be the w -sector containing a . Suppose w is anticomplete to $\{p_1, p_k\}$ and $p_1, p_k \notin H$. Then, p_k is anticomplete to Q^* .

Proof. Since p_1 has a neighbor in the interior of a w -sector, $H \cup \{w, p_1\}$ is not MNC configuration (6). By Lemma 2.10, it follows that $N_H(p_1)$ is contained in an extended neighborhood of w . Note that, by definition, a w -sector of length greater than one is contained in exactly one extended neighborhood of w . Let \overline{Q} be the extended neighborhood of w containing Q , and suppose for sake of contradiction that p_k has a neighbor in Q^* . Since p_k has a neighbor in the interior of a w -sector, $H \cup \{w, p_k\}$ is not MNC configuration (6), so by Lemma 2.10, $N_H(p_k)$ is contained in an extended neighborhood of w . Because p_1 has a neighbor a in Q^* , it follows that $N_H(p_1) \subseteq \overline{Q}$. Similarly, because p_k has a neighbor in Q^* , it follows that $N_H(p_k) \subseteq \overline{Q}$. Then, a and b are contained in an extended neighborhood of w , a contradiction. \square

Lemma 2.12. *Let $G \in \mathcal{C}$, let H be a hole in G of length greater than six, and let w be a major vertex for H such that either w is not a hub or every major vertex for H is a hub. Let $P = p_1 \dots p_k$ be (H, w) -significant with $a, b \in V(H)$ as in the definition of a significant path. Assume that P^* is anticomplete to H , w is anticomplete to $\{p_1, p_k\}$, and $p_1, p_k \notin H$. Then, p_1 and p_k have a common neighbor in H .*

Proof. Assume for a contradiction that p_1 and p_k do not have a common neighbor in H . In particular, $p_1 \neq p_k$. Suppose that p_1 and p_k are nested. Then, they are strictly nested. By Lemma 2.9, p_1 and p_k are pendants of H with adjacent neighbors in H . It follows that a and b are adjacent, contradicting that a and b are distant in H with respect to w . Therefore, p_1 and p_k cross, and hence they are clone or major for H . Since p_1 and p_k cross and they do not have a common neighbor in H , it follows that p_1 and p_k are both major. If w and p_1 are nested, then by Lemma 2.11, p_1 and p_k are nested. Hence, w and p_1 cross. By Lemma 2.8, H contains a $\{w, p_1\}$ -complete edge. Let $Q = x \dots y$ be the w -sector containing a . Let x' and y' be the neighbors of x and y in $H \setminus Q$, respectively. Since p_1 has a neighbor in the interior of a w -sector, $H \cup \{w, p_1\}$ is not MNC configuration (6). By Lemma 2.10, it follows that $N_H(p_1) \subseteq Q \cup (\{x', y'\} \cap N_H(w))$, and by Lemma 2.11, it follows that $N_H(p_k) \subseteq H \setminus Q^*$. Up to symmetry, suppose $\{w, p_1\}$ is complete to $\{x, x'\}$. Since p_1 and p_k cross and p_1 and p_k have no common neighbor in H , $p_1 y', p_k y \in E(G)$ and $p_1 y, p_k y' \notin E(G)$. Also, p_k has another neighbor in $H \setminus (Q \cup \{x', y'\})$ and in particular, $x' y' \notin E(G)$. Because $p_1 y' \in E(G)$, it follows that $w y' \in E(G)$. Let p' be the neighbor of p_1 in Q^* closest to y . Then, G contains a pyramid from p_1 to $w y'$ through $p_1 y', p_1 x' w$, and $p_1 p' Q y$, a contradiction. \square

We can now prove the main result of this section.

Theorem 2.13. *Let $G \in \mathcal{C}$, let H be a hole in G of length greater than six, and let w be a major vertex for H such that either w is not a hub or every major vertex for H is a hub. Let $P = p_1 \dots p_k$ be (H, w) -significant. Then, w has a neighbor in P .*

Proof. We may assume that no subpath of P is (H, w) -significant. Suppose that w is anticomplete to P . Let $a, b \in V(H)$ be as in the definition of a significant path, i.e., $a \in N_H(p_1) \setminus N_H(w)$, $b \in N_H(p_k)$, and a and b are distant in H with respect to w . Let $Q = x \dots y$ be the w -sector containing a , and let x' and y' be the neighbors of x and y in $H \setminus Q$, respectively. Possibly $x' = b$ (resp. $y' = b$), in which case w is not adjacent to x' (resp. y') since a and b are distant in H with respect to w .

Suppose that $k = 1$. Since a and b are distant in H , $p_1 \notin V(H)$. So, by Lemma 2.11 and Lemma 2.2, p_1 is major but not a hub. It follows that w is not a hub. But this contradicts Lemma 2.10. Therefore, $k > 1$.

(1) P is disjoint from H .

Suppose that $p_i \in P$ is in H . Since no subpath of P is (H, w) -significant, it follows that a and p_i are not distant in H with respect to w . Then, either $p_i \in Q^*$ or $p_i w \in E(G)$. Since w is anticomplete to P , $p_i w \notin E(G)$, so $p_i \in Q^*$. Note that since $p_i \in Q^*$ and $b \notin Q$, $i < n$. Then, $p_{i+1} \dots p_k$ is (H, w) -significant, a contradiction. This proves (1).

(2) Either $N_H(P^*) \subseteq \{x\} \cup (N_H(w) \cap \{x'\})$ or $N_H(P^*) \subseteq \{y\} \cup (N_H(w) \cap \{y'\})$.

No vertex $p_i \in P \setminus \{p_1\}$ can have a neighbor in Q^* , otherwise $p_i \dots p_k$ is (H, w) -significant. Similarly, no vertex $p_i \in P \setminus \{p_k\}$ can be adjacent to a vertex v such that a and v are distant in H with respect to w , otherwise $p_1 \dots p_i$ is (H, w) -significant. It follows that $N_H(P^*) \subseteq \{x, y\} \cup (\{x', y'\} \cap N_H(w))$.

Consider a vertex $p_i \in P^*$ such that p_i has neighbors in H . Suppose p_i is not a cap or a pendant, so by Lemmas 2.1 and 2.2, $N_H(p_i) = \{x, y, x', y'\}$. Then, since $Q = x \dots a \dots y$ is a w -sector and $p_i x, p_i y \in E(G)$, by Lemma 2.4 it follows that $xy \in E(G)$, a contradiction. Therefore, p_i is a cap or a pendant for every $p_i \in P^*$ that has a neighbor in H .

Now, assume that there exist $p_i, p_j \in P^*$ such that $N_H(p_i) \subseteq \{x\} \cup (N_H(w) \cap \{x'\})$ and $N_H(p_j) \subseteq \{y\} \cup (N_H(w) \cap \{y'\})$. Consider the shortest path R from $\{x\} \cup (N_H(w) \cap \{x'\})$ to $\{y\} \cup (N_H(w) \cap \{y'\})$ through P^* . By Lemma 2.9, the endpoints of R^* have a common neighbor in H or are pendants of H with adjacent neighbors, a contradiction since $b \in H \setminus (Q \cup (\{x', y'\} \cap N_H(w)))$. This proves (2).

In view of (2), we assume from now on that $N_H(P^*) \subseteq \{x\} \cup (N_H(w) \cap \{x'\})$. Let H_x and H_y be the paths in H from a to b through x and y , respectively. Since p_1 has a neighbor in the interior of a w -sector, $H \cup \{w, p_1\}$ is not MNC configuration (6). By Lemma 2.10, it follows that $N_H(p_1) \subseteq Q \cup (\{x', y'\} \cap N_H(w))$, and by Lemma 2.11, it follows that $N_H(p_k) \subseteq H \setminus Q^*$. Therefore, if p_1 and p_k have a common neighbor v in H_y , then $v \in \{y, y'\}$. Let $r_1 = r_s = v$ if p_1 and p_k have a common neighbor v in H_y . Otherwise, let r_1 be the neighbor of p_1 in H_y that is furthest from a , and let r_s be the neighbor of p_k in H_y that is furthest from b . Let $R = r_1 \dots r_s$ be the path from r_1 to r_s through H_y . Note that r_1 is between a and r_s unless $r_1 = y'$ and $r_s = y$, in which case $p_k y', p_1 y \notin E(G)$. It follows that $P \cup R$ is a hole when P has length at least two.

(3) w has a neighbor in $R \cap \{y, y'\}$.

Because $N_H(p_1) \subseteq Q \cup (\{x', y'\} \cap N_H(w))$, it follows that $N_{H_y}(p_1) \subseteq Q \cup (\{y'\} \cap N_H(w))$. If $r_1 \neq y'$, then $y \in R$ and $wy \in E(G)$, so w has a neighbor in $R \cap \{y, y'\}$. If $r_1 = y'$, then $p_1 y' \in E(G)$, so $wy' \in E(G)$, and w has a neighbor in $R \cap \{y, y'\}$. This proves (3).

(4) If $P \cup R$ is a hole, then x is anticomplete to $P \cup R$.

Let J be the hole given by $P \cup R$. We prove a number of subclaims.

(4.1) $x' \notin J$.

Suppose $x' \in J$. Then $x' = r_s = b$, and so x' is non-adjacent to w , since otherwise a and b are not distant in H with respect to w . By Lemma 2.9 applied to J and the path $x-w$, it follows that w and x are strictly nested with respect to the hole J , and so x and w are both pendants of J with adjacent neighbors in J . Let $N_J(w) = \{x''\}$, then x'' is the neighbor of x' in $H \setminus x$. Since w has a unique neighbor in J , it follows from (3) that $x'' \in \{y, y'\}$. If $x'' = y'$, then $H = y-Q-x-x'-x''-y$ and $N_H(w) = \{x, y, y'\}$, so $H \cup \{w\}$ is a pyramid, a contradiction. So $x'' = y$. But now $H = y-Q-x-x'-y$ and $N_H(y) = \{x, y\}$, so $H \cup \{w\}$ is a theta, a contradiction. This proves (4.1).

(4.2) If w is a hub, then p_1 is anticomplete to $\{x', y'\}$.

Suppose w is a hub and p_1 is adjacent to y' (the argument is similar if p_1 is adjacent to x'). Since p_1 is adjacent to a and to y' , we deduce that p_1 is either a clone of y or p_1 is major for H . Since w is a hub, it follows that if p_1 is major for H , then p_1 is a hub for H . In both cases, $N_H(p_1) \setminus Q = \{y'\}$.

Then, G contains a pyramid from y' to $xx'w$ through the paths $y'-w$, $y'-p_1-Q-x$, and the path from y' to x' with interior in $H \setminus Q$, a contradiction. This proves (4.2).

Suppose that x has a neighbor in J . Since, by (4.1), $x' \notin J$, it follows that x is anticomplete to R , so x has a neighbor in P . We apply Lemma 2.9 to J and the path $x-w$. Since w is anticomplete to P , we have that x and w have no common neighbor in J . Consequently, w and x are strictly nested with respect to J , and so x and w are both pendants of J with adjacent neighbors in J . Since x is anticomplete to R , and w is anticomplete to P , there are two possibilities:

1. $N_J(x) = \{p_k\}$, $N_J(w) = \{r_s\}$, or
2. $N_J(x) = \{p_1\}$, $N_J(w) = \{r_1\}$.

Suppose the former holds. By (3), $r_s \subseteq \{y, y'\} \cap N(w)$. If p_k is adjacent to y (so $r_s = y$), then G contains a theta between x and r_s given by the paths $x-w-r_s$, $x-p_k-r_s$ and $x-Q-r_s$, a contradiction. It follows that p_k is non-adjacent to y , and so $r_s = y'$. Then, G contains a pyramid from x to $yy'w$ through the paths $x-w$, $x-p_k-r_s$ and $x-Q-y$, a contradiction. This proves that the former case does not hold, and therefore the latter holds.

By (3), $r_1 \in \{y, y'\}$. If $r_1 = y$, let $r'_1 = y'$, and if $r_1 = y'$ let r'_1 be the neighbor of r_1 in $H \setminus \{y\}$. Let M be the subpath of $H \setminus \{a\}$ from r'_1 to x' . Suppose that both w and p_k have neighbors in M^* . Then there is a path M' from w to p_k with $M'^* \subseteq M^*$. Now, G contains a theta between p_1 and w through the paths p_1-r_1-w , p_1-x-w and $p_1-P-p_k-M'-w$, a contradiction. This proves that either p_k or w is anticomplete to M^* .

(4.3) w has no neighbor in M^* .

Suppose that w has a neighbor in M^* . Then, p_k is anticomplete to M^* . Since w has a neighbor in M^* , and $N_R(w) = \{r_1\}$, it follows that $r_s \neq x'$ and r_s is non-adjacent to x' . Consequently, $r_s \in \{y, y', r'_1\}$. Since $r_1 \in \{y, y'\} \cap N(w)$ and a and b are distant in H with respect to w , it follows that either $b = r'_1$, or $b = x'$ and x' is non-adjacent to w . Also, since w has a unique neighbor in J , it holds that if $r_s = y$ then $r_1 = y$.

Suppose x' has a neighbor in $P \setminus p_1$. Then, there is a path P' from x' to p_k with interior in P^* . It follows from the minimality of k that P' is not (H, w) -significant. If x' is non-adjacent to w , then x' and r_s are distant in H with respect to w since w has a neighbor in M^* , and so P' is (H, w) -significant, a contradiction. It follows that x' is adjacent to w , and so $x' \neq b$ and $b = r'_1$. Suppose $r'_1 \in R$. Then, w is non-adjacent to r'_1 since $N_R(w) = \{r_1\}$. Now, we get a contradiction applying Lemma 2.9 to the path $x'-w$ and the hole J . This implies that $r'_1 \notin R$, and so $r_1 = r_s$. (Indeed, if $r_1 \neq r_s$, then $r_1 = y$, $r_s = y'$, $p_k y \notin E(G)$, $wy' \notin E(G)$, $r'_1 = b = y'$, and hence the path $x'-w$ and the hole J contradict Lemma 2.9.) Since J is a hole, we have $k > 2$. Again, by Lemma 2.9 applied to the path $x'-w$ and the hole J , it follows that x' is a pendant for J , and $N_J(x') = p_k$. But now G contains a pyramid from r_1 to $xx'w$ with paths r_1-p_1-x , r_1-w and r_1-p_k-x' , a contradiction. This proves that x' is anticomplete to $P \setminus p_1$.

Since $N_H(P^*) \subseteq \{x\} \cup (N_H(w) \cap \{x'\})$, x' is anticomplete to $P \setminus p_1$, and p_k is anticomplete to M^* , it follows that $P \setminus p_1$ is anticomplete to $M \setminus r'_1$. Since x' is non-adjacent to p_k , it follows that $b = r'_1$. Since $k > 1$ and by minimality of k , we deduce that p_1 is non-adjacent to r'_1 . If $r'_1 \in R$, then G contains a theta between p_1 and r'_1 through the paths $p_1-r_1-r'_1$, $p_1-x-H_x-r'_1$ (possibly shortcutting through the edge p_1x'), and $p_1-P-p_k-r'_1$, a contradiction. This proves that $r'_1 \notin R$, and so $r_s \in \{y, r_1\}$. Since $N_J(x) = \{p_1\}$, x' is anticomplete to $P \setminus \{p_1\}$, and $N_H(P^*) \subseteq \{x, x'\}$, it follows that P^* is anticomplete to H . By Lemma 2.12, p_1 and p_k have a common neighbor in H . It follows that either $r_1 = r_s = y$ or $r_1 = r_s = y'$. Since $P \cup R$ is a hole, it follows that $k > 2$. But now G contains a pyramid from p_1 to $r_1r'_1p_k$ through the paths p_1-r_1 , p_1-P-p_k , and $p_1-x-H_x-r'_1$ (possibly shortcutting through p_1x'). This proves (4.3).

(4.4) w is a hub for H .

If w has no neighbor in $M \setminus \{x', y'\}$, then by Lemma 2.1, w is a hub for H . Suppose w has a neighbor in $M \setminus \{x', y'\}$. Since, by (4.3), w is anticomplete to M^* , it follows that the only neighbor of w in $M \setminus \{x', y'\}$ is r'_1 . Thus, $r'_1 \neq y'$, and so $r_1 = y'$. Since $r_1 = y'$ (and thus p_1 is adjacent to y'), it follows that w is adjacent to y' . Since $N_J(w) = \{r_1\}$, it holds that $r_1 = r_s = y'$, and since J is a hole, P has length at least two. If w is non-adjacent to x' , then p_1 is non-adjacent to x' (as $N_H(p_1) \subseteq Q \cup (\{x', y'\} \cap N_H(w))$) and hence G contains a pyramid from x to $y'r'_1w$ through $x-p_1-y'$, $x-w$, and $x-x'-M^*-r'_1$, a contradiction. Therefore, w is adjacent to x' . Suppose that the only neighbor of p_k in M is r'_1 . Then, G contains a pyramid from p_1 to $p_k r_1 r'_1$ through p_1-r_1 , p_1-P-p_k (recall that P has length at least two), and $p_1-x-x'-H_x-r'_1$ (possibly shortcutting through the edge p_1-x'), a contradiction. So p_k has a neighbor in M different from r'_1 . Let b' be the neighbor of p_k in M closest to x' . Now, G contains a pyramid from y' to $xx'w$ through $y'-w$, $y'-p_1-x$, and $y'-p_k-b'-M-x'$, a contradiction. This proves (4.4).

It follows that w is a hub and $N_H(w) = \{x, x', y, y'\}$. Consequently, by (4.2), p_1 is anticomplete to $\{x', y'\}$, and so $N_H(p_1) \subseteq Q$. Since $r_1 \in \{y, y'\}$ and p_1 is non-adjacent to y' , we deduce that $r_1 = y$. Since $N_J(w) = \{r_1\}$, it follows that $y' \notin R$, and so $r_s = y$. Since $(H \setminus \{y'\}) \cup \{w, p_k\}$ is not a pyramid, it follows that p_k is not a clone of y' . Since a and b are distant in H with respect to w , it follows that $b \in H \setminus (Q \cup \{x', y'\})$. Since p_k is adjacent to y and to b , it holds that p_k is a major vertex for H , and so p_k is a hub by the assumption of the theorem. Consequently, p_k is adjacent to y' , and p_k is non-adjacent to x . Since $P \cup R$ is a hole and $r_1 = r_s$, it follows that $k > 2$. But now G contains a pyramid from y to wxx' given by paths $y-w$, $y-p_1-x$ and $y-p_k-H_x-x'$. This proves that x is anticomplete to J and completes the proof of (4).

(5) If x' is not anticomplete to $P \setminus p_k$, then $p_1x \in E(G)$.

Assume x' has a neighbor in $P \setminus p_k$, but $p_1x \notin E(G)$. By our assumption, $N_H(P^*) \subseteq \{x\} \cup (N_H(w) \cap \{x'\})$, and by Lemma 2.10, $N_H(p_1) \subseteq Q \cup (\{x', y'\} \cap N_H(w))$. Hence, $wx' \in E(G)$. Let z be the neighbor of x' in P closest to p_1 . Note that if $z \neq p_1$, then P is of length at least two, so $P \cup R$ is a hole and by (4), x is anticomplete to P . In particular, x is anticomplete to p_1-P-z . Consider the triangle given by wxx' . If $N_Q(p_1) = a$, then G contains a pyramid from a to wxx' through $a-Q-y-w$, $a-Q-x$, and $a-P-z-x'$, a contradiction. Suppose p_1 has two non-adjacent neighbors in Q and let q and q' be the neighbors of p_1 in Q closest to x and y , respectively. Then, G contains a pyramid from p_1 to wxx' through $p_1-q'-Q-y-w$, $p_1-q-Q-x$, and $p_1-P-z-x'$, a contradiction. Finally, suppose p_1 has exactly two adjacent neighbors in Q and let $N_H(p_1) = \{q, q'\}$, where q is between x and q' in Q . Then, G contains a prism between p_1qq' and $x'xw$ through $p_1-P-z-x'$, $q-Q-x$, and $q'-Q-y-w$, a contradiction. This proves (5).

(6) If $P \cup R$ is a hole, then $\{x, x'\}$ is anticomplete to $P \setminus p_k$. In particular, $N_H(P^*) = \emptyset$.

Suppose $P \cup R$ is a hole. By (4), x is anticomplete to P . If x' has neighbors in $P \setminus p_k$, then, by (5), $p_1x \in E(G)$, contradicting that x is anticomplete to P . This proves the first assertion. Next, suppose that $N_H(P^*) \neq \emptyset$. Then, $P^* \neq \emptyset$, and so $P \cup R$ is a hole. Now, by the first assertion, $\{x, x'\}$ is anticomplete to $P \setminus p_k$. But $N_H(P^*) \subseteq \{x, x'\}$, a contradiction. This proves (6).

By (6), $N_H(P^*) = \emptyset$, and so the symmetry between x and y is restored. Let $T = N_H(p_1) \cap N_H(p_k)$. By Lemma 2.12, $T \neq \emptyset$. Because $N_H(p_1) \subseteq Q \cup \{x', y'\}$ and $N_H(p_k) \subseteq H \setminus Q^*$, it follows that $T \subseteq \{x, x', y, y'\}$. Suppose first that one of $T \cap \{x, x'\}$ and $T \cap \{y, y'\}$ is empty. We may assume up to symmetry that $T \subseteq \{x, x'\}$. Because p_1 and p_k do not have a common neighbor in $\{y, y'\}$, it follows that $P \cup R$ is a hole. Then, by (6), $\{x, x'\}$ is anticomplete to p_1 , a contradiction. Therefore, we may assume that $T \cap \{x, x'\} \neq \emptyset$ and $T \cap \{y, y'\} \neq \emptyset$. By Lemma 2.4 and since p_k is anticomplete to Q^* , it follows that p_k is adjacent to at most one of x and y , and so not both x and

y are in T . Suppose that x' and y are both in T . Then, w has three neighbors in the hole given by $x'-x-Q-y-p_k-x'$ and w is not a clone or a major vertex for this hole, contradicting Lemmas 2.1 and 2.2. This proves that not both x' and y are in T . By symmetry, not both y and x' are in T . It follows that $T = \{x', y'\}$. Because p_1 and w are major and non-adjacent, and p_1 is adjacent to x' and y' , it follows by Lemma 2.7 that $H \cup \{p_1, w\}$ is MNC configuration (5). Therefore, p_1 is adjacent to x and y . Since $T = \{x', y'\}$, it follows that $\{x, y\}$ is anticomplete to $V(P) \setminus \{p_1\}$. Further, because p_1 is not a hub, it follows that w is not a hub, so w has neighbors in $H \setminus (Q \cup \{x', y'\})$. Note also that $b \in H \setminus (Q \cup \{x', y'\})$. Then, G contains a theta between p_1 and w , through x, y , and $P \cup (H \setminus (Q \cup \{x', y'\}))$, a contradiction. \square

Let $H = h_1-h_2-\dots-h_k-h_1$ be a hole in a graph $G \in \mathcal{C}$ and let $v \in V(G)$. We say that v is a *gem-center* if $k \geq 5$ and $N_H(v) = \{h_1, h_2, h_3, h_4\}$.

Corollary 2.14. *Let $G \in \mathcal{C}$ and let H be a hole in G of length greater than six. Let w be a major vertex for H such that w is not complete to H , w is not a gem-center, and either w is not a hub or every major vertex for H is a hub. Then, w is the center of a star cutset in G .*

Proof. Let $u \in H$ such that $uw \notin E(G)$. We claim that there exists a vertex $v \in H$ such that u and v are distant in H with respect to w . Suppose otherwise. Let Q be the w -sector containing u and let \overline{Q} be the extended neighborhood of w containing Q . It follows that $\overline{Q} = H$, so $N_H(w)$ is contained in a subpath of H of length at most three. Since w is major, it follows that $N_H(w)$ is a subpath of H of length exactly three, so w is a gem-center, a contradiction. Let $v \in H$ be such that u and v are distant in H with respect to w . It follows from Theorem 2.13 that w has a neighbor in the interior of every path from u to v . Therefore, u and v are in different components of $G \setminus (N[w] \setminus v)$, so w is the center of a star cutset in G . \square

3 Structure of proper separators

In this section, we consider minimal separators of graphs in \mathcal{C} . We start with the following result concerning minimal separators that are cliques.

Lemma 3.1 ([4]). *For every graph G , there are at most $\mathcal{O}(|V(G)|)$ minimal clique separators of G and they can be enumerated in time $\mathcal{O}(|V(G)||E(G)|)$.*

A separator in a graph is *proper* if it is minimal and not a clique. By Lemma 3.1, we restrict our attention here to proper separators. Our goal is to prove that graphs in \mathcal{C} have polynomially many proper separators.

Let C be a minimal separator of a graph G . A connected component D of $G \setminus C$ is a *full component for C* if every vertex of C has a neighbor in D , i.e., $N(D) = C$. Recall that there are at least two full components for every minimal separator. The next lemma, while not necessary for our results, is a convenient observation about full components for proper separators of graphs in \mathcal{C} .

Lemma 3.2. *If C is a proper separator of a graph $G \in \mathcal{C}$, then there are exactly two full components for C .*

Proof. Let c_1c_2 be a non-edge in C , and suppose that there are three full components for C . Then, G contains a path from c_1 to c_2 through each of the three full components, and so G contains a theta between c_1 and c_2 , a contradiction. \square

For the rest of this section, we let C be a proper separator of a graph $G \in \mathcal{C}$, and we denote by L and R the two full components for C . Let H be a hole with $V(H) \cap V(C) = \{c_1, c_2\}$, and let H_L and H_R be the two paths of H between c_1 and c_2 . We say that H is a (C, c_1, c_2) -hole if $H_L^* \subseteq L$ and $H_R^* \subseteq R$. A vertex $v \in V(G)$ is (c_1, c_2) -heavy with respect to H if v is major for H and c_1, c_2 are distant in H with respect to v . Note that if $v \in V(G)$ is (c_1, c_2) -heavy with respect to H , then v has a neighbor in H_L^* and a neighbor in H_R^* , and therefore $v \in C$. The frame of H is given by $F(H) = (c_1, c_2, \ell'_1, \ell_1, r_1, r'_1, \ell'_2, \ell_2, r_2, r'_2)$, where ℓ_1 is the neighbor of c_1 in H_L , ℓ'_1 is the neighbor of ℓ_1 in H_L^* if $\ell_1 \neq \ell_2$, and otherwise $\ell'_1 = \ell_1 = \ell_2$. We define similarly $\ell_2, \ell'_2, r_1, r'_1, r_2, r'_2$. We denote by $V(F)$ the vertices of F . We call F a (C, c_1, c_2) -frame if F is the frame of a (C, c_1, c_2) -hole. A hole H is an F -hole if H is a (C, c_1, c_2) -hole with frame F .

Lemma 3.3. *Let H be a (C, c_1, c_2) -hole with frame $F = (c_1, c_2, \ell'_1, \ell_1, r_1, r'_1, \ell'_2, \ell_2, r_2, r'_2)$. Assume that $v \in V(G) \setminus V(H)$ has a neighbor both in $H_L^* \setminus \{\ell_1, \ell_2\}$ and in $H_R^* \setminus \{r_1, r_2\}$. Then, v is (c_1, c_2) -heavy with respect to H .*

Proof. Suppose that v is not (c_1, c_2) -heavy with respect to H . Then, c_1 and c_2 are in an extended neighborhood \overline{Q} of v . Let $\overline{Q} = Q \cup (N_H(v) \cap \{x', y'\})$, where $Q = x \dots y$ is a v -sector, and x' and y' are the neighbors of x and y in $H \setminus Q^*$, respectively. Then, c_1 and c_2 are either in $V(Q)$ or have a neighbor in $V(Q)$. Since v has a neighbor in $H_L^* \setminus \{\ell_1, \ell_2\}$, it follows that $H_L^* \setminus V(Q) \neq \emptyset$. Similarly, $H_R^* \setminus V(Q) \neq \emptyset$.

Suppose first that c_1 is not adjacent to v . Let S be the v -sector of H that contains c_1 . Since $c_1 v \notin E(G)$ and $c_1 \in \overline{Q}$, it follows that $S = Q$. Since $c_2 \in \overline{Q}$, either v has no neighbor in $H_L^* \setminus \{\ell_1, \ell_2\}$ or v has no neighbor in $H_R^* \setminus \{r_1, r_2\}$, a contradiction. Thus, $c_1 v \in E(G)$, and similarly $c_2 v \in E(G)$. But then, since $c_1, c_2 \in \overline{Q}$, either $H_L^* \setminus V(Q) = \emptyset$ or $H_R^* \setminus V(Q) = \emptyset$, a contradiction. \square

The potential of a (C, c_1, c_2) -hole H is the total number of (c_1, c_2) -heavy vertices with respect to H . The following lemma shows that the potential of a (C, c_1, c_2) -hole only depends on its frame.

Lemma 3.4. *Let H_1 and H_2 be (C, c_1, c_2) -holes with the same frame, given by $F(H_1) = F(H_2) = (c_1, c_2, \ell'_1, \ell_1, r_1, r'_1, \ell'_2, \ell_2, r_2, r'_2)$. Then, $v \in V(G)$ is (c_1, c_2) -heavy with respect to H_1 if and only if v is (c_1, c_2) -heavy with respect to H_2 . In particular, the potential of H_1 and the potential of H_2 are equal.*

Proof. Suppose $v \in V(G)$ is (c_1, c_2) -heavy with respect to H_1 and not with respect to H_2 .

(1) *If v has no neighbor in $H_{2L}^* \setminus \{\ell_1, \ell_2\}$, then $N(v) \cap H_{1L}^* \subseteq \{\ell_1, \ell_2\}$. Similarly, if v has no neighbor in $H_{2R}^* \setminus \{r_1, r_2\}$, then $N(v) \cap H_{1R}^* \subseteq \{r_1, r_2\}$.*

By symmetry, it suffices to prove the first statement. So assume that v has no neighbor in $H_{2L}^* \setminus \{\ell_1, \ell_2\}$. We may assume that $\ell_1, \ell'_1, \ell_2, \ell'_2$ are all distinct, since otherwise the result clearly holds. In particular, H_1 and H_2 are both of length greater than six.

Since v is (c_1, c_2) -heavy with respect to H_1 , v has a neighbor in both H_{1L}^* and H_{1R}^* . Suppose v is anticomplete to $\{\ell_1, \ell_2\}$. Then, there exists a path $P = p_1 \dots p_k$ in $(H_1 \setminus \{\ell_1, \ell_2\}) \cup \{v\}$ such that $P \cap H_2 = \emptyset$, p_1 has a neighbor in H_{2L}^* , p_k has a neighbor in H_{2R} , and P^* is anticomplete to H_2 . Note that $v \in P$ and $p_1 \in L$ (i.e. $p_1 \neq v$), so P is of length at least 1. But then P and H_2 contradict Lemma 2.9. So v is not anticomplete to $\{\ell_1, \ell_2\}$. Thus, we may assume that v is adjacent to ℓ_1 . Let Q be the v -sector of H_1 that contains ℓ'_1 . Then, $\ell_1 \in V(Q)$. Since c_1 and c_2 are distant in H_1 with respect to v , it follows that v is a major vertex for H_1 . We claim that v is not a hub for H_1 . Suppose v is a hub for H_1 . Since $\ell_1 \in N(v)$ and $\ell'_1 \notin N(v)$, it follows that $c_1 \in N(v)$. But then c_1 and c_2 are not distant in H_1 with respect to v , a contradiction. This proves that v is not a hub for H_1 . Now, since v is anticomplete to ℓ'_1 - H_{2L} - ℓ'_2 , it follows from Theorem 2.13 that ℓ'_1 and

ℓ'_2 are not distant in H_1 with respect to v . Since v is not adjacent to ℓ'_2 , it follows that $\ell'_2 \in Q$, so $N(v) \cap H_{1L}^* \subseteq \{\ell_1, \ell_2\}$. This proves (1).

By Lemma 3.3, we may assume that v has no neighbor in $H_{2L}^* \setminus \{\ell_1, \ell_2\}$. By (1), it follows that $N(v) \cap H_{1L}^* \subseteq \{\ell_1, \ell_2\}$.

(2) v has a neighbor in $H_{2R}^* \setminus \{r_1, r_2\}$.

Assume that v has no neighbor in $H_{2R}^* \setminus \{r_1, r_2\}$. Then, by (1), $N(v) \cap H_{1R}^* \subseteq \{r_1, r_2\}$. But now, $N(v) \cap H_1 = N(v) \cap H_2$, and so c_1 and c_2 are distant in H_2 with respect to v , a contradiction. This proves (2).

Since c_1 and c_2 are not distant in H_2 with respect to v , there exists an extended neighborhood \overline{Q} of v in H_2 such that c_1 and c_2 are both in \overline{Q} . Let $\overline{Q} = Q \cup (N_{H_2}(v) \cap \{x', y'\})$ where $Q = x \dots y$ is a v -sector in H_2 and x' and y' are the neighbors of x and y in $H_2 \setminus Q$, respectively. Since \overline{Q} contains c_1 and c_2 , it follows that either $H_{2L} \subseteq \overline{Q}$ or $H_{2R} \subseteq \overline{Q}$. Suppose that $H_{2L} \subseteq \overline{Q}$. Since c_1 and c_2 are distant in H_1 with respect to v and $N(v) \cap H_{1L}^* \subseteq \{\ell_1, \ell_2\}$, we may assume that v is adjacent to ℓ_1 . Since c_1 is in \overline{Q} , it follows that v is also adjacent to c_1 . Because \overline{Q} is an extended neighborhood of v in H_2 that contains c_1 and c_2 , v is either non-adjacent to ℓ_2 , or v is adjacent to ℓ_2 and c_2 . But now c_1 and c_2 are not distant in H_1 with respect to v , a contradiction. Therefore, $H_{2R} \subseteq \overline{Q}$. However, by (2), v has a neighbor in $H_{2R}^* \setminus \{r_1, r_2\}$, a contradiction. \square

Let $F = (c_1, c_2, \ell'_1, \ell_1, r_1, r'_1, \ell'_2, \ell_2, r_2, r'_2)$ be a (C, c_1, c_2) -frame. We say that a vertex $v \in V(G)$ is F -heavy if there exists an F -hole H such that v is (c_1, c_2) -heavy with respect to H . Note that Lemma 3.4 implies that an F -heavy vertex v is (c_1, c_2) -heavy with respect to every hole H with frame F . A vertex v that is not F -heavy is said to be F -light. The *potential* of F is the total number of F -heavy vertices.

Let $c_1, c_2 \in C$. We denote by $\text{dist}_L(c_1, c_2)$ and $\text{dist}_R(c_1, c_2)$ the length of the shortest path from c_1 to c_2 through L and R , respectively, and we let $\text{dist}(c_1, c_2) = \min(\text{dist}_R(c_1, c_2), \text{dist}_L(c_1, c_2))$. We say that (c_1, c_2) is a *long pair* of C if $\text{dist}(c_1, c_2) \geq 4$. A (C, c_1, c_2) -frame F is *long* if (c_1, c_2) is a long pair of C . A proper separator C is *rich* if there exist $c_1, c_2 \in C$ such that (c_1, c_2) is a long pair, and *poor* otherwise.

Lemma 3.5. *Suppose F is a (C, c_1, c_2) -frame, H is an F -hole, and $c_3 \in C \setminus \{c_1, c_2\}$ is F -light. Let $P = p_k \dots p_1 c_3 q_1 \dots q_j$ be a path such that $c_3 p_1 \dots p_k$ is a path from c_3 to H_L^* through L and $c_3 q_1 \dots q_j$ is a path from c_3 to H_R^* through R (possibly $c_3 = p_k$ or $c_3 = q_j$), and assume P has length at least two. Then, up to symmetry between c_1 and c_2 , one of the following holds:*

- (i) c_1 and c_2 are anticomplete to P^* , $N_H(p_k) = \{\ell_1, c_1\}$, and $N_H(q_j) = \{c_1, r_1\}$,
- (ii) c_2 is anticomplete to P^* , c_1 has neighbors in P^* , p_k is either adjacent to c_1 or a pendant of H with neighbor ℓ_1 , and q_j is either adjacent to c_1 or a pendant of H with neighbor r_1 .

Proof. If both c_1 and c_2 have neighbors in P^* , then G contains a theta between c_1 and c_2 through H_L , H_R , and P^* , so we may assume that c_2 is anticomplete to P^* .

Suppose c_1 is also anticomplete to P^* . By Lemma 2.9, either p_k and q_j have a common neighbor in H , or their neighbors in H form an edge. Since p_k has a neighbor in H_L^* and q_j has a neighbor in H_R^* , it follows that the neighbors of p_k and q_j in H do not form an edge. Hence, we may assume that p_k and q_j are both adjacent to c_1 . If p_k and q_j both have neighbors in $H \setminus \{c_1\}$ other than ℓ_1 and r_1 , respectively, then G contains a theta between p_k and q_j through P , c_1 , and $H \setminus \{\ell_1, c_1, r_1\}$, a contradiction. Suppose $N_H(q_j) = \{c_1, r_1\}$ and p_k has a neighbor in H_L other than ℓ_1 . Let s be the neighbor of p_k in H_L closest to c_2 . Then, G contains a pyramid from p_k to $q_j c_1 r_1$ through p_k - P - q_j ,

p_k-c_1 , and $p_k-s-H \setminus \{c_1\}-r_1$, a contradiction. By definition, p_k and q_j have neighbors in H_L^* and H_R^* , respectively, and so $N_H(p_k) = \{\ell_1, c_1\}$ and $N_H(q_j) = \{c_1, r_1\}$, and outcome (i) holds.

Next, suppose c_1 has neighbors in P^* . Let r be the closest neighbor of c_1 to p_k in P^* . By Lemma 2.9 applied to the path p_k-P-r , either p_k and r have a common neighbor in H , or p_k and r are pendants of H with adjacent neighbors in H . Since $N_H(r) = \{c_1\}$, either p_k is adjacent to c_1 or $N_H(p_k) = \{\ell_1\}$. By symmetry, either q_j is adjacent to c_1 or $N_H(q_j) = \{r_1\}$, and outcome (ii) holds. \square

Let H be an F -hole and let $c_3 \in C \setminus \{c_1, c_2\}$ be F -light. A c_3 -butterfly is a path $P = p_k-\dots-p_1-c_3-q_1-\dots-q_j$, where $c_3-p_1-\dots-p_k$ is a shortest path from c_3 to H_L^* through L and $c_3-q_1-\dots-q_j$ is a shortest path from c_3 to H_R^* through R (possibly $p_k = c_3$ or $c_3 = q_j$). We call the path $c_3-p_1-\dots-p_k$ the *left wing* of P , and the path $c_3-q_1-\dots-q_j$ the *right wing* of P . We say that c_3 is a *central vertex* of P if $c_3 \neq p_k, p_{k-1}, q_j, q_{j-1}$.

The following results deal with the structure of c_3 -butterflies.

Lemma 3.6. *Suppose F is a (C, c_1, c_2) -frame, H is an F -hole, and $c_3 \in C \setminus \{c_1, c_2\}$ is F -light. Suppose further that if C is a rich separator, then F is long, and if C is a poor separator, then $\text{dist}(c_1, c_2)$ is maximum over all non-adjacent pairs in C . Let P be a c_3 -butterfly and assume c_2 is anticomplete to P^* . Suppose that c_3 is a central vertex of P . Then, (c_3, c_2) is a long pair of C . In particular, C is a rich separator.*

Proof. Assume for a contradiction that (c_3, c_2) is not a long pair of C . Then, there exists a path from c_3 to c_2 of length less than or equal to three through L or through R . First, assume that there exists a path of length two from c_3 to c_2 , say c_3-x-c_2 , and without loss of generality let $x \in L$. Because P is a butterfly and c_3 is a central vertex of P , neither c_3 nor c_3-x is the left wing of a c_3 -butterfly, so $x \notin H$ and x is anticomplete to H_L^* . It follows that $N_H(x) \subseteq \{c_1, c_2\}$. If $N_H(x) = \{c_1, c_2\}$, then G contains a theta between c_1 and c_2 through H_L, H_R , and x , so $N_H(x) = \{c_2\}$. If c_1 has neighbors in P^* , then G contains a theta between c_1 and c_2 through H_L, H_R , and $P^* \cup \{x\}$, so c_1 is anticomplete to P^* . It follows from Lemma 3.5 that $N_H(p_k) = \{\ell_1, c_1\}$. Now, G contains a pyramid from c_2 to $p_k \ell_1 c_1$ through $c_2-x-c_3-P-p_k, c_2-H_L-\ell_1$, and $c_2-H_R-c_1$, a contradiction. Therefore, there is no path of length two from c_3 to c_2 .

Next, let $c_3-x-y-c_2$ be a path of length three from c_3 to c_2 , and without loss of generality let $x, y \in L$. Since $\text{dist}(c_3, c_2) = 3$, it follows that $\text{dist}(c_1, c_2) \geq 3$. In particular, c_1 is not adjacent to y . Because c_3 is a central vertex of P , it follows that neither c_3 nor c_3-x is the left wing of a c_3 -butterfly. Therefore, $x, y \notin H$ and $N_H(x) \subseteq \{c_1\}$. Suppose x is adjacent to c_1 . Then, x and y are strictly nested with respect to H . By Lemma 2.9, x and y are pendants of H with adjacent neighbors in H , so c_1 is adjacent to c_2 , a contradiction. Hence, x is anticomplete to H .

Suppose first that c_1 is adjacent to c_3 . Consider the path $y-x-c_3$. By Lemma 2.9, either y and c_3 have a common neighbor in H , or y and c_3 are pendants of H with adjacent neighbors in H . Since $N_H(c_3) = \{c_1\}$ and y is not adjacent to c_1 , it follows that y is a pendant with $N_H(y) = \{\ell_1\}$. But $c_2 \in N_H(y)$, a contradiction. This shows that c_1 is not adjacent to c_3 . Next, suppose that c_1 has a neighbor in $\{q_1, q_2, \dots, q_j\}$. Let t be minimum such that c_1 is adjacent to q_t . Let Q be a path from y to q_t with $Q^* \subseteq \{x, c_3, q_1, \dots, q_{t-1}\}$. By Lemma 2.9, either y and q_t have a common neighbor in H , or y and q_t are pendants of H with adjacent neighbors in H . Suppose $t \neq j$, so $N_H(q_t) = \{c_1\}$. Since y is not adjacent to c_1 , it follows that y is a pendant of H and $N_H(y) = \{\ell_1\}$. But $c_2 \in N_H(y)$, a contradiction. Therefore, $t = j$. Since q_j is adjacent to c_1 and q_j has a neighbor in H_R^* , q_j is not a pendant of H . Therefore, q_j and y have a common neighbor in H . Since $y \in L$ and $q_j \in R$, the common neighbor of y and q_j is c_2 . Then, q_j is a common neighbor of c_1 and c_2 , contradicting that $\text{dist}(c_1, c_2) \geq 3$. This proves that c_1 is anticomplete to $\{c_3, q_1, \dots, q_j\}$.

Since q_j is not adjacent to c_1 , by Lemma 3.5, $N_H(q_j) = \{r_1\}$. Now, consider the path $Q = y-x-c_3-\dots-q_j$. By Lemma 2.9, either y and q_j are pendants of H with adjacent neighbors in H , or y and q_j have a common neighbor in H . Since $y \in L$ and $N_H(q_j) = \{r_1\}$, y and q_j do not have a common neighbor in H . Therefore, r_1 is adjacent to c_2 , contradicting that $\text{dist}(c_1, c_2) \geq 3$. \square

By Lemma 3.5 and Lemma 3.6, if C is a poor separator and $\text{dist}(c_1, c_2)$ is maximum over all non-adjacent pairs in C , then c_3 is not a central vertex of P . The following two lemmas prove a similar result for rich separators.

Lemma 3.7. *Suppose C is a rich separator, (c_1, c_2) is a long pair of C , F is a (C, c_1, c_2) -frame, H is an F -hole, $c_3 \in C \setminus \{c_1, c_2\}$ is F -light, and $P = p_k-\dots-c_3-\dots-q_j$ is a c_3 -butterfly with c_2 anticomplete to P^* . Assume that c_3 is a central vertex of P and let w be an F -heavy vertex. Let s be the neighbor of p_k in H_L closest to c_2 , and let t be the neighbor of q_j in H_R closest to c_2 . Let S be the path from s to t in $H \setminus \{c_1\}$, and consider the (C, c_3, c_2) -hole given by $J = P \cup S$. Then, w is a (c_3, c_2) -heavy vertex with respect to J .*

Proof. By Lemma 3.5, p_k is either adjacent to c_1 or $N_H(p_k) = \{\ell_1\}$. Since (c_1, c_2) is a long pair of C , it follows that $s \neq \ell_2$. By symmetry, $t \neq r_2$. Assume for a contradiction that w is not (c_3, c_2) -heavy with respect to J .

(1) *If p_k is adjacent to c_1 , then w has a neighbor in $p_k-s-H_L-\ell_2$.*

Because (c_1, c_2) is a long pair of C , H has length at least eight. By Lemma 3.4, w is (c_1, c_2) -heavy with respect to H . Suppose w is anticomplete to the path given by $p_k-s-H_L-\ell_2$. If w is not complete to $\{c_1, c_2\}$, then since c_1 and c_2 are distant in H with respect to w , the path $p_k-s-H_L-\ell_2$ is (H, w) -significant and w is not a hub for H , a contradiction to Theorem 2.13. So w is complete to $\{c_1, c_2\}$. Let w' be the neighbor of w in H_L^* that is closest to ℓ_2 . Note that w' exists and $w' \neq \ell_1$ since w is (c_1, c_2) -heavy with respect to H . Also, observe that there is a path Q from w to p_k through $P \cup H_R \setminus \{c_1, r_1\}$: either w has a neighbor in $P \setminus \{p_k\}$, or there is a path $Q' = q_j-t-H_R-w''-w$, where w'' is the neighbor of w in $t-H_R-c_2$ closest to t (possibly $w'' = c_2$). Let s' be the neighbor of p_k in $w'-H_L-c_2$ closest to w' ; note that s' exists since s is in $w'-H_L-c_2$ (possibly $s' = w'$). Now, G contains a theta between p_k and w through $p_k-s'-H_L-w'-w$, p_k-c_1-w , and p_k-Q-w , a contradiction. This proves (1).

(2) *w has a neighbor in $J_L^* \setminus N_J(c_3)$ and a neighbor in $J_R^* \setminus N_J(c_3)$.*

By symmetry, it suffices to show that w has a neighbor in $J_L^* \setminus N_J(c_3)$. Assume first that $s = \ell_1$. Then, $H_L^* \subseteq J_L^*$. Further, c_3 does not have a neighbor in H_L^* , otherwise $c_3 = p_k$, contradicting that c_3 is a central vertex of P . Finally, since w is (c_1, c_2) -heavy with respect to H , w has a neighbor in H_L^* , and we are done. Hence, we may assume that $s \neq \ell_1$. By Lemma 3.5, p_k is adjacent to c_1 , so by (1), w has a neighbor in $p_k-s-H_L-\ell_2$. Note that $p_k-s-H_L-\ell_2 \subseteq J_L^*$, so w has a neighbor in J_L^* . Further, c_3 does not have a neighbor in $p_k-s-H_L-\ell_2$, otherwise c_3 is adjacent to p_k , contradicting that c_3 is a central vertex of P . Therefore, w has a neighbor in $J_L^* \setminus N_J(c_3)$. This proves (2).

By Lemma 3.3, w does not have a neighbor in both $J_L^* \setminus N_J(\{c_3, c_2\})$ and $J_R^* \setminus N_J(\{c_3, c_2\})$, so by (2) we may assume that ℓ_2 is the only neighbor of w in $J_L^* \setminus N_J(c_3)$. By (2) w has a neighbor in $J_R^* \setminus N_J(c_3)$ and since c_3 and c_2 are not distant in J with respect to w , it follows that w is also adjacent to c_2 . Because c_1 and c_2 are distant in H with respect to w , it follows that $s \neq \ell_1$ and w has neighbors in the interior of c_1-H_L-s . Since $s \neq \ell_1$, by Lemma 3.5, it follows that p_k is adjacent to c_1 . Thus, w and p_k cross with respect to H . There are two cases: either p_k is a clone of ℓ_1 , or p_k is major and by Lemma 2.7 $H \cup \{w, p_k\}$ is MNC configuration (4), (5), (6), (7), or (8). Suppose the first case holds, so p_k is a clone of ℓ_1 . Since p_k is a clone of ℓ_1 , it holds that $s = \ell'_1$. By Lemma 2.8, H contains a $\{w, p_k\}$ -complete edge, so w is adjacent to ℓ_1 and c_1 (note that w is not adjacent

to $s = \ell'_1$ since ℓ_2 is the only neighbor of w in $J_L^* \setminus N_J(c_3)$. Then, $N(w) \cap H_L = \{c_1, \ell_1, \ell_2, c_2\}$, so c_1 and c_2 are not distant in H with respect to w , a contradiction. Therefore, p_k is major and $H \cup \{w, p_k\}$ is MNC configuration (4), (5), (6), (7), or (8). It follows that there is a $\{w, p_k\}$ -complete edge in c_1 - H_L - s . Let $e = v_1v_2$ be a $\{w, p_k\}$ -complete edge in c_1 - H_L - s such that v_2 is between v_1 and s in H_L . Suppose that w is not a cap with respect to J and let u be the neighbor of w in the $p_k c_2$ -subpath of $J \setminus \ell_2$ that is closest to p_k . Then G contains a theta between p_k and w through p_k - v_1 - w , p_k - s - H_L - ℓ_2 - w , and p_k - $J \setminus \{\ell_2\}$ - u - w . Hence, $N_J(u) = \{\ell_2, c_2\}$. Then, G contains a pyramid from p_k to $w\ell_2c_2$ through p_k - v_1 - w , p_k - s - H_L - ℓ_2 , and p_k - P - q_j - t - H_R - c_2 , a contradiction. \square

Lemma 3.8. *Suppose C is a rich separator, F is a (C, c_1, c_2) -frame with maximum potential over all long frames, and H is an F -hole. Let $c_3 \in C \setminus \{c_1, c_2\}$ be F -light, and let $P = p_k \dots -c_3 \dots -q_j$ be a c_3 -butterfly. Then, c_3 is not a central vertex of P .*

Proof. Suppose that c_3 is a central vertex of P . By Lemma 3.5, we may assume that c_2 is anti-complete to P^* . Furthermore, by Lemma 3.5, p_k is adjacent to c_1 or $N_H(p_k) = \{\ell_1\}$. Since (c_1, c_2) is a long pair of C , it follows that p_k is anticomplete to $\{c_2, \ell_2\}$. By symmetry, q_j is anticomplete to $\{c_2, r_2\}$. Let r be the neighbor of p_k in H_L closest to c_2 , and let s be the neighbor of q_j in H_R closest to c_2 . Let S be the path in H from r to s through c_2 . Let J be the (C, c_3, c_2) -hole given by $P \cup S$, and let F' be the (C, c_3, c_2) -frame of J . By Lemma 3.7 it follows that every F -heavy vertex is (c_2, c_3) -heavy with respect to J , and therefore is F' -heavy. Now, consider c_1 with respect to the hole J . Either c_1 is adjacent to p_k , or ℓ_1 is in J and c_1 is adjacent to ℓ_1 . Similarly, either c_1 is adjacent to q_j , or r_1 is in J and c_1 is adjacent to r_1 . Then, c_1 has a neighbor in $J_L^* \setminus N_J(\{c_3, c_2\})$ and a neighbor in $J_R^* \setminus N_J(\{c_3, c_2\})$, so by Lemma 3.3, c_1 is a (c_3, c_2) -heavy vertex of J . Finally, it follows from Lemma 3.6 that (c_3, c_2) is a long pair of C . Then, F' is a long (C, c_3, c_2) -frame with higher potential than F , a contradiction. \square

We call a (C, c_1, c_2) -frame F *optimal* if one of the following holds:

- (i) C is a rich separator and F has maximum potential over all long frames of C
- (ii) C is a poor separator and $\text{dist}(c_1, c_2)$ is maximum over all non-adjacent pairs of vertices in C .

The following theorem combines the results of Lemma 3.5, Lemma 3.6, and Lemma 3.8.

Theorem 3.9. *Let F be an optimal (C, c_1, c_2) -frame, H be an F -hole, $c_3 \in C \setminus \{c_1, c_2\}$ be F -light, and $P = p_k \dots -c_3 \dots -q_j$ be a c_3 -butterfly. Then, c_3 is not a central vertex of P .*

4 Constructing proper separators

In this section, we show how to use the structure results from previous sections to prove the main result of the paper. Our goal is to reconstruct proper separators C given only an optimal frame F of C , and two 4-tuples $M_1(C), M_2(C)$ of vertices in C . We first show that we can construct an F -hole H , and then show that we can construct three sets C_1, C_2, C_3 such that $C = C_1 \cup C_2 \cup C_3$.

We begin with a key observation about the structure of graphs in \mathcal{C} .

Lemma 4.1. *If $G \in \mathcal{C}$, then G does not contain a 3-creature.*

Proof. Assume that G contains a 3-creature with notation as in the definition of a k -creature. Suppose first that x_3 is adjacent to x_1 and x_2 . Let Q_A be a path from x_1 to x_2 through A , and let Q_B be a path from y_1 to y_2 through B . Then, x_1 - Q_A - x_2 - y_2 - Q_B - y_1 - x_1 is a hole H in G . Let $R_B = y_3 \dots -b$ be a path from y_3 to Q_B through B . Consider the path $R = x_3$ - y_3 - R_B - b . Since x_3

and b are strictly nested with respect to H , by Lemma 2.9, it follows that x_3 and b are pendants of H with adjacent neighbors in H . However, x_3 is adjacent to x_1 and x_2 , a contradiction.

We may therefore assume that x_1 is not adjacent to x_2 and y_1 is not adjacent to y_2 . Let Q_A be a path from x_1 to x_2 through A , and let Q_B be a path from y_1 to y_2 through B . Then, $x_1-Q_A-x_2-y_2-Q_B-y_1-x_1$ is a hole H in G . Let $R_A = x_3-\dots-a$ be a path from x_3 to Q_A^* through A and let $R_B = y_3-\dots-b$ be a path from y_3 to Q_B^* through B . Consider the path $R = a-R_A-x_3-y_3-R_B-b$. If $\{x_1, y_1, x_2, y_2\}$ is anticomplete to R^* , then a and b are strictly nested with respect to H , and a and b are not pendants of H with adjacent neighbors in H , contradicting Lemma 2.9. Hence, one of x_1, y_1, x_2, y_2 has a neighbor in R^* . In particular, R^* is not empty. Suppose x_1 and x_2 both have neighbors in R^* . Then, G contains a theta between x_1 and x_2 through Q_A , Q_B , and R^* , a contradiction. Therefore, not both x_1 and x_2 have neighbors in R^* . Similarly, not both y_1 and y_2 have neighbors in R^* . Since $\{x_1, y_1, x_2, y_2\}$ is not anticomplete to R^* , we may assume that x_1 has a neighbor in R^* . If y_2 also has a neighbor in R^* , then G contains a theta between x_1 and y_2 through Q_A , Q_B , and R^* , a contradiction. Therefore, y_2 is anticomplete to R^* .

Let c be the closest neighbor of x_1 to x_3 in R_A . Suppose y_1 is anticomplete to R^* and consider the path $c-R_A-x_3-y_3-R_B-b$. Then, c and b are strictly nested with respect to H . Since b has a neighbor in Q_B^* , c and b are not pendants of H with adjacent neighbors in H , contradicting Lemma 2.9. Hence, y_1 has a neighbor in R^* . Let a' be the neighbor of a in Q_A closest to x_2 , and let b' be the neighbor of b in Q_B closest to y_2 . Let H' be the hole given by $H' = x_3-R_A-a-a'-Q_A-x_2-y_2-Q_B-b'-b-R_B-y_3-x_3$. Since x_1 and y_1 are strictly nested with respect to H' , by Lemma 2.9, x_1 and y_1 are pendants of H' with adjacent neighbors in H' . Therefore, $N_{H'}(x_1) = \{x_3\}$ and $N_{H'}(y_1) = \{y_3\}$. In particular, $a'x_1, b'y_1 \notin E(G)$. Since $R^* \neq \emptyset$, without loss of generality $y_3 \neq b$. Now, G contains a theta between x_3 and y_1 through y_3 , x_1 , and $x_3-R_A-a-a'-Q_A-x_2-y_2-Q_B-y_1$, a contradiction. \square

For the rest of the section, unless otherwise specified, let C be a proper separator of $G \in \mathcal{C}$ and let $F = (c_1, c_2, \ell'_1, \ell_1, r_1, r'_1, \ell'_2, \ell_2, r_2, r'_2)$ be an optimal (C, c_1, c_2) -frame. We denote by G_F the graph $G \setminus (N(\{c_1, c_2, \ell_1, r_1, \ell_2, r_2\}) \setminus \{\ell'_1, \ell'_2, r'_1, r'_2\})$. The following two lemmas show that we can construct a set $W = W(F)$ containing every F -heavy vertex v such that $v \in V(G_F)$.

Lemma 4.2. *Let H be an F -hole and let $v \in C \cap V(G_F)$ be major for H . Then, v is F -heavy.*

Proof. Assume that v is not F -heavy, and let P be a v -butterfly. Since v is major for H , v must be an endpoint of P . Since $v \in V(G_F)$, v is anticomplete to $\{c_1, c_2, \ell_1, r_1, \ell_2, r_2\}$, and so by Lemma 3.5, P is of length at most one. Since v is F -light, by Lemma 3.3, P is of length exactly one. But then since v is anticomplete to $\{c_1, c_2, \ell_1, r_1, \ell_2, r_2\}$, P and H contradict Lemma 2.9. \square

We call $v \in C$ a (c_1, c_2) -strong vertex of G if c_1 and c_2 belong to different components of $G \setminus N[v]$. Note that given a graph G , and $v, c_1, c_2 \in V(G)$, one can determine if v is (c_1, c_2) -strong in time $\mathcal{O}(|V(G)|^2)$.

Lemma 4.3. *One can construct in polynomial time a set $W = W(F)$ that contains all F -heavy vertices v such that v is anticomplete to $\{c_1, c_2, \ell_1, \ell_2, r_1, r_2\}$ and $W \subseteq C$.*

Proof. Let H be an F -hole where the path from ℓ'_1 to ℓ'_2 through H_L is a shortest path from ℓ'_1 to ℓ'_2 in L , and the path from r'_1 to r'_2 through H_R is a shortest path from r'_1 to r'_2 through R . We may assume that H has length greater than six since otherwise W is empty. Let X_1 be the set of all (c_1, c_2) -strong vertices of G_F , and let X_2 be the set of all (c_1, c_2) -strong vertices of $G_F \setminus X_1$. Note that X_1 and X_2 can be constructed in time $\mathcal{O}(|V(G)|^3)$. If v is (c_1, c_2) -strong, then v has a neighbor in H_L^* and a neighbor in H_R^* , so $v \in C$. It follows that $X_1 \cup X_2 \subseteq C$. We claim that $W = X_1 \cup X_2$ contains all F -heavy vertices v such that v is anticomplete to $\{c_1, c_2, \ell_1, r_1, \ell_2, r_2\}$.

By Theorem 2.13, X_1 contains all F -heavy vertices v in G_F such that v is not a hub of H . Now, consider $G_F \setminus X_1$. Every F -heavy vertex in $G_F \setminus X_1$ is a hub. Suppose $v \in V(G_F \setminus X_1)$ is a major vertex for H and v is F -light. By Lemma 4.2, $v \in L$ or $v \in R$. Without loss of generality suppose $v \in L$. Since v is a major vertex for H and $v \in V(G_F \setminus X_1)$, it follows that $N(v) \cap (H_L^* \setminus \{\ell_1, \ell_2\})$ is not contained in a path of length three, so there exists a shorter path from ℓ'_1 to ℓ'_2 in L through v , a contradiction. Therefore, every major vertex for H in $G_F \setminus X_1$ is F -heavy, so every major vertex for H in $G_F \setminus X_1$ is a hub. Then, it follows from Theorem 2.13 that X_2 contains every F -heavy vertex of H in $G_F \setminus X_1$.

Finally, let v be an F -heavy vertex in G such that v is anticomplete to $\{c_1, c_2, \ell_1, r_1, \ell_2, r_2\}$. If v is not a hub, then v is an F -heavy vertex in G_F , so $v \in X_1$. If v is a hub and $v \notin X_1$, then v is an F -heavy vertex of $G_F \setminus X_1$, so $v \in X_2$. \square

Lemma 4.4. *Given an optimal frame F of C , one can construct in polynomial time an F -hole H .*

Proof. By Lemma 4.3, we can construct the set $W = W(F) \subseteq C$ of all F -heavy vertices v such that v is anticomplete to $\{c_1, c_2, \ell_1, \ell_2, r_1, r_2\}$. Let H be the graph given by the union of $V(F)$, a shortest path Q_L from ℓ'_1 to ℓ'_2 through $G_F \setminus W$, and a shortest path Q_R from r'_1 to r'_2 through $G_F \setminus W$. We claim that H is an F -hole.

If $Q_L \subseteq L$ and $Q_R \subseteq R$, then clearly H is an F -hole, so assume without loss of generality that $Q_L \not\subseteq L$. Let ℓ^* be the vertex of $Q_L \setminus L$ closest to ℓ'_1 on Q_L . Since ℓ^* has a neighbor in L and $\ell^* \notin L$, it follows that $\ell^* \in C$. Suppose ℓ^* is F -heavy. Since W contains all F -heavy vertices anticomplete to $\{c_1, c_2, \ell_1, \ell_2, r_1, r_2\}$, it follows that ℓ^* has a neighbor in $\{c_1, c_2, \ell_1, \ell_2, r_1, r_2\}$, a contradiction. Therefore, ℓ^* is F -light. Let J be an F -hole. Let P_R be a path from ℓ^* to J_R^* through R , and let P_L be a path from ℓ^* to J_L^* contained in $\ell^*Q_L\ell'_1$. Consider the path $P = P_L\ell^*P_R$ and let p_k be the end of P_L with neighbors in J_L^* . Suppose P is of length at least two. By Lemma 3.5, it follows that either p_k is adjacent to c_1 or p_k is a pendant with $N_J(p_k) = \{\ell_1\}$. Since $P_L \subseteq V(G_F \setminus W)$, p_k is not adjacent to c_1 or ℓ_1 , a contradiction. Therefore, P is of length at most one. Because ℓ^* is F -light and ℓ^* is anticomplete to $\{c_1, c_2, \ell_1, \ell_2, r_1, r_2\}$, it follows that ℓ^* does not have a neighbor in both J_L^* and J_R^* . Therefore, P has length exactly one, and P and J contradict Lemma 2.9. \square

By Lemma 4.4, we can construct an F -hole H . Let $c_3 \in C$ be F -light, and let $P = p_k \dots p_1 c_3 q_1 \dots q_j$ be a c_3 -butterfly for H . By Theorem 3.9, c_3 is not a central vertex of P . We call c_3 an L -end vertex if $c_3 = p_k$, and an L -adjacent vertex if $c_3 = p_{k-1}$. We define similarly R -end and R -adjacent. The following lemma shows that every L -adjacent vertex is in the neighborhood of two vertices in L and that every R -adjacent vertex is in the neighborhood of two vertices in R .

Lemma 4.5. *Let $X \subseteq N(H_L^*) \cap L$ be a minimal subset of $N(H_L^*) \cap L$ such that every L -adjacent vertex has a neighbor in X . Then, $|X| \leq 2$. Similarly, let $Y \subseteq N(H_R^*) \cap R$ be a minimal subset of $N(H_R^*) \cap R$ such that every R -adjacent vertex has a neighbor in Y . Then, $|Y| \leq 2$.*

Proof. Suppose $|X| > 2$ and let $x_1, x_2, x_3 \in X$. It follows from the minimality of X that for every $x_i \in X$ there exists $y_i \in C$ such that y_i is L -adjacent and $N_X(y_i) = \{x_i\}$. For $i = 1, 2, 3$, let P_i be the right wing of a y_i -butterfly. Let $A = H_L^*$ and let $B = (P_1 \setminus \{y_1\}) \cup (P_2 \setminus \{y_2\}) \cup (P_3 \setminus \{y_3\}) \cup H_R^*$. Then, A is anticomplete to B , $G[A]$ and $G[B]$ are connected, and for $i = 1, 2, 3$, x_i has a neighbor in A and is anticomplete to B , and y_i has a neighbor in B and is anticomplete to A . It follows that $A \cup B \cup \{x_1, x_2, x_3\} \cup \{y_1, y_2, y_3\}$ is a 3-creature, contradicting Lemma 4.1. \square

Let $X = \{x_1, x_2\}$ and $Y = \{y_1, y_2\}$ be as in Lemma 4.5 (so possibly $x_1 = x_2$ or $y_1 = y_2$). Let $M_1(C) = (x_1, x_2, y_1, y_2)$. Let $C_L = N(H_L^* \cup \{x_1, x_2\})$ and $C_R = N(H_R^* \cup \{y_1, y_2\})$. Note that C_L and C_R depend only on H and $M_1(C)$.

Lemma 4.6. $C_L \cap C_R \subseteq C \subseteq C_L \cup C_R$.

Proof. It follows from the definition of C_L that $C_L \subseteq L \cup C$. Similarly, $C_R \subseteq R \cup C$. Since L and R are disjoint, it follows that $C_L \cap C_R \subseteq C$. Next, suppose $c \in C$. Since $\{c_1, c_2\} \subseteq C_L \cap C_R$, we may assume that $c \in C \setminus \{c_1, c_2\}$. If c is F -heavy, then by Lemma 3.4 c is (c_1, c_2) -heavy with respect to H , and hence c has neighbors in both H_L^* and H_R^* , so $c \in C_L \cap C_R$. Therefore, we may assume that c is F -light. By Theorem 3.9, c is either L -end, L -adjacent, R -end, or R -adjacent. If c is L -end, then c has a neighbor in H_L^* . If c is L -adjacent, it follows from Lemma 4.5 that c has a neighbor in $\{x_1, x_2\}$. Therefore, if c is L -end or L -adjacent, then $c \in C_L$. By symmetry, if c is R -end or R -adjacent, $c \in C_R$. Hence, $C \subseteq C_L \cup C_R$. \square

Let $C_1 = C_L \cap C_R$. Note that for every $s \in C$, if there exists an s -butterfly P of length zero or one, then $s \in C_1$. Let $D = V(G) \setminus (H \cup C_L \cup C_R)$. The following lemmas show how to identify the vertices of $C \setminus C_1$.

Lemma 4.7. *Let $S \subseteq C_R \cap R$ be a minimal subset of $C_R \cap R$ such that for every vertex $z \in (C_L \setminus C_R) \cap C$, there exists a path from z to S through D . Then, $|S| \leq 2$. Similarly, let $T \subseteq C_L \cap L$ be a minimal subset of $C_L \cap L$ such that for every vertex $z \in (C_R \setminus C_L) \cap C$, there exists a path from z to T through D . Then, $|T| \leq 2$.*

Proof. First, note that for every vertex $z \in (C_L \setminus C_R) \cap C$ there exists a path from z to $C_R \cap R$ through D given by a subpath of the right wing of a z -butterfly. Suppose $|S| > 2$ and let $s_1, s_2, s_3 \in S$. By the minimality of S , it follows that there exist $z_1, z_2, z_3 \in (C_L \setminus C_R) \cap C$ such that there exists a path P_i from z_i to s_i through D for $i = 1, 2, 3$, and there does not exist a path from z_i to s_j through D for $1 \leq i \neq j \leq 3$. Let z'_1, z'_2, z'_3 be the neighbors of z_1, z_2, z_3 in P_1, P_2, P_3 , respectively. Let $A = H_L^* \cup \{x_1, x_2\}$ and let $B = (P_1 \setminus \{z_1, z'_1\}) \cup (P_2 \setminus \{z_2, z'_2\}) \cup (P_3 \setminus \{z_3, z'_3\}) \cup (H_R^* \cup \{y_1, y_2\})$. Then, A is anticomplete to B , $G[A]$ and $G[B]$ are connected, and for $i = 1, 2, 3$ z_i has a neighbor in A and is anticomplete to B , and z'_i has a neighbor in B and is anticomplete to A . It follows that $A \cup B \cup \{z_1, z_2, z_3\} \cup \{z'_1, z'_2, z'_3\}$ is a 3-creature, contradicting Lemma 4.1. \square

Let $S = \{r_a, r_b\}$ and $T = \{\ell_a, \ell_b\}$ be as in Lemma 4.7 (possibly $r_a = r_b$ or $\ell_a = \ell_b$). Let $M_2(C) = (\ell_a, \ell_b, r_a, r_b)$. Let C_2 be the set of all vertices $c \in C_L$ such that there exists a path P from c to $\{r_a, r_b\}$ through D . Similarly, let C_3 be the set of all vertices $c \in C_R$ such that there exists a path P from c to $\{\ell_a, \ell_b\}$ through D . Note that C_2 and C_3 depend only on H, W, C_L, C_R , and $M_2(C)$.

Lemma 4.8. $C_2 \cup C_3 \subseteq C$.

Proof. Suppose $c \in C_L$ such that there exists a path P from c to $\{r_a, r_b\}$ through D . Since $c \in C_L$, c has a neighbor in L , so some vertex of P belongs to C . Since, by Lemma 4.6, $C \subseteq C_L \cup C_R$, no vertex of $P \setminus \{c\}$ is in C . It follows that $c \in C$. Therefore, $C_2 \subseteq C$. By symmetry, $C_3 \subseteq C$. \square

Lemma 4.9. $C = C_1 \cup C_2 \cup C_3$. *In particular, C is uniquely determined by $F, M_1(C)$, and $M_2(C)$.*

Proof. By Lemmas 4.6 and 4.8, it follows that $C_1 \cup C_2 \cup C_3 \subseteq C$. Consider $c \in C$. We may assume $c \notin C_1$. Then, by Lemma 4.6, either $c \in (C_L \setminus C_R) \cap C$ or $c \in (C_R \setminus C_L) \cap C$. If $c \in (C_L \setminus C_R) \cap C$, it follows from Lemma 4.7 that there is a path P from c to $\{r_a, r_b\}$, so $c \in C_2$. Similarly, if $c \in (C_R \setminus C_L) \cap C$, then $c \in C_3$. Therefore, $C \subseteq C_1 \cup C_2 \cup C_3$. \square

Let $C(F, M_1(C), M_2(C)) = C_1 \cup C_2 \cup C_3$ be the set constructed from $F, M_1(C)$, and $M_2(C)$, as described in this section. We proved that if C is a proper separator and F is an optimal frame of C , then $C(F, M_1(C), M_2(C)) = C$. The following corollary is a summary of the results presented in Section 4 so far.

Corollary 4.10. *Given the tuples F , M_1 , and M_2 , one can construct $C(F, M_1, M_2)$ in polynomial time. Further, if F is an optimal frame of C , then $C(F, M_1(C), M_2(C)) = C$.*

Finally, we prove Theorem 1.2, which we restate here for convenience. Recall that by [3], to construct a list of all minimal separators of a graph, it suffices to prove that it has polynomially many minimal separators. We prove that \mathcal{C} has the polynomial separator property and provide in addition a polynomial-time algorithm to construct the minimal separators of graphs in \mathcal{C} , which follows naturally from the results in this section.

Theorem 4.11. *Let $G \in \mathcal{C}$. One can construct a set \mathcal{S} of size at most $|V(G)|^{18}$ in polynomial time such that \mathcal{S} is the set of all minimal separators of G .*

Proof. Let $\mathcal{S} = \{\}$. By Lemma 3.1, we add to \mathcal{S} all minimal clique separators of G . Next, we list the proper separators of G . Let $T = (c_1, c_2, \ell'_1, \ell_1, r_1, r'_1, \ell'_2, \ell_2, r_2, r'_2, x_1, x_2, y_1, y_2, \ell_a, \ell_b, r_a, r_b)$ be an 18-tuple consisting of vertices in $V(G)$. Let $F^T = (c_1, c_2, \ell'_1, \ell_1, r_1, r'_1, \ell'_2, \ell_2, r_2, r'_2)$, $M_1^T = (x_1, x_2, y_1, y_2)$, and $M_2^T = (\ell_a, \ell_b, r_a, r_b)$. For every 18-tuple T , let $C^T = C(F^T, M_1^T, M_2^T)$. By Corollary 4.10, C^T can be constructed in polynomial time. We can test in time $\mathcal{O}(|E(G)||V(G)|)$ whether C^T is a minimal separator of G . We add C^T to \mathcal{S} if and only if C^T is a minimal separator of G . Clearly, \mathcal{S} has size at most $|V(G)|^{18}$ and can be constructed in polynomial time.

It remains to show that \mathcal{S} contains every minimal separator of G . Let C be a minimal separator of G . We may assume that C is proper. Let F be an optimal frame of C and let T be the 18-tuple given by the union of F , $M_1(C)$, and $M_2(C)$, in that order. It follows from Corollary 4.10 that $C^T = C$, so $C \in \mathcal{S}$. \square

References

- [1] T. Abrishami, M. Chudnovsky, M. Pilipczuk, P. Rzażewski, P. Seymour. “Induced subgraphs of bounded treewidth and the container method”, *arXiv preprint*, arXiv:2003.05185 (2020).
- [2] I. Adler, N.K. Le, H. Müller, M. Radovanović, N. Trotignon, K. Vušković. “On rank-width of (diamond, even hole)-free graphs”, *Discrete Mathematics and Theoretical Computer Science*, 19(1) (2017).
- [3] A. Berry, J.P. Bordat, O. Cogis. “Generating all the minimal separators of a graph”, In *Lecture Notes in Computer Science*, Workshop on Graph-theoretic Concepts in Computer Science, WG (1999), Vol. 1665, Springer, Berlin, 1999.
- [4] A. Berry, R. Pogorelcnik, G. Simonet. “An introduction to clique minimal separator decomposition”, *Algorithms*, 3(2) (2010), 197–215.
- [5] V. Bouchitté, I. Todinca. “Treewidth and minimum fill-in: Grouping the minimal separators”, *SIAM J. Comput.*, 31 (1)(2001), 212–232.
- [6] V. Bouchitté, I. Todinca. “Listing all potential maximal cliques of a graph”, *Theor. Comput. Sci.*, 276 (1-2)(2002), 17–32.
- [7] M. Chudnovsky, S. Thomassé, N. Trotignon, K. Vušković. “Maximum independent sets in (pyramid, even hole)-free graphs”, *arXiv preprint*, arXiv:1912.11246 (2019).
- [8] M. Conforti, B. Gerards, K. Pashkovich. “Stable sets and graphs with no even holes”, *Mathematical Programming*, 153(1) (2015), 13–39.
- [9] F.V. Fomin, Y. Villanger. “Finding induced subgraphs via minimal triangulations”, In *27th International Symposium on Theoretical Aspects of Computer Science*, STACS (2010), 383–394. Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2010.
- [10] A. Grzesik, T. Klimošová, M. Pilipczuk, M. Pilipczuk. “Polynomial-time algorithm for maximum weight independent set on P_6 -free graphs”, In *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA (2019), 1257–1271.
- [11] D. Lokshantov, M. Vatschelle, Y. Villanger. “Independent set in P_5 -free graphs in polynomial time”, In *Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA (2014), 570–581.