



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/208104/>

Version: Accepted Version

Conference abstract:

Forbes, K. (2023) Interaction between Glucose, Extracellular Vesicles and Placenta Function in Gestational Diabetes. [Conference abstract]

<https://doi.org/10.1016/j.jri.2023.104001>

© 2023, Elsevier. This is an author produced version of a conference abstract published in Journal of Reproductive Immunology. Uploaded in accordance with the publisher's self-archiving policy. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Interaction between Glucose, Extracellular Vesicles and Placenta Function in Gestational Diabetes

Karen Forbes

Leeds Institute for Cardiovascular and Metabolic Medicine, University of Leeds, UK

Pregnancies affected by gestational diabetes (GDM) commonly result in large-for-gestational-age (LGA) infants; these infants have an increased risk of stillbirth and developing cardiometabolic complications in later life. Women with GDM who deliver LGA infants have subtle (1-1.5mM) differences in temporal maternal glucose control detectable by continuous glucose monitoring (CGM), compared to women who deliver AGA infants. It is unclear how these subtle changes contribute to LGA.

This talk will present several data sets using blood and placental samples collected from women with GDM; novel in-vitro and ex-vivo models of human placenta; and in-vivo models of pregnancy in mice, that begin to unravel the complex link between maternal glucose and LGA in pregnancies complicated by GDM. Our data provides evidence that pathological fetal growth in pregnancies complicated by GDM, may be attributed to both the direct -and indirect- (via extracellular vesicles) actions of temporal changes in maternal glucose levels on placental growth, metabolism, inflammatory and angiogenic processes.