



UNIVERSITY OF LEEDS

This is a repository copy of *Characterising Trends in the Initiation, Timing, and Completion of Recommended Summary Plan for Emergency Care and Treatment (ReSPECT) Plans: Retrospective Analysis of Routine Data from a Large UK Hospital Trust*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/209876/>

Version: Accepted Version

---

**Article:**

Anik, E., Hurlow, A., Azizoddin, D. et al. (5 more authors) (2024) Characterising Trends in the Initiation, Timing, and Completion of Recommended Summary Plan for Emergency Care and Treatment (ReSPECT) Plans: Retrospective Analysis of Routine Data from a Large UK Hospital Trust. *Resuscitation*. ISSN 0300-9572

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

1 **Title:** Characterising Trends in the Initiation, Timing, and Completion of Recommended  
2 Summary Plan for Emergency Care and Treatment (ReSPECT) Plans: Retrospective  
3 Analysis of Routine Data from a Large UK Hospital Trust  
4

5 **Authors:** Evrim Anik<sup>1,2</sup>, Adam Hurlow<sup>3</sup>, Desiree Azizoddin<sup>4</sup>, Robert West<sup>1</sup>, Felix  
6 Muehlensiepen<sup>5</sup>, Gemma Clarke<sup>1</sup>, Sarah Mitchell<sup>1</sup>, Matthew Allsop<sup>1\*</sup>  
7

8 **Affiliations:**  
9

10 1 = Leeds Institute of Health Sciences, University of Leeds, UK

11 2 = Leeds Dental Institute, University of Leeds, UK

12 3 = Leeds Teaching Hospitals NHS Trust, Leeds, UK

13 4 = Health Promotion Research Center, University of Oklahoma, USA

14 5 = Center for Health Services Research, Brandenburg Medical School Theodor Fontane,  
15 Germany  
16

17 **Email addresses:**

18 Evrim Anik: E.Anik@leeds.ac.uk

19 Adam Hurlow: a.hurlow@nhs.net

20 Desiree Azizoddin: desiree-azizoddin@ouhsc.edu

21 Robert West: R.M.West@leeds.ac.uk

22 Felix Muehlensiepen: Felix.Muehlensiepen@mhb-fontane.de

23 Gemma Clarke: G.C.Clarke@leeds.ac.uk

24 Sarah Mitchell: S.Mitchell2@leeds.ac.uk

25 Matthew Allsop: m.j.allsop@leeds.ac.uk  
26

27 Corresponding author: Matthew J Allsop. Email: [m.j.allsop@leeds.ac.uk](mailto:m.j.allsop@leeds.ac.uk). Postal address:

28 Leeds Institute of Health Sciences, Level 10, Worsley Building, Clarendon Way, Leeds, LS2

29 9LU, United Kingdom.

30 **ABSTRACT**

31

32 **Aim:** To assess patient socio-demographic and disease characteristics associated with the  
33 initiation, timing, and completion of emergency care and treatment planning in a large UK-  
34 based hospital trust.

35

36 **Methods:** Secondary retrospective analysis of data across 33 months extracted from digitally  
37 stored Recommended Summary Plan for Emergency Care and Treatment (ReSPECT) plans  
38 within the electronic health record system of an acute hospital trust in England, UK.

39

40 **Results:** Data analysed from ReSPECT plans (n=23,729), indicate an increase in the  
41 proportion of admissions having a plan created from 4.2% in January 2019 to 6.9% in  
42 September 2021 (mean = 8.1%). Forms were completed a median of 41 days before death (a  
43 median of 58 days for patients with capacity, and 21 days for patients without capacity). Do  
44 not attempt resuscitation was more likely to be recorded for patients lacking capacity, with  
45 increasing age (notably for patients aged over 74 years), being male, having ethnicity recorded  
46 as 'Asian or Asian British' or 'Black or Black British', and the absence of multiple disease  
47 groups. Having a preferred place of death recorded as 'hospital' led to a five-fold increase in  
48 the likelihood of dying in hospital.

49

50 **Conclusion:** Variation in the initiation, timing, and completion of ReSPECT plans was  
51 identified by applying an evaluation framework. Digital storage of ReSPECT plan data  
52 presents opportunities for assessing trends and completion of the ReSPECT planning process  
53 and benchmarking across sites. Further research is required to monitor and understand any  
54 inequity in the implementation of the ReSPECT process in routine care.

55

56

57 **Keywords:** acute setting, emergency care, treatment planning, advance care planning,  
58 routine data

59 **INTRODUCTION**

60 In recent years, an approach to inform emergency care and treatments of adults and children  
61 across health and care settings in the UK has been developed, known as the Recommended  
62 Summary Plan for Emergency Care and Treatment (ReSPECT) process. The ReSPECT  
63 process was developed by the Resuscitation Council UK and is designed to facilitate proactive  
64 discussions about a person's options and preferences for care and treatment in the event of  
65 serious illness. This would inform decisions should they lack the mental capacity to engage in  
66 decision-making. This includes decisions relating to cardiopulmonary resuscitation (CPR).  
67 Historically, UK localities relied on standalone do-not-attempt resuscitation orders, but an  
68 increasing number of organisations are adopting ReSPECT or similar treatment escalation  
69 planning processes<sup>1</sup>. The ReSPECT process is underpinned by conversations between the  
70 patient and clinicians to inform realistic treatment preferences, leading to clear  
71 recommendations recorded by clinicians<sup>2</sup>. The approach is intended to guide care and  
72 treatment in the event of serious clinical deterioration across all settings, including palliative  
73 and end-of-life care<sup>3</sup>. The ReSPECT process however exists in a noisy landscape of  
74 approaches for discussing and documenting advance and future care planning<sup>4</sup>. There is  
75 limited evidence reported on its use and recognised variation in the way that the ReSPECT  
76 process influences practice, including inconsistent conversations surrounding the process<sup>5</sup>.

77  
78 ReSPECT sits within the envelope of broader Advance Care Planning (ACP), focusing on  
79 context-specific clinical recommendations, such as emergency care, treatment, and  
80 cardiopulmonary resuscitation. ACP is '...a process that supports adults at any age or stage  
81 of health in understanding and sharing their personal values, life goals and preferences  
82 regarding future medical care'<sup>6</sup>. ACP involves determining what people want to happen in the  
83 delivery of their future care (advance status of wishes and preferences), what people do not  
84 want to happen (including advance decisions to refuse treatment), and who will speak on their  
85 behalf if needed (proxy or lasting power of attorney). Internationally, different approaches to  
86 support documentation and sharing of advance care plans have been explored, including  
87 Portable Medical Orders, formerly Physician Orders for Life-Sustaining Treatment (POLST),  
88 in the United States<sup>7</sup>. Care is largely concordant with preferences documented on POLST  
89 forms<sup>8</sup>, but there remains a limited evidence base underpinning their use, including how they  
90 influence care delivery and any potential unintended consequences<sup>9</sup>.

91  
92 The rollout and implementation of ReSPECT in the UK has occurred in more than one-quarter  
93 of acute hospitals<sup>10</sup>, with increasing coverage across geographical regions<sup>11</sup>. ReSPECT seeks  
94 to improve the identification of people, including children and young people, at risk of life-  
95 threatening clinical deterioration to offer relevant ACP<sup>12</sup>. It is therefore important to monitor

96 and appraise its implementation. This need was emphasised during the first wave of the  
97 COVID-19 pandemic when in the context of rapidly developed new guidance regarding  
98 escalation, evidence indicated increased documentation of recommendations to not  
99 administer CPR for some ethnic minority groups with approximately one in five patients not  
100 being consulted regarding CPR recommendations in hospital settings<sup>13</sup>. The primary aim of  
101 this study was to assess patient socio-demographic and disease characteristics associated  
102 with the initiation, timing, and completion of emergency care and treatment planning in a large  
103 UK-based hospital trust. A secondary aim was to develop a systematic framework that  
104 identifies patterns of the completion, timing, and characteristics of patients with ReSPECT  
105 plans.

106

## 107 **METHODS**

### 108 **Study design**

109 A secondary retrospective analysis using data extracted from ReSPECT plans stored within  
110 the electronic health record system of an acute hospital trust in England, UK.

111

### 112 **Study setting**

113 The study took place using data collected from Leeds Teaching Hospital Trust (LTHT), one of  
114 the largest acute hospital trusts in the UK. The Trust comprises five hospitals and provides  
115 healthcare and specialist services for the population of Leeds and the surrounding region of  
116 Yorkshire and the Humber, alongside specialist services that can be accessed nationally.  
117 Details of ReSPECT plan implementation at LTHT is available in Appendix A. Under a data-  
118 sharing agreement, deidentified data for all ReSPECT plans recorded between 1st January  
119 2019 to 30<sup>th</sup> September 2021 were extracted by a data quality officer in the informatics team  
120 at LTHT and shared in a secure data environment at the University of Leeds. Details of data  
121 extraction, preparation and cleaning can be found in Appendix A.

122

### 123 **Statistical analysis**

124 Five questions guided analysis of data: 1) How many ReSPECT plans are being created and  
125 how has the proportion of plans changed over time? 2) What are the characteristics of people  
126 who receive a ReSPECT plan? 3) Which sociodemographic characteristics are associated  
127 with the completeness of ReSPECT plans? 4) Which patient sociodemographic and clinical  
128 characteristics are associated with documented resuscitation decisions? 5) Is documentation  
129 of the hospital as a preferred place of death associated with dying in hospital? For the data  
130 analysis, both Microsoft Excel (Office Professional 2016) and R version 4.2.3 were used. Data  
131 was explored using descriptive statistics to report sociodemographic characteristics of the

132 patients (see Appendix A for the list of variables), recorded preferences for cardiopulmonary  
133 resuscitation, and information regarding who administered ReSPECT stratified by recorded  
134 capacity (questions 1 and 2). Chi-squared tests were used for categorical variables and t-tests  
135 for the continuous variable to assess whether patients with capacity and without capacity  
136 statistically differ from each other in terms of sociodemographic characteristics and recorded  
137 preferences.

138

139 The completeness of each ReSPECT plan was coded into one of four categories based on  
140 the extent of fields completed (see Appendix B for further detail): Mandatory (minimal data  
141 required to record a ReSPECT plan (i.e., nature of patient (or proxy) involvement in agreeing  
142 on the plan, and a documented cardiopulmonary resuscitation recommendation), Level 1  
143 (minimally useful additional content recorded), Level 2 (intermediate completion), and Level 3  
144 (comprehensive completion) (See Appendix C). Binary logistic regression was used to assess  
145 which variables predicted completeness of records (as completed at a Level (Level 1, Level 2  
146 or Level 3) versus mandatory level) (question 3).

147

148 Binary logistic regression was used to assess which variables predicted a recording of Do Not  
149 Attempt CPR (DNACPR) (question 4), where a subset of the data was used for this analysis.  
150 This subset of data included patients who were at least 18 years old and for whom a CPR  
151 decision was recorded (question 4). Different regression models were used to assess greatest  
152 model fit. The best fit was chosen based on lower Akaike Information Criteria (AIC)<sup>14</sup>. Some  
153 regression models included interaction terms to assess whether there was interaction between  
154 independent variables used. In regression models, cases with missing values for the variables  
155 used were deleted.

156

157 Binary logistic regression was also used to assess which variables predicted hospital death  
158 (died in LTHT- yes/no) with the data from all patients who had died and had a documented  
159 place of death (question 5). Missing values are presented as counts in Appendix B. All  
160 regression modelling adjusted for age and collinearity checks were performed to suggest that  
161 any confounding would have very little impact.

162 The reporting of the study is aligned with the STROBE checklist<sup>15</sup>. This research was  
163 undertaken as part of service evaluation work to inform a programme of work within LTHT and  
164 Leeds Palliative Care Network relating to ReSPECT implementation. As a service evaluation,  
165 and with data sharing agreements in place, the project was excluded from Health Research  
166 Authority approval.

167

## 168 **RESULTS**

169 In total, content from 33,895 ReSPECT plans was received. Of these, plans that were either  
170 the sole existing or most recent active plan for a patient were used (n=23,729). Most records  
171 contained one, single plan (n=18,640), with fewer patients having two or more saved iterations  
172 of the plan.

173

### 174 **Trends in the creation of ReSPECT plans over 33 months**

175 Trends in the number of records created each month over 33 months can be seen in Figure  
176 1a. There were consistently over 600 records created each month, with an increase in the  
177 number of ReSPECT plans documented in the early months of the COVID-19 pandemic from  
178 March to June 2020. As shown in Figure 1b, there was also an increase in the number of  
179 recorded ReSPECT plans as a proportion of all admissions (see Appendix D for monthly  
180 values). Whilst there was a surge during 2020, the proportion of plans increased from 4.2% in  
181 January 2019 to 6.9% in September 2021 ( $\bar{x} = 8.1\%$ ). Figure 1c indicates that the number of  
182 records created during weekdays is consistently higher than those created during weekends,  
183 with most plans recorded during daytime working hours. For decedents (n=16,156/23,729;  
184 68.0% of all patients), plans were recorded a median of 41 days (IQR 7, 206.5) before death.

185

### 186 **Recipients of ReSPECT plans and recorded preferences**

187 Sociodemographic information on patients and documentation of items is presented in Table  
188 1. More than half of all recorded ReSPECT plans (61.8%) were for patients aged over 74  
189 years, with 207 patients (0.9%) under the age of 20. There were more females (52.6%) than  
190 males (47.4%), with most plans created by White British patients (88.9%). Compared to all  
191 admissions during the same period (Appendix E), there was a greater proportion of patients  
192 with a ReSPECT plan that were older (61.3% with ReSPECT plan aged 75+ vs 19.3% of all  
193 admissions) and recorded as White ethnicity (89.2% vs 77.4%). When compared to all  
194 admissions that died within one year (Appendix E) there is alignment for age and ethnicity, but  
195 differences by sex for females (47.3% died vs 52.6% with ReSPECT plans) and males (52.7%  
196 died vs 47.4% with ReSPECT plans).

197

198 A total of 50.3% of patients with ReSPECT plans were living in areas of most or high levels of  
199 deprivation. The majority of patients (64.4%) had mental capacity. A small proportion (1.2%)  
200 of patients with capacity declined direct involvement in a ReSPECT discussion, accepting  
201 recommendations made by a clinician in conjunction with identified advocates. The existence  
202 of LPA for health and welfare was “unknown” for most plans (75.9%).

203

204 When compared to patients with mental capacity to agree on recommendations, ReSPECT  
205 plans for those who lacked capacity were more likely to be for people who were: over the age

206 of 74 years (72.3 versus 56.7% who had capacity), female (54.3 versus 51.8% with capacity),  
207 had non-White ethnicity recorded (12.4 versus 10% with capacity), not an inpatient at the time  
208 the plan was created (10.6 versus 7.6% with capacity), have an LPA documented (6.6 versus  
209 1.8% with capacity), and have an undecided preferred place of care (49.1 versus 36.5% with  
210 capacity) and death (57.6 versus 51.2% with capacity). Plans for those without capacity were  
211 more likely to document not for attempted CPR (97.9 versus 79.4% with capacity), with a  
212 shorter median number of days from creating a plan to death (21 versus 58 days with  
213 capacity). For those with DNACPR recorded, 22.4% (n=4,511) were alive or discharged at the  
214 time of the study (see Appendix F).

215

216 Recording of broader ACP information was limited. Preferred place of care was documented  
217 in 37.0% (8,775/23,729) of plans, and preferred place of death in 33.4% (7,917/23,729). For  
218 both categories, 'undecided' was most recorded (e.g., 53.5% (4,233/7,917) for preferred place  
219 of death). DNACPR in the event of cardiac arrest was recorded in 85.9% (20,153/23,729) of  
220 plans. There was variation across each month with >35% of forms recommending CPR during  
221 the first three months of the COVID-19 pandemic (see Appendix G). Free-text treatment  
222 escalation recommendations commonly related to interventions within the hospital setting (i.e.,  
223 58.5% of all plans). Treatment escalation information was documented less for treatment  
224 outside the hospital setting (4.7%) and for hospital readmission (16.4%).

225

### 226 **Completeness of recorded ReSPECT plans**

227 Figure 2 reflects data on levels of completion across all iterations of ReSPECT plans. A total  
228 of 14,138 (41.2%) contained only mandatory information. Most others were at Level 1  
229 (minimally useful) (n=16,220; 47.1%). Fewer than 10% of plans included Level 2  
230 (intermediate) (n=2,878; 8.2%) or Level 3 (comprehensive) (n=1,161; 3.4%).

231

### 232 **Factors influencing completeness**

233 Demographic information stratified according to the completeness of ReSPECT plans can be  
234 found in Appendix H. Logistic regression (see Table 2) was used to explore the relationship  
235 between sociodemographic variables and completeness (complete at any level versus  
236 mandatory items only). Table 2 shows that there was a significant increase in the likelihood of  
237 having only mandatory level items completed in plans for patients in age categories of either  
238 18 to 49 or over 74 years compared to being 50-74 years, being female, having ethnicity  
239 recorded "Other" compared to White, and living in less deprived areas.

240

### 241 **Factors influencing cardiopulmonary resuscitation recommendations**



242 Logistic regression was used to explore factors influencing the likelihood of a DNACPR  
243 recommendation being recorded. There was a significant increase in the likelihood of having  
244 a DNACPR recommendation for patients lacking capacity, older patients (notably for patients  
245 aged over 74 years), being male, and recorded ethnicity 'Asian or Asian British' or 'Black or  
246 Black British'. For most records with a specific disease recorded, there was an increased  
247 likelihood of recording not for CPR. The exception was COVID-19, where having COVID-19  
248 documented was associated with an increased likelihood of a recommendation for CPR (OR  
249 = 2.14, 95 CI [1.77, 2.59]).

250

### 251 **Relationship between documented place of death and likelihood of hospital death**

252 Most patients who had a ReSPECT plan died (n=16,154;68.4%), with the majority dying in  
253 hospital (n=8,467;52.4%). For patients who died with a ReSPECT plan, 62.2% (n=10,045)  
254 were missing a preferred place of death. For patients who died with a ReSPECT plan and  
255 without a preferred place of death recorded, 65.5% (n=5,546) died in hospital. A sub-analysis  
256 was conducted on data for all patients who had died and had a documented place of death  
257 (n=6,109) (see Table 4). Having a preferred place of death recorded as hospital leads to a  
258 five-fold increase in the likelihood of dying in hospital. Other factors that may increase the  
259 likelihood of a hospital death are for patients where their plan records that a carer does not  
260 have insight into the patient's illness or the carer's insight is unknown.

261

## 262 **DISCUSSION**

263 Use of ReSPECT plans increased since initial implementation with most patients having a  
264 single iteration of a plan, suggesting that either plans were not reviewed after creation or were  
265 reviewed but not amended. When compared to all admissions, patients with ReSPECT plans  
266 were older with a larger proportion of White ethnicity recorded, although similar to admissions  
267 that died within one year. Among patients who had died, plans were created a median of 41  
268 days before death ( $\bar{x}$ =58 days for people with capacity,  $\bar{x}$ =21 days for those without capacity).  
269 The majority (>80%) of plans recommended DNACPR. Lacking capacity, increasing age  
270 (notably aged over 74 years), being male, and having any ethnicity other than 'White'  
271 recorded, significantly increased the likelihood of a DNACPR recommendation. Plans with  
272 COVID-19 recorded as a disease were twice as likely to recommend CPR. Lower  
273 completeness of plans was more likely for patients aged 18 to 50, or over 74 years, having  
274 'Other' as recorded ethnicity, being female, and from least deprived areas. Only a third of all  
275 plans had ACP information, including preferences around place of care and death. Plans for  
276 patients aged 18 to 50, or over 74 years, with 'other' as recorded ethnicity, who are female,  
277 and from least deprived areas were less comprehensively completed. LPA for health and  
278 welfare were "unknown" for the majority of plans.

279

280 Around 5 – 10% of all admissions had a ReSPECT plan created, with more than half recorded  
281 for patients in the age group 70 to 90 years old, aligned with findings on the increasing  
282 relevance of advance care planning for hospital-based acute medical care with increasing  
283 age<sup>16</sup>. During the first wave of the COVID-19 pandemic, there was a spike in the proportion of  
284 admissions with a completed ReSPECT plan. Having COVID-19 recorded as a diagnosis  
285 increased the likelihood of having 'For attempted CPR' recorded as opposed to DNACPR.  
286 Documentation of DNACPR preferences during the pandemic has subsequently been  
287 scrutinised, with mixed findings relating to the influence of socio-demographic factors such as  
288 ethnicity and deprivation. In the US, for example, white race and higher deprivation were  
289 associated with having a do not resuscitate order<sup>17</sup>. In the UK, Asian ethnicity was associated  
290 with a lower use of early DNACPR<sup>18</sup>. This study found that patients with 'other' ethnicity were  
291 less likely to have comprehensively completed ReSPECT plans. 'Asian or Asian British' or  
292 'Black or Black British' ethnicity had an increased likelihood of having not for CPR  
293 documented. The latter finding is in contrast to international evidence indicating that people  
294 from minoritised ethnic groups have a preference for life-sustaining and aggressive treatments  
295 at the end of life, and are less likely to have formally documented advance care plans<sup>19, 20</sup>.  
296 Clinicians may believe there are a greater number of barriers to conducting discussions with  
297 patients from minoritised ethnic groups about CPR preferences (e.g., a patient's previous  
298 experiences of racism and discrimination, language and communication, uncertainty over  
299 family decision-making processes, and religious imperatives to preserve life at all costs)<sup>21</sup>.  
300 This study additionally identified that patients with the highest levels of deprivation formed the  
301 largest proportion of people with a ReSPECT form and an increased likelihood of more  
302 complete forms. Greater representation of high deprivation across those with a ReSPECT  
303 plan however may reflect known associations between greater deprivation and the likelihood  
304 of dying in hospital<sup>22</sup>. Further exploration of trends relating to ReSPECT plans and their  
305 association with socio-demographic factors is needed to determine gaps in the aims of the  
306 ReSPECT approach and its implementation in practice<sup>5</sup>.

307

308 Multiple benefits were derived from digital-stored ReSPECT plans in this research project,  
309 incorporated into the hospital's electronic health record system<sup>23</sup>. This study was able to apply  
310 a replicable and scalable framework whilst addressing a need for better utilisation of routine  
311 data to understand care delivery<sup>24</sup>. Previous research has largely relied on qualitative, survey,  
312 case note and mixed-method approaches<sup>10, 25-32</sup>. Digital ReSPECT data enables comparisons  
313 across other hospital and community settings to be developed to enable benchmarking (e.g.,  
314 the proportion of patients with ReSPECT forms, equitable creation of ReSPECT forms  
315 irrespective of socio-demographic and disease characteristics) which in turn may stimulate

316 quality improvement<sup>33</sup>. It also provides opportunities to explore low-cost and readily scalable  
317 approaches to promote conversations about future care and to inform targeted interventions  
318 for groups who have lower completion rates (e.g., 'other' ethnicity, female, lower age) such as  
319 conversation prompts and suggested topics and phrases for conversations<sup>34</sup>.

320

321 This study had limitations, with some missing data, and being limited to one site, albeit one of  
322 the largest acute trusts in England. In this study, the majority of plans were created within  
323 weekday working hours compared to out-of-hours. This may be more conducive to a planned  
324 approach to ReSPECT conversations supported by a team that is familiar with the patient,  
325 senior clinicians, and those people important to the patient. Nonetheless, it has been  
326 questioned whether acute hospital admissions provide an acceptable setting for ACP  
327 discussions. Evidence is mixed, with patients finding ACP discussions difficult during an acute  
328 illness, while others find them more relevant during this time <sup>35</sup>. This has led to increasing  
329 interest in the implementation of ReSPECT in community settings, enabling the facilitation of  
330 conversations during admission or shortly after discharge when a patient may be more  
331 clinically stable and less likely to have an acute compromise of mental capacity. Future  
332 research is required to reflect activity in the community setting and to develop a more  
333 comprehensive understanding of the uptake and completion of plans created across different  
334 settings.

335

## 336 **CONCLUSIONS**

337 Multiple sociodemographic and disease characteristics are associated with the initiation,  
338 timing, and completion of ReSPECT plans. These reflect variations in how the approach is  
339 being used in routine practice. At an early stage of implementation, the framework used within  
340 the study provides a means of determining engagement with and completion of the ReSPECT  
341 planning process across sites, monitoring any disparities in their use, and potentially enabling  
342 benchmarking of practice. This may help to inform how the ReSPECT process is implemented  
343 to support timely access to information to guide the care and treatment of people in the event  
344 of serious clinical deterioration across all settings.

345 **Conflicts of Interest**

346 The authors declare no conflict of interest.

347

348 **Funding**

349 This project was supported by the Leeds Palliative Care Network as part of a service  
350 evaluation project to determine how the ReSPECT process had been implemented within the  
351 hospital trust in Leeds. The funder had no role in the study design, collection, analysis and  
352 interpretation of data, or in the writing of the report or decision to submit the article for  
353 publication.

354

355 **Acknowledgements**

356 The authors would like to thank Martin Portman, Senior Information Analyst, and Ele Chan,  
357 Senior Project Manager, both based at Leeds Teaching Hospitals Trust for their support in  
358 extracting and formatting the data used in this study.

359

360 **Data availability statement**

361 Pseudonymized data are stored on a secure server at the University of Leeds. Due to the  
362 sensitive information contained, restrictions to data have been mandated through the data  
363 sharing agreement agreed between Leeds Teaching Hospitals Trust and the University of  
364 Leeds. Requests for de-identified metadata can be made to the corresponding author.

365 **Legends to figures**

366

367 **Figure 1a:** Trends in the number of new ReSPECT plans, update of records and the total  
368 number of records by each month over 33 month period

369 **Figure 1b:** Trends in the percentage of people with a ReSPECT plan within the total number  
370 of people admitted to the hospital trust over 33 months

371 **Figure 1c:** Trends in the number of ReSPECT plans created during normal hours across  
372 weekdays and weekends, alongside out-of-hours on weekdays and weekends over 33 months

373

374 **Figure 2:** Completeness of ReSPECT plan content across levels of Mandatory (minimal data  
375 required to record a ReSPECT plan (i.e., nature of patient involvement in content recorded in  
376 the plan, and a documented cardiopulmonary resuscitation recommendation), Level 1  
377 (minimally useful additional content recorded), Level 2 (intermediate completion), and Level 3  
378 (comprehensive completion).

<b>Data category</b>	<b>Level</b>	<b>Overall</b>	<b>Had capacity</b>	<b>Lacked capacity</b>	<b>p</b>
<i>n</i>		N = 23,729	15,278	8,274	
<i>Age (%)</i>	0-49	1933 ( 8.1)	1391 ( 9.1)	370 ( 4.5)	<0.001
	50-74	7143 (30.1)	5221 (34.2)	1922 (23.2)	
	75+	14653 (61.8)	8666 (56.7)	5982 (72.3)	
<i>Sex (%)</i>	Female	12470 (52.6)	7911 (51.8)	4491 (54.3)	<0.001
	Male	11259 (47.4)	7367 (48.2)	3783 (45.7)	
<i>Ethnicity (%)</i>	White	21089 (88.9)	13754 (90.0)	7248 (87.6)	<0.001
	Mixed	121 ( 0.5)	87 ( 0.6)	27 ( 0.3)	
	Asian or Asian British	815 ( 3.4)	438 ( 2.9)	326 ( 3.9)	
	Black or Black British	422 ( 1.8)	269 ( 1.8)	145 ( 1.8)	
	Other	1282 ( 5.4)	730 ( 4.8)	528 ( 6.4)	
<i>Life status (%)</i>	Alive	7575 (31.9)	5955 (39.0)	1596 (19.3)	<0.001
	Died	16154 (68.1)	9323 (61.0)	6678 (80.7)	
<i>Indices of multiple deprivation quintiles (%)</i>	1 (most deprived)	8582 (36.2)	5406 (35.4)	3089 (37.4)	0.003
	2	3341 (14.1)	2172 (14.2)	1138 (13.8)	
	3	4096 (17.3)	2688 (17.6)	1387 (16.8)	
	4	4579 (19.3)	2922 (19.1)	1631 (19.7)	
	5 (least deprived)	3116 (13.1)	2083 (13.6)	1021 (12.4)	
<i>Died in hospital (%)</i>	Yes	8467 (52.4)	4218 (45.2)	4173 (62.5)	<0.001
	No	7689 (47.6)	5106 (54.8)	2506 (37.5)	
<i>Inpatient (%)</i>	Yes	21658 (91.3)	14118 (92.4)	7397 (89.4)	<0.001
	No	2071 ( 8.7)	1160 ( 7.6)	877 (10.6)	
<i>Lasting power of attorney (LPA) for health and welfare documented (%)</i>	Yes	824 ( 3.5)	275 ( 1.8)	549 ( 6.6)	<0.001
	No	4710 (20.0)	2670 (17.5)	2040 (24.7)	
	Clinician recorded as unknown	18022 (76.5)	12333 (80.7)	5685 (68.7)	
<i>Patient has insight into their illness (%)</i>	Yes	9224 (39.3)	8752 (57.4)	472 ( 5.7)	<0.001
	No	3445 (14.7)	240 ( 1.6)	3205 (39.0)	
	Clinician recorded as unknown	10799 (46.0)	6261 (41.0)	4537 (55.2)	
<i>Carer has insight into patient illness (%)</i>	Yes	7268 (31.0)	3584 (23.5)	3684 (44.9)	<0.001
	No	401 ( 1.7)	120 ( 0.8)	281 ( 3.4)	
	Clinician recorded as unknown	15787 (67.3)	11542 (75.7)	4244 (51.7)	
<i>Preferred place of Care (%)</i>	Care-home	267 ( 3.0)	101 ( 1.7)	166 ( 5.5)	<0.001
	Home	2848 (32.5)	2161 (37.4)	687 (22.9)	
	Hospice	363 ( 4.1)	293 ( 5.1)	70 ( 2.3)	
	Hospital	1505 (17.2)	1015 (17.6)	490 (16.3)	
	Other	211 ( 2.4)	98 ( 1.7)	113 ( 3.8)	
	Undecided	3581 (40.8)	2107 (36.5)	1474 (49.1)	
<i>Preferred place of death (%)</i>	Care-home	220 ( 2.8)	74 ( 1.4)	146 ( 5.2)	<0.001

	Home	2161 (27.3)	1590 (31.0)	571 (20.5)	
	Hospice	600 ( 7.6)	507 ( 9.9)	93 ( 3.3)	
	Hospital	535 ( 6.8)	256 ( 5.0)	279 (10.0)	
	Other	168 ( 2.1)	74 ( 1.4)	94 ( 3.4)	
	Undecided	4233 (53.5)	2628 (51.2)	1605 (57.6)	
<i>CPR Recommendation (%)</i>	Yes for CPR	3237 (13.8)	3078 (20.2)	159 ( 1.9)	<0.001
	No formal decision made	83 ( 0.4)	69 ( 0.5)	14 ( 0.2)	
	No for CPR	20153 (85.9)	12110 (79.4)	8042 (97.9)	
<i>CPR Discussed with (%)</i>	Carer or family member	8038 (34.3)	876 ( 5.8)	7161 (87.3)	<0.001
	Patient	14539 (62.1)	14250 (93.7)	289 ( 3.5)	
	Urgent decision	838 ( 3.6)	83 ( 0.5)	755 ( 9.2)	
<i>Days from ReSPECT plan creation to death (median [IQR])</i>		41.00 [7.00, 206.50]	58.00[14.00,239.00]	21.00[3.00,160.00]	<0.001
<i>Number of diseases recorded (%)</i>	0	1894 (10.4)	1207 (10.5)	680 (10.2)	<0.001
	1	8119 (44.4)	5314 (46.1)	2768 (41.5)	
	2	4534 (24.8)	2761 (23.9)	1750 (26.2)	
	3	2398 (13.1)	1446 (12.5)	943 (14.1)	
	4	951 (5.2)	579 (5.0)	369 (5.5)	
	5	305 (1.7)	179 (1.6)	123 (1.8)	
	6+	88 (0.5)	50 (0.5)	38 (0.5)	
<b>Documentation of treatment escalation plans (%)</b>					
<i>Treatment escalation relating to hospital readmission (%)</i>	Blank	19828 (83.6)	12583 (82.4)	7166 (86.6)	<0.001
	Free-text data present	3900 (16.4)	2694 (17.6)	1108 (13.4)	
<i>Treatment escalation relating care within hospital (%)</i>	Blank	9853 (41.5)	6265 (41.0)	3554 (43.0)	0.004
	Free-text data present	13876 (58.5)	9013 (59.0)	4720 (57.0)	
<i>Treatment escalation relating 'Other' (%)</i>	Blank	22614 (95.3)	14588 (95.5)	7904 (95.5)	0.901
	Free-text data present	1115 ( 4.7)	690 ( 4.5)	370 ( 4.5)	
<i>Treatment escalation relating to care outside hospital (%)</i>	Blank	22608 (95.3)	14519 (95.0)	7966 (96.3)	<0.001
	Free-text data present	1121 ( 4.7)	759 ( 5.0)	308 ( 3.7)	
<i>Treatment escalation relating potentially reversible conditions (%)</i>	Blank	22103 (93.1)	14181 (92.8)	7802 (94.3)	<0.001
	Free-text data present	1626 ( 6.9)	1097 ( 7.2)	472 ( 5.7)	
<i>Treatment escalation relating intensive care unit (%)</i>	Blank	14075 (59.3)	8923 (58.4)	5053 (61.1)	<0.001
	Free-text data present	9654 (40.7)	6355 (41.6)	3221 (38.9)	
<i>Completion levels (N = 23,729) (%)</i>	Mandatory fields only	17908 (75.5)	11434 (74.8)	6297 (76.1)	<0.001
	Level 1 (Minimally useful)	3592 (15.1)	2269 (14.9)	1323 (16.0)	
	Level 2 (Intermediate)	1425 (6.0)	1000 (6.5)	425 (5.1)	
	Level 3 (Comprehensive)	804 (3.4)	575 (3.8)	229 (2.8)	

**Table 1:** Sociodemographic information of patients and documentation of items relating to the ReSPECT stratified by recorded mental capacity

<i>Dependent: Completeness</i>		<b>At Levels 1, 2 or 3</b>	<b>Mandatory</b>	<b>OR (univariable)</b>	<b>OR (multivariable)</b>
<i>Age Category</i>	50-74	4402 (66.3)	2237 (33.7)		
	18-49	976 (58.3)	698 (41.7)	1.41 (1.26 - 1.57, p < 0.001)	1.39 (1.24 - 1.55, p < 0.001)
	75+	7849 (59.6)	5315 (40.4)	1.33 (1.25 - 1.42, p < 0.001)	1.31 (1.23 - 1.39, p < 0.001)
<i>Sex</i>	Male	6529 (63.8)	3709 (36.2)		
	Female	6698 (59.6)	4541 (40.4)	1.19 (1.13 - 1.26, p < 0.001)	1.17 (1.10 - 1.23, p < 0.001)
<i>Ethnicity</i>	White	11846 (61.9)	7304 (38.1)	-	-
	Mixed	63 (58.3)	45 (41.7)	1.16 (0.79-1.70, p=0.452)	1.16 (0.79-1.71, p=0.445)
	Asian or Asian British	423 (60.3)	279 (39.7)	1.07 (0.92-1.25, p=0.391)	1.10 (0.94-1.29, p=0.217)
	Black or Black British	237 (63.5)	136 (36.5)	0.93 (0.75-1.15, p=0.508)	0.99 (0.79-1.22, p=0.901)
	Other	658 (57.5)	486 (42.5)	1.20 (1.06-1.35, p=0.003)	1.24 (1.10-1.40, p=0.001)
<i>IMD</i>	1 (Most deprived)	4857 (63.0)	2857 (37.0)	-	-
	2	1832 (61.2)	1163 (38.8)	1.08 (0.99-1.18, p=0.085)	1.08 (0.99-1.17, p=0.101)
	3	2305 (61.7)	1433 (38.3)	1.06 (0.98-1.15, p=0.178)	1.04 (0.96-1.13, p=0.314)
	4	2533 (60.6)	1647 (39.4)	1.11 (1.02-1.19, p=0.011)	1.08 (1.00-1.17, p=0.043)
	5 (Least deprived)	1693 (59.7)	1144 (40.3)	1.15 (1.05-1.25, p=0.002)	1.13 (1.04-1.24, p=0.005)

**Table 2: Logistic regression exploring factors influencing completeness only at a mandatory level.** Number in data frame = 21477, Number in model = 21464, Missing = 13, AIC = 28471.6, C-statistic = 0.55, H&L = Chi-sq (8) 12.22 (p=0.141)



<b>Dependent: CPR Recommendation</b>	<b>Levels</b>	<b>For attempted CPR</b>	<b>Do Not Attempt CPR (DNACPR)</b>	<b>OR (univariable)</b>	<b>OR (multivariable)</b>
<i>Capacity</i>	Had capacity	2880 (20.6)	11124 (79.4)	-	-
	Lacked capacity	142 (1.9)	7184 (98.1)	13.10 (11.08-15.61, p<0.001)	12.99 (10.47-16.29, p<0.001)
<i>Age</i>	18-49	1006 (60.9)	645 (39.1)	-	-
	50-74	1557 (23.6)	5037 (76.4)	5.05 (4.50-5.66, p<0.001)	4.65 (3.95-5.49, p<0.001)
	75+	459 (3.5)	12626 (96.5)	42.90 (37.48-49.19, p<0.001)	33.10 (27.35-40.15, p<0.001)
<i>Sex</i>	Female	1306 (11.7)	9856 (88.3)	-	-
	Male	1716 (16.9)	8452 (83.1)	0.65 (0.60-0.71, p<0.001)	0.82 (0.73-0.92, p=0.001)
<i>Ethnicity</i>	White	2385 (12.5)	16643 (87.5)	-	-
	Mixed	42 (39.6)	64 (60.4)	0.22 (0.15-0.33, p<0.001)	0.57 (0.30-1.07, p=0.076)
	Asian or Asian British	222 (31.9)	473 (68.1)	0.31 (0.26-0.36, p<0.001)	0.47 (0.36-0.62, p<0.001)
	Black or Black British	129 (34.8)	242 (65.2)	0.27 (0.22-0.34, p<0.001)	0.49 (0.34-0.70, p<0.001)
	Other	244 (21.6)	886 (78.4)	0.52 (0.45-0.60, p<0.001)	0.87 (0.68-1.11, p=0.245)
<i>Dementia</i>	Yes	121 (5.2)	2206 (94.8)	-	-
	No	1594 (11.3)	12566 (88.7)	0.43 (0.36-0.52, p<0.001)	0.64 (0.51-0.80, p<0.001)
<i>Cancer</i>	Yes	242 (6.1)	3734 (93.9)	-	-
	No	1473 (11.8)	11038 (88.2)	0.49 (0.42-0.56, p<0.001)	0.24 (0.20-0.28, p<0.001)
<i>Haematological Disease</i>	Yes	60 (10.3)	523 (89.7)	-	-
	No	1655 (10.4)	14249 (89.6)	0.99 (0.75-1.29, p=0.929)	0.62 (0.44-0.84, p=0.003)
<i>COPD</i>	Yes	128 (7.1)	1679 (92.9)	-	-
	No	1587 (10.8)	13093 (89.2)	0.63 (0.52-0.76, p<0.001)	0.59 (0.48-0.73, p<0.001)
<i>Heart Failure</i>	Yes	148 (6.9)	1995 (93.1)	-	-
	No	1567 (10.9)	12777 (89.1)	0.60 (0.51-0.72, p<0.001)	0.63 (0.52-0.77, p<0.001)
<i>Frailty</i>	Yes	189 (5.2)	3468 (94.8)	-	-
	No	1526 (11.9)	11304 (88.1)	0.40 (0.34-0.47, p<0.001)	0.50 (0.42-0.60, p<0.001)
<i>Neurological Diseases</i>	Yes	26 (8.3)	286 (91.7)	-	-

	No	1689 (10.4)	14486 (89.6)	0.78 (0.51-1.14, p=0.228)	0.59 (0.37-0.93, p=0.027)
<i>Covid19</i>	Yes	288 (27.6)	756 (72.4)	-	-
	No	1427 (9.2)	14016 (90.8)	3.74 (3.23-4.33, p<0.001)	2.14 (1.77-2.59, p<0.001)
<i>Liver Disease</i>	Yes	15 (7.0)	198 (93.0)	-	-
	No	1700 (10.4)	14574 (89.6)	0.65 (0.37-1.06, p=0.109)	0.34 (0.18-0.60, p<0.001)

**Table 3 Logistic regression exploring factors influencing recording not for CPR.** Number in data frame = 23462, Number in model = 16487, Missing = 6975, AIC = 7484.3, C-statistic = 0.876, H&L = Chi-sq(8) 39.88 (p<0.001)

<b>Dependent: Died in LTHT</b>	<b>Levels</b>	<b>Died in hospital</b>		<b>OR (univariable)</b>	<b>OR (multivariable)</b>
		<b>No</b>	<b>Yes</b>		
<i>Place</i>	Other	3121 (55.0)	2556 (45.0)	-	-
	hospital	69 (16.0)	363 (84.0)	6.42 (4.97-8.42, p<0.001)	5.18 (3.95-6.88, p<0.001)
<i>Capacity</i>	Had capacity	2271 (59.4)	1555 (40.6)	-	-
	Lacked capacity	919 (40.3)	1364 (59.7)	2.17 (1.95-2.41, p<0.001)	1.21 (0.96-1.52, p=0.111)
<i>Age Category</i>	0-49	134 (47.2)	150 (52.8)	-	-
	50-74	1082 (51.8)	1008 (48.2)	0.83 (0.65-1.07, p=0.147)	1.09 (0.83-1.44, p=0.519)
	75+	1974 (52.9)	1761 (47.1)	0.80 (0.63-1.01, p=0.066)	1.10 (0.84-1.44, p=0.505)
<i>Sex</i>	Female	1674 (54.9)	1377 (45.1)	-	-
	Male	1516 (49.6)	1542 (50.4)	1.24 (1.12-1.37, p<0.001)	1.21 (1.09-1.35, p=0.001)
<i>Ethnicity</i>	White	2921 (53.0)	2594 (47.0)	-	-
	Mixed	10 (50.0)	10 (50.0)	1.13 (0.46-2.75, p=0.791)	1.09 (0.40-2.95, p=0.868)
	Asian or Asian British	67 (43.5)	87 (56.5)	1.46 (1.06-2.03, p=0.021)	1.15 (0.81-1.64, p=0.424)
	Black or Black British	41 (44.1)	52 (55.9)	1.43 (0.95-2.17, p=0.091)	1.28 (0.82-2.00, p=0.276)
	Other	151 (46.2)	176 (53.8)	1.31 (1.05-1.64, p=0.017)	1.05 (0.82-1.35, p=0.699)
<i>Index of Multiple Deprivation (IMD)</i>	1-Most deprived	1072 (49.5)	1092 (50.5)	-	-
	-2	444 (51.6)	417 (48.4)	0.92 (0.79-1.08, p=0.314)	0.94 (0.79-1.11, p=0.467)
	-3	542 (53.5)	471 (46.5)	0.85 (0.73-0.99, p=0.037)	0.87 (0.74-1.02, p=0.094)
	-4	673 (54.2)	568 (45.8)	0.83 (0.72-0.95, p=0.008)	0.87 (0.75-1.02, p=0.078)
	5-Least deprived	459 (55.6)	367 (44.4)	0.78 (0.67-0.92, p=0.003)	0.77 (0.64-0.91, p=0.003)
<i>Inpatient when ReSPECT form created</i>	Yes	2914 (52.9)	2592 (47.1)	-	-
	No	276 (45.8)	327 (54.2)	1.33 (1.13-1.58, p=0.001)	1.25 (1.04-1.50, p=0.018)
<i>LPA for health and welfare recorded or unknown</i>	yes	181 (60.5)	118 (39.5)	-	-
	no	814 (46.3)	945 (53.7)	1.78 (1.39-2.29, p<0.001)	1.77 (1.34-2.33, p<0.001)
	unknown	2195 (54.2)	1856 (45.8)	1.30 (1.02-1.65, p=0.034)	1.41 (1.09-1.85, p=0.010)
<i>Recorded that patient has insight into their illness</i>	yes	1948 (59.8)	1310 (40.2)	-	-
	no	489 (39.3)	755 (60.7)	2.30 (2.01-2.63, p<0.001)	1.32 (1.07-1.63, p=0.008)
	unknown	753 (46.9)	852 (53.1)	1.68 (1.49-1.90, p<0.001)	1.30 (1.10-1.53, p=0.002)
<i>Recorded that carer has insight into their illness</i>	yes	1644 (53.6)	1421 (46.4)	-	-
	no	42 (25.8)	121 (74.2)	3.33 (2.35-4.82, p<0.001)	2.11 (1.43-3.16, p<0.001)
	unknown	1500 (52.2)	1375 (47.8)	1.06 (0.96-1.17, p=0.259)	1.34 (1.18-1.51, p<0.001)

<i>Resuscitation Recommendation</i>	Yes for CPR	38 (62.3)	23 (37.7)	-	-
	No formal decision made	9 (75.0)	3 (25.0)	0.55 (0.11-2.07, p=0.405)	0.25 (0.01-3.90, p=0.348)
	No for CPR	3143 (52.1)	2893 (47.9)	1.52 (0.91-2.59, p=0.114)	0.89 (0.50-1.61, p=0.683)
<i>Who resuscitation recommendation was discussed with</i>	Carer or family member	955 (42.1)	1312 (57.9)	-	-
	Patient	2169 (59.8)	1457 (40.2)	0.49 (0.44-0.54, p<0.001)	0.69 (0.55-0.86, p=0.001)
	Urgent decision	59 (28.6)	147 (71.4)	1.81 (1.33-2.50, p<0.001)	1.50 (1.07-2.12, p=0.021)
<i>Days to death</i>	Mean (SD)	178.4 (230.7)	77.6 (177.4)	1.00 (1.00-1.00, p<0.001)	1.00 (1.00-1.00, p<0.001)

**Table 4: Factors influencing the likelihood of hospital death.** Number in data frame = 6109, Number in model = 6074, Missing = 35, AIC = 7570, C-statistic = 0.728, H&L = Chi-sq(8) 125.42 (p<0.001). CPR = cardiopulmonary resuscitation.

## References

1. Hawkes CA, Griffin J, Eli K, et al. Implementation of ReSPECT in acute hospitals: A retrospective observational study. *Resuscitation* 2022; 178: 26-35. DOI: <https://doi.org/10.1016/j.resuscitation.2022.06.020>.
2. Hawkes CA, Fritz Z, Deas G, et al. Development of the Recommended Summary Plan for eEmergency Care and Treatment (ReSPECT). *Resuscitation* 2020; 148: 98-107. DOI: 10.1016/j.resuscitation.2020.01.003.
3. James Lind Alliance. Emergency Medicine. <https://www.jla.nihr.ac.uk/priority-setting-partnerships/emergency-medicine/>: James Lind Alliance, 2022.
4. Ramirez-Valdez EA, Leong C, Wu F, et al. Towards cataloguing and characterising advance care planning and end-of-life care resources. *BMC Palliative Care* 2022; 21: 211. DOI: 10.1186/s12904-022-01102-3.
5. Eli K, Hawkes CA, Ochieng C, et al. Why, when and how do secondary-care clinicians have emergency care and treatment planning conversations? Qualitative findings from the ReSPECT Evaluation study. *Resuscitation* 2021; 162: 343-350. DOI: <https://doi.org/10.1016/j.resuscitation.2021.01.013>.
6. Sudore RL, Lum HD, You JJ, et al. Defining Advance Care Planning for Adults: A Consensus Definition From a Multidisciplinary Delphi Panel. *Journal of pain and symptom management* 2017; 53: 821-832.e821. DOI: 10.1016/j.jpainsymman.2016.12.331.
7. Hickman SE, Sabatino CP, Moss AH, et al. The POLST (Physician Orders for Life-Sustaining Treatment) Paradigm to Improve End-of-Life Care: Potential State Legal Barriers to Implementation. *Journal of Law, Medicine & Ethics* 2008; 36: 119-140. 2021/01/01. DOI: 10.1111/j.1748-720X.2008.00242.x.
8. Tark A, Song J, Parajuli J, et al. Are We Getting What We Really Want? A Systematic Review of Concordance Between Physician Orders for Life-Sustaining Treatment (POLST) Documentation and Subsequent Care Delivered at End-of-Life. *American Journal of Hospice and Palliative Medicine* 2020; 38: 1142-1158. DOI: 10.1177/1049909120976319.
9. Vranas KC, Plinke W, Bourne D, et al. The influence of POLST on treatment intensity at the end of life: A systematic review. *Journal of the American Geriatrics Society* 2021; 69: 3661-3674. DOI: <https://doi.org/10.1111/jgs.17447>.
10. Perkins GD, Hawkes CA, Eli K, et al. Health and Social Care Delivery Research. *Recommended summary plan for emergency care and treatment: ReSPECT a mixed-methods study*. Southampton (UK): National Institute for Health and Care Research, 2022.
11. UK RC. Where in the UK has adopted the ReSPECT process?, <https://www.resus.org.uk/respect/respect-patients-and-carers/respect-adoption-uk> (2024, accessed 15th January 2024).
12. Hopper A. *Geriatric Medicine: GIRFT Programme National Specialty Report*. 2021. London, UK Getting It Right First Time.
13. Bows H and Herring J. DNACPR decisions during Covid-19: An empirical and analytical study. *Med Law Rev* 2022; 30: 60-80. DOI: 10.1093/medlaw/fwab047.
14. Stoica P and Selen Y. Model-order selection: a review of information criterion rules. *IEEE Signal Processing Magazine* 2004; 21: 36-47. DOI: 10.1109/MSP.2004.1311138.
15. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008; 61: 344-349. DOI: 10.1016/j.jclinepi.2007.11.008.
16. Knight T, Malyon A, Fritz Z, et al. Advance care planning in patients referred to hospital for acute medical care: Results of a national day of care survey. *eClinicalMedicine* 2020; 19. DOI: 10.1016/j.eclinm.2019.12.005.
17. Olds PK, Musinguzi N, Geisler BP, et al. Evaluating disparities in code status designation among patients admitted with COVID-19 at a quaternary care center early in the pandemic. *Medicine (Baltimore)* 2023; 102: e34447. DOI: 10.1097/md.00000000000034447.

- 55 18. Sutton L, Goodacre S, Thomas B, et al. Do not attempt cardiopulmonary  
56 resuscitation (DNACPR) decisions in people admitted with suspected COVID-19: Secondary  
57 analysis of the PRIEST observational cohort study. *Resuscitation* 2021; 164: 130-138. DOI:  
58 <https://doi.org/10.1016/j.resuscitation.2021.04.028>.
- 59 19. Lenko R. Differences in Informal and Formal Advance Care Planning Use by  
60 Race/Ethnicity Among U.S. Older Adults. *Journal of pain and symptom management* 2022;  
61 63: 849. DOI: <https://doi.org/10.1016/j.jpainsymman.2022.02.026>.
- 62 20. Crooks J, Trotter S, Obe RB, et al. How does ethnicity affect presence of advance  
63 care planning in care records for individuals with advanced disease? A mixed-methods  
64 systematic review. *BMC Palliative Care* 2023; 22: 43. DOI: 10.1186/s12904-023-01168-7.
- 65 21. Islam Z, Taylor L and Faull C. Thinking ahead in advanced illness: Exploring  
66 clinicians' perspectives on discussing resuscitation with patients and families from ethnic  
67 minority communities. *Future Healthc J* 2021; 8: e619-e624. DOI: 10.7861/fhj.2021-0012.
- 68 22. Macfarlane M and Carduff E. Does place of death vary by deprivation for patients  
69 known to specialist palliative care services? *BMJ supportive & palliative care* 2018; 8: 428-  
70 430. 20161201. DOI: 10.1136/bmjspcare-2016-001099.
- 71 23. Crossfield S, Johnson O and Fleming T. Large Scale Infrastructure for Health Data  
72 Analytics. In: *2016 IEEE International Conference on Healthcare Informatics (ICHI)* 4-7 Oct.  
73 2016 2016, pp.306-306.
- 74 24. Davies JM, Gao W, Sleeman KE, et al. Using routine data to improve palliative and  
75 end of life care. *BMJ Supportive & Palliative Care* 2016; 6: 257-262. DOI:  
76 10.1136/bmjspcare-2015-000994.
- 77 25. Kesten JM, Redwood S, Pullyblank A, et al. Using the recommended summary plan  
78 for emergency care and treatment (ReSPECT) in care homes: a qualitative interview study.  
79 *Age Ageing* 2022; 51. DOI: 10.1093/ageing/afac226.
- 80 26. Hartanto M and Suthantirakumar R. Comparison of clinicians' perceptions of the  
81 Recommended Summary Plan for Emergency Care and Treatment (ReSPECT) before and  
82 during the COVID-19 pandemic. *Resusc Plus* 2022; 9: 100206. 20220114. DOI:  
83 10.1016/j.resplu.2022.100206.
- 84 27. Huxley CJ, Eli K, Hawkes CA, et al. General practitioners' experiences of emergency  
85 care and treatment planning in England: a focus group study. *BMC Fam Pract* 2021; 22:  
86 128. 20210624. DOI: 10.1186/s12875-021-01486-w.
- 87 28. Eli K, Ochieng C, Hawkes C, et al. Secondary care consultant clinicians' experiences  
88 of conducting emergency care and treatment planning conversations in England: an  
89 interview-based analysis. *BMJ Open* 2020; 10: e031633. 20200120. DOI: 10.1136/bmjopen-  
90 2019-031633.
- 91 29. Eli K, Hawkes CA, Ochieng C, et al. Why, when and how do secondary-care  
92 clinicians have emergency care and treatment planning conversations? Qualitative findings  
93 from the ReSPECT Evaluation study. *Resuscitation* 2021; 162: 343-350. 20210119. DOI:  
94 10.1016/j.resuscitation.2021.01.013.
- 95 30. Hawkes CA, Griffin J, Eli K, et al. Implementation of ReSPECT in acute hospitals: A  
96 retrospective observational study. *Resuscitation* 2022; 178: 26-35. 20220630. DOI:  
97 10.1016/j.resuscitation.2022.06.020.
- 98 31. Eli K, Huxley CJ, Hawkes CA, et al. Why are some ReSPECT conversations left  
99 incomplete? A qualitative case study analysis. *Resusc Plus* 2022; 10: 100255. 20220614.  
100 DOI: 10.1016/j.resplu.2022.100255.
- 101 32. Eli K, Hawkes CA, Fritz Z, et al. Assessing the quality of ReSPECT documentation  
102 using an accountability for reasonableness framework. *Resusc Plus* 2021; 7: 100145.  
103 20210729. DOI: 10.1016/j.resplu.2021.100145.
- 104 33. Willmington C, Belardi P, Murante AM, et al. The contribution of benchmarking to  
105 quality improvement in healthcare. A systematic literature review. *BMC health services*  
106 *research* 2022; 22: 139. DOI: 10.1186/s12913-022-07467-8.
- 107 34. Curtis JR, Lee RY, Brumback LC, et al. Intervention to Promote Communication  
108 About Goals of Care for Hospitalized Patients With Serious Illness: A Randomized Clinical  
109 Trial. *Jama* 2023; 329: 2028-2037. DOI: 10.1001/jama.2023.8812.

110 35. Peck V, Valiani S, Tanuseputro P, et al. Advance care planning after hospital  
111 discharge: qualitative analysis of facilitators and barriers from patient interviews. *BMC Palliat*  
112 *Care* 2018; 17: 127. 20181205. DOI: 10.1186/s12904-018-0379-0.