



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/210765/>

Version: Published Version

Article:

Coakley, M., Villacampa, G., Sritharan, P. et al. (2024) Comparison of circulating tumor DNA assays for molecular residual disease detection in early-stage triple-negative breast cancer. *Clinical Cancer Research*, 30 (4). pp. 895-903. ISSN: 1078-0432

<https://doi.org/10.1158/1078-0432.ccr-23-2326>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Comparison of Circulating Tumor DNA Assays for Molecular Residual Disease Detection in Early-Stage Triple-Negative Breast Cancer



Maria Coakley¹, Guillermo Villacampa², Prithika Sritharan¹, Claire Swift³, Kathryn Dunne³, Lucy Kilburn², Katie Goddard², Christodoulos Pipinikas⁴, Patricia Rojas⁴, Warren Emmett⁴, Peter Hall⁵, Catherine Harper-Wynne⁶, Tamas Hickish⁷, Iain Macpherson⁸, Alicia Okines⁹, Andrew Wardley¹⁰, Duncan Wheatley¹¹, Simon Waters¹², Carlo Palmieri¹³, Matthew Winter¹⁴, Rosalind J. Cutts¹, Isaac Garcia-Murillas¹, Judith Bliss², and Nicholas C. Turner^{1,3,9}

ABSTRACT

Purpose: Detection of circulating tumor DNA (ctDNA) in patients who have completed treatment for early-stage breast cancer is associated with a high risk of relapse, yet the optimal assay for ctDNA detection is unknown.

Experimental Design: The cTRAK-TN clinical trial prospectively used tumor-informed digital PCR (dPCR) assays for ctDNA molecular residual disease (MRD) detection in early-stage triple-negative breast cancer. We compared tumor-informed dPCR assays with tumor-informed personalized multimitation sequencing assays in 141 patients from cTRAK-TN.

Results: MRD was first detected by personalized sequencing in 47.9% of patients, 0% first detected by dPCR, and 52.1% with both assays simultaneously ($P < 0.001$; Fisher exact test). The median lead time from ctDNA detection to relapse was 6.1 months with personalized sequencing and 3.9 months with dPCR ($P = 0.004$, mixed-effects Cox model). Detection of MRD at the first time point was associated with a shorter time to relapse compared with detection at subsequent time points (median lead time 4.2 vs. 7.1 months; $P = 0.02$).

Conclusions: Personalized multimitation sequencing assays have potential clinically important improvements in clinical outcome in the early detection of MRD.

Introduction

Substantial advances have been made in the adjuvant treatment of patients with early-stage breast cancer, with approximately 85% of patients cured by current treatment. Further improvements in outcomes for breast cancer, without substantial overtreatment of patients already cured by current therapy will require improved ways of identifying those patients with a residual risk of relapse. Detection of circulating tumor DNA (ctDNA) in plasma following the completion

of treatment, often referred to as molecular residual disease (MRD) detection, is associated with a high risk of future relapse. Detection of ctDNA is most frequently done with tumor-informed assays, with sequencing of the tumor tissue to identify somatic variants used to develop personalized assays that track either a few mutations with digital PCR (dPCR), or multiple mutations with error-corrected sequencing. Tumor agnostic assays that utilize detection of aberrant tumor-specific methylation, without needing tumor tissue sequencing, are also in development. With the increasing development of multiple ctDNA methods, comparison studies are urgently required to identify the characteristics that are clinically important and identify the optimal assays to implement in clinical trials and future clinical practice.

The c-TRAK-TN prospective clinical trial identified patients with MRD following treatment for early-stage moderate to high-risk triple-negative breast cancer (TNBC) and assessed the potential activity of further adjuvant therapy with pembrolizumab after MRD detection. Other clinical trials with a similar design are ongoing, such as IMvigor011 (NCT04660344), which aims to identify and treat patients with MRD following therapy for high-risk muscle-invasive bladder cancer (1). There was a higher rate of metastatic disease at the point of MRD detection with dPCR in the c-TRAK TN clinical trial than anticipated (2), emphasizing the need to assess whether ctDNA assays with better sensitivity may lengthen the lead time from MRD detection to clinical relapse and facilitate clinical trials designed to improve patient outcomes from interventions at the point of MRD detection.

Detection of MRD with ctDNA assays is challenging, as the levels of ctDNA in these patients may be very low, requiring ultrasensitive and highly specific assays (3). A broad range of ctDNA assays are currently available, and with only a limited cross-platform comparison of these technologies (4–7), it is in general unknown whether evidence from one assay can be safely applied to other assays. An American Society of Clinical Oncology position paper has highlighted the need for

¹Breast Cancer Now Research Centre, The Institute of Cancer Research, London, United Kingdom. ²Clinical Trials and Statistics Unit, The Institute of Cancer Research, London, United Kingdom. ³Ralph Lauren Centre for Breast Cancer Research, London, United Kingdom. ⁴NeoGenomics Ltd, Glenn Berge Building, Babraham Research Park, Cambridge, United Kingdom. ⁵University of Edinburgh, Edinburgh, United Kingdom. ⁶Maidstone Hospital, Maidstone and Tunbridge Wells NHS Trust, Maidstone, United Kingdom. ⁷University Hospitals Dorset NHS Foundation Trust, Bournemouth, United Kingdom. ⁸The Beatson West of Scotland Cancer Centre, Glasgow. ⁹Breast Unit, Royal Marsden Hospital, London, United Kingdom. ¹⁰Outreach Research & Innovation Group Ltd, Manchester, United Kingdom. ¹¹Royal Cornwall Hospitals NHS Trust, Truro, United Kingdom. ¹²Velindre Cancer Centre, Velindre University NHS Trust, Cardiff, United Kingdom. ¹³Clatterbridge Cancer Centre NHS Foundation Trust, Liverpool, United Kingdom. ¹⁴Weston Park Cancer Centre, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, United Kingdom.

Corresponding Author: Nicholas C. Turner, The Institute of Cancer Research and Royal Marsden, Fulham Road, London SW3 6JB, United Kingdom. E-mail: nick.turner@icr.ac.uk

Clin Cancer Res 2024;30:895–903

doi: 10.1158/1078-0432.CCR-23-2326

This open access article is distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) license.

©2023 The Authors; Published by the American Association for Cancer Research

Translational Relevance

Key to the implementation of ctDNA assays for cancer MRD detection is an understanding of which ctDNA assay characteristics are important. Through interassay comparison in a prospective trial, we show that increased analytical sensitivity—in this case via personalized multimitation sequencing—results in clinically meaningful improvements in clinical performance. These findings emphasize the importance of implementing the most sensitive ctDNA detection assays in MRD detection clinical trials.

standardization in techniques and reporting, along with more studies on cross-assay comparisons (8).

Here, we performed an initial pilot to assess the ability of the tumor-informed Residual Disease and Recurrence (RaDaR) personalized sequencing assays (NeoGenomics) to detect MRD in patients with breast cancer. Next, we compared RaDaR with prospectively conducted tumor-informed dPCR assays in the c-TRAK TN clinical trial.

Materials and Methods

Samples

ChemoNEAR sample collection study (REC ID: 11/EE/0063) was a biological sample collection study sponsored by the Royal Marsden Hospital and approved by a research ethics committee (11/EE/0063). All patients gave written informed consent. Patients were eligible who had early breast cancer and were scheduled to receive neoadjuvant chemotherapy. Blood samples were collected at baseline, after the first cycle of chemotherapy, post surgery, every 3 months for the first year, and then every 6 months up to 5 years (9, 10).

The c-TRAK-TN clinical trial was a phase II multicenter clinical trial sponsored by the Institute of Cancer Research (NCT03145961) and approved by a research ethics committee (17/SC/0090). All patients gave written informed consent. Patients were eligible with early-stage TNBC with either residual disease following neoadjuvant chemotherapy and surgery or with a tumor size > 20 mm and/or axillary lymph node involvement prior to primary surgery and adjuvant chemotherapy. Following written informed consent, a tumor tissue sample was sequenced with a targeted assay, and personalized dPCR assays designed for one to two mutations per patient. Following completion of standard therapy, patients had prospective dPCR testing every 3 months for 2 years. If patients had ctDNA detected during the first 12 months of testing (or extended to 15- or 18-month time points if patients missed samples due to the COVID-19 pandemic), they were randomized between observation and pembrolizumab. The trial is described in greater detail in the primary trial report (2). The coprimary endpoints of the clinical trial were: (i) ctDNA detection rate at 12 months and (ii) sustained ctDNA clearance rate on pembrolizumab (2).

Both studies were conducted in accordance with the Declaration of Helsinki and Good Clinical Practice.

Tissue whole-exome sequencing

DNA was extracted from formalin-fixed, paraffin-embedded (FFPE) tissue after microdissection using the QIAamp DNA Investigator Kit (Qiagen, Manchester UK, catalog No. 56504), and from buffy coat using QIAamp DNA Blood Mini Kit (Qiagen, Manchester UK, catalog No. 51104). Sequencing libraries were prepared using KAPA HyperPlus kit with IDT UDI 8nt Adaptors (Integrated DNA Technologies). Adapter-ligated DNA fragments were validated using

Agilent TapeStation (Agilent Technologies, Palo Alto, CA; RRID: SCR_019547), and quantified using High Sensitivity Qubit assay on Qubit 3.0 Fluorometer (ThermoFisher Scientific, Waltham, MA). Libraries were pooled and hybridized with SureSelectXT Human All Exon V6 Kit (Agilent, Santa Clara, CA) at 65°C for 24 hours. Sequencing libraries were multiplexed and sequenced on a NovaSeq6000 (Illumina, San Diego, CA).

dPCR assay

DNA was extracted utilizing the QIAamp Circulating Nucleic Acid Kit (Qiagen, Manchester, United Kingdom). ctDNA testing via dPCR assay was performed as previously described (2–10). In brief, targeted-panel sequencing from two FFPE samples was performed using the ABC-Bio or RMH-200 gene panels and one to two variants were selected for the dPCR assay (11, 12). The Thermo Scientific Custom TaqMan SNP Genotyping Assay design tool was employed for assay design. Assay optimization was performed with a ProFlex Thermal Cycler (Applied Biosystems), Automated Droplet Generator (Bio-Rad, Pleasanton) and Droplet Reader (Bio-Rad, Pleasanton). Two or more FAM-positive droplets were required for the sample to be called positive. If a positive result was obtained, this was independently confirmed on a second sample. Copies/mL and allele fraction were calculated as previously described (13).

RaDaR personalized sequencing analysis

As previously described (14), somatic variants from whole-exome sequencing (WES) were prioritized using proprietary algorithms to build patient-specific primer panels of up to 48 primer pairs and combined with 21 common population SNP to facilitate panel QC. Where available, an aliquot of tumor DNA from FFPE tissue or pre-capture sequencing library for progression plasma that was subjected to WES was used to validate variants.

Plasma DNA extracted from a minimum of 2-mL banked plasma was sequenced with personalized RaDaR assays. Plasma DNA samples were run alongside a buffy coat DNA control sample from the patient, which was used to identify and remove germline variants, remove variants due to clonal hematopoiesis of indeterminate potential, and as a positive amplification control. RaDaR libraries were sequenced using a Nova-Seq 6000 system (Illumina Inc., San Diego) and sequencing data analyzed in a multistep process: fastq files were demultiplexed using Illumina bcl2fastq 2.20 software. Reads were aligned using the Burrows–Wheeler Aligner (15) and processed using proprietary software to identify primer pairs and count mutant and reference bases. Variants present in the buffy coat material or absent in the tumor tissue DNA were excluded from further analysis.

Proprietary methods were used to determine if MRD was present or absent, from the primary MRD panel variants. As previously summarized, (14) a statistical model was used to assess the statistical significance of the observed mutant counts for each variant, and the information was integrated over the entire set of personalized variants to obtain evidence of tumor presence or absence at the sample level. A sample was determined to be positive for residual disease if its cumulative statistical score was above a preset threshold. The tumor fraction estimated from the model was then reported [estimated variant allele frequency (VAF)] along with the mean VAF across all variants. These values were closely correlated (Supplementary Fig. S1, Pearson correlation $R = 0.99$; $P < 0.001$).

Statistical analysis

Survival endpoints were calculated using the Kaplan–Meier method. Time to positive ctDNA detection was calculated as a survival endpoint

from the start of ctDNA surveillance to the first ctDNA-positive result, with patients being censored after discontinuation of ctDNA surveillance. The proportion of patients with positive ctDNA detection by 12 months from study entry was calculated with the Kaplan–Meier method including all patients with information in both assays ($n = 141$). A separate sensitivity analysis was performed using calendar windows from the end of treatment (date of surgery for patients with neoadjuvant chemotherapy, or date of last chemotherapy with adjuvant chemotherapy). Lead time between ctDNA detection and disease recurrence was calculated in the subset of patients with a ctDNA-positive result, from the date of the first ctDNA-positive result to the date of relapse. Patients were censored at date of withdrawal or at their latest follow-up date. Median lead time and recurrence rate at 12 months were reported. A mixed-effect Cox regression with patient ID as a random effect to adjust by repeated measurements, was fitted to compare the time to relapse between the two assays. As a complementary analysis, the nonparametric paired-sample Wilcoxon test was calculated only in patients with relapse to evaluate whether the differences between the time to relapse according to each assay was different from 0.

The Fisher exact test was used to compare the proportion of patients with MRD first detected using the RaDaR or the dPCR assay. To evaluate the diagnostic performance of RaDaR to predict relapses at 24-months, the sensitivity, specificity, and positive and negative predictive values were calculated. To compare the RaDaR VAF values, a linear mixed model using patient ID as random intercept was used to deal with the repeated measurement data. To compare the maximum RaDaR VAF values across sites of relapse, pairwise comparisons using the Wilcoxon test and adjusting for multiple comparisons with the Benjamini & Hochberg method were reported. No data imputation was performed. The median follow-up was calculated using the reverse Kaplan–Meier method. The threshold for statistical significance was defined as 0.05 (two-sided). Statistical analyses were performed with the R statistical software version 4.1.2 and GraphPad Prism version 6.0 (RRID:SCR_002798).

Data availability

Deidentified individual participant data, together with a data dictionary defining each field in the set, will be made available to other researchers on request to the corresponding author, or by contacting c-trak-tn-icrctsu@icr.ac.uk. Trial documentation including the protocol are available on request by contacting c-trak-tn-icrctsu@icr.ac.uk.

The ICR-CTSUS supports the wider dissemination of information from the research it conducts, and increased cooperation between investigators. Trial data are collected, managed, stored, shared, and archived according to ICR-CTSUS Standard Operating Procedures to ensure the enduring quality, integrity, and utility of the data. Formal requests for data sharing are considered in line with ICR-CTSUS procedures, with due regard given to funder and sponsor guidelines. Requests are via a standard proforma describing the nature of the proposed research and extent of data requirements.

Data recipients are required to enter a formal data-sharing agreement, which describes the conditions for release and requirements for data transfer, storage, archiving, publication, and intellectual property. Requests are reviewed by the Trial Management Group (TMG) in terms of scientific merit and ethical considerations including patient consent. Data sharing is undertaken if proposed projects have a sound scientific or patient benefit rationale, as agreed by the TMG and approved by the Independent Data Monitoring and Steering Committee, as required. Restrictions relating to patient confidentiality and consent will be limited by aggregating and anonymizing identifiable patient data. In addition, all indirect identifiers that may lead to

deductive disclosures will be removed in line with Cancer Research UK Data Sharing Guidelines.

Results

Pilot study to assess the ability of RaDaR to detect MRD in patients with breast cancer


We initially conducted a pilot study to assess the potential of personalized sequencing with RaDaR in early-stage breast cancer, using a cohort of 170 patients recruited into a tissue collection study, to identify 17 patients who had relapsed, and five patients had no known relapse as controls. These patients included those with TNBC, hormone receptor-positive (HR+) and HER2 overexpressed breast cancer (Supplementary Table S1). Using a WES approach, archival tissue tumor samples were sequenced to median coverage 116x (13–227x), and matched germline buffy coat sequence to median coverage of 42x (8–100x). Personalized sequencing panels with a median of 43 (9–61) variants were designed, and a median of eight plasma time points (2–12) were tested per patient, with 141 plasma time points in total (Fig. 1A).

In this pilot study, ctDNA was detected in 39.7% (56/141) of plasma time points and was not detected in 60.3% (85/141) of plasma time points. As expected, the mean VAF was significantly higher in samples where MRD was detected ($P < 0.001$) but with some overlap at lower VAF ranges, reflecting detection of MRD with an error model specific to the individual mutations tracked (Fig. 1B). From patients with a known clinical relapse, RaDaR detected MRD in 100% of patients (17/17) prior to clinical relapse, and MRD was not detected in 100% (5/5) of the patients with no known clinical relapse (Fig. 1C). The median relapse-free survival (RFS) was 16.8 months [95% confidence interval (CI), 14.7–28.9 months] for patients who had MRD detected and was not reached in patients who did not have MRD detected ($P < 0.001$; Fig. 1D). The three patients who had central nervous system (CNS)-only relapse, had a shorter time from ctDNA detection to relapse (4.1, 5.1, and 5.7 months). We concluded that the high sensitivity and lead time of RaDaR had the characteristics to take forward to formal comparison in the cTRAK-TN trial.

Cross-assay comparison in the c-TRAK TN clinical trial cohort

c-TRAK-TN enrolled 161 patients into prospective dPCR MRD surveillance (2). For 87.5% (141/161) of patients on MRD surveillance, plasma samples were available for retrospective orthogonal testing with the RaDaR personalized sequencing ctDNA assay. The orthogonal testing set was representative of the overall study population with a median follow-up of 32.5 months (95% CI, 27.5–34.8; Supplementary table S2–4). A median of two (range, 1–3) FFPE samples per patient underwent WES (226 samples in total), 28.3% (64/226) diagnostic biopsies and 71.7% (162/226) surgical specimens, with a median coverage of 155x (43–361x). Matched germline buffy coat was sequenced with a median coverage of 52x (20–127x). All patients had successful assays designed, tracking a median of 47 variants (range, 33–56) per patient, and a total of 899 plasma time points were analyzed [median of seven time points per patient (range, 1–11)] by both RaDaR and dPCR assays (Fig. 2A).

In total, 839/899 of plasma time points tested had concordant test results, giving an overall test agreement between RaDaR and dPCR assays of 93.3% (95% CI, 91.4%–94.8%; Fig. 2B). Similar rates of agreement were found between RaDaR and dPCR assays that tracked one or two variants (Supplementary table S5). From a per-patient perspective, concordant test results occurred in 92.9% (131/141) of patients, with 58.9% (83/141) of patients not having MRD detected by either assay, and 34.0% (48/141) had MRD detected by both assays,

**Figure 1.**

Assessment of personalized multimutation sequencing in a pilot study. **A**, CONSORT diagram of samples selected in the pilot study to assess the ability of RaDaR to detect MRD in early-stage breast cancer. **B**, RaDaR ctDNA mean VAF in samples with MRD detected and MRD not detected, comparison with Mann-Whitney U test, and results on the y-axis log transformed. **C**, Plasma time points and point of clinical relapse colored by ctDNA detected, from start of ctDNA surveillance for patients shown for patients with HER2⁺ breast cancer ($n = 8$), HR⁺HER2⁻ ($n = 9$), and TNBC ($n = 5$). MRD was not detected in any patient who did not have a clinical relapse reported ($n = 5$) and was detected prior to relapse in all patients who had a clinical relapse reported ($n = 17$). **D**, RFS in patients with and without MRD detected. The MRD-detected status was treated as a time-dependent covariate to avoid the immortal time bias. The log-rank test was used to estimate survival differences.

Comparison of Circulating Tumor DNA Assays




Figure 2.

Comparison of ctDNA detection between dPCR and RaDaR in cTRAK- TN. **A**, CONSORT diagram of the samples available to perform ctDNA MRD cross-assay comparison in the c-TRAK TN clinical trial. **B**, Comparison of ctDNA detection in all samples analyzed with dPCR and RaDaR. (i) Of 899 samples, 113 were concordant positive and 726 were concordant negative, leading to a test agreement of 93.3% (95% CI, 91.4%–94.8%). In total, 60 samples were discordant, where 58 samples had MRD detected by RaDaR, but not by dPCR, and two samples had MRD detected by dPCR and not by RaDaR. (ii) Of 114 patients, 48 were concordant positive and 83 were concordant negative, leading to a test agreement rate of 92.2% (95% CI, 87.0%–96.4%). Ten patients had discordant test results, with MRD detected by RaDaR only in eight cases and by dPCR only in two cases. **C**, Plasma time points and point of clinical relapse colored by ctDNA detected, from start of ctDNA surveillance for patients with no plasma time points with MRD detected (left), which includes one patient who had a clinical relapse and for patients who had any plasma time point that tested positive by either assay (right).

although potentially at different time points (Fig. 2B). Of the 48 patients who had MRD detected by both assays, MRD was detected by RaDaR at an earlier time point than dPCR in 47.9% (23/48) patients, MRD was first detected at the same time point in 52.1% (25/48) of patients, and there were no cases where MRD was first detected by dPCR (Fig. 2C; $P < 0.001$ Fisher exact test).

The rate of ctDNA detection by 12 months after the end of treatment (date of surgery for neoadjuvant chemotherapy, date of last cycle of chemotherapy for adjuvant chemotherapy) with RaDaR was 36.9% (95% CI, 28.3%–45.5%) and with dPCR assays was 29.8% (21.5%–37.1%; Fig. 3A). By 24 months, the total duration of ctDNA testing, 39.7% (56/141) of patients had MRD detected by a RaDaR assay and 35.5% (50/141) had MRD detected by a dPCR assay. First detection of ctDNA after 12 months after the end of treatment was rare with RaDaR, and more frequent with a dPCR assay (Fig. 3B). For all patients where ctDNA was detected at a plasma time point prior to clinical relapse, the median lead time from ctDNA detection to relapse was 6.1 (95% CI, 4.2–9.0) months with RaDaR and 3.9 (95% CI, 2.8–6.5) months with dPCR ($P = 0.004$, mixed-effects Cox model;

Fig. 3C and Supplementary Fig. S2). With the RaDaR assay, detection of ctDNA at the first time point was associated with shorter time to relapse, than detection at a subsequent time point (median lead time 4.2 vs. 7.1 months; $P = 0.02$; Fig. 3D).

RaDaR assay to predict relapse at 24 months

To evaluate the diagnostic performance of RaDaR personalized sequencing assays, we assess its ability to predict clinical relapses at 24 months. Patients confirmed to be free of relapse at least 24 months from the start of the study, and patients with a relapse before 24 months were included in this analysis (80.9%, 114/141). Among those 114 patients, 47 (41.2%) had a clinical relapse, and 67 (58.8%) were relapse-free at 24 months. The sensitivity of RaDaR was 95.7% (95% CI, 84.3%–99.3%), with 45 of 47 patients with relapse having a previous ctDNA-positive result. The specificity was 91.0% (95% CI, 80.9%–96.3%); 61 of 67 relapse-free patients did not have ctDNA detection during the follow-up, although assessment of specificity is limited by follow-up. The positive and negative predictive values were 88.2% and 96.8%, respectively (Supplementary Table S6).




Figure 3. Clinical validity of ctDNA detection with dPCR and RaDaR in cTRAK-TN. **A**, Time to ctDNA detection from the start of ctDNA surveillance. **B**, ctDNA detection rates in calendar/time windows from the date of completion of standard treatment. The first window was for 0 to 1.5 months, with intervals spanning 3 months thereafter (1.5–4.5, 4.5–7.5, etc.). **C**, Time from first ctDNA detection to relapse with dPCR and RaDaR assays. **D**, Time from first RaDaR ctDNA detection to relapse, comparing patients with ctDNA detected at first time point and subsequent time points.

Downloaded from <http://aacrjournals.org/clinccancerres/article-pdf/30/4/895/3414869/895.pdf> by guest on 22 March 2024


Analysis of discordant test results

Discordant test results between RaDaR and dPCR assays were found in 7.1% (10/141) of patients, where eight patients had MRD detected by RaDaR but not by dPCR, and two patients had MRD detected by a

dPCR but not by RaDaR. Relapses were reported within the follow-up period for six of the eight patients who had MRD detected by RaDaR only and for one of the patients who had MRD detected by a dPCR assay only. The median RaDaR VAF of variants with MRD detected by

Figure 4.

Analysis of factors contributing to discordance and ctDNA detection. **A**, RaDaR mean VAF split by time points, positive by dPCR only, positive by RaDaR only, or concordant results. Comparison between groups was performed using linear mixed models. Mean VAF is demonstrated on a log10 transformed axis. **B**, RaDaR mean VAF by sites of relapse, in patients with site of relapse data (n = 49), in patients with local-only, lung-only, CNS-only, other single sites of metastatic disease (including bone, liver, and pleural), and multiple sites of disease. Pairwise comparisons using the Wilcoxon test and adjusting for multiple comparisons with the Benjamini & Hochberg method were performed. For each patient, the highest mean VAF across all plasma time points was used. Mean VAF is demonstrated on a log10-transformed axis. **C**, Anatomical sites of disease at the time of relapse for patients with MRD was not detected by either assay (n = 1, local relapse), MRD detected by dPCR only (n = 1, CNS relapse), and MRD detected by RaDaR only (n = 6), known sites of disease include lung, liver, and local relapses.



Downloaded from <http://aacrjournals.org/clinccancerres/article-pdf/30/4/895/3414869/895.pdf> by guest on 22 March 2024

both assays was higher than those samples where MRD was detected by RaDaR only (0.28% vs. 0.01%; $P < 0.001$; Fig. 4A).

Sites of relapse and ctDNA detection

We investigated whether the location of relapse affected ctDNA levels and the ability of ctDNA to detect relapse. Patients with local recurrences or lung-only relapse had lower levels of ctDNA (mean VAF in RaDaR) detected prior to relapse compared with patients with multiple sites of disease ($P < 0.001$; adjusted pairwise comparisons). Patients with multiple sites of relapse had the highest ctDNA levels (median value of the maximum VAF, 0.0256%), followed by CNS only (Fig. 4B).

In patients with relapse detected only by RaDaR prior to relapse, both local and distant recurrences were reported at relapse. In the one patient in whom MRD was detected before relapse by a dPCR assay and not by RaDaR, the patient had CNS disease only at relapse. One patient had a local recurrence in the absence of ctDNA detection by dPCR or RaDaR assay before relapse (Fig. 4C).

Discussion

Multiple ctDNA assays are in clinical development for the detection of MRD, and it is vital to understand whether differences between assays have clinical consequences and whether assays give similar or different results. Here, we show that personalized multi-mutation sequencing with the RaDaR assay frequently detects ctDNA at an earlier time point than dPCR, resulting in improved lead time over clinical relapse, without discernable impact on specificity. The improved clinical impact of RaDaR likely reflected the ability, through tracking a larger number of variants, to confidently detect ctDNA at substantially lower ctDNA levels, below the limit of detection of dPCR.

We demonstrate that improved sensitivity of RaDaR also reduced the required duration of ctDNA surveillance in moderate to high-risk TNBC, with most relapses detected within the first 12 months of completing standard therapy, potentially negating the need for ctDNA surveillance beyond this point in this clinical setting. This has both the potential for cost savings, with fewer time points needing testing and allows a patient to complete ctDNA surveillance earlier, potentially achieving an earlier time to achieving an “all clear.”

The cTRAK-TN trial demonstrated that at the time of ctDNA detection with dPCR, most patients had already asymptotically relapsed when imaged. In aggressive tumor types such as high-risk TNBC, we demonstrated that RaDaR could achieve early ctDNA assay detection in patients with TNBC. In addition, in the RaDaR pilot study that included patients with HR⁺ disease, the median time from ctDNA detection to relapse was long at 16.8 months. However, only a future prospective study, or studies in clinical settings where serial imaging is routinely performed, could ascertain whether imaging at the point of ctDNA detection with RaDaR will identify patients without radiological recurrence.

Emerging data suggest that sites of relapse may differ in their release of ctDNA. Patients with lung-only metastases had lower levels of ctDNA detection than other sites of disease. However, it must be highlighted that in this study, imaging was conducted on those randomized to pembrolizumab, at the point of ctDNA detection via a dPCR assay. This may reflect the increased ability to detect low levels of recurrence in the lungs, with similarly low levels of disease less easily detectable in other sites, or could reflect lower shedding into the circulation from lung metastases. Interestingly, this finding concurs

with the relatively low rates of ctDNA detection in patients with stage I/II primary non-small cell lung cancer (16), suggesting that reduced shedding is a contributory factor. A relatively high number of patients in the study had apparently local-only relapse, which may reflect the ability of imaging to detect local relapse, even though these patients conceivably had imaging occult distant disease that also contributed to ctDNA detection.

We report here a longer follow-up of ctDNA detection in the cTRAK-TN study than previously reported. With this longer follow-up, we find that patients with ctDNA not detected at the first time point, but detected at a subsequent time point, had a longer lead time from ctDNA detection to relapse. Intriguingly, this group also included a set of patients with a long time to relapse, with 34% of patients not having progressed at 12 months from ctDNA detection. This finding identifies a subgroup of TNBC that is more indolent, at least in the MRD setting and a heterogeneity in clinical behavior that was not previously appreciated. However, the non-uniform approach to imaging in those randomized to pembrolizumab and observation limits the study of unique characteristics of patients who did not relapse during the study. It will be interesting in future research to understand the distinct biology of this group of patients, including assessing the gene-expression profiles of their primary tumors, although we speculate this may reflect immune surveillance of micrometastases.

In summary, we have demonstrated that the increased analytical sensitivity of personalized multimutation sequencing assays translated to detection of MRD with a longer median time to relapse than dPCR assays in the c-TRAK TN clinical trial. This also resulted in substantially shorter required durations of ctDNA surveillance to detect future recurrences. These findings strongly support the implementation of personalized sequencing assays in clinical trials but also emphasize the potential benefits of further assay development to further improve analytical sensitivity and detect lower levels of ctDNA and further improve ctDNA-based MRD detection.

Authors' Disclosures

M. Coakley reports grants from the Wellcome Trust during the conduct of the study. G. Villacampa reports personal fees from MSD, Pfizer, Pierre Fabre, GSK, and Reveal Genomics outside the submitted work. L. Kilburn reports grants from Cancer Research UK during the conduct of the study. K. Goddard reports grants from Cancer Research UK Core grant (C1491/A15955) and Merck Sharpe & Dohme (MSD) c-TRAK TN Grant during the conduct of the study. P. Hall reports grants from SeaGen, Lilly, AbbVie, Eisai, Novartis, Gilead, and Sanofi outside the submitted work. C. Harper-Wynne reports personal fees from Pfizer Pharmaceuticals, Roche, Lilly, Novartis, Gilead, Seagen, and AstraZeneca pharma outside the submitted work. T. Hickish reports grants from University Hospitals Dorset during the conduct of the study and personal fees and other support from iQ HealthTech outside the submitted work. I. Macpherson reports personal fees and nonfinancial support from Roche, Novartis, Gilead, Eli Lilly, Daiichi Sankyo, and Eisai and personal fees from Pfizer, AstraZeneca, MSD, Pierre Fabre, Stemline Therapeutics, and Owkin outside the submitted work. A. Okines reports grants and personal fees from Roche and Pfizer and personal fees from Lilly, Seagen, Esai, Gilead, and AstraZeneca outside the submitted work. A. Wardley reports other support from CRUK, NIHR, and Seagen during the conduct of the study and personal fees and other support from Roche, Novartis, and Daiichi Sankyo outside the submitted work. D. Wheatley reports personal fees from Roche, Novartis, AstraZeneca, and MSD outside the submitted work. S. Waters reports personal fees from Novartis, Sanofi UK, Daiichi Sankyo UK Ltd., Gilead Sciences, and Eli Lilly & Co. outside the submitted work. C. Palmieri reports grant funding from Pfizer and Daiichi Sankyo and honoraria from Pfizer, Roche, Daiichi Sankyo, Novartis, Exact Sciences, Gilead, SeaGen, and Eli Lilly. M. Winter reports nonfinancial support from Novartis and personal fees from Eli Lilly, Gilead, Roche, Eisai, and Pfizer outside the submitted work. J. Bliss reports grants from

Merck Sharp & Dohme, ICR NIHR Biomedical Research Centre, AstraZeneca, Puma Biotechnology, Clovis Oncology, Pfizer, Janssen-Cilag, Novartis, Eli Lilly, and Roche and grants and other support from Cancer Research UK during the conduct of the study; grants from Merck Sharpe & Dohme outside the submitted work. N.C. Turner reports advisory board honoraria from AstraZeneca, Lilly, Pfizer, Roche/Genentech, Novartis, GlaxoSmithKline, Repare Therapeutics, Relay Therapeutics, Zentalis, Gilead, Inivata, Guardant, and Exact Sciences; and research funding from AstraZeneca, Pfizer, Roche/Genentech, Merck Sharpe & Dohme, Guardant Health, Invitae, Inivata, Personalis, and Natera. No disclosures were reported by the other authors.

Authors' Contributions

M. Coakley: Data curation, investigation, visualization, methodology, writing—original draft. **G. Villacampa:** Formal analysis, visualization, methodology, writing—original draft. **P. Sritharan:** Data curation. **C. Swift:** Writing—review and editing. **K. Dunne:** Writing—review and editing. **L. Kilburn:** Methodology, project administration, writing—review and editing. **K. Goddard:** Data curation, project administration, writing—review and editing. **C. Pipinikas:** Writing—review and editing. **P. Rojas:** Writing—review and editing. **W. Emmett:** Writing—review and editing. **P. Hall:** Writing—review and editing. **C. Harper-Wynne:** Writing—review and editing. **T. Hickish:** Writing—review and editing. **I. Macpherson:** Writing—review and editing. **A. Okines:** Writing—review and editing. **A. Wardley:** Writing—review and editing. **D. Wheatley:** Writing—review and editing. **S. Waters:** Writing—review and editing. **C. Palmieri:** Writing—review and editing. **M. Winter:** Writing—review and editing. **R.J. Cutts:** Data curation, validation, writing—review and editing. **I. Garcia-Murillas:** Supervision, methodology, project administration, writing—review and editing. **J. Bliss:** Supervision, methodology, writing—review and editing. **N.C. Turner:** Conceptualization, resources, supervision, validation, writing—original draft, project administration.

References

- Powles T, Gschwent J, Bracarda S, Castellano D, Gross-Goupil M, Jensen J, et al. 716TIP IMvigor011: a global, double-blind, randomized phase III study of atezolizumab (atezo; anti-PD-L1) vs placebo (pbo) as adjuvant therapy in patients (pts) with high-risk muscle-invasive bladder cancer (MIBC) who are circulating tumor (ct)DNA+ post cystectomy. *Ann Oncol* 2021;32.
- Turner N, Swift C, Jenkins B, Kilburn L, Coakley M, Beaney M, et al. Results of the c-TRAK TN trial: a clinical trial utilizing ctDNA mutation tracking to detect molecular residual disease and trigger intervention in patients with moderate- and high-risk early-stage triple-negative breast cancer. *Ann Oncol* 2023;34:200–11.
- Pascual J, Attard G, Bidard F, Curigliano G, De Mattos-Arruda L, Diehn M, et al. ESMO recommendations on the use of circulating tumor DNA assays for patients with cancer: a report from the ESMO Precision Medicine Working Group. *Ann Oncol* 2022;33:750–68.
- Kenneth S, Brant R, Carr T, Dearden S, Jenkins S, Brown H, et al. EGFR mutation detection in ctDNA from NSCLC patient plasma: a cross-platform comparison of leading technologies to support the clinical development of AZD9291. *Lung Cancer* 2015;90:509–15.
- Xu T, Kang X, You X, Dai L, Tian D, Yan D, et al. Cross-platform comparison of four leading technologies for detecting EGFR mutations in circulating tumor DNA from non-small cell lung carcinoma patient plasma. *Theranostics* 2016;7:1437–46.
- Stetson D, Ahmed A, Xu X, Nuttal B, Lubinski T, Johnson J, et al. Orthogonal comparison of four plasma NGS tests with tumor suggests technical factors are a major source of assay discordance. *JCO Precision Oncology* 2019;3:1–9.
- Yu L, Lopez G, Rassa J, Wang Y, Basavanahally T, Browne A, et al. Direct comparison of circulating tumor DNA sequencing assays with targeted large gene panels. *PLoS One* 2022;17:e0266889.
- Merker JD, Oxnard GR, Compton C, Diehn M, Hurley P, Lazar AJ, et al. Circulating tumor DNA analysis in patients with cancer: American Society of Clinical Oncology and College of American Pathologists Joint Review. *J Clin Oncol* 2018;36:1631–41.
- Garcia-Murillas I, Chopra N, Comino-Mendez I, Beaney M, Tovey H, Cutts R, et al. Assessment of molecular relapse detection in early-stage breast cancer. *JAMA Oncol* 2019;5:1473–8.
- Garcia-Murillas I, Schiavon G, Weigelt B, Ng C, Hrebien S, Cutts R, et al. Mutation tracking in circulating tumor DNA predicts relapse in early breast cancer. *Sci Transl Med* 2015;7:302ra133.
- Pearson A, Proszek P, Pascual J, Fribbens C, Shamsher M, Kingston B, et al. Inactivating NF1 mutations are enriched in advanced breast cancer and contribute to endocrine therapy resistance. *Clin Cancer Research* 2020;26:608–22.
- Pascual J, Lim J, Macpherson I, Armstrong A, Ring A, Okines A, et al. Triplet therapy with palbociclib, taselisib, and fulvestrant in PIK3CA mutant breast cancer and doublet palbociclib and taselisib in pathway mutant solid cancers. *Cancer Discov* 2021;11:92–107.
- O'Leary B, Hrebien S, Morden J, Beaney M, Fribbens C, Huang X, et al. Early circulating tumor DNA dynamics and clonal selection with palbociclib and fulvestrant for breast cancer. *Nat Commun* 2018;9:896.
- Flach S, Howarth K, Hackinger S, Pipinikas C, Ellis P, McLay K, et al. Liquid BiOpsy for MiNimal RESidual DiSease detection in head and neck squamous cell carcinoma (LIONESS)—a personalized circulating tumor DNA analysis in head and neck squamous cell carcinoma. *Br J Cancer* 2022;126:1186–95.
- Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* 2009;25:1754–60.
- Gale D, Heider K, Ruiz-Valdepenas A, Hackinger S, Perry M, Marsico G, et al. Residual ctDNA after treatment predicts early relapse in patients with early-stage non-small cell lung cancer. *Ann Oncol* 2022;33:500–10.

Acknowledgments

This work was supported in part by Le Cure, the Royal Marsden Cancer Charity. It was endorsed by Cancer Research UK (CRUKE/16/024) and the ICR Clinical Trials & Statistics Unit is supported by a core program grant (grant number C1491/A15955) from Cancer Research UK. This study represents independent research funded by the National Institute for Health Research (NIHR) Biomedical Research Centre at The Royal Marsden NHS Foundation Trust and The Institute of Cancer Research, London. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care. This work was supported in part by provision of drug by the Investigator Initiated Studies Program of Merck Sharp & Dohme Limited. The opinions expressed in this article are those of the authors and do not necessarily represent those of Merck Sharp & Dohme Limited. This work uses data provided by patients and collected by the NHS as part of their care and support. We thank the patients and all investigators and research support staff at the participating centres. We also thank our patient representatives from the Independent Cancer Patients' Voice patient advocate group, members of the TMG, Trial Steering Committee, and Independent Data Monitoring Committee.

The publication costs of this article were defrayed in part by the payment of publication fees. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

Note

Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

Received August 11, 2023; revised October 16, 2023; accepted December 6, 2023; published first December 11, 2023.