

This is a repository copy of *Unraveling Island Economies through Organic Residue Analysis: The Case of Mocha Island (Southern Chile)*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/211331/>

Version: Published Version

---

**Article:**

Montalvo-Cabrera, Javier A., Colonese, André C. orcid.org/0000-0002-0279-6634, Campbell, Roberto et al. (5 more authors) (2024) Unraveling Island Economies through Organic Residue Analysis: The Case of Mocha Island (Southern Chile). *Latin American Antiquity*. pp. 1-15. ISSN: 1045-6635

<https://doi.org/10.1017/laq.2023.58>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

ARTICLE

# Unraveling Island Economies through Organic Residue Analysis: The Case of Mocha Island (Southern Chile)

Javier A. Montalvo-Cabrera<sup>1</sup> , André C. Colonese<sup>2</sup>, Roberto Campbell<sup>3</sup>, Helen M. Talbot<sup>1</sup>, Alexandre Lucquin<sup>1</sup>, Marjolein Admiraal<sup>1</sup>, Gabriela Palma<sup>4</sup>, and Oliver E. Craig<sup>1</sup>

<sup>1</sup>BioArCh, Department of Archaeology, University of York, United Kingdom, <sup>2</sup>Departament de Prehistòria, Institut de Ciència i Tecnologia Ambientals (ICTA), Universitat Autònoma de Barcelona, Spain, <sup>3</sup>Escuela de Antropología, Pontificia Universidad Católica de Chile, Santiago, Chile, and <sup>4</sup>Independent researcher

**Corresponding author:** Javier A. Montalvo-Cabrera; Email: [jam603@york.ac.uk](mailto:jam603@york.ac.uk)

(Received 24 October 2022; revised 7 July 2023; accepted 25 October 2023)

## Abstract

Biophysical conditions played a fundamental role in early human colonization of insular territories, particularly in food-producing societies dealing with limited resources and the challenges of maintaining a sustainable carrying capacity. Studies on past human colonization of small oceanic islands thus offer insights into economic plasticity, ecological impacts, and adaptation of early food-producing groups. On the coast of southern Chile, early evidence is dated to 950 cal BP of island colonization by coastal populations with mainland subsistence systems based on the exploitation of marine resources, along with gathering, managing, and cultivating plants and hunting terrestrial animals. Strikingly, the extent to which these mixed economies contributed to insular colonization efforts is largely unknown. Here we used organic residue analysis of ceramic artifacts to shed light on the subsistence of populations on Mocha Island in southern Chile. We extracted and analyzed lipids from 51 pottery sherds associated with the El Vergel cultural complex that flourished in southern Chile between 950 and 400 cal BP. Chemical and stable isotope analysis of the extracts identified a range of food products, including C<sub>3</sub> and C<sub>4</sub> plants and marine organisms. The results reveal the central role of mixed subsistence systems in fueling the colonization of Mocha Island.

## Resumen

Las condiciones biofísicas desempeñaron un papel fundamental en la temprana colonización humana de territorios insulares, sobre todo en sociedades productoras de alimentos que se enfrentaban a recursos limitados y a los desafíos de mantener una capacidad de carga sostenible. Los estudios sobre la colonización humana de pequeñas islas oceánicas en el pasado ofrecen, por lo tanto, información sobre la plasticidad económica, los impactos ecológicos y la adaptación de los primeros grupos productores de alimentos. En la costa del sur de Chile, los primeros indicios de colonización de islas datan de hace 950 años cal aP por parte de poblaciones costeras con sistemas de subsistencia en tierra firme basados en la explotación de los recursos marinos, además de la recolección, manejo y cultivo de plantas, y la caza de animales terrestres. Sorprendentemente, se desconoce en qué medida estas economías mixtas contribuyeron a los esfuerzos de colonización insular. En este trabajo, empleamos el análisis de residuos orgánicos de artefactos cerámicos para elucidar la subsistencia de las poblaciones de la isla Mocha, en el sur de Chile. Extrajimos y analizamos lípidos de 51 fragmentos cerámicos asociados al complejo cultural El Vergel, que se desarrolló en el sur de Chile entre 950 y 400 años cal aP. El análisis químico y de isótopos estables de los extractos identificó una serie de productos alimenticios, incluyendo plantas C<sub>3</sub> y C<sub>4</sub>, así como organismos marinos. Los resultados revelan el papel central de los sistemas mixtos de subsistencia en el impulso de la colonización de la isla Mocha.

**Keywords:** pottery; organic residue analysis; island colonization; southern Chile; El Vergel; mixed economies

**Palabras clave:** cerámica; análisis de residuos orgánicos; colonización de islas; sur de Chile; El Vergel; economías mixtas

© The Author(s), 2024. Published by Cambridge University Press on behalf of Society for American Archaeology. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

The socioecological drivers of early human colonization of oceanic islands have been the subject of a contentious debate in archaeology (Braje et al. 2017; Terrell 2023). Insular biogeographic conditions, distance from mainland enclaves, seafaring technology, and the flexibility of a subsistence economy were some of the interplaying factors conditioning island-colonization efforts (Crowther et al. 2016; Leppard 2014; O'Connor et al. 2019; Takamiya 2006). Over the preceding several decades, genetic, archaeological, and linguistic studies have increased our understanding of these processes in regions such as the west and northeast Pacific Ocean (Horsburgh and McCoy 2017; Lepofsky et al. 2015; McFadden et al. 2021; Potter and White 2009; Walworth 2014). Despite extensive archaeological research along the western coast of South America (Beresford-Jones et al. 2018; King et al. 2018; Reyes et al. 2022; Stothert et al. 2003), large geographic areas of the Chilean coast remain poorly understood.

On the Pacific coast of South America, the productive Humboldt upwelling system (3°24'S–54°55'S) has sustained coastal adapted populations since the Late Pleistocene. Throughout the Holocene, coastal communities developed mixed economies with distinct degrees of reliance on marine and terrestrial-based resources (Dillehay et al. 2022; Knudson et al. 2015; Pearsall et al. 2020). In southern Chile the intensification of food production during the Late Ceramic period (950–400 cal BP) led to sedentarism and a remarkable demographic upsurge (Campbell and Quiroz 2015). During this time, farming groups from the El Vergel cultural complex migrated and permanently settled on small offshore islands (Campbell 2015; Campbell and Quiroz 2015). From here, the occupation of Mocha Island, the largest, most isolated island, lying 35 km offshore, is an intriguing case. The proliferation of El Vergel residential sites and the construction of monumental architecture by a large workforce seem to suggest that a dense, complex, and permanently settled society was established here (Campbell 2015; Campbell and Pfeiffer 2017). However, Mocha Island's biogeographic characteristics denote limited terrestrial resources, making for unsuitable conditions for supporting large groups of people (Campbell 2015; Pefaur and Yáñez 1980).

The translocation of domestic plants and tamed camelids is argued to have been a prerequisite for the permanent settlement of Mocha Island by the El Vergel groups (Becker 1997a; Campbell 2015; Godoy-Aguirre 2018; Roa et al. 2015, 2021). Preliminary results on lipid distributions extracted from Mocha Island pottery fragments indicated a preferential use of terrestrial resources, including the exploitation of terrestrial animals (Roa et al. 2021). Carbon ( $\delta^{13}\text{C}$ ) and nitrogen ( $\delta^{15}\text{N}$ ) stable isotope analyses of El Vergel individuals from Mocha Island suggest a diet largely based on terrestrial resources with only a low contribution of marine proteins (Campbell et al. 2020). The results clearly show the predominance of a mixed diet.  $^{13}\text{C}$  enrichment in the individuals is attributed to the consumption of locally cultivated maize (Campbell et al. 2020), while marine resources are claimed to have played a marginal role in the foodways and adaptive process of El Vergel farmers—despite the island's location within one of the most productive upwelling systems of the world (Campbell et al. 2020; Roa et al. 2021).

Attributing human bone collagen  $^{13}\text{C}$  enrichment to the sustained consumption of  $\text{C}_4$  plants such as maize is problematic when marine resources are involved because of the possible overlapping of  $\delta^{13}\text{C}$  values (Corr et al. 2005). Moreover, mollusk remains were found at every residential site on Mocha Island, along with marine mammal and fish bones; in some cases, they outnumbered terrestrial animal remains (Power 2013; Roa et al. 2021). The identification of net sinkers and fishing hooks in some sites indicates the use of specialized technology for exploiting local marine resources (Martínez 2013, 2015; Quiroz and Sánchez 2005). The clear evidence for the use of marine commodities and the possible methodological biases suggest that the role of marine products in El Vergel farmers' foodways on Mocha Island is not well understood and needs further assessment.

Here we present the first organic residue analysis from the Pacific coast of South America using single-compound stable carbon isotope composition and molecular evidence extracted from pottery sherds to evaluate the use of marine resources by El Vergel coastal farmers settling on Mocha Island. This analysis enabled us to assess local foodways, the use pottery might have had for processing marine resources, and whether this reflects the adaptive and socioeconomic strategies developed during Mocha Island's colonization.

### Mocha Island: Environmental Setting

Mocha Island is an offshore small insular territory of 52 km<sup>2</sup> that is located 35 km from the mainland (Figure 1). It originates as the highest section of an emerging ridge from the continental plaque, continuously rising above sea level for the last 6,000 years (Melnick et al. 2003; Nelson and Manley 1992). It comprises two main mountain ranges of 390 m asl and is covered by a dense evergreen forest that is surrounded by a coastal plain of beaches and meadows (Bahlburg and Spiske 2015; Prieto 1997). Annual precipitation of around 1,350 mm forms small rivers and lakes supplying the island with fresh drinkable water (Pefaur and Yáñez 1980; Prieto 1997).

The composition of its flora has remained relatively unchanged for the last 2,000 years: it is mainly dominated by evergreen species such as *Aextoxicon punctatum*, *Drimys winteri*, and *Azara lanceolata* and a variety of Myrtaceae (Lequesne et al. 1999). This landscape also contains a range of wild edible plants, mainly *Aristotelia chilensis* (Chilean wineberry), *Fragaria chiloensis* (Chilean strawberry), *Rubus geoides* (wild raspberry), *Gevuina avellana* (Chilean hazelnut), and *Madia sativa* (Roa et al. 2015, 2021).

The native fauna comprised various small rodents, birds, reptiles, and amphibian species; no large terrestrial mammals were present before the translocation of mainland animals by humans (Pefaur and Yáñez 1980; Reiche 1903; Saavedra et al. 2003). Local marine fauna are abundant and include mollusks, crustaceans, marine mammals, and different fish species (Báez 1997; Gálvez 1997; Rebolledo 2013; Reiche 1903).


### The El Vergel Cultural Complex and Pottery Production

The El Vergel cultural complex consisted of farming groups that cultivated domestic crops, tamed wild animals, and engaged in hunting-and-gathering practices (Becker 1997b; Contreras et al. 2005; Roa et al. 2018). The main characteristic element of the El Vergel cultural complex is its pottery. Decoration using red or black geometric motifs over a white slip surface is a distinctive feature of the El Vergel pottery found in funerary contexts (Adán et al. 2005; Aldunate 1989; Bahamondes et al. 2006). Large ceramic containers were common, some of which functioned as funerary urns; however, these containers were also found in residential settings (Bullock 1970; Gordon 1978; Navarro and Aldunate 2002). Ethnographic investigations within the Mapuche people, descendants of the El Vergel cultural complex, described the use of similar large containers for culinary purposes, mainly the storage and preparation of fermented maize or wheat beverages known as *chicha* or *muday* (Alvarado 1997). Other types, such as nondecorated monochrome vessels and pots with a red slip surface, are frequently found in residential sites (Adán et al. 2005; Quiroz 2001). These pots correspond to various vessel types such as jars, cooking pots, mugs, and dishes, presenting specific morphological characteristics that denote a long-standing pottery tradition in southern Chile (Adán et al. 2005; Aldunate 1989; Dillehay 1990; López 2017; Palma 2013).

On Mocha Island, diagnostic elements found in residential sites confirm the presence of the El Vergel cultural complex (Campbell 2020; Campbell and Pfeiffer 2017; Roa et al. 2021). Its population comprised autonomous communities with equal access to local and exotic resources (Campbell 2020). Camelids or *chilihueque*, the name for local guanaco (*Lama guanicoe*), were translocated from the mainland and exploited at every residential site for fur and meat (Becker 1997a; Roa et al. 2021; Westbury et al. 2016). Domesticated crops, specifically quinoa (*Chenopodium quinoa*), maize (*Zea mays*), and beans (*Phaseolus vulgaris*), were cultivated along with gathering wild local plants (Roa et al. 2015, 2021; Rojas and Cardemil 1995). The presence of mollusks, marine birds and mammals, and fish remains at every residential site confirms generalized access to coastal foraging, hunting, and fishing, mainly nearshore fishing (Gálvez 1997; Martínez 2014, 2015; Power 2013; Rebolledo 2013; Roa et al. 2021).

### Materials and Methods

We selected 51 pottery sherds chronologically associated with the Late Ceramic period from three residential sites located at different points of the island (Table 1). These sherds comprised polished, smoothed, and eroded monochrome pottery fragments from the rim, neck, body, and base (Supplemental Table 1). Only samples with a thickness greater than 7.25 mm were selected. The



**Figure 1.** (A) Mocha Island in South America; (B) archaeological sites at Mocha Island; (C) Mocha Island facing the coast of Chile.

**Table 1.**  $^{14}\text{C}$  Dates from Associated Material.

Site	Number of Samples	Laboratory Code	Excavation Unit	Material	Taxa	$^{14}\text{C}$ Age BP	Median Probability cal yrs BP
P5-1	17	UB-26214	05.02.03	Kernel	<i>Zea mays</i>	552 $\pm$ 26	530
		UB-26216	05.02.03	Bone (Phalange)	Camelidae	605 $\pm$ 26	554
		UB-26215	05.02.03	Cobb	<i>Zea mays</i>	635 $\pm$ 25	607
P23-2	15	UB-29290	23.03.01	Kernel	<i>Zea mays</i>	1108 $\pm$ 28	961
		UB-29289	23.03.01	Seed	<i>Phaseolus vulgaris</i>	679 $\pm$ 25	603
P25-1	19	UB-29287	25.03.04	Kernel	<i>Zea mays</i>	644 $\pm$ 35	605
		UB-29288	25.03.04	Kernel	<i>Zea mays</i>	661 $\pm$ 30	606

Source: Data from Campbell and Pfeiffer (2017:Supplemental Tables 10–15).

identification of soot marks in some of them potentially indicates their exposure to fire. Based on these characteristics, the selected fragments likely represent cooking pots and large containers, mainly used for processing food resources (Albán et al. 2013).

To obtain molecular and  $\delta^{13}\text{C}$  references of local resources, cooking experiments were conducted using seven nontempered replica vessels (Bondetti et al. 2020; see Supplemental Table 2 and Supplemental Text 1). Each commodity was cooked in a specific container, with no mixing between resources.

### Organic Residues Analysis

A small section of the inner surface of the 51 pottery fragments was cleaned before drilling 2–3 mm into the potsherds to obtain 1 g of pottery powder for acid extraction (AE). A one-step acidified

methanol protocol was followed (Craig et al. 2013). A blank was included as a control to assess contamination throughout the extraction process. An internal standard (*n*-tetratriacontane (C<sub>34</sub>): 10 µg) was incorporated in each tube. Methanol (4 mL) and sulphuric acid (800 µL) were added and the mixture was subsequently sonicated and centrifuged before being placed on a heating block for four hours at 70°C. Three successive extractions using *n*-hexane allowed the lipids' separation before their neutralization using potassium carbonate. Subsequently, lipids were concentrated and resuspended in *n*-hexane at appropriate dilution for gas chromatography-mass spectrometry (GC-MS) analysis. An internal standard (*n*-hexatriacontane (C<sub>36</sub>): 10 µg) was added at the end of the extraction to infer lipid yield and concentration.

Another 1 g was prepared from nine potsherds and subjected to solvent extraction (SE). The extraction procedure followed a protocol similar to that described by Colonese and coworkers (2017). Lipids were extracted by sonication from the ceramic powder using dichloromethane:methanol DCM:MeOH 2/1 v/v three times. A blank was also prepared to assess contamination. After the centrifugation, the liquid fraction was transferred into clean and labeled hydrolysis vials, and their content was evaporated to complete dryness under a gentle stream of nitrogen. One aliquot of the sample was resuspended in 50 µl of *n*-hexane and derivatized by adding four drops of N, O-bis (trimethylsilyl) trifluoroacetamide with 1% trimethylchlorosilane (BSTFA+TMCS, 99:1). Samples were placed on a heating block for one hour at 70°C and then dried under a gentle stream of nitrogen. Subsequently, samples were resuspended in *n*-hexane for their analysis by GC-MS.

Seven acid extracted samples were also derivatized with BSTFA+TMCS, 99:1, to identify very long-chain fatty alcohols. Samples were transferred from their respective GC-vial inserts by adding 100 µl of *n*-hexane into each vial and transferring it into a clean hydrolysis vial (x3). Samples were dried and resuspended in 50 µl of hexane, and four drops of BSTFA+TMCS, 99:1, and a flush of nitrogen were added before they were placed on a heating block at 70°C for one hour. Samples were concentrated and then transferred to a new GC-vial with a conical insert for their analysis within 48 hours.

Around 1 g of pottery powder was obtained from each of the seven replica vessels used in the cooking experiments. An extraction protocol using acidified methanol was also followed.

Lipid extracts were screened and quantified using an Agilent 7890B high-temperature gas chromatograph (Agilent Technologies, Cheadle, Cheshire, UK) equipped with a flame ionization detector (HT-GC-FID). A 100% Dimethylpolysiloxane DB-1 column (15 m × 320 µm × 0.1 µm; J&W Scientific, Folsom, California, USA) was used. A splitless injector was used to inject 1 µL of acid-extracted samples into the GC at 300°C. The carrier gas was helium, with a constant flow rate of 2 mL min<sup>-1</sup>. The temperature of the oven was set at 100°C for two minutes and then increased to 20°C min<sup>-1</sup> up to 325°C, holding for three minutes.

A splitless injection was also used to inject 1 µL of solvent-extracted samples at 350°C. The temperature of the oven was set at 50°C for two minutes and then increased 10°C min<sup>-1</sup> up to 375°C, settling for 10 minutes.

Acid-extracted and solvent-extracted samples were analyzed using an Agilent 7890A series chromatograph attached to an Agilent 5975C inert XL mass selective detector with a quadrupole mass analyzer (Agilent Technologies). The column used was a methylpolysiloxane (5%-phenyl) DB-5ms (30 m × 0.25 mm × 0.25 µm; J&W Scientific). Samples were injected (1 µL) using a splitless injector at 300°C. Helium was used as the carrier gas, at a constant flow rate of 2 mL min<sup>-1</sup>. The temperature of the oven was set at 50°C for two minutes and then rose 10°C min<sup>-1</sup> to 325°C, where it settled for 15 minutes. The mass spectrometer ionization energy was 70 eV, obtaining a spectrum by scanning ions between m/z 50 and 800.

Preliminary results using HT-GC-FID indicated the possible presence of triacylglycerols in some samples (Roa et al. 2021). We used high-temperature gas chromatography-mass spectrometry (HT-GC-MS) to assess their presence in the pots. TMS solvent extracts were analyzed using a Perkin Elmer Clarus 690 gas chromatograph coupled to a SQ8-T mass spectrometer. The column was a DB5-HT column (30 m × 0.25 mm × 0.1 µm). Helium was used as the carrier gas at a constant flow rate of 1 mL min<sup>-1</sup>. The temperature of the oven was set at 50°C for two minutes and then increased to 10°C min<sup>-1</sup>, reaching a maximum temperature of 375°C held for 15 minutes.



The identification of isoprenoid acids and  $\omega$ -(*o*-alkylphenyl) alkanolic acids (APAAs) as aquatic biomarkers was performed using a GC-MS equipped with a 50% cyanopropyl-methylpolysiloxane DB-23 column (60 m  $\times$  0.25 mm  $\times$  0.25  $\mu$ m; J&W Scientific) as in the study by Shoda and coworkers (2017). A splitless injector was used to inject 1  $\mu$ L of the acidified methanol extracts into the GC at 300°C. Helium was used as the carrier gas, with a flow rate of 1.5 mL min<sup>-1</sup>. The temperature of the oven was set at 50°C for two minutes and then rose 10°C min<sup>-1</sup> to 100°C, after which it increased 4°C min<sup>-1</sup> up to 140°C then 0.5°C min<sup>-1</sup> up to 160°C, and finally by 20°C min<sup>-1</sup>, reaching 250°C where it settled for 20 minutes. The mass spectrometer was operated in single ion monitoring (SIM) mode to achieve higher sensitivity of target compounds. Selected ions allowed the identification of the three main isoprenoid acids (*m/z* 74, 101, 171, and 326 for phytanic acid; *m/z* 74, 88, 101, and 312 for pristanic acid; and *m/z* 74, 87, 213, and 270 for 4, 8, 12-trimethyltridecanoic acid [4, 8, 12-TMTD]), and APAAs of carbon length C<sub>16</sub>–C<sub>22</sub> (*m/z* 74, 105, 262, 290, 318, 346).

MSD Chemstation F.01.03.2357 software was used to compute the GC-MS results. Compounds were identified according to retention time and mass spectrum and by comparison with the National Institute of Standards and Technology (NIST) library. Peaks were integrated using Agilent Mass Hunter Quantitative Analysis software version B.07.01/ Build7.1.524.0 for GC-MS.

### GC-c-IRMS and Single-Compound $\delta^{13}\text{C}$ Analysis

We selected 49 samples for  $\delta^{13}\text{C}$  analysis of the main alkanolic acids—C<sub>16:0</sub> and C<sub>18:0</sub>—based on a minimal alkanolic acid quantity needed for injection. These samples were analyzed using a Delta V Advantage Isotope Ratio Mass Spectrometer (Thermo Fisher Scientific, Bremen, Germany), coupled to a Trace Ultra 1310 Gas Chromatograph (Thermo Fisher) with a GC IsoLink II interface (with a CuO combustion reactor held at 850°C). The column was an ultra-inert fused-silica DB-5 ms UI (60 m  $\times$  0.25 mm  $\times$  0.25  $\mu$ m; J&W Scientific), into which 1  $\mu$ L of each sample was injected for analysis. Ultra-high-purity grade helium was used as the carrier gas with a flow rate of 2 mL min<sup>-1</sup>, from which a small part of the flow was diverted to an ISQ mass spectrometer (Thermo Fisher) for parallel acquisition of molecular data. The temperature of the oven was set at 50°C for 0.5 min, and then increased 25°C min<sup>-1</sup> to 175°C and then 8°C min<sup>-1</sup> until it reached 325°C and then settled for 20 minutes.

Eluted compounds were ionized in the mass spectrometer through electron impact. The intensities of ions *m/z* 44, 45, and 46 were recorded to automatically compute the <sup>13</sup>C/<sup>12</sup>C ratios of each peak in the extracts. The software used for the computation were Isodat (Thermo Fisher Scientific) and LyticOS (Isoprime, Cheadle, UK). Computation was based on the comparison with the repeatedly measured standard reference gas (CO<sub>2</sub>). Calculated  $\delta^{13}\text{C}$  values are presented in parts per mil (‰), relative to the Vienna PeeDee Belemnite (V-PDB) international standard.

Batches were calibrated using a linear calibration curve (average  $R^2 = 0.996 \pm 0.002$  in 16 batches) based on expected versus measured  $\delta^{13}\text{C}$  values of *n*-alkanes and *n*-alkanoic acid esters from international standards (Indiana A7 and F8-3 mixtures). The accuracy of the instrument was determined on *n*-alkanoic acid methyl esters of known isotopic composition (Indiana F8-3, 18 measurements). The mean and standard deviation of these compounds were  $-29.89 \pm 0.1\text{‰}$  for C<sub>16:0</sub> (reported mean value vs. V-PDB  $-29.90 \pm 0.03\text{‰}$ ) and  $-23.41 \pm 0.06\text{‰}$  for C<sub>18:0</sub> (reported mean value vs. V-PDB  $-23.24 \pm 0.13\text{‰}$ ). The precision was based on a laboratory standard mixture regularly injected between samples (108 measurements). Alkanolic acids mean  $\pm$  SD values were  $-30.70 \pm 0.15\text{‰}$  for C<sub>16:0</sub> methyl ester and  $-26.44 \pm 0.15\text{‰}$  for C<sub>18:0</sub> methyl ester. After this analysis, values were corrected for the methylation of the alkanolic acids that occurred during acid extractions. Corrections were performed using a mass balance formula to compare the values with a standard mixture of C<sub>16:0</sub> and C<sub>18:0</sub> fatty acids of known isotopic composition, which was already included in each batch during the acid extractions. Modern references were adjusted according to the atmospheric  $\delta^{13}\text{C}$  variations between the Early Holocene Epoch and the present to better interpret the archaeological data (Hellevang and Aagaard 2015).

## Results

Lipid concentrations from the acid extracted fraction ranged from 3 to 360  $\mu\text{g g}^{-1}$ , with a mean of 57.4  $\mu\text{g g}^{-1}$ . Only one sample (P25-1.19) yielded lipids below the interpretable limit of  $>5 \mu\text{g g}^{-1}$ .

(Evershed 2008) and was excluded from further analysis (Supplemental Table 3). Lipid profiles exhibited a mixture of aliphatic compounds comprising a distribution of saturated fatty acids ( $C_{8:0}$ – $C_{30:0}$ ), monounsaturated fatty acids ( $C_{14:1}$ – $C_{24:1}$ ), fatty alcohols ( $C_{12}$ – $C_{32}$ ),  $\alpha$ ,  $\omega$ -dicarboxylic acids ( $C_8$ – $C_{14}$ ), and branched and linear fatty acids  $C_{15:0}$  and  $C_{17:0}$ . Overall, from the lipid distributions and biomarker identification it was possible to distinguish between the processing of aquatic and plant commodities in the El Vergel pots.

### *The El Vergel Pottery and Its Use for Processing Aquatic Organisms*

Four El Vergel pots (8%) had a full range of aquatic biomarkers comprising at least one of the three main isoprenoid acids (4, 8, 12-trimethyltridecanoic acid, pristanic, or phytanic acid), and a homologous series of  $C_{18}$ ,  $C_{20}$ , and, in one sample,  $C_{22}$  APAAs. These biomarkers are established indicators for the processing of aquatic resources (Evershed et al. 2008; Lucquin et al. 2016). APAAs are only formed by the protracted heating  $\geq 200^\circ\text{C}$  of  $C_{18:x}$ ,  $C_{20:x}$ , and  $C_{22:x}$  unsaturated fatty acids that are found in plants and terrestrial animals but mainly in aquatic commodities (Supplemental Figure 1; Bondetti et al. 2020; Evershed et al. 2008; Hansel et al. 2004) and therefore must be derived from cooking events. The ratio between the two phytanic acid diastereomers, 3S, 7R, 11R, 15-phytanic acid (SRR), and 3R, 7R, 11R, 15-phytanic acid (RRR), helps discriminate between ruminant and aquatic products. A high contribution of the SRR-diastereomer (i.e.,  $76\% \pm 16.6$ ) is likely derived from aquatic animal tissue (Supplemental Table 4; Lucquin et al. 2016). An SRR-phytanic acid diastereomer contribution higher than 70% in three of these four samples further supports an aquatic origin. The absence of freshwater commodities on the Island during the Late Ceramic period likely reflects a marine origin for the aquatic resources.

### *Molecular Evidence for Terrestrial Commodities*


Plant commodities were the most representative resources identified in the El Vergel pots. Nineteen samples (38%) had a distribution of even over odd-numbered very long-chain fatty acids and odd over even-numbered middle- and long-chain alkanes, characteristic of the epicuticular wax of plant leaves and stems (Diefendorf et al. 2011; Dove and Mayes 2006; Dunne 2021). Very long-chain fatty alcohols with an even over odd-numbered carbon chain distribution, characteristic of plant commodities (Dove and Mayes 2006), and plant sterols were identified in several samples.

Several pots ( $n = 16$ , 32%) contained APAAs with 18 carbon atoms (APAA- $C_{18}$ ) but lacked the  $C_{20}$  and  $C_{22}$   $\omega$ -(*o*-alkylphenyl) alkanolic acids. APAA- $C_{18}$  can form from a wide range of plant and animal products, but the distribution of isomers (A-I) can provide further discrimination (Supplemental Figure 2; Bondetti et al. 2020). Samples in Figure 2 had an APAA- $C_{18}$  E/H isomer ratio, possibly reflecting (1) cereals, fruits, and non-leafy vegetables; (2) terrestrial and aquatic animals; and (3) leafy plants (Bondetti et al. 2020; Dolbunova et al. 2023). Mixing cereals, fruits, and non-leafy vegetables with leafy plants may produce an intermediate E/H ratio similar to an animal origin (Bondetti et al. 2020). Only one of six samples with intermediate E/H values had a complete set of aquatic biomarkers; the rest lacked any evidence for processing aquatic or terrestrial animal products. The identification of plant epicuticular wax lipid distributions in four of these samples might indicate a plant origin.

Three samples with a full set of aquatic biomarkers had high APAA- $C_{18}$  E/H ratios similar to those found in heated local reference samples of cereals, fruits, and non-leafy vegetables (Figure 2). This evidence may indicate the use of aquatic and plant products in the same container.

A palmitic over stearic acid ratio (P/S)  $\leq 1$  identified in 46% of the samples may indicate that terrestrial animals were exploited in the El Vergel pots (Romanus et al. 2007). However, the higher solubility of  $C_{16:0}$  compared to  $C_{18:0}$  may alter the P/S ratio in animal tissue (Whelton et al. 2021). Monoacylglycerols and diacylglycerols were present in a small number of solvent-extracted samples, whereas no triacylglycerols were identified. A distribution of  $C_{15:0}$  and  $C_{17:0}$  branched and linear fatty acids was found in most of the El Vergel samples. Their presence is associated with bacterial lipids from the rumen of polygastric animals (Evershed et al. 2002). However, they can also be linked to soil bacteria contamination, making them unspecific to ruminant fat (Dudd et al. 1998). An





**Figure 2.** Boxplots of the APAA-C<sub>18</sub> E/H ratio of Mocha Island samples. Black circles represent samples with a full set of aquatic biomarkers. The black triangles are samples showing <sup>13</sup>C enrichment of their residues consistent with a C<sub>4</sub> origin. Black square, asterisks, and diamonds correspond to modern references for Chilean hazelnut, maize, and quinoa, respectively. References are based on Bondetti and others (2020) and this study (see Supplemental Table 5).

SRR-diastereomer contribution below aquatic values may indicate the processing of ruminants in only one El Vergel pot (Lucquin et al. 2016). Cholesterol was detected in some samples in small amounts; however, we cannot ascertain its archaeological origin due to the absence of clear bio-hydrogenated and oxidation derivatives (Hammann et al. 2018).


### Single-Compound Stable Carbon Isotope Analysis

To further identify the commodities' origin, we measured the stable carbon isotope values of the palmitic ( $\delta^{13}\text{C}_{16:0}$ ) and stearic ( $\delta^{13}\text{C}_{18:0}$ ) alkanolic acids extracted from the pots and then compared these values to modern and archaeological references of marine, C<sub>3</sub> plants, maize, and ruminant animals, including South American camelid *Lama glama* (Supplemental Tables 6 and 7).

Most pots (74%) had  $\delta^{13}\text{C}_{16:0}$  and  $\delta^{13}\text{C}_{18:0}$  values plotting within the ranges for C<sub>3</sub> plants. Three of the four samples showing a complete set of aquatic biomarkers were more depleted in <sup>13</sup>C relative to marine references, plotting close to C<sub>3</sub> plant values (Figure 3). The E/H ratios of these samples coincided with the use of cereals, fruits and non-leafy vegetables, probably indicating that plants and aquatic products were used in these pots.

It is likely that maize was processed in some pots with high  $\delta^{13}\text{C}$  values relative to C<sub>3</sub> plant references (Figure 3). Extracts from these pots had high APAA-C<sub>18</sub> E/H ratios coinciding with cereals, fruits, and non-leafy vegetables; a high P/S ratio; and a fatty alcohol distribution characteristic of plant commodities. The identification of C<sub>32</sub> alcohol in a high relative abundance can be associated with panicoid grasses like maize (Reber and Evershed 2004). Although C<sub>32</sub> was found in low abundance in one sample, its association with panicoid plants is debatable (Supplemental Figure 3). The different  $\delta^{13}\text{C}$  values in these samples might indicate the incorporation of other commodities, possibly C<sub>3</sub> plants, in varying proportions.

Nine samples had  $\Delta^{13}\text{C}_{(\text{C}_{18:0}-\text{C}_{16:0})}$  offsets coinciding with ruminant carcass fat references (Copley et al. 2003; Dolbunova et al. 2023). These samples yielded plant lipids and some aquatic biomarkers;



**Figure 3.** Scatterplot of compound-specific  $\delta^{13}\text{C}$  analysis of the main alkanolic acids (X-axis =  $\delta^{13}\text{C}_{16:0}$ ; Y-axis =  $\delta^{13}\text{C}_{18:0}$ ) extracted from the El Vergel pots. Blue circles indicate the four samples with a full range of aquatic biomarkers. The red circles highlight samples with a high APAA- $\text{C}_{18}$  E/H associated with cereals, fruits, and non-leafy vegetables (Bondetti et al. 2020). The 68% confidence ellipses are based on reference values published in the literature and from this study (see Supplemental Tables 6 and 7). (Color online)

however, they lacked compounds typically documented in ruminant fats, such as SRR% <64 and acylglycerol distributions. They possibly reflect the mixing of plant and animal products that can result in low  $\Delta^{13}\text{C}_{(\text{C}_{18:0}-\text{C}_{16:0})}$  values (Cramp et al. 2019; Hendy et al. 2018; Taché et al. 2021). To investigate further, we applied a simple linear mixing model based on averaged  $\delta^{13}\text{C}$  values of modern reference endpoints, adopting a progressive 10% increment (Figure 4). The hypothetical mixing lines indicate the possible use of marine and terrestrial animal products or  $\text{C}_3$  and  $\text{C}_4$  plants in these pots.


## Discussion

Organic residues from pottery artifacts demonstrated that local marine products were certainly part of the El Vergel farmers' foodways, which aligns with other evidence of marine resources exploitation on the island, such as faunal remains and fishing gear (fishhooks and net sinkers). This coastal adaptation gave the inhabitants access to a rich source of animal protein, compensating for the lack of large mammals on the island. Together our results highlight the overall importance of marine resources in the region and as a critical component of early island colonization efforts. Coastal adaptation was essential for island colonization and perhaps was more widespread than previously thought, as evidenced in the mainland El Arenal-1 site in Punta Lavapié (Contreras et al. 2005).

The single-compound  $\delta^{13}\text{C}$  analysis and more robust molecular evidence allowed us to reinterpret previous results indicating the use of pots for processing terrestrial animal fat (Roa et al. 2021). The exploitation of translocated ruminants was underrepresented in the El Vergel pots, contradicting the abundance of camelid remains found on Mocha Island (Roa et al. 2021). According to the chronicles of the first Europeans arriving on Mocha Island and southern Chile, camelids were important for local communities, which would have limited their availability for trading and consumption (Rosales 1877 [1674]:324–325; Van Meurs 1993; Van Noort [1600] quoted in Ijzerman 1926:57–58). Their exploitation as a source of animal protein cannot be ruled out because other culinary practices, such as roasting and drying the meat, were possibly used.

Marine products were processed in the pots alone or mixed with  $\text{C}_3$  plants. The  $\delta^{13}\text{C}$  values of these samples demonstrate that  $\text{C}_3$  plants were exploited intensively, given that oily and lipid-rich marine products are likely to increase  $\delta^{13}\text{C}$  values. Our results thus reveal that plants were the basis of the foodways of Mocha Island inhabitants. The use of leaves and stems and non-leafy plants, such as cereals and fruits, and non-leafy vegetables in the pots are concordant with the diversity of plants exploited by local Mocha Island farmers (Godoy-Aguirre 2018; Roa et al. 2015). Evidence for the epicuticular wax of leaves and stems was present in most El Vergel pots. The Mapuche people from southern Chile traditionally use the leaves and stems of some plant species found on Mocha Island for medicinal purposes. However, the culinary use of leaves with other commodities such as cereals and animal meat has also been documented in the past, including the preparation of various stews and other local dishes (Gay 2018 [1873]:197; Rosales 1877 [1674]:153–154).

**Figure 4.** Compound-specific  $\delta^{13}\text{C}$  analysis of Mocha Island samples plotted according to their  $\delta^{13}\text{C}_{16:0}$  value against  $\Delta^{13}\text{C}_{(C18:0 - C16:0)}$ . Blue circles indicate the four samples with a full set of aquatic biomarkers. The red circles highlight the samples with an APAA- $\text{C}_{18}$  E/H ratio associated with cereals, fruits, and non-leafy vegetables (Bondetti et al. 2020). Average  $\delta^{13}\text{C}$  endpoints were built using modern and archaeological references from published data and this study (see Supplemental Tables 6 and 7). Mixing lines with 10% increments connecting the endpoints were calculated based on the mean relative amount of each alkanoic acid present in the products. Data were gathered from the USDA database. The dashed line indicates a  $-1.26\text{‰}$  threshold set for the identification of ruminant carcass fat, as in Dolbunova and colleagues (2023). (Color online)



The lipid distribution of grains, seeds, and fruits is characterized by a low abundance of very long-chain fatty acids and alkanes, along with a P/S ratio  $>1.3$  (Dunne 2021; Dunne et al. 2016). Mixing these products with leaves and stems will likely produce a distribution similar to that seen for plant epicuticular wax, hence overshadowing their presence. The APAA- $\text{C}_{18}$  E/H ratio and  $\delta^{13}\text{C}$  values revealed the exploitation of  $\text{C}_3$  cereals, fruits, and non-leafy vegetables and their mixing with leafy plants; one may presume that the former group was overshadowed in the pots by the latter. The whole spectrum of cereals, fruits, and non-leafy vegetables may have included domestic and wild plants traditionally used in southern Chile. Chilean wineberry, Chilean strawberries, Chilean hazelnut, and *Madia sativa* seeds are wild plants used in pottery containers for roasting, preparing alcoholic beverages, and extracting oil (Godoy-Aguirre 2018; Pino Ramos et al. 2019; Schmeda-Hirschmann 1995). In the case of domestic plants, this group was dominated by beans, maize, and quinoa. However, chronicles mentioned the consumption of potatoes by Mocha Island inhabitants (Van Meurs 1993; Van Noort [1600] quoted in Ijzerman 1926:56–58), although this tuber has not yet been identified in the archaeological record of the island.

Quinoa seeds were the most abundant plant remains found at every Mocha Island site (Roa et al. 2021), and quinoa likely comprises a major part of the  $\text{C}_3$  plant residues found in the pots. Its use in southern Chile was reported since the Early Ceramic period (1550–950 cal BP) from wild-type quinoa managed by hunter-gatherer groups (Adán and Mera 2011; Campbell and Quiroz 2015; Roa et al. 2018). Its ability to adapt to different environments without much maintenance investment may have facilitated its early management and incorporation into local foodways (Planella 2019).

The recent arrival of maize on Mocha Island and southern Chile (around 950 BP) would have required a period of experimentation and adaptation to the local temperate conditions. This might have influenced the degree to which the El Vergel farmers exploited maize; it was possibly restricted to the production of fermented beverages. The two pots used for processing maize likely also included  $\text{C}_3$  plants. The different  $\Delta^{13}\text{C}$  offsets between the two samples possibly indicate the mixing of maize with  $\text{C}_3$  plants with varying proportions of  $\text{C}_{16:0}$  and  $\text{C}_{18:0}$  alkanoic acids. Incorporating other starchy plants into the pot may have enhanced the fermentation process for producing maize-based beverages (Arriaza et al. 2016). From the residue data, we can argue that maize was not consumed as a staple food by Mocha island farmers; otherwise, we would expect more samples showing high  $\delta^{13}\text{C}$  values, possibly with a negative  $\Delta^{13}\text{C}$  offset due to the mixing of  $\text{C}_3$  and  $\text{C}_4$  plants, as indicated by our simple linear model. Therefore, we suggest that the high  $\delta^{13}\text{C}$  and relatively low  $\delta^{15}\text{N}$  values for

human bone collagen described in previous studies might not necessarily represent a preferential consumption of maize over aquatic resources. Low trophic-level marine organisms such as mollusks might also have been preferentially consumed, an interpretation supported by the faunal evidence from the island.

## Conclusion

The permanent settlement of Mocha Island by the El Vergel farmers required the translocation of domestic crops and tamed animals. Previous studies referred to the exploitation of marine resources as only a secondary activity and stressed the continuity of farming to support the dense populations on the island. Our organic residue analysis revealed that the El Vergel pottery of Mocha Island was mainly used for exploiting plant commodities and marine products. Plant commodities were the basis of the local diet; however, maize was not a staple and might have been mainly destined for the production of fermented beverages. Plant products were mixed with aquatic resources, although different types of plants were also simultaneously exploited. Marine commodities were a source of animal protein and may have compensated for the limited availability of terrestrial animal products during certain seasons and complemented the exploitation of translocated ruminants.

The use of pottery indicates that local and translocated resources were an important part of the foodways of the El Vergel coastal farmers from Mocha Island. Local marine resources complemented translocated domestic plant commodities such as maize and quinoa, sustaining a mixed economy where farming, coastal foraging, hunting, and fishing were practiced. Perhaps it was only when this mixed economy developed that the dense populations could be supported on Mocha Island following its settlement by the El Vergel farmers. More studies focusing on mainland sites are needed to characterize better the foodways and socioeconomic strategies of the El Vergel farmers inhabiting coastal and inland areas.

**Acknowledgments.** Archaeological potsherds and maize kernels were analyzed under the permit ORD No. 2477 and decree CVE 2292672 issued by the Consejo de Monumentos Nacionales de Chile and the Ministerio de las Culturas, las Artes y el Patrimonio de Chile. We want to thank the Department of Archaeology, University of York, for funding this research. We are grateful for the staff and students from BioArch, University of York, and especially Matthew Von Tersch and Jasmine Lundy for their unconditional support. Professor Guillermo Schmeda-Hirschmann from the Universidad de Talca, Chile, provided valuable guidance on local flora. FONDECYT Project No11150397, “Trayectoria socio-política y complejización en la Araucanía septentrional: El Complejo El Vergel en la región de Angol (1000–1550 d.C.)” led by Dr. Roberto Campbell, supplied the archaeological material. FONDECYT Project No1181829, “Monumentos arqueológicos y memorias materiales: Historias andinas de larga duración en Pampa Iluga, Tarapacá (900 AC–1600 DC),” led by Professor Mauricio Uribe, supplied the archaeobotanical references. We are grateful for the support given by the ERC Consolidator project TRADITION, which has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program under Grant Agreement No. 817911. This article contributes to the “María de Maeztu” program for Units of Excellence of the Spanish Ministry of Science and Innovation (CEX2019-000940-M).

**Funding statement.** This project was funded by the Department of Archaeology, University of York.

**Data Availability Statement.** Potsherds are stored at the Escuela de Antropología, Pontificia Universidad Católica de Chile. For future analyses, permission should be requested from the Consejo de Monumentos Nacionales, Chile. The generated isotopic and molecular data are stored by the corresponding author and at the BioArch database. Access to this database is restricted to the Department of Archaeology, University of York.

**Competing Interests.** The authors declare none.

**Supplemental Material.** For supplemental material accompanying this article, visit <https://doi.org/10.1017/laq.2023.58>.

Supplemental Text 1. Cooking experiments and archaeological maize for  $\delta^{13}\text{C}$  references.

Supplemental Table 1. Mocha Island Potsherds.

Supplemental Table 2. List of Products.

Supplemental Table 3. Summary of the Molecular and Isotopic Data.

Supplemental Table 4. Phytanic acid SRR-Isomer References.

Supplemental Table 5. Modern References for the APAA-C18 E/H Ratio.

Supplemental Table 6. Archaeological Maize from the Iluga Túmulos Site.

Supplemental Table 7.  $\delta^{13}\text{C}$  References.

Supplemental Figure 1. Boxplot for APAA C20/C18 ratio of samples P25-1.13, P25-1.10, P23-2.2, and P5-1.13 (references from Bondetti et al. 2020).

Supplemental Figure 2. Partial chromatogram of Sample P25-1.10 showing APAAs C18, C20, and C22. APAA-C18 isomers are labeled A-I.

Supplemental Figure 3. Partial ion chromatogram (m/z 75) of the total lipid extract (solvent extract) of sample P25-1.5 showing a distribution of fatty alcohols and fatty acids.

## References Cited

- Adán, Leonor, and Rodrigo Mera. 2011. Variabilidad interna en el Alfarero Temprano del Centro-Sur de Chile: El complejo Pitrén en el valle central del Cautín y el sector lacustre andino. *Chungara* 43:3–24.
- Adán, Leonor, Rodrigo Mera, Mauricio Uribe, and Margarita Alvarado. 2005. La tradición cerámica bícroma rojo sobre blanco en la región Sur de Chile: Los estilos decorativos Valdivia y Vergel. In *Actas del XVI Congreso Nacional de Arqueología Chilena*, edited by Mauricio Massone Mezzano, pp. 399–410. Escaparate Ediciones, Concepción, Tomé, Chile.
- Albán, María, Gabriela Palma, and Ayelén Delgado. 2013. Informe de análisis cerámico sitio P5 Isla Mocha, comuna de Lebu, provincia de Arauco, VIII Región del Biobío, Fondecyt 3130515. Report on file at the Agencia Nacional de Investigación y Desarrollo (ANID), Santiago.
- Aldunate, Carlos. 1989. Estadio alfarero en el sur de Chile. In *Culturas de Chile: Prehistoria, desde sus orígenes hasta los albores de la conquista*, edited by Jorge Hidalgo, pp. 329–348. Editorial Andrés Bello, Santiago.
- Alvarado, Margarita. 1997. La tradición de los grandes cántaros: Reflexiones para una estética del “envase.” *Aisthesis* 30:105–124.
- Arriaza, Bernardo, Juan Ogalde, Juan Chacama, Vivien Standen, Luis Huaman, Fiorella Villanueva, Natalia Aravena, Pablo Méndez-Quiros, and Pedro Tapia. 2016. Microscopic Analysis of Botanical Residues from Cerro Esmeralda Burial in Northern Chile: State and Death Ritual Implications. *Interciencia* 41(12):844–850.
- Báez, Pedro. 1997. Crustáceos en excavaciones arqueológicas de Isla Mocha. In *La isla de las palabras rotas*, edited by Daniel Quiroz and Marco Sánchez, pp. 209–214. Dirección de Bibliotecas, Archivos y Museos (DIBAM), Santiago.
- Bahamondes, Francisco, Claudia Silva, and Roberto Campbell. 2006. La Candelaria: Un yacimiento funerario del complejo El Vergel en el curso inferior del río Bío-Bío. *Boletín de la Sociedad Chilena de Arqueología* 39:69–85.
- Bahlburg, Heinrich, and Michaela Spiske. 2015. Styles of Early Diagenesis and the Preservation Potential of Onshore Tsunami Deposits—A Re-Survey of Isla Mocha, Central Chile, Two Years after the February 27, 2010, Maule Tsunami. *Sedimentary Geology* 326:33–44.
- Becker, Cristian. 1997a. Zooarqueología y etnohistoria: Un contraste en Isla Mocha. In *La isla de las palabras rotas*, edited by Daniel Quiroz and Marco Sánchez, pp. 71–86. Dirección de Bibliotecas, Archivos y Museos (DIBAM), Santiago.
- Becker, Cristian. 1997b. Los antiguos mochanos, cómo interactuaron con la fauna que hallaron y llevaron a la isla. In *La isla de las palabras rotas*, edited by Daniel Quiroz and Marco Sánchez, pp. 159–168. Dirección de Bibliotecas, Archivos y Museos (DIBAM), Santiago.
- Beresford-Jones, David, Alexander Pullen, George Chauca, Lauren Cadwallader, Maria García, Isabel Salvatierra, Oliver Whaley, et al. 2018. Refining the Maritime Foundations of Andean Civilization: How Plant Fiber Technology Drove Social Complexity during the Preceramic Period. *Journal of Archaeological Method and Theory* 25(2):393–425.
- Bondetti, Manon, Erin Scott, Blandine Courel, Alexandre Lucquin, Shinya Shoda, Jasmine Lundy, Catalina Labra-Odde, Léa Drieu, and Oliver E. Craig. 2020. Investigating the Formation and Diagnostic Value of  $\omega$ -(o-Alkylphenyl) Alkanoic Acids in Ancient Pottery. *Archaeometry* 63(3):594–608.
- Braje, Todd J., Thomas P. Leppard, Scott M. Fitzpatrick, and Jon M. Erlandson. 2017. Archaeology, Historical Ecology and Anthropogenic Island Ecosystems. *Environmental Conservation* 44(3):286–297.
- Bullock, Dillman. 1970. La cultura Kofkeche. *Boletín de la Sociedad de Biología de Concepción* 43:1–203.
- Campbell, Roberto. 2015. So Near, so Distant: Human Occupation and Colonization Trajectories on the Araucanian Islands (37° 30' S. 7000–800 cal BP [5000 cal BC–1150 cal AD]). *Quaternary International* 373:117–135.
- Campbell, Roberto. 2020. Diferenciación social en Isla Mocha durante el complejo El Vergel (1000–1550 D.C., sur de Chile). In *Iguals pero diferentes: Trayectorias históricas prehispánicas en el cono sur*, edited by Lorena Sanhueza, Andrés Troncoso, and Roberto Campbell, pp. 17–44. Social Ediciones, Santiago.
- Campbell, Roberto, and Marco Pfeiffer. 2017. Early Public Architecture in Southern Chile: Archaeological and Pedological Results from the Mocha Island Mounds-and-Platform Complex. *Latin American Antiquity* 28(4):495–514.
- Campbell, Roberto, and Daniel Quiroz. 2015. Chronological Database for Southern Chile (35°30'–42° S), ~33000 BP to Present: Human Implications and Archaeological Biases. *Quaternary International* 356:39–53.
- Campbell, Roberto, Francisca Santana-Sagredo, Doina Munita, Rodrigo Mera, Mauricio Massone, Pedro Andrade, Marco Sánchez, and Tatiana Márquez. 2020. Diet in Southern Chile (36°–42°S); A Synthesis from the Isotopic Data. *Quaternary International* 548:109–123.
- Colonese, André C., Alexandre Lucquin, Eduardo P. Guedes, Richard Thomas, Julia Best, B. Tyr. Fothergill, Naomi Sykes, et al. 2017. The Identification of Poultry Processing in Archaeological Ceramic Vessels Using in-situ Isotope References for Organic Residue Analysis. *Journal of Archaeological Science* 78:179–192.
- Contreras, Lino, Daniel Quiroz, Marco Sánchez, and Claudia Caballero. 2005. Ceramios, maíces y ranas . . . un campamento El Vergel en las costas de Arauco. In *Actas del XVI Congreso Nacional de Arqueología Chilena*, edited by Mauricio Massone Mezzano, pp. 357–367. Escaparate Ediciones, Concepción, Tomé, Chile.
- Copley, Mark S., Robert Berstan, Stephanie N. Dudd, Gordon Docherty, Anna J. Mukherjee, Vanessa Straker, Sebastian Payne, and Richard P. Evershed. 2003. Direct Chemical Evidence for Widespread Dairying in Prehistoric Britain. *PNAS* 100(4):1524–1529.



- Corr, Lorna T., Judith C. Sealy, Mark C. Horton, and Richard P. Evershed. 2005. A Novel Marine Dietary Indicator Utilising Compound-Specific Bone Collagen Amino Acid  $\delta^{13}\text{C}$  Values of Ancient Humans. *Journal of Archaeological Science* 32(3):321–330.
- Craig, Oliver E., Hayley Saul, Alexandre Lucquin, Yastami Nishida, Karine Taché, Leon Clarke, Anu Thompson, et al. 2013. Earliest Evidence for the Use of Pottery. *Nature* 496(7445):351–354.
- Cramp, Lucy J. E., Jonathan Ethier, Dushka Urem-Kotsou, Clive Bonsall, Dušan Borić, Adina Boroneanț, Richard P. Evershed, et al. 2019. Regional Diversity in Subsistence among Early Farmers in Southeast Europe Revealed by Archaeological Organic Residues. *Proceedings of the Royal Society B* 286(1894):20182347.
- Crowther, Alison, Patrick Faulkner, Mary E. Prendergast, Eréndira M. Quintana Morales, Mark Horton, Edwin Wilmsen, and Anna M. Kotarba-Morley, et al. 2016. Coastal Subsistence, Maritime Trade, and the Colonization of Small Offshore Islands in Eastern African Prehistory. *Journal of Island and Coastal Archaeology* 11(2):211–237.
- Diefendorf, Aaron F., Katherine H. Freeman, Scott L. Wing, and Heather V. Graham. 2011. Production of N-Alkyl Lipids in Living Plants and Implications for the Geologic Past. *Geochimica et Cosmochimica Acta* 75(23):7472–7485.
- Dillehay, Tom. 1990. *Araucanía: Presente y pasado*. Editorial Andrés Bello, Santiago.
- Dillehay, Tom D., Teresa Rosales Tham, Víctor Vázquez, Steve Goodbred, Elizabeth Chamberlain, and Gabino Rodríguez. 2022. Emergent Consilience among Coeval Fishing and Farming Communities of the Middle Holocene on the North Peruvian Coast. *Frontiers of Earth Science in China* 10. <https://doi.org/10.3389/feart.2022.939214>.
- Dolbunova, Ekaterina, Alexandre Lucquin, T. Rowan McLaughlin, Manon Bondetti, Blandine Courel, Ester Oras, Henny Piezonka, et al. 2023. The Transmission of Pottery Technology among Prehistoric European Hunter-Gatherers. *Nature Human Behaviour* 7(2):171–183.
- Dove, Hugh, and Robert W. Mayes. 2006. Protocol for the Analysis of N-Alkanes and Other Plant-Wax Compounds and for Their Use as Markers for Quantifying the Nutrient Supply of Large Mammalian Herbivores. *Nature Protocols* 1(4):1680–1697.
- Dudd, Stephanie N., Martine Regert, and Richard P. Evershed. 1998. Assessing Microbial Lipid Contributions during Laboratory Degradations of Fats and Oils and Pure Triacylglycerols Absorbed in Ceramic Potsherds. *Organic Geochemistry* 29(5–7):1345–1354.
- Dunne, Julie. 2021. Gone to Seed? Early Pottery and Plant Processing in Holocene North Africa. *Quaternary International* 608–609(2022):178–193.
- Dunne, Julie, Anna M. Mercuri, Richard P. Evershed, Silvia Bruni, and Savino di Lernia. 2016. Earliest Direct Evidence of Plant Processing in Prehistoric Saharan Pottery. *Nature Plants* 3(16194):1–6.
- Evershed, Richard P. 2008. Experimental Approaches to the Interpretation of Absorbed Organic Residues in Archaeological Ceramics. *World Archaeology* 40(1):26–47.
- Evershed, Richard P., Stephanie N. Dudd, Mark S. Copley, Robert Berstan, Andrew W. Stott, Hazel Mottram, Stephen A. Buckley, and Zoe Crossman. 2002. Chemistry of Archaeological Animal Fats. *Accounts of Chemical Research* 35(8):660–668.
- Evershed, Richard P., Mark S. Copley, Luke Dickson, and Fabricio A. Hansel. 2008. Experimental Evidence for the Processing of Marine Animal Products and Other Commodities Containing Polyunsaturated Fatty Acids in Pottery Vessels. *Archaeometry* 50(1):101–113.
- Gálvez, Óscar. 1997. Análisis de restos malaco-arqueológicos de la Isla Mocha. In *La isla de las palabras rotas*, edited by Daniel Quiroz and Marco Sánchez, pp. 195–208. Dirección de Bibliotecas, Archivos y Museos (DIBAM), Santiago.
- Gay, Claudio. 2018 [1873]. *Usos y costumbres de los araucanos*. Edited by Diego Milos. Penguin Random House, Santiago.
- Godoy-Aguirre, Carolina. 2018. Elaboración de bebidas fermentadas en un contexto El Vergel de Isla Mocha (1000–1300 d.C.). *Chungara* 50:107–120.
- Gordon, Américo. 1978. Urna y canoa funerarias: Una sepultura doble excavada en Padre las Casas, Provincia de Cautín, IX Región, Chile. *Revista Chilena de Antropología* 1:60–80.
- Hammann, Simon, Lucy J. E. Cramp, Mathilda Whittle, and Richard P. Evershed. 2018. Cholesterol Degradation in Archaeological Pottery Mediated by Fired Clay and Fatty Acid Pro-Oxidants. *Tetrahedron Letters* 59:4401–4404.
- Hansel, Fabricio A., Mark S. Copley, Luiz A. S. Madureira, and Richard P. Evershed. 2004. Thermally Produced  $\omega$ -(o-Alkylphenyl) Alkanoic Acids Provide Evidence for the Processing of Marine Products in Archaeological Pottery Vessels. *Tetrahedron Letters* 45(14):2999–3002.
- Hellevang, Helge, and Per Aagaard. 2015. Constraints on Natural Global Atmospheric CO<sub>2</sub> Fluxes from 1860 to 2010 Using a Simplified Explicit Forward Model. *Scientific Reports* 5:17352.
- Hendy, Jessica, Andre C. Colonese, Ingmar Franz, Ricardo Fernandes, Roman Fischer, David Orton, Alexandre Lucquin, et al. 2018. Ancient Proteins from Ceramic Vessels at Çatalhöyük West Reveal the Hidden Cuisine of Early Farmers. *Nature Communications* 9(1):4064.
- Horsburgh, K. Ann, and Mark D. McCoy. 2017. Dispersal, Isolation, and Interaction in the Islands of Polynesia: A Critical Review of Archaeological and Genetic Evidence. *Diversity* 9(3):37.
- Ijzerman, Jan. 1926. *De Reis om de Wereld door Olivier van Noort, 1598–1601*. Martinus Nijhoff, The Hague, Netherlands.
- King, Charlotte L., Andrew R. Millard, Darren R. Gröcke, Vivien G. Standen, Bernardo T. Arriaza, and Siân E. Halcrow. 2018. Marine Resource Reliance in the Human Populations of the Atacama Desert, Northern Chile—A View from Prehistory. *Quaternary Science Reviews* 182:163–174.
- Knudson, Kelly J., Ann H. Peters, and Elsa Tomasto Cagigao. 2015. Paleodiet in the Paracas Necropolis of Wari Kayan: Carbon and Nitrogen Isotope Analysis of Keratin Samples from the South Coast of Peru. *Journal of Archaeological Science* 55:231–243.
- Lepofsky, Dana, Nicole F. Smith, Nathan Cardinal, John Harper, Mary Morris, Gitla Elroy White, Randy Bouchard, et al. 2015. Ancient Shellfish Mariculture on the Northwest Coast of North America. *American Antiquity* 80(2):236–259.



- Leppard, Thomas P. 2014. Similarity and Diversity in the Prehistoric Colonization of Islands and Coasts by Food-Producing Communities. *Journal of Island and Coastal Archaeology* 9(1):1–15.
- Lequesne, Carlos, Carolina Villagrán, and Rodrigo Villa. 1999. Historia de los bosques relictos de “olivillo” (*Aextoxicon punctatum*) y Mirtáceas de la Isla Mocha, Chile, durante el Holoceno tardío. *Revista Chilena de Historia Natural* 72:31–47.
- López, Manuela. 2017. Integración social a nivel supra doméstico de las comunidades presentes en Isla Mocha durante el período Alfarero Tardío: Una aproximación a partir de los estilos tecnológicos de producción cerámica. Undergraduate thesis, Department of Anthropology, Universidad de Chile, Santiago.
- Lucquin, Alexandre, André C. Colonese, Thomas F. G. Farrell, and Oliver E. Craig. 2016. Utilising Phytanic Acid Diastereomers for the Characterisation of Archaeological Lipid Residues in Pottery Samples. *Tetrahedron Letters* 57(6):703–707.
- Martínez, Ismael. 2013. Informe de análisis zooarqueológico de excavaciones de sondeo, enero 2013, Isla Mocha, Provincia de Arauco, Región del Bío-Bío. Proyecto Fondecyt 3130515. Report on file at the Agencia Nacional de Investigación y Desarrollo (ANID), Santiago.
- Martínez, Ismael. 2014. Informe de análisis zooarqueológico de excavaciones de sondeo 2014, Isla Mocha, Provincia de Arauco, Región del Bío-Bío. Proyecto Fondecyt 3130515. Report on file at the Agencia Nacional de Investigación y Desarrollo (ANID), Santiago.
- Martínez, Ismael. 2015. Informe de análisis zooarqueológico de excavaciones de sondeo 2015, Isla Mocha, Provincia de Arauco, Región del Bío-Bío. Proyecto Fondecyt 3130515. Report on file at the Agencia Nacional de Investigación y Desarrollo (ANID), Santiago.
- McFadden, Clare, Richard Walter, Hallie Buckley, and Marc F. Oxenham. 2021. Temporal Trends in the Colonisation of the Pacific: Palaeodemographic Insights. *Journal of World Prehistory* 34(1):47–73.
- Melnick, Daniel, Marco Sánchez, Helmut P. Echlert, and Verónica Pineda. 2003. Geología estructural de la Isla Mocha, Centro-Sur de Chile (38°30'S, 74°W): Implicancias en la tectónica regional. In *Actas del 10° Congreso Geológico Chileno*, pp. 1–9. Servicio Nacional de Geología y Minería, Santiago.
- Navarro, Ximena, and Carlos Aldunate. 2002. Un contexto funerario de la cultura El Vergel (la Araucanía-Chile). *Gaceta Arqueológica Andina* 26:207–229.
- Nelson, Alan R., and William F. Manley. 1992. Holocene Coseismic and Aseismic Uplift of Isla Mocha, South-Central Chile. *Quaternary International* 15–16:61–76.
- O'Connor, Sue, Mahirta, Shimona Kealy, Clara Boulanger, Tim Maloney, Stuart Hawkins, Michelle C. Langley, et al. 2019. Kisar and the Archaeology of Small Islands in the Wallacean Archipelago. *Journal of Island and Coastal Archaeology* 14(2):198–225.
- Palma, Gabriela. 2013. *Inversión de trabajo en la cerámica El Vergel del sitio P-5 de Isla Mocha*. Report on file at the Departamento de Antropología, Universidad de Chile, Santiago.
- Pearsall, Deborah M., Neil A. Duncan, Karol Chandler-Ezell, Douglas H. Ubelaker, and James A. Zeidler. 2020. Food and Society at Real Alto, an Early Formative Community in Southwest Coastal Ecuador. *Latin American Antiquity* 31(1):122–142.
- Pefaur, Jaime E., and José Yáñez. 1980. Ecología descriptiva de la isla Mocha (Chile), en relación al poblamiento de vertebrados. *Boletín del Museo Nacional de Historia Natural* 37:103–112.
- Pino Ramos, Liudis Leidy, Felipe Jiménez-Aspee, Cristina Theoduloz, Alberto Burgos-Edwards, Raúl Domínguez-Perles, Camille Oger, et al. 2019. Phenolic, Oxylipin and Fatty Acid Profiles of the Chilean Hazelnut (*Gevuina avellana*): Antioxidant Activity and Inhibition of Pro-inflammatory and Metabolic Syndrome-Associated Enzymes. *Food Chemistry* 298:125026.
- Planella, María Teresa. 2019. Quinoa in Pre-Hispanic Central Chile: Contributions from Archaeology and Cultural Processes. *International Journal of Agriculture and Natural Resources* 46(2):69–81.
- Potter, Amiee B., and P. Scott White. 2009. The Mitochondrial DNA Affinities of the Prehistoric People of San Clemente Island: An Analysis of Ancient DNA. *Journal of California and Great Basin Anthropology* 29(2):163–182.
- Power, Ximena. 2013. *Informe arqueomalacológico sitio P5-1, Isla Mocha, VIII Región del Bío Bío*. Proyecto FONDECYT 3130515. Report on file at the Agencia Nacional de Investigación y Desarrollo (ANID), Santiago.
- Prieto, Ximena. 1997. Evolución geomorfológica de Isla Mocha durante el Holoceno. In *La isla de las palabras rotas*, edited by Daniel Quiroz and Marco Sánchez, pp. 87–102. Dirección de bibliotecas, archivos y museos (DIBAM), Santiago.
- Quiroz, Daniel. 2001. Ocupaciones El Vergel en las costas de la Araucanía. In *Actas del IV Congreso Chileno de Antropología*, pp. 1456–1465. Colegio de Antropólogos de Chile, Santiago.
- Quiroz, Daniel, and Marcos Sánchez. 2005. La secuencia Pitren-El Vergel en Isla Mocha: Soluciones de continuidad y discontinuidades culturales. In *Actas del XVI Congreso Nacional de Arqueología Chilena*, edited by Mauricio Massone Mezzano, pp. 369–378. Escaparte Ediciones, Concepción, Tomé, Chile.
- Reber, Eleanor A., and Richard P. Evershed. 2004. Identification of Maize in Absorbed Organic Residues: A Cautionary Tale. *Journal of Archaeological Science* 31:399–410.
- Rebolledo, Sandra. 2013. *Informe ictioarqueológico Proyectos NSF BCS-0956229 y FONDECYT 3130515, Isla Mocha, Provincia de Arauco, Región del Bío-Bío*. Report on file at the Agencia Nacional de Investigación y Desarrollo (ANID), Santiago.
- Reiche, Carlos. 1903. Reseña zoológica de la isla. *Anales del Museo Nacional de Chile* 16–18:46–48.
- Reyes, Omar, César Méndez, Manuel San Román, Carolina Belmar, and Amalia Nuevo-Delaunay. 2022. Biogeographic Barriers in the Circulation and Interaction of Hunter-Gatherer Marine Fishers: The Role of the Taitao Peninsula and the Gulf of Penas (~47°S) in the Differentiation of the Cultural Trajectories of West Patagonia. *Frontiers of Earth Science in China* 10. <https://doi.org/10.3389/feart.2022.946732>.
- Roa, Constanza, Daniela Bustos, Horacio Ramírez, and Roberto Campbell. 2018. Entre la Pampa y el Pacífico Sur: Evaluando la dispersión más austral de cultígenos en el cono sur Americano desde la evidencia arqueobotánica y radiométrica de Isla Mocha y Cueva de los Catalanes (Sur de Chile). *Anales de Arqueología y Etnología* 73(2):189–219.

- Roa, Constanza, Ismael Martínez, Javier Montalvo-Cabrera, Ximena Power, Sandra Rebolledo, André C. Colonese, Daniela Bustos, Francisca Santana-Sagredo, and Roberto Campbell. 2021. Apuntes sobre comidas y dietas en Isla Mocha: Integrando resultados de recursos vegetales, animales, residuos orgánicos e isótopos estables (provincia de Arauco, Chile). *Boletín de la Sociedad Chilena de Arqueología* Número Especial 2021:327–360.
- Roa, Constanza, Claudia Silva, and Roberto Campbell. 2015. El aporte de la Isla Mocha al conocimiento sobre el aprovechamiento de plantas con valor alimenticio en el sur de Chile (1000–1700 D.C). In *Actas del XIX Congreso Nacional de Arqueología Chilena*, edited by Marcela Sepúlveda Retamal, Camila Alday Mamani, Camila Castillo Fuentes, Adrián Oyaneder Rodríguez, pp. 549–559. Andros Impresores, Santiago.
- Rojas, Gloria, and Angélica Cardemil. 1995. Estudios arqueobotánicos en Isla Mocha. *Museos* 20:16–17.
- Romanus, Kerlijne, Jeroen Poblome, Kristin Verbeke, Anja Luybaerts, Jacobs Pierre, Dirk De Vos, and Marc Waelkens. 2007. An Evaluation of Analytical and Interpretative Methodologies for the Extraction and Identification of Lipids Associated with Pottery Sherds from the Site of Sagalassos, Turkey. *Archaeometry* 49(4):729–747.
- Rosales, R. P. Diego. 1877 [1674]. *Historia general de el Reyno de Chile*. Edited by Benjamín Vicuña Mackenna. Imprenta del Mercurio, Valparaíso, Chile.
- Saavedra, Bárbara, Daniel Quiroz, and José Iriarte-Díaz. 2003. Past and Present Small Mammals of Isla Mocha (Chile). *Mammalian Biology* 68(6):365–371.
- Schmeda-Hirschmann, Guillermo. 1995. Madia sativa, a Potential Oil Crop of Central Chile. *Economic Botany* 49(3):257–259.
- Shoda, Shinya, Alexandre Lucquin, Jae ho Ahn, Chul Joo Hwang, and Oliver E. Craig. 2017. Pottery Use by Early Holocene Hunter-Gatherers of the Korean Peninsula Closely Linked with the Exploitation of Marine Resources. *Quaternary Science Reviews* 170:164–173.
- Stothert, Karen E., Dolores R. Piperno, and Thomas C. Andres. 2003. Terminal Pleistocene/Early Holocene Human Adaptation in Coastal Ecuador: The Las Vegas Evidence. *Quaternary International* 109–110:23–43.
- Taché, Karine, Yizchak Jaffe, Oliver E. Craig, Alexandre Lucquin, Jing Zhou, Hui Wang, Shengpeng Jiang, Edward Standall, and Rowan K. Flad. 2021. What Do “Barbarians” Eat? Integrating Ceramic Use-Wear and Residue Analysis in the Study of Food and Society at the Margins of Bronze Age China. *PLoS ONE* 16(4):e0250819.
- Takamiya, Hiroto. 2006. An Unusual Case? Hunter-Gatherer Adaptations to an Island Environment: A Case Study from Okinawa, Japan. *Journal of Island and Coastal Archaeology* 1(1):49–66.
- Terrell, John Edward. 2023. Metaphor and Theory in Island Archaeology. *Journal of Island and Coastal Archaeology* 18(4):682–692.
- Van Meurs, Marijke. 1993. Isla Mocha: Un aporte etnohistórico. In *Actas del XII Congreso Nacional de Arqueología Chilena*, Vol.1, edited by Hans Niemeyer F, pp. 193–197. Dirección de Bibliotecas, Archivos y Museos, Santiago.
- Walworth, Mary. 2014. Eastern Polynesian: The Linguistic Evidence Revisited. *Oceanic Linguistics* 53(2):256–272.
- Westbury, Michael, Stefan Prost, Andrea Seelenfreund, José Miguel Ramírez, Elizabeth A. Matisoo-Smith, and Michael Knapp. 2016. First Complete Mitochondrial Genome Data from Ancient South American Camelids: The Mystery of the Chilihueques from Isla Mocha (Chile). *Scientific Reports* 6:1–7.
- Whelton, Helen L., Simon Hammann, Lucy J. E. Cramp, Julie Dunne, Mélanie Roffet-Salque, and Richard P. Evershed. 2021. A Call for Caution in the Analysis of Lipids and Other Small Biomolecules from Archaeological Contexts. *Journal of Archaeological Science* 132:105397.