



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/212458/>

Version: Published Version

Article:

Arblaster, G., Buckley, D., Barnes, S. et al. (2024) Strabismus surgery for psychosocial reasons—a literature review. *British and Irish Orthoptic Journal*, 20 (1). pp. 107-132. ISSN: 2516-3590

<https://doi.org/10.22599/bioj.352>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Strabismus Surgery for Psychosocial Reasons—A Literature Review

REVIEW

GEMMA ARBLASTER

DAVID BUCKLEY

SARAH BARNES

HELEN DAVIS

*Author affiliations can be found in the back matter of this article

WHITE ROSE
UNIVERSITY PRESS
Universities of Leeds, Sheffield & York

ABSTRACT

Introduction: Strabismus surgery may be undertaken for visual benefit, to improve or eliminate diplopia symptoms, or to restore or improve binocular single vision (BSV). In patients without visual symptoms or expected visual benefit, strabismus surgery may still be undertaken if the presence of strabismus causes the patient psychosocial symptoms. To evaluate strabismus surgery undertaken for psychosocial reasons, evidence of postoperative outcomes in this specific cohort is needed.

Methods: A systematic search of the literature was conducted (1946–2023) to identify evidence where postoperative outcomes were reported for adult patients (age 18 years and above) who had undergone strabismus surgery for psychosocial reasons.

Results: Sixty-nine papers were included in the literature review. Most sources of evidence included patients within heterogeneous cohorts of strabismus surgery outcomes, with a range of symptoms and differing surgical aims.

Discussion: In adults who underwent strabismus surgery for psychosocial reasons, improved postoperative ocular alignment and/or improved health related quality of life (HRQoL) were common. Strabismus surgery outcomes appeared to be measured satisfactorily at three months postoperatively. Additional surgical outcomes, including an expanded field of vision, unexpected BSV, improved binocular summation, improved task performance and improved eye movements have been reported, but not fully investigated. There was a lack of consensus on how postoperative success should be defined and measured. A core outcome set for strabismus has been suggested and there is potential to add to the available evidence by investigating which outcome measures are most relevant to those with strabismus and psychosocial symptoms. There is a growing need for robust evidence in this specific subgroup of patients due to a lack of evidence specifically reporting postoperative outcomes in adults with strabismus and psychosocial symptoms.

CORRESPONDING AUTHOR:

Gemma Arblaster

Division of Ophthalmology and Orthoptics, School of Allied Health Professions, Nursing and Midwifery, University of Sheffield, UK; Orthoptic Department, Sheffield Teaching Hospitals NHS Foundation Trust, UK
g.arblaster@sheffield.ac.uk

KEYWORDS:

strabismus; strabismus surgery; psychosocial; outcomes; quality of life; health related quality of life

TO CITE THIS ARTICLE:

Arblaster, G, Buckley, D, Barnes, S and Davis, H. 2024. Strabismus Surgery for Psychosocial Reasons—A Literature Review. *British and Irish Orthoptic Journal*, 20(1), pp. 107–132. DOI: <https://doi.org/10.22599/bioj.352>

INTRODUCTION

Strabismus affects 4–5% of the population (Beauchamp et al. 2003; Goseki & Ishikawa 2017; Hashemi et al. 2017). The aim of strabismus management is to reduce or eliminate the visual and/or psychosocial symptoms caused by strabismus by realigning the eyes into a straighter position. Postoperative restoration of binocular single vision (BSV) or improved diplopia (or confusion) symptoms are considered functional aims of surgery that give the patient visual benefit. If there are no visual symptoms and no potential BSV was predicted, surgery may still be considered if the strabismus caused psychosocial symptoms (Beauchamp et al. 2003).

‘Psychosocial symptoms’ describe the impact of having strabismus on all aspects of the patient’s life. They include lower health-related quality of life (HRQoL) and quality of life (QoL) (Adams et al. 2016; Durnian et al. 2010; Fieß et al. 2020; McBain et al. 2014b; Sah et al. 2017; Wang et al. 2014) and worse self-reported visual function than other ocular diseases (Chang et al. 2015). Patients with strabismus were 10 times more likely to suffer with clinical depression or anxiety (McBain et al. 2014b). Worse depression was associated with reduced HRQoL (Hatt et al. 2014) and in children, strabismus was linked to anxiety, depression, drinking alcohol (Lin et al. 2014), and mental illness (Hassan et al. 2015; Mohney et al. 2008). Psychosocial symptoms caused by strabismus also include social phobia, social fear, social avoidance (Bez et al. 2009; Xu et al. 2012), and difficulty making eye contact and interacting with people (Ghiasi et al. 2013; Xu et al. 2012), leading to hiding strabismus from others (Ghiasi et al. 2013; Menon et al. 2002), embarrassment, negative self-esteem, and avoiding activities (Ghiasi et al. 2013). Strabismus is also reported to interfere with friendships and relationships (Burke et al. 1997), leading to feeling different, having low self-confidence, poor self-image (Satterfield et al. 1993; Xu et al. 2012), and receiving ridicule throughout life (Satterfield et al. 1993). Patients with strabismus report their psychological symptoms are not affected by diplopia or vision in the poorer eye (Ritchie et al. 2013). Patients without diplopia typically report more psychosocial symptoms (McBain

et al. 2014a) and perceive their strabismus to be more noticeable and severe (Jackson et al. 2006).

Negative perceptions of strabismus have been identified in children as young as five years old (Paysse et al. 2001). Adults with strabismus were perceived negatively by others (Kothari & Joshi 2014; Olitsky et al. 1999), as significantly less intelligent, as worse at communication (Olitsky et al. 1999), as less suitable for promotion (Goff et al., 2006), and as less able in the workplace (Coats et al. 2000; Mojon-Azzi & Mojon 2009). Negative perceptions of strabismus negatively affect employment and dating opportunities (Mojon-Azzi & Mojon 2009; Mojon-Azzi et al. 2008).

Strabismus surgery for psychosocial reasons is considered low cost, relatively low risk (Bradbury & Taylor 2013; Ritchie & Ali 2019), highly cost effective (Beauchamp et al. 2005; Beauchamp et al. 2006; Fujiike et al. 2011) and beneficial for patients (Das et al. 2017; Royal College of Ophthalmologists 2016). Whilst the NHS has not withdrawn funding for strabismus surgery, some areas of England were no longer funding strabismus surgery, unless the patient has visual symptoms (such as diplopia) or proven visual benefit from treatment (such as regaining BSV). There was concern that not enough patient benefit was proven in those without expected functional visual gains from surgery (Bristol North Somerset and South Gloucestershire Clinical Commissioning Group 2019). There was therefore a need to increase the evidence of strabismus surgery outcomes specifically in patients with psychosocial symptoms (Durnian et al. 2011). This literature review aimed to evaluate the evidence of outcomes from strabismus surgery when undertaken for psychosocial reasons.

METHODS

A systematic search of the literature was undertaken. Search terms are shown in Table 1. The following databases and repositories were searched: PubMed, Scopus, Cochrane Library, NICE, PsycINFO, Web of Science, Google Scholar, the British and Irish Orthoptic Journal online and an EndNote database of non- or pre-

| TERMS | BOOLEAN OPERATOR | FILTERS USED |
|---|------------------|--|
| Strabismus, Adult, Surgery, Outcomes | AND | English Humans |
| Thyroid, Graves, Myasthenia, Nerve palsy, Myopia, Fracture, Intermittent, Duane | NOT | All adult age categories From 1/1/1946–31/12/2022 |
| Additional search performed using the MeSH terms: Strabismus AND Surgery including the term AND psychosocial (all fields) | | |
| Additional search performed using the terms: outcome AND functional AND eye alignment AND squint | | |
| Initial search performed 1/11/20 (date range 1946–2020). Search updated 7/8/23 (date range 2020–2023). | | |

Table 1 Literature search terms used.

Medline indexed sources (American Orthoptic Journal, Australian Orthoptic Journal, British Orthoptic Journal, Strabismus, Binocular Vision, Journal of AAPOS, and the Transactions of the International Orthoptic Congress, the International Strabismological Association, and the European Strabismological Association). Of specific interest were the treatment outcomes in strabismus in patients with psychosocial symptoms, but no diplopia and no demonstrable BSV. The search was purposely broadened to include larger strabismus cohorts, where

surgery for psychosocial reasons may have been a subgroup.

RESULTS

The results of the literature search are shown in Figure 1. Sixty-nine papers were included in the literature review reporting the outcomes of strabismus surgery undertaken for psychosocial reasons (see Table 2).




Figure 1 Flow chart illustrating the results of the literature search.

*No automated tools were used, all records were excluded by GA.

**Exclusions due to:

- Strabismus surgery planned for visual benefit (to gain BSV or to eliminate diplopia), or to investigate outcomes in patients with potential BSV (for example prism adaptation to restore BSV prior to strabismus surgery).
- Strabismus secondary to or associated with other aetiologies such as neurogenic palsy, mechanical condition (for example Duane syndrome), high myopia, retinal detachment, orbital fractures, congenital fibrosis of the extraocular muscles, age related distance ET (with diplopia).
- Other strabismus diagnoses reported only (for example acute acquired concomitant esotropia, DVD, double elevator palsy).
- Strabismus surgery outcomes in co-existing ocular pathology (for example glaucoma).
- Strabismus surgery anaesthetic techniques.
- Strabismus surgery but without strabismus outcome data reported or where it was unclear which patients, within a larger cohort, had undergone surgery for psychosocial reasons.
- Strabismus surgery techniques and outcomes following specific vertical muscle procedures for a vertical or torsional deviation (for example Harada-Ito procedure).
- Intermittent strabismus or heterophoria only.
- Paediatric patients only (with the following exceptions: childhood strabismus that had recurred in adulthood and childhood onset strabismus that had received the primary surgical treatment in adulthood).
- Other surgical outcomes (for example refractive surgery outcomes performed in patients with strabismus).
- Treatments for diplopia (with the exception of diplopia resulting from psychosocial strabismus surgery, which was included).
- Slipped extraocular muscles during surgery (for example, description of surgical technique but no reported strabismus outcome).
- Outcomes from Botulinum Toxin (BT) injections.
- Poster abstracts.
- Review papers reporting no original data.
- Editorial articles.

**Exclusions due to:

- Strabismus surgery outcomes reported in a heterogeneous cohort and not possible to extract outcomes in those undergoing strabismus surgeries for psychosocial reasons only.
- Insufficient evidence reported to be able to determine postoperative outcomes of strabismus surgery in those undergoing strabismus surgery for psychosocial reasons.
- Cohort already reported in an earlier study.

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|----------------------------|--|---|--|---|---------------|
| (Adams et al. 2016) | Investigating psychological issues in patients before and after strabismus surgery | All strabismus patients (n = 220) Age 17–88 No diplopia (n = 96) | Clinical assessment of success, partial success or failure using criteria 1 = largest angle of deviation <12PD (for ET, XT and HT), <20PD HoT; 2 = no (or rare) diplopia or visual confusion in primary and reading position; and 3 = no prisms or Bangarter foil occlusion Success = 3/3 criteria met Partial success = 1 or 2/3 criteria met Failure = 0/3 criteria met Psychological questionnaires (QoL: Adult Strabismus quality of life questionnaire AS-20, Mood: Hospital Anxiety & Depression Scale (HADS), Appearance related social anxiety and social avoidance: The Derriford Appearance Scale (DAS24), Beliefs about strabismus: Revised Illness Perception Questionnaire (IPQ-R), Beliefs about strabismus surgery: Treatment Representations Inventory (TRI), Fear of negative evaluation: Fear of Negative Evaluation (FNE), Perceived visibility: 7-point Likert scale from 1 (not at all visible) to 7 (extremely visible), Importance of appearance: The Centre of Appearance Research Salience Scale (CARSAL), Perception of their appearance: The Centre of Appearance Research Valence Scale (CARVAL), Satisfaction with social support: Multidimensional Scale of Perceived Social Support (MSPSS), Expectations about the outcome of surgery: designed by psychology team (ESSQ) Reasons for having surgery: designed by psychology team (RSSQ), Satisfaction with surgery: designed by psychology team | 3 months clinical 3 and 6 months psychological | Prospective |
| (Akbari et al. 2015) | Persian version of AS-20 pre and postoperatively | All types of strabismus N = 112 Age 15–43 years | AS-20 (Persian version) VFQ-25 (Persian version) Diplopia (yes/no) PCT <10PD and ≥10PD | 3 months | Prospective |
| (Alam et al. 2014) | Investigating AS-20 outcomes in those considered surgical success | Concomitant manifest strabismus >15PD (preop) successfully aligned within 10PD orthotropia N = 30 Age 11–34 years | AS-20 | 6 weeks 3 months | Prospective |
| (Aletaha et al. 2016) | Comparison of surgical techniques | Horizontal strabismus N = 54 Age 2–50 years | PCT Number of reoperations | 3 months | Prospective |
| (Alkharashi & Hunter 2017) | Comparison of surgical techniques | All rectus strengthening procedures (resection or plication) N = 72 Age 1–86 years | Success = distance PCT ≤10 PD horizontal deviation and ≤6 PD vertical deviation Reoperation rate Postoperative alignment drift (change from immediate postoperative measurement to final visit measurement) | 6–12 weeks | Retrospective |
| (Al-Wadaani 2017) | Retrospective review of all strabismus surgery | All non-adjustable strabismus surgery N = 96 Age 16–61 years | Improvement in deviation postoperatively | 6–47 months | Retrospective |

(Contd.)

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|---------------------------|--|---|---|-----------------------------------|---------------------------|
| (Ball et al. 1993) | Case series of unexpected stereopsis postoperatively | N = 8 | BSV tests | | Retrospective |
| (Bayramlar & Gunduz 2006) | Review of long term outcome of strabismus surgery in dense amblyopes (6/60 or worse) | N = 33 Age 8–61 years | Krimsky measurement of deviation Success ± 12 PD deviation | 2 months and 24–108 months | Retrospective |
| (Beauchamp et al. 2003) | Review of strabismus outcomes (all patients combined) | All patients who had strabismus surgery (6 centres) N = 299 Age 16 years + | Success alignment = ≤ 8 PD horizontal deviation and ≤ 2 PD vertical deviation Success motor = $\leq +1$ overaction Success sensory = no diplopia | 1 day–19 months | Multicentre retrospective |
| (Berland et al. 1998) | Patients undergoing 8–9mm bilateral LR recession for XT | N = 30 | Abduction limitation Reoperation rate | 3–30 months | Retrospective |
| (Biglan et al. 1994) | Comparison of surgical procedures | All strabismus patients (all aetiologies) N = 24 adjustable N = 113 nonadjustable Mean age 43 and 42 | Success = ± 8 PD horizontal deviation and ± 4 PD vertical deviation % success BSV Correction of diplopia | 1 week and 6 weeks | Retrospective |
| (Bucci et al. 2009) | Horizontal saccades and vergence pre and postoperatively | With and without BSV N = 9 Age 8–20 years | PCT BSV Saccades (measured onset: time to reach 5% of peak velocity, offset: time when velocity reduced to < 10 degrees/sec, gain, mean velocity) Vergence (convergence and divergence) (measured onset: time when velocity reached > 5 degrees/sec, offset: time when velocity reduced to 5 degrees/sec, gain, mean velocity). Saccades combined with vergence | 2 weeks–2 months and 3–10 months | Prospective |
| (Burke et al. 1997) | Psychosocial implications of strabismus and surgery | All had surgery for alignment N = 31 Age 18–68 years | Self-reporting repertory grid – self rating psychosocial issues (pre op and post op) PCT | 3 months | Prospective |

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|-------------------------|--|---|---|---|---|
| (Chang et al. 2017) | Binocular summation in strabismic amblyopia and effect of surgery | N = 15 strabismic amblyopia & Sx N = 30 normal N = 30 strabismus but no amblyopia | VA at 100%, 2.5% and 1.25% contrast (BEO & monocularly) Calculation of BiS Stereopsis PCT | 6–10 weeks | Prospective |
| (Cifuentes et al. 2018) | Outcomes after 3 muscle surgery for large angle horizontal deviations | Consecutive patients having 3 muscle surgery for large angle horizontal strabismus patients N = 28 Age 1–79 years | Motor alignment success criteria: Dist = Primary position 10PD residual deviation – 4PD consecutive deviation and no induced lateral incomitance 5PD between lateral gazes Nr = Primary position 10PD residual deviation – 4PD consecutive deviation Sensory success: improvement in stereopsis of 2 octaves Overcorrection >4PD consecutive deviation Dist & Near (primary position) Undercorrection >10PD deviation Dist & Nr (primary position) | 6 weeks–57 months | Retrospective |
| (Currie et al. 2003) | Outcomes after surgery for large angle XT | Consecutive patients having surgery for large angle XT N = 26 Age 14–68 years | PCT Success criteria Dist = ≤ 10 PD heterotropia or phoria BSV Subjective question – Happy? Yes/No | 8–12 months 18–36 months | Retrospective |
| (Dadeya et al. 2002) | Use of a drug during surgery to reduce restrictions postoperatively | Strabismus patients having a second surgery, +ve FDT but ≤ 25 PD N = 20 Age 6–25 years | PCT FDT score Success criteria Satisfactory = ± 5 PD of orthophoria Undercorrection Overcorrection | 1, 4 and 8 weeks then monthly for 12 months Outcome at 12 months | Prospective RCT |
| (Daga et al. 2022) | Comparison of surgical techniques in XT | N = 80 Intermittent or constant XT Mean age = 23 (range unclear) Two groups of different surgical techniques | PCT Exo drift Slit lamp assessment (ocular surface changes, muscle lump related changes) | 1 day, 1 week, 1 month, 3 months and 6 months | Prospective randomised intervention |
| (Dawson et al. 2013) | Outcomes of strabismus treatment with poor VA (6/24 – PL) | Strabismus treatment outcomes in patients with reduced VA BT n = 11 (n = 2 then Sx) Sx (n = 8 total) N = 17 Age 19–74 years | PCT Comments documented in clinical notes about patient satisfaction postoperatively | 2 weeks | Retrospective |
| (Dotan et al. 2014) | Strabismus surgery in patients with unilateral vision loss and horizontal strabismus | Horizontal strabismus and unilateral VA in worst eye 1.0 or worse, VA in better seeing eye 0.3 or better N = 21 Age 3–64 years | PCT Success ≤ 10 PD horizontal deviation and 1 surgical procedure was required Not success if >10PD or if >1 surgical procedure required | 6–60 months | Retrospective |

(Contd.)

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|------------------------------|--|---|---|--|---------------------------------------|
| (Eino & Kraft 1997) | Adjustable surgery for horizontal deviation | Compared predetermined target angle (after adjustment) to deviation at 6–8 months N = 109 Age 15–72 years | PCT Drift from final alignment to 6–8 month measurement (in PCT and direction) Success if <10PD | Final alignment after adjustment 1–2 weeks 6–8 weeks 6–8 months | Retrospective |
| (Elkamshoushy & Langué 2019) | biLR recession for recurrent XT (prev biMR resect) | Previous biMR resection for XT, but recurrent XT N = 15 Age 20–31 years | PCT OM limitation of ABDuction Success 8PD ET – 10PD XT | 6 months | Retrospective |
| (Estes et al. 2020) | Strabismus surgery, social anxiety and self consciousness | N = 95 >18 years old | Questionnaire to evaluate self-consciousness (private and public) and social anxiety (self-consciousness survey instrument) Pre-op and post-op | 6 months | Prospective |
| (Faridi et al. 2007) | All surgery for primary XT, no previous surgery | Intermittent or constant XT N = 124 Mdn age at surgery 13 years (IQR 6–34 years) | Good motor outcome = \pm 10PD orthotropia (SPCT) BSV | 1–79 months | Retrospective |
| (Fatima et al. 2009) | Report postoperative BSV when none predicted preoperatively | Constant strabismus with no predicted BSV (free space with prisms) N = 15 Age 12–40 years | BSV Success = \leq 10 PD horizontal deviation and \leq 4 PD vertical deviation | 6 weeks | Retrospective |
| (Feliuss et al. 2001) | Re-recession of MR for recurrent ET | N = 115 Age 11 months–77 years | PCT Success ET \leq 10 PD or XT \leq 8 PD OM on versions (underaction of MR) | 4 weeks–8 months Long-term follow-up 8–120 months | Retrospective |
| (Frangouli & Adams 2013) | Amniotic membrane in complex repeat strabismus surgery | Strabismus surgery complicated by fibrosis, range of aetiology N = 8 Age 10–70 years | PCT Objective improvement Subjective improvement in patient symptoms (mainly relating to diplopia, but also includes report of binocular field of vision) Need for further interventions | 9–24 months | Retrospective |
| (Ganesh et al. 2011) | Long-term follow-up of patients who had surgery for childhood ET | Surgery for ET until aligned to 0–10PD ET. Review 32–44 years later N = 85 Age 2–24 at surgery | Initial surgery success = 0–10PD ET Incidence of consecutive XT = \geq 10PD XT Near and Dist Reoperations OM restriction of ADDuction BSV | 32–44 years | Prospective long term follow up study |
| (Ghiasi et al. 2013) | Psychosocial improvement after strabismus surgery | N = 124 Age 15 years+ (71% no diplopia) | Used questionnaires from (Nelson et al, 2008) translated (Iranian population) | 3 months | Prospective |

(Contd.)

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|-------------------------------|--|--|--|--|---------------------------------|
| (Gigante et al. 2018) | 10-year follow-up after monocular surgery for large angle ET | Range of aetiologies of large angle ET N = 36 Age at surgery 4–58 years | PCT Good ≤15PD Fair 16–20PD Poor >20PD Rate of consecutive XT | 6 months 10 years | Prospective long-term follow-up |
| (Glasman et al. 2013) | QoL following strabismus surgery – all patients with complete data | Horizontal and vertical deviations N = 86 Age 17–76 years | PCT AS-20 (total, function subscale and psychosocial subscale) | 12 days–1 year | Prospective |
| (Gusek-Schneider & Boss 2010) | Secondary sensory strabismus surgery outcomes | All patients having surgery for secondary sensory strabismus N = 26 Age 3–45 years | PCT Dist VA BSV Diplopia yes/no Patient satisfaction with surgery yes/no | 3 months Last follow up (1 year 8 m – 13 years 3 m) | Retrospective |
| (Hatt et al. 2010) | HRQoL questionnaires in strabismus surgery | All strabismus, with diplopia (n = 80) and without diplopia (n = 26) N = 106 Age 18–84 years | AS-20 VFQ-25 PCT (SPCT) Success criteria <ol style="list-style-type: none"> 1. no diplopia/visual confusion in primary position or for reading 2. <10PD heterotropia primary position Near or Dist 3. No prism/Bangerter foil/occlusion 4. No symptoms relating to misalignment or strabismus surgery Partial success <ol style="list-style-type: none"> 1. No diplopia/visual confusion in primary position or reading 2. <20PD heterotropia in primary position at Dist and Near 3. No prism/Bangerter foil/occlusion 4. Mild/intermittent symptoms relating misalignment or strabismus surgery (eyestrain/blur/photophobia/suture reaction) Failure <ol style="list-style-type: none"> 1. Diplopia/visual confusion in primary position and reading 2. ≥20PD heterotropia in primary position at Dist or Near 3. Using prism/Bangerter foil/occlusion 4. Moderate/severe symptoms related to misalignment or strabismus surgery | 4–13 weeks | Prospective |

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|---------------------|--|---|--|--|---------------|
| (Hatt et al. 2018) | Identify factors associated with failure of AS-20 scores to improve following strabismus surgery | All strabismus patients – looked at failure to improve on each of the 4 AS-20 domains N = 276 Age 18–91 years | PCT (SPCT) Near 1/3m and Dist 3m AS-20 (4 domains) Diplopia questionnaire Center for Epidemiologic Studies Depression Scale–Revised (CESD-R) (depressive symptoms) Type-D Scale 14 questionnaire (type-Distressed [type-D] personality) | 6 weeks | Prospective |
| (Hatt et al. 2012a) | Changes in HRQoL 1 year after successful strabismus Sx | All strabismus patients included, all aetiologies N = 73 Age 18–88 years | PCT (SPCT & PACT, but SPCT used in criteria) AS-20 Change in AS-20 psychosocial score Change in AS-20 function score Revised diplopia questionnaire Success: no/rare diplopia/visual confusion straight ahead at distance and for reading, <10PD heterotropia in primary position at distance and near Partial success: diplopia/visual confusion ‘sometimes’ or less straight ahead distance and for reading (with or without prism), and <15PD heterotropia Failure: either diplopia/visual confusion was ‘often’ or ‘always’ straight ahead distance or for reading, >15PD heterotropia at distance or near, or the patient was using a Bangerter foil/occlusion | 6 weeks (but between 4–14 weeks) 1 year (but between 5–22 months) | Retrospective |
| (Hatt et al. 2016) | Incorporating HRQoL into the assessment of outcome after strabismus surgery | Assess ‘failures’ by motor and diplopia criteria and evaluate change in HRQoL. Any strabismus type with and without diplopia. All aetiologies. N = 227 Failures (n = 40) Age 18–88 years | PCT (SPCT) Dist 3m and Near 1/3m Diplopia questionnaire AS-20 Motor criteria Diplopia criteria Failure: if 1 of the following criteria was met: (1) SPCT \geq 15 PD (horizontal or vertical) at distance or near; (2) diplopia or visual confusion was present more than “sometimes” straight ahead at distance or for reading (unless atypical diplopia due to decompensated childhood strabismus was present preoperatively, in which case diplopia was allowed postoperatively); (3) occlusive patch/Bangerter foil needed. Partial success: SPCT \leq 15 PD (horizontal and vertical) at distance and near, and diplopia/visual confusion was present never/rarely/sometimes. Correction of diplopia with prism was allowed. Success: if SPCT <10 PD (horizontal and vertical) at distance and near, and diplopia/visual confusion was present never or only rarely. | 1 year (but between 5 months – 2 years) | Prospective |

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|-----------------------|--|--|---|--|---------------|
| (Hertle 1998) | Compare clinical characteristics of strabismus surgery with different onset | Compared strabismus onset before visual maturation (BVM) and after visual maturation (AVM). All surgery and all patients reported. N = 255 Age 14–72 years | PCT BSV Subjective report Success – sensory: restoration of function field of BSV (>20°), regaining central or peripheral fusion, orthotropia or heterophoria in primary position and at near Success – motor: absence of binocular function without diplopia, horizontal alignment <12PD and vertical alignment <5PD in primary position and near Success – subjective: subjective interpretation on improved eye position, binocular function and appearance (including happy/unhappy with eye position, tolerant/intolerant of residual diplopia, happy/unhappy with eye movement) Incomitance = difference ≥8PD | 6 months–5 years | Retrospective |
| (Jackson et al. 2006) | What are the psychosocial benefits of strabismus surgery | All strabismus patients. N = 46 Age 16–61 years (40% diplopia 60% no diplopia) | PCT 1/3m Visual Analogue Scales (VAS) (0–10) for 5 questions on coping, lifestyle, worry, noticeable strabismus, strabismus severity Derriford Appearance Scale (DAS-24) Hospital Anxiety and Depression Scale (HADS) WHOQoLBref (four quality of life domains: physical, psychological, social, and environmental) BSV | 3 months (but between 1–6 months) | Prospective |
| (Ji et al. 2020) | Self-reported sense of deviation in adults successfully aligned with surgery | All deviations N = 91 | PCT EOM BSV AS-20 (Chinese version) Self-report of deviation: no deviation/still have some deviation/still have obvious deviation (some and obvious were classed as self-reported sense of deviation) Success: (>1 year of follow up) no/rare diplopia/visual confusion in primary position and for reading, <10PD horizontal deviation, <5PD vertical deviation at near or dist | Follow up >1 year Last postoperative visit (12–42 months) | Retrospective |
| (Jung & Kim 2018) | Surgical outcomes in sensory XT | Unilateral visual loss and constant horizontal strabismus VA <6/30 (0.7) N = 64 Age 18–71 years | Success = <10PD dist Failure = recurrence or overcorrection Recurrence ≥10PD XT Overcorrection ≥10PD ET | 1 year | Retrospective |
| (Kannam et al. 2021) | Surgical outcomes of horizontal rectus muscle transplantation in recurrent and residual strabismus | Case series N = 7 Age 16–40 years (N = 6 XT, N = 1 ET) | PCT Good alignment (cosmetic success) <12PD Duction limitation | 1 week, 6 weeks and final visit (3–6 months) | Retrospective |

(Contd.)

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|----------------------------|--|---|--|---|-------------------------|
| (Kattan et al. 2016) | Binocular summation and stereoacuity after strabismus surgery | All types of strabismus and surgery N = 130 Age 20–60 years | VA 100% contrast VA reduced contrast 2.5%, 1.25% in dimly lit room Binocular summation Stereoacuity near and dist Diplopia Measures only taken postoperatively | 2 months | Prospective case series |
| (Keskinbora et al. 2011) | Long standing infantile ET – outcomes in late surgery | Alignment and BSV despite late surgery and early onset ET N = 21 Age 8–26 years | PCT BSV <5PD heterotropia = orthotropia Residual ET ≥5PD ET Exotropia ≥5PD XT | 3–9 years | Retrospective |
| (Kim et al. 2008) | Reoperation in sensory strabismus | N = 11 Age 4–33 years | PCT Success = 0–10PD | 1 month Last visit (1–48 months) | Retrospective |
| (Kim et al. 2016) | Self-identity in strabismus and after surgery | N = 351 Age 19 years + 3 groups Strabismus (n = 96) Surgery age 4–15 years (n = 108) No strabismus (n = 147) | Korean self-identity scale (subscales: subjectivity, self-acceptance, future confidence, goal orientation, initiative, and familiarity) | 3 independent groups – not before and after surgery | Retrospective |
| (Kishimoto & Ohtsuki 2012) | VF14 in different ophthalmic conditions | Concomitant and incomitant strabismus N = 625 Age 40–85 years | VF-14 questionnaire PCT BSV (Concomitant group) | 3 months | Prospective |
| (Koc et al. 2013) | Strabismus surgery outcomes – does binocular vision make a difference to QoL | N = 61 Age ≥18 years | AS-20 A&SQ (Amblyopia and strabismus questionnaire) BSV Diplopia score (from A&SQ) Motor success <10PD horizontal deviation and <5PD vertical deviation Sensory results BVP (binocular vision positive) and BVN (binocular vision negative) | 3 months | Prospective |
| (Kushner 1994) | Visual field (binocular or BEO) after surgery for ET | ET Sx N = 37 Age 16–62 years | PCT Binocular VF (BEO) BSV (BG) | 6 weeks | Prospective |

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|--------------------------|---|--|---|---|-------------------------------|
| (Kutschke & Scott 2004) | PAT in ET (childhood onset, but Sx when visually mature) | All types of ET N = 85 Age 9–70 years | Success 0–8PD SPCT at near and dist + peripheral fusion Those with no BSV postoperatively are reported | 6 weeks to 13.7 years | Retrospective |
| (Lee et al. 2013) | Postoperative change in spatial localisation after XT surgery | XT N = 60 Age 4–43 years | PCT Computer touch screen – spatial localisation (pointing errors) | 1 day 1 month | Prospective |
| (Liebermann et al. 2013) | Compare long-term outcomes in reoperation of horizontal strabismus-adjustment vs. no adjustment following surgery | ET and XT With and without potential BSV N = 89 Age 12–83 years | Success: <10PD dist deviation (primary and near), no/rare diplopia (primary and reading), no prism or occlusion Partial success: ≤15PD dist deviation (primary and near) without prism, diplopia none/rare/sometimes in primary and reading, prism allowed, no occlusion Failure: if any of these are met >15PD dist deviation in primary or reading, diplopia always/often in primary and reading, needs occlusion | 6 weeks (but 3–21 weeks) 1 year (but 23 weeks–2 years) | Retrospective |
| (Liebermann et al. 2014) | Improvement in specific function HRQoL concerns after strabismus surgery in nondiplopic adults | N = 20 Age 22–79 years | Same success criteria as Liebermann et al. (2013) AS-20 PCT BSV | 1 year (but 6–19 months) | Retrospective |
| (Lipton & Willshaw 1995) | Comparison of surgery accuracy – specialist centre compared to general | N = 205 Age ? | PCT Success: Grade 1 within 0–5 PD of surgical goal Grade 2 within 6–10 PD of surgical goal Grade 3 >10PD of surgical goal | 6 months | Prospective multicentre study |

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|-----------------------|--|---|---|---|---------------|
| (McBain et al. 2014b) | QoL and mood postoperatively | Range of aetiologies N = 210 Age 17–88 years | PCT (APCT 6m) Self-reports of pain, swelling, scarring, redness 0–10 scale At 3 months: Success: 3 out of 3 criteria met: <12PD ET/ XT/HT <20PD HoT, no/rare diplopia/visual confusion in primary position and reading, no prism/occlusion needed Partial success: 1 of the 3 criteria met Failure: 0 out of 3 criteria met AS-20 Success AS-20: >17.7-point increase in psychosocial subscale and >19.5-point increase in function subscale (>95% LOA) Psychosocial measures: Revised Illness Perception Questionnaire (IPQ-R) Treatment Representations Inventory (TRI) Fear of Negative Evaluation (FNE) scale The Derriford Appearance Scale (DAS24) Perceived Visibility of Strabismus Saliency of Appearance scale (CARSAL) Valence of Appearance scale (CARVAL) Multidimensional Scale of Perceived Social Support (MSPSS) Hospital Anxiety and Depression Scale (HADS) Questionnaires: Reasons for strabismus surgery (RSSQ) Expectations of strabismus surgery (ESSQ) Additional questions: Do you regret having strabismus surgery: Yes definitely 1 – Not at all 4 Would you go through the surgery again: No hesitation at all 1 – Certainly not 4 | 3 months 6 months | Prospective |
| (Menon et al. 2002) | Psychosocial aspects of strabismus | All having surgery for alignment N = 40 Age 15–25 years | Semi-structured interview to complete questionnaire and score questionnaire items (pre-op and post-op) Neuroticism questionnaire | 3 months | Prospective |
| (Murray et al. 2007) | Changes in binocular status after late surgery for infantile ET | N = 17 (if aligned 0–8PD at 1 day post op) | BSV (Worth 4 dot test, BG, Titmus, fusion on Synoptophore) Visual field BEO | Last follow-up N = 6 < 1 month N = 5 < 3 months N = 6 > 1 year | Retrospective |
| (Natung et al. 2022) | Evaluation of surgical dose calculation for horizontal, concomitant strabismus | N = 38 Age 18–47 years N = 19 sensory strabismus | Measurement of deviation only (PCT or Krimsky) Compared correction achieved to correction expected from surgical dose | 3 months | Retrospective |

(Contd.)

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|-----------------------------|---|---|--|------------------------------------|---------------|
| (Nelson et al. 2008) | Psychosocial impact of strabismus and surgery | N = 128 Age ≥ 15 years N = 20 teenagers N = 108 adults | Postoperative telephone interviews to complete questionnaire about psychosocial issues (1–10) and postoperative outcome (1–7) | Unclear | Retrospective |
| (Ozates et al. 2019) | Psychological impact of strabismus surgery | N = 83 Age 14–21 years XT & X(T) | Grouped by constant/manifest deviation XT or X(T) Turkish versions of: Social Appearance Anxiety Scale (SAAS) Depression subscale of the HADS (HAD-D) Brief Fear of Negative Evaluation Scale (BFNE) state anxiety subscale of State-Trait Anxiety Inventory (STAI-S) trait anxiety subscale of State-Trait Anxiety Inventory (STAI-T) | 1 year | Prospective |
| (Pineles et al. 2015) | Binocular summation after strabismus surgery | All strabismus types N = 97 Age 2.5–90 years | VA high contrast (100%) VA low contrast (2.5% and 1.25%) Binocular summation calculation PCT Diplopia Success = 0–10PD horizontal strabismus and 0–4PD vertical strabismus | 6–10 weeks | Prospective |
| (Ribeiro et al. 2014) | QoL in strabismus | N = 101 Age 7–67 years 75% no surgery 25% had surgery | Semi-structured interviews to complete questionnaire (own modified version of AS-20) | ? | Prospective |
| (Sandercoe et al. 2014) | Retrospective review of strabismus surgery | Categorised reasons for surgery (78% for psychosocial reasons) N = 83 Mean age 37 years | PCT BSV Diplopia Objective criteria for success <10PD and acceptable 10–20PD results Subjective criteria = satisfaction with surgical outcome (very satisfied/satisfied/neutral/unsatisfied/very dissatisfied) | Mean 16 weeks | Retrospective |
| (Sefi-Yurdakul et al. 2022) | Comparison of surgical techniques for consecutive XT | N = 49 Age 5–50 years Four groups of different surgical procedures compared | PCT Success <10PD | Last follow up visit (7–17 months) | Retrospective |
| (Sim et al. 2018) | Factors associated with patient perception of success | N = 87 Age 16–83 years 35% had no diplopia | AS-20 (used >95% limits of agreement as evidence of change) Diplopia PCT | 24–126 days | unclear |

| AUTHOR | STUDY PURPOSE | PATIENTS | OUTCOME CRITERIA | TIME POSTOPERATIVE OUTCOME JUDGED | STUDY DESIGN |
|----------------------------|---|---|--|--|---------------|
| (Tibrewal et al. 2021) | Surgical outcomes of primary EOM transplantation for large angle XT | N = 10 Large angle XT ≥ 60 PD Age 2–30 years | Measurement (PCT or modified Krimsky test) Motor success ≤ 10 PD Restorative success ≤ 12 PD ABDuction limitation | 1 week, 6 weeks, final visit (8.6–38.8 months) | Retrospective |
| (Wang & Nelson 2011) | Sm-mod ET surgery outcomes | N = 123 Age 11 months–48 years | Success 0–5PD (PCT near and dist, primary position and lateral gaze) | 6 months Last follow-up (6 months–8 years) | Retrospective |
| (Wortham & Greenwald 1989) | Binocular visual field in ET | N = 10 Age 22–49 years | PCT Visual field BEO BSV | 1–2 months | Retrospective |
| (Xu et al. 2012) | Psychosocial effect of strabismus surgery | N = 56 Age 16–49 years No diplopia pre-op 64% surgery for BSV 36% had surgery for alignment | Own questionnaire (social function and psychological function scores) CT = fair alignment (small manifest deviation) or excellent alignment (no manifest deviation) | 2–3 months | Prospective |
| (Xu et al. 2016) | Long-term follow-up and HRQoL following strabismus surgery | N = 122 Compared AS-20 results to control group without strabismus N = 89 | AS-20 (Chinese version) PCT OM BSV Sense of deviation (no deviation/still have some deviation/still have obvious deviation) Diplopia | Last follow up 12–24 months) | Prospective |

Table 2 Displaying the evidence included in the literature review.

BEO both eyes open, BiS binocular summation, BSV binocular single vision, CT cover test, E esophoria, ET esotropia, HRQoL health related quality of life, OM ocular movements, PCT prism cover test, PD prism dioptres, VA visual acuity, X exophoria, XT exotropia.

DISCUSSION

Evidence of surgical outcomes in adults undergoing strabismus surgery for psychosocial reasons was needed to improve our understanding of the potential risks and benefits of undergoing strabismus surgery. Postoperative outcomes from strabismus surgery were typically reported within a heterogeneous cohort.

DELPHI STUDY AND CORE OUTCOME SETS

A Delphi study attempted to identify areas of consensus and disagreement amongst Ophthalmologists when defining success following strabismus surgery (Serafino et al. 2019). A range of different strabismus types and aetiologies were included, however some of the questions included in the Delphi study were pertinent to adults with strabismus and psychosocial symptoms. There was a lack of consensus reached on the time point at which postoperative outcomes should be evaluated, the deviation size considered successful postoperatively and how the deviation should be measured (simultaneous PCT, alternate PCT or both). Consensus was reached in support of some strabismus conditions having unique outcome criteria (for example, sixth cranial nerve palsy) and for BSV outcomes to be included in the definition of success for some strabismus types. Al Jabri et al. (2019) also highlighted the difficulty in comparing studies reporting strabismus outcomes due to a lack of 'core outcome measures' used. The COMET Initiative ('COMET Initiative: Core Outcome Measures in Effectiveness Trials') aims to encourage core outcome set development and use in clinical trials. A core outcome set is the minimum set of measurements that should be taken and reported in a clinical trial of a specific condition. Core outcome sets are therefore useful as they allow comparison of study results and outcomes across different studies. Al Jabri et al. (2019) identified the outcome measurements most commonly used and reported in amblyopia, strabismus, and ocular motility disorder studies, as well as highlighting that consensus was required to develop core outcome sets for trials and research into these conditions. Of note in the strabismus studies were the most commonly reported core outcome measurements of a near and distance measurement of the deviation, binocularity, HRQoL, and adverse events, with some studies additionally reporting visual acuity (VA) and control of the deviation.

EYE ALIGNMENT

Overwhelmingly the most commonly reported strabismus surgery outcome was the primary position angle of deviation, usually in the distance, measured by the prism cover test (PCT) and reported in prism dioptres (PD). Additionally, stating criteria for 'success' based on the strabismus size postoperatively was common. These had the advantage of allowing comparison between the

percentage successfully aligned with surgery, even when different procedures or techniques were compared. Typically, a target angle considered surgical 'success' was stated and a success rate or percentage achieving success postoperatively was reported. A successful angle was often 0–10PD horizontal deviation (Alkharashi & Hunter 2017), with some specifying 0–5PD (Wang & Nelson 2011), 0–8PD (Beauchamp et al. 2003), or 0–15PD (Gigante et al. 2018). Vertical angles considered successful were 0–2PD (Beauchamp et al. 2003), 0–4PD (Biglan et al. 1994), 0–5PD, 0–6PD (Alkharashi & Hunter 2017), although vertical deviations as large as 12PD hypertropia (HT) and 20PD hypotropia (HoT) were also considered successful (Adams et al. 2016).

Additional factors could be included in the definition of success. For example, a large prospective multicentre study compared outcomes between different centres (specialist or general) and success was graded based on the preoperative surgical aim. Postoperatively success was graded as within 0–5PD (grade 1 success), 6–10PD (grade 2) or greater than 10PD (grade 3) compared to the surgical goal (Lipton & Willshaw 1995). The original angle of deviation may be included, for example Cifuentes et al. (2018) reported success criteria of residual deviation up to 10PD and consecutive deviation up to 4PD, with no induced lateral incomitance after surgery for large angle horizontal strabismus. A difference in the esotropia (ET) and exotropia (XT) angle, depending on the strabismus type or aim of the procedure may also be specified. For example, in a large retrospective study reporting re-recessions for recurrent ET, success was considered to be 0–10PD residual ET or 0–8PD consecutive XT (Felius et al. 2001). Outcome measures relating to the specific surgical procedure may also be included. For example, the amount of abduction limitation was an outcome of bilateral lateral rectus (LR) recessions for recurrent XT (Elkamshoushy & Langué 2019) and incidence of consecutive XT and reoperation rate were outcomes in a long-term follow-up of surgery for childhood-onset ET. Postoperative drift (Alkharashi & Hunter 2017; Eino & Kraft 1997), whether reoperation was required (Aletaha et al. 2016; Alkharashi & Hunter 2017) and complications (Faridi et al. 2007) have also been reported as outcome measures, with some including need for reoperation as failure (Dotan et al. 2014).

DIPLOPIA AND BSV

Surgical procedures for planned visual benefit typically included the aim of surgery as an outcome, for example the percentage achieving BSV or improvement in BSV postoperatively (Cifuentes et al. 2018). Surgery for strabismus and psychosocial symptoms would not typically include visual symptoms as outcomes, unless postoperative BSV (Ball et al. 1993) or diplopia occurred. Gusek-Schneider and Boss (2010) included diplopia (yes/no), PCT, VA, BSV, and patient satisfaction (yes/no) when

reporting postoperative outcomes in secondary sensory strabismus (n = 26). The challenge of different outcome measures for different patients was recognised in a retrospective study that grouped patients by strabismus onset, before or after visual maturation (n = 255) (Hertle 1998). Success criteria were divided into sensory and motor success. Sensory success included restoration of BSV or functional field of BSV. Motor success included orthotropia or heterophoria in primary position and at near. In the absence of BSV and diplopia, motor success included alignment, with a less than 12PD horizontal and less than 5PD vertical deviation considered successful.

DEFINING SUCCESS

Increasingly a range of factors have been included in a definition of success to reflect the view that eye alignment is not the only important outcome measure. Hatt et al. (2010) reported success, partial success, and failure outcomes, although their cohort included patients both with diplopia and BSV, and without. Success included no diplopia or visual confusion in primary position or when reading, less than 10PD heterotropia in primary position at both near and distance, no prism or occlusion, and no symptoms relating to strabismus or strabismus surgery. Partial success included the same criteria, but with a less than 20PD deviation and mild or intermittent symptoms (relating to the strabismus or surgery). Failure included diplopia or visual confusion in primary position and when reading, 20PD heterotropia or larger, using prism or occlusion, and moderate or severe symptoms (relating to strabismus or surgery). Their criteria were later refined to include success as having no or rare diplopia, partial success as less than 15PD with diplopia sometimes, with and without a prism, and failure as greater than 15PD heterotropia and diplopia often or always at distance or reading (Hatt et al. 2012a, 2016; Liebermann et al. 2013, 2014). In a large prospective cohort study (n = 210), patients with all types of strabismus were recruited to a study investigating QoL and mood, before and after strabismus surgery. Deviation size less than 12PD ET, XT, or HT and less than 20PD HoT, no or rare diplopia in primary position and reading, and no prism or occlusion needed were used as the criteria determining success, partial success, and failure. Success required all three criteria, partial success required one of three criteria and failure required none of the criteria were met (Adams et al. 2016; McBain et al. 2016).

PATIENT PERCEPTION OF THE POSTOPERATIVE OUTCOME

Success from the patient's perspective may be different to the clinician's perspective. In recognition of this, some studies included objective and subjective outcomes postoperatively (Frangouli & Adams 2013) or asked patients to report their eye alignment, binocular

function, and appearance subjectively (happy/unhappy) (Hertle 1998). In a retrospective study (n = 83) 78% underwent surgery for psychosocial reasons (without diplopia) and both objective and subjective success criteria were used to report the outcomes. Eighty-three percent of all patients had a successful outcome, both objectively (deviation less than 10PD) and subjectively (very satisfied) (Sandercoe et al. 2014).

QUESTIONNAIRES

Increasingly, questionnaires for patients to self-report visual function, QoL, HRQoL, and patient reported outcome measures (PROMs), both generic and those developed specifically for strabismus, have been used pre- and postoperatively (Hatt et al. 2016). Using telephone interviews to complete questionnaires postoperatively (n = 128), patients reported satisfactory eye position (98%) and improved self-esteem (85%), abilities to meet new people (65%), interpersonal relationships (27%), and abilities to try new activities (16%). Younger patients reported greater improvements postoperatively and a larger preoperative deviation was associated with greater improvements in self-esteem and self-image postoperatively (Nelson et al. 2008). Interviews have been used to complete questionnaires rather than explore patient perceptions of postoperative outcome (Menon et al. 2002; Ribeiro et al. 2014). Menon et al. (2002) reported 97.5% of their cohort (n = 40) had improved appearance, relationships with others, self-esteem, and self-confidence postoperatively. Postoperatively 37.5% changed future plans, and 95% reported trying new activities or things that had previously been avoided.

Ghiasi et al. (2013) used a similar questionnaire to Nelson et al. (2008) to prospectively evaluate changes three months after strabismus surgery. All aspects of the questionnaire were reported as improved postoperatively. A high percentage of patients reported improved self-esteem (89%), improved relationships (82%), being able to meet new people (79%), and being better at their job or work (76%) postoperatively. A smaller percentage of patients also reported having improved chances of employment (53%) and being able to try new activities (36%) postoperatively. Gender and direction of strabismus did not affect the results.

Burke et al. (1997) asked patients (n = 31) seeking surgery for alignment only to complete questions about psychosocial issues, rating themselves on a five-point scale preoperatively and three months postoperatively. Patients reported significantly improved psychosocial functioning postoperatively. However, they also reported less than 'ideal world' results and that others would rate them less highly than they rated themselves postoperatively. Age did not affect the results, but females and ETs reported greater improvements in psychosocial functioning compared to males and XTs. Greater

improvements in HRQoL in females postoperatively has also been reported using the AS-20 (Akbari et al. 2015; Alam et al. 2014; Glasman et al. 2013).

Xu et al. (2012) used their own questionnaire to investigate social and psychological effects of strabismus and surgical correction. None of the cohort ($n = 56$) had diplopia preoperatively and 36% had surgery for alignment only (psychosocial reasons). The most common postoperative outcomes (and the percentage of respondents reporting that outcome) were change in appearance (96%), change in self-esteem or self-confidence (96%), change in relationships with friends (91%), trying activities previously avoided (82%), and changing plans for the future (68%). However, it is unclear which outcomes were gained by those having surgery for alignment only.

VISUAL FUNCTION

The VFQ-25 questionnaire is used to measure self-reported visual function and the AS-20 questionnaire is reported to measure HRQoL. Visual functioning questionnaires have measured improved visual function after strabismus surgery (VF-14) (Kishimoto & Ohtsuki 2012). The VFQ-25 was compared to the AS-20 in a prospective study ($n = 106$). In those without diplopia ($n = 26$), the AS-20 was better able to discriminate between surgical success (total or partial) and failure than the VFQ-25; however, VFQ-25 scores did improve. In those without diplopia, successful outcomes had significantly higher VFQ-25 scores (composite score, all vision-specific subscales, driving subscale, and colour vision subscale) (Hatt et al. 2010). Akbari et al. (2015) reported good correlation between the AS-20 and VFQ-25 (Persian versions) but did not analyse their results based on surgical success. Jackson et al. (2006) used visual analogue scales to report coping, lifestyle, worry, noticeable strabismus, and strabismus severity on a 0–10 scale, as well as the DAS-24, HADS and the WHOQoL-Bref. Strabismus surgery ($n = 46$) resulted in significant improvements in QoL, psychological and physical functioning, which were greater in those without diplopia.

AS-20

The AS-20 (Hatt et al. 2009) has become the most commonly used HRQoL questionnaire in strabismus (Adams et al. 2016; Alam et al. 2014; Glasman et al. 2013; Hatt et al. 2010; Hatt et al. 2012a, 2016; Hatt et al. 2018; Ji et al. 2020; Koc et al. 2013; Liebermann et al. 2014; McBain et al. 2016; Sim et al. 2018). Despite not being specific to strabismus with psychosocial symptoms, surgery in these patients has improved both psychosocial and functional aspects of the AS-20 (Alam et al. 2014; Hatt et al. 2010; Hatt et al. 2012a; Koc et al. 2013). Liebermann et al. (2014) reported all AS-20 functional elements improved postoperatively in patients without diplopia ($n = 20$), with the greatest

improvements in stress, worry, needing to take breaks, enjoying hobbies, depth perception, and eye strain items. Significant improvements in self-reported visual function after strabismus surgery for psychosocial reasons were difficult to explain, as no visual change was measured using standard clinical vision tests. However, it is possible that a change in binocular field of vision may have occurred as this was not tested (Kushner 1994; Wortham & Greenwald 1989).

The AS-20 and A&SQ were used in a prospective study of adult strabismus surgery outcomes ($n = 61$) (Koc et al. 2013). Both questionnaires measured significant improvements in HRQoL three months postoperatively. Those with BSV postoperatively had significantly greater improvements in HRQoL scores on the functional subscales than those without BSV, but only when amblyopes were removed from the analysis. The change in overall scores and psychosocial scores (using both questionnaires) were not significantly different between those with and without BSV, highlighting that visual benefit postoperatively was not required for improvement in HRQoL.

Alam et al. (2014) used the AS-20 in a cohort of older children and adults undergoing first strabismus surgery ($n = 30$). None had diplopia, but it is unclear whether any had BSV. Significant improvements in AS-20 HRQoL were measured six weeks and three months postoperatively, with a greater improvement in females. Glasman et al. (2013) reported larger improvements in HRQoL (AS-20) in females, those with larger changes of the deviation and those with smaller strabismus postoperatively. Their prospective study of 17- to 76-year-olds ($n = 86$) found surgery led to improvements in all aspects of the AS-20; however, BSV and diplopia were not reported. Their cohort may therefore have included some surgery for visual benefit.

Adams et al. (2016) used the AS-20, as well as a large battery of QoL and psychosocial measures in a prospective study of patients aged 17–88 years ($n = 210$). A range of aetiologies of strabismus were included and it is unclear how many had surgery for psychosocial reasons; however, 44% had no diplopia. Postoperatively there was a reduction in the number of patients reporting poor AS-20 HRQoL, from 85% to 68% at three and six months postoperatively. Other measures of social anxiety and avoidance, clinical anxiety, and depression also improved significantly. In a study reporting the same cohort ($n = 210$) McBain et al. (2016) used the AS-20 as the primary outcome measure. Strabismus surgery resulted in significantly improved HRQoL three months postoperatively, with no further improvements at six months. Improvements in HRQoL were not associated with clinical judgements of success, highlighting that clinical definitions of success may not adequately capture the postoperative result from the patient's perspective. Postoperatively there were improvements

in a wide range of psychosocial domains, as well as all aspects of the AS-20 (McBain et al. 2016). Using a questionnaire to evaluate self-consciousness, Estes et al. (2020) reported improved public (but not private) self-consciousness and improved social anxiety six months postoperatively. It was unclear how many of their cohort (n = 95) had surgery for psychosocial reasons, as some had diplopia (66%) and depth perception (62%) preoperatively. Using a range of psychological measures Ozates et al. (2019) demonstrated that constant XT (and no BSV) was significantly worse than intermittent XT and BSV. Surgery resulted in significant improvements in all psychological measures for the constant XT group, to the extent that there was no difference between constant and intermittent XT postoperatively.

Patients without diplopia reported significantly lower AS-20 psychosocial subscale scores preoperatively compared to those with diplopia. Interestingly, AS-20 function subscale scores were not significantly different. Postoperatively psychosocial and function subscale scores improved in all patients. Although the improvement in psychosocial subscale score was higher in those without diplopia, they continued to report lower postoperative psychosocial subscale scores than those with diplopia initially. The only factor predictive of a greater improvement in AS-20 HRQoL was socioeconomic status. Those from a more deprived area had a higher rate of success postoperatively (Sim et al. 2018)

Hatt et al. (2010) reported strabismus patients without diplopia gained significant improvements in AS-20 HRQoL, particularly if they had a successful result. Even those with 'failure' postoperatively reported AS-20 improvements, leading Hatt et al. (2016) to suggest success should include HRQoL improvements (beyond test-retest variability), in addition to improved alignment and diplopia. Having a distressed personality type, worse diplopia or depressive symptoms postoperatively, and coexisting facial abnormalities were associated with postoperatively failure, using the AS-20 as the outcome measure (Hatt et al. 2018). These results highlight that mental health as well as clinical factors influence the outcomes from strabismus surgery, a view shared by others (Adams et al. 2016; McBain et al. 2016). Hatt et al. (2012a) retrospectively reported outcomes in adults between 5–22 months postoperatively (n = 73), described as one-year results. Those who continued to meet success criteria (less than 10PD alignment and no or rare symptoms) maintained improved AS-20 results at one year compared to six weeks postoperatively. From six weeks to one year, those without diplopia showed stable function subscale results and further improved psychosocial subscale results. Ji et al. (2020) used the AS-20 (Chinese version) to investigate successful outcomes one year postoperatively, using similar success criteria to Hatt et al. (2016). Patients with BSV and diplopia were included. Despite successful strabismus surgery, 24% of

their cohort (n = 91) still reported they had strabismus. Those who perceived a deviation postoperatively reported lower AS-20 scores and were more likely to have a larger vertical deviation (Ji et al. 2020).

Whilst motor outcomes (strabismus size) may be more likely to define surgery as successful, the method of AS-20 analysis has been shown to affect the results (Leske et al. 2010). Change in either AS-20 subscale, greater than 95% limits of agreement, was considered difficult to achieve (Hatt et al. 2012b), but relying on motor outcomes only may fail to capture improved symptoms or HRQoL (Hatt et al., 2016). The AS-20 was considered to have excellent test-retest variability and a low chance of a ceiling effect. A change in overall score of 14, psychosocial subscale score of 17.7, and function subscale of 19.5 were considered evidence of real change. Whilst different results were provided for those with and without diplopia (Leske et al. 2010), it was unclear whether the 'without diplopia' subgroup was strabismus with psychosocial symptoms only, as it may have included strabismus with BSV. A later evaluation of the AS-20 using Rasch analysis suggested refining the questions, the response options and the subscales to increase responsiveness to change in QoL. This resulted in removal of two questions from the previous function subscale (Leske et al. 2012).

Surgery has been reported to improve and normalise symptoms of anxiety and depression, HRQoL, daily functioning, and psychological adjustment postoperatively (Jackson et al. 2006); however, others report improved but not normalised HRQoL (Xu et al. 2016). Patients who perceived they had no strabismus postoperatively achieved greater HRQoL improvement (Xu et al. 2016). Kim et al. (2016) used a self-identity questionnaire to evaluate young adult males at a military service examination. Having strabismus negatively affected self-identity compared to those with no strabismus and those who had previously undergone strabismus surgery in childhood. There was no difference in self-identity between those who had previous strabismus surgery and those without strabismus. With the recognition that QoL is an important outcome from strabismus surgery, focus has shifted to consider whether psychosocial interventions preoperatively could improve QoL and psychosocial outcomes. No trials have yet been undertaken and this is an area for future research (Mackenzie et al. 2016).

TIMING OF POSTOPERATIVE OUTCOME

Clinical care of patients following strabismus surgery varied among different clinicians, hospitals, healthcare systems, and countries. Patients may be discharged at a specific time point if they are asymptomatic and happy with the surgical result, yet others may be kept under longer review. Strabismus surgery outcomes were reported at one week (Berland et al. 1998), two weeks (Dawson et al. 2013),

one month (Kim et al. 2008), six weeks (Alam et al. 2014; Fatima et al. 2009), three months (Adams et al. 2016; Alam et al. 2014), six months (Adams et al. 2016; Lipton & Willshaw 1995), one year (Currie et al. 2003; Dadeya et al. 2002; Jung & Kim 2018) and later than one year (Currie et al. 2003; Felius et al. 2001). In some studies, the time at which outcome is being reported was unclear (Nelson et al. 2008). Reporting one-year postoperative outcomes had the advantage of providing longer-term data, yet many patients had been discharged and less data available for analysis (Liebermann et al. 2013). Longer-term postoperative outcomes may therefore be biased and include a greater proportion of poorer outcomes that have not been discharged.

The last available follow-up (Al-Wadaani 2017; Beauchamp et al. 2003; Berland et al. 1998; Faridi et al. 2007) was commonly used to report postoperative outcomes, but this was also variable. Kim et al. (2008) reported postoperative outcomes following reoperation for sensory strabismus one month postoperatively and at the final postoperative visit, which ranged from 1–48 months. In contrast, the last available follow up visit ranged from six weeks to 13 years in a study of later surgery for childhood onset ET (Kutschke & Scott, 2004). Specific longer-term studies reporting outcomes after more than one year were less common but offered a unique view of postoperative stability and change over time. For example, 2–9-year follow-up (Bayramlar & Gunduz 2006), 3–9-year follow-up (Keskinbora et al. 2011) and 10-year follow-up (Gigante et al. 2018) have been reported. A unique Swedish prospective study invited adults who had surgery for childhood ET for review and reported 32–44-year follow-ups (Ganesh et al. 2011).

On balance, evaluation of strabismus surgery outcomes at, or later than, three months represented a useful and achievable time point, unless measuring longer term outcomes was the specific aim. For most patients this was thought to allow sufficient time for healing (Escardó-Paton & Harrad 2009), for eye alignment to stabilise and for the patient to adapt to their eye position. Measuring QoL outcomes at six months postoperatively was not significantly different to three months (McBain et al. 2016).

ADDITIONAL OUTCOMES FROM STRABISMUS SURGERY

Patients undergoing strabismus surgery for psychosocial reasons may achieve more than just psychosocial benefit, as shown by QoL or HRQoL improvements. Observational studies reporting additional postoperative changes are discussed in detail below.

Visual field

Patients have gained an enlarged peripheral visual field following surgery to reduce ET (Kushner 1994; Murray et al. 2007; Wortham & Greenwald 1989). Wortham and Greenwald (1989) reported ten patients with ET who

postoperatively gained peripheral visual field, gaining a mean 16 degrees horizontally (range 5–30 degrees). Visual field size was measured using the Goldmann perimeter, I4e target. The gain in peripheral visual field was ipsilateral to the strabismic eye and occurred even in the presence of amblyopia ($n = 3$). Three patients gained some stereopsis postoperatively (range 80" of arc to Titmus fly). This suggested in patients without BSV the suppressed eye contributed to the peripheral field of vision. It also suggested that aligning the strabismic changed the amount, or extent, it contributed to the peripheral visual field. Anecdotally four patients reported visual improvement; however, patients were not asked to subjectively report visual change postoperatively. No follow up data were presented and comparisons with other patients were lacking. Murray et al. (2007) reported older children and adults ($n = 17$) with untreated infantile ET gained an expanded field of binocular vision (mean 32 degrees) postoperatively. However, in contrast to Wortham and Greenwald (1989), sensory fusion was always achieved in addition to binocular field expansion (Murray et al. 2007). Kushner (1994) reported that 34 of 35 patients (age 16–62 years) gained an expanded field of binocular vision postoperatively. The patient that did not gain field of binocular vision ($n = 1$) had unilateral poor vision and retinal abnormalities secondary to uveitis, which may have affected the postoperative outcome. Of those who gained field of binocular vision ($n = 34$), 29 had sensory fusion and 5 had suppression postoperatively (Bagolini glasses). Most patients that gained field of vision postoperatively were aware they had improved peripheral vision.

Unexpected binocular vision

Despite surgery for planned psychosocial benefit, unexpected BSV may occur postoperatively. For example, patients with longstanding large angle strabismus ($n = 8$) have achieved good stereopsis, mean 45" of arc (Titmus) (Ball et al. 1993). Eight patients (out of 20) achieved 60–400" of arc (Frisby Near Stereotest (FNS)) or 40–80" of arc (Frisby Davis distance stereotest (FD2)) one year postoperatively (Liebermann et al. 2014). Detailed reports of pre- and postoperative investigations of BSV in patients with strabismus are lacking. Retrospective studies aiming to identify factors that predict BSV postoperatively can lack complete outcome data (Umazume et al. 1997), leading to difficulty providing data on the proportion of patients who may achieve unexpected BSV or factors that may predict BSV postoperatively. These factors highlight the importance of assessing potential BSV preoperatively (Ball et al. 1993) and BSV outcomes postoperatively, even when it is assumed no BSV is possible (Murray et al. 2007).

Binocular summation

Strabismus surgery has been reported to improve binocular summation, with a greater effect measured using lower

contrast (1.25%) acuity charts. This improvement can mean binocular summation is measured postoperatively, despite binocular inhibition preoperatively. Successful surgical alignment and later onset strabismus have both been associated with greater improvements in binocular summation postoperatively (Pineles et al. 2015). Yet, other studies have shown highly variable changes in binocular summation following strabismus surgery (Chang et al. 2017). Interpreting postoperative binocular summation data only, rather than change as a result of surgery, and interpretation of binocular summation data in isolation, rather than as part of an investigation of pre and postoperative BSV may also be misleading. Further evidence is required to establish whether binocular summation improves following all strabismus surgery, or whether BSV and stereopsis (pre- and postoperatively) affect the binocular summation outcome (Kattan et al. 2016).

Task performance

Patients have reported improved ability to perform daily activities (Nelson et al. 2008) and being able to work better (Ghiasi et al. 2013) when completing questionnaires postoperatively. Improved AS-20 function subscale results have been measured postoperatively even though patients have undergone surgery specifically for psychosocial symptoms or had no measurable visual change postoperatively (Alam et al. 2014; Hatt et al. 2010; Hatt et al. 2012a; Koc et al. 2013; Liebermann et al. 2014). Few studies have measured task performance before and after strabismus surgery. Lee et al. (2013) used a spatial localisation pointing task presented on a touch screen to measure pointing accuracy in patients pre- and post-XT surgery. Pointing accuracy was reduced one day postoperatively, but accuracy improved to preoperative levels at one month postoperatively (Lee et al. 2013).

Eye movements

Using a photoelectric eye tracker, Bucci et al. (2009) measured the accuracy and mean velocity of saccades, convergence and divergence, and combined saccades and vergence eye movements, pre- and postoperatively. Nine subjects (children and adults) with strabismus were included, six with no BSV pre- and postoperatively, although diplopia was not mentioned. Preoperatively, compared to normative data, accuracy was reduced for vergences and combined saccades and vergence; and mean velocity was reduced for saccades and convergence. Postoperatively, accuracy improved for saccades (at near), vergences and combined saccades and vergence; and mean velocity improved for convergence and combined saccades and divergence.

Limitations

Some studies included a heterogeneous cohort and a wide range of patient ages. It is possible this may have

introduced bias or variability in the interpretation of surgical outcome, particularly in studies where both children and adults were reported.

CONCLUSION

Most of the evidence describing the outcomes of strabismus surgery in patients without visual symptoms reported improved postoperative ocular alignment and/or improved HRQoL. Yet, QoL and HRQoL measures were not used consistently, and different questionnaires were used. None of the questionnaires were exclusively for strabismus with psychosocial symptoms; however, the AS-20 was developed for adults with strabismus and was the most commonly used HRQoL questionnaire and PROM. There were variable reports of the outcomes of surgery, the time at which outcomes are measured and a lack of consensus on how success should be defined and measured after strabismus surgery. Strabismus surgery outcomes appeared to be measured satisfactorily at, or around, three months postoperatively. However, there is acknowledgement that the postoperative outcome at three months may differ from the longer-term outcome. Additional surgical outcomes, including an expanded field of vision, unexpected BSV, improved binocular summation, improved task performance and improved eye movements have been suggested, but have not been fully investigated. A core outcome set for strabismus has been suggested and there is potential to add to the available evidence by investigating which outcome measures are most relevant to those with strabismus and psychosocial symptoms. Criteria for 'success, partial success, and failure' have been used by several studies attempting to categorise and compare surgical outcomes. However, there is the potential to improve these categories, as patients categorised as a failure postoperatively can still report significant improvements postoperatively.

Overall, there was a lack of evidence specifically reporting the outcomes of strabismus surgery in adults with psychosocial symptoms. Large heterogeneous cohorts of strabismus patients were often reported, typically with a range of symptoms and differing surgical aims. There is a growing need for robust evidence in this specific subgroup of patients with strabismus and psychosocial symptoms.

FUNDING INFORMATION

Gemma Arblaster was funded by an NIHR Clinical Doctoral Research Fellowship (ICA-CDRF-2016-02-063). The views expressed are those of the authors and not necessarily those of NIHR or the Department of Health and Social Care.

COMPETING INTERESTS

At the time of submission, Gemma Arblaster was the Editor of the *British and Irish Orthoptic Journal*. She had no role in the peer review process of this article.

AUTHOR AFFILIATIONS

Gemma Arblaster  orcid.org/0000-0002-3656-3740

Division of Ophthalmology and Orthoptics, School of Allied Health Professions, Nursing and Midwifery, University of Sheffield, UK; Orthoptic Department, Sheffield Teaching Hospitals NHS Foundation Trust, UK

David Buckley  orcid.org/0000-0001-9140-8543

Division of Ophthalmology and Orthoptics, School of Allied Health Professions, Nursing and Midwifery, University of Sheffield, UK

Sarah Barnes  orcid.org/0000-0003-3279-6368

School of Medicine and Population Health, University of Sheffield, UK

Helen Davis  orcid.org/0000-0002-4138-5969

Division of Ophthalmology and Orthoptics, School of Allied Health Professions, Nursing and Midwifery, University of Sheffield, UK

REFERENCES

- Adams, GG, McBain, H, MacKenzie, K, Hancox, J, Ezra, DG, and Newman, SP. 2016. Is strabismus the only problem? Psychological issues surrounding strabismus surgery. *J AAPOS*, 20(5): 383–386. DOI: <https://doi.org/10.1016/j.jaapos.2016.07.221>
- Akbari, MR, Masoomian, B, Ameri, A, Keshtkar Jaafari, A and Mirmohammadsadeghi, A. 2015. Development of a Persian version of the adult strabismus questionnaire and evaluating the effect of strabismus surgery on health-related quality of life. *Strabismus*, 23(2): 66–72. DOI: <https://doi.org/10.3109/09273972.2015.1025986>
- Alam, D, Khan, AA, Bani, SA, Sharma, R and Amitava, AK. 2014. Gain beyond cosmesis: Demonstration of psychosocial and functional gains following successful strabismus surgery using the adult strabismus questionnaire adult strabismus 20. *Indian J Ophthalmol*, 62(7): 799–803. DOI: <https://doi.org/10.4103/0301-4738.138620>
- Aletaha, M, Bagheri, A, Gholipour, HM and Kheiri, B. 2016. Effect of limited tenon capsule and intermuscular membranes dissection on the outcome of surgery in patients with horizontal strabismus. *Strabismus*, 24(1): 12–15. DOI: <https://doi.org/10.3109/09273972.2015.1130065>
- Al Jabri, S, Kirkham, J and Rowe, FJ. 2019. Development of a core outcome set for amblyopia, strabismus and ocular motility disorders: A review to identify outcome measures. *BMC Ophthalmol*, 19(1): 47. DOI: <https://doi.org/10.1186/s12886-019-1055-8>
- Alkharashi, M and Hunter, DG. 2017. Reduced surgical success rate of rectus muscle plication compared to resection. *J AAPOS*, 21(3): 201–204. DOI: <https://doi.org/10.1016/j.jaapos.2017.05.012>
- Al-Wadaani, FA. 2017. Outcome of strabismus surgery by nonadjustable suture among adults attending a university hospital of Saudi Arabia. *Niger J Clin Pract*, 20(3): 335–340. DOI: <https://doi.org/10.3109/09273972.2015.1025986>
- Ball, A, Drummond, GT and Pearce, WG. 1993. Unexpected stereoacuity following surgical correction of long-standing horizontal strabismus. *Can J Ophthalmol*, 28(5): 217–220.
- Bayramlar, H and Gunduz, A. 2006. Medium term outcomes of strabismus surgery in patients with monocular dense amblyopia. *Saudi Med J*, 27(2): 276–278.
- Beauchamp, CL, Beauchamp, GR, Stager Sr, DR, Brown, MM, Brown, GC and Felius, J. 2006. The cost utility of strabismus surgery in adults. *J AAPOS*, 10(5): 394–399. DOI: <https://doi.org/10.1016/j.jaapos.2006.06.004>
- Beauchamp, GR, Black, BC, Coats, DK, Enzenauer, RW, Hutchinson, AK, Saunders, RA, ... Felius, J. 2003. The management of strabismus in adults-I. Clinical characteristics and treatment. *J AAPOS*, 7(4): 233–240. DOI: [https://doi.org/10.1016/S1091-8531\(03\)00112-5](https://doi.org/10.1016/S1091-8531(03)00112-5)
- Beauchamp, GR, Felius, J, Stager, DR and Beauchamp, C. 2005. The utility of strabismus in adults. *Trans Am Ophthalmol Soc*, 103: 164–172.
- Berland, JE, Wilson, ME and Saunders, RB. 1998. Results of large (8–9 mm) bilateral lateral rectus muscle recessions for exotropia. *Binocul Vis Strabismus Q*, 13(2): 97–104.
- Bez, Y, Coskun, E, Erol, K, Cingu, AK, Eren, Z, Topcuoglu, V and Ozerterk, Y. 2009. Adult strabismus and social phobia: A case-controlled study. *J AAPOS*, 13(3): 249–252. DOI: <https://doi.org/10.1016/j.jaapos.2009.02.010>
- Biglan, AW, Davis, JS, Day, R and Landsittel, D. 1994. Prospective investigation of the effectiveness of intraoperative adjustable sutures for correction of strabismus. *Trans Am Ophthalmol Soc*, 92: 325–347.
- Bradbury, JA and Taylor, RH. 2013. Severe complications of strabismus surgery. *J AAPOS*, 17(1): 59–63. DOI: <https://doi.org/10.1016/j.jaapos.2012.10.016>
- Bristol North Somerset and South Gloucestershire Clinical Commissioning Group. 2019. *Surgical Correction of Strabismus or Amblyopia in Adults Policy (version 1920.01.02)*. Available at https://bnssgccg-media.ams3.cdn.digitaloceanspaces.com/attachments/Strabismus_and_Amblyopia_Policy_PA_v1920.01.02_draft.pdf [Last accessed 7 November 2020].
- Bucci, MP, Brémond-Gignac, D and Kapoula, Z. 2009. Speed and accuracy of saccades, vergence and combined eye movements in subjects with strabismus before and after eye surgery. *Vision Res*, 49(4): 460–469. DOI: <https://doi.org/10.1016/j.visres.2008.12.006>
- Burke, JP, Leach, C and Davis, H. 1997. Psychosocial implications of strabismus surgery in adults. *J Pediatr Ophthalmol Strabismus*, 34(3): 159–164. DOI: <https://doi.org/10.3928/0191-3913-19970501-06>
- Chang, MY, Demer, JL, Isenberg, SJ, Velez, FG and Pineles, SL. 2017. Decreased binocular summation in strabismic

- amblyopes and effect of strabismus surgery. *Strabismus*, 25(2): 73–80. DOI: <https://doi.org/10.1080/09273972.2017.1318153>
- Chang, MY, Velez, FG, Demer, JL, Isenberg, SJ, Coleman, AL and Pineles, SL.** 2015. Quality of life in adults with strabismus. *Am J Ophthalmol*, 159(3): 539–544 e532. DOI: <https://doi.org/10.1016/j.ajo.2014.12.003>
- Cifuentes, DL, Pineles, SL, Demer, JL and Velez, FG.** 2018. Surgical success and lateral incomitance following three-muscle surgery for large-angle horizontal strabismus. *J AAPOS*, 22(1): 17–21. DOI: <https://doi.org/10.1016/j.jaapos.2017.10.005>
- Coats, DK, Paysse, EA, Towler, AJ and Dipboye, RL.** 2000. Impact of large angle horizontal strabismus on ability to obtain employment. *Ophthalmol*, 107(2): 402–405. DOI: [https://doi.org/10.1016/S0161-6420\(99\)00035-4](https://doi.org/10.1016/S0161-6420(99)00035-4)
- COMET Initiative: Core Outcome Measures in Effectiveness Trials.** Available at comet-initiative.org [Last accessed 17 September 2021].
- Currie, ZI, Shipman, T and Burke, JP.** 2003. Surgical correction of large-angle exotropia in adults. *Eye*, 17(3): 334–339. DOI: <https://doi.org/10.1038/sj.eye.6700347>
- Dadeya, S and Kamlesh, SF.** 2002. Preliminary results of intraoperative daunorubicin in strabismus surgery. *J Pediatr Ophthalmol Strabismus*, 39(6): 340–344. DOI: <https://doi.org/10.3928/0191-3913-20021101-08>
- Daga, D, Pujari, A, Sharma, P, Phuljhele, S and Saxena, R.** 2022. Evaluation of reinforced plication as an alternative to resection in exotropia. *Indian J Ophthalmol*, 70(3): 948–951. DOI: https://doi.org/10.4103/ijjo.IJO_2884_20
- Das, A, Hancox, J and RCOphth Quality and Safety Group.** 2017. *Strabismus surgery for adults in the United Kingdom: indications, evidence base and benefits. Commissioning Guidance.* Available at <https://www.rcophth.ac.uk/standards-publications-research/commissioning-in-ophthalmology/> [Last accessed 6 March 2021].
- Dawson, EL, Leung, H, Webster, A and Lee, JP.** 2013. Beneficial effect of treatment for strabismus in patients with bilaterally poor vision, who cannot see their strabismus. *Strabismus*, 21(1): 33–36. DOI: <https://doi.org/10.3109/09273972.2012.742119>
- Dotan, G, Nelson, LB, Mezad-Koursh, D, Stolovitch, C, Cohen, Y and Morad, Y.** 2014. Surgical outcome of strabismus surgery in patients with unilateral vision loss and horizontal strabismus. *J Pediatr Ophthalmol Strabismus*, 51(5): 294–298. DOI: <https://doi.org/10.3928/01913913-20140709-02>
- Durnian, JM, Noonan, CP and Marsh, IB.** 2011. The psychosocial effects of adult strabismus: A review. *Br J Ophthalmol*, 95(4): 450–453. DOI: <https://doi.org/10.1136/bjo.2010.188425>
- Durnian, JM, Owen, ME, Baddon, AC, Noonan, CP and Marsh, IB.** 2010. The psychosocial effects of strabismus: Effect of patient demographics on the AS-20 score. *J AAPOS*, 14(6): 469–471. DOI: <https://doi.org/10.1016/j.jaapos.2010.08.013>
- Elkamshoushy, A and Langué, MA.** 2019. Outcomes of bilateral lateral rectus recession in treatment of recurrent exotropia after bilateral medial rectus resection. *Eur J Ophthalmol*, 29(4): 402–405. DOI: <https://doi.org/10.1177/1120672118795066>
- Eino, D and Kraft, SP.** 1997. Postoperative drifts after adjustable-suture strabismus surgery. *Can J Ophthalmol*, 32(3): 163–169.
- Escardó-Paton, JA and Harrad, RA.** 2009. Duration of conjunctival redness following adult strabismus surgery. *J AAPOS*, 13(6): 583–586. DOI: <https://doi.org/10.1016/j.jaapos.2009.09.013>
- Estes, KJ, Parrish, RK, Sinacore, J, Mumby, PB and McDonnell, JF.** 2020. Effects of corrective strabismus surgery on social anxiety and self-consciousness in adults. *J AAPOS*, 24(5): 280.e281–280.e284. DOI: <https://doi.org/10.1016/j.jaapos.2020.05.017>
- Faridi, UA, Saleh, TA, Ewings, P and Twomey, JM.** 2007. Factors affecting the surgical outcome of primary exotropia. *Strabismus*, 15(3): 127–131. DOI: <https://doi.org/10.1080/09273970701506086>
- Fatima, T, Amitava, AK, Siddiqui, S and Ashraf, M.** 2009. Gains beyond cosmesis: Recovery of fusion and stereopsis in adults with longstanding strabismus following successful surgical realignment. *Indian J Ophthalmol*, 57(2): 141–143. DOI: <https://doi.org/10.4103/0301-4738.45505>
- Felius, J, Stager, DR, Jr, Beauchamp, GR and Stager, DR.** 2001. Re-recession of the medial rectus muscles in patients with recurrent esotropia. *J AAPOS*, 5(6): 381–387. DOI: <https://doi.org/10.1067/mpa.2001.119784>
- Fieß, A, Elflein, HM, Urschitz, MS, Pesudovs, K, Münzel, T, Wild, PS, ... Schuster, AK.** 2020. Prevalence of strabismus and its impact on vision-related quality of life – Results from the German population-based Gutenberg Health Study (GHS). *Ophthalmol*, 127(8): 1113–1122. DOI: <https://doi.org/10.1016/j.ophtha.2020.02.026>
- Frangouli, O and Adams, GG.** 2013. The use of amniotic membrane for the management of fibrosis in complex strabismus surgery. *Strabismus*, 21(1): 13–22. DOI: <https://doi.org/10.3109/09273972.2012.762531>
- Fujiike, K, Mizuno, Y, Hiratsuka, Y and Yamada, M.** 2011. Quality of life and cost-utility assessment after strabismus surgery in adults. *Jpn J Ophthalmol*, 55(3): 268–276. DOI: <https://doi.org/10.1007/s10384-011-0022-6>
- Ganesh, A, Pirouznia, S, Ganguly, SS, Fagerholm, P, and Lithander, J.** 2011. Consecutive exotropia after surgical treatment of childhood esotropia: a 40-year follow-up study. *Acta Ophthalmol*, 89(7): 691–695. DOI: <https://doi.org/10.1111/j.1755-3768.2009.01791.x>
- Ghiasi, G, Shojaei, A, Soltan-Sanjari, M, Kosari, M and Aslani, M.** 2013. The psychosocial improvement after strabismus surgery in Iranian patients. *Iranian J Ophthalmol*, 25(3): 211–215.
- Gigante, E, Romão, RA and Valério, FJP.** 2018. Monocular surgery to correct large-angle esotropia: A 10-year follow-up study. *Arq Bras Oftalmol*, 81(3): 232–238. DOI: <https://doi.org/10.5935/0004-2749.20180046>

- Glasman, P, Cheeseman, R, Wong, V, Young, J and Durnian, JM.** 2013. Improvement in patients' quality-of-life following strabismus surgery: Evaluation of postoperative outcomes using the Adult Strabismus 20 (AS-20) score. *Eye*, 27(11): 1249–1253. DOI: <https://doi.org/10.1038/eye.2013.174>
- Goff, MJ, Suhr, AW, Ward, JA, Croley, JK and O'Hara, MA.** 2006. Effect of adult strabismus on ratings of official U.S. Army photographs. *J AAPOS*, 10(5): 400–403. DOI: <https://doi.org/10.1016/j.jaapos.2006.04.011>
- Goseki, T and Ishikawa, H.** 2017. The prevalence and types of strabismus, and average of stereopsis in Japanese adults. *Jpn J Ophthalmol*, 61(3): 280–285. DOI: <https://doi.org/10.1007/s10384-017-0505-1>
- Gusek-Schneider, G and Boss, A.** 2010. Results following eye muscle surgery for secondary sensory strabismus. *Strabismus*, 18(1): 24–31. DOI: <https://doi.org/10.3109/09273971003623582>
- Hashemi, H, Nabovati, P, Yekta, A, Ostadimoghaddam, H, Behnia, B and Khabazkhoob, M.** 2017. The prevalence of strabismus, heterophorias, and their associated factors in underserved rural areas of Iran. *Strabismus*, 25(2): 60–66. DOI: <https://doi.org/10.1080/09273972.2017.1317820>
- Hassan, MB, Hodge, DO and Mohney, BG.** 2015. Prevalence of mental health illness among patients with adult-onset strabismus. *Strabismus*, 23(3): 105–110. DOI: <https://doi.org/10.3109/09273972.2015.1070881>
- Hatt, SR, Leske, DA, Bradley, EA, Cole, SR and Holmes, JM.** 2009. Development of a quality-of-life questionnaire for adults with strabismus. *Ophthalmol*, 116(1): 139–144 e135. DOI: <https://doi.org/10.1016/j.ophtha.2008.08.043>
- Hatt, SR, Leske, DA and Holmes, JM.** 2010. Responsiveness of health-related quality-of-life questionnaires in adults undergoing strabismus surgery. *Ophthalmol*, 117(12): 2322–2328.e2321. DOI: <https://doi.org/10.1016/j.ophtha.2010.03.042>
- Hatt, SR, Leske, DA, Liebermann, L and Holmes, JM.** 2012a. Changes in health-related quality of life 1 year following strabismus surgery. *Am J Ophthalmol*, 153(4): 614–619. DOI: <https://doi.org/10.1016/j.ajo.2011.10.001>
- Hatt, SR, Leske, DA, Liebermann, L and Holmes, JM.** 2012b. Comparing outcome criteria performance in adult strabismus surgery. *Ophthalmol*, 119(9): 1930–1936. DOI: <https://doi.org/10.1016/j.ophtha.2012.02.035>
- Hatt, SR, Leske, DA, Liebermann, L and Holmes, JM.** 2016. Incorporating health-related quality of life into the assessment of outcome following strabismus surgery. *Am J Ophthalmol*, 164: 1–5. DOI: <https://doi.org/10.1016/j.ajo.2015.12.029>
- Hatt, SR, Leske, DA, Liebermann, L, Philbrick, KL and Holmes, JM.** 2014. Depressive symptoms associated with poor health-related quality of life in adults with strabismus. *Ophthalmol*, 121(10): 2070–2071. DOI: <https://doi.org/10.1016/j.ophtha.2014.04.031>
- Hatt, SR, Leske, DA, Philbrick, KL and Holmes, JM.** 2018. Factors associated with failure of adult strabismus-20 questionnaire scores to improve following strabismus surgery. *JAMA Ophthalmol*, 136(1): 46–52. DOI: <https://doi.org/10.1001/jamaophthalmol.2017.5088>
- Hertle, RW.** 1998. Clinical characteristics of surgically treated adult strabismus. *J Pediatr Ophthalmol Strabismus*, 35(3): 138–145. DOI: <https://doi.org/10.3928/0191-3913-19980501-04>
- Jackson, S, Harrad, RA, Morris, M and Rumsey, N.** 2006. The psychosocial benefits of corrective surgery for adults with strabismus. *Br J Ophthalmol*, 90(7): 883–888. DOI: <https://doi.org/10.1136/bjo.2005.089516>
- Ji, N, Xu, M, Yu, H, Xu, J and Yu, X.** 2020. Factors influencing the self-reported sense of deviation in adults with successful surgical outcomes for strabismus. *BMC Ophthalmol*, 20(1): 9. DOI: <https://doi.org/10.1186/s12886-019-1299-3>
- Jung, EH and Kim, SJ.** 2018. Surgical results and factors affecting outcome in adult patients with sensory exotropia. *Eye*, 32(12): 1851–1857. DOI: <https://doi.org/10.1038/s41433-018-0189-x>
- Kannam, M, Sutrave, J, Kapoor, R, Tibrewal, S, Kekunnaya, R and Sachdeva, V.** 2021. Horizontal rectus muscle transplantation for recurrent and residual strabismus. *Strabismus*, 29(1): 42–50. DOI: <https://doi.org/10.1080/09273972.2020.1871378>
- Kattan, JM, Velez, FG, Demer, JL and Pineles, SL.** 2016. Relationship between binocular summation and stereoacuity after strabismus surgery. *Am J Ophthalmol*, 165: 29–32. DOI: <https://doi.org/10.1016/j.ajo.2016.02.020>
- Keskinbora, KH, Gonen, T and Horozoglu, F.** 2011. Outcome of surgery in long-standing infantile esotropia with cross fixation. *J Pediatr Ophthalmol Strabismus*, 48(2): 77–83. DOI: <https://doi.org/10.3928/01913913-20110303-03>
- Kim, MJ, Khwarg, SI, Kim, SJ and Chang, BL.** 2008. Results of re-operation on the deviated eye in patients with sensory heterotropia. *Korean J Ophthalmol*, 22(1): 32–36. DOI: <https://doi.org/10.3341/kjo.2008.22.1.32>
- Kim, Y, Kim, C, Kim, S, Han, Y, Chung, I, Seo, S, ... Yoo, J.** 2016. Difference of self-identity levels between strabismus patients and normal controls. *Korean J Ophthalmol*, 30(6): 410–415. DOI: <https://doi.org/10.3341/kjo.2016.30.6.410>
- Kishimoto, F and Ohtsuki, H.** 2012. Comparison of VF-14 scores among different ophthalmic surgical interventions. *Acta Med Okayama*, 66(2): 101–110. DOI: <https://doi.org/10.18926/amo/48260>
- Koc, F, Erten, Y and Yurdakul, NS.** 2013. Does restoration of binocular vision make any difference in the quality of life in adult strabismus. *Br J Ophthalmol*, 97(11): 1425–1430. DOI: <https://doi.org/10.1136/bjophthalmol-2013-303704>
- Kothari, M and Joshi, V.** 2014. The perceived personality traits of adults with digitally induced large angle strabismus and the impact of its correction. *Indian J Ophthalmol*, 62(7): 773–776. DOI: <https://doi.org/10.4103/0301-4738.138617>
- Kushner, BJ.** 1994. Binocular field expansion in adults after surgery for esotropia. *Arch Ophthalmol*, 112(5): 639–643. DOI: <https://doi.org/10.1001/archophth.1994.01090170083027>

- Kutschke, PJ and Scott, WE.** 2004. Prism adaptation in visually mature patients with esotropia of childhood onset. *Ophthalmol*, 111(1): 177–179. DOI: <https://doi.org/10.1016/j.ophtha.2003.04.010>
- Lee, TE, Kim, SH and Cho, YA.** 2013. Postoperative changes in spatial localization following exotropia surgery. *Curr Eye Res*, 38(1): 210–214. DOI: <https://doi.org/10.3109/02713683.2012.713151>
- Leske, DA, Hatt, SR and Holmes, JM.** 2010. Test-retest reliability of health-related quality-of-life questionnaires in adults with strabismus. *Am J Ophthalmol*, 149(4): 672–676. DOI: <https://doi.org/10.1016/j.ajo.2009.11.004>
- Leske, DA, Hatt, SR, Liebermann, L and Holmes, JM.** 2012. Evaluation of the Adult Strabismus-20 (AS-20) questionnaire using Rasch analysis. *Invest Ophthalmol Vis Sci*, 53(6): 2630–2639. DOI: <https://doi.org/10.1167/iovs.11-8308>
- Liebermann, L, Hatt, SR, Leske, DA and Holmes, JM.** 2013. Adjustment versus no adjustment when using adjustable sutures in strabismus surgery. *J AAPOS*, 17(1): 38–42. DOI: <https://doi.org/10.1016/j.jaapos.2012.10.017>
- Liebermann, L, Hatt, SR, Leske, DA and Holmes, JM.** 2014. Improvement in specific function-related quality-of-life concerns after strabismus surgery in nondiplopic adults. *J AAPOS*, 18(2): 105–109. DOI: <https://doi.org/10.1016/j.jaapos.2013.11.018>
- Lin, S, Congdon, N, Yam, JC, Huang, Y, Qiu, K, Ma, D, ... Zhang, M.** 2014. Alcohol use and positive screening results for depression and anxiety are highly prevalent among Chinese children with strabismus. *Am J Ophthalmol*, 157(4): 894–900. DOI: <https://doi.org/10.1016/j.ajo.2014.01.012>
- Lipton, JR and Willshaw, HE.** 1995. Prospective multicentre study of the accuracy of surgery for horizontal strabismus. *Br J Ophthalmol*, 79(1): 10–11. DOI: <https://doi.org/10.1136/bjo.79.1.10>
- MacKenzie, K, Hancox, J, McBain, H, Ezra, DG, Adams, G and Newman, S.** 2016. Psychosocial interventions for improving quality of life outcomes in adults undergoing strabismus surgery. *Cochrane Database Syst Rev*. 2016(5): CD010092. DOI: <https://doi.org/10.1002/14651858.CD010092.pub4>
- McBain, HB, Au, CK, Hancox, J, MacKenzie, KA, Ezra, DG, Adams, GGW and Newman, SP.** 2014a. The impact of strabismus on quality of life in adults with and without diplopia: A systematic review. *Surv Ophthalmol*, 59: 185–191. DOI: <https://doi.org/10.1016/j.survophthal.2013.04.001>
- McBain, HB, MacKenzie, KA, Au, C, Hancox, J, Ezra, DG, Adams, GG and Newman, SP.** 2014b. Factors associated with quality of life and mood in adults with strabismus. *Br J Ophthalmol*, 98(4): 550–555. DOI: <https://doi.org/10.1136/bjophthalmol-2013-304220>
- McBain, HB, MacKenzie, KA, Hancox, J, Ezra, DG, Adams, GG, and Newman, SP.** 2016. Does strabismus surgery improve quality and mood, and what factors influence this? *Eye*, 30(5): 656–667. DOI: <https://doi.org/10.1038/eye.2016.70>
- Menon, V, Saha, J, Tandon, R, Mehta, M and Khokhar, S.** 2002. Study of the psychosocial aspects of strabismus. *J Pediatr Ophthalmol Strabismus*, 39(4): 203–208. DOI: <https://doi.org/10.3928/0191-3913-20020701-07>
- Mohney, BG, McKenzie, JA, Capo, JA, Nusz, KJ, Mrazek, D and Diehl, NN.** 2008. Mental illness in young adults who had strabismus as children. *Pediatrics*, 122(5): 1033–1038. DOI: <https://doi.org/10.1542/peds.2007-3484>
- Mojon-Azzi, SM and Mojon, DS.** 2009. Strabismus and employment: The opinion of headhunters. *Acta Ophthalmol*, 87(7): 784–788. DOI: <https://doi.org/10.1111/j.1755-3768.2008.01352.x>
- Mojon-Azzi, SM, Potnik, W and Mojon, DS.** 2008. Opinions of dating agents about strabismic subjects' ability to find a partner. *Br J Ophthalmol*, 92(6): 765–769. DOI: <https://doi.org/10.1136/bjo.2007.128884>
- Murray, AD, Orpen, J and Calcutt, C.** 2007. Changes in the functional binocular status of older children and adults with previously untreated infantile esotropia following late surgical realignment. *J AAPOS*, 11(2): 125–130. DOI: <https://doi.org/10.1016/j.jaapos.2006.10.016>
- Natung, T, Pandey, I and Lyngdoh, LA.** 2022. Outcomes of a simplified, fixed surgical dosage calculation for uncomplicated, horizontal, concomitant strabismus in adults. *Indian J Ophthalmol*, 70(4): 1321–1326. DOI: https://doi.org/10.4103/ijo.IJO_2399_21
- Nelson, BA, Gunton, KB, Lasker, JN, Nelson, LB and Drohan, LA.** 2008. The psychosocial aspects of strabismus in teenagers and adults and the impact of surgical correction. *J AAPOS*, 12(1): 72–76. DOI: <https://doi.org/10.1016/j.jaapos.2007.08.006>
- Olitsky, SE, Sudesh, S, Graziano, A, Hamblen, J, Brooks, SE, and Shaha, SH.** 1999. The negative psychosocial impact of strabismus in adults. *J AAPOS*, 3(4): 209–211. DOI: [https://doi.org/10.1016/S1091-8531\(99\)70004-2](https://doi.org/10.1016/S1091-8531(99)70004-2)
- Ozates, S, Ezerbolat Ozates, M, Can, CU, Polat, S, Yasar, HH, Taskale, B and Gogus, AK.** 2019. Improvement in psychiatric symptoms after strabismus surgery in adolescent patients in long-term follow-up. *Br J Ophthalmol*, 103(7): 966–970. DOI: <https://doi.org/10.1136/bjophthalmol-2018-312510>
- Paysse, EA, Steele, EA, McCreery, KM, Wilhelmus, KR and Coats, DK.** 2001. Age of the emergence of negative attitudes toward strabismus. *J AAPOS*, 5(6): 361–366. DOI: <https://doi.org/10.1067/mpa.2001.119243>
- Pineles, SL, Demer, JL, Isenberg, SJ, Birch, EE and Velez, FG.** 2015. Improvement in binocular summation after strabismus surgery. *JAMA Ophthalmol*, 133(3): 326–332. DOI: <https://doi.org/10.1001/jamaophthalmol.2014.5265>
- Ribeiro G de, B, Bach, AG, Faria, CM, Anastásia, S and Almeida, HC.** 2014. Quality of life of patients with strabismus. *Arq Bras Oftalmol*, 77(2): 110–113. DOI: <https://doi.org/10.5935/0004-2749.20140027>
- Ritchie, A, Colapinto, P and Jain, S.** 2013. The psychological impact of strabismus: does the angle really matter?

- Strabismus*, 21(4): 203–208. DOI: <https://doi.org/10.3109/09273972.2013.833952>
- Ritchie, AE and Ali, N.** 2019. The incidence and clinical outcome of complications in 4,000 consecutive strabismus operations. *J AAPOS*, 23(3): 140.e141–140.e146. DOI: <https://doi.org/10.1016/j.jaapos.2018.08.015>
- Royal College of Ophthalmologists.** 2016. *Strabismus (squint) surgical intervention is not a cosmetic procedure*. Available at <https://www.rcophth.ac.uk/2016/09/squint-surgical-intervention-is-not-a-cosmetic-procedure/> [Last accessed 7 November 2020].
- Sah, SP, Sharma, IP, Chaudhry, M and Saikia, M.** 2017. Health-related quality of life (HRQoL) in young adults with strabismus in India. *J Clin Diagn Res*, 11(2): NC01–NC04. DOI: <https://doi.org/10.7860/JCDR/2017/24541.9389>
- Sandercoe, T, Beukes, S and Martin, F.** 2014. Adults with strabismus seek surgery for psychosocial benefits. *Taiwan J Ophthalmol*, 4(1): 17–20. DOI: <https://doi.org/10.1016/j.tjo.2013.10.004>
- Satterfield, D, Keltner, JL and Morrison, TL.** 1993. Psychosocial aspects of strabismus study. *Arch Ophthalmol*, 111: 1100–1105. DOI: <https://doi.org/10.1001/archophth.1993.01090080096024>
- Sefi-Yurdakul, N, Oto, S and Pelit, A.** 2022. Surgical treatment of consecutive exotropia: Comparison of different surgical methods applied to one eye in one session. *Eur J Ophthalmol*, 32(3): 1411–1416. DOI: <https://doi.org/10.1177/11206721211034288>
- Serafino, M, Granet, DB, Kushner, BJ, Dagi, LR, Kekunnaya, R and Nucci, P.** 2019. Use of the Delphi process for defining successful outcomes for strabismus surgery. *J AAPOS*, 23(6): 309–312. DOI: <https://doi.org/10.1016/j.jaapos.2019.07.006>
- Sim, PY, Cleland, C, Dominic, J and Jain, S.** 2018. Investigation of factors associated with the success of adult strabismus surgery from the patient's perspective. *J AAPOS*, 22(4): 266–271. DOI: <https://doi.org/10.1016/j.jaapos.2018.03.006>
- Tibrewal, S, Kapoor, R, Rath, SS, Sachdeva, V and Kekunnaya, R.** 2021. Extraocular Muscle Transplantation for Primary Treatment of Large-Angle Exotropia. *Journal of Binocular Vision and Ocular Motility*, 71(4): 150–160.
- Umazume, F, Ohtsuki, H and Hasebe, S.** 1997. Predictors of postoperative binocularity in adult strabismus. *Jpn J Ophthalmol*, 41(6): 414–421. DOI: [https://doi.org/10.1016/S0021-5155\(97\)00076-2](https://doi.org/10.1016/S0021-5155(97)00076-2)
- Wang, L and Nelson, LB.** 2011. Outcome study of graded unilateral medial rectus recession for small to moderate angle esotropia. *J Pediatr Ophthalmol Strabismus*, 48(1): 20–24. DOI: <https://doi.org/10.3928/01913913-20100818-05>
- Wang, Z, Ren, H, Frey, R, Liu, Y, Raphael, D, Bian, W and Wang, X.** 2014. Comparison of the Adult Strabismus Quality of Life Questionnaire (AS-20) with the Amblyopia and Strabismus Questionnaire (ASQE) among adults with strabismus who seek medical care in China. *BMC Ophthalmol*, 14: 139. DOI: <https://doi.org/10.1186/1471-2415-14-139>
- Wortham, EV and Greenwald, MJ.** 1989. Expanded binocular peripheral visual fields following surgery for esotropia. *J Pediatr Ophthalmol Strabismus*, 26(3): 109–112. DOI: <https://doi.org/10.3928/0191-3913-19890501-04>
- Xu, J, Yu, X, Huang, Y, Chen, J, Yu, H, Wang, Y and Zhang, F.** 2012. The psychosocial effects of strabismus before and after surgical correction in Chinese adolescents and adults. *J Pediatr Ophthalmol Strabismus*, 49(3): 170–175. DOI: <https://doi.org/10.3928/01913913-20110920-02>
- Xu, M, Yu, H, Chen, Y, Xu, J, Zheng, J and Yu, X.** 2016. Long-term quality of life in adult patients with strabismus after corrective surgery compared to the general population. *PLoS One*, 11(11): e0166418. DOI: <https://doi.org/10.1371/journal.pone.0166418>

TO CITE THIS ARTICLE:

Arblaster, G, Buckley, D, Barnes, S and Davis, H. 2024. Strabismus Surgery for Psychosocial Reasons—A Literature Review. *British and Irish Orthoptic Journal*, 20(1), pp. 107–132. DOI: <https://doi.org/10.22599/bioj.352>

Submitted: 09 November 2023

Accepted: 04 March 2024

Published: 22 April 2024

COPYRIGHT:

© 2024 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

British and Irish Orthoptic Journal is a peer-reviewed open access journal published by White Rose University Press.