

This is a repository copy of *Chain-length dependent organisation in mixtures of hydrogenous and fluorous ionic liquids*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/212776/>

Version: Accepted Version

---

**Article:**

Elstone, Naomi Sarah, Shaw, Emily, Shimizu, Karina et al. (10 more authors) (2024) Chain-length dependent organisation in mixtures of hydrogenous and fluorous ionic liquids. FARADAY DISCUSSIONS. pp. 55-78. ISSN: 1364-5498

<https://doi.org/10.1039/D4FD00047A>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:


<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Faraday Discussions

Accepted Manuscript



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

This article can be cited before page numbers have been issued, to do this please use: N. Elstone, E. Shaw, K. Shimizu, J. Lai, B. Demé, P. Lane, M. L. Costen, K. G. McKendrick, S. E. Youngs, S. E. Rogers, J. N. Canongia Lopes, D. W. Bruce and J. M. Slattery, *Faraday Discuss.*, 2024, DOI: 10.1039/D4FD00047A.

# Chain-length dependent organisation in mixtures of hydrogenous and fluoruous ionic liquids

View Article Online  
DOI: 10.1039/D4FD00047A

Naomi S. Elstone,<sup>a</sup> Emily V. Shaw,<sup>a</sup> Karina Shimizu\*,<sup>b</sup> Joshua Lai,<sup>a</sup> Bruno Demé,<sup>c</sup>  
Paul D. Lane,<sup>d</sup> Matthew L. Costen,<sup>d</sup> Kenneth G. McKendrick,<sup>d</sup> Sarah Youngs,<sup>e</sup>  
Sarah E. Rogers,<sup>e</sup> Jose N. Canongia Lopes,<sup>b</sup>  
Duncan W. Bruce\*<sup>a</sup> and John M. Slaterry\*<sup>a</sup>

<sup>a</sup> Department of Chemistry, University of York, Heslington, York YO10 5DD, UK.

<sup>b</sup> Centro de Química Estrutural, Institute of Molecular Sciences, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais 1049 001 Lisboa, Portugal.

<sup>c</sup> Institut Laue-Langevin, 38000 Grenoble, France.

<sup>d</sup> Institute of Chemical Sciences, School of Engineering and Physical Sciences, Heriot-Watt University, Edinburgh, EH14 4AS, U.K.

<sup>e</sup> ISIS Neutron and Muon Source, Harwell Campus, Didcot, OX11 0QX, UK.

## Abstract

As part of an ongoing study of the structure and properties of mixtures of ionic liquids in which one component has a hydrocarbon chain and the other a semiperfluorocarbon chain, we now report a study of the mixtures  $[C_8MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$ ,  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  and  $[C_{10}MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$ , where  $[C_8MIM][Tf_2N]$  is 1-methyl-3-octylimidazolium bis(trifluoromethylsulfonyl)imide,  $[C_{10}MIM][Tf_2N]$  is 1-decyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide,  $[C_8MIM-F_{13}][Tf_2N]$  is 1-(1*H*,1*H*,2*H*,2*H*-perfluorooctyl)-3-methylimidazolium bis(trifluoromethylsulfonyl)imide and  $[C_{10}MIM-F_{17}][Tf_2N]$  is 1-(1*H*,1*H*,2*H*,2*H*-perfluorodecyl)-3-methylimidazolium bis(trifluoromethylsulfonyl)imide. The mixtures were investigated using small-angle X-ray (SAXS) and neutron (SANS) scattering complemented by molecular dynamics simulations (with viscosity and surface tension measurements also possible for the mixtures  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$ ). Unlike previous studies of  $[C_8MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$ , where no strong evidence of alkyl/fluoroalkyl chain segregation or triphilic behaviour was seen (Elstone *et al.*, *J. Phys. Chem. B*, 2023, **127**, 7394-7407), these new mixtures showed the



formation of small aggregates of varying sizes of each component, even though all were co-miscible across the full range of compositions. Thus, while a clear polar non-polar peak (PNPP) was observed at large or small values of  $x$ , at intermediate compositions the small-angle neutron scattering at low  $q$  was dominated by scattering from these small aggregates, with little or no evidence of the PNPP at some compositions. The origins of this behaviour are discussed in terms of inter-chain interactions.

## Introduction

The preparation of mixtures of ionic liquids (ILs) represents a time- and cost-effective way to generate families of materials whose properties can be tuned by simply changing the proportion of each component.<sup>1, 2</sup> This offers some significant advantages to the alternative approach of synthesising many individual, bespoke ILs to access a range of liquids with different properties. The field of supported ionic liquid phase (SILP) catalysis is one where mixtures of ILs has the possibility to have significant impact.<sup>3-7</sup> SILPs are generally prepared by coating the surface of porous, solid supports such as silica with an IL film containing a catalyst, which can be either a homogeneous or nanoparticle system.<sup>8-16</sup> In both cases, the solvent environment around the active site(s) of the catalyst and the gas/liquid interface (where gaseous reagents are employed) will have a profound effect on reactivity, and both naturally depend upon the properties of the IL in which the catalyst is dissolved/suspended. As such, methods to prepare, understand and tune the behaviour of IL mixtures will provide important approaches to allow the rational design of more effective SILP catalyst systems.

We have reported previously on the surface and bulk composition of alkyl-functionalised imidazolium ILs and their mixtures (**1**) using a combination of reactive-atom scattering (RAS) in conjunction with laser-induced fluorescence (RAS-LIF),<sup>17-23</sup> and with mass spectrometry (RAS-MS) along with surface tension to characterise the gas-liquid interface,<sup>24-28</sup> while using small-angle neutron (SANS) and X-ray (SAXS) scattering alongside viscometry and conductivity to characterise the bulk material.<sup>22, 25, 29, 30</sup> These studies were complemented by molecular dynamics (MD) simulations, which in each case brought valuable additional understanding. Thus, for example, it



was possible to demonstrate the preference of long-chain ILs to concentrate at the gas-liquid or vacuum-liquid interface when dissolved in a short-chain host,<sup>22</sup> and percolation behaviour as a function of chain length was established for a wide range of mixtures.<sup>29</sup> Similar behaviour in related mixtures has been explored in detail by others using SAXS,<sup>31-33</sup> second harmonic generation,<sup>34</sup> NMR spectroscopy,<sup>35</sup> angle-resolved XPS,<sup>36</sup> X-ray reflectivity,<sup>37</sup> and combinations of other techniques.<sup>38-41</sup> Other recent studies on IL mixture systems have explored concepts such as hydrogen-bonding in protic IL mixtures,<sup>42</sup> machine learning for IL mixture property prediction,<sup>43</sup> and the pressure-dependent properties of IL mixtures.<sup>44</sup>




Figure 1: Hydrocarbon mixtures (**1**) and hydrocarbon/fluorocarbon mixtures (**2**) studied previously.

More recently, our attention has turned to binary mixtures where one component has a hydrocarbon chain and the other a fluorocarbon chain.<sup>23, 30</sup> Interest in such mixtures arises from preliminary observations that show preferential location of fluorocarbons at the surface of such mixtures, combined with the possibilities offered by local organisation given the predisposition of at least longer hydrocarbon and fluorocarbon chains to de-mix in the bulk,<sup>26, 45</sup> and at the vacuum-liquid interface.<sup>46-48</sup> Recent work has shown that these effects can be applied to metal complexes of relevance to SILP systems to affect surface enrichment of metal complexes that may prove beneficial in catalysis.<sup>28, 49, 50</sup> As part of our studies in the area, we recently reported on a combined viscometry, SANS, SAXS and MD study of the binary mixtures **2** ( $n = 8$ ), which we found to be miscible continuously across the range of compositions.<sup>26, 30</sup> Although hinted at in previous data on a related system,<sup>45</sup> we observed no strong evidence for triphasic, i.e. segregation of fluoroalkyl and alkyl chains, behaviour in these mixtures like that seen in related ILs.<sup>51-53</sup> This may



have been due to the length of the fluoroalkyl chains - only 6 of the carbon atoms in the chain were fluorinated, as a C<sub>2</sub>H<sub>4</sub> unit is included in between the fluoroalkyl chain and the imidazolium ring to aid the synthesis of these salts. It has been recognised in previous work that so called 'fluorous' effects are often stronger with increasing fluoroalkyl chain length.<sup>54</sup> Therefore, in order to investigate the effect of chain length on such mixtures, we now report an analogous study using longer-chain-length materials (**2**) in which both chains are the same overall length ( $n = 10$ ,  $m = 8$ ) or where they differ by two carbon atoms ( $n = 8$ ,  $m = 8$  and  $n = 10$ ,  $m = 6$ ), plus some limited data where  $n = 12$ ,  $m = 10$ .

## Results

### Synthesis

The ILs were prepared as described previously,<sup>23, 26, 30</sup> shown in Figure 2, and full details are given in the Supporting Information (SI). The hydrogenous bromide salts were prepared in yields that were very close to quantitative, while anion metathesis showed yields >90% after rigorous purification and drying. In order to minimise side reactions, which are promoted at higher temperatures, the quaternisation reactions to prepare the intermediate semiperfluoroalkyl iodides were run at moderate temperatures (60 °C) with long reaction times of around 3 weeks. Yields were nonetheless good at >70%. Metathesis to the [Tf<sub>2</sub>N]<sup>-</sup> salts was performed in a similar way to the hydrogenous materials, although with lower isolated yields (typically *ca.* 60%) and the final products invariably retained some amber colouration after purification. Despite this, the products showed no evidence of iodine/iodide impurities, based on AgNO<sub>3</sub> tests, combustion analysis results were within the expected tolerance for C, H and N and NMR spectra showed no evidence of impurities, within the sensitivity of the technique. Presumably, a very low concentration of a coloured impurity with a large molar absorption coefficient in the visible region is present that does not otherwise affect the analytical data, or in our experience the scattering and physical property data for these materials.






Figure 2 Synthesis of the salts used in this study.

### Mixtures Prepared for Study

Studies were undertaken on the mixtures  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$ ,  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  and  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  over the compositional range where  $x = 0, 0.05, 0.2, 0.35, 0.5, 0.65, 0.8, 0.95$  and  $1$ .  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures were liquid at room temperature for all of these compositions, while mixtures of  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  and  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  with the highest proportion of  $[\text{C}_{10}\text{MIM-F}_{17}][\text{Tf}_2\text{N}]$  tended to be solid at room temperature. Thus, the higher melting point of  $[\text{C}_{10}\text{MIM-F}_{17}][\text{Tf}_2\text{N}]$  precluded accurate measurement of its density with the vibrating tube densitometer used for the other ILs, which in turn precluded determination of surface tension *via* the pendant drop method, which requires the density of the liquid to be known. Likewise, instrumental constraints meant that viscosity data could not be acquired across the full compositional range in these mixtures.

### Surface Tension and Viscosity for the Mixtures $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$

As shown in Figure 3, the viscosity of  $[\text{C}_8\text{MIM-F}_{13}][\text{Tf}_2\text{N}]$  (1083 cP at 25 °C) is significantly higher than that of  $[\text{C}_{10}\text{MIM}][\text{Tf}_2\text{N}]$  (120 cP at 25 °C). The relationship between composition and viscosity for mixtures of these two ILs is not linear, rather it broadly follows the logarithmic form of the Arrhenius mixing rule, albeit with some deviation from this.<sup>55</sup> Interestingly, and in common with data from  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures,<sup>30</sup> the deviation from the Arrhenius mixing rule is negative, such that the increased viscosity of the fluoro component is somehow damped compared to what would be expected based on the simple mixing relationship. This is compared with data from  $[\text{C}_{12}\text{MIM}]_{1-x}[\text{C}_2\text{MIM}]_x[\text{Tf}_2\text{N}]$  mixtures, where the deviation from the Arrhenius mixing rule was in fact positive.<sup>22</sup> This was discussed in previous work in terms of the formation of more organised non-polar domains with increasing  $x$  in  $[\text{C}_{12}\text{MIM}]_{1-x}[\text{C}_2\text{MIM}]_x[\text{Tf}_2\text{N}]$ , which resist shear stress, and disruption of ion packing when rigid fluoroalkyl chains are introduced in  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$ , which does the opposite.<sup>22, 30</sup>






Figure 3 a) Viscosity measured at 25 °C for the mixture series  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$ , showing the actual data, the Arrhenius fit and the hypothetical linear relationship with composition. b) Percentage deviation from Arrhenius model for  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  and related mixtures.<sup>30</sup>

Figure 4 shows the surface tension data for the  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  mixtures and reveals both the lower surface tension of the fluorinated IL and, through the non-linear variation with composition, the fact that the fluorinated component preferentially accumulates at the air-liquid interface. Once more, this is similar to the behaviour reported for the mixture series  $[C_8MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$ . Measurement of the density as a function of composition showed that the variation in density is linear, based on composition, for these mixtures (Table S1).




Figure 4 Surface tension data for the mixture series  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  recorded at 20 °C.





## Scattering Experiments

Scattering experiments were performed on all IL mixture systems, at 20 °C for  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  and 60 °C for the other mixture systems, due to the higher melting points of some compositions. Small-angle neutron scattering (SANS) measurements were carried out on beamline D16 at the ILL in Grenoble and full experimental details are contained in the SI. Small-angle X-ray scattering (SAXS) measurements were carried out on a Bruker D8 Discover diffractometer equipped with a bespoke temperature-controlled, bored-graphite rod furnace, custom built at the University of York. Cu-K $\alpha$  ( $\lambda = 0.154056$  nm) radiation was used, generated from a 1 mS microfocus source. As described previously, X-ray and neutron scattering data for ILs often display three key features in different regions of the scattering vector  $q$ .<sup>56</sup> At higher  $q$  (around 1.3 to 1.4 Å<sup>-1</sup>) the so called contact peak (CP) is related to anion-cation, but also chain-chain, correlations.<sup>30</sup> At intermediate values of  $q$  (around 0.8 to 0.9 Å<sup>-1</sup>) the charge-ordering peak (COP) is often interpreted in terms of cation-anion-cation and anion-cation-anion correlations. In addition, in ILs containing ions with sufficiently long (typically > C6) alkyl chains an additional peak, the polar non-polar peak (PNPP) is often seen at lower  $q$  (around 0.4 Å<sup>-1</sup>, but strongly dependent on chain length). This is indicative of the formation of a local bilayer structure in the IL with short-range order resulting from separation of the polar (anion and cationic headgroup) and non-polar (alkyl chain) components and accompanies the formation of a percolated structure. This nanostructure in ILs has been known for some time and is often studied using small-angle scattering and molecular dynamics (MD) simulations.<sup>57-62</sup>

### $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$

Considering the  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  mixtures first (Figure 5), the SAXS data show a CP for all mixtures at  $q \approx 1.3$  Å<sup>-1</sup>, which moves to lower  $q$  as the proportion of the fluorinated component increases. Thus, for pure  $[\text{C}_{10}\text{MIM}][\text{Tf}_2\text{N}]$  it is centred at  $q = 1.34$  Å<sup>-1</sup> corresponding to a length scale of 4.7 Å, while for pure  $[\text{C}_{10}\text{MIM-F}_{17}][\text{Tf}_2\text{N}]$  it has moved to  $q = 1.15$  Å<sup>-1</sup>, which corresponds to a length scale of 5.5 Å. The mixtures also show a COP in the SAXS data around 0.8 Å<sup>-1</sup> and as the proportion of  $[\text{C}_{10}\text{MIM-F}_{17}][\text{Tf}_2\text{N}]$  increases, the COP moves to higher  $q$ , from 0.83



$\text{\AA}^{-1}$  ( $7.6 \text{ \AA}$ ) to  $0.91 \text{ \AA}^{-1}$  ( $6.9 \text{ \AA}$ ). These trends were also observed in the study of  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures and will be discussed in the context of this work below.<sup>30</sup>

View Article Online

DOI: 10.1039/D4FD00047A




Figure 5 a) SAXS data and b) SANS data collected for  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  mixtures at  $60 \text{ }^\circ\text{C}$ . Data for each composition are shifted along the  $y$  axis for clarity. Note the different scales used for the  $x$ -axes.

To lower  $q$  than both the CP and the COP is the PNPP and while this is observed in the SAXS data at higher proportions of  $[\text{C}_{10}\text{MIM}][\text{Tf}_2\text{N}]$  in the mixture, it gradually disappears as  $x$  increases, as there is effectively no contrast arising from the scattering length densities (SLDs) of the fluorocarbon chains (X-ray  $\text{SLD} \sim 13.7 \times 10^{-6} \text{ \AA}^{-2}$ ) compared with the ionic components (X-ray  $\text{SLD} \sim 13.2 \times 10^{-6} \text{ \AA}^{-2}$ ) of the system.<sup>30</sup> Due to the differences in neutron and X-ray SLDs we previously observed that in SANS experiments the PNPP was visible at all compositions of  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$ . However, in the present case the PNPP was observed at the highest proportions of  $[\text{C}_{10}\text{MIM}][\text{Tf}_2\text{N}]$  and for pure  $[\text{C}_{10}\text{MIM-F}_{17}][\text{Tf}_2\text{N}]$ , but not at intermediate concentrations (Figure 5b).

Closer examination of the SANS data shows significant differences to those reported previously for  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures. For the shorter-chain system, very little low  $q$  (below the PNPP) scattering was observed.<sup>30</sup> Figure 5b shows that there is a good deal of scattering at low  $q$  for the  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  mixtures. Indeed, it is likely that this is important in



understanding the behaviour of the PNPP and differences in the liquid structure with increasing alkyl and fluoroalkyl chain lengths in general. We considered whether these differences might relate to the temperature of the experiments, so collected the SANS data at 20 °C, to match the temperature used for the previous measurements on the shorter-chained system (see SI for details, Fig S40-S41). This showed that the temperature change had a negligible effect upon the scattering features, with data being well fitted using the same parameters, so changes are likely to be due to differences in liquid structure.




Figure 6 SANS data for  $[C_{10}MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$  with different models applied for a)  $x = 0.2$ , b)  $x = 0.5$  and c)  $x = 0.8$ .



The low- $q$  data were fitted to various scattering models to better understand the system. The form of the scattering at low- $q$  is consistent with scattering from discrete scattering objects, e.g. spherical aggregates, with different SLDs to the bulk liquid. There is a significant variation in the intensity of the scattering from these objects across the compositions, with the strongest scattering observed around  $x = 0.35$ . In order to account for both the low- $q$  scattering and the PNPP present in the data three fitting methodologies were attempted: 1) A combination of spherical scattering objects for the low  $q$  data and a Lorentzian peak for the PNPP. 2) An Ornstein-Zernike model for the low  $q$  data in combination with a Lorentzian peak for the PNPP. 3) The Teubner Strey model, which was developed to describe scattering in microemulsions that can form discrete scattering objects, bicontinuous networks or lamellar structures, depending on composition.<sup>63</sup> Figure 6 shows the three models applied to three mixtures with different compositions ( $x = 0.2, 0.5$  and  $0.8$ ), chosen to demonstrate different scattering characteristics (full fitting information can be found in the SI, pages S8-S18). The combination of the spherical model and Lorentzian peak gave the best fit across the whole compositional range and the parameters extracted from these fits are shown along with the peak positions in Table 1, noting that the position of the PNPP has been fixed to the position corresponding to the closest pure IL when its position could not be clearly identified in the mixtures. In our previous fitting of SANS data for  $[\text{C}_2\text{MIM}]_{1-x}[\text{C}_{12}\text{MIM}]_x[\text{Tf}_2\text{N}]$  mixtures,<sup>22</sup> the Teubner Strey and Ornstein–Zernike/Lorentzian peak models also fitted the data well across the whole compositional range. For the  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  system, however, these models proved less satisfactory (see SI for details, pages S8-S16). The Ornstein–Zernike/Lorentzian peak model fits are similar, but not quite as good, as the sphere/Lorentzian peak model. The Teubner Strey fits are able to describe either the low- $q$  scattering, or the PNPP, but in compositions where both are present do not capture the balance between them well. As such, only the sphere/Lorentzian peak model parameters will be discussed further.

View Article Online  
DOI: 10.1039/D4FD00047A



Table 1 Parameters used for fitting  $[C_{10}MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$  data to a sphere + Lorentzian model, along with the peak positions of the three peaks seen in the scattering data.

View Article Online

DOI: 10.1039/D4FD00047A

$x$	Radius / Å	PNPP / Å	COP / Å	CP / Å
$[C_{10}MIM][Tf_2N]$	-	$22 \pm 1$	$7.6 \pm 0.1$	$4.69 \pm 0.04$
<b>0.05</b>	-	$22 \pm 2$	$7.6 \pm 0.2$	$4.72 \pm 0.04$
<b>0.2</b>	$12.0 \pm 0.3$	$22 \pm 2$	$7.5 \pm 0.2$	$4.99 \pm 0.08$
<b>0.35</b>	$12.0 \pm 0.4$	$22 \pm 2$	$7.3 \pm 0.1$	$5.15 \pm 0.08$
<b>0.5</b>	$10.8 \pm 0.1$	$22 \pm 2$	$7.3 \pm 0.2$	$5.19 \pm 0.09$
<b>0.65</b>	$7.8 \pm 0.2$	$23 \pm 1$	$7.1 \pm 0.1$	$5.37 \pm 0.05$
<b>0.8</b>	$7.7 \pm 0.1$	$26 \pm 1$	$7.0 \pm 0.1$	$5.46 \pm 0.05$
<b>0.95</b>	$7.6 \pm 0.6$	$26 \pm 1$	$7.0 \pm 0.1$	$5.51 \pm 0.10$
$[C_{10}MIM-F_{17}][Tf_2N]$	-	$26 \pm 1$	$6.9 \pm 0.2$	$5.46 \pm 0.09$

The data in Table 1 quantify the changes in CP, COP and PNPP positions described above. In addition, for compositions where  $0.2 \leq x \leq 0.95$  the radius of the scattering objects that describe the low- $q$  scattering are seen to decrease in size with increasing  $x$ , from 12.0 to 7.6 Å. For reference, the length of a fully extended  $C_{10}H_{21}$  alkyl chain is  $\sim 13$  Å and of a  $C_8F_{17}$  chain is  $\sim 12$  Å and so the diameter of the scattering objects is on the order of two chains, with some interdigitation and/or folding. However, the necessity to fix the position of the PNPP to that observed in one of the pure components gives uncertainty in its value in the mixtures. The interpretation of this will be discussed further in the discussion section below.

### $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$ and $[C_8MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$

It was reported recently that the presence of significant low- $q$  scattering was not observed in  $[C_8MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  mixtures,<sup>30</sup> which is a significant difference to the behaviour described above. Therefore, in order to investigate the relationship between increases in chain length in either of the components of the mixture, and the impact of this on the presence of low- $q$  scattering from chain-chain aggregates, mixtures with combinations of C8 and C10 chain lengths were explored.






Figure 7 a) SAXS data and b) SANS data collected for  $[C_8MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$  mixtures at 60 °C. Data for each composition are shifted along the  $y$  axis for clarity. Note the different scales used for the  $x$ -axes.

SAXS and SANS data for  $[C_8MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$  mixtures are shown in Figure 7 and the peak positions and fitted parameters are shown in Table 2. These mixtures show similar features to those seen in the  $[C_{10}MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$  mixtures in terms of variation of both the CP (moves to lower  $q$ , from 1.33 (4.7 Å) to 1.15 (5.5 Å) Å<sup>-1</sup>, with increasing  $x$ ) and the COP (moves from 0.83 (7.6 Å) to 0.91 (6.9 Å) Å<sup>-1</sup> with increasing  $x$ ). These are consistent with similar changes in the polar network and chain-chain correlations in the mixtures, which appear to be insensitive to aliphatic chain length. The PNPP for  $[C_8MIM][Tf_2N]$  is weaker in intensity than that seen in  $[C_{10}MIM][Tf_2N]$ , as the C8 chains are relatively short and the polar/non-polar separation is not as well established compared to the longer alkyl chains. The low- $q$  scattering, fitted above as spherical scattering objects in the  $[C_{10}MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$  mixtures, is still present when the aliphatic chain length is reduced from C10 to C8. The size of the scattering objects follows a broadly similar pattern, again suggesting aggregates that are approximately two chains in diameter. Further discussion follows below.





Table 2 Parameters used for fitting  $[C_8MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$  data to a sphere + Lorentzian model, along with the peak positions of the three peaks seen in the scattering data.

$x$	Radius / Å	PNPP / Å	COP / Å	CP / Å
$[C_8MIM][Tf_2N]$	-	$18 \pm 1$	$7.6 \pm 0.1$	$4.72 \pm 0.04$
0.05	-	$18 \pm 1$	$7.6 \pm 0.2$	$4.80 \pm 0.04$
0.2	$13.7 \pm 0.4$	$20 \pm 1$	$7.4 \pm 0.1$	$4.99 \pm 0.04$
0.35	$10.8 \pm 0.7$	-	$7.2 \pm 0.1$	$5.24 \pm 0.04$
0.5	$8.1 \pm 0.2$	-	$7.2 \pm 0.1$	$5.37 \pm 0.09$
0.65	$6.8 \pm 0.2$	-	$7.1 \pm 0.1$	$5.46 \pm 0.04$
0.8	$5.9 \pm 0.4$	$27 \pm 1$	$7.0 \pm 0.1$	$5.42 \pm 0.05$
0.95	-	$26 \pm 1$	$6.9 \pm 0.2$	$5.46 \pm 0.05$
$[C_{10}MIM-F_{17}][Tf_2N]$	-	$26 \pm 1$	$6.9 \pm 0.2$	$5.46 \pm 0.09$

SAXS and SANS data for  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  mixtures are shown in Figure 8 and the peak positions and fitted parameters are shown in Table 3.




Figure 8 a) SAXS data and b) SANS data collected for  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  mixtures at 20 °C. Data for each composition are shifted along the y axis for clarity. Note the different scales used for the x-axes.

Changes in the CP and COP positions are similar to those described above. However, given the lack of significant low- $q$  scattering in  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures,<sup>30</sup> it was surprising to see that the scattering data for  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  more closely resembled those where the two chain lengths are C10, than where they are both C8. Thus, for  $0.20 \leq x \leq 0.80$  there is significant low- $q$  scattering that is similar in form to that described above and was also fitted using a spherical scattering object model. This suggests that the origin of this organisation of these liquids is not simply a function of the length of the fluoroalkyl chain. When considering the sphere radii in Table 3, there are trends similar to those found in the other liquids reported here. However, in the case of the  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures, the chain-chain aggregates represented by the spherical scattering objects are smaller at higher values of  $x$  than in mixtures containing the  $\text{C}_{10}\text{H}_4\text{F}_{17}$  chains. This is discussed in more detail below.

View Article Online  
DOI: 10.1039/D4FD00047A





Table 3 Parameters used for fitting  $[C_8MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$  data to a sphere + Lorentzian model, along with the peak positions of the three peaks seen in the scattering data.

View Article Online

DOI: 10.1039/D4FD00047A

$x$	Radius / Å	PNPP / Å	COP / Å	CP / Å
$[C_{10}MIM][Tf_2N]$	-	$22 \pm 1$	$7.6 \pm 0.1$	$4.69 \pm 0.04$
<b>0.05</b>	-	$22 \pm 1$	$7.5 \pm 0.1$	$4.72 \pm 0.07$
<b>0.2</b>	$13.7 \pm 0.4$	$24 \pm 1$	$7.3 \pm 0.1$	$4.80 \pm 0.04$
<b>0.35</b>	$7.5 \pm 0.2$	$26 \pm 1$	$7.1 \pm 0.1$	$4.91 \pm 0.04$
<b>0.5</b>	$6.8 \pm 0.1$	-	$7.0 \pm 0.1$	$4.99 \pm 0.04$
<b>0.65</b>	$7.4 \pm 0.2$	-	$6.9 \pm 0.1$	$5.03 \pm 0.08$
<b>0.8</b>	$7.3 \pm 0.2$	-	$6.7 \pm 0.1$	$5.07 \pm 0.04$
<b>0.95</b>	$8.9 \pm 0.2$	$22 \pm 1$	$6.6 \pm 0.1$	$5.15 \pm 0.08$
$[C_8MIM-F_{13}][Tf_2N]$	-	$22 \pm 1$	$6.6 \pm 0.1$	$5.11 \pm 0.08$

MD Simulations of the Bulk

Details of the MD simulations are found in the SI and calculations were carried out for the three series under investigation, namely  $[C_8MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$ ,  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  and  $[C_{10}MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$ . In addition, the data obtained from the simulations of  $[C_8MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$ , published previously,<sup>30</sup> were interrogated further to provide comparative data in relation to aggregation behaviour. These latter data were obtained at the slightly lower simulation temperature of 300 K, but it is not believed that this has a significant bearing on the comparisons made.

Figure 9 illustrates the structure factor functions,  $S(q)$ , for the  $[C_8MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$ ,  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  and  $[C_{10}MIM]_{1-x}[C_{10}MIM-F_{17}]_x[Tf_2N]$  mixtures at 333 K, and  $[C_8MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$  mixtures at 300 K, for  $0 \leq x \leq 1$ . The analysis focuses primarily on the low- $q$  regions of the  $S(q)$  functions ( $0.1 < q < 1.7 \text{ \AA}^{-1}$ ) to capture the structural features at the intermolecular level, specifically the PNPP, COP, and CP. The precise  $q$ -values of these primary peaks were determined through the deconvolution of their respective  $S(q)$  functions, noting that the very broad PNPP observed in some mixtures required up to four Gaussian curves for this. The deconvolution results are presented in Table S6.






Figure 9. Total structure factor functions,  $S(q)$ : (a)  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$ ; (b)  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$ ; (c)  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$ ; (d)  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  mixtures. Data for (a) at 300 K and for (b) to (c) at 330 K.

The  $q$ -values of the PNPP for all new mixtures with a component with a  $\text{C}_{10}$  chain (Table S6) represent characteristic distances of approximately 25  $\text{\AA}$ , indicating localised segregation between the polar and non-polar domains. The distance in the mixtures tends to be slightly greater than that of either pure component, with the effective distance greater in the pure fluorinated component. Likewise, the data show that the COP spacing decreases (*ca* 7.5 to 7.3  $\text{\AA}$ ) and the CP spacing increases (*ca* 4.6 to *ca* 5.2  $\text{\AA}$ ) as the proportion of the fluorinated component in the mixture increases. This overall behaviour has a direct parallel in the behaviour of the  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures and the details will be discussed below. Pleasingly, there is



good agreement between the spacings for the CP and the COP obtained from both experiment and theory, as illustrated in Figure S43.

View Article Online  
DOI: 10.1039/D4FD00047A

As noted above, the major difference in the data presented here compared with the previous study of the mixtures  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  is the presence of significant low- $q$  scattering. MD trajectories were therefore analysed thoroughly to identify possible aggregation patterns within the non-polar domains of the different systems, starting with mixtures where the chain length of just one component was increased. The baselines for these comparisons are represented by the pure materials  $[\text{C}_8\text{MIM}][\text{Tf}_2\text{N}]$  and  $[\text{C}_8\text{MIM-F}_{13}][\text{Tf}_2\text{N}]$ , data for which show each to be percolated (Figure 10a and Figure 10b, respectively). Thus, in this figure and those that follow, the  $y$ -axis represents the probability of finding a chain belonging to an aggregate of a particular size  $n_a$ , illustrate the distribution of chain-chain aggregates in the mixtures. The  $y$ -axis is normalised by the total number of corresponding monomers in the MD simulation box. Note that aggregation data were not reported in the earlier publication of the study of the  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures.<sup>30</sup>






Figure 10 Discrete probability distribution function,  $P(n_a)$ , of  $[C_8MIM]$  chains belonging to a non-polar aggregate for pure samples of (a)  $[C_8MIM][Tf_2N]$ ; (b)  $[C_8MIM-F_{13}][Tf_2N]$ ; (c)  $[C_{10}MIM][Tf_2N]$  and (d)  $[C_{10}MIM-F_{17}][Tf_2N]$ .

On addition of  $[C_8MIM-F_{13}][Tf_2N]$  to give  $[C_8MIM]_{0.65}[C_8MIM-F_{13}]_{0.35}[Tf_2N]$  (Figure 11a), the alkyl chains remain close to percolation with a cluster of aggregates around  $n_a \approx 0.7$ ; a number of smaller aggregates observed in addition to this. At this composition, the fluorocarbon chains form a range of rather small aggregates (Figure 11b). By the time the mixture has an equimolar composition, the situation has changed and now it is the hydrocarbon chains that are behaving as small aggregates, while the fluorocarbon chains are significantly concentrated into large aggregates around  $n_a \approx 0.7$  en route to percolation (Figure 11c and 11d), which is established by  $x = 0.8$  (Figure S44b). If the longer-chain  $[C_{10}MIM-F_{17}][Tf_2N]$  is now used, at  $x = 0.35$  both the hydrocarbon and fluorocarbon chains are dispersed in the medium across a range of aggregate sizes up to around  $n_a \approx 0.6$  and the form of the distribution is remarkably similar (Figure 12a and 12b). However, once the mixture is equimolar in composition then  $[C_{10}MIM-F_{17}][Tf_2N]$  is all but



percolated with a very small number of very small aggregates, while all of the hydrocarbon chains are concentrated also into very small aggregates (Figure 12c and 12d).

View Article Online  
DOI: 10.1039/D4FD00047A




Figure 11 Discrete probability distribution function,  $P(n_a)$ , for  $[C_8MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]$ :  $[C_8MIM]$  chains at (a)  $x = 0.35$ ; (c)  $x = 0.5$ ;  $[C_8MIM-F_{13}]$  chains at (b)  $x = 0.35$ ; (d)  $x = 0.5$ .






Figure 12 Discrete probability distribution function,  $P(n_a)$ , for  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F17}]_x[\text{Tf}_2\text{N}]: [\text{C}_8\text{MIM}]$  chains at (a)  $x = 0.35$ ; (c)  $x = 0.5$ ;  $[\text{C}_{10}\text{MIM-F17}]$  chains at (b)  $x = 0.35$ ; (d)  $x = 0.5$ .

Now adding  $[\text{C}_8\text{MIM-F13}][\text{Tf}_2\text{N}]$  to  $[\text{C}_{10}\text{MIM}][\text{Tf}_2\text{N}]$  at 35 mol%, the percolated nature of  $[\text{C}_{10}\text{MIM}][\text{Tf}_2\text{N}]$  largely remains with a small proportion of very small clusters also seen, while the fluorocarbon chains are shown to be arranged in very small aggregates (Figures 13a & 13b). However, at the equimolar composition, the situation is very similar indeed to that observed for  $[\text{C}_8\text{MIM}]_{0.5}[\text{C}_8\text{MIM-F13}]_{0.5}[\text{Tf}_2\text{N}]$ , where both chain types are contained in a range of aggregates of varying size up to  $n_a \approx 0.6-0.7$ , with a slightly greater proportion of larger hydrocarbon-based aggregates (Figures 13c & 13d). Once more, the fluorous component is effectively percolated at  $x = 0.85$  (Figure S44d). Then finally considering the mixtures  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F17}]_x[\text{Tf}_2\text{N}]$ , at  $x = 0.35$  the hydrocarbon chains remain percolated, while the fluorocarbon chains form a range of aggregates with a probability that falls off from a high value for small aggregates to close to zero by  $n_a \approx 0.5$  (Figures 14a & 14b). However, in the equimolar mixture it is evident that the



fluorocarbon chains are close to full percolation, while the hydrocarbon chains are forming a range of aggregates up to around  $n_a \approx 0.4$  (Figures 14c & 14d).

View Article Online  
DOI: 10.1039/D4FD00047A




Figure 13 Discrete probability distribution function,  $P(n_a)$ , for  $[C_{10}MIM]_{1-x}[C_8MIM-F_{13}]_x[Tf_2N]:[C_{10}MIM]$  chains at (a)  $x = 0.35$ ; (c)  $x = 0.5$ ;  $[C_8MIM-F_{13}]$  chains at (b)  $x = 0.35$ ; (d)  $x = 0.5$ .






Figure 14 Discrete probability distribution function,  $P(n_a)$ , for  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$ :  $[\text{C}_{10}\text{MIM}]$  chains at (a)  $x = 0.35$ ; (c)  $x = 0.5$ ;  $[\text{C}_{10}\text{MIM-F}_{17}]$  chains at (b)  $x = 0.35$ ; (d)  $x = 0.5$ .

To understand the aggregation behaviour better, pair radial distribution functions –  $g(r)$  – were examined considering interactions between the terminal carbon atoms of the alkyl chains (CT and CTF for the hydrogenated and fluorinated chains, respectively). These are displayed in Figure 15. For  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures (Figure 15a), the probability of finding a 'mixed' CT–CTF pair is the simple average of the probabilities of finding 'pure' CT–CT or CTF–CTF pairs, whereas this is not the case for all of the mixtures  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  (Figure 15b) and  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  (Figure 15c). At first sight and given the relatively inflexible nature of the perfluorocarbon segment, this may be thought to be attributable to the unequal chain lengths, but the fact that the observation persists for  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  (Figure 15d) suggests an alternative observation. This is that the  $[\text{C}_{10}\text{MIM-F}_{17}]^+$  fluorinated chains have a higher probability of being found in close contact with each other than do the alkyl chains on the other





component i.e. that there is some preferential aggregation of the fluororous chains when they are longer than C8.

View Article Online  
DOI: 10.1039/D4FD00047A




Figure 15. Five selected radial distribution functions (RDFs),  $g(r)$ , as a function of distance,  $r$ , for (a)  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$ ; (b)  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$ ; (c)  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  and d)  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$ , each showing data for the two pure components and the equimolar mixture. Orange lines represent pure ILs, and dark green lines represent a mole fraction of 0.50: RDFs between the terminal carbon atoms of  $[\text{C}_8\text{MIM}]$  or  $[\text{C}_{10}\text{MIM}]$ , CT; light orange lines represent pure ILs, and medium green lines represent a mole fraction of 0.50: RDFs between the terminal carbon atoms of  $[\text{C}_8\text{MIM-F}_{13}]$  or  $[\text{C}_{10}\text{MIM-F}_{17}]$ , CTF; light green lines represent RDFs between the terminal carbon atoms of  $[\text{C}_8\text{MIM}]$  or  $[\text{C}_{10}\text{MIM}]$  and the terminal carbon atoms of  $[\text{C}_8\text{MIM-F}_{13}]$  or  $[\text{C}_{10}\text{MIM-F}_{17}]$ , CT-CTF. Note the superposition between pure ILs and the mole fraction 0.50 for CT-CT and CTF-CTF RDFs.



## Discussion

It has been noted before that ILs with chain lengths greater than around C<sub>6</sub>,<sup>60-62, 64</sup> form nanostructured liquids with both polar (ionic) and non-polar (alkyl or fluoroalkyl) percolated domains. The question posed by the current study is what happens to this nanostructure when these ILs are mixed, and in particular what is the impact of both the alkyl and fluoroalkyl chain lengths on the liquid structure? The [C<sub>8</sub>MIM]<sub>1-x</sub>[C<sub>8</sub>MIM-F<sub>13</sub>]<sub>x</sub>[Tf<sub>2</sub>N] system, discussed above and in a previous publication,<sup>30</sup> showed little evidence of low-*q* scattering, rather a PNPP was observed for all compositions, with the lengthscale of the PNPP increasing only slightly with *x*, as fluoroalkyl chains are slightly longer than their alkyl equivalents. This is consistent with little or no triphasic behaviour *i.e.* little or no preferential separation of the alkyl and fluoroalkyl chains, consistent with the observed radial distribution function *g(r)* reproduced again in Figure 15a. This is supported by aggregate analysis showing a transition between an effectively percolated alkyl chain network and a percolated fluoroalkyl chain network as a function of composition (Figure 11). Thus, the alkyl and fluoroalkyl chains are generally well mixed in [C<sub>8</sub>MIM]<sub>1-x</sub>[C<sub>8</sub>MIM-F<sub>13</sub>]<sub>x</sub>[Tf<sub>2</sub>N] mixtures, consistent with the absence of low-*q* X-ray or neutron scattering.

When considering the new mixtures described here, it is first worth noting similarities between these and the [C<sub>8</sub>MIM]<sub>1-x</sub>[C<sub>8</sub>MIM-F<sub>13</sub>]<sub>x</sub>[Tf<sub>2</sub>N] system. Thus, the CP moves to lower *q* as the proportion of fluorinated IL increases in all mixtures, with the lengthscale associated with this peak changing from 4.7 to ~5.3 Å. As noted previously, this occurs because the CP contains information not only related to nearest neighbour anion-cation distances, as previously assumed, but crucially also on lateral chain-chain correlations. The lengthscale of these correlations is greater in systems containing fluoroalkyl rather than alkyl chains, as the fluorinated chains are bulkier (*i.e.* C–F bonds are longer than C–H bonds and F is larger than H).<sup>30</sup> In addition, the COP shows similar behaviour in all of these mixtures, moving to higher *q* with increasing proportion of the fluorinated IL, with associated changes in lengthscale from around 7.5 to 6.8 Å. As described before,<sup>30</sup> this is linked to the increased rigidity of the fluoroalkyl chains, which leads to the population of fewer gauche conformations in the fluoroalkyl chains (Figure S45). In turn, this



allows for more efficient packing of the ions and a corresponding tightening of the ion-ion distances in the polar network within the ILs.

View Article Online  
DOI: 10.1039/D4FD00047A

While there are then similarities in the medium- and wide-angle scattering for all of the mixtures under consideration, in the new mixtures, which contain C10 or C10-F17 chains, the scattering at lower  $q$  is quite different. Thus, while all three new mixture systems show a PNPP reflection when the mixtures are rich in either the alkyl or fluoroalkyl IL, at intermediate compositions there is significant low- $q$  scattering and a PNPP is not observed.

Considering first the  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  mixtures, it can be seen that increasing the length of the fluoroalkyl chain segment from C6 to C8 leads to changes in scattering (Figure 7 and Table 2) and the observation of aggregation in the MD simulations (Figures 12 and 15b). Figures 12a&b show that when 35 mol% of  $[\text{C}_{10}\text{MIM-F}_{17}][\text{Tf}_2\text{N}]$  is present in the mixture, there is no percolation of either alkyl or fluoroalkyl chains, rather the different chains form aggregates within the mixture over a range of sizes. This is consistent with the observation of significant low- $q$  scattering that was modelled as spherical scattering objects with radii of around 8 Å (the approximate length of a  $\text{C}_8\text{F}_{17}$  chain), assuming that this is a time averaged picture where small aggregates are most common. However, it is important to note that these aggregates will be composed of ILs in bilayer organisation, but the wide range of sizes will reduce the effective correlation length between them, effectively broadening the PNPP and making it difficult to observe experimentally. This is reflected to an extent in the structure factor calculations, for while a PNPP is observed therein, the broader peaks are consistent with the fact that they were better modelled by the use of two rather than one Gaussian function (see SI for details). Note also that the  $S(q)$  calculations did not attempt to incorporate features from low- $q$  scattering.

It is important to note that these mixtures still contain polar (ions) and non-polar (chains) domains, and the scattering and MD simulation data can be rationalised in terms of the degree of homogeneity of the non-polar domains. Thus, in  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures, the alkyl and fluoroalkyl chains are well mixed in the non-polar domains leading to a relatively homogenous



non-polar sub phase. When  $[\text{C}_{10}\text{MIM-F}_{17}]^+$  ions are introduced in  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  mixtures, there is some segregation of the alkyl and fluoroalkyl chains, which leads to the observation of small aggregates that indicate inhomogeneity in the non-polar sub phase, although evidently this is insufficient to promote demixing. In terms of scattering, the non-polar domains in the IL mixtures no longer approximate a uniform SLD, rather they contain regions of differing SLDs making the observation of a simple, discrete PNPP difficult. In a simple model, this can be thought of as introducing 'patchiness' or 'stripiness' to the non-polar parts of the mixtures (Figure 16). With increasing  $x$  in  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  the fluoroalkyl chains percolate by  $x = 0.5$  (Figure 12d) and we see a PNPP begin to emerge (Figure 7) at larger  $x$ , although the presence of small alkyl chain aggregates means that low- $q$  scattering persists until the mixture is almost purely composed of  $[\text{C}_{10}\text{MIM-F}_{17}][\text{Tf}_2\text{N}]$ .

This behaviour is reflected also in the radial distribution functions and Figure 15b shows that while the probability of finding self interactions between the terminal carbons of hydrogenated chains (CT-CT) and the probability of finding self interactions between the terminal carbons of fluorinated chains (CTF-CTF) appear essentially independent of composition, the probability of finding interactions between the terminal carbons of chains of different types (CT-CTF) is not linear with composition (as it is in the  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures) and remains appreciably lower than the probability of CTF-CTF interactions. This is consistent with the observations from both scattering data and aggregation analysis.

View Article Online  
DOI: 10.1039/D4FD00047A






Figure 16. Cartoon representations of nanoscale organisation proposed in the liquid structure of a) pure  $[\text{C}_8\text{MIM}][\text{Tf}_2\text{N}]$  and b) a  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  mixture of intermediate composition. Blue areas represent the polar (ionic) network present in all ILs. a) Grey areas represent well-mixed alkyl and fluoroalkyl chains approximating a uniform SLD. b) Striped areas represent greater segregation between alkyl and fluoroalkyl chains leading to a non-uniform SLD in the non-polar network.

Consideration of the various data for the  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  mixtures (Figure 5, Table 1, Figure 14, Figure 15d) shows that the behaviour can be rationalised in the same way. Aggregate analysis for this system (Figure 14) shows that the percolated sub-phase formed from the C10 alkyl chains is more resistant to disruption than that formed by the C8 alkyl chains (Figure 12) as the C10 chain network remains percolated at  $x = 0.35$ , whereas it had disappeared at this composition in the  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_{10}\text{MIM-F}_{17}]_x[\text{Tf}_2\text{N}]$  system. Nonetheless, significant  $\text{C}_{10}\text{H}_4\text{F}_{17}$  chain aggregates are observed at this composition (Figure 14b). This system can also be described as having a non-uniform distribution of the two chain types within a non-polar network leading to low- $q$  scattering that can be described by spherical aggregates with average diameters of around two  $\text{C}_8\text{F}_{17}$  chains (Table 1).

The final mixture system to consider is  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$ . If the above effects were simply driven by preferential interactions between the longer fluoroalkyl chains we would perhaps expect the  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  system to behave similarly to  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$ . However, things are not so simple. Thus, there is significant low- $q$  scattering from the  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures (Figure 8, Table 3) and the aggregate analyses also differ



(Figure 13). As above, the C10 alkyl chains remain percolated at  $x = 0.35$ , but small C<sub>6</sub>F<sub>13</sub> aggregates are seen. By  $x = 0.5$  neither the alkyl nor fluoroalkyl chains are percolated, instead both form predominantly small aggregates, but with a significant distribution of larger aggregates, reminiscent of the results from [C<sub>8</sub>MIM]<sub>0.65</sub>[C<sub>10</sub>MIM-F<sub>17</sub>]<sub>0.35</sub>[Tf<sub>2</sub>N] (Figure 12a&b). This is consistent with the radial distribution functions (Figure 15), which do not show such a strong preference for CTF-CTF interactions over their CT-CT equivalents. As such, these data are similar to those for the [C<sub>8</sub>MIM]<sub>1-x</sub>[C<sub>8</sub>MIM-F<sub>13</sub>]<sub>x</sub>[Tf<sub>2</sub>N] mixtures.

The contrast between this system and the [C<sub>8</sub>MIM]<sub>1-x</sub>[C<sub>8</sub>MIM-F<sub>13</sub>]<sub>x</sub>[Tf<sub>2</sub>N] mixtures can then be rationalised by the stronger tendency of the C10 alkyl chains to associate with each other than the C8 alkyl chains. This leads to a greater resistance to disruption of the C10 non-polar network and, when disrupted, a greater tendency for C10 aggregates to form. As with the systems above, this leads to a non-uniform distribution of chains in the non-polar sub phase, leading to 'stripiness' in the SLDs and thus to low- $q$  scattering arising from these aggregates, which are time averaged to approximate spherical scattering objects.

## Conclusions

Previous studies of the mixtures [C<sub>8</sub>MIM]<sub>1-x</sub>[C<sub>8</sub>MIM-F<sub>13</sub>]<sub>x</sub>[Tf<sub>2</sub>N] had shown a smooth evolution of liquid structure across the range of compositions, as evidenced by X-ray and neutron scattering data. When at least one of the chain lengths increases to contain ten carbon atoms, this situation changes. While reflections in the mid- and low-angle show systematic changes in the COP (decreases in size with  $x$ ) and the CP (increases in size with  $x$ ), respectively, for all mixtures studied, at low angle the PNPP is lost, with this region being dominated by scattering from small scattering objects, fitted as spheres. Analysis shows that these are composed of fluorocarbon chain segments dispersed in a hydrocarbon background, which matches data from MD simulations showing, in particular in the [C<sub>8</sub>MIM]<sub>1-x</sub>[C<sub>10</sub>MIM-F<sub>17</sub>]<sub>x</sub>[Tf<sub>2</sub>N] and [C<sub>10</sub>MIM]<sub>1-x</sub>[C<sub>8</sub>MIM-F<sub>13</sub>]<sub>x</sub>[Tf<sub>2</sub>N] mixtures, formation of a large number of aggregates across a range of sizes. While all of the pure components are percolated, this is disrupted by the addition of the other component and the re-formation of a percolated network of the added chain depends upon the combination of chain lengths. Thus, a



well-percolated mixture of fluorocarbon chains is found at  $x = 0.5$  for  $[\text{C}_{10}\text{MIM-F}_{17}][\text{Tf}_2\text{N}]$  mixed with either  $[\text{C}_8\text{MIM}][\text{Tf}_2\text{N}]$  or  $[\text{C}_{10}\text{MIM}][\text{Tf}_2\text{N}]$ . With  $[\text{C}_8\text{MIM-F}_{13}][\text{Tf}_2\text{N}]$ , percolation is fairly well developed in mixtures with  $[\text{C}_8\text{MIM}][\text{Tf}_2\text{N}]$  at  $x = 0.5$ , whereas in  $[\text{C}_{10}\text{MIM}][\text{Tf}_2\text{N}]$  it is not re-established until  $x = 0.85$ . However, while  $[\text{C}_{10}\text{MIM-F}_{17}][\text{Tf}_2\text{N}]$  disrupts the percolation in  $[\text{C}_8\text{MIM}][\text{Tf}_2\text{N}]$  by  $x = 0.35$ , it does not do so in  $[\text{C}_{10}\text{MIM}][\text{Tf}_2\text{N}]$  until  $x = 0.5$ .

Taken together, the data can be understood with reference to an extensive computational study of hydrocarbon/fluorocarbon mixing by Pollice and Chen.<sup>54</sup> They showed that inter-chain HH interactions, while weak, have a close to ideal geometry for interaction, while inter-chain FF interactions, while stronger, have a non-ideal geometry. Inter-chain HF interactions also have a very non-ideal geometry, but are intermediate in strength. As such, they found that one HH and one FF interaction were generally preferred over two HF interactions. Clearly this will be influenced by the chain length and so can start to account for the observed immiscibility of longer chain hydrocarbons and fluorocarbons around room temperature (shorter chain components are immiscible at sub-ambient temperatures).

Of course, intermolecular interactions in the IL mixtures under consideration here also have a strong electrostatic component and so the potential immiscibility of the different ILs is appreciably modulated compared to pure hydrocarbons and fluorocarbons. However, the way in which percolation is disrupted and the observation by both scattering and MD simulation of aggregates in the mixtures shows that the inter-chain interactions have a clear influence on the behaviour. Returning to the physical properties of the mixtures, surface-tension measurements on  $[\text{C}_{10}\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures highlight that the  $[\text{C}_8\text{MIM-F}_{13}]^+$  ions are clearly enriched at the gas-liquid interface. However, there is no evidence of a CMC in the system – it is different from the behaviour of a surfactant in water. As we have seen before for alkyl/fluoroalkyl IL mixtures, the viscosity of the mixtures is lower than would be expected based on simple Arrhenius-type mixing. This is even more pronounced than the changes seen for  $[\text{C}_8\text{MIM}]_{1-x}[\text{C}_8\text{MIM-F}_{13}]_x[\text{Tf}_2\text{N}]$  mixtures and suggests that addition of the fluorocarbon-functionalised cations disrupt the IL structure in such



a way as to allow for increased ion mobility. Such understanding is crucial as predictive capability in mixture properties continues to evolve.

View Article Online  
DOI: 10.1039/D4FD00047A

### Experimental and computational methods

All experimental and computational methodologies are described in the SI.

### Conflict of Interest

The authors declare no conflicts of interest.

### Supporting Information

Experimental and computational details. Key spectroscopic data.

### Authors Contribution Statement

The study was conceived by JMS, DWB and NSE. Grant funding was secured by MLC, KGM, JMS and DWB, while the beamtime allocation was secured by NSE, SER, JMS and DWB. ILs were prepared by NSE and EVS, while the perdeuterated precursors were prepared by SY. Experimental work was undertaken by NSE, PDL, EVS, JL and JMS and simulations were carried out by KS. Data interpretation was done by NSE, EVS, KS, BD, JNCL, MLC, KGM, JMS and DWB and the manuscript was written and edited by NSE, KS, NM, SER, BD, JNCL, MLC, KGM, JMS and DWB.

### Acknowledgements

The authors thank the EPSRC (NSE: EP/T031174/1, PDL: EP/T03114X/1 & EP/T021675/1) and the University of York Wild Scholars Fund (JL) for support, the ILL for beamtime (9-11-2147) and the ISIS Deuteration Facility (SY) for some deuterated precursors. KS and JNCL thank Fundação para a Ciência e Tecnologia, FCT/MCTES (Portugal) for financial support through CEEC contract (IST-ID/100/2018 to KS), and through projects UIDB/00100/2020 and LA/P/0056/2020.





## References

1. H. Niedermeyer, J. P. Hallett, I. J. Villar-Garcia, P. A. Hunt and T. Welton, *Chem. Soc. Rev.*, 2012, **41**, 7780-7802.   
View Article Online  
DOI: 10.1039/D4FD00047A
2. G. Chatel, J. F. B. Pereira, V. Debbeti, H. Wang and R. D. Rogers, *Green Chem.*, 2014, **16**, 2051-2083.
3. A. Riisager, R. Fehrmann, M. Haumann and P. Wasserscheid, *Eur. J. Inorg. Chem.*, 2006, 695-706.
4. A. Riisager, R. Fehrmann, M. Haumann and P. Wasserscheid, *Top. Catal.*, 2006, **40**, 91-102.
5. T. Selvam, A. Machoke and W. Schwieger, *Appl. Catal. A*, 2012, **445**, 92-101.
6. M. Abai, M. P. Atkins, A. Hassan, J. D. Holbrey, Y. Kuah, P. Nockemann, A. A. Oliferenko, N. V. Plechkova, S. Rafeen, A. A. Rahman, R. Ramli, S. M. Shariff, K. R. Seddon, G. Srinivasan and Y. R. Zou, *Dalton Trans.*, 2015, **44**, 8617-8624.
7. J. M. Marinkovic, A. Riisager, R. Franke, P. Wasserscheid and M. Haumann, *Ind. Eng. Chem. Res.*, 2019, **58**, 2409-2420.
8. L. Offner-Marko, A. Bordet, G. Moos, S. Tricard, S. Rengshausen, B. Chaudret, K. L. Luska and W. Leitner, *Angew. Chem. Int. Ed.*, 2018, **57**, 12721-12726.
9. D. Geier, P. Schrnitz, J. Walkowiak, W. Leitner and G. Francio, *ACS Catal.*, 2018, **8**, 3297-3303.
10. J. Brunig, Z. Csendes, S. Weber, N. Gorgas, R. W. Bittner, A. Limbeck, K. Bica, H. Hoffmann and K. Kirchner, *ACS Catal.*, 2018, **8**, 1048-1051.
11. S. More, S. Jadhav, R. Salunkhe and A. Kumbhar, *Molecular Catalysis*, 2017, **442**, 126-132.
12. S. G. Khokarale, E. J. Garcia-Suarez, R. Fehrmann and A. Riisager, *Chemcatchem*, 2017, **9**, 1824-1829.
13. J. Zhao, S. C. Gu, X. L. Xu, T. T. Zhang, Y. Yu, X. X. Di, J. Ni, Z. Y. Pan and X. N. Li, *Catal. Sci. Technol.*, 2016, **6**, 3263-3270.
14. K. L. Luska, P. Migowski, S. El Sayed and W. Leitner, *ACS Sustain. Chem. Eng.*, 2016, **4**, 6186-6192.
15. O. D. Pavel, P. Goodrich, L. Cristian, S. M. Coman, V. I. Parvulescu and C. Hardacre, *Catal. Sci. Technol.*, 2015, **5**, 2696-2704.
16. K. L. Luska, P. Migowski, S. El Sayed and W. Leitner, *Angew. Chem. Int. Ed.*, 2015, **54**, 15750-15755.
17. C. Waring, P. A. J. Bagot, J. M. Slattery, M. L. Costen and K. G. McKendrick, *J. Phys. Chem. A*, 2010, **114**, 4896-4904.
18. C. Waring, P. A. J. Bagot, J. M. Slattery, M. L. Costen and K. G. McKendrick, *J. Phys. Chem. Lett.*, 2010, **1**, 429-433.
19. M. A. Tesa-Serrate, B. C. Marshall, E. J. Smoll, S. M. Purcell, M. L. Costen, J. M. Slattery, T. K. Minton and K. G. McKendrick, *J. Phys. Chem. C*, 2015, **119**, 5491-5505.
20. S. M. Purcell, M. A. Tesa-Serrate, B. C. Marshall, D. W. Bruce, L. D'Andrea, M. L. Costen, J. M. Slattery, E. J. Smoll, T. K. Minton and K. G. McKendrick, *Langmuir*, 2016, **32**, 9938-9949.
21. M. A. Tesa-Serrate, E. J. Smoll, L. D'Andrea, S. M. Purcell, M. L. Costen, D. W. Bruce, J. M. Slattery, T. K. Minton and K. G. McKendrick, *J. Phys. Chem. C*, 2016, **120**, 27369-27379.



22. D. W. Bruce, C. P. Cabry, J. N. C. Lopes, M. L. Costen, L. D'Andrea, I. Grillo, B. C. Marshall, K. G. McKendrick, T. K. Minton, S. M. Purcell, S. Rogers, J. M. Slattery, K. Shimizu, E. Smoll and M. A. Tesa-Serrate, *J. Phys. Chem. B*, 2017, **121**, 6002-6020. View Article Online  
DOI: 10.1039/D4FD00047A
23. S. M. Purcell, P. D. Lane, L. D'Andrea, N. S. Elstone, D. W. Bruce, J. M. Slattery, E. J. Smoll, S. J. Greaves, M. L. Costen, T. K. Minton and K. G. McKendrick, *J. Phys. Chem. B*, 2022, **126**, 1962-1979.
24. B. H. Wu, J. M. Zhang, T. K. Minton, K. G. McKendrick, J. M. Slattery, S. Yockel and G. C. Schatz, *J. Phys. Chem. C*, 2010, **114**, 4015-4027.
25. C. P. Cabry, L. D'Andrea, K. Shimizu, I. Grillo, P. X. Li, S. Rogers, D. W. Bruce, J. N. C. Lopes and J. M. Slattery, *Faraday Discuss.*, 2018, **206**, 265-289.
26. E. J. Smoll, M. A. Tesa-Serrate, S. M. Purcell, L. D'Andrea, D. W. Bruce, J. M. Slattery, M. L. Costen, T. K. Minton and K. G. McKendrick, *Faraday Discuss.*, 2018, **206**, 497-522.
27. E. J. Smoll, S. M. Purcell, L. D'Andrea, J. M. Slattery, D. W. Bruce, M. L. Costen, K. G. McKendrick and T. K. Minton, *J. Phys. Chem. Lett.*, 2019, **10**, 156-163.
28. E. J. Smoll, X. M. Chen, L. M. Hall, L. D'Andrea, J. M. Slattery and T. K. Minton, *J. Phys. Chem. C*, 2020, **124**, 382-397.
29. C. P. Cabry, L. D'Andrea, N. S. Elstone, S. Kirchhecker, A. Riccobono, I. Khazal, P. X. Li, S. E. Rogers, D. W. Bruce and J. M. Slattery, *Phys. Chem. Chem. Phys.*, 2022, **24**, 15811-15823.
30. N. S. Elstone, K. Shimizu, E. V. Shaw, P. D. Lane, L. D'Andrea, B. Demé, N. Mahmoudi, S. E. Rogers, S. Youngs, M. L. Costen, K. G. McKendrick, J. C. Lopes, D. W. Bruce and J. M. Slattery, *J. Phys. Chem. B*, 2023, **127**, 7394-7407.
31. C. C. Weber, N. J. Brooks, F. Castiglione, M. Mauri, R. Simonutti, A. Mele and T. Welton, *Phys. Chem. Chem. Phys.*, 2019, **21**, 5999-6010.
32. D. Pontoni, M. DiMichiel and M. Deutsch, *J. Mol. Liq.*, 2020, **300**, 112280.
33. D. Pontoni, M. DiMichiel and M. Deutsch, *J. Mol. Liq.*, 2021, **338**, 11.
34. R. Costa, C. M. Pereira, A. F. Silva, P. F. Brevet and E. Benichou, *J. Phys. Chem. B*, 2020, **124**, 3954-3961.
35. M. E. Di Pietro, F. Castiglione and A. Mele, *J. Phys. Chem. B*, 2020, **124**, 2879-2891.
36. B. S. J. Heller, C. Kolbeck, I. Niedermaier, S. Dommer, J. Schatz, P. Hunt, F. Maier and H.-P. Steinrück, *ChemPhysChem*, 2018, **19**, 1733-1745.
37. F. Wu, W. V. Karunaratne and C. J. Margulis, *J. Phys. Chem. C*, 2019, **123**, 4914-4925.
38. T. Cosby, U. Kapoor, J. K. Shah and J. Sangoro, *J. Phys. Chem. Lett.*, 2019, **10**, 6274-6280.
39. I. J. Villar-Garcia, S. Fearn, N. L. Ismail, A. J. S. McIntosh and K. R. J. Lovelock, *Chem. Commun.*, 2015, **51**, 5367-5370.
40. M. Chakraborty, T. Ahmed, R. S. Dhale, D. Majhi and M. Sarkar, *J. Phys. Chem. B*, 2018, **122**, 12114-12130.
41. L. F. Lepre, M. C. Gomes, A. A. H. Padua, R. A. Ando and M. C. C. Ribeiro, *J. Phys. Chem. B*, 2019, **123**, 6579-6587.



42. B. Golub, D. Ondo, V. Overbeck, R. Ludwig and D. Paschek, *Phys. Chem. Chem. Phys.*, 2022, **24**, 14740-14750.
43. Y. Q. Chen, S. L. Ma, Y. Lei, X. D. Liang, X. Y. Liu, G. M. Kontogeorgis and R. Gani, *AIChE J.*, 2024, **70**, 13. View Article Online  
DOI: 10.1002/aiche.1747A
44. Z. W. Zhai, U. Paap, A. Gezmis, F. Maier, H. P. Steinrueck and T. M. Koller, *J. Mol. Liq.*, 2023, **386**, 9.
45. O. Holloczki, M. Macchiagodena, H. Weber, M. Thomas, M. Brehm, A. Stark, O. Russina, A. Triolo and B. Kirchner, *ChemPhysChem*, 2015, **16**, 3325-3333.
46. B. S. J. Heller, M. Lexow, F. Greco, S. Shin, G. Partl, F. Maier and H. P. Steinrueck, *Chem. Eur. J.*, 2020, **26**, 1117-1126.
47. B. S. J. Heller, U. Paap, F. Maier and H. P. Steinrueck, *J. Mol. Liq.*, 2020, **305**, 9.
48. T. M. Koller, F. D. Lenahan, P. S. Schmidt, T. Klein, J. Mehler, F. Maier, M. H. Rausch, P. Wasserscheid, H. P. Steinrueck and A. P. Fröba, *Int. J. Thermophys.*, 2020, **41**, 24.
49. D. Hemmeter, D. Kremitzl, P. S. Schulz, P. Wasserscheid, F. Maier and H. P. Steinrueck, *Chem. Eur. J.*, 2022, **29**, e202204023.
50. D. Hemmeter, U. Paap, N. Wellenhofer, A. Gezmis, D. Kremitzl, P. Wasserscheid, H. P. Steinrueck and F. Maier, *ChemPhysChem*, 2023, **24**, e202300612.
51. A. Riccobono, R. R. Parker, A. C. Whitwood, J. M. Slattey, D. W. Bruce, I. Pibiri and A. Pace, *Chem. Commun.*, 2018, **54**, 9965-9968.
52. A. Riccobono, G. Lazzara, S. E. Rogers, I. Pibiri, A. Pace, J. M. Slattey and D. W. Bruce, *J. Mol. Liq.*, 2021, **321**, 114758.
53. A. Riccobono, A. C. Whitwood, R. R. Parker, S. Hart, A. Pace, J. M. Slattey, I. Pibiri and D. W. Bruce, *CrystEngComm*, 2022, **24**, 7852-7860.
54. R. Pollice and P. Chen, *J. Am. Chem. Soc.*, 2019, **141**, 3489-3506.
55. S. Arrhenius, *Z. Phys. Chem.*, 1887, **1**, 285.
56. K. Shimizu, C. E. S. Bernardes and J. N. C. Lopes, *J. Phys. Chem. B*, 2014, **118**, 567-576.
57. J. N. C. Lopes, M. F. C. Gomes and A. A. H. Padua, *J. Phys. Chem. B*, 2006, **110**, 16816-16818.
58. H. V. R. Annapureddy, H. K. Kashyap, P. M. De Biase and C. J. Margulis, *J. Phys. Chem. B*, 2010, **114**, 16838-16846.
59. H. K. Kashyap, J. J. Hettige, H. V. R. Annapureddy and C. J. Margulis, *Chem. Commun.*, 2012, **48**, 5103-5105.
60. A. Triolo, O. Russina, H. J. Bleif and E. Di Cola, *J. Phys. Chem. B*, 2007, **111**, 4641-4644.
61. R. Hayes, G. G. Warr and R. Atkin, *Chem. Rev.*, 2015, **115**, 6357-6426.
62. Y. L. Wang, B. Li, S. Sarman, F. Mocci, Z. Y. Lu, J. Y. Yuan, A. Laaksonen and M. D. Fayer, *Chem. Rev.*, 2020, **120**, 5798-5877.
63. M. Teubner and R. Strey, *J. Chem. Phys.*, 1987, **87**, 3195-3200.
64. J. N. A. C. Lopes and A. A. H. Padua, *J. Phys. Chem. B*, 2006, **110**, 3330-3335.

