



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/214998/>

Version: Accepted Version

Article:

Liu, Y., Bowyer, F.T., Zhu, M. et al. (2024) Marine redox and nutrient dynamics linked to the Cambrian radiation of animals. *Geology*, 52 (9). pp. 729-734. ISSN: 0091-7613

<https://doi.org/10.1130/G52220.1>

© 2024 Geological Society of America. This is an author produced version of an article accepted for publication in *Geology*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Marine redox and nutrient dynamics linked to the Cambrian radiation of animals

Yang Liu^{1,2*}, Fred T. Bowyer², Maoyan Zhu^{3,4}, Yijun Xiong², Tianchen He⁵,
Meiling Han³, Xuan Tang⁶, Jinchuan Zhang⁶, Simon W. Poulton²

¹*College of Environmental Science and Engineering, Dalian Maritime University, Dalian 116026, China*

²*School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK*

³*State Key Laboratory of Palaeobiology and Stratigraphy, Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences, Nanjing 210008, China*

⁴*College of Earth and Planetary Sciences, University of Chinese Academy of Sciences, Beijing 100049, China*

⁵*College of Oceanography, Hohai University, Nanjing 210024, China*

⁶*School of Energy Resource, China University of Geosciences (Beijing), Beijing 100083, China*

ABSTRACT

The early Cambrian witnessed an increase in metazoan ecosystem complexity, likely linked to enhanced oxygen and nutrient availability. However, while improved stratigraphic and geochemical records suggest that the Cambrian explosion occurred under highly dynamic redox conditions, mechanistic links to nutrient availability and early Cambrian evolutionary innovations are poorly constrained. Here, we report paleoredox and nutrient data for two drill cores documenting late Cambrian Stage 2 to Stage 3 (~522 to 514 Ma) strata from the Yangtze Block, South China. The development of water column euxinia during Cambrian Stage 2 led to extensive recycling of P, fueling elevated primary production and hence an increase in atmospheric and shallow marine oxygen concentrations. The resulting expansion of oxygenated shelf area promoted sedimentary P retention, and in combination with a diminished supply of P from upwelling, drove a transition to oligotrophic shallow marine conditions during Cambrian Stage 3. Reduced primary production and limited water column oxygen consumption allowed for the stabilization of oxygenated continental shelf habitats that supported a burgeoning biotic complexity.

INTRODUCTION

The early Cambrian (Fortunian–end Stage 3; ca. 538.8–514 Ma) witnessed the abrupt appearance and rapid diversification of most modern animal phyla during the ‘Cambrian Explosion’ (Knoll and Carroll, 1999; Erwin et al., 2011; Wood et al., 2019). A wide range of potential triggers for the Cambrian Explosion have been proposed, including intrinsic developmental and ecological innovations (Erwin et al., 2011), and extrinsic changes (particularly oxygen levels) to environmental habitability (Smith and Harper, 2013; Knoll and Carroll, 1999; Lenton et al., 2014). The first animals likely had low oxygen demands (Mills et al., 2014), however the larger and more metabolically active skeletonizing animals that appeared during the early Cambrian would have required considerably higher oxygen levels (Sperling et al., 2013). Indeed, potential links between pulses of shallow-ocean oxygenation and biodiversity have been recorded on the early Cambrian Siberian platform (He et al.,

2019; Zhuravlev et al., 2022) and in South China (Chen et al., 2015; Li et al., 2017).

The availability of the bio-limiting nutrients phosphorus (P) and nitrogen (N) likely also exerted an important control on environmental habitability, with P generally considered to be the ultimate limiting nutrient for primary production on geological timescales (Tyrell, 1999). These nutrients control the degree of regional oxygen production and consumption, as well fueling more complex life (Brocks et al., 2017). Depositional redox conditions also control P recycling, resulting in feedbacks that dictate the evolution of regional marine redox state (Schobben et al., 2020; Bowyer et al., 2023). However, despite their fundamental importance for ecosystem habitability and animal evolution, integrated records of redox and P cycling in the early Cambrian are highly limited (see Creveling et al., 2014).

Here, we report a geochemical study of shelf and slope successions from the Yangtze Block, South China. We utilize Fe speciation, redox sensitive trace element systematics and P phase partitioning to evaluate contemporaneous marine redox conditions and controls on primary productivity. These data allow a detailed assessment of redox and nutrient controls on the major biological events that occurred during the early Cambrian.

GEOLOGIC SETTING AND MATERIALS

We investigate two drill cores from the Yangtze Block, deposited in an open-marine continental shelf (ZK 402) and slope (CY 1) setting (see Supplemental Material¹ for full details of the geologic setting and methods). Both cores preserve a relatively continuous record spanning the latter part of Cambrian Stage 2 through Cambrian Stage 3, offering a prime opportunity to evaluate biogeochemical changes across an interval that captures a major evolutionary event.

RESULTS AND DISCUSSION

Oceanic redox dynamics during the early Cambrian

We employ a combination of Fe speciation and redox-sensitive trace metal (RSTM; Mo, U, V) systematics (see Text S3) to provide a robust framework for interpreting depositional

redox conditions (e.g., Tribovillard et al., 2006; Poulton, 2021). In modern and ancient marine sediments, a highly reactive (Fe_{HR}) to total Fe (Fe_{T}) ratio of >0.38 provides an indication of anoxia, while $\text{Fe}_{\text{HR}}/\text{Fe}_{\text{T}} < 0.22$ suggests oxic conditions, with values between considered equivocal. When anoxia is indicated, the extent of pyritization (Fe_{Py}) of the Fe_{HR} pool is used to differentiate between ferruginous ($\text{Fe}_{\text{Py}}/\text{Fe}_{\text{HR}} < 0.6$) or euxinic ($\text{Fe}_{\text{Py}}/\text{Fe}_{\text{HR}} > 0.8$) conditions (Poulton, 2021). Generally, RSTMs tend to be less soluble under reducing conditions, leading to authigenic enrichments in the sediment, with Mo in particular requiring the presence of dissolved sulfide (Tribovillard et al., 2006).

During late Cambrian Stage 2 to early Stage 3 (interval I), the development of persistent euxinia at both sites is suggested by a combination of high $\text{Fe}_{\text{HR}}/\text{Fe}_{\text{T}}$ (>0.7) and high $\text{Fe}_{\text{Py}}/\text{Fe}_{\text{HR}}$ ratios (0.69 ± 0.11 for ZK 402 and 0.92 ± 0.04 for CY 1; Fig. 1; Table S1). This interpretation is supported by high enrichment factors (EFs; see Text S2) for the RSTMs and high $\text{Mo}_{\text{EF}}/\text{U}_{\text{EF}}$ ratios. During the initial stages of interval II (middle to late Stage 3), a decline in $\text{Fe}_{\text{HR}}/\text{Fe}_{\text{T}}$ ratios, EF values and $\text{Mo}_{\text{EF}}/\text{U}_{\text{EF}}$ ratios occurs in both drill cores (Fig. 1). Subsequently, on the shelf (ZK 402), $\text{Fe}_{\text{HR}}/\text{Fe}_{\text{T}}$ ratios and RSTM enrichments remain persistently low. This suggests a transition to dominantly oxic conditions, although occasional minor increases in $\text{Fe}_{\text{HR}}/\text{Fe}_{\text{T}}$ ratios, EF values and $\text{Mo}_{\text{EF}}/\text{U}_{\text{EF}}$ ratios may indicate transient, short-lived anoxia. On the slope (CY 1), despite decreasing trends in redox proxy data from intervals I to II, values remain elevated, indicating continued euxinia, perhaps punctuated by episodic ferruginous intervals (lower $\text{Mo}_{\text{EF}}/\text{U}_{\text{EF}}$ ratios; Fig. 1).

A compilation of global Fe speciation data (Text S4; Table S2) provides further insight into spatio-temporal variability in marine redox state (Fig. 2A). While the majority of data are from South China, a stratified early Cambrian ocean is apparent, with oxic surface waters overlying ferruginous deeper waters (with occasional euxinia), prior to the latter part of Stage 2 (Fig. 2). Euxinia then became common in outer shelf to slope settings, before extending into shallow shelf settings at the Stage 2-Stage 3 boundary. A clear transition to expanded shelf oxygenation then occurred, while deeper waters remained largely anoxic during Stage 3. The local marine redox state inferred from Fe speciation data is consistent with estimates of global marine redox from U and Mo isotopes (Fig. 2). Carbonates from South China,

103 Siberia and Morocco in the latest Fortunian and Stage 2 document negative $\delta^{238}\text{U}$
104 excursions (Wei et al., 2018; Dahl et al., 2019), suggesting episodic expansions of seafloor
105 anoxia during these intervals. By contrast, consistently higher $\delta^{238}\text{U}$ data recorded in
106 carbonates from Siberia and shales from South China in Stage 3 (Dahl et al., 2019; Wei et
107 al., 2020) provide evidence for more widespread oxygenation. Furthermore, modern-like
108 $\delta^{98}\text{Mo}$ signatures in Cambrian Stage 3 also suggest a major expansion of oxic seafloor
109 (Chen et al., 2015).

110

111 **Marine eutrophication and animal diversification**

112 High total organic carbon (TOC) and total P (P_{tot}) concentrations recorded in the lower
113 part of interval I in both drill cores (Fig. 3) potentially indicate high marine primary productivity
114 and nutrient levels. The greenhouse climate of the early Cambrian (Hearing et al., 2018),
115 likely drove enhanced chemical weathering (Peters and Gaines, 2012; Lipp et al., 2021) and
116 hence a high terrigenous nutrient influx. Alternatively, or in addition, high nutrient levels could
117 have occurred via upwelling of nutrient-replete deep ocean waters (e.g., Behrenfeld et al.,
118 2006; Bowyer et al., 2017). Upwelling can be evaluated using Co (ppm)×Mn (wt%) values,
119 whereby upwelling systems are characterized by low sedimentary Co and Mn
120 concentrations due to their depletion in upwelled deep-ocean waters, in addition to recycling
121 of Mn back to the water column under reducing conditions. However, restricted basins are
122 characterized by relatively high Co and Mn concentrations due to relatively high riverine
123 inputs and limited water exchange with the open ocean (Sweere et al., 2016).

124 The low Co×Mn values in the lower part of interval I are similar to those of modern
125 upwelling systems (<0.4 ppm%; Sweere et al., 2016), suggesting strong upwelling (Text S5).
126 This internal marine nutrient supply was augmented by nutrient recycling from the sediment,
127 which we evaluate via the phase partitioning of P. This approach quantifies four
128 operationally-defined P pools (Thompson et al., 2019), including Fe-bound P (P_{Fe}),
129 authigenic carbonate fluorapatite, biogenic apatite and CaCO_3 -bound P (P_{auth}), detrital
130 apatite (P_{det}), and organic-bound P (P_{org}). Reactive P (P_{reac}) represents potentially mobile P
131 during transport and early diagenesis, and is calculated as the sum of P_{Fe} , P_{auth} and P_{org} .

132 P_{org} is preferentially released from organic matter (C_{org}) via anaerobic remineralization,
133 resulting in C_{org}/P_{org} ratios above the Redfield ratio of 106/1 (Ingall et al., 1993). In addition,
134 P_{Fe} may be released during reductive dissolution of Fe (oxyhydr)oxide minerals (Slomp et
135 al., 1996). A proportion of this P may undergo 'sink switching' to authigenic phases in the
136 sediment (Slomp et al., 1996). However, P may also be recycled back to the water column,
137 particularly under sulfidic conditions, potentially driving a positive productivity feedback (Van
138 Cappellen and Ingall, 1996).

139 In samples from interval I, molar C_{org}/P_{org} and C_{org}/P_{reac} ratios are significantly greater than
140 the Redfield ratio of 106/1 (Fig. 3), demonstrating release of P from organic matter (C_{org}/P_{org})
141 and recycling back to the water column (C_{org}/P_{reac}). This recycling occurs alongside high P_{tot}
142 contents relative to average shale (0.009; Turekian and Wedepohl, 1961), suggesting
143 particularly high levels of bioavailable phosphate in the overlying water column. Expanded
144 euxinia (Fig. 2) would have resulted in a high degree P recycling in continental margin
145 settings, promoting a strong positive productivity feedback following upwelling to the surface
146 ocean (Van Cappellen and Ingall, 1996), and providing a food supply for early mobile
147 animals (Fig. 4). Furthermore, increased TOC burial in shelf and slope environments as a
148 consequence of enhanced primary productivity likely contributed to gradual atmospheric and
149 shallow ocean oxygenation (He et al., 2019). Hence, increased oxygen availability and food
150 supply in the shallow ocean drove the major evolutionary event recorded by the Chengjiang
151 and Qingjiang lagerstätte, which occur coincident with the first appearance of mineralized
152 trilobites in Cambrian Stage 3 (Fig. 2; Sun et al., 2022). The gradual decline in TOC and P
153 in the upper part of interval I may be related to the development of weakly euxinic conditions
154 and weakened upwelling, leading to a decrease in bioavailable P and hence lower TOC
155 burial.

156

157 **Oligotrophic stabilization of shelf oxygenation**

158 Total P (as P_{tot} and P_{tot}/Al ratios) and TOC remain low throughout interval II in both cores,
159 suggesting reduced levels of productivity and nutrient availability (Fig. 3). Variability in TOC
160 and P may also be influenced by changes in redox conditions, lithology and sedimentation

161 rate. However, the decrease in TOC and P in both cores initially occurs in shale in the upper
162 part of interval I, which was deposited under euxinic conditions. Furthermore, despite the
163 lithological differences between the two cores in interval II (and possible corresponding
164 changes in sedimentation rate), a synchronous decline in TOC and P is observed in both
165 cores, suggesting that productivity was the dominant control on TOC and P variability (Text
166 S6).

167 Given the generally enhanced continental weathering flux indicated for the early Cambrian
168 (including during intervals I and II; Peters and Gaines, 2012; Lipp et al., 2021), the decline
169 in nutrient levels likely reflects a reduction in sedimentary P recycling and P upwelling from
170 deeper waters. During interval II, P_{tot}/Al ratios remain below average shale in both drill cores
171 (Fig. 3). On the shelf (ZK 402), ferruginous samples in the lower part of interval II have
172 elevated $C_{\text{org}}/P_{\text{org}}$ and $C_{\text{org}}/P_{\text{reac}}$ ratios relative to the Redfield ratio, but with values that are
173 considerably lower than those of interval I (Fig. 3). This suggests that the extent of P
174 recycling was considerably diminished, with lower P_{tot}/Al ratios implying a major overall
175 decrease in bioavailable P. Furthermore, while the dominantly oxic samples from the rest of
176 interval II also have high $C_{\text{org}}/P_{\text{org}}$ ratios, $C_{\text{org}}/P_{\text{reac}}$ ratios plot below the Redfield ratio. This
177 suggests that the P released during early diagenesis was fixed in the sediment as authigenic
178 phases (supported by an increased proportion of P_{auth} ; Fig. 3). This C/P pattern implies very
179 low bioavailable P, consistent with most modern oligotrophic settings, which are
180 characterized by higher $C_{\text{org}}/P_{\text{org}}$ ratios and lower $C_{\text{org}}/P_{\text{reac}}$ ratios (Slomp et al., 2013).

181 In the slope setting (CY 1), $C_{\text{org}}/P_{\text{org}}$ ratios are elevated and $C_{\text{org}}/P_{\text{reac}}$ ratios scatter close
182 to the Redfield ratio during interval II, with higher values for euxinic samples and lower
183 values for ferruginous samples (Fig. 3 and Fig. S4). These are consistent with a dynamic P
184 response to the prevailing water column redox conditions, with intermittently enhanced P
185 recycling back to the water column under euxinic conditions, although the lower P_{tot} contents
186 relative to interval I (Fig. 3) suggest that water column P concentrations were likely greatly
187 diminished overall. In addition, elevated Co×Mn values suggest that this limited flux of
188 regenerated P was likely not efficiently transported to the photic zone, due to much lower
189 rates of upwelling, thereby resulting in the development of oligotrophic conditions on the

shelf (Fig. 4). This would have reduced primary productivity in surface waters, thus decreasing the amount of dissolved O₂ consumed during the oxidation of sinking organic matter, and thereby stabilizing oxic conditions in shelf environments.

Our data demonstrate that variability in ocean circulation and sedimentary P recycling played a critical role in marine redox and nutrient dynamics during the early Cambrian. Prevailing euxinia along continental margins and active ocean circulation during late Cambrian Stage 2 to early Stage 3 resulted in intensified sedimentary P recycling in deep waters and enhanced nutrient supply to the surface ocean (Fig. 4). Increased primary production would have led to higher atmospheric and shallow marine oxygen levels, and hence an expansion of oxic waters. The enhanced oxygen availability and food supply likely facilitated the early Cambrian bioradiation that occurred at this time. The expanded oxic seafloor area and diminished ocean circulation that occurred during middle to late Cambrian Stage 3 then promoted the retention of sedimentary P in shelf areas, and along with reduced upwelling, resulted in a decrease in shallow marine nutrient levels. Under the ensuing oligotrophic conditions, stable oxygenated continental shelves developed, thereby creating habitats that supported a burgeoning diversity in animal life.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China (42102171, 41921002, 42073006, 41927801). FB and SWP acknowledge funding from the Natural Environment Research Council, UK (NE/R010129/1). TH acknowledges support from a Jiangsu Distinguished Professor Award. We thank Eva Stüeken, Miquela Ingalls and an anonymous reviewer for constructive comments that significantly improved the manuscript.

Data Availability

All data that support the findings of this study are provided in Supplementary Tables S1 and S2.

REFERENCES CITED

- Behrenfeld, M. J., O'Malley, R. T., Siegel, D. A., McClain, C. R., Sarmiento, J. L., Feldman, G. C., Milligan, A. J., Falkowski, P. G., Letelier, R. M., and Boss, E. S., 2006, Climate-driven trends in contemporary ocean productivity: *Nature*, v. 444, no. 7120, p. 752-755, <https://doi.org/10.1038/nature05317>.
- Bowyer, F. T., Krause, A. J., Song, Y. F., Huang, K. J., Fu, Y., Shen, B., Li, J., Zhu, X. K., Kipp, M. A., van Maldegem, L. M., Brocks, J. J., Shields, G. A., Le Hir, G., Mills, B. J. W., and Poulton, S. W., 2023, Biological diversification linked to environmental stabilization following the Sturtian Snowball glaciation: *Science Advances*, v. 9, no. 34, <https://doi.org/10.1126/sciadv.adf9999>.
- Bowyer, F., Wood, R. A., and Poulton, S. W., 2017, Controls on the evolution of Ediacaran metazoan ecosystems: A redox perspective: *Geobiology*, v. 15, no. 4, p. 516-551, <https://doi.org/10.1111/gbi.12232>.
- Brocks, J. J., Jarrett, A. J. M., Sirantoine, E., Hallmann, C., Hoshino, Y., and Liyanage, T., 2017, The rise of algae in Cryogenian oceans and the emergence of animals: *Nature*, v. 548, no. 7669, p. 578+, <https://doi.org/10.1038/nature23457>.
- Chen, X., Ling, H. F., Vance, D., Shields-Zhou, G. A., Zhu, M. Y., Poulton, S. W., Och, L. M., Jiang, S. Y., Li, D., Cremonese, L., and Archer, C., 2015, Rise to modern levels of ocean oxygenation coincided with the Cambrian radiation of animals: *Nature Communications*, v. 6, <https://doi.org/10.1038/ncomms8142>.
- Creveling, J. R., Johnston, D. T., Poulton, S. W., Kotrc, B., März, C., Schrag, D. P., and Knoll, A. H., 2014, Phosphorus sources for phosphatic Cambrian carbonates: *Geological Society of America Bulletin*, v. 126, no. 1-2, p. 145-163, <https://doi.org/10.1130/B30819.1>.
- Dahl, T. W., Connelly, J. N., Li, D., Kouchinsky, A., Gill, B. C., Porter, S., Maloof, A. C., and Bizzarro, M., 2019, Atmosphere-ocean oxygen and productivity dynamics during early animal radiations: *Proc Natl Acad Sci U S A*, v. 116, no. 39, p. 19352-19361, <https://doi.org/10.1073/pnas.1901178116>.
- Erwin, D. H., Laflamme, M., Tweedt, S. M., Sperling, E. A., Pisani, D., and Peterson, K. J.,

247 2011, The Cambrian conundrum: early divergence and later ecological success in the
 248 early history of animals: *Science*, v. 334, no. 6059, p. 1091-1097,
 249 <https://doi.org/10.1126/science.1206375>.

250 He, T. C., Zhu, M. Y., Mills, B. J. W., Wynn, P. M., Zhuravlev, A. Y., Tostevin, R., von
 251 Strandmann, P. A. E. P., Yang, A. H., Poulton, S. W., and Shields, G. A., 2019, Possible
 252 links between extreme oxygen perturbations and the Cambrian radiation of animals:
 253 *Nature Geoscience*, v. 12, no. 6, p. 468-+, <https://doi.org/10.1038/s41561-019-0357-z>.

254 Hearing, T. W., Harvey, T. H. P., Williams, M., Leng, M. J., Lamb, A. L., Wilby, P. R., Gabbott,
 255 S. E., Pohl, A., and Donnadieu, Y., 2018, An early Cambrian greenhouse climate:
 256 *Science Advances*, v. 4, no. 5, <https://doi.org/10.1126/sciadv.aar5690>.

257 Ingall, E. D., Bustin, R. M., and Vancappellen, P., 1993, Influence of Water Column Anoxia
 258 on the Burial and Preservation of Carbon and Phosphorus in Marine Shales:
 259 *Geochimica Et Cosmochimica Acta*, v. 57, no. 2, p. 303-316,
 260 [https://doi.org/10.1016/0016-7037\(93\)90433-W](https://doi.org/10.1016/0016-7037(93)90433-W).

261 Knoll, A. H., and Carroll, S. B., 1999, Early animal evolution: emerging views from
 262 comparative biology and geology: *Science*, v. 284, no. 5423, p. 2129-2137,
 263 <https://doi.org/10.1126/science.284.5423.2129>.

264 Lenton, T. M., Boyle, R. A., Poulton, S. W., Shields-Zhou, G. A., and Butterfield, N. J., 2014,
 265 Co-evolution of eukaryotes and ocean oxygenation in the Neoproterozoic era: *Nature*
 266 *Geoscience*, v. 7, no. 4, p. 257-265, <https://doi.org/10.1038/ngeo2108>.

267 Li, C., Jin, C. S., Planavsky, N. J., Algeo, T. J., Cheng, M., Yang, X. L., Zhao, Y. L., and Xie,
 268 S. C., 2017, Coupled oceanic oxygenation and metazoan diversification during the
 269 early-middle Cambrian?: *Geology*, v. 45, no. 8, p. 743-746,
 270 <https://doi.org/10.1130/G39208.1>.

271 Lipp, A. G., Shorttle, O., Sperling, E. A., Brocks, J. J., Cole, D. B., Crockford, P. W., Del
 272 Mouro, L., Dewing, K., Dornbos, S. Q., Emmings, J. F., Farrell, U. C., Jarrett, A.,
 273 Johnson, B. W., Kabanov, P., Keller, C. B., Kunzmann, M., Miller, A. J., Mills, N. T.,
 274 O'Connell, B., Peters, S. E., Planavsky, N. J., Ritzer, S. R., Schoepfer, S. D., Wilby, P.
 275 R., and Yang, J., 2021, The composition and weathering of the continents over geologic

time: Geochemical Perspectives Letters, v. 17, p. 21-26,
<https://doi.org/10.7185/geochemlet.2109>.

Mills, D. B., Ward, L. M., Jones, C., Sweeten, B., Forth, M., Treusch, A. H., and Canfield, D. E., 2014, Oxygen requirements of the earliest animals: Proceedings of the National Academy of Sciences of the United States of America, v. 111, no. 11, p. 4168-4172, <https://doi.org/10.1073/pnas.1400547111>.

Peters, S. E., and Gaines, R. R., 2012, Formation of the 'Great Unconformity' as a trigger for the Cambrian explosion: Nature, v. 484, no. 7394, p. 363-366, <https://doi.org/10.1038/nature10969>.

Poulton, S. W., 2021, The Iron Speciation Paleoredox Proxy: Cambridge, Cambridge University Press.

Schlesinger, W. H., and Bernhardt, E. S., 2013, Biogeochemistry: An analysis of global change. 3rd edition. .

Schobben, M., Foster, W. J., Sleveland, A. R. N., Zuchuat, V., Svensen, H. H., Planke, S., Bond, D. P. G., Marcelis, F., Newton, R. J., Wignall, P. B., and Poulton, S. W., 2020, A nutrient control on marine anoxia during the end-Permian mass extinction: Nature Geoscience, v. 13, no. 9, p. 640-+, <https://doi.org/10.1038/s41561-020-0622-1>.

Slomp, C. P., Mort, H. P., Jilbert, T., Reed, D. C., Gustafsson, B. G., and Wolthers, M., 2013, Coupled Dynamics of Iron and Phosphorus in Sediments of an Oligotrophic Coastal Basin and the Impact of Anaerobic Oxidation of Methane: Plos One, v. 8, no. 4, <https://doi.org/10.1371/journal.pone.0062386>.

Slomp, C. P., VanderGaast, S. J., and VanRaaphorst, W., 1996, Phosphorus binding by poorly crystalline iron oxides in North Sea sediments: Marine Chemistry, v. 52, no. 1, p. 55-73, [https://doi.org/10.1016/0304-4203\(95\)00078-X](https://doi.org/10.1016/0304-4203(95)00078-X).

Smith, M. P., and Harper, D. A. T., 2013, Causes of the Cambrian Explosion: Science, v. 341, no. 6152, p. 1355-1356, <https://doi.org/10.1126/science.1239450>.

Sperling, E. A., Frieder, C. A., Raman, A. V., Girguis, P. R., Levin, L. A., and Knoll, A. H., 2013, Oxygen, ecology, and the Cambrian radiation of animals: Proceedings of the National Academy of Sciences of the United States of America, v. 110, no. 33, p. 13446-

13451, <https://doi.org/10.1073/pnas.1312778110>.

Sun, Z. X., Zhao, F. C., Zeng, H., Luo, C., Van Iten, H., and Zhu, M. Y., 2022, The middle Cambrian Linyi Lagerstätte from the North China Craton: a new window on Cambrian evolutionary fauna: National Science Review, v. 9, no. 7, <https://doi.org/10.1093/nsr/nwac069>.

Sweere, T., van den Boorn, S., Dickson, A. J., and Reichart, G. J., 2016, Definition of new trace-metal proxies for the controls on organic matter enrichment in marine sediments based on Mn, Co, Mo and Cd concentrations: Chemical Geology, v. 441, p. 235-245, <https://doi.org/10.1016/j.chemgeo.2016.08.028>.

Thompson, J., Poulton, S. W., Guilbaud, R., Doyle, K. A., Reid, S., and Krom, M. D., 2019, Development of a modified SEDEX phosphorus speciation method for ancient rocks and modern iron-rich sediments: Chemical Geology, v. 524, p. 383-393, <https://doi.org/10.1016/j.chemgeo.2019.07.003>.

Tribovillard, N., Algeo, T. J., Lyons, T., and Riboulleau, A., 2006, Trace metals as paleoredox and paleoproductivity proxies: An update: Chemical Geology, v. 232, no. 1-2, p. 12-32, <https://doi.org/10.1016/j.chemgeo.2006.02.012>.

Turekian, K. K., and Wedepohl, K. H., 1961, Distribution of the Elements in Some Major Units of the Earth's Crust: GSA Bulletin, v. 72, no. 2, p. 175-192, [https://doi.org/10.1130/0016-7606\(1961\)72\[175:Doteis\]2.0.Co;2](https://doi.org/10.1130/0016-7606(1961)72[175:Doteis]2.0.Co;2).

Tyrrell, T., 1999, The relative influences of nitrogen and phosphorus on oceanic primary production: Nature, v. 400, no. 6744, p. 525-531, <https://doi.org/10.1038/22941>.

Van Cappellen, P., and Ingall, E. D., 1996, Redox Stabilization of the Atmosphere and Oceans by Phosphorus-Limited Marine Productivity: Science, v. 271, no. 5248, p. 493-496, <https://doi.org/doi:10.1126/science.271.5248.493>.

Wei, G. Y., Planavsky, N. J., Tarhan, L. G., He, T. C., Wang, D., Shields, G. A., Wei, W., and Ling, H. F., 2020, Highly dynamic marine redox state through the Cambrian explosion highlighted by authigenic $\delta^{238}\text{U}$ records: Earth and Planetary Science Letters, v. 544, <https://doi.org/10.1016/j.epsl.2020.116361>.

Wei, G.-Y., Planavsky, N. J., Tarhan, L. G., Chen, X., Wei, W., Li, D., and Ling, H.-F., 2018,

Marine redox fluctuation as a potential trigger for the Cambrian explosion: *Geology*, v. 46, no. 7, p. 587-590, <https://doi.org/10.1130/g40150.1>.

Wood, R., Liu, A. G., Bowyer, F., Wilby, P. R., Dunn, F. S., Kenchington, C. G., Cuthill, J. F. H., Mitchell, E. G., and Penny, A., 2019, Integrated records of environmental change and evolution challenge the Cambrian Explosion: *Nat Ecol Evol*, v. 3, no. 4, p. 528-538, <https://doi.org/10.1038/s41559-019-0821-6>.

Zhuravlev, A. Y., Mitchell, E. G., Bowyer, F., Wood, R., and Penny, A., 2022, Increases in reef size, habitat and metacommunity complexity associated with Cambrian radiation oxygenation pulses: *Nat Commun*, v. 13, no. 1, p. 7523, <https://doi.org/10.1038/s41467-022-35283-5>.

FIGURE CAPTIONS

Fig. 1. Stratigraphic plots of Fe speciation, RSTM EFs and Mo_{EF}/U_{EF} ratios for drill cores ZK402 and CY 1. Fm, Formation; Ed, Ediacaran; Fo, Fortunian; DY, Dengying; LCP, Liuchapo.

Fig. 2. Marine redox conditions and biodiversity patterns during the early Cambrian. A. Compilation of Fe speciation data for platform, shelf, slope and basin environments from 541 Ma to 514 Ma. Blue and orange lines represent the LOESS curves. B. Global diversity of phyla and classes (Erwin et al., 2011) and major Cambrian lagerstätten (Sun et al., 2022). C. Schematic evolution of local marine redox conditions based on iron speciation data. D. Schematic evolution of global marine redox conditions based on Mo and U isotope records (Chen et al., 2015; Dahl et al., 2019; Wei et al., 2020).

Fig. 3. Stratigraphic distribution of TOC, $\delta^{13}C_{org}$, P_{tot} , P_{tot}/Al , C_{org}/P_{org} , C_{org}/P_{reac} , $Co \times Mn$ and P phase partitioning.

Fig. 4. Summary of marine redox and nutrient cycling during the early Cambrian. A.

363 Persistently euxinic conditions on the shelf and slope promoted P regeneration into the water
364 column. Recycled P was transported to the photic zone by active upwelling, resulting in a
365 positive productivity feedback that led to increased C_{org} burial, and ultimately atmospheric
366 and shallow marine oxygenation. B. P was effectively fixed in shelf sediments. Continued
367 euxinia in slope environments may have resulted in intermittent P recycling to the water
368 column, but P in deep waters was not effectively transported to the photic zone due to
369 decreased upwelling, which limited primary production and O₂ consumption.

370

371 ¹Supplemental Material. Detailed geological setting, methods, supplemental notes, Figures
372 S1–S4 and Tables S1–S2. Please visit <https://doi.org/10.1130/XXXX> to access the
373 supplemental material, and contact editing@geosociety.org with any questions.