



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/220502/>

Version: Published Version

Article:

Rahim, Abid, Hassan, Sabreen, Ullah, Naeem et al. (2023) Association and comparison of periodontal and oral hygiene status with serum HbA1c levels: a cross-sectional study. BMC Oral Health. 442. ISSN: 1472-6831

<https://doi.org/10.1186/s12903-023-03042-7>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown


If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

RESEARCH

Open Access



Association and comparison of periodontal and oral hygiene status with serum HbA1c levels: a cross-sectional study

Abid Rahim^{1,2}, Sabreen Hassan³, Naeem Ullah⁴, Nawal Noor¹, Ahmed¹, Rimsha Rafique⁵, Farhad Ali Khattak^{6*}  and Saima Afaq^{7,8*}

Abstract

Background Diabetes Mellitus and periodontitis are chronic diseases with known reciprocal association. Studies have shown that uncontrolled diabetes increases the risk of development and progression of periodontal disease. This study aimed to explore the association and severity of periodontal clinical parameters and oral hygiene with HbA1c levels in non-diabetics and T2DM patients.

Materials and methods In this cross-sectional study, the periodontal status of 144 participants, categorized into non-diabetics, controlled T2DM, and uncontrolled T2DM and were assessed via the Community Periodontal Index (CPI), Loss of Attachment Index (LOA index), and the number of missing teeth, while oral hygiene was measured by utilizing the Oral Hygiene Index Simplified (OHI-S). SPSS was used for data analysis. Chi-square test was used to find out the association of different independent variables with HbA1c groups, while ANOVA and post-hoc tests were run for inter-group and intra-group comparison respectively.

Results Out of 144 participants, the missing dentition was prevalent in uncontrolled T2DM with mean 2.64 ± 1.97 (95% CI 2.07–3.21; $p=0.01$) followed by controlled T2DM 1.70 ± 1.79 (95% CI 1.18–2.23; $p=0.01$) and non-diabetics 1.35 ± 1.63 (95% CI 0.88–1.82; $p=0.01$) respectively. Furthermore, non-diabetics had a higher proportion of CPI score 0 (Healthy) [30 (20.8%); $p=0.001$] as compared to uncontrolled T2DM [6 (4.2%); $p=0.001$], while CPI score 3 was more prevalent in uncontrolled T2DM in comparison to non-diabetics. Loss of attachment (codes-2,3 and 4) was also frequently observed in uncontrolled T2DM compared to non-diabetics ($p=0.001$). Similarly, based on Oral Hygiene Index- Simplified (OHI-S), the result showed that poor oral hygiene was most commonly observed in uncontrolled T2DM 29 (20.1%) followed by controlled T2DM patients 22 (15.3%) and non-diabetic [14 (9.7%); $p=0.03$].

Conclusion This study showed that periodontal status and oral hygiene status were deteriorated in uncontrolled T2DM patients compared to non-diabetic participants and controlled T2DM.

Keywords Periodontitis, Diabetes mellitus, Probe, Bleeding on probing, Pocket depth, Loss of attachment

*Correspondence:

Farhad Ali Khattak

farhadkcd@gmail.com

Saima Afaq

s.afaq11@imperial.ac.uk; saima.iph@kmu.edu.pk

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Diseases of the periodontium (i.e., gingivitis and periodontitis) are multifactorial bacterial diseases of the soft and hard tissues encompassing and supporting the teeth. It is initiated via the aggregation of a pathogenic dental plaque on the tooth surface, and inside which microbial dysbiosis prompts a constant non-resolving damaging provocative reaction [1]. The global prevalence of periodontitis ranges from 15–47% and in its most severe form affects 10.8% of the population [2]. It has been ranked seventh and 32nd worldwide for prevalence and incidence respectively [3]. Chronic periodontal disease not only affects the oral health but also systemic health [4]. Studies by Isola et al. found that periodontitis patients showed a higher risk of developing endothelial dysfunction and cardiovascular disease [5, 6]. The development and progression of periodontal diseases are directly linked to oral hygiene, and maintaining good oral hygiene reduces the risk of periodontitis [7, 8]. Fair and poor oral hygiene maximize the risk of periodontitis by two to five times respectively [9].

Diabetes mellitus (DM) portrayed by hyperglycemia, is a metabolic disorder caused by insufficient insulin production, insufficient insulin activity, or both. Diabetes Type-1 is characterized by an autoimmune attack on the pancreatic insulin-producing β cells, which leads to insufficient insulin synthesis. Type-2 diabetes mellitus is brought about by a blend of insulin opposition and insulin emission disability. Diabetes is currently thought to affect over 10% of people worldwide. According to estimates, 462 million people worldwide or 6.28% of the world's population have T2DM [10, 11].

The relationship between diabetes and periodontitis is symbiotic because persistent hyperglycemia has been shown to negatively influence oral health, and severe periodontitis can have a negative impact on both glycaemic control and diabetic complications [12]. Advanced glycation end-products are produced by type 2 diabetes and impaired insulin sensitivity, which leads to the generation of inflammatory cytokines and predisposes individuals to inflammatory conditions like periodontitis [13]. Diabetic individuals with periodontitis exhibited higher amounts of inflammatory mediators in their saliva and their crevicular fluid than non-diabetics with periodontitis, including numerous kinds of cytokines [14]. There is a proportionate association between HbA1c and periodontitis [15, 16]. On the other hand, periodontitis adds to the overall inflammatory load in the body by the transfer of bacteria and their products, cytokines, and inflammatory mediators via breached pocket epithelium and higher GCF miRNAs expression, exacerbating the development of complications in diabetic patients [5, 14, 17].

Given the well-established link between periodontitis and type 2 diabetes mellitus, this cross-sectional study

aims to evaluate and compare the periodontal and oral hygiene status of individuals having different levels/categories of HbA1c. Unlike previous studies, this research simultaneously investigates periodontitis, oral hygiene, and diabetes in three subgroups: non-diabetics, controlled, and uncontrolled type 2 diabetes patients. By investigating this triad, this study provides a comprehensive understanding of the relationship between oral health and diabetes, which has a significant impact on overall well-being. Moreover, it aims to provide valuable insights into the extent of periodontal disease and oral hygiene status in a population specific to low and middle-income countries. The results of this study may have important implications for the prevention and management of periodontitis and type 2 diabetes, particularly in resource-limited settings.

Materials and methods

Study participants

This cross-sectional study was conducted on patients visiting the outpatient department of Qazi Hussain Ahmed Medical Complex (Medical Teaching Institute/MTI), Khyber Pakhtunkhwa - Pakistan from March 2021 to June 2021, after ethical approval from Khyber Medical University (DIR/KMU-EB/EC/00080/DR). Using the WHO sample size calculator, considering the frequency of poor oral hygiene status among diabetics P1 (22% compared to poor oral hygiene among non-diabetics P2 (37%), taking 80% power of the test and 5% margin of error, a total of 144 patients were recruited [18]. Based on the HbA1c levels, all of the participants were divided into three equal groups. *Group A*: Non-diabetic (HbA1c \leq 5.7%) *Group B*: Controlled T2DM (HbA1c 6.0–6.9%) *roup C*: Uncontrolled T2DM (HbA1c \geq 7%) [19]. The enrolment criteria were (i) Dentate patients with age 20–60 years. (ii) Subjects diagnosed with T2DM for more than one year (groups B and C). The exclusion criteria were (i) Patients having any other systemic disease/s other than diabetes mellitus. (ii) individuals who have undergone periodontal therapy in the past three months (iii) Patients who have taken antibiotics for the past month. (iv) Patients having 3rd molar impaction, endodontic problems, or limited mouth opening. (v) Female participants that are expecting/lactating.

Interview questionnaire

Each participant gave written informed consent, after being informed about the pertinent interview and oral examination required for carrying out the study. An organized form was used to obtain the necessary information like age, gender, education level, occupation, frequency of tooth cleaning and tool used for cleaning, last dental visit, the reason for a dental visit, frequency and type of tobacco consumed, duration of diabetes,

and complications of diabetes, number of missing teeth. The HbA1c level for each T2DM patient was obtained from his/her medical record. Similarly, the HbA1c level for non-diabetics was determined before the oral examination.

Oral examination

All the subjects underwent oral examination to evaluate their periodontal status (i.e., bleeding upon probing, depth of the pocket, and loss of attachment), and oral hygiene status (dental plaque and calculus). The examination for each participant was done via mouth mirror, CPITN-C probe, and No. 5 explorer (Shepherd's Crook). The periodontal assessment was done using the Community Periodontal Index (CPI index) (with scoring criteria as code-0, code-1, code-2, code-3, code-4, or code-x) and the Loss of Attachment Index. Oral hygiene for each patient was assessed according to Oral Hygiene Index-Simplified and categorized into Good oral hygiene, Fair, and Poor oral hygiene. Indexed teeth: 17/16, 11, 26/27 (buccal surfaces) and 47/46, 31, 36/37 (lingual surfaces) were examined during the oral examination. Oral examination was carried out by a single and trained periodontist (AR) to prevent differences in measurements.

Data analysis

All the relevant data were collected and recorded on a preformed structured questionnaire. Statistical analysis was done via SPSS version 25. The Chi-square test/Fisher exact test was used to compare non-diabetics, controlled T2DM, and uncontrolled T2DM with other explanatory variables like sociodemographic traits, diabetes-associated factors, oral hygiene status, and periodontal status, while one-way ANOVA was carried out to determine any statistical significance among the three groups in relation to other continuous variables. For those with significant P value, Post Hoc (Tukey HSD) Test was run to explore the mean differences between the pair of groups. The threshold for statistical significance was set at $p \leq 0.05$.

Results

Baseline characteristics

A total of 144 participants voluntarily took part in this study with 87 (59.7%) males, with a mean age of 42.5 ± 12.3 years for non-diabetic and 43.7 ± 10.2 years for controlled diabetics while 49.2 ± 8.8 years for uncontrolled T2DM ($p=0.002$) and mean serum HbA1c levels of 5.2 ± 1.09 , 6.7 ± 0.50 and 8.3 ± 1.09 for the three groups (non-diabetics, controlled T2DM, uncontrolled T2DM) respectively ($p=0.001$) (Tables 1 and 2). The prevalence of missing dentition was comparatively higher in the uncontrolled T2DM group with mean of 2.64 ± 1.9 and 1.3 ± 1.6 (Table 2). The mean difference of various variables was notably different between non-diabetics and

uncontrolled T2DM (Table 3). The majority of the participants were illiterate 44 (30.6%), while only 32 (22.2%) had completed their graduation. The majority of the participants 47 (32.6%) were nonoccupational, followed by employed participants 41 (28.55). Most of the participants 48 (33.3%) cleaned their teeth once a day, twice daily 32 (22.2%), and once a week by 33 (22.9%) participants. Toothbrush was used as a cleaning tool by 101 (70.1%), miswak was used by 22 (15.3%), and 21 (14.6%) participants used other tools (*danddassa/Juglans regia* linn) for cleaning their teeth. A minor number of participants 13 (9.0%) also had no dental check-up throughout their life. Most of the participants 91 (63.2%) did not use any tobacco product, and 22 (15.3%) consumed smoked tobacco (cigarette), while 31 (21.5%) used smokeless tobacco (*naswar*). Among the tobacco consumers, 44 (30.6%) consumed tobacco ≥ 3 times a day (Table 1).

Periodontal Status among Non-Diabetics, controlled diabetics, and uncontrolled diabetics

Periodontal status as clinically determined via the Community Periodontal Index, Loss of Attachment Index, and missing teeth, varied distinctly among the three groups. Healthy periodontium (code-0) was found in 30 (20.8%) of non-diabetics while this ratio was 6 (4.2%) in uncontrolled T2DM ($p=0.001$). Similarly, code-4 was most prevalent in uncontrolled T2DM 19 (13.2%); $p=0.01$ and least in the non-diabetic group 1 (0.7%) (Table 1 and Fig. 1). Loss of attachment (code-0) was prevalent in non-diabetics [27 (18.8%); $p=0.001$] followed by the controlled T2DM group [17 (11.8%); $p=0.001$] and infrequent in uncontrolled T2DM [5 (3.5%); $p=0.001$]. Loss of attachment (code-4) was more common in the uncontrolled T2DM group [9 (6.3%); $p=0.001$] and least common in non-diabetics [1 (0.7%); $p=0.001$] (Table 1 and Fig. 2). The uncontrolled T2DM group showed the highest mean for missing teeth [2.64 ± 1.97 ; $p=0.01$] followed by controlled T2DM [1.70 ± 1.79 ; $p=0.01$] and non-diabetics [1.35 ± 1.63 ; $p=0.01$] (Table 1).

Oral hygiene status among non-diabetics, controlled diabetics, and uncontrolled diabetics

Oral hygiene status was designated as Good, Fair, and Poor according to Oral Hygiene Index-Simplified. Good Oral Hygiene was observed in [27 (18.8%); $p=0.03$] non-diabetics followed by controlled T2DM [19 (1.2%); $p=0.03$] and uncontrolled T2DM [09 (6.3%); $p=0.03$]. Poor oral hygiene was more prevalent in uncontrolled T2DM [29 (20.1%); $p=0.03$] and least prevalent in non-diabetics [14 (9.7%); $p=0.03$] (Tables 1 and Fig. 3).

Table 1 Association of Diabetic and Non-diabetic group(s) with different explanatory variables

Characteristics	Categories	Groups			P-value
		Non-diabetics (n = 48)	Controlled T2DM (n = 48)	Uncontrolled T2DM (n = 48)	
Gender	Male	26 (18.1%)	32 (22.2%)	28 (19.4%)	0.44
	Female	22 (15.3%)	16 (11.1%)	20 (13.9%)	
Education level	Illiterate	11 (7.6%)	14 (9.7%)	19 (13.2%)	0.46*
	Primary school	8 (5.6%)	8 (5.6%)	9 (6.3%)	
	Secondary school graduation	11 (7.6%)	11 (7.6%)	13 (9.0%)	
	More than graduation	15 (10.4%)	12 (8.3%)	5 (3.5%)	
Occupation	Professional	3 (2.1%)	3 (2.1%)	2 (1.4%)	0.44*
	Employment	2 (1.4%)	3 (2.1%)	1 (0.7%)	
	Business	15 (10.4%)	16 (11.1%)	10 (6.9%)	
	Vocational	8 (5.6%)	13 (9.0%)	17 (11.8%)	
Frequency of teeth cleaning	None	6 (4.2%)	3 (2.1%)	3 (2.1%)	0.39*
	Once a day	17 (11.8%)	13 (9.0%)	17 (11.8%)	
	Twice a day	16 (11.1%)	16 (11.1%)	16 (11.1%)	
	Once a week	15 (10.4%)	12 (8.3%)	5 (3.5%)	
	2–3 times a week	9 (6.3%)	11 (7.6%)	13 (9.0%)	
Tool used for teeth cleaning	Rarely	6 (4.2%)	6 (4.2%)	8 (5.6%)	0.05*
	Toothbrush	2 (4.2%)	3 (2.1%)	6 (4.2%)	
	Miswak	39 (27.1%)	36 (25.0%)	26 (18.1%)	
Last dental visit	Others	5 (3.5%)	6 (4.2%)	11 (7.6%)	0.02*
	≤ 1 year	4 (2.8%)	6 (4.2%)	11 (7.6%)	
	≥ 1 year	20 (13.9%)	12 (8.3%)	20 (13.9%)	
Reason for a dental visit	None	27 (18.8%)	29 (20.1%)	35 (24.3%)	0.04*
	Pain	1 (0.7%)	7 (4.9%)	5 (3.5%)	
	Sensitivity	8 (5.6%)	4 (2.8%)	3 (2.1%)	
	Scaling and polishing	11 (7.6%)	8 (5.6%)	3 (2.1%)	
Tobacco consumption and types used	Others	1 (0.7%)	0 (0.0%)	3 (2.1%)	0.75
	None	1 (0.7%)	7 (4.9%)	5 (3.5%)	
	Smoked tobacco	7 (4.9%)	9 (6.3%)	6 (4.2%)	
Frequency of tobacco consumption	Smokeless tobacco	8 (5.6%)	11 (7.6%)	12 (8.3%)	0.86*
	None	33 (22.9%)	28 (19.4%)	30 (20.8%)	
	1–2 times a day	3 (2.1%)	3 (2.1%)	3 (2.1%)	
Duration of diabetes	≥ 3 times a day	12 (8.3%)	17 (11.8%)	15 (10.4%)	0.001*
	None	33 (22.9%)	28 (19.4%)	30 (20.8%)	
	≤ 5 years	0 (0.0%)	24 (16.7%)	11 (7.6%)	
Diabetes medication	6–9 years	0 (0.0%)	16 (11.2%)	15 (10.5%)	0.001*
	≥ 10 years	0 (3.3%)	8 (5.6%)	22 (15.4%)	
	None	48 (33.3%)	13 (9.0%)	1 (0.7%)	
Community Peri-odontal Index (CPI)	Oral	0 (0.0%)	34 (23.6%)	34 (23.6%)	0.001*
	Combination (oral + inj insulin)	0 (0.0%)	1 (0.7%)	13 (9.0%)	
	Healthy-0	30 (20.8%)	15 (10.4%)	6 (4.2%)	
	Bleeding on probing – 1	4 (2.8%)	5 (3.5%)	4 (2.8%)	
	Calculus detected- 2	3 (2.1%)	6 (4.2%)	8 (5.6%)	
Loss of Attachment Index	Pocket 4–5 mm (gingival margin within the black band of probe) – 3	8 (5.6%)	12 (8.3%)	11 (7.6%)	0.001*
	Pocket ≥ 6 mm (black band not visible)-4	3 (2.1%)	10 (6.9%)	19 (13.2%)	
	Loss of attachment 0-3 mm (CEJ not visible) Code-0	27 (18.8%)	17 (11.8%)	5 (3.5%)	
	Loss of attachment 4-5 mm (CEJ within the black band) Code – 1	17 (11.8%)	15 (10.4%)	11 (7.6%)	
	Loss of attachment 6-8 mm (CEJ between 5-8 mm ring) Code-2	2 (1.4%)	7 (4.9%)	15 (10.4%)	
Loss of Attachment Index	Loss of attachment 9-11 mm (CEJ between 8.5-11.5 mm ring) Code- 3	1 (0.7%)	6 (4.2%)	8 (5.6%)	0.001*
	Loss of attachment ≥ 12 mm (CEJ beyond 11 mm ring)- 4	1 (0.7%)	3 (2.1%)	9 (6.3%)	

Table 1 (continued)

Characteristics	Categories	Groups			P-value
		Non-diabetics (n = 48)	Controlled T2DM (n = 48)	Uncontrolled T2DM (n = 48)	
Oral Hygiene Index- Simplified (OHI-S)	Good oral hygiene	27 (18.8%)	19 (13.2%)	10 (6.9%)	0.03
	Fair oral hygiene	7 (4.9%)	7 (4.9%)	9 (6.3%)	
	Poor oral hygiene	14 (9.7%)	22 (15.3%)	29 (20.1%)	

Chi-square test/Fisher Exact* | p≤0.05=statistically significant | p=0.001=highly significant

Table 2 Comparative analysis of continuous variables with non-diabetic/diabetic groups

Variable	Groups	Mean	Std.Deviation	95% Confidence Interval for Mean		F	P Value
				Lower	Upper		
Age	Non-diabetic	42.54	12.33	38.95	46.12	5.46	0.005
	Controlled diabetic	43.72	10.23	40.75	46.69		
	Uncontrolled diabetic	49.22	8.87	46.65	51.8		
HbA1c level(%)	Non-diabetic	5.24	0.41	5.12	5.36	207.26	0.001
	Controlled diabetic	6.72	0.5	6.57	5.3		
	Uncontrolled diabetic	8.3	1.09	7.98	8.61		
Number of missing teeth (except 3rd molar)	Non-diabetic	1.35	1.63	0.88	1.82	6.55	0.002
	Controlled diabetic	1.7	1.79	1.18	2.23		
	Uncontrolled diabetic	2.64	1.97	2.07	3.21		

One Way ANOVA| p≤0.05=statistically significant

Table 3 Post Hoc (Tukey HSD) Test with Multiple comparisons

Dependent Variable	Type of groups	Intra groups Comparison	Mean Difference	P value	95% Confidence Interval	
					Lower Bound	Upper Bound
Age	Controlled T2DM	Uncontrolled T2DM	-5.50	0.03*	-10.61	-0.39
		Non-diabetic	1.18	0.85	-3.93	6.30
	Uncontrolled T2DM	Controlled T2DM	5.500	0.03*	0.39	10.61
		Non-diabetic	6.68	0.01*	1.57	11.80
	Non-diabetic	Controlled T2DM	-1.18	0.85	-6.30	3.93
		Uncontrolled T2DM	-6.68	0.01*	-11.80	-1.57
Level of HbA1c (%)	Controlled Diabetes	Uncontrolled T2DM	-1.57	0.00*	-1.93	-1.22
		Non-diabetic	1.47	0.00*	1.12	1.83
	Uncontrolled T2DM	Controlled T2DM	1.57	0.00*	1.22	1.93
		Non-diabetic	3.05	0.00*	2.70	3.41
	Non-diabetic	Controlled T2DM	-1.47	0.00*	-1.83	-1.12
		Uncontrolled T2DM	-3.05	0.00*	-3.41	-2.70
Number of missing teeth (except 3rd molar)	Controlled T2DM	Uncontrolled T2DM	-0.93	0.03*	-1.81	-0.06
		Non-diabetic	0.35	0.60	-0.52	1.23
	Uncontrolled T2DM	Controlled T2DM	0.93	0.03*	0.06	1.81
		Non-diabetic	1.29	0.00*	0.42	2.16
	Non-diabetic	Controlled T2DM	-0.35	0.60	-1.23	0.52
		Uncontrolled T2DM	-1.29	0.00*	-2.16	-0.42

*. The mean difference is significant at the 0.05 level

Discussion

By utilizing a representative sample of adults with type II diabetes mellitus and those without the disease, a statistically significant association was found between periodontal and oral hygiene status and the HbA1c levels. The results of this study showed that patients with

uncontrolled T2DM had worse periodontal health as revealed by the increase in bleeding on probing and pocket depth (assessed by CPI Index) and increased clinical attachment loss (assessed by Loss of Attachment Index) as compared to their healthy counterpart and controlled T2DM.




Fig. 1 Community Periodontal Index




Fig. 2 Loss of Attachment Index

CPI code 0 (healthy) was more prevalent in non-diabetics and controlled T2DM as compared to uncontrolled T2DM which are close to the observations of the other studies comparing the same groups [18–20]. Similarly, code 1 (BOP) was almost similar among the three groups, which opposes the findings of other cross-sectional studies, using the CPI index [21–23] and coincides with the study of Kim et al. [19]. CPI code 2 (calculus) was higher in uncontrolled T2DM in comparison to non-diabetics and controlled T2DM which is consistent with the results of Apporva et al. [20]. The code 2 scores of our study contradict the results of other cross-sectional studies involving T2DM patients [19, 24]. The three groups of the present study shared almost the same scores for CPI code 3, which contradict the results of studies done

by other investigators involving CPI usage [18, 20]. CPI code 4 was highest in uncontrolled T2DM as compared to non-diabetics and controlled T2DM, which is alike the findings of other studies [18–20, 22]. The tentative propositions for these similarities and discrepancies could be the same/different sociodemographic characteristics, oral hygiene behavior, education level, access to dental services, and discrepancies/similarities in the measuring. On a universally agreeable scale, the possible phenomenon resulting in these grievous periodontal clinical parameters is the diabetes-periodontitis reciprocity. T2DM impacts periodontitis development, progression, and severity by the production of advanced glycation end products, triggering a hyperinflammatory response,




Fig. 3 Oral Hygiene Index -Simplified

modulation of the periodontal microbiota, and impaired alveolar bone healing [25–28].

Clinical attachment loss, which is another prime indicator of periodontal disease progression, was also assessed among the three groups in this study. Loss of attachment code 0 (Loss of attachment 0–3 mm) was highest in non-diabetics (18.8%) compared to uncontrolled T2DM patients (3.5%), while code 4 (Loss of attachment 9–11 mm) which is the worst score, was highest in uncontrolled T2DM (6.3%) compared to non-diabetics in which it was prevalent in just 0.7%. These findings are supported by other studies which compared the same among T2DM patients [28, 29]. The possible mechanisms which affect the clinical level are AGEs which initiate and progress a cascade of inflammatory events resulting in the degradation of the attachment apparatus. Moreover, the fibroblasts' function is impaired by the hyperglycemic environment, predisposing the collagen to degrade, which inhibits the repair and regeneration [29, 30].

Another key finding of this study was the increased number of tooth loss in uncontrolled T2DM. This finding is in accordance with previous studies which proved that tooth loss due to periodontitis is closely influenced by uncontrolled T2DM. This is supported by other studies [31, 32]. The possible explanation for this augmented tooth loss in uncontrolled T2DM is the defective bone composition and structure as well as increased severity of alveolar bone loss [33, 34]. Tooth loss due to periodontitis not only jeopardizes the masticatory function but also negatively affects the levels of progenitor cells which appear to increase the susceptibility of endothelial cell dysfunction and chronic inflammation [35].

Oral hygiene was another prime aspect of this cross-sectional study. Oral hygiene was determined with the

Oral Hygiene Index Simplified which is a simple and time-saving tool with three categories of Good, Fair, and Poor Oral hygiene. Non-diabetics had good oral hygiene as compared to uncontrolled T2DM, while poor oral hygiene was more prevalent in uncontrolled T2DM in contrast to non-diabetics. Other studies comparing oral hygiene status are consistent with the finding of these studies [6, 31]. These findings contradict the results of a cross-sectional study done in India, in which only 22% of diabetics showed poor oral hygiene while 37% of non-diabetics showed poor oral hygiene [24]. The possible justification for this contrast could be differences in oral hygiene behavior like less frequency of tooth brushing, improper tooth brushing technique or lack of proper oral hygiene knowledge, or compromised access to dental care services. Poor oral hygiene and low frequency of toothbrushing not only enhance the prevalence of periodontal disease but also increase the risk of T2DM [36, 37].

This study attempted to comprehend the two key risk factors (periodontitis and T2DM) of oral and systemic health. Both these conditions, sharing a bidirectional relationship, require a multidisciplinary approach to minimize and prevent the adverse effects of these two chronic diseases on oral health and overall well-being. Dentists should be aware of the diabetic status of their patients and the same goes for the physicians/diabetologist to be aware of the oral complication of diabetes mellitus. Access to dental care should be made easier and regular professional dental care should be encouraged in T2DM patients specifically, which will not only minimize the diabetes' oral complication but also will improve the serum HbA1c levels [27, 38–40].

The limitation of this study was the sample size which was limited to a specific geographic area that cannot be generalized. A cross-sectional study could be the other limitation. Using the WHO-validated tool and oral examination by a single examiner were the strengths of this study. Future studies may be carried out on a large sample size using different study designs to find out a temporal relationship between periodontal health and T2DM.

Conclusion

In light of the results of this cross-sectional study, it is concluded that periodontal parameters (bleeding on probing, pocket depth, clinical attachment loss, and tooth loss) and oral hygiene status (plaque and calculus) were worsened in uncontrolled in T2DM in comparison to non-diabetic participants. A multidisciplinary approach is needed to maintain periodontal health and overall well-being in diabetic patients.

List of Abbreviations

HbA1c	Hemoglobin A1c
T2DM	Type 2 diabetes mellitus
WHO	World Health Organization
CPITN-C	Community Periodontal Index of Treatment Needs - C
CPI	Community Periodontal Index
OHI- S	Oral Hygiene Index – Simplified
BOP	Bleeding of probing
AGEs	Advanced Glycation End Products

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12903-023-03042-7>.

Supplementary Material 1

Supplementary Material 2

Acknowledgements

The authors extend their gratitude to Abdul Basit for his technical assistance.

Authors' contributions

Design of the study – AR, FAK, SA; Data collection – AR; Analysis – AR, FAK, Data interpretation – all authors, Draft the work – all authors, Revision – AR, FAK, SA.

Funding

No funding was obtained from any institution/organization.

Data Availability

All data generated or analyzed during this study are included in these published articles & supporting materials (Supporting data_S1 and Supporting data_S2).

Declarations

Ethics approval and consent to participate

Ethical approval from the Ethics Board of Khyber Medical University, Peshawar - Pakistan (KMU/IPHSS/Ethics/2021/CO/090). Each participant gave written informed consent for participation in the study and relevant data sharing. In addition to this, all the methods/procedures were performed in accordance with the relevant guidelines and regulations (Declaration of Helsinki).

Consent for publication

Not applicable.

Competing interests

No financial and non-financial competing interests.

Author details

¹Sardar Begum Dental College, Gandhara University, Peshawar, Pakistan

²Dental Surgery Department, Qazi Hussain Ahmed Medical Complex (MTI), Nowshera, Pakistan

³Department of Oral Pathology, Saidu College of Dentistry, Swat, Pakistan

⁴Department of Community Medicine, Saidu Medical College, Swat, Pakistan

⁵Superior University, Lahore, Pakistan

⁶Research & Development Cell, Khyber College of Dentistry, Peshawar, Pakistan

⁷Institute of Public Health and Social Sciences, Khyber Medical University, Peshawar, Pakistan

⁸School of Public Health Faculty of Medicine, Imperial College London, London, UK

Received: 16 February 2023 / Accepted: 13 May 2023

Published online: 02 July 2023

References

- Slots J. Periodontitis: facts, fallacies and the future. *Periodontol* 2000. 2017;75(1):7–23.
- Thornton-Evans G, Eke P, Wei L, Palmer A, Moeti R, Hutchins S, et al. Periodontitis among adults aged ≥ 30 years - United States, 2009–2010. *MMWR Suppl*. 2013;22(3):129–35.
- The Institute for Health Metrics and Evaluation (IHME). Periodontal diseases — Level 4 cause [Internet]. [cited 2023 Apr 4]. Available from: https://www.healthdata.org/results/gbd_summaries/2019/periodontal-diseases-level-4-cause.
- Hajishengallis G, Chavakis T. Local and systemic mechanisms linking periodontal disease and inflammatory comorbidities. *Nat Rev Immunol* 2021;21(7):426–40.
- Isola G, Santonocito S, Distefano A, Polizzi A, Vaccaro M, Raciti G et al. Impact of periodontitis on gingival crevicular fluid miRNAs profiles associated with cardiovascular disease risk. *J Periodontol Res* 2023 Feb 8;58(1):165–74.
- Isola G, Giudice A, Lo, Polizzi A, Alibrandi A, Patini R, Ferlito S. Periodontitis and Tooth Loss Have Negative Systemic Impact on Circulating Progenitor Cell Levels: A Clinical Study. *Genes (Basel)*. 2019;10(12):01–12.
- Ronnie Levine, Catherine Stillman-Lowe. The scientific basis of oral Health Education (BDJ Clinician's Guides). 8th ed. Springer; 2018.
- Petersen PE, Baez RJ, Organization WH. Oral health surveys: basic methods [Internet]. 5th ed. Geneva: World Health Organization; 2013. Available from: <https://apps.who.int/iris/handle/10665/97035>.
- Lertpimonchai A, Rattanasiri S, Arj-Ong Vallibhakara S, Attia J, Thakkinian A. The association between oral hygiene and periodontitis: a systematic review and meta-analysis. *Int Dent J* [Internet]. 2017 Dec 1 [cited 2023 Apr 9];67(6):332–43. Available from: <https://pubmed.ncbi.nlm.nih.gov/28646499/>.
- Sun H, Saeedi P, Karuranga S, Pinkepank M, Ogurtsova K, Duncan BB et al. IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res Clin Pract* [Internet]. 2022;183:109119. Available from: <https://www.sciencedirect.com/science/article/pii/S0168822721004782>.
- Khan MAB, Hashim MJ, King JK, Govender RD, Mustafa H, Al Kaabi J. Epidemiology of type 2 diabetes – global burden of Disease and Forecasted Trends. *J Epidemiol Glob Health*. 2019;10(1):107–11.
- Khader YS, Dauod AS, El-Qaderi SS, Alkafajei A, Batayha WQ. Periodontal status of diabetics compared with nondiabetics: a meta-analysis. *J Diabetes Complications*. 2006;20(1):59–68.
- Genco RJ, Grossi SG, Ho A, Nishimura F, Murayama Y. A proposed model linking inflammation to obesity, diabetes, and Periodontal Infections. *J Periodontol*. 2005;76(11–s):2075–84.

14. Preshaw PM, Taylor JJ, Jaedicke KM, De Jager M, Bikker JW, Selten W et al. Treatment of periodontitis reduces systemic inflammation in type 2 diabetes. *J Clin Periodontol*. 2020;47(6):737–46.
15. Pattayil S, Vadakkekuttikal RJ, Radhakrishnan C, Kanakkath H, Hrishi TS. Proportional relationship between periodontal inflamed surface area, clinical attachment loss, and glycated hemoglobin level in patients with type 2 diabetes mellitus on insulin therapy and on oral antidiabetic therapy. *J Periodontol*. 2023;3(1):31–40.
16. Banjar A, Alyafi R, AlGhamdi A, Assaggaf M, Almarghani A, Hassan S et al. The relationship between glycated hemoglobin level and the stage of periodontitis in individuals without diabetes. *PLoS One* 2023;18(1):e0279755.
17. Zhang X, Wang M, Wang X, Qu H, Zhang R, Gu J et al. Relationship between periodontitis and microangiopathy in type 2 diabetes mellitus: a meta-analysis. *J Periodontal Res* 2021;56(6):1019–27.
18. Kesavan R, Chaly P, Reddy Vc M, Av. Periodontal status among type II diabetic and nondiabetic individuals in Chennai, India: a comparative study. *J Indian Association Public Health Dentistry*. 2015;13(4):393–8.
19. SingleCare Team. The 411 on A1C: Normal A1C levels and 15 ways to lower high A1C [Internet]. 2020 [cited 2023 Apr 16]. Available from: <https://www.singlecare.com/blog/normal-a1c-levels/>.
20. Reddy CVK, Maurya M. A comparative study to assess the oral health status and treatment needs of diabetics and non-diabetic population attending some of the hospitals in Mysore City. *Journal of Indian Association of Public Health Dentistry* [Internet]. 2023 [cited 2023 Apr 14];6(12):1–14. Available from: <https://journals.lww.com/aphd/pages/default.aspx/article.asp?issn=2319-5932;year=2008;volume=6;issue=12;spage=1;epage=14;aulast=Reddy;type=0>.
21. Kim EK, Lee SG, Choi YH, Won KC, Moon JS, Merchant AT et al. Association between diabetes-related factors and clinical periodontal parameters in type-2 diabetes mellitus. *BMC Oral Health* 2013;13(1):1–8.
22. Apoorva S, Sridhar N, Suchetha A. Prevalence and severity of periodontal disease in type 2 diabetes mellitus (non-insulin-dependent diabetes mellitus) patients in Bangalore city: an epidemiological study. *J Indian Soc Periodontol*. 2013;17(1):25–9.
23. Khader YS, Albashaireh ZSM, Hammad MM. Periodontal status of type 2 diabetics compared with nondiabetics in north Jordan. *East Mediterr Health J*. 2008;14(3):654–61.
24. Mohamed HG, Idris SB, Ahmed MF, Bøe OE, Mustafa K, Ibrahim SO et al. Association between Oral Health Status and Type 2 Diabetes Mellitus among Sudanese Adults: A Matched Case-Control Study. *PLoS One* [Internet]. 2013 Dec 11 [cited 2023 Apr 14];8(12):e82158. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0082158>.
25. Javed F, Näsström K, Benchimol D, Altamash M, Klinge B, Engström PE. Comparison of Periodontal and Socioeconomic Status between subjects with type 2 diabetes Mellitus and non-diabetic controls. *J Periodontol*. 2007;78(11):2112–9.
26. Chee B, Park B, Bartold MP. Periodontitis and type II diabetes: a two-way relationship. *Int J Evid Based Healthc*. 2013;11(4):317–29.
27. Lalla E, Papapanou PN. Diabetes mellitus and periodontitis: a tale of two common interrelated diseases. *Nat Rev Endocrinol* 2011;7(12):738–48.
28. Teles F, Wang Y, Hajishengallis G, Hasturk H, Marchesan JT. Impact of systemic factors in shaping the periodontal microbiome. *Periodontol* 2000. 2021;85(1):126–60.
29. Wu C, Yuan Y hang, Liu H hang, Li S sui, Zhang B wen, Chen W et al. Epidemiologic relationship between periodontitis and type 2 diabetes mellitus. *BMC Oral Health*. 2020;20(1):01–15.
30. Preshaw PM, Alba AL, Herrera D, Jepsen S, Konstantinidis A, Makrilakis K et al. Periodontitis and diabetes: a two-way relationship. *Diabetologia* 2012;55(1):21–31.
31. Buranasin P, Mizutani K, Iwasaki K, Pawaputanon Na Mahasarakham C, Kido D, Takeda K et al. High glucose-induced oxidative stress impairs proliferation and migration of human gingival fibroblasts. *PLoS One*. 2018;13(8):e0201855.
32. Sandini Trentin M, De João; P, De M, Ferreira C, Diego ;, et al. Prevalence and Severity of Periodontal Disease in Type 2 Diabetes Mellitus Patients: a Cross-Sectional Study. *Biosci J*. 2018;4(34):1114–23.
33. Kaur G, Holtfreter B, Rathmann WG, Schwahn C, Wallaschofski H, Schipf S et al. Association between type 1 and type 2 diabetes with periodontal disease and tooth loss. *J Clin Periodontol* [Internet]. 2009 Sep 1;36(9):765–74. Available from: <https://doi.org/10.1111/j.1600-051X.2009.01445.x>.
34. Pabisch S, Akabane C, Wagermaier W, Roschger A, Ogura T, Hyodo R, et al. The nanostructure of murine alveolar bone and its changes due to type 2 diabetes. *J Struct Biol*. 2016;196(2):223–31.
35. Farooq A, Riasat M, Baloch HKN, Shah SGS, Shehzad S, Shah SZEHS. Severity of alveolar bone loss in control and uncontrolled type II diabetics. *Pakistan J Med Health Sci* 2021;15(6):1459–62.
36. Hintao J, Teanpaisan R, Chongsuvivatwong V, Dahlen G, Rattarasarn C. Root surface and coronal caries in adults with type 2 diabetes mellitus. *Community Dent Oral Epidemiol*. 2007;35(4):302–9.
37. Chang Y, Lee JS, Lee KJ, Woo HG, Song TJ. Improved oral hygiene is associated with decreased risk of new-onset diabetes: a nationwide population-based cohort study. *Diabetologia*. 2020;63(2):924–33.
38. Fu W, Lv C, Zou L, Song F, Zeng X, Wang C et al. Meta-analysis on the association between the frequency of tooth brushing and diabetes mellitus risk. *Diabetes Metab Res Rev* 2019;35(5):e3141.
39. Nishihara U, Tanabe N, Nakamura T, Okada Y, Nishida T, Akihara S. A periodontal disease care program for patients with type 2 diabetes: a randomized controlled trial. *J Gen Fam Med*. 2017;18(5):249–57.
40. Greenblatt AP, Salazar CR, Northridge ME, Kaplan RC, Taylor GW, Finlayson TL et al. Association of diabetes with tooth loss in Hispanic/Latino adults: findings from the Hispanic Community Health Study/Study of Latinos. *BMJ Open Diabetes Res Care*. 2016;4(1):e000211.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.