



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/221319/>

Version: Published Version

Article:

Delanerolle, G., Phiri, P., Elneil, S. et al. (2025) Menopause: a global health and wellbeing issue that needs urgent attention. *The Lancet Global Health*, 13 (2). e196-e198. ISSN: 2214-109X

[https://doi.org/10.1016/s2214-109x\(24\)00528-x](https://doi.org/10.1016/s2214-109x(24)00528-x)

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Menopause: a global health and wellbeing issue that needs urgent attention



Menopause, whether natural (ie, the permanent cessation of an individual's menstrual cycle for at least 12 consecutive months, without any other obvious cause), surgical, or induced by medical treatments, is characterised by the permanent cessation of menstruation. Menopause affects all those assigned female at birth, hereafter referred to as "women". Common symptoms include hot flushes, night sweats, joint pain, low mood, vaginal dryness, and sleep disturbances. Menopause is also associated with several long-term health risks, perhaps the most notable of which is cardiovascular disease. Cardiovascular disease is the leading cause of death in women and an important concern during menopause, when low oestrogen concentrations increase risks of coronary artery disease and atherosclerosis. A systematic review from five countries and regions reported that—compared with women who had menopause aged 50–51 years—women with premature menopause (ie, aged <40 years) had a 55% increased risk of a non-fatal cardiovascular disease event, and women with early menopause (ie, aged 40–44 years) had a 30% increased risk.¹ Osteoporosis is another major health concern, leading to fractures and increased rates of disability, particularly within health-care systems that have poor access to bone health services. A study from Sweden reported that women with early menopause (ie, aged ≤47 years) are at an 83% increased risk of osteoporosis, 68% increased risk of fracture, and 59% higher mortality compared with women who had menopause after 47 years.² Urogenital atrophy is another common complication of menopause, and it can lead to vaginal dryness, urinary incontinence, and a heightened risk of urinary tract infections.³

Despite the severity of acute symptoms and the serious implications of menopause for women's long-term health, access to menopause care varies widely and is particularly scarce in low-income and middle-income countries (LMICs) with little access to health-care professionals with specialist knowledge of menopause.⁴ Additionally, cultural and societal factors in some regions can prevent individuals from seeking care, further complicating the management of menopause-related

health issues and leading to inadequate support in the workplace and even early retirement.⁵ The stigma is such that open dialogue is inhibited in the majority of LMICs and the first-generation migrants from these LMICs to high-income countries (HICs); the paucity of broader understanding and acceptance in society exacerbates the mental and emotional strain of menopause and can lead to social isolation, particularly in these stigmatised contexts.^{6,7} The various factors in LMICs that contribute to adverse health outcomes in women with menopause—for effective and timely intervention, to prevent these outcomes—are important to understand.

As life expectancy rises globally, more individuals spend more of their lives in the post-menopausal phase.⁸ According to WHO, by 2030, over 1.2 billion women worldwide will be menopausal or post-menopausal.⁸ This demographic shift, along with the range and impact of the consequences of menopause, amplify menopause as a global health and wellbeing issue and clarify the need for more effective management and equitable access to care.⁸

The future of menopausal care needs to be inclusive, personalised, and gender-sensitive, addressing unique experiences across all ethnicities and races. Historically, research and clinical guidelines have predominantly focused on cisgender, White women residing in HICs, overlooking the distinct needs of transgender men and women and non-binary individuals, First Nation populations, and people residing in LMICs, particularly those living in rural areas. Given the rapid increase in migration, HIC health-care systems also need to be prepared to better understand the needs of first-generation immigrants. To provide comprehensive support, there must be a greater understanding of intersectional care that accounts for diverse social and cultural influences and considers disease sequelae and their impact on health outcomes. For individuals who are transgender, especially those having long-term hormone therapy or gender-affirming surgeries, the management of hormone levels is a crucial aspect of menopause care.⁹ Hormone replacement therapy in all groups must be carefully tailored to balance the physiological and psychological impacts of menopause.

Lancet Glob Health 2024

Published Online
December 18, 2024
[https://doi.org/10.1016/S2214-109X\(24\)00528-X](https://doi.org/10.1016/S2214-109X(24)00528-X)

We provide eight key recommendations, supported by latest evidence, for the consideration of policy makers, commissioners, researchers, and clinicians (panel).

See Online for appendix

Panel: Recommendations to improve health outcomes for individuals with menopause

Lifestyle modifications

Encouraging regular weight-bearing exercise, such as walking or strength training, to help reduce the risk of osteoporosis. A healthy diet can further protect bone and cardiovascular health.¹⁰

Mental health support

Mental health should be a core component of menopause care given the risk of negative psychological impacts that it confers. Cognitive-behavioural therapy has shown initial evidence of helping with management of vasomotor symptoms, mood fluctuations, and sleep issues related to menopause,⁷ although further research is required.

Holistic approaches

Alternative therapies, including mindfulness, yoga, and acupuncture, have been studied for their potential to reduce hot flushes, anxiety, and sleep disturbances. Although more research is needed to determine whether these practices provide clinically significant benefits, these approaches can be considered as part of a holistic care plan alongside a health-care plan or as a stand-alone approach.

Culturally competent care

Health-care providers should be trained appropriately, particularly to prevent misgendering and to prevent barriers to accessing menopause care. Culturally competent care should include hormone management, mental health support, and reproductive health counselling.

Personalisation of hormone replacement therapy care

For many individuals experiencing menopause, hormone replacement therapy remains an effective treatment, but the choice of treatment should consider individual factors, such as family history, lifestyle, severity of symptoms, and cultural practices (eg, the use of primrose oil or other non-pharmacological remedies). For transgender men, adjustments to testosterone therapy might be required to manage menopause-like symptoms, such as hot flushes, mood changes, and bone loss. Similarly, transgender women who are taking oestrogen therapy should have their hormone levels monitored to ensure adequate cardiovascular and bone protection.

Peer support groups or social prescribing

Creating safe spaces for all individuals to share their experiences of menopause can provide emotional support and reduce isolation. Peer support groups offer a sense of community and help while navigating the complexities of hormone management and menopausal symptoms.

Conduct comprehensive research

The development of proactive, evidence-based interventions are crucial to improve menopause care relevant to all genders, ethnicities, and races.

Improve accessibility

Improve access of hard-to-reach populations (eg, those in rural regions) with technology, such as artificial intelligence, awareness programmes using electronic health tools, and improved monitoring of symptoms and treatment effects using telemedicine and personalised devices.

Understanding various risk factors, such as lifestyles, environment, and climate

Risk factors, such as lifestyles, environmental exposures, and adverse climatic conditions, aggravate adverse health outcomes in individuals experiencing menopause. Any intervention for individuals with menopause can only be effective if the local risk factors of individuals living in unfavourable conditions are taken into account while designing and implementing the intervention.

Menopause care must adopt an inclusive and personalised approach, as has been done for other conditions, to address the unique needs of both cisgender and transgender populations across global settings and to integrate diverse cultural practices. However, to truly improve menopause care, we must also work towards destigmatising menopause. Destigmatisation requires a global, multisectoral approach with comprehensive research, improved health-care access, practical workplace policies, and coherent messaging and communication to enable a more open and supportive society.

We declare no competing interests. Members of the MARIE collaborative are listed in the appendix.

Copyright © 2024 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Gayathri Delanerolle, Peter Phiri, Sohler Elneil, Vikram Talaulikar, George U Eleje, Rabia Kareem, Ashish Shetty, Lucky Saraswath, *Om Kurmi, Cristina Laguna Benetti-Pinto, Ifran Muhammad, Nirmala Rathnayake, Teck-Hock Toh, Ieera Madan Aggarwal, Jian Qing Shi, Julie Taylor, Kathleen Riach, Kristina Potocnik, Ian Litchfield, Helen Felicity Kemp, Paula Briggs, and the MARIE collaborative
om.kurmi@coventry.ac.uk

Institute of Applied Health Sciences, University of Birmingham, Birmingham, UK (GD, JT, IL); Hampshire and Isle of Wight Healthcare NHS Foundation Trust, Southampton, UK (PP); EGA Institute for Women's Health, University College London, London, UK (SE, VT, AS); University College London NHS Foundation Trust, London, UK (SE, VT, AS); Allied Health Sciences, University of Ruhuna, Mataara, Sri Lanka (NR); Liverpool Women's NHS Trust, Liverpool, UK (PB); University of Edinburgh, Edinburgh, UK (KP); Southern University of Science and Technology, Shenzhen, China (JQS); Sibiu Hospital, Sibiu, Malaysia (T-HT); Department of Gynaecology, Universidade Estadual de Campinas, São Paulo, Brazil (CLB-P); Department of Obstetrics and Gynaecology, Nnamdi Azikiwe University, Awka, Nigeria (JUE); KK Women's and Children Hospital, Singapore (IMA); Peshawar Medical College, Riphah International University, Islamabad, Pakistan (RK, IM); Dugald Baird Centre for Research on Women's Health, University of Aberdeen, Aberdeen, UK (LS); Centre for Healthcare and Communities, Coventry University, Coventry CV1 5FB, UK (OK); Division of Respiriology, Department of Medicine, McMaster University, Hamilton, ON, Canada (OK); Adam Smith Business School, University of Glasgow, Glasgow, UK (KR); Edinburgh, UK (HFK)

- Zhu D, Chung H-F, Dobson AJ, et al. Age at natural menopause and risk of incident cardiovascular disease: a pooled analysis of individual patient data. *Lancet Public Health* 2019; **4**: e553-64.
- Svejme O, Ahlborg HG, Nilsson J-Å, Karlsson MK. Early menopause and risk of osteoporosis, fracture and mortality: a 34-year prospective observational study in 390 women. *BJOG* 2012; **119**: 810-16.
- Briggs P, Delanerolle G, Burton R, Shi JQ, Hamoda H, Hapangama DK. The silent epidemic of urogenital atrophy. *Br J Gen Pract* 2021; **71**: 538-39.
- Sriram V, Bennett S. Strengthening medical specialisation policy in low-income and middle-income countries. *BMJ Glob Health* 2020; **5**: e002053.
- Huang DR, Goodship A, Webber I, et al. Experience and severity of menopause symptoms and effects on health-seeking behaviours: a cross-sectional online survey of community dwelling adults in the United Kingdom. *BMC Womens Health* 2023; **23**: 373.

- 6 Klotzbaugh R, Fawcett J. Gender minority persons' perceptions of peer-led support groups: a Roy Adaptation Model interpretation. *ANS Adv Nurs Sci* 2023; **46**: 59–74.
- 7 Hunter MS. Cognitive behavioral interventions for the treatment of menopausal symptoms. *Expert Rev Obstet Gynecol* 2012; **7**: 321–26.
- 8 WHO. Menopause. Oct 16, 2024. <https://www.who.int/news-room/fact-sheets/detail/menopause#:~:text=The%20regularity%20and%20length%20of,55%20years%20for%20women%20worldwide> (accessed Oct 25, 2024).
- 9 Deutsch MB. Guidelines for the primary and gender-affirming care of transgender and gender nonbinary people. June 17, 2016. <https://transcare.ucsf.edu/guidelines> (accessed Oct 28, 2024).
- 10 Elavsky S, McAuley E. Physical activity and mental health outcomes during menopause: a randomized controlled trial. *Ann Behav Med* 2007; **33**: 132–42.