



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/221422/>

Version: Published Version

Article:

Pokotheoane, Retselisitsoe, Argefa, Terefe, Miderho, Christus Cito et al. (2025) Prevalence and determinants of tobacco use among school-going adolescents in 53 African countries: Evidence from global youth tobacco surveys. *Addictive Behaviors Reports*. 100581. ISSN: 2352-8532

<https://doi.org/10.1016/j.abrep.2024.100581>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Prevalence and determinants of tobacco use among school-going adolescents in 53 African countries: Evidence from the Global Youth Tobacco Survey

Retselisitsoe Pokothoane^{a,b,*} , Terefe Gelibo Agerfa^{a,c} , Christus Cito Miderho^d ,
Noreen Dadirai Mdege^{a,e,f} 

^a Development Gateway: an IREX Venture, Washington, DC, United States

^b Research Unit on the Economics of Excisable Products (REEP), School of Economics, University of Cape Town, Cape Town, South Africa

^c Public Health, ICAP at Columbia University Mailman School of Public Health, Addis Ababa, Ethiopia

^d Département d'Agro-Vétérinaire, Institut Supérieur Pédagogique de Bukavu (ISP/Bukavu), BP 854 Bukavu, Democratic Republic of the Congo

^e Department of Health Sciences, University of York, York, United Kingdom

^f Centre for Research in Health and Development, York, United Kingdom

ARTICLE INFO

Keywords:

Adolescents
Africa
Cigarettes
GYTS
Shisha
Tobacco

ABSTRACT

Introduction: Tobacco use typically begins during adolescence. There is a lack of comprehensive evidence on the use of different tobacco products among adolescents in Africa.

Aims and Methods: We used the most recent Global Youth Tobacco Surveys from 53 African countries, covering 2003–2020, to estimate the overall and gender-specific prevalence of each type of tobacco product by country, Africa region, World Bank income group, and age group among adolescents aged 11–17 years. We further used Logit regressions to assess the determinants of using different tobacco products.

Results: The overall prevalence of any tobacco use among adolescents was 14.3 % [95 % CI: 13.5, 15.3]. Specifically, the prevalence for cigarettes was 6.4 % [95 % CI: 5.9, 7.0], for other smoked tobacco was 6.7 % [95 % CI: 6.0, 7.4], for smokeless tobacco use was 6.4 % [95 % CI: 5.9, 6.9], and for shisha smoking was 5.2 % [95 % CI: 4.4, 6.1]. The prevalence of dual use of smoked and smokeless tobacco was 3.0 % [95 % CI: 2.8, 3.2], and that of shisha and cigarettes was 1.5 % [95 % CI: 1.2, 2.0]. Any tobacco use prevalence was higher among boys (17.4 %) than girls (10.6 %). Seeing health warnings about tobacco dangers, exposure to smoking at home and school, the age restriction to tobacco purchases, and peer pressure were positively associated with the use of all tobacco products. Being a female was negatively associated with tobacco use across all products.

Conclusions: Policymakers should prioritize implementing large pictorial health warnings about tobacco dangers covering the entire packaging of different products.

1. Introduction

There are over 8 million deaths globally each year due to tobacco use and second hand tobacco smoke exposure (World Health Organization, 2023c). If no effective tobacco control measures are taken, the number of deaths will increase to as high as 1 billion this century (Drope et al., 2018). Tobacco use typically starts during adolescence (i.e., 10–19 years age period) (Sawyer et al., 2018), and adolescents are easily addicted and ultimately exposed to long-term nicotine addiction and use. Thus, adolescent smoking often leads to prolonged adult smoking tobacco use

(Nonnemaker & Farrelly, 2011; World Health Organization, 2016). If individuals do not initiate tobacco use during the adolescent stage, they are more likely not to even start using tobacco in adulthood (Mayhew et al., 2000). The Theory of Triadic Influence (TTI) states that tobacco use among adolescents can be explained by factors that fall within three tiers. The first tier argues that demographic factors such as age and gender are associated with adolescent's decisions to use tobacco. The second tier states that social environmental factors such as friends, family members, neighbours, and schoolmates play a crucial role in influencing adolescents to use tobacco products. The last tier states that

* Corresponding author at: Retselisitsoe Pokothoane, Research Unit on the Economics of Excisable Products (REEP), School of Economics, University of Cape Town, Cape Town, South Africa.

E-mail address: retselisitsoepokothoane@gmail.com (R. Pokothoane).

<https://doi.org/10.1016/j.abrep.2024.100581>

Received 31 May 2024; Received in revised form 16 December 2024; Accepted 30 December 2024

Available online 2 January 2025

2352-8532/© 2025 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

broader factors, such as government policy on tobacco control (e.g., tobacco taxation and educational interventions) and tobacco industry advertising are candidate explanations for adolescent tobacco use (Flay et al., 1995).

In Africa, the tobacco industry is powerful and utilizes various tactics that increasingly target adolescents through the use of role models or attractive flavours, for example (Drope et al., 2018). The industry also employs sales strategies like the sale of single cigarette sticks or disposable electronic cigarettes to make them more affordable and easily accessible to adolescents. The tobacco industry also engages in corporate social responsibility activities, such as offering sports sponsorships, academic scholarships, and youth entrepreneurial schemes to promote brand recognition and create a positive perception among minors (World Health Organization, 2024).

It is, therefore, important to monitor the use of tobacco products among young people to have the data needed to inform effective tobacco control policies. For low- and middle-income countries, the school-based Global Youth Tobacco Survey (GYTS) provides most of the available national-level data on tobacco use among adolescents (World Health Organization, 2023a). The GYTS is a school-based survey conducted among primary/secondary school-going adolescents who are in grades that are associated with the age group 13–15 years. It collects information on tobacco use among adolescents across the globe using a two-stage cluster sample design. However, despite most GYTS datasets covering the age group 11–17 years, the most recent cross-country analyses that include datasets from African countries have focused on the 13–15 or 12–16 years age groups (James et al., 2022; Ma et al., 2021, 2022; Yang et al., 2022). In addition, most of these analyses either consider the tobacco products together without disaggregated data on the different tobacco products (James et al., 2022), or consider only a few products (e.g., just cigarettes and other smoked tobacco (Ma et al., 2021), smokeless tobacco (Yang et al., 2022) or shisha (waterpipe) (Ma et al., 2022)). In addition, these studies only include datasets from 11 to 38 African countries.

Our study, therefore, aims to, for the first time, provide information that covers the 11–17 years age group, disaggregated by tobacco product, and covers all African countries where at least one GYTS has been conducted. We focus on the 11–17-year age group because this is a broader subset of the official adolescent age group (10–19), which has not been examined before. We also assess the dual use of smoked and smokeless tobacco products, as well as of cigarettes and shisha, and investigate the factors associated with the use of different tobacco products. To the best of our knowledge, this is the most comprehensive study to date to explore the prevalence of the use of tobacco products and their associated factors in Africa.

2. Methods

2.1. Data sources

This study uses the latest GYTS data from each of the 53 African countries that have conducted at least one round of the survey (World Health Organization, 2023b). These datasets covered the period 2003–2020. South Sudan is the only African country that has not conducted a GYTS. The GYTS is collected through a two-stage cluster sampling design. In the first stage, schools are selected proportional to the school enrolment sizes. The second stage involves the random sampling of classes, and all students within the selected classes are eligible to participate in the self-administered questionnaire. The GYTS covers many topics related to tobacco use: 1) the prevalence of cigarette smoking and other tobacco use, 2) knowledge and attitudes of young people towards smoking, 3) the role of the media and advertising in your people's use of tobacco products, 4) access to tobacco products, 5) tobacco-related school curriculum, and 6) second-hand smoke of tobacco, and 7) cessation of cigarette smoking (World Health Organization, 2023b).

Most of the 53 datasets comprise school-going adolescents in the age group, 11–17 years, with the exception of those for Carbo Verde, Chad, Ethiopia, and South Africa, which also include those outside this age group. We therefore excluded any respondents older than 17 years from the analysis for consistency across countries.

2.2. Outcomes of interest

Our outcome variables of interest were current:

- Cigarette smoking status: this was obtained from the responses to one of the following questions: “During the past 30 days, on how many days did you smoke cigarettes?” or “During the past 30 days, did you smoke any cigarettes?”.
- Shisha smoking status: this was obtained from the responses to one of the following questions: “During the past 30 days, on how many days did you smoke shisha?” or “During the past 30 days, did you smoke shisha?”
- Use of other smoked tobacco products: this was obtained from the responses to one of the following questions: “During the past 30 days, on how many days did you smoke any smoked tobacco products other than cigarettes?” or “During the past 30 days, did you smoke any of the smoked tobacco products other than cigarettes?”
- Smokeless tobacco use status: this was obtained from the responses to one of the following questions: “During the past 30 days, on how many days did you use any smokeless tobacco products?” or “During the past 30 days, did you use any smokeless tobacco products?”
- Dual use of smoked and smokeless tobacco: individuals who had been classified as current smokers of cigarettes, shisha or other smoked tobacco products, AND current users of smokeless tobacco products were dual users of smoked and smokeless tobacco. Those who only currently used smoked products, as well as those who only currently used smokeless tobacco products were not dual users of smoked and smokeless tobacco.
- Dual use of cigarettes and shisha: individuals who had been classified as current smokers of cigarettes, AND current smokers of shisha were current dual users of cigarettes and shisha. Those who only currently used one of these products were not dual users of cigarettes and shisha.
- Use of any tobacco product: individuals who had been classified as current users of any tobacco product, either smoked or smokeless, were classified as current users of any tobacco product. Those who did not report using any tobacco product were classified as non-users of tobacco.

For current cigarette smoking, shisha smoking, use of other smoked tobacco products and smokeless tobacco use, those whose response was one day or more to the question on the number of days they used a product in the past 30 days were classified as current smokers/users of the product, whilst those whose response was zero were current non-smokers/users. When using the question that asked whether they used a product during the past 30 days, those who answered ‘yes’ were current smokers/users of the product, whilst those who answered ‘no’ were current non-smokers/users.

Each of the outcomes of interest was defined as a dummy variable, where 1 denoted a current user and 0 a current non-user. We were also interested in heated tobacco products (HTPs) where tobacco is heated without combustion to produce aerosols containing nicotine and other chemicals, which are then inhaled by users (World Health Organization, 2020). However, none of the surveys had data on these products.

2.3. Potential predictors

2.3.1. Demographic variables

Guided by the Theory of Triadic Influence (Flay et al., 1995) and the empirical literature on significant factors associated with tobacco use

and the GYTS questionnaires, our potential predictors included demographic, pro-tobacco, and anti-tobacco-use variables (James et al., 2022; Ma et al., 2021, 2022; Yang et al., 2022). The demographic variables include gender, age, age squared, Africa region, and World Bank income classification. In line with (de la Torre-Luque et al., 2021), we also include the predominant religion in each African country (Pew Research Center Religion & Public Life, 2015).

2.3.2. Anti-tobacco use variables

We use 4 types of anti-tobacco use variables:

- Age restriction: this dummy variable was generated from the question, “During the past 30 days, did anyone refuse to sell you cigarettes because of your age?”
- Exposure to anti-tobacco media messages – a dummy variable created from the question, “During the past 30 days, did you use or hear any anti-tobacco messages on TV, radio, billboards, posters, newspapers, magazines or movies?”
- School teaching on tobacco use: this was a dummy variable generated from the question, “During the past 12 months, were you taught in any of your classes about the dangers of tobacco use?”
- Health warnings on the dangers of tobacco use: this was a dummy variable generated from the question, “During the past 30 days, did you see any health warnings on cigarette/tobacco packages?”

2.3.3. Pro-tobacco use variables

We use the following 7 pro-tobacco use variables:

- Media exposure to smoking: a dummy variable generated from the question, “During the past 30 days, did you see any people using tobacco when you watched TV, videos or movies?”
- Tobacco adverts: a dummy variable generated from the question, “During the past 30 days, did you see any advertisements or promotions for tobacco products at points of sale (such as street vendors/aprons, shops, kiosks, travelling carts, etc)?”
- Offer of free tobacco products: a dummy variable generated from the question, “Has a person working for the tobacco product company ever offered you a free tobacco product?”
- Public exposure to tobacco use/smoking: a dummy variable generated from the question, “During the past 7 days, on how many days has anyone smoked in your presence, at any outdoor public place such as football or basketball, fields, sidewalks, entrances to buildings, malls, parking lots, beaches, stadiums, etc?”
- School exposure to tobacco smoking: a dummy variable generated from the question, “During the past 30 days, did you see anyone smoke inside the school building or outside on school property?”
- Home exposure to tobacco smoking: a dummy variable is generated from the question, “During the past 7 days, on how many days has anyone smoked inside your home in your presence?”
- Peer/friendship smoking: a dummy variable generated from the question, “If one of your best friends offered you a tobacco product, would you use it?”

2.4. Data analysis

The GYTS datasets for the 53 countries were retrieved and homogenized in terms of key study variables, and combined into one dataset with a total sample of 204,537 school-going adolescents aged 11–17 years. We used individual-level data to compute descriptive statistics and prevalence estimates. We computed the prevalence estimates for each of the outcomes of interest for Africa, as well as by country, African region (i.e., Central, East, North, Southern, and West) (African-Union, 2023), World Bank income group classification (Hamadeh et al., 2023) as well as by gender and age group. Reported prevalence estimates include the associated 95 % confidence intervals (CIs). We adjusted for the cluster variable (schools) and stratification in the variance

estimation to account for the complex sampling design, and applied sample weights to account for differential probabilities of selection and participation. Data analyses were performed in Stata version 17 (StataCorp, 2021).

To assess the factors associated with tobacco use, we use data from the GYTS surveys conducted from 2013 to 2020 to ensure that we have consistent information for all variables used, including the potential predictors. This is because the GYTS questionnaire changed after 2012. Given that our outcomes of interest are binary, we estimate Logit regression models separately for each of the 6 products and product combinations (cigarette smoking, other smoked tobacco use, smokeless tobacco use, shisha smoking, dual cigarettes and shisha smoking, and dual smokeless and smoked tobacco). We present the odds ratios from the Logit models. We also provide the Logit model’s marginal effects in [Supplementary Table 9](#).

2.5. Sensitivity analysis for prevalence estimates

Data used in this study spans the period 2003 – 2020. It is possible that, for countries with older datasets, the prevalence estimates generated might not reflect the present scenario. In order to account for this, we performed a sensitivity analysis where we excluded countries with surveys conducted before 2010.

2.6. Ethics and information approval

The GYTS data is freely available on the WHO microdata repository, and the public is permitted to use it. However, we acknowledge that the original data is owned by Centres for Disease Control and Prevention (CDC) and WHO. We have also referenced the data source appropriately.

3. Results

3.1. Sample characteristics

The mean age of adolescents in the sample was 14.3 years (standard deviation = 2.2 years). About 42 % of the adolescents were aged either 14 or 15 years, and there was a 50/50 gender split between boys and girls. Close to one-third (31.89 %) of the sample was from West Africa, and approximately half (48.59 %) came from lower-middle-income African countries. About 20 % of the sample was from surveys conducted in 2008 ([Table 1](#)). The highest proportion of observations was from Zimbabwe (6.3 %), and the least was from Guinea-Bissau (0.7 %) ([Supplementary Table 1](#)). All 53 surveys had data on cigarette smoking status, but only 16 had specific data on shisha smoking status. Forty-one (41) surveys also had aggregated data on ‘other smoked tobacco products’ that are not cigarettes, and 45 surveys reported on smokeless tobacco use. Notably, no Southern African country had data on shisha smoking ([Supplementary Table 2](#)).

3.2. Prevalence of tobacco use

The overall prevalence of any tobacco use was 14.3 % [95 % CI: 13.5, 15.3] ([Table 2](#)). This was higher for boys (17.4 % [95 % CI: 16.3, 18.6]) than for girls (10.6 % [95 % CI: 9.5, 11.7]). The overall prevalence, as well as those for boys and girls, were highest in Southern Africa and lowest in East Africa. Upper middle-income countries had the highest, and low-income countries had the lowest overall prevalence of the use of any tobacco product. Prevalences of any tobacco use by country and gender are provided in [Supplementary Table 3](#). The lowest prevalence was in Angola (1.2 % [95 % CI: 0.7, 2.0]) and the highest was in South Africa (35.2 % [95 % CI: 32.8, 37.8]). For boys, this ranged from 1.7 % [95 % CI: 0.9, 3.1] in Angola to 40.4 % [95 % CI: 36.9, 44.1] in South Africa. Among girls, the prevalence ranged from 0.4 % [95 % CI: 0.1, 1.4] in Angola to 30 % [95 % CI: 27.8, 32.3] in South Africa. Product-specific prevalence estimates by region, World Bank income group

Table 1
Sample characteristics.

Variable	Number of observations (n)	Percent (%)
All	204, 537	100 %
Gender		
Boys	99,153	49.75 %
Girls	100,159	50.25 %
Age		
11 years	10,905	5.44 %
12 years	18,526	9.24 %
13 years	34,678	17.30 %
14 years	42,868	21.39 %
15 years	40,538	20.23 %
16 years	28,566	14.25 %
17 years	24,340	12.14 %
African Region		
Central Africa	33,135	16.20 %
East Africa	40,841	19.97 %
North Africa	20,814	10.18 %
Southern Africa	44,529	21.77 %
West Africa	65,218	31.89 %
World Bank Income Group		
Low income	78,520	38.39 %
Lower middle	99,384	48.59 %
Upper middle	24,148	11.81 %
High income	2,485	1.21 %
Survey Year		
2003	5,959	2.94 %
2006	9,639	4.76 %
2007	3,965	1.96 %
2008	38,209	18.86 %
2009	16,140	7.96 %
2010	11,964	5.90 %
2011	11,837	5.84 %
2013	13,645	6.73 %
2014	20,028	9.88 %
2015	5,295	2.61 %
2016	11,896	5.87 %
2017	27,377	13.51 %
2018	10,118	4.99 %
2019	12,250	6.05 %
2020	4,320	2.13 %

Source: Authors' computations from African GYTS (2003–2020)

classification, and age group are also provided in [Table 3](#).

The overall prevalence of use of the different products in Africa was similar for cigarettes (6.4 % [95 % CI: 5.9, 7.0]), shisha (5.2 % [95 % CI: 4.4, 6.1]), other smoked tobacco products (6.7 % [95 % CI: 6.0, 7.4]), and smokeless tobacco products (6.4 % [95 % CI: 5.9, 6.9]). The overall prevalence of the dual use of smokeless & smoked tobacco products was 3.0 % [95 % CI: 2.8, 3.2], and that of the dual use of cigarettes and shisha was 1.5 % [95 % CI: 1.2, 2.0]. Southern Africa had the highest prevalence across all product types, except for shisha smoking where no country in that region has data. North Africa had the highest prevalence for shisha smoking. East Africa had the lowest prevalences for cigarette smoking, smokeless tobacco use and the use of other smoked tobacco products. Central Africa had the lowest prevalence of shisha smoking. Irrespective of gender, the age-specific prevalence estimates for different tobacco products showed a similar pattern: they decreased from age 11 to 13 years, and started to increase from age 13 to 17 years ([Table 3](#)). The overall product-specific prevalence rates by country are provided in [Supplementary Table 4](#).

[Fig. 1](#) presents the regional prevalence estimates for the use of different tobacco products by gender. The prevalence of cigarette smoking was significantly higher among boys than girls in all regions. The prevalence of smokeless tobacco use was higher among boys than girls in 4 regions except Southern Africa, where it was similar. The prevalence rates among girls in Southern Africa were higher than that of boys in other regions for almost all products. Gender-specific country-level estimates for different tobacco products are provided in [Supplementary Table 5](#) for boys and [Supplementary Table 6](#) for girls. For Africa as a whole, tobacco use prevalence was higher among boys than for girls

Table 2
Prevalence of use of any tobacco product by African region, World Bank income group, age group, and gender.

	Use of any tobacco products		
	Boys (95 % CI)	Girls (95 % CI)	Total (95 % CI)
African region			
Central Africa	17.2 % (14.7, 19.9)	10.5 % (8.8, 12.4)	14.5 % (12.5, 16.7)
East Africa	10.9 (9.7, 12.2)	6.3 % (5.5, 7.2)	8.9 % (8.0, 9.9)
North Africa	18.4 % (15.3, 22.0)	7.3 % (4.9, 10.9)	12.9 % (10.7 to 15.5)
Southern Africa	29.7 % (27.7, 32.0)	23.6 % (22.1, 25.1)	27.1 % (25.6, 28.6)
West Africa	15.9 % (14.6, 17.3)	10.4 % (8.8, 12.1)	13.7 % (12.5, 15.0)
Income Group			
High income	24.1 % (20.9, 27.5)	13.6 % (11.2, 16.3)	18.9 % (16.5, 21.5)
Upper middle income	36.2 % (33.4, 39.1)	26.7 % (24.9, 28.5)	31.5 % (29.6, 33.4)
Lower middle income	15.1 % (13.5, 16.9)	7.8 % (6.5, 9.5)	11.8 % (10.6, 13.1)
Low income	14.4 % (13.3, 15.5)	8.7 % (7.6, 9.8)	12.0 % (11.1, 13.0)
Age Group			
11 years	28.9 % (24.3, 34.0)	13.7 % (9.3, 19.9)	21.1 % (17.8, 25.4)
12 years	13.5 % (10.5, 17.3)	7.4 % (4.9, 10.8)	10.3 % (7.9, 13.2)
13 years	12.4 % (10.3, 14.9)	7.0 % (5.8, 8.4)	9.9 % (8.6 %, 11.4)
14 years	14.2 % (12.6, 15.9)	10.7 % (8.7, 13.1)	12.7 % (11.4, 14.1)
15 years	15.8 % (14.1, 17.7)	10.6 % (9.1, 12.2)	13.4 % (12.1, 14.8)
16 years	21.2 % (18.9, 23.7)	12.4 % (10.9, 14.1)	17.3 % (15.8, 19.0)
17 years	27.2 % (24.9, 29.6)	16.7 % (14.7, 19.0)	23.1 % (21.4, 25.0)
Overall prevalence	17.4 % (16.3, 18.6)	10.6 % (9.5, 11.7)	14.3 % (13.5, 15.3)

Source: Authors' computations from African GYTS (2003–2020).

across all product types. Among boys, cigarette smoking was more common (9.2 %), followed by other smoked tobacco (8.3 %), smokeless (7.3 %), and shisha (6.5 %), in that order. Among girls, the prevalence of smokeless tobacco was more common (5.1 %), followed by other smoked (4.8 %), shisha (3.6 %), and cigarettes (3.4 %), in that order.

3.3. Factors associated with the use of tobacco products

3.3.1. Cigarette smoking

For cigarettes, girls were significantly less likely to smoke than boys (odds ratio (OR) 0.3, [95 % CI: 0.2, 0.3]). Adolescents from East Africa (OR 0.4, [95 % CI = 0.3, 0.7]), Southern Africa (OR 0.5, [95 % CI = 0.3, 0.9]), and West Africa (OR 0.5, [95 % CI = 0.3, 1.0]) were less likely to smoke than those from Central Africa. There was no statistically significant difference between North Africa and Central Africa in the likelihood of an adolescent being a cigarette smoker. Adolescents from lower-middle-income African countries were less likely to smoke (OR 0.5, [95 % CI = 0.4, 0.7]) than those from low-income countries. There was no statistically significant difference between upper-middle and low-income countries on the likelihood of an adolescent being a current cigarette smoker ([Table 4](#)).

Adolescents who were refused the purchase of cigarettes because of their age (aged below 18) (either for themselves or on behalf of other people) were more likely to smoke cigarettes (OR 4.7, [95 % CI = 3.6, 6.2]) than those who were not. Those who were exposed to anti-tobacco smoking messages were less likely to smoke (OR 0.8, [95 % CI = 0.6, 0.8]) than those who were not. Adolescents who had seen the health warnings about the dangers of tobacco use were more likely to use

Table 3
Prevalence of use of different tobacco products, by African region, World Bank income group, and age group.

Location	Cigarettes (95 % CI)	Smokeless (95 % CI)	Other smoked (95 % CI)	Shisha (95 % CI)	Dual smokeless & smoked tobacco (95 % CI)	Dual cigarettes & shisha (95 % CI)
African Region						
Central Africa	6.9 % (5.4, 8.7)	7.8 % (6.8, 9.0)	6.7 % (5.5, 8.2)	3.1 % (2.3, 4.1)	3.2 % (2.5, 3.9)	1.0 % (0.8, 1.5)
East Africa	4.2 % (3.6, 4.8)	4.1 % (3.6, 4.7)	3.4 % (2.8, 4.2)	4.5 % (3.7, 5.5)	1.5 % (1.2, 1.8)	1.5 % (1.1, 1.9)
North Africa	5.6 % (4.3, 7.3)	4.5 % (3.4, 5.9)	5.6 % (3.9, 7.9)	6.0 % (4.6, 7.8)	2.0 % (1.7, 2.5)	1.8 % (1.2, 2.9)
Southern Africa	12.7 % (11.4, 14.0)	12.8 % (12.0, 13.7)	14.0 % (13.0, 15.0)	X	7.3 % (6.6, 8.0)	X
West Africa	5.5 % (4.9, 6.1)	6.3 % (5.5, 7.16)	6.2 % (5.5, 6.9)	4.7 % (3.6, 6.1)	2.8 % (2.4, 3.3)	1.0 % (0.7, 1.4)
Income Group						
High income	15.6 % (13.4, 18.2)	1.9 % (1.3, 2.6)	9.8 % (8.2, 11.7)	13.6 % (11.8, 15.6)	1.3 % (0.9, 2.0)	5.9 % (4.8, 7.2)
Upper middle income	15.0 % (13.4, 16.7)	14.3 % (13.3, 15.3)	17.6 % (16.2, 19.1)	4.7 % (3.7, 5.9)	8.4 % (7.6, 9.4)	1.8 % (1.3, 2.6)
Lower middle income	5.1 % (4.4, 6.0)	4.8 % (4.2, 5.5)	4.9 % (4.1, 6.0)	5.6 % (4.6, 6.8)	2.1 % (1.8, 2.3)	1.6 % (1.1, 2.2)
Low income	5.3 % (4.7, 6.0)	6.0 % (5.4, 6.8)	5.8 % (5.0, 6.6)	3.9 % (3.0, 5.0)	2.4 % (2.0, 2.8)	1.2 % (0.9, 1.7)
Age Group						
11 years	7.3 % (5.4, 9.8)	12.6 % (9.3, 16.9)	11.3 % (8.3, 15.4)	9.3 % (6.6, 12.9)	7.0 % (4.7, 10.5)	2.2 % (0.7, 6.1)
12 years	3.4 % (2.6, 4.5)	5.0 % (3.8, 6.7)	5.3 % (3.6, 7.7)	3.3 % (2.0, 5.6)	2.1 % (1.5, 2.8)	1.0 % (0.5, 2.1)
13 years	3.7 % (3.2, 4.3)	3.7 % (3.1, 4.5)	4.7 % (3.4, 6.3)	3.7 % (2.8, 4.9)	1.4 % (1.5, 2.8)	0.5 % (0.4, 0.8)
14 years	5.0 % (4.4, 5.8)	5.5 % (4.5, 6.8)	5.5 % (4.7, 6.5)	4.8 % (3.9, 5.8)	2.2 % (1.9, 2.6)	1.1 % (0.8, 1.6)
15 years	6.5 % (5.4, 7.6)	5.7 % (5.0, 6.4)	6.0 % (5.2, 6.9)	6.0 % (4.5, 8.1)	2.8 % (2.3, 3.3)	2.0 % (1.2, 3.2)
16 years	8.0 % (7.0, 9.1)	8.2 % (7.1, 9.5)	8.1 % (7.2, 9.0)	5.8 % (4.6, 7.3)	3.8 % (3.2, 4.4)	2.5 % (1.6, 3.7)
17 years	14.0 % (12.6, 15.5)	10.0 % (9.0, 11.0)	11.6 % (10.3, 13.0)	6.5 % (5.0, 8.3)	6.1 % (5.4, 6.9)	2.8 % (1.8, 4.2)
Overall Prevalence	6.4 % (5.9, 7.0)	6.4 % (5.9, 6.9)	6.7 % (6.0, 7.4)	5.2 % (4.4, 6.1)	3.0 % (2.8, 3.2)	1.5 % (1.2, 2.0)

Source: Authors' computations from African GYTS (2003–2020). Notes: X indicates that data was not available.




Fig. 1. Regional prevalence of the use of different tobacco products among adolescents (%) by gender. Source: Authors' computations from African GYTS (2003–2020).

cigarettes (OR 2.0, 95 % CI = 1.5, 2.7]) than those who had not (Table 4).

When considering the pro-tobacco use variables, adolescents who had been offered free tobacco products were more likely to smoke (OR

1.8, [95 % CI = 1.3, 2.5]) than those who had not. Exposure to tobacco use in public places (OR 2.2, [95 % CI= (1.7, 2.9)]), at school (OR 2.0, [95 % CI: 1.5, 2.7]), and at home (OR 2.5, [95 % CI: 1.8, 3.4]) had higher of odds of cigarette smoking. Having smoking friends/peers was

Table 4
Factors associated with cigarette smoking, other smoked and smokeless tobacco use.

	Cigarettes	Other Smoked Tobacco	Smokeless Tobacco
	Odds Ratio	Odds Ratio	Odds Ratio
VARIABLES	(95 % CI)	(95 % CI)	(95 % CI)
Age	0.4 (0.1, 1.4)	0.3 (0.1, 1.3)	0.7 (0.1, 5.1)
Age Squared	1.0 (0.9, 1.1)	1.0 (1.0, 1.1)	1.0 (1.0, 1.1)
Female (Baseline: Male)	0.3* (0.2, 0.3)	0.4* (0.3, 0.6)	0.9 (0.6, 1.3)
<i>Region (Base: Central Africa)</i>			
East Africa	0.4* (0.3, 0.7)	0.8 (0.5, 1.3)	0.7* (0.5, 1.0)
North Africa	0.6 (0.3, 1.1)	1.1 (0.6, 2.0)	0.9 (0.6, 1.4)
Southern Africa	0.5* (0.3, 0.9)	1.1 (0.7, 1.7)	0.9 (0.6, 1.3)
West Africa	0.5* (0.3, 1.0)	1.3 (0.9, 1.8)	0.7 (0.4, 1.2)
<i>World Bank Income Group (Base: Low income)</i>			
Lower middle income	0.5* (0.4, 0.7)	0.9 (0.7, 1.3)	0.9 (0.7, 1.3)
Upper middle income	1.4 (0.9, 2.0)	1.5 (1.0, 2.4)	0.7 (0.4, 1.1)
<i>Christian (Base: Muslim & Hindu)</i>			
	0.8 (0.6, 1.2)	0.8* (0.6, 1.0)	1.0 (0.7, 1.5)
Age Restriction	4.7* (3.6, 6.2)	2.4* (1.6, 3.5)	2.4* (1.7, 3.5)
Anti-tobacco use messages	0.7* (0.6, 0.8)	1.183 (0.9, 1.5)	1.0 (0.7, 1.3)
School teaching on tobacco use	0.9 (0.7, 1.1)	1.0 (0.7, 1.5)	1.2 (0.8, 1.6)
Health warnings	2.0* (1.5, 2.7)	1.5* (1.2, 1.9)	1.8* (1.3, 2.4)
Media exposure to smoking	1.1 (0.9, 1.3)	0.9 (0.6, 1.2)	0.8 (0.6, 1.1)
Tobacco adverts	0.9 (0.7, 1.2)	1.2 (0.8, 1.8)	1.1 (0.8, 1.6)
Free tobacco product offer	1.8* (1.3, 2.5)	3.7* (2.6, 5.3)	4.6* (3.3, 6.4)
Public exposure to smoking	2.2* (1.7, 2.9)	0.9 (0.6, 1.5)	1.1 (0.7, 1.6)
School exposure to smoking	2.0* (1.5, 2.7)	1.6* (1.1, 2.3)	1.7* (1.3, 2.3)
Home exposure to smoking	2.5* (1.8, 3.4)	1.9* (1.2, 2.8)	1.2 (0.9, 1.7)
Peer smoking	8.7* (6.9, 11.0)	2.7* (2.1, 3.4)	3.0* (2.3, 3.9)
Observations	82,424	81,272	83,649

Notes: The 95 % Confidence Intervals (CI) of the Odds Ratios are in parentheses. * indicates statistical significance, that is, p-value < 0.05. Missing observations from all variables on interest were dropped.

associated with higher odds of current smoking among adolescents (OR 8.7, [95 % CI: 6.9, 11.0]) compared to not having any smoking friends (Table 4).

3.3.2. Other smoked tobacco products

For other smoked tobacco products, girls were less likely to smoke them (OR 0.4, [95 % CI = 0.3, 0.6]) than boys. Adolescents from countries that predominantly have Christians were less likely to smoke (OR 0.8, 95 % CI = [0.6, 1.0]) compared to those that predominantly have Muslims and Hindus. Those who were refused the purchase of cigarettes because of their age were more likely to smoke other smoked products (OR 2.4, [95 % CI = 1.6, 3.5]) than those who were not. Those

who had seen health warnings about the dangers of tobacco were more likely to smoke (OR 1.5, [95 % CI: 1.2, 1.9]) than those who had not (Table 4).

Regarding pro-tobacco variables, adolescents who had been offered free tobacco products were more likely to use other smoked tobacco (OR 3.7 [95 % CI = 2.6, 5.3]) than those who had not. Those who had been exposed to tobacco use at school (OR 1.6, [95 % CI: 1.1, 2.3]) and at home (OR 1.9, [95 % CI: 1.2, 2.8]) were more likely to use other smoked tobacco than those who had not been exposed to smoking in any of these places. Having smoking peers was associated with higher odds of other smoked tobacco use (OR 2.7, [95 % CI: 2.1, 3.4]) compared to not having any smoking friends (Table 4).

3.3.3. Smokeless tobacco use

For smokeless tobacco products, adolescents who reside in East Africa (OR 0.7, [95 % CI: 0.5, 1.0]) were less likely to use smokeless tobacco than those in Central Africa. Residing in other African regions (North, West, and South) was not significantly associated with smokeless tobacco use. Only two anti-tobacco use variables were significantly associated with consuming smokeless tobacco. Those who were restricted from purchasing cigarettes because of their age were more likely (OR 2.4, [95% CI = 1.7, 3.5]) to use smokeless tobacco than those who were not. Those who had seen health warnings about the dangers of tobacco were more likely to consume smokeless tobacco (OR 1.8, [95 % CI: 1.3, 2.4]) than those who had not. Concerning the pro-tobacco variables, those who had been offered free tobacco products were likelier to use smokeless tobacco (OR 4.6, [95 % CI = 3.3, 6.4]) than those who had not. Being exposed to tobacco use at school was associated with higher odds of using smokeless tobacco (OR 4.6, [95 % CI = 3.3, 6.4]) compared to not being exposed (Table 4).

3.3.4. Shisha smoking

Girls were less likely to smoke shisha (OR 0.7, [95 % CI: 0.5, 0.9]) than boys. Adolescents from North Africa (OR 3.2, [95 % CI: 1.7, 5.8]) and West Africa (OR 2.1, [95 % CI = 1.3, 3.5]) were more likely to smoke than those from Central Africa. The association was statistically insignificant between residing in East Africa and shisha smoking. Adolescents from lower-middle-income countries were more likely to use shisha (OR 1.4, [95 % CI = 1.1, 2.0]) than those from low-income countries. Students from countries dominated by Christianity were more likely to consume shisha (OR 1.7, [95 % CI: 1.2, 2.3]) than those dominated by Islam and Hinduism.

Anti-tobacco use factors indicate that adolescents who were refused to buy cigarettes because of their age (aged below 18) (either for themselves or on behalf of other people) were more likely to smoke shisha (OR 5.2, [95 % CI = 3.4, 7.7]) than those who were not. Those who were taught about tobacco dangers at school were less likely to use shisha (OR 0.6, [95 % CI = 0.5, 0.8]) than those who were not. Adolescents who had seen the health warnings about the dangers of tobacco use were more likely to use shisha (OR 1.7, [95 % CI = 1.1, 2.7]) than those who had not (Table 5).

In terms of pro-tobacco use variables, those who had been offered free tobacco products were more likely to consume shisha (OR 2.9, [95 % CI = 2.0, 4.2]) than those who had not. Exposure to tobacco use in public places (OR 1.4, [95 % CI= (1.0, 2.0)]), at school (OR 1.3, [95 % CI: 1.0, 1.7]), and at home (OR 2.0, [95 % CI: 1.5, 2.6]) had a significant positive association with shisha consumption. Having smoking friends/peers was associated with higher odds of current shisha smoking among adolescents (OR 3.7, [95 % CI: 2.5, 5.4]) compared to not having any smoking friends.

3.3.5. Dual use

For the factors associated with the dual use of tobacco products, dual smoked and smokeless, and dual shisha and cigarettes, the same anti-tobacco and pro-tobacco factors that were significantly associated with the use of single products were associated with the dual use (see

Table 5
Factors associated with shisha smoking, dual smoked and smokeless tobacco use, and dual cigarette and shisha smoking.

VARIABLES	Shisha Odds Ratio (95 % CI)	Dual smoked and smokeless tobacco Odds Ratio (95 % CI)	Dual cigarettes and shisha Odds Ratio (95 % CI)
Age	2.3 (0.3, 17.2)	0.0* (0.0, 0.4)	2.2 (0.1, 37.2)
Age Squared	1.0 (0.9, 1.0)	1.1* (1.0, 1.2)	1.0 (0.9, 1.1)
Female (Baseline: Male)	0.7* (0.5, 0.9)	0.5* (0.3, 0.8)	0.4* (0.2, 0.9)
<i>Region (Base: Central Africa)</i>			
East Africa	0.8 (0.4, 1.6)	0.7 (0.4, 1.2)	0.7 (0.2, 2.2)
North Africa	3.2* (1.7, 5.8)	1.0 (0.5, 2.3)	1.3 (0.3, 4.6)
Southern Africa	X	1.1 (0.7, 1.8)	X
West Africa	2.1* (1.2, 3.5)	1.1 (0.5, 2.3)	1.3 (0.4, 4.6)
<i>World Bank Income Group (Base: Low income)</i>			
Lower middle income	1.4* (1.1, 2.0)	1.5* (1.0, 2.1)	0.9 (0.4, 2.1)
Upper middle income	1.1 (0.4, 2.5)	0.9 (0.4, 1.7)	0.6 (0.1, 2.4)
<i>Christian (Base: Muslim & Hindu)</i>			
	1.7* (1.2, 2.3)	1.2 (0.7, 2.1)	1.6 (0.8, 3.2)
Age Restriction	5.1* (3.4, 7.7)	4.1* (2.6, 6.6)	10.6* (6.3, 17.8)
Anti-tobacco use messages	1.1 (0.7, 1.5)	1.2 (0.9, 1.7)	0.8 (0.4, 1.8)
School teaching on tobacco use	0.6* (0.5, 0.8)	0.9 (0.7, 1.3)	0.6* (0.3, 1.0)
Health warnings	1.7* (1.1, 2.7)	1.4* (1.0, 2.0)	1.8* (1.1, 3.0)
Media exposure to smoking	1.2 (0.9, 1.8)	0.7 (0.5, 1.0)	2.2* (1.1, 4.3)
Tobacco adverts	0.9 (0.6, 1.3)	1.1 (0.8, 1.5)	0.9 (0.5, 1.5)
Free tobacco product offer	2.9* (2.0, 4.2)	4.8* (2.7, 8.5)	2.6* (1.5, 4.6)
Public exposure to smoking	1.4* (1.0, 2.0)	1.4* (1.0, 1.9)	3.0* (1.6, 5.7)
School exposure to smoking	1.3* (1.0, 1.7)	2.7* (2.0, 3.7)	2.4* (1.6, 3.6)
Home exposure to smoking	2.0* (1.0, 1.7)	1.7* (1.0, 1.7)	3.3* (1.6, 3.6)

Table 5 (continued)

	Shisha	Dual smoked and smokeless tobacco	Dual cigarettes and shisha
	(1.5, 2.6)	(1.2, 2.3)	(2.0, 5.2)
Peer smoking	3.7* (2.5, 5.4)	4.1* (2.8, 6.1)	14.4* (9.3, 22.1)
Observations	37,527	83,261	35,875

Notes: X indicates that data was unavailable. The 95 % Confidence Intervals (CI) of the Odds Ratios are in parentheses. * indicates statistical significance, that is, p-value < 0.05. The regressions on shisha smoking and its dual use were performed using 16 countries, with no country from Southern Africa. Missing observations from all variables of interest were dropped.

Table 5). Exposure to tobacco smoking on any media platform within a month had a significant positive association only with dual cigarette and shisha smoking (OR 2.2, [95 % CI = 1.1, 4.3]) but a statistically insignificant association with other products. Being taught about tobacco dangers at school had a significant negative association with dual cigarettes and shisha smoking (OR 0.6, [95 % CI: 0.1, 1.0]) but an insignificant association with other products (Table 5).

3.4. Sensitivity analysis results

The prevalence estimates generated with the full dataset were similar to those generated when datasets collected before 2010 were excluded. In the instance that there were differences in the prevalence estimates, the difference was not more than 2 percentage points. Thus, overall, our results are robust to excluding the data that is at least 10 years old (Supplementary Table 7 and Table 8).

4. Discussion

4.1. On the prevalence of tobacco use

We found a prevalence of current use of any tobacco of 14.3 % [95 % CI: 13.5, 15.3]. For cigarettes, the prevalence of current use was 6.4 % [95 % CI: 5.9, 7.0], and this was 5.2 % [95 % CI: 4.4, 6.1] for shisha, 6.7 % [95 % CI: 6.0, 7.4] for other smoked tobacco products, and 6.4 % [95 % CI: 5.9, 6.9] for smokeless tobacco products. The prevalence of tobacco use, including for many of the individual tobacco product types, was highest in Southern Africa and lowest in East Africa, except for shisha, where it was highest in North Africa and lowest in Central Africa. The high prevalence in Southern Africa could be explained by the low tobacco excise taxes in the region and high tobacco industry interference among the youth. This is because tobacco taxes and prices are primary determinants of adolescent smoking and are considered one of the effective tobacco control measures (Filby & van Walbeek, 2022). For instance, the 2022 data from the Tobacconomics Cigarette Tax Scorecard, which measures the performance of cigarette tax structures, shows that on average, out of 5 points, Southern Africa scores (1.90) lower than East Africa (2.08) (Drope et al., 2024). Similarly, available data from a few countries within Southern Africa and East Africa shows that, on average, the tobacco industry interference is higher in Southern Africa than in East Africa, with industry interference scores of 60.25 and 52, respectively (Assunta, 2023). The high prevalence of shisha use in North Africa could be because Shisha may be an emerging or re-emerging product in that region. In fact, across the world, shisha use is increasing among adolescents (Hirpa et al., 2021).

Tobacco use prevalence was higher for boys than for girls, both overall and across all product types. The age-specific prevalence estimates for the different tobacco products decreased from age 11 to 13 years, and then increased from age 13 to 17 years. Our overall prevalence rates for Africa, including the product-, gender- and age-specific prevalence rates, align with those from other studies. For example,

another study reported Africa-specific prevalence rates of 16.2 % for any tobacco use, 10.7 % for other smoked products, and 8.6 % for cigarettes among boys; and 10.3 for any tobacco use, 7.9 % for other smoked products, and 3.7 % for cigarettes among girls (Ma et al., 2021). Other studies have also reported a prevalence of smokeless tobacco use of 4.1 % and 5.1 % in Africa for the 13–15 (James et al., 2022) and 12–16 years (Yang et al., 2022) age groups, respectively. Another analysis of data from 22 African countries reported cigarette smoking and any tobacco use prevalence estimates that are higher than those obtained in the current study (10.9 % and 19.1 %, respectively) (James et al., 2022). The difference in these prevalence rates is because our study includes 31 more African countries that were not included in this study.

In our study, we explored the dual use of tobacco products among adolescents in Africa for the first time and found a prevalence of 3.0 % for the dual use of smoked and smokeless tobacco products. Additionally, we found a prevalence of 1.5 % for the dual use of cigarettes and shisha. The availability and heavy marketing of more tobacco and nicotine products could result in a continued increase in dual tobacco use among the youth (Evans et al., 2015), and result in high exposure to addiction and lower chances of quitting among young people in Africa (Chen et al., 2021). In addition, tobacco use and alcohol consumption in adolescents often co-occur, and both are strongly and positively associated with the use of other psychoactive substances such as marijuana (Myers & Kelly, 2006). Understanding this phenomenon, including the interplay among these products in terms of initiation and continued use, will be important for effective tobacco control efforts. For example, there might be a need for the development of tobacco cessation programmes that account for the dual use of tobacco products and their co-consumption with other psychoactive substances.

4.2. On the factors associated with tobacco use

Our study confirms findings from other African studies that girls are currently less likely to use smoked tobacco products than boys (James et al., 2022; Oyewole et al., 2018). However, this might change in the future because of changing social and cultural norms. There is also an emergence of tobacco products that are viewed as more acceptable for women to use. For example, although we did not find this in our current study, some studies in Nigeria have shown that among the youth, females are as likely to be current shisha smokers as males (Otakhoigbogie et al., 2022) and that shisha use among females is perceived as more accepted by society, and more comfortable to females when compared to cigarette smoking (Mdege et al., 2024). In addition, with the plateauing or decline of tobacco use in many high-income countries, there are indications that the tobacco industry is now aggressively marketing to women and girls in low and middle-income countries (Amos & Haglund, 2000). Our analysis showed that for some countries, the gap between boys and girls in terms of overall tobacco use prevalence has closed: for example, Zimbabwe with a prevalence of 22.6 % [95 % CI: 17.4, 28.8] among boys and 16.9 % [95 % CI: 12.9, 21.8] among girls.

In our study, those who had been denied the purchase of cigarettes because of their age were more likely to be current tobacco users than those who had not, which contradicts evidence from another study focusing on South Asia, where age restriction was found to be protective of adolescents from smoking (Mishu et al., 2021). Second, the results for the health warnings also suggested that seeing health warnings about tobacco dangers is a positive predictor of tobacco use across all products. For policy, this implies tobacco control policymakers should consider larger textual and graphic/pictorial health warnings, as they are considered one of the effective tools to reduce tobacco use (White et al., 2008). Currently, many African have “text only” health warnings. Sixteen countries also require pictorial health warnings on tobacco products packs, but they have not necessarily implemented them (Campaign for Tobacco-Free Kids, 2021).

The unexpected results of health warnings and age restrictions may be because of how the GYTS questionnaire was designed. It is plausible

that this question was asked even to those who did not smoke. Specifically, in the case of health warnings, adolescents may be confusing the text health warnings with tobacco advertisements. That is, the fact that they have access to the products may make them misinterpret them as advertisements. In other words, for this group, it may not make sense they are sold these products and at the same time given health warnings against them. Together, the surprising results of the relationship between health warnings and age restriction from purchasing tobacco may be because adolescents are sometimes naturally curious and risk-takers (Duell & Steinberg, 2021). Thus, when advised against something, they do the opposite. That is, when they are restricted from purchasing tobacco because of their age, they seem to be more curious about purchasing and using tobacco. Similarly, when given health warnings against tobacco use, they do obey them and instead seem to show more interest in tobacco use.

School teaching about the dangers of tobacco use had a negative association with tobacco use, especially shisha smoking and dual shisha and cigarette smoking. This aligns with evidence from countries like Lesotho (Pokothoane, 2021), and cross-country findings from 73 countries (Ma et al., 2022), 4 South Asian countries (Mishu et al., 2021), and 138 countries (Yang et al., 2022). In fact, our results about school pro- and anti-tobacco use factors suggest that the school environment plays a role in shaping the odds of cigarette smoking and dual shisha and cigarette smoking.

In contrast, exposure to anyone smoking at home and peer smoking have strong positive associations with tobacco use, irrespective of tobacco product type. Public exposure to smoking was also a statistically highly significant positive predictor of specific tobacco products: cigarette smoking, shisha smoking, and the dual use of smoked and smokeless tobacco. This supports existing evidence on the importance of the home and social environment in shaping adolescents' tobacco use behaviours (James et al., 2022; Ma et al., 2022; Oyewole et al., 2018; Pokothoane, 2021). As with other studies (Yang et al., 2022), our results also show that tobacco industry interference through the offer of free tobacco products can significantly increase the likelihood of use across all tobacco product types. This is critical for African tobacco control policy to fight tobacco industry influence since the industry targets African adolescents to retain them as their lifetime consumers (Drope et al., 2018).

4.3. On policy implications and limitations

In Africa, the age of initiation among young people ranges from as low as 7 years to about 16 years (Chido-Amajuoyi et al., 2021; Mdege et al., 2024). African countries and regions need to strengthen their tobacco control responses to curb tobacco initiation and use among young people and to reduce tobacco use prevalence. Evidence of the prevalence of both the single and dual use of tobacco products suggests that effective policy tools should be implemented. Overall, our results on the factors of tobacco use show that school teaching about tobacco dangers is negatively associated with tobacco use, while health warnings, free offer of tobacco products and peer influence are positive correlates. This suggests a policy mechanism that can complement the school teaching on tobacco use and potentially reduce the effectiveness of the positive factors is needed. Thus, we recommend implementing pictorial health warnings on the packs of cigarettes and other products. The pictorial warnings would educate the entire peer group about the dangers of tobacco, and possibly make them resist the temptation of using any products when offered freely.

It is also essential to have comprehensive surveillance systems that monitor the use of the whole range of tobacco and nicotine products overtime. Data has to cover all adolescents (10–19 year olds) not only those that are 13–15 years old. Data that includes adolescents who are out of school is also needed: some studies suggest that they might be more vulnerable to tobacco use than those who are in school (Desai et al., 2019; Otakhoigbogie et al., 2022).

This study is not without limitations. The datasets used for some countries are old and are from different years. Further, the GYTS questionnaire is self-reported, such that adolescents may underreport tobacco use because of social desirability or cultural issues. The results might not be generalizable to all the adolescents at the country or regional levels, particularly in the context of high out-of-school rates; and might not reflect the current use situation accurately. In our Logistic regressions, because of data limitations, we did not account for significant factors associated with tobacco use, such as the price of tobacco products and adolescents' income (pocket money). We also did not account for taxes on different products and the tobacco industry interference indices due to a lack of data across the included countries. Nonetheless, our study provides comprehensive estimates, using currently available data, for different types of tobacco products in Africa, and contributes to the evidence needed for tobacco control policy decisions in Africa.

5. Conclusion

The prevalence of tobacco use among African adolescents is high and similar for different tobacco products. Some demographic factors only matter for using specific products, while anti-tobacco factors, although counterintuitive, matter for tobacco use irrespective of product type. Peer influence, school and home environment exposure to smoking, and health warnings about tobacco use are positive factors of tobacco use, irrespective of product type among adolescents. Policymakers should develop pictorial and graphical health warnings about tobacco dangers to discourage tobacco use and educate the entire adolescent peer group.

Author statement

All authors acknowledge that the material presented in this manuscript has not been previously published, except in abstract form, nor is it simultaneously under consideration by any other journal.

Funding

This work was supported by the Bill & Melinda Gates Foundation (INV-048743).

CRediT authorship contribution statement

Retselisitsoe Pokothoane: Writing – review & editing, Writing – original draft, Visualization, Software, Methodology, Investigation, Formal analysis, Conceptualization. **Terefe Gelibo Agerfa:** Writing – review & editing, Visualization, Validation, Software, Formal analysis, Conceptualization. **Christus Cito Miderho:** Writing – review & editing, Conceptualization. **Noreen Dadirai Mdege:** Writing – review & editing, Supervision, Methodology, Formal analysis, Conceptualization.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: The Bill & Melinda Gates Foundation had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results. The findings and conclusions contained within are those of the authors and do not necessarily reflect positions or policies of the Bill & Melinda Gates Foundation. The authors declare that they have no conflicts of interest related to publication of this manuscript.

Acknowledgments

This study was conducted as part of the Data on Youth and Tobacco in Africa (DaYTA) program. The DaYTA program seeks to empower

stakeholders to make timely, data-driven decisions by using evidence to inform policy and advance tobacco control efforts in sub-Saharan Africa. The program achieves this by addressing data gaps related to tobacco use among 10- to 17-year-olds in Kenya, Nigeria, and the Democratic Republic of Congo. The DaYTA program is led by Development Gateway: an IREX Venture (DG), a global non-profit organization that specializes in data for development. We thank Josiane Djiofack Tsague, the associate editor, and three anonymous reviewers for their helpful comments.

Data availability

All the data are publicly available on the WHO Microdata repository and can be accessed [here](#).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.abrep.2024.100581>.

References

- African-Union. (2023). *Member states*. African Union.
- Amos, A., & Haglund, M. (2000). From social taboo to "torch of freedom": The marketing of cigarettes to women. *Tobacco Control*, 9(1), 3–8. <https://doi.org/10.1136/tc.9.1.3>
- Assunta, M. (2023). *Global Tobacco Industry Interference Index 2023*. Global Center for Good Governance in Tobacco Control (GGTC). <https://globaltobaccoindex.org/report-summary>.
- Campaign for Tobacco-Free Kids. (2021). *Countries and Jurisdictions with Pictorial Health Warning Labels, by Size*. Campaign for Tobacco-Free Kids. https://assets.tobaccofreekids.org/global/pdfs/en/WL_country_size.pdf.
- Chen, D.-T.-H., Girvalaki, C., Mechili, E. A., Millett, C., & Filippidis, F. T. (2021). Global patterns and prevalence of dual and poly-tobacco use: A systematic review. *Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco*, 23(11), 1816–1820. <https://doi.org/10.1093/ntr/ntab084>
- Chido-Amajuoyi, O. G., Fueta, P., & Mantey, D. (2021). Age at smoking initiation and prevalence of cigarette use among youths in Sub-Saharan Africa, 2014–2017. *JAMA Network Open*, 4(5), e218060.
- de la Torre-Luque, A., Ozeylem, F., & Essau, C. A. (2021). Prevalence of addictive behaviours among adolescents from 73 low-and middle-income countries. *Addictive Behaviors Reports*, 14, Article 100387. <https://doi.org/10.1016/j.abrep.2021.100387>
- Desai, R., Mercken, L. A. G., Ruiter, R. A. C., Schepers, J., & Reddy, P. S. (2019). Cigarette smoking and reasons for leaving school among school dropouts in South Africa. *BMC Public Health*, 19(1), 130. <https://doi.org/10.1186/s12889-019-6454-5>
- Drope, J., Oo, S., Chaloupka, F., Dorokhina, M., Guerrero-López, C., Rodriguez-Iglesias, G., Mugosa, A., Mirza, M., Bontu, A., & Chaloupka, F. (2024). *Tobaccconomics Cigarette Tax Scorecard* (3rd Edition). Baltimore, MD: Bloomberg School of Public Health. <https://tobaccconomics.org/files/research/919/tobacco-scorecard-report-3rd-ed-eng-v5.0.pdf>.
- Drope, J., Schluger, N., Cahn, Z., Drope, J., Hamill, S., Islami, F., Liber, A., Nargis, N., & Stoklosa, M. (2018). *The Tobacco Atlas*. Atlanta: American Cancer Society and Vital Strategies. https://pngcancerfoundation.org/wp-content/uploads/2021/06/TobaccoAtlas_6thEdition_LoRes.pdf.
- Duell, N., & Steinberg, L. (2021). Adolescents take positive risks, too. *Developmental Review*, 62. <https://doi.org/10.1016/j.dr.2021.100984>
- Evans, W. D., Horn, K. A., & Gray, T. (2015). Systematic review to inform dual tobacco use prevention. *Pediatric Clinics of North America*, 62(5), 1159–1172. <https://doi.org/10.1016/j.pcl.2015.05.003>
- Filby, S., & van Walbeek, C. (2022). Cigarette prices and smoking among youth in 16 African countries: Evidence from the global youth Tobacco Survey. *Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco*, 24(8), 1218–1227. <https://doi.org/10.1093/ntr/ntac017>
- Flay, B. R., Petraitis, J., & Hu, F. B. (1995). The theory of triadic influence: Preliminary evidence related to alcohol and tobacco use. In *Alcohol and tobacco: From basic science to clinical practice*. NIAAA, Bethesda, MD. https://www.researchgate.net/profile/Brian-Flay/publication/224942198/The-theory-of-triadic-influence-Preliminary_evidence_related_to_alcohol_and_tobacco_use/links/55a2ca808a9ca1e64fe7f/The-theory-of-triadic-influence-Preliminary-evidence-related-to-alcohol-and-tobacco-use.pdf.
- Hamadeh, N., Van Rompaey, C., & Metreau, E. (2023). *June 30*. World Bank Group country classifications by income level for FY24. World Bank Blogs.
- Hirpa, S., Fogarty, A., Addissie, A., Bauld, L., Frese, T., Unverzagt, S., Kantelhardt, E. J., Getachew, S., & Deressa, W. (2021). An Emerging Problem of Shisha Smoking among High School Students in Ethiopia. *International Journal of Environmental Research and Public Health*, 18(13). <https://doi.org/10.3390/ijerph18137023>
- James, P. B., Bah, A. J., Kabba, J. A., Kassim, S. A., & Dalinjong, P. A. (2022). Prevalence and correlates of current tobacco use and non-user susceptibility to using tobacco products among school-going adolescents in 22 African countries: A secondary analysis of the 2013–2018 global youth tobacco surveys. *Archives of Public Health* =

- Archives Belges de Sante Publique, 80(1), 121. <https://doi.org/10.1186/s13690-022-00881-8>
- Ma, C., Xi, B., Li, Z., Wu, H., Zhao, M., Liang, Y., & Bovet, P. (2021). Prevalence and trends in tobacco use among adolescents aged 13–15 years in 143 countries, 1999–2018: Findings from the Global Youth Tobacco Surveys. *The Lancet Child & Adolescent Health*, 5(4), 245–255. [https://doi.org/10.1016/S2352-4642\(20\)30390-4](https://doi.org/10.1016/S2352-4642(20)30390-4)
- Ma, C., Yang, H., Zhao, M., Magnussen, C. G., & Xi, B. (2022). Prevalence of waterpipe smoking and its associated factors among adolescents aged 12–16 years in 73 countries/territories. *Frontiers in Public Health*, 10, 1052519. <https://doi.org/10.3389/fpubh.2022.1052519>
- Mayhew, K. P., Flay, B. R., & Mott, J. A. (2000). Stages in the development of adolescent smoking. *Drug and Alcohol Dependence*, 59(Suppl 1), S61–S81. [https://doi.org/10.1016/S0376-8716\(99\)00165-9](https://doi.org/10.1016/S0376-8716(99)00165-9)
- Mdege, N. D., Ekpo, R., Ogolla, S., Ali, S. J., Camara, A., & Mugweni, E. (2024). Reasons for shisha smoking: Findings from a mixed methods study among adult shisha smokers in Nigeria. *PLoS Global Public Health*, 4(2), e0002853.
- Mishu, M. P., Siddiqui, F., Shukla, R., Kanaan, M., Dogar, O., & Siddiqi, K. (2021). Predictors of Cigarette Smoking, Smokeless Tobacco Consumption, and Use of both forms in Adolescents in South Asia: A Secondary Analysis of the Global Youth Tobacco Surveys. *Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco*, 23(6), 956–965. <https://doi.org/10.1093/ntr/ntaa202>
- Myers, M. G., & Kelly, J. F. (2006). Cigarette smoking among adolescents with alcohol and other drug use problems. *Alcohol Research & Health: The Journal of the National Institute on Alcohol Abuse and Alcoholism*, 29(3). <https://pubmed.ncbi.nlm.nih.gov/17373413/>.
- Nonnemaker, J. M., & Farrelly, M. C. (2011). Smoking initiation among youth: The role of cigarette excise taxes and prices by race/ethnicity and gender. *Journal of Health Economics*, 30(3), 560–567. <https://doi.org/10.1016/j.jhealeco.2011.03.002>
- Otakhoigbogbe, U., Osagbemi, B. B., & Akaji, E. A. (2022). Knowledge and sociodemographic predictors of shisha smoking among students in a Nigerian university. *Nigerian Journal of Clinical Practice*, 25(6). https://doi.org/10.4103/njcp.njcp_1424_21
- Oyewole, B. K., Animasahun, V. J., & Chapman, H. J. (2018). Tobacco use in Nigerian youth: A systematic review. *PloS One*, 13(5), e0196362.
- Pew Research Center Religion & Public Life. (2015). *The Future of World Religions: Population Growth : Projections, 2010-2050*. <https://www.pewforum.org/2015/04/02/religious-projections-2010-2050/>.
- Pokothoane, R. (2021). *Econometric Analysis of Tobacco Use: Evidence from Lesotho Global Youth Tobacco Surveys* [Master's Thesis, University of Cape Town]. <https://open.uct.ac.za/server/api/core/bitstreams/b0eb0e37-b3b3-4e1a-ac7d-bc6bca92e89e/content>.
- Sawyer, S. M., Azzopardi, P. S., Wickremarathne, D., & Patton, G. C. (2018). The age of adolescence. *The Lancet. Child & Adolescent Health*, 2(3), 223–228. [https://doi.org/10.1016/S2352-4642\(18\)30022-1](https://doi.org/10.1016/S2352-4642(18)30022-1)
- StataCorp. (2021). *Stata Statistical Software Release 17* (Version 17).
- White, V., Webster, B., & Wakefield, M. (2008). Do graphic health warning labels have an impact on adolescents' smoking-related beliefs and behaviours? *Addiction*, 103(9), 1562–1571. <https://doi.org/10.1111/j.1360-0443.2008.02294.x>
- World Health Organization. (2016). *Fact Sheet: Tobacco Use in Adolescence*. *World Health Organization Region Office for Europe*. https://who-sandbox.squiz.cloud/_data/assets/pdf_file/0016/303532/HBSC-No.7_factsheet_Tobacco.pdf.
- World Health Organization. (2020). *Heated Tobacco Products: A Brief*. *World Health Organization Regional Office for Europe*. <https://iris.who.int/bitstream/handle/10665/350470/WHO-EURO-2020-4571-44334-64934-eng.pdf?sequence=3>.
- World Health Organization. (2023a). *Noncommunicable disease surveillance, monitoring and reporting: Global Youth Tobacco Survey*. *World Health Organization*. <https://www.who.int/teams/noncommunicable-diseases/surveillance/systems-tools/global-youth-tobacco-survey>.
- World Health Organization. (2023b). *WHO NCD Microdata Repository*. *World Health Organization*. <https://extranet.who.int/ncdsmicrodata/index.php/home>.
- World Health Organization. (2023c). *WHO report on the global tobacco epidemic, 2023: Protect people from tobacco smoke*. *World Health Organization*. <https://www.who.int/publications/i/item/9789240077164>.
- World Health Organization. (2024). *Hooking the next generation: How the tobacco industry captures young customers*. *World Health Organization*. <https://www.who.int/publications/i/item/9789240094642>.
- Yang, H., Ma, C., Zhao, M., Magnussen, C. G., & Xi, B. (2022). Prevalence and trend of smokeless tobacco use and its associated factors among adolescents aged 12–16 years in 138 countries/territories, 1999–2019. *BMC Medicine*, 20(1), 460. <https://doi.org/10.1186/s12916-022-02662-0>