



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/222348/>

Version: Published Version

Article:

Schmoeckel, M., Thielmann, M., Vitanova, K. et al. (2025) Direct-acting oral anticoagulant removal by intraoperative hemoadsorption in CABG and/or single valve surgery: interim analysis of the International Safe and Timely Antithrombotic Removal (STAR) registry. Journal of Cardiothoracic Surgery, 20. 74. ISSN: 1749-8090

<https://doi.org/10.1186/s13019-024-03326-1>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

RESEARCH

Open Access



Direct-acting oral anticoagulant removal by intraoperative hemoadsorption in CABG and/or single valve surgery: interim analysis of the International Safe and Timely Antithrombotic Removal (STAR) registry

Michael Schmoeckel^{1,15*}, Matthias Thielmann², Keti Vitanova³, Thomas Eberle⁴, Nandor Marczin⁵, Kambiz Hassan⁶, Andreas Liebold⁷, Sandra Lindstedt⁸, Georg Mächler⁹, Marijana Matejic-Spasic¹⁰, Daniel Wendt^{10,11}, Efthymios N. Deliargyris¹² and Robert F. Storey^{13,14}

Abstract

Objective Patients on direct-acting oral anticoagulants (DOACs) are at high risk of perioperative bleeding complications. Intraoperative hemoadsorption is a novel strategy to reduce perioperative bleeding in patients on DOACs undergoing non-deferable cardiac surgery. The international STAR-registry reports real-world clinical outcomes associated with this application.

Methods The hemoadsorption device was incorporated into the cardiopulmonary bypass (CPB) circuit and active for the duration of the pump run. Patients on DOACs undergoing CABG and/or single valve surgery before completing the recommended washout were included. Outcome measurements included bleeding events according to standardized definitions and 24-hour chest-tube-drainage (CTD).

Results A total of 62 patients were included from 7 institutions in Austria, Germany, Sweden, and the UK (mean age 69.9 ± 7.5 years, 71% male). Approximately half were on apixaban and the other half was split between rivaroxaban and edoxaban with 21% of patients also on aspirin. Surgery occurred at a median time of 28.9 h since the last DOAC dose with single valve surgery accounting for 2/3 of cases. Mean CPB duration was 118.6 ± 46.4 min. Severe bleeding (UDP ≥ 3) occurred in 4.8%, and BARC-4 bleeding occurred in 3.2% of the patients. Only one patient (1.6%) required reoperation for bleeding control. The mean 24-hour CTD was 771.3 ± 482.79 mL. No device-related adverse events were reported.

Presented at the 38th EACTS Annual Meeting in Lisbon, Portugal, Oct 9–12, 2024.

*Correspondence:

Michael Schmoeckel
michael.schmoeckel@med.uni-muenchen.de

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Conclusions This interim report of the ongoing STAR-registry shows that, in patients on DOAC undergoing non-deferable CABG and/or single valve surgery, the use of intraoperative hemoadsorption is associated with low rates of severe perioperative bleeding complications. Further prospective studies in larger cohorts are needed to validate the efficacy of this method.

Clinical registration number ClinicalTrials.gov identifier: NCT05077124.

Keywords Hemoadsorption, Antithrombotic removal, Cardiac surgery, CytoSorb, DOAC

Introduction

Antithrombotic drugs are cornerstone treatments for patients with cardiovascular disease [1]. Direct-acting oral anticoagulants (DOACs) are the first-line therapy for millions of patients around the world for lifelong stroke prevention in patients with atrial fibrillation (AF) and for prevention and management of deep vein thrombosis [2–4]. In this context, DOACs have been shown to be safe and effective, but also easier to use since they do not require strict and frequent laboratory monitoring or dose adjustments compared to warfarin. The main safety risk associated with DOACs is increased bleeding, which can be spontaneous or iatrogenic, such as when these patients require urgent or emergency interventions, including cardiac surgery [1].

Current guidelines recommend that DOACs should be discontinued at least 48 h before surgery to minimize the risk of bleeding [5]. However, clinical scenarios often preclude this waiting period, necessitating alternative strategies to reduce the risk of bleeding in cases of non-deferable surgery. Removal of DOACs from the circulation by intraoperative polymer bead hemoadsorption has emerged as a novel strategy to reduce perioperative bleeding in patients on antithrombotic drugs undergoing urgent or emergency on-pump cardiac surgery [6, 7]. The international Safe and Timely Antithrombotic Removal (STAR) registry is designed to collect real-world clinical outcomes associated with this application (NCT05077124).

The current interim analysis of the international STAR Registry reports perioperative bleeding rates in patients on DOACs undergoing non-deferable CABG and/or single valve surgery with the use of intraoperative hemoadsorption.

Methods

The international Safe and Timely Antithrombotic Removal (STAR) registry is designed to collect high-fidelity data on patients who underwent antithrombotic removal (ATR) with CytoSorb® as part of their routine care. Participating centers can include both prospective and retrospective patients depending on the date of ethics approval: patients who received hemoadsorption for ATR as part of their clinical routine prior to the registry's protocol ethics approval are entered as retrospective

patients, whereas patients enrolled after ethics approval are entered as prospective. Collected clinical and resource utilization data are entered in an electronic case report form (Medpace Inc., Ohio, Cincinnati, USA). Safety is assessed by collection of definite, probable, or possible device-related investigator-reported adverse events according to the latest ISO 14155:2020 classification. Data collection is done up to 30 days post ATR. The sponsor and funding source of the registry is CytoSorbents Inc., Princeton, NJ, USA. The ongoing registry is designed to enroll up to 1,000 real-world patients undergoing cardiac surgery with intraoperative drug removal by hemoadsorption.

Ethical statement

This registry complies with the Declaration of Helsinki. National central or local approvals of respective Ethics committees were granted for the STAR registry according to local regulations (complete list of all ethics available within the Supplement). Written consent for data collection was obtained before or after surgery from prospective patients and was waived for retrospective patients.

Inclusion criteria

CytoSorb® utilization for antithrombotic removal and informed consent for prospective registry participation. The CE-marked CytoSorb adsorber was used as part of routine clinical care.

Exclusion criteria

Use of CytoSorb® for a purpose other than antithrombotic removal.

Patients

This interim analysis of the STAR Registry included all enrolled patients undergoing ATR for DOAC removal with intraoperative hemoadsorption during CABG and/or single valve surgery.

Hemoadsorption therapy

Antithrombotic removal via hemoadsorption therapy was performed with the CytoSorb® adsorber (CytoSorbents Inc., Princeton, NJ, USA). The device is based on a proprietary extracorporeal blood purification and is

CE-mark approved to remove ticagrelor and rivaroxaban during on-pump cardiac surgery. The cartridge is filled with highly biocompatible, porous polymer beads covered with a divinylbenzene coating and can be easily integrated into various extracorporeal circuits, such as e.g., continuous renal replacement therapy, extracorporeal membrane oxygenation (ECMO), or cardio-pulmonary bypass (CPB), as shown in Fig. 1. Each polymer bead is between 300 μm and 800 μm in size and has pores and channels, giving it a large ($>40,000 \text{ m}^2$) effective surface area for binding hydrophobic small and medium-sized molecules up to 60 kDa of molecular weight [8]. DOACs have a molecular weight ranging between ~ 400 –600 Daltons [9] and can therefore easily pass into the pores of the beads and readily adsorb onto the internal polymer surface through a combination of non-polar interactions, hydrogen bonding, and van der Waals forces.

Outcome measures

Perioperative bleeding complications were evaluated according to two standardized bleeding definitions: (a) *Bleeding Academic Research Consortium* (BARC), and (b) *Universal Definition of Perioperative Bleeding* (UDPB). Additional outcomes included 24-hour chest tube drainage (CTD), blood product transfusions, re-operation for bleeding, and in-hospital mortality. Safety of the device was assessed by investigator-reported adverse device events, including severity and related classifications (**Supplement**).

Results

For the present analysis, a total of 62 patients on DOACs were included from 7 institutions in Austria, Germany, Sweden, and UK. All patients underwent on-pump isolated CABG and/or single valve surgery and received intraoperative hemoadsorption as part of their clinical routine.

Baseline characteristics are presented in Table 1. Patients on all 4 available DOACs (rivaroxaban, apixaban, edoxaban and dabigatran) were included; however, almost half of the patients were on apixaban, approximately one-quarter each were on rivaroxaban and edoxaban, and only few were on dabigatran (Fig. 2). The washout period from the last DOAC dose (median 28.9 h [Q1: 24.0, Q3: 50.9]) was less than 48 h in all patients (Table 1). Patients with either longer or shorter wash-out periods according to the overall cohort's median and interquartile range for washout did not differ in terms of demographics or bleeding outcomes according to the UDPB ($p=0.562$) or BARC-4 ($p=1.000$) definition (See Fig. 3).

Procedural and postoperative outcomes are summarized in Table 2. A total of 23 patients underwent isolated CABG and 39 underwent single valve surgery, of whom 16 received concomitant CABG surgery. The duration of CPB (a direct measure also of hemoadsorption device exposure) and aortic cross clamp duration were 118.6 ± 46.4 and 80.9 ± 32.8 min, respectively (See Table 3).





Fig. 1 Hemoadsorption device (CytoSorb®) incorporated in the cardiopulmonary (CPB) circuit. The device remained active for the duration of the CPB-run

Table 1 Demographics

Variable	Isolated CABG or single valve surgery n = 62
Age, years	69.9 ± 7.5
Gender, male	44 (71.0%)
Weight, kg	84.8 ± 18.9
Acetylsalicylic acid use before surgery	13 (21.0)
Acute coronary syndrome	17 (27.4%)
Atrial fibrillation	56.7% (34/60)
History of deep vein thrombosis	7.1% (4/56)
Prior Stroke	6.7% (4/60)
Current smoker	39.3% (11/28)
Time since last dose of DOAC, hours	28.9 (24.0, 50.9)
NYHA functional class III/IV	61.3% (19/31)
Hypertension	52 (85.2%)
Diabetes	20 (32.3%)
Hyperlipidemia	42.4% (25/59)
Prior myocardial infarction	11.5% (7/61)
Peripheral artery disease	6.8% (4/59)
Renal dysfunction (creatinine > 1.3 mg/dL / failure (dialysis))	31.1% (19/61)
Hepatic Impairment	6.8% (4/59)
Prior sternotomy	6.6% (4/61)
EuroSCORE-II, %	6.9 ± 12.8

Data are presented as number (%), or % (n/N), mean ± SD or median (Q1, Q3)

NYHA - New York Heart Association, EuroSCORE - European System for Cardiac Operative Risk Evaluation

**Fig. 2** Type of DOAC drug

Serious postoperative bleeding complications were infrequent and included one re-operation for bleeding control, one case of CTD > 2 L and one patient receiving 8 units of fresh-frozen plasma. These events resulted in rates of 4.8% and 3.2% according to the UDPB and

BARC-4 definitions, respectively. Mean 24-hour CTD was 771.3 ± 482.79 mL with 10 patients exceeding 1 L (16.1%) and 1 patient exceeding 2 L (1.6%).

Packed red blood cells (pRBC) were not transfused during the first postoperative day in 75.8% of the patients. Platelets and fresh frozen plasma were administered in 12.9% and 9.7% of the patients, respectively.

Mortality at 30 days in the overall cohort was 6.5% (4/62) and all deaths were unrelated to bleeding. There were no device-related adverse events reported by investigators.

Discussion

This interim analysis from the ongoing international STAR registry reports rates of serious perioperative bleeding complications in patients on DOACs undergoing non-deferable on-pump CABG and/or single-valve surgery with intraoperative hemoadsorption. The main observation of this report is that the intraoperative use of a hemoadsorption device resulted in low rates of serious perioperative bleeding complications in patients on DOACs undergoing isolated CABG and/or single valve surgery before completing the recommended washout period. In addition, participating investigators reported that incorporation of intra-operative hemoadsorption in their routine surgical protocol was simple and safe since there were no device-related adverse events observed.

Tens of millions of patients are on DOACs worldwide and a significant number of them will require urgent surgical intervention during the course of their therapy [1, 2, 10]. In such a scenario, high perioperative bleeding rates of up to 23% have been reported [2]. Accordingly, current guidelines recommend that patients discontinue DOACs at least 48 h prior to surgery whenever possible with even longer periods required for patients with renal insufficiency [1, 5, 11].

The CytoSorb hemoadsorption device (CytoSorbents, Inc., Princeton, NJ, USA) has been shown to effectively remove all DOACs from blood in the in-vitro setting [7]. CytoSorb is currently CE mark approved for clinical use for removal of cytokines, myoglobin, bilirubin, ticagrelor and rivaroxaban and can be easily integrated into any extracorporeal hemoperfusion circuit such as cardiopulmonary bypass, extracorporeal membrane oxygenation, continuous renal replacement therapy/continuous venovenous hemofiltration or simple hemoperfusion [12]. CytoSorb is meanwhile used in various clinical settings in cardiac surgery [13].

The international STAR Registry is designed to capture high-fidelity data with the use of CytoSorb for anti-thrombotic removal in everyday clinical practice [6]. The current analysis focused on bleeding outcomes after CABG and/or single valve surgery in order to minimize any impact on postoperative bleeding by more complex




Fig. 3 Perioperative outcomes according to UDPB and BARC-4, the surgical revision rate and 30-day mortality

Table 2 Procedural & postoperative outcomes

Outcome parameters	Isolated CABG or single valve surgery n=62
Procedural	
Operation duration, hours	3.8 ± 1.2
CPB time (device exposure), min.	118.6 ± 46.4
ACC time, min.	80.9 ± 32.8
Isolated CABG surgery	23 (37.1)
Single valve surgery	39 (62.9)
Single valve surgery and concomitant CABG surgery	16/39
Cerebral perfusion performed	0 (0)
Postoperative	
24-hour CTD, mL	771.3 ± 482.79
0-600mL	44.6%
601-800mL	19.6%
801-1,000mL	17.9%
1,001–2,000mL	16.1%
>2,000mL	1.6%
UDPB ≥ 3, %	4.8
BARC-4 within 48 h	3.2
Re-operation for bleeding at any time, %	1.6
Re-operation for bleeding by POD1, %	1.6
ICU-stay, hours	70.8 (26.2, 163.0)
Hospital stay, days	13 (9, 17)
30-day mortality, %	6.5

Data are presented as number (%), mean ± SD or median (Q1, Q3)
CPB - cardiopulmonary bypass, ACC - aortic cross clamp, CTD - chest tube drainage, BARC - Bleeding Academic Research Consortium, UDPB - Universal Definition of Perioperative Bleeding, POD - postoperative day

Table 3 Blood product consumption (until postoperative day 1)

Variable	Isolated CABG or single valve surgery n= 62
Patients receiving pRBC transfusion	15 (24.2)
pRBC units	2.5 ± 1.6
Patients receiving platelet transfusion	8 (12.9)
platelet units	2.3 ± 1.6
Patients receiving FFP transfusion	6 (9.7)
FFP units	5.2 ± 2.9

Data are presented as number (%) or mean ± SD. pRBC - packed red blood cells, FFP - fresh frozen plasma

surgery (i.e. aortic surgery or multiple valve procedures). In a recently published first analysis of the STAR registry, all types of surgery were analyzed, including complex aortic surgery, heart transplantation or combination procedures, resulting in higher UDPB and BARC-4 bleeding rates compared with the current analysis, which was intended to evaluate the impact of the device on antithrombotic related bleeding without any potential confounding due to the additional bleeding risk associated with more complex and higher risk procedures [6]. Consistent with global prescribing trends, patients in the STAR Registry were more frequently on apixaban followed by rivaroxaban and edoxaban. The overall median time from last dose to surgery in our cohort was 28.9 h, which is substantially shorter than the guideline recommendation of at least 48 h [5]. However, surgeons routinely report that their standard washout period is longer as many of their elderly patients have concomitant renal or hepatic disease, resulting in slower drug metabolism and washout. It is worth noting that in a previous analysis

of cardiac surgery patients on DOACs, a longer withdrawal period, particularly in patients with reduced renal function, was essential for safe on-pump cardiac surgery [14]. The authors concluded that, despite official recommendations, patients should not be considered for elective cardiac surgery within 10 days of stopping DOAC treatment [14]. The current report was representative of an all-comer population that included approximately 1/3 of patients with some degree of renal insufficiency and to a certain extent, liver disease. Of note, for dabigatran compared to other DOACs, renal excretion is the dominant elimination pathway, but all DOACs are eliminated to a certain extent by the renal pathway as mentioned by Chen et al. All DOAC therapies are eliminated by the kidneys to varying degrees, and changes in renal clearance must be taken into account when dosing these agents. Dabigatran has the highest fraction eliminated renally at 80%, of its clearance pathway, followed by edoxaban, rivaroxaban, apixaban, and betrixaban: 50%, 35%, 27%, and 11%, respectively [15].

A study published by Hassan et al. analyzed 25 consecutive patients on apixaban who underwent cardiac surgery, 13 with CytoSorb and 12 without. In their analysis, the preoperative washout interval was 0.6 ± 1.2 days in the CytoSorb vs. 1.3 ± 0.9 days in the control group ($p < 0.001$). Despite the shorter washout interval, the authors concluded that no BARC-4 bleeding complications did occur in the CytoSorb group and chest tube drainage was significantly lower [10].

The bleeding rates observed in the current analysis (UDPB ≥ 3 : 4.8% and BARC-4: 3.2%) are similar to rates observed among patients not on DOACs and compare favorably to rates of bleeding complications reaching 23% that have been reported in the literature for patients on DOACs [2]. In addition, we observed a low surgical revision rate of 1.6% compared to a recent analysis by Hassan et al., who reported a surgical revision rate of 11.4% (4/35) in a cohort of DOAC-treated patients undergoing cardiac surgery prior to recommended washout compared to less than 2% in the control group [16]. In the current literature, incidence of surgical revision in DOAC patients varies between 6.28% and 11.8% [14, 17].

By adopting checklists for hemostasis and a dedicated algorithm for surgical preoperative planning in elective cases, bleeding rates can be further reduced [18]. Similarly, the same authors reported an increased red blood cell (RBC) transfusion rate of 39.5% in DOAC patients compared to less than 20% in controls [16]. Therefore, the results of the present analysis, with less than 2% of surgical revisions and 24.2% of patients receiving RBC transfusion, are almost comparable to control patients not on DOACs, suggesting a significantly lower incidence of major postoperative bleeding complications with the use of intraoperative hemoadsorption.

To date, only a few studies with smaller single-center cohorts have been published on this topic. Only one study reported bleeding outcomes in isolated CABG patients with prior rivaroxaban treatment. The authors reported no surgical revisions in the CytoSorb treated group ($n = 7$) compared to 2 surgical revisions in the control group ($n = 5$). All patients treated with rivaroxaban and the use of adsorption showed a significant reduction in drug levels in the anti-factor Xa assay [16]. In another observational case series from 2 centers, the use of intraoperative hemoadsorption in patients on apixaban undergoing different types of cardiac surgery (there were no BARC-4 bleeding events and no surgical revisions in the CytoSorb group compared with 3 and 1 in the control group [10].

It is also noteworthy that no serious device-related adverse events were observed. This important observation is consistent with other published reports and may provide an important advantage for hemoadsorption over the use of reversal agents that may be associated with serious thrombotic complications in this setting. Moreover, DOAC reversal agents are expensive: Andexanet al.pha ranges from \$10,000 to \$12,000, idarucizumab from \$5,000 to \$6,000 [2]. Specifically, for andexanet al.fa, the approved reversal agent for apixaban and rivaroxaban, there are published reports of significant intraoperative complications associated with heparin reversal and resistance during CPB and high risk for thrombotic postoperative complications. Various reports [19–22] warn about the off-label use of andexanet al.fa in CPB-assisted cardiac surgery due to the anticipated development of heparin resistance and anticoagulant rebound. Similarly, a Direct Healthcare Professional Communication letter issued by the European Medicines Agency advises against the use of this reversal agent prior to heparinization since it causes unresponsiveness to the anticoagulant effects of heparin [23]. A recently published comment discussed the problems associated with reversal agents such as andexanet al.pha and the authors concluded that it seems advisable to avoid preoperative administration of andexanet al.pha in cardiothoracic procedures involving the use of a heart–lung machine; they also mentioned alternative methods for eliminating DOACs, such as hemoadsorption [24]. Intraoperative hemoadsorption is increasingly recognized as a valid alternative with a favorable safety profile and recently the latest ESAIC guidelines for the management of major perioperative bleeding [25] include a Class 2 C recommendation for the use of hemoadsorption as an adjunct in patients on ticagrelor or rivaroxaban undergoing emergency cardiac/aortic surgery on cardiopulmonary bypass to reduce bleeding complications. Moreover, Mair et al. recently published their experience with off-pump

CABG and antithrombotic removal by using a stand-alone hemoperfusion pump intraoperatively [26].

Just recently, a group from Oslo published their experience with intraoperative hemoadsorption in 8 emergent apixaban-treated patients presenting with acute type-A aortic dissection. Røed-Undlien and colleagues were able to prove a significant removal of apixaban by intraoperative hemoadsorption as reflected by a significant decrease in apixaban concentrations. Interestingly the authors reported that 50% of the patients received 2,000IE of prothrombin complex concentrate (PCC) and one patient received recombinant activated factor VII (rFVIIa). Intraoperative administration of factor VII per se represents a severe bleeding event as per the UDPB definition. Of note, no surgical revisions have been reported by the authors [27].

In conclusion, these interim results from the ongoing International Safe and Timely Antithrombotic Removal (STAR) Registry show lower rates of major bleeding compared to historical cohort rates in patients on DOACs undergoing cardiac surgery before completion of the recommended washout period. Hemoadsorption appears to be a feasible, safe and potentially effective solution for reducing perioperative bleeding in this patient population.

Limitations

Our study has some major limitations that must be considered when interpreting the results. First, since the current dataset is derived from a single-arm observational registry, a DOAC control group (without adsorber use) is missing. Therefore, definitive conclusions about the effectiveness of the device cannot be made, but only contextualized with existing literature. Second, although the current analysis reflects the largest dataset on this topic, it is still limited by the fairly small number of patients and therefore results should be interpreted with the appropriate caution. Third, safety reporting was not according to GCP-level assessment of all adverse events, but rather via investigator surveillance and oversight for device related adverse events. Fourth, coagulation testing (anti-factor Xa or thromboelastometry) was not performed and therefore specific analyses based on the preoperative degree of anticoagulant DOAC effect cannot be performed. Finally, local institutional transfusion policies may have had an impact on the overall administration of blood products. Future larger studies also including control patients will be required to validate the preliminary results presented in this report to better define the efficacy of intraoperative hemadsorption to reduce perioperative bleeding in these patients.

Abbreviations

ACC	Aortic Cross Clamp
BARC	Bleeding Academic Research Consortium

CABG	Coronary Artery Bypass Grafting
CPB	Cardio-pulmonary bypass
CTD	Chest tube drainage
DOAC	Direct-acting oral anticoagulants
ECMO	Extracorporeal membrane oxygenation
ESAIC	European Society of Anaesthesiology and Intensive Care
STAR	Safe and Timely Antithrombotic Removal
UDPB	Universal Definition of Perioperative Bleeding

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13019-024-03326-1>.

Supplementary Material 1

Acknowledgements

We would like to thank Weihong Fan for her important statistical support that made this analysis possible. We also want to thank Cindy Rechner, Robert Wilke and Antje Fechner for the administrative organization of the registry. Moreover, we thank Harriet Adamson for medical librarian assistance.

Author contributions

Data akquisition was performed (reflecting all local principal investigators) by: M.S., M.T., K.V., T.E., N.M., K.H., A.L., S.L., G.M., R.S. Data cruration and analysis was performed by: M.S., M.T., M.S., D.W., E.D., R.S. The main manuscript was written by M.S., M.T., D.W., E.D., R.S. and all authors reviewed the manuscript including substantial contributions. Figure were prepared by D.W.

Data availability

Data availabilityThe data underlying this article were provided by CytoSorbents Inc., Princeton, NJ, USA. Data will be shared on request to the corresponding author with permission of CytoSorbents.

Declarations

Competing interests

Michael Schmoeckel, Matthias Thielmann, Kambiz Hassan, Stephan Geidel, Sandra Lindstedt, received speaker honoraria and travel fees. Marijana Matejic-Spasic, Daniel Wendt are full-time employees of CytoSorbents Europe GmbH, Berlin, Germany, Efthymios Deliargyris is a full-time employee of CytoSorbents Inc., Princeton, NJ, USA. Robert Storey reports institutional research grants/ support from Cytosorbents and AstraZeneca; and personal fees from Alfagamma, AstraZeneca, Boehringer Ingelheim/Lilly, Chiesi, Cytosorbents, Daiichi Sankyo, Idorsia, Novartis, Novo Nordisk, Pfizer, PhaseBio and Tabuk.

Author details

¹Department of Cardiac Surgery, Klinikum Grosshadern, Ludwig-Maximilians-University, Munich, Germany

²Department of Thoracic- and Cardiovascular Surgery, West German Heart and Vascular Center, Essen, Germany

³Department of Cardiovascular Surgery, German Heart Centre, Munich, Germany

⁴Anesthesiology & Intensive Care Medicine, MediClin Herzzentrum Coswig, Coswig, Germany

⁵Department of Anaesthesia, Royal Brompton Hospital, Royal Brompton & Harefield Hospitals, Part of Guy's and St Thomas' NHS Foundation Trust, London, UK

⁶Department of Cardiac Surgery, Asklepios Klinik St. Georg, Hamburg, Germany

⁷Department of Cardiothoracic and Vascular Surgery, Ulm University Medical Center, Ulm, Germany

⁸Department of Cardiothoracic Surgery and Transplantation, Skåne University Hospital, Lund, Sweden

⁹Department of Cardiac Surgery, Medical University Graz, Graz 8036, Austria

¹⁰CytoSorbents Europe GmbH, Berlin, Germany

¹¹Faculty of Medicine, University Hospital Essen, Essen, Germany

¹²CytoSorbents Inc., Princeton, NJ, USA

¹³Division of Clinical Medicine, University of Sheffield, Sheffield, UK

¹⁴NIHR Sheffield Biomedical Research Centre, Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK

¹⁵Dept. of Cardiac Surgery, LMU Munich, Klinikum Grosshadern, Marchioninstr. 15, D-81377 Munich, Germany

Received: 5 November 2024 / Accepted: 26 December 2024

Published online: 20 January 2025

References

1. Cao D, Amabile N, Chiarito M, Lee VT, Angiolillo DJ, Capodanno D, et al. Reversal and removal of oral antithrombotic drugs in patients with active or perceived imminent bleeding. *Eur Heart J*. 2023;44:1780–94.
2. Douketis JD, Spyropoulos AC. Perioperative Management of patients taking direct oral anticoagulants: a review. *JAMA*. 2024;332:825–34.
3. Budd AN, Wood B, Zheng W, Rong LQ. Perioperative Management of direct oral anticoagulants in cardiac surgery: practice recommendations based on current evidence. *J Cardiothorac Vasc Anesth*. 2022;36:4141–49.
4. Erdoes G, Martinez Lopez De Arroyabe B, Bolliger D, Ahmed AB, Koster A, Agarwal S, et al. International consensus statement on the peri-operative management of direct oral anticoagulants in cardiac surgery. *Anaesthesia*. 2018;73:1535–45.
5. Task Force on Patient Blood Management for Adult Cardiac Surgery of the European Association for Cardio-Thoracic S, the European Association of Cardiothoracic, Boer A, Meesters C, Milojevic MI, Benedetto M. 2017 EACTS/EACTA guidelines on patient blood management for adult cardiac surgery. *J Cardiothorac Vasc Anesth*. 2018;32:88–120.
6. Schmoeckel M, Thielmann M, Hassan K, Geidel S, Schmitt J, Meyer AL, et al. Intraoperative haemoadsorption for antithrombotic drug removal during cardiac surgery: initial report of the international safe and timely antithrombotic removal (STAR) registry. *J Thromb Thrombolysis*. 2024;57:888–97.
7. Tripathi R, Morales J, Lee V, Gibson CM, Mack MJ, Schneider DJ, et al. Antithrombotic drug removal from whole blood using haemoadsorption with a porous polymer bead sorbent. *Eur Heart J Cardiovasc Pharmacother*. 2022;8:847–56.
8. Nierhaus A, Morales J, Wendt D, Scheier J, Gutzler D, Jarczak D, et al. Comparison of the CytoSorb(R) 300 mL and Jafron HA380 hemoabsorption devices: an in vitro study. *Minim Invasive Ther Allied Technol*. 2022;31:1058–65.
9. Nutescu EA, Burnett A, Fanikos J, Spinler S, Wittkowsky A. Pharmacology of anticoagulants used in the treatment of venous thromboembolism. *J Thromb Thrombolysis*. 2016;41:15–31.
10. Hassan K, Thielmann M, Easo J, Kamler M, Wendt D, Haidari Z, et al. Removal of apixaban during emergency cardiac surgery using hemoabsorption with a porous polymer bead sorbent. *J Clin Med*. 2022;11:5889.
11. Matejic-Spasic M, Hassan K, Thielmann M, Geidel S, Storey RF, Schmoeckel M, et al. Management of perioperative bleeding risk in patients on antithrombotic medications undergoing cardiac surgery—a systematic review. *J Thorac Dis*. 2022;14:3030–44.
12. Hassan K, Geidel S, Zamvar V, Tanaka K, Knezevic-Woods Z, Wendt D, et al. Intraoperative ticagrelor removal via hemoabsorption during on-pump coronary artery bypass grafting. *JTCVS Open*. 2023;15:190–96.
13. Matejic-Spasic M, Lindstedt S, Lebreton G, Dzemali O, Suwalski P, Folliquet T, et al. The role of hemoabsorption in cardiac surgery - a systematic review. *BMC Cardiovasc Disord*. 2024;24:258.
14. Hassan K, Bayer N, Schlingloff F, Oberhoffer M, Wohlmuth P, Schmoeckel M, Geidel S. Bleeding complications after use of novel oral anticoagulants in patients undergoing cardiac surgery. *Ann Thorac Surg*. 2018;105:702–08.
15. Chen A, Stecker E. Direct oral anticoagulant use: a practical guide to Common Clinical challenges. *J Am Heart Assoc*. 2020;9:e017559.
16. Hassan K, Kannmacher J, Wohlmuth P, Budde U, Schmoeckel M, Geidel S. Cytosorb Adsorption during Emergency Cardiac operations in patients at high risk of bleeding. *Ann Thorac Surg*. 2019;108:45–51.
17. Fox V, Kleikamp A, Dittrich M, Zittermann A, Flieder T, Knabbe C, et al. Direct oral anticoagulants and cardiac surgery: a descriptive study of preoperative management and postoperative outcomes. *J Thorac Cardiovasc Surg*. 2021;161:1864–74. e2.
18. Ellassal AA, Al-Ebrahim KE, Debs RS, Ragab ES, Faden MS, Fatani MA, et al. Re-exploration for bleeding after cardiac surgery: reevaluation of urgency and factors promoting low rate. *J Cardiothorac Surg*. 2021;16:166.
19. Akhrass R, Gillinov M, Bakaeen F, Akas D, Cameron SJ, Bishop J, et al. Emergency cardiac surgery in patients on oral anticoagulants and antiplatelet medications. *J Card Surg*. 2022;37:214–22.
20. Suzuki Y, Kikura M, Kawashima S, Kimura T, Nakajima Y. Andexanet alpha-induced heparin resistance treated by nafamostat mesylate in a patient undergoing total aortic arch repair for Stanford type A acute aortic dissection: a case report. *JA Clin Rep*. 2024;10:6.
21. Muther M, Schwindt W, Mesters RM, Minnerup J, Stracke P, Holling M, et al. Andexanet-Alfa-Associated Heparin Resistance in the context of hemorrhagic stroke. *Neurocrit Care*. 2022;37:372–76.
22. Apostel H, Winckers K, Bidar E, Schreiber JU. Successful Antithrombin Administration in Andexanet Alfa-Associated Heparin Resistance. *J Cardiothorac Vasc Anesth*. 2021;35:904–07.
23. European Medicines Agency. Direct Healthcare Professional Communication; Ondexxya (andexanet alfa): Avoid use of andexanet prior to heparinization et al. 2020:1–3.
24. Dohle DS, Pfeiffer P, Probst C, Treede H. Andexanet Alfa and heparin resistance in cardiac surgery: experiences and risks associated with the reversal of direct oral anticoagulants. *Eur J Cardiothorac Surg*. 2023;63:ezad100.
25. Kietaibl S, Ahmed A, Afshari A, Albaladejo P, Aldecoa C, Barauskas G, et al. Management of severe peri-operative bleeding: guidelines from the European Society of Anaesthesiology and Intensive Care: second update 2022. *Eur J Anaesthesiol*. 2023;40:226–304.
26. Mair H, Ulrich S, Rosenzweig D, Goeppel J, Jurma C, Vogt F, et al. Antithrombotic drug removal with hemoabsorption during off-pump coronary artery bypass grafting. *J Cardiothorac Surg*. 2024;19:246.
27. Røed-Undlien H, Schultz NH, Husebråten IM, Wollmann BM, Akerkar RR, Molden E, et al. Apixaban removal during emergency surgery for type a acute aortic dissection. A prospective cohort study. *International Journal of Surgery*; 2024.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.