



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/222811/>

Version: Published Version

---

**Article:**

Oxenham, A.F., Braune, T., van Sluijs, E. et al. (2025) New job, new habits? A multilevel interrupted time series analysis of changes in diet, physical activity and sleep among young adults starting work for the first time. *International Journal of Behavioral Nutrition and Physical Activity*, 22. 10.

<https://doi.org/10.1186/s12966-024-01682-8>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**


If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

RESEARCH

Open Access



# New job, new habits? A multilevel interrupted time series analysis of changes in diet, physical activity and sleep among young adults starting work for the first time

Alena F. Oxenham<sup>1\*</sup> , Tanya Braune<sup>1</sup>, Esther van Sluijs<sup>1</sup>, Hannah Fairbrother<sup>3</sup>, Adam Martin<sup>4</sup> and Eleanor M. Winpenny<sup>1,2</sup>

## Abstract

**Background** The workplace is an important determinant of health that people are exposed to for the first-time during adolescence or early adulthood. This study investigates how diet, physical activity, and sleep change as people aged 16–30 years transition into work and whether this varies for different individuals and job types.

**Methods** Multilevel linear regression models assessed changes in fruit and vegetable intake, sleep duration, and physical activity among 3,302 UK Household Longitudinal Study (UKHLS) participants aged 16–30 years, who started work for the first time between 2015 and 2023. In line with interrupted time series analysis, models assessed behavioural trends in the period before starting work, the immediate effect of starting work, and changes in behaviour over time after employment. Stratified analyses examined differences by selected individual and job characteristics, adjusted for covariates. All analyses were conducted in R v.4.3.2.

**Results** Sleep duration was stable over the years before and after starting work, but starting work was associated with an immediate reduction in sleep duration ( $\beta = -9.74$  [95% CI: -17.32 to -2.17 min/night]). Physical activity, measured in Metabolic Equivalent Tasks (METs), increased immediately after starting work ( $\beta = 113.3$ , [95% CI: 80.49 to 146.11] MET-min/day), but subsequently decreased over time after starting work ( $\beta = -26.7$ , [95% CI: -40.75 to -12.66] MET-min/day/year). The increase in physical activity was greater among males, among those with no degree and among those starting lower socioeconomic classification jobs. Starting a “work from home” job had an immediate negative effect on physical activity ( $\beta = -126.42$  [95% CI: -264.45 to 11.61] MET-min/day), whereas those who worked at their employer’s premises showed an initial increase ( $\beta = 128.81$  [95% CI: 89.46 to 168.16] MET-min/day). Starting work had little influence on fruit and vegetable consumption.

**Conclusions** This is the first study to examine how diet, physical activity, and sleep in young adults change as they start employment in the UK. Starting work is associated with decreased sleep time and increased physical activity,

\*Correspondence:  
Alena F. Oxenham  
[alena.oxenham@mrc-epid.cam.ac.uk](mailto:alena.oxenham@mrc-epid.cam.ac.uk)

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

with differences based on sociodemographic and job characteristics. Future research should consider these potential influences of the work environment when developing interventions to promote healthy behaviour in the workplace.

**Keywords** Early adulthood, Employment, Life transitions

## Background

Healthy lifestyles, including behaviours such as adequate sleep, regular physical activity, and a balanced diet, are fundamental to long-term well-being and good physical and mental health [1]. Young adulthood, defined here as ages 16 to 30 is a critical period in health development. Whilst this is typically a period of peak physical health, young adults may also be developing risk factors for chronic disease that arise later in life [2–5]. It is recommended that young adults get between 7 and 9 h of sleep per night [6], engage in at least 150 min of moderate physical activity per week [7], and consume at least 5 portions of fruit and vegetables per day [8]. Notably, obesity prevalence rises sharply during young adulthood, with the majority of the United Kingdom (UK) population living with overweight or obesity by their early thirties [9]. This transitional phase can also involve significant life changes, including transitioning from education to employment, and from familial homes to independent living, all of which might have a detrimental effect on health behaviours including diet, physical activity, and sleep patterns [10]. Understanding the unique challenges and opportunities of the transition from adolescence to young adulthood for the development of long-term health behaviour habits is crucial to supporting healthy adult lifestyles [2, 11, 12].

Starting work is one key transition of young adulthood, encompassing changes in physical and social environments, daily routines and activities, and resources such as time and money, all of which are determinants of health behaviours and later life health [13]. Transitions into employment are becoming increasingly more complex, as young people are spending more time in education, have longer spells of unemployment and spend more time working in part-time jobs [14]. The employment conditions experienced by those starting work are also rapidly changing, with rises in remote and hybrid working since the COVID-19 pandemic, and increases in precarious employment and zero-hour contracts, all of which have impacts on health behaviours [15–17].

A number of previous studies have investigated changes in health behaviours on starting employment, but no studies have been conducted in the UK context. A recent systematic review on changes in physical activity and diet through young adulthood transitions found that two out of three longitudinal studies reported decreases in physical activity through the transition into employment, and one reported no change [18]. Another scoping review reported that entering employment was

associated with decreases in physical activity across several studies, although this was mostly in leisure time physical activity. Four studies in the review did not find any changes in physical activity, and one study reported differences based on sex such that males were less likely to show decreases in physical activity after starting work [19]. The Project Eating and Activity over Time (EAT) study, based in the United States (US), followed up participants aged 11–18 at baseline over four timepoints across fifteen years and assessed life events as well as moderate and vigorous activity. The authors reported that starting work in a full-time job (more than 30 h) was not associated with any changes in physical activity [20]. Overall, the current literature depicts mixed results of decreases or no changes in physical activity after starting work.

Studies examining changes in diet across the transition into employment have also found mixed results. A systematic review conducted in 2020 included only one study which examined entering employment and changes in diet, which reported no significant change after adjusting for confounders [18]. A recent latent growth analysis using Project EAT data found that starting full-time work was associated with increases in diet quality, especially among males [21]. Meanwhile, another analysis of Project EAT data found that starting work was associated with an increase in fast-food intake [22].

To our knowledge, the effect of starting work on sleep duration in young adults has not been studied, but previous longitudinal studies using cohort data from Australia and Brazil have found that a majority of young adults have a negative trajectory of sleep duration and sleep disturbances through the transition into young adulthood from adolescence [23, 24].

Previous studies in adult populations have demonstrated the workplace as an important determinant of health [25–29], with factors such as working shifts, and irregular or long hours having a negative impact on sleep, diet, and physical activity [30–34]. However, few studies have focused on young adults. One cross-sectional study showed that the healthfulness of the work food- and physical activity- environment (i.e. access to fast-food, sugar-sweetened beverages, gyms, walkability etc.) was positively associated with diet and physical activity levels in young adults [35]. However, changes in health behaviours across the transition into the workplace are likely to depend on the characteristics of the job itself.


This study responds to the lack of evidence exploring changes in health behaviours on starting employment in

the United Kingdom (UK). We examine the longitudinal relationship between entering employment and changes in young adults' health behaviours, and how these differ by individual and job characteristics. Looking at individual-level change in health behaviours over time, and in response to the transition of starting work, provides a more causal understanding of how work and work-related factors may be contributing to the development of young adult health behaviour patterns. The aim of this paper is therefore to investigate how starting work for the first time is associated with changes in health behaviours (physical activity, diet and sleep duration), and explore the role of individual (sex, education level and parental social economic position) and job characteristics (shift times, commute mode, job location, work hours) in moderating this relationship.

## Methods

### Study overview and data collection participants and procedures

The UK Household Longitudinal Study (UKHLS) is a panel survey that has collected data annually since 2009 on approximately 40,000 UK households [36], with



**Fig. 1** Inclusion Criteria Flowchart

the most recent data available (wave 13) collected in 2022/2023. Data on all four health behaviours; physical activity, sleep, vegetable consumption, and fruit consumption, were first consistently collected in the adult surveys from wave 7 onwards (2015–2017).

We included in these analyses all UKHLS participants who started work between the ages of 16 and 30, and responded at least once before and once after starting work to the health behaviour questions in waves 7–13. We did not exclude participants based on health status, or family status, to allow our findings to represent the entire young adult population.

All data collection and survey methods were approved by the University of Essex Ethics Committee and informed consent was received prior to completing any interviews or surveys. Detailed information on consent procedures can be found at the UKHLS webpage [37].

### Exposure variables

#### Starting work

Participants were asked to identify their main occupation, from a list including “Self-employed”, “Paid Employment”, “Unemployed”, “On maternity leave”, “Family care or home”, “Full-time student”, “LT sick or disabled”, “Government training scheme”, “Unpaid, family business”, “On apprenticeship”, “On furlough”, “Temporarily laid off/short term working”, “On shared parental leave”, and “Doing something else”. Those identifying as working self-employed, full-time or part-time, or as an apprentice were defined as ‘working’. Those who reported that their main occupation as “Full-time student” were not defined as ‘working’ as working was not their main occupation. A binary variable was generated indicating whether or not the participant had yet started work, coded as 0 for all waves of data collected before starting work and 1 for all waves of data collected after starting work ( $X_{ti}$ ), where starting work was defined as the participant first reporting working self-employed, full-time, part-time, or as an apprentice as their main occupation.

#### Time relative to starting work

Using information on survey completion date (MM/YY) which is reported in UKHLS at every wave for all participants, a continuous variable was generated measuring time (years) in relation to the recorded date of the first wave of data collection after starting work ( $Z_{ti}$ ), such that  $Z_{ti} < 0$  when  $X_{ti} = 0$  and  $Z_{ti} > 0$  when  $X_{ti} = 1$ . In a sensitivity analysis, this variable ( $Z_{ti}$ ) was altered to instead measure time (months) since the reported job start date (MM/YY), a variable which is only available in UKHLS for a subsample of participants (see Fig. 1).

**Table 1** Fruit and Vegetable questions and response options

	Question	Response Options
Fruit frequency	"Including tinned, frozen, dried and fresh fruit, on how many days in a usual week do you eat fruit?"	Never [coded as 0] 1–3 Days [2] 4–6 Days [5] Every day [7]
Fruit amount	"On the days when you eat fruit, how many portions (e.g. an apple, an orange, some grapes) do you eat?"	Values >= 0
Vegetable frequency	"Including tinned, frozen and fresh vegetables, on how many days in a usual week do you eat vegetables? Do not include potatoes, crisps or chips."	Never [coded as 0] 1–3 Days [2] 4–6 Days [5] Every day [7]
Vegetable amount	"On the days when you eat vegetables, how many portions (i.e. 3 heaped tablespoons) do you eat? Please do not include potatoes."	Values >= 0

**Table 2** Job characteristics response options and categorisations

	Response options	Coded categorisations
National Statistics	Lower supervisory & technical, and semi-routine & routine	Low NS-Sect. [0]
Socio-economic classification (NS-SEC)	Management & professional, intermediate, small employers & own account	High NS-Sect. [1]
Shift Times	Mornings, Afternoons, daytimes, Evenings, nighttime shifts, No usual shift pattern, rotating shifts, daytime and evenings, both lunchtimes and evenings	Daytime only Shifts [0] Night-time only shifts [1] Rotating/Varying shifts [2]
Commute Behaviours	"Drive myself by car or van", "Get a lift with someone", "Motorcycle/moped/scooter", "Taxi/minicab" "Bus/coach", "Train", "Underground/Metro/Tram/Light railway", "Cycle", "Walk" NA due to work from home	Inactive commute [0] Active commute [1] No commute [2]
Work Location	"At home" "At your employer's premises" "Driving or travelling around, At one or more other places, or Other"	Work from Home [0] Office Work [1] Travelling/ Varying Work-places [2]
Work Hours	FT employee (>30 h) PT employee (<30 h)	Full-time [0] Part-time [1]

## Outcome variables

### Diet

Vegetable and fruit consumption were measured in two separate questions for each participant every other wave (7, 9, 11, and 13) as shown in Table 1. The questions were adapted from the Household, Income and Labour Dynamics in Australia (HILDA) Survey [38] and the Eating Choices Index (ECI) [39].

A daily vegetable portion measure was calculated by multiplying responses to vegetable frequency and vegetable portion questions and dividing by 7. Daily fruit consumption was assessed in a similar manner.

### Sleep

Participants were asked every three waves [7, 10, 13]: "How many hours of actual sleep did you usually get per night during the last month?" They were advised to indicate the most accurate reply for the majority of days and nights and that this may differ from the actual number of hours spent in bed. The question was taken from the validated Pittsburgh Sleep Quality Index [40].

### Physical activity

Three variables from the validated International Physical Activity Questionnaire (IPAQ) [41] were used to assess participants' weekly physical activity habits in waves 7, 9, 11, 12, and 13: moderate, vigorous and walking activity.

These responses were combined using the formula provided by the IPAQ guidelines to generate a weekly Metabolic Equivalent Tasks (MET)-minutes/week score [42]. The final MET-minutes/week variable was then used to generate a daily MET-min/day estimate.

### Moderators

Data on individual and job characteristics were self-reported at all survey waves. We used data reported at the first wave of data collection after participants started work. Data were recoded to give the following individual characteristics, with job characteristics presented in Table 2.

### Individual characteristics (coded as time invariant)

**Sex** ("Male" [0], "Female" [1]); **Education** (collapsed into categories: "no university degree"[0] or "at least a

university degree”[1]); **Parent’s Education status** (collapsed into categories: (“no university degree”[0] or “at least a university degree”[1]); **Ethnicity** (collapsed into categories: (“White”[0], “Asian” [1], “Black” [2], “Multiple” [3], “Other” [4]).

## Statistical analyses

### Descriptive statistics

Analyses were performed using R Version 2023.12.0 + 369. Descriptive data on outcome and moderator variables were reported by sex.

Models were built up in line with interrupted time series regression, using the following formula:

$$Y_{ti} = \beta_0 + \beta_1 T_{ti} + \beta_2 X_{ti} + \beta_3 Z_{ti} + u_{0i} + T_{ti}u_{1i} + \varepsilon_{ti}$$

Multilevel linear models were fitted separately for each of the health behaviour outcomes ( $Y_{ti}$ ). Weights were not used due to the complexity of the model and due to the focus on within-person changes in outcomes. Models were built up sequentially, adding first the random intercept ( $u_{0i}$ ), then change in outcome over time in years ( $T_{ti}$ ); starting work ( $X_{ti}$ ) to assess an immediate effect of starting work, and time relative to starting work ( $Z_{ti}$ ), indicating time passed since starting work. The model assesses the sustained effect of entering the workforce and how this evolves over time, by allowing an additional intercept and a change in slope in response to starting work. Likelihood ratio tests were conducted to determine the best model fit for each outcome (Supplementary Table 1).

Data on fruit and vegetable consumption and sleep were roughly normally distributed, however physical activity data showed strong positive skew and heteroskedasticity of residuals. We considered applying various transformations but these limited interpretability and a log transformation would have implied an exponential growth curve, which was not a good fit with our data. We therefore decided to model the untransformed data using robust standard errors, benefiting from the robustness of multilevel models to violation of distributional assumptions [43].

We added moderator terms between time in years ( $T_{ti}$ ), starting work ( $X_{ti}$ ) and time relative to starting work ( $Z_{ti}$ ), to explore differences in intercept and slopes between different groups. The interaction with job characteristic models were adjusted for confounders including sex, ethnicity, parental education status, NS-SEC of their first job, and own education status. The interaction with education models were adjusted for sex, ethnicity, and parental education status. The interaction with NS-SEC models were adjusted for education, sex, ethnicity, and parental education status (see Supplementary Fig. 1 for Directed Acyclic Graphs). Multicollinearity was assessed using the `check_collinearity()` function in R [44]

and correlation between the confounders was low (variance inflation factor < 5).

Robust standard errors were used to generate 95% Confidence Intervals for all reported outcomes.

A sensitivity analysis using only participants who had exact job start dates from the wave they first started working was conducted to get a more accurate estimation of changes in behaviour before and after starting employment.

Graphs were generated using the “ggplot” and “predictions” packages in R to generate predicted values from the models to visualise changes in behaviour through the transition into employment. The full Rmarkdown html files for this analysis are available at: [https://osf.io/wbscr/?view\\_only=fa9d385bb81e4c709fcb3fc11b0896e3](https://osf.io/wbscr/?view_only=fa9d385bb81e4c709fcb3fc11b0896e3).

## Results

### Descriptive statistics

A total of 3,302 participants met the inclusion criteria and were included in the analyses (Fig. 1). The mean age of participants when first starting working as their main occupation was 21.5 years (s.d.: 3.3), 57.2% were female, and 38.8% had a degree (Table 1). 184 people were included in a Supplementary Analysis where an exact job start date was reported (Supplementary Table 3) (Table 3).

### Interrupted time series analysis

The main results are reported in Tables 4, Supplementary Tables 3–5; Fig. 2. The unadjusted models are in the Supplementary Tables 6–9 (Supplementary Material) and were not dissimilar to the main results. Due to issues with singularity, random slope terms ( $T_{ti}u_{1i}$ ) were not included in the vegetable intake models. Random slope terms were also not included in the physical activity or sleep models as they did not improve model fit, but were included in the fruit intake model. The addition of quadratic time variables was tested to allow the change in exposure over time to be non-linear. Addition of these quadratic terms did improve model fit, but the coefficients of these terms were very small and not significant, therefore they were dropped from the model to aid easy interpretation (Supplementary Table 1).

### Changes in physical activity through the employment transition

Physical activity increased as people started work ( $\beta = 113.3$ , [95% CI: 80.49 to 146.11] MET-min/day) and then decreased over time after starting work ( $\beta = -26.7$ , [95% CI: -40.75 to -12.66] MET-min/day/year) (Fig. 2; Table 4).

Physical activity increased more in males ( $\beta = 180.97$  [95% CI: 124.39 to 237.55] MET-min/day) than in females ( $\beta = 62.58$  [95% CI: 24.66 to 100.5] MET-min/

**Table 3** Descriptive statistics at first instance of reported employment (n; (%))

Variable	Male n = 1412 (42.8%)	Female n = 1890 (57.2%)
<b>Outcome Variables (Mean (SD))</b>		
Physical Activity (MET-min/day)	650.1 (643.3)	439.1 (475.5)
Sleep (hours)	7.3 (1.2)	7.3 (1.2)
Vegetable Portions	1.7 (1.6)	1.9 (1.6)
Fruit Portions	1.2 (1.2)	1.4 (1.3)
<b>Individual Characteristics</b>		
Age (Mean (SD))	21.3 (3.1)	21.6 (3.5)
Ethnicity (n (%))		
White	958 (67.8)	1347 (71.3)
Asian	283 (20.1)	331 (17.5)
Black	90 (6.4)	104 (5.5)
Multiple	66 (4.7)	81 (4.3)
Other	13 (0.9)	27 (1.4)
Missing	2 (0.3)	0 (0)
Wave first started work (n (%))		
8 (2016–2018)	273 (19.3)	303 (16.0)
9 (2017–2019)	288 (20.3)	484 (25.6)
10 (2018–2020)	282 (20.0)	337 (17.8)
11 (2019–2021)	202 (14.3)	308 (16.3)
12 (2020–2022)	181 (12.8)	227 (12.0)
13 (2021–2023)	186 (13.2)	231 (12.2)
Education (n (%))		
No University Degree	885 (62.7)	1097 (58.0)
University Degree or higher	510 (36.1)	766 (40.5)
Missing	17 (1.2)	27 (1.4)
NS-SEC (n (%))		
Low	718 (50.9)	983 (52.0)
High	482 (34.1)	718 (38.0)
Missing	212 (15.0)	189 (10.0)
Parental Education (n (%))		
No University Degree	688 (48.7)	823 (43.5)
University Degree or higher	635 (45.0)	826 (43.7)
Missing	89 (6.3)	241 (12.8)
<b>Job Characteristics</b>		
Commute Behaviour (n (%))		
Inactive	551 (39.0)	914 (48.4)
Active	508 (36.0)	656 (34.7)
No Commute	66 (4.7)	62 (3.3)
Missing	287 (20.3)	258 (13.7)
Work Location (n (%))		
Office Work	1073 (76.0)	1590 (84.1)
Work from home	66 (4.7)	62 (3.3)
Driving around/ Travel Work	137 (9.7)	110 (5.8)
Missing	136 (9.6)	128 (6.8)
Shift Times (n (%))		
Daytime	527 (37.0)	688 (36.4)
Night-time	90 (6.4)	101 (5.3)
Rotating	308 (21.8)	517 (27.4)
Missing	487 (34.5)	584 (30.9)
Work Hours (n (%))		
Full Time (30+ hours)	841 (59.6)	956 (50.6)

**Table 3** (continued)

Variable	Male n = 1412 (42.8%)	Female n = 1890 (57.2%)
Part Time (less than 30)	444 (31.4)	830 (43.9)
Missing	127 (9.0)	104 (5.5)


**Table 4** Changes in MET-min per day over time before and after starting work, showing adjusted interactions with different individual and job characteristics. Initial slope shows baseline trend per year before employment transition, transition into work shows change in intercept immediately after starting work, and change in slope after starting work shows longer-term trends in MET-min per day each year after starting work

		Initial slope [95% CI]	Transition into work [95% CI]	Change in slope after starting work [95% CI]
Overall Effect [n = 3,080]		8.15 [-1.53, 17.84]	113.30 [80.49, 146.11]	-26.70 [-40.75, -12.66]
Sex [n = 3,080]	Male	-1.95 [-18.39, 14.49]	180.97 [124.39, 237.55]	-24.92 [-49.04, -0.80]
	Female	17.07 [5.97, 28.17]	62.58 [24.66, 100.50]	-29.33 [-45.62, -13.04]
	<i>Interaction p-value</i>	0.060	0.001	0.766
University Education [n = 2,753]	Attended university	11.91 [0.15, 23.67]	58.98 [14.18, 103.78]	-39.03 [-56.92, -21.15]
	No university	5.19 [-12.86, 23.24]	168.66 [116.03, 221.28]	-21.40 [-45.13, 2.33]
	<i>Interaction p-value</i>	0.541	0.002	0.245
NSSEC [n = 2,440]	High	13.76 [0.73, 26.80]	21.56 [-24.11, 67.23]	-21.55 [-39.98, -3.12]
	Low	0.06 [-17.54, 17.67]	210.47 [156.76, 264.18]	-32.17 [-56.97, -7.37]
	<i>Interaction p-value</i>	0.218	< 0.001	0.500
Work Hours [n = 2,397]	Full-time	1.60 [-11.64, 14.84]	128.25 [80.84, 175.65]	-25.10 [-44.87, -5.33]
	Part-time	7.25 [-11.68, 26.17]	132.19 [74.92, 189.45]	-25.07 [-50.86, 0.71]
	<i>Interaction p-value</i>	0.630	0.917	0.999
Shift Times [n = 1,793]	Day time	-4.11 [-21.90, 13.68]	125.60 [70.04, 181.16]	-10.36 [-33.67, 12.95]
	Night	-8.70 [-79.01, 61.62]	211.20 [51.64, 370.77]	-35.40 [-125.65, 54.86]
	<i>Interaction p-value</i>	0.901	0.320	0.598
	Rotate	15.16 [-8.21, 38.53]	90.22 [13.11, 167.32]	-37.79 [-71.26, -4.32]
	<i>Interaction p-value</i>	0.195	0.462	0.187
Commute [n = 2,183]	Inactive	14.00 [-1.66, 29.67]	119.61 [65.17, 174.05]	-35.11 [-58.43, -11.80]
	Active	0.97 [-16.56, 18.49]	143.41 [89.91, 196.91]	-23.12 [-47.83, 1.59]
	<i>Interaction p-value</i>	0.272	0.538	0.487
	No Commute	7.04 [-25.62, 39.70]	-125.64 [-263.64, 12.36]	40.63 [-13.16, 94.43]
	<i>Interaction p-value</i>	0.706	0.001	0.011
Work Location [n = 2,356]	Office	6.25 [-5.93, 18.43]	128.81 [89.46, 168.16]	-27.39 [-44.58, -10.19]
	Home	6.91 [-25.77, 39.59]	-126.42 [-264.45, 11.61]	39.85 [-13.89, 93.59]
	<i>Interaction p-value</i>	0.970	< 0.001	0.019
	Travel	-3.76 [-44.06, 36.54]	243.82 [106.57, 381.08]	-47.50 [-104.96, 9.96]
	<i>Interaction p-value</i>	0.641	0.114	0.510

day), (Fig. 3). Participants who did not have a university degree showed a greater increase in physical activity ( $\beta = 168.66$  [95% CI: 116.03 to 221.28] MET-min/day) compared to those with a university degree ( $\beta = 58.98$  [95% CI: 14.18 to 103.78] MET-min/day). Working from home was associated with an initial decrease in physical activity ( $\beta = -126.42$ , [-264.45 to 11.61] MET-min/day), whereas those who worked in an office/employer's premises showed an initial increase ( $\beta = 128.81$  [95% CI: 89.46 to 168.16] MET-min/day), although this difference was not maintained.

#### Changes in sleep duration through the employment transition

Usual minutes of sleep per night stayed stable over time ( $\beta = -1.05$  [95% CI: -3.44 to 1.33] min/night/year), and then decreased immediately after starting work ( $\beta = -9.74$  [95% CI: -17.32 to -2.17], min/night). After starting work, sleep did not change over time ( $\beta = 0.97$ , [95% CI: -1.77 to 3.71] min/night/year) (Fig. 2, Supplementary Table 3). Sleep duration did not differ by sex or NS-SEC status. While participants both with and without a university degree showed an initial decrease in sleep



**Fig. 2** Predicted trajectories of physical activity, sleep, vegetable and fruit intake across the transition into employment. Description: Changes in four outcome behaviours through transition into employment. 95% CIs are represented by grey bands. **A)** shows change in physical activity, **b)** shows change in sleep, **c)** shows change in vegetable intake, and **d)** show change in fruit intake


duration after starting work, trends in behaviour differed over time. Participants who did not have a university degree showed a negative change in slope after starting work ( $\beta = -3.05$  [95% CI: -7.74 to 1.64] min/night/year) resulting in decreasing sleep over time, whereas those with a university degree showed a positive change in slope ( $\beta = 3.49$  [95% CI: 0.00 to 6.98] min/night/year) leading to an increase in sleep over time (Fig. 4b). There were no differences in sleep duration based on job characteristics (Supplementary Table 3).

#### Changes in diet through the employment transition

Daily vegetable portions showed a slight 0.05 portion/day increase over time [95% CI: 0.01 to 0.08] but starting work ( $\beta = 0.02$  [95% CI: -0.07 to 0.12] vegetable portions/day) and time in employment ( $\beta = -0.04$  [95% CI: -0.08 to 0.00] vegetable portions/day/year) had no significant effect on vegetable intake (Fig. 2, Supplementary Table 4). The transition into work was not associated with a significant change based on any individual or job characteristics.

Daily fruit portions showed a decrease of -0.04 portions/day each year before starting work [95% CI: -0.07 to -0.01 portions/day/year]. Starting work and time in employment had no significant effect on fruit intake (Fig. 2, Supplementary Table 5). There were no differences in fruit intake based on sex. There was a significant interaction effect based on education status ( $p=0.037$ ), such that participants who did not attend university showed a decrease in daily fruit intake after starting work ( $\beta = -0.12$ , [95%CI: -0.24 to 0.00] portions/day) and those who had a university degree showed an increase after starting work ( $\beta = 0.06$  [95% CI: -0.06 to 0.18] portions/day). No differences in fruit consumption were found based on job characteristics.

A supplementary analysis of changes in health behaviours through the employment transition using participants who reported an exact job start date ( $n=184$ ) is reported in Supplementary Table 2 in the Appendix. The results do not differ directionally from our main results.



**Fig. 3** Physical activity (MET-min/day) trajectories by individual and job characteristics. Description: Changes in physical activity through the transition into employment. 95% CIs are represented by grey bands. Change in physical activity by **(a)** sex, **(b)** education status, adjusted for sex, ethnicity, and parent education, **(c)** NS-SEC, adjusted for sex, ethnicity, parent education and own education, **(d)** Commute, adjusted for sex, ethnicity, parent education, own NS-SEC, and own education


## Discussion

Analysis of changes in physical activity, sleep duration, and diet suggest that starting work for the first time has important implications for developing health behaviours in young adults. Our findings showed that the greatest change in response to starting work was seen in physical activity, and moderate changes were seen in sleep duration. We found little association between starting work and changes in diet. Physical activity showed an initial increase on starting work of 113.3 MET-min/day, comparable to about 30 min of moderate activity per day (e.g. cycling at a regular pace) [32]. The increase in physical activity was greatest among those of lower socioeconomic position, e.g. lower education and lower NS-SEC, as well as among males. Sleep patterns displayed a modest decline, on average 9.7 min per night, as young people entered the workforce, and then remained stable over the following years. After adjusting for confounders, only education status interacted with changes in sleep, such that people without a degree showed a continuing decline

of about 3 minutes of sleep per night each year after starting work, while those with a degree showed a positive trajectory of usual minutes of sleep per night after starting work. Starting work showed limited associations with changes in diet. There was no overall effect of entering employment on vegetable intake, and no differences were seen based on any individual or job characteristics. No overall changes in fruit intake were seen after starting work, although those with no higher education showed a short-term decrease in daily intake after starting work compared to those with a university degree or higher.

## Implications of the findings and comparison with previous research

Increases in physical activity were more pronounced among individuals who are more likely to work in manual labour jobs, such as males and those with lower socioeconomic positions, likely due to their work requiring more physical activity [45]. The differences in physical activity due to education status persisted over time, with people



**Fig. 4** Sleep trajectories by individual and job characteristics. Figure Description: Changes in sleep duration through transition into employment. 95% CIs are represented by grey bands. **A)** Shows changes in behaviour by sex, **b)** By education status adjusted for sex, ethnicity, and parent education **c)** By shift time adjusted for sex, ethnicity, parent education, own nssec, and own education, **c)** By Commute behaviour adjusted for sex, ethnicity, parent education, own nssec, and own education

with a university degree consistently reporting lower levels of activity than those with no degree after starting work. Although stratifying by participants' NS-SEC status showed similar differences immediately after starting work, these differences did not remain over time. People with a low NS-SEC job showed a steeper decline in physical activity each year after starting work, such that the difference in activity levels between those low NS-SEC and high NS-SEC groups grew smaller over time. This suggests that education status and NS-SEC have separate mechanisms of action on physical activity levels. The NS-SEC effect may be related to job role, which may well change over time as people get more senior.

Although physical activity increased more after starting employment among groups more likely to encounter physically demanding work [45–47], a previous meta-analysis found that occupational physical activity may not have the same benefits in terms of cardiovascular outcomes compared to leisure-time physical activity [48]. More research is needed to differentiate between the

longitudinal health effects of occupational physical activity and 'leisure-time' physical activity, as there may be negative long-term effects of work-related stressors and work quality related to occupational physical activity.

Commuting behaviours also seemed to play a role in changes in physical activity through the transition of starting work. Those with no commute showed a non-significant decrease in physical activity after starting work compared to increases in activity in both active and inactive commuters. Relatedly, working from home also showed a decrease in physical activity compared to those who worked in offices or worked elsewhere out of the home. Previous studies have found that active commuting is associated with positive outcomes in physical activity and physical fitness compared to no active commutes [49, 50].

While usual hours of sleep only showed a slight decrease after starting work, people without a university degree seemed to show more long-term negative effects after starting work. This is consistent with previous

findings from the US Behavioral Risk Factor Surveillance study that found associations between low occupational social class and sleep disturbances [51]. While our study did not find any associations between shift work and sleep duration, a systematic review reported that both psychosocial factors and shift times are associated with sleep disturbances [25]. It may be that sleep quality is more strongly influenced by the workplace than sleep duration. Sleep and work have previously been connected in a negative cycle where poor sleep quality is associated with more negative events and perceived stress the following day, and these negative events are also associated with poorer sleep quality the following night [52–54].

We found very limited changes in fruit and vegetable intakes after starting work, which does not align with previous work on workplace environments and diet, as well as other longitudinal studies assessing life transitions and diet [21, 22, 31, 35, 55]. However, these previous studies were all conducted in different contexts (mainly the U.S. and Norway).

### Strengths and limitations

This the first study to examine how health behaviours change in young adults when they start employment in the UK. The study design using a multilevel interrupted-time series approach improves causal inference compared to other study designs by including a baseline trend before starting work and allowing modelling of both immediate and long-term effects of starting work.

We use data from a national panel survey, representative of the UK population, increasing the generalizability of the findings within the UK context. The survey included detailed questions on job characteristics as well as health behaviour measures, over many years, allowing for longitudinal analysis. However, all the variables used were self-reported, as there was no objective measurement of physical activity, sleep, or diet in this dataset, providing opportunity for reporting bias. Additionally, the diet variables only assessed fruit and vegetable intake, and while increased fruit and vegetable consumption have been linked to a better overall diet [56], they do not provide a comprehensive insight into overall diet quality. Given the limited availability of longitudinal surveys that provide high-quality, repeated measures of diet quality over the early adulthood period, the available fruit and vegetable intake variables were considered appropriate for this analysis.

Our study included a large sample size of over 3,000 participants, providing data across multiple time points. Despite this, some groups (e.g., work-from-home or night-time and rotating shift workers) had small numbers, which may contribute to small effect sizes and large confidence intervals. Additionally, due to missing data regarding exact job start dates, the interview date from

participants' first wave of reported working was used instead, which may limit the accuracy and interpretation of results. We may not see the true immediate effect, as participants could have up to 2 years from when the transition takes place to their first recorded data point. However, a sensitivity analysis that only included participants with an exact job start date showed similar direction and magnitude of findings.

### Conclusions

This analysis revealed a clear impact of starting work on changes in physical activity and sleep among young adults. A positive impact of starting work was seen on physical activity, and a negative impact through reduction in sleep duration. Changes in physical activity in response to starting work differed by job characteristics, highlighting the potential influence of the work environment on health behaviours. This suggests that the workplace could be a good opportunity for health promotion. Workplace health promotion interventions promoting healthier diets, physical activity, and sleep in young adults could result in healthier employees and fewer sick days. Targeting young adulthood for health behaviour interventions could prevent future health issues, combat rising obesity rates, and promote positive generational changes. This study highlights the importance of understanding how employment transitions impact young adults' health behaviours and the potential for policy interventions to support healthier lifestyles among young working adults. Future research should focus on the interplay between individual characteristics, job attributes, and job quality in shaping long-term health outcomes.

### Abbreviations

UKHLS	UK Household Longitudinal Study
UK	United Kingdom
US	United States
Project EAT	The Project Eating and Activity across Time
IPAQ	International Physical Activity Questionnaire
MET	Metabolic Equivalent Tasks
NSSEC	National Statistics Socio-economic classification
HILDA Survey	Household, Income and Labour Dynamics in Australia
ECI	Development of the Eating Choices Index

### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12966-024-01682-8>.

#### Supplementary Material 1: Supplementary Table 1: Likelihood

**Ratio Tests** Description: Additional file 1: A table depicting likelihood ratio tests conducted to determine the best model fit for each outcome.

#### Supplementary Material 2: Supplementary Table 2: Sensitivity

**Analysis** Description: Additional file 2: Table of changes in health behaviours among participants with an exact job start date.

#### Supplementary Material 3: Supplementary Figure 1: DAGs

Description: Additional file 3: Directional Acyclic Graphs to show how each interaction model was adjusted for.

**Supplementary Material 4: Supplementary Tables 3-5** Description: Additional file 4: Tables of overall and adjusted interaction models for sleep, vegetable & fruit intake models.

**Supplementary Material 5: Supplementary Tables 6-9: Unadjusted models** Description: Additional file 5: Tables of unadjusted interaction models for sleep, vegetable & fruit intake models.

### Acknowledgements

We would like to acknowledge the contributions of Dr. Yinhua Tao for guidance on statistical analysis and visualisation methods, as well as thank the study participants for their contribution in providing the data used in this analysis. We would also like to thank the members of our Young Persons Advisory Group and project stakeholders for their feedback. Understanding Society is an initiative funded by the Economic and Social Research Council and various Government Departments, with scientific leadership by the Institute for Social and Economic Research, University of Essex, and survey delivery by the National Centre for Social Research (NatCen) and Verian (formerly Kantar Public). The research data are distributed by the UK Data Service. The COVID-19 study (2020–2021) was funded by the Economic and Social Research Council and the Health Foundation. Serology testing was funded by the COVID-19 Longitudinal Health and Wealth – National Core Study. Fieldwork for the web survey was carried out by Ipsos MORI and for the telephone survey by Kantar.

### Author contributions

AFO conducted the analysis and drafted the manuscript. EMW, AM, EvS and HF supported the conceptualization of the study, interpreting results and drafting, reviewing and editing the manuscript. TB provided additional guidance on coding in R. All authors reviewed drafts of the manuscript.

### Funding

This work was funded by the NIHR [Work and Health Research Programme, grant number NIHR206285]. The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care. EMW is funded by a Career Development Award from the UK Medical Research Council [grant number MR/T010576/1]. TB was funded by the Cambridge MRC Doctoral Training Partnership [grant number MR/N013433/1] and the Elizabeth McDowell Studentship at Newnham College, Cambridge. The contribution of EvS was supported by the Medical Research Council [grant number MC\_UU\_00006/5].

### Data availability

The data that support the findings of this study are available on request from the UK Data Service <https://doi.org/10.5255/UKDA-SN-6614-19>, study number (SN) 6614. The data are publicly available, however, they are considered safeguarded and therefore require users to register and accept the End User Licence. All Rmarkdown files are available on OSF at: [https://osf.io/wbscr/?view\\_only=fa9d385bb81e4c709fcb3fc11b0896e3](https://osf.io/wbscr/?view_only=fa9d385bb81e4c709fcb3fc11b0896e3).

### Declarations

#### Ethics approval and consent to participate

The University of Essex Ethics Committee has approved all data collection on Understanding Society main study and innovation panel waves, including asking consent for all data linkages except to health records. Requesting consent for health record linkage was approved at Wave 1 by the National Research Ethics Service (NRES) Oxfordshire REC A (08/H0604/124), at BHPS Wave 18 by the NRES Royal Free Hospital & Medical School (08/H0720/60) and at Wave 4 by NRES Southampton REC A (11/SC/0274). Approval for the collection of biosocial data by trained nurses in Waves 2 and 3 of the main survey was obtained from the National Research Ethics Service (Understanding Society – UK Household Longitudinal Study: A Biosocial Component, Oxfordshire A REC, Reference: 10/H0604/2). For further details on the various committees which have provided ethical approval of the Understanding Society study and its components as appropriate see below:

#### Main survey: Ethics approval was received from the University of Essex Ethics Committee

By letter dated 6 July 2007 for Waves 1 and 2. By letter dated 17 December 2010 for Waves 3 to 5. By letter dated 20 August 2013 for Waves 6 to 8. By letter dated 4 October 2016 for Waves 9–11. Ethics Approval number ETH1920-0123 for Wave 12. number ETH2021-0015 for Wave 13. Ethics Approval number ETH2122-0246 for Wave 14. Ethics Approval number ETH2223-0264 for Wave 15. Ethics approval number 22/EE/0260 was received from the NHS Research Ethics Committee for Wave 16.

#### Consent for publication

Not Applicable.

#### Competing interests

The authors have declared no competing interest.

#### Author details

<sup>1</sup>MRC Epidemiology Unit, University of Cambridge Level 3 Institute of Metabolic Science, University of Cambridge School of Clinical Medicine, Cambridge CB2 0SL, UK

<sup>2</sup>Mohn Centre for Children's Health and Wellbeing, School of Public Health, Imperial College London, London, UK

<sup>3</sup>Faculty of Health, University of Sheffield, Sheffield, UK

<sup>4</sup>Academic Unit of Health Economics, Leeds Institute of Health Sciences, University of Leeds, Leeds LS2 9TJ, UK

Received: 19 July 2024 / Accepted: 5 November 2024

Published online: 28 January 2025

### References

1. Dale H, Brassington L, King K. The impact of healthy lifestyle interventions on mental health and wellbeing: a systematic review. *Ment Health Rev J*. 2014;19(1):1–26.
2. Nelson MC, Story M, Larson NI, Neumark-Sztainer D, Lytle LA. Emerging Adulthood and College-aged youth: an overlooked age for weight-related Behavior Change. *Obesity*. 2008;16(10):2205–11.
3. Alberga AS, Sigal RJ, Goldfield G, Prud'homme D, Kenny GP. Overweight and obese teenagers: why is adolescence a critical period? *Pediatr Obes*. 2012;7(4):261–73.
4. Kimm SY, Glynn NW, Kriska AM, Fitzgerald SL, Aaron DJ, Similo SL, et al. Longitudinal changes in physical activity in a biracial cohort during adolescence. *Med Sci Sports Exerc*. 2000;32(8):1445–54.
5. Lien N, Lytle LA, Klepp KI. Stability in consumption of fruit, vegetables, and sugary foods in a cohort from age 14 to age 21. *Prev Med*. 2001;33(3):217–26.
6. Hirshkowitz M, Whitton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health J Natl Sleep Found*. 2015;1(1):40–3.
7. WHO guidelines on physical activity and sedentary behaviour [Internet]. [cited 2024 Oct 28]. <https://www.who.int/publications/i/item/9789240015128>
8. nhs.uk [Internet]. 2022 [cited 2024 Oct 28]. Why 5 A Day? <https://www.nhs.uk/live-well/eat-well/5-a-day/why-5-a-day/>
9. Digital NHS, NHS Digital. 2022 [cited 2023 Sep 20]. Health Survey for England, 2021 part 1: Overweight and obesity in adults. <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/2021/overweight-and-obesity-in-adults>
10. Koehn S, Gillison F, Standage M, Bailey J. Life transitions and relevance of healthy living in late adolescence. *J Health Psychol*. 2016;21(6):1085–95.
11. Johnson W, Li L, Kuh D, Hardy R. How has the age-related process of overweight or obesity development changed over Time? Co-ordinated Analyses of Individual Participant Data from Five United Kingdom Birth Cohorts. *PLOS Med*. 2015;12(5):e1001828.
12. Patton GC, Sawyer SM, Santelli JS, Ross DA, Afifi R, Allen NB, et al. Our future: a Lancet commission on adolescent health and wellbeing. *Lancet*. 2016;387(10036):2423–78.
13. Dahlgren G, Whitehead M. Policies and strategies to promote social equity in health [Internet]. Institute for Future Studies; 1991. (Background document to WHO – Strategy paper for Europe). <https://core.ac.uk/download/pdf/6472456.pdf>

14. Pelikh A, Rowe F. Increasing diversity, precarity and prolonged periods of education in the transition from school to work in Britain. *Popul Space Place*. n/a(n/a):e2771.
15. Oddo VM, Zhuang CC, Andrea SB, Eisenberg-Guyot J, Peckham T, Jacoby D, et al. Changes in precarious employment in the United States: a longitudinal analysis. *Scand J Work Environ Health*. 2021;47(3):171–80.
16. Koumenta M, Williams M. An anatomy of zero-hour contracts in the UK. *Ind Relat J*. 2019;50(1):20–40.
17. Bick A, Blandin A, Mertens K. Work from Home before and after the COVID-19 outbreak. *Am Econ J Macroecon*. 2023;15(4):1–39.
18. Winpenny EM, Smith M, Penney T, Foubister C, Guagliano JM, Love R, et al. Changes in physical activity, diet, and body weight across the education and employment transitions of early adulthood: a systematic review and meta-analysis. *Obes Rev*. 2020;21(4):e12962.
19. Gropper H, John JM, Sudeck G, Thiel A. The impact of life events and transitions on physical activity: a scoping review. *PLoS ONE*. 2020;15(6):e0234794.
20. Miller J, Nelson T, Barr-Anderson DJ, Christoph MJ, Winkler M, Neumark-Sztainer D. Life events and Longitudinal effects on physical activity: adolescence to Adulthood. *Med Sci Sports Exerc*. 2019;51(4):663.
21. Tao Y, Wall M, Larson N, Neumark-Sztainer D, Winpenny EM. Changes in diet quality across life transitions from adolescence to early adulthood: a latent growth analysis [Internet]. 2024 [cited 2024 Apr 26]. <https://doi.org/10.1101/2024.02.14.24302819>
22. Winpenny EM, Winkler MR, Stochl J, Van Sluijs EMF, Larson N, Neumark-Sztainer D. Associations of early adulthood life transitions with changes in fast food intake: a latent trajectory analysis. *Int J Behav Nutr Phys Act*. 2020;17(1):130.
23. McVeigh JA, Smith A, Howie EK, Stamatakis E, Ding D, Custilli PA, et al. Developmental trajectories of sleep during childhood and adolescence are related to health in young adulthood. *Acta Paediatr*. 2021;110(8):2435–44.
24. Kramer Fiala Machado A, Wendt A, Baptista Menezes AM, Gonçalves H, Wehrmeister FC. Sleep duration trajectories from adolescence to emerging adulthood: findings from a population-based birth cohort. *J Sleep Res*. 2021;30(3):e13155.
25. Linton SJ, Kecklund G, Franklin KA, Leissner LC, Sivertsen B, Lindberg E, et al. The effect of the work environment on future sleep disturbances: a systematic review. *Sleep Med Rev*. 2015;23:10–9.
26. Clohessy S, Walasek L, Meyer C. Factors influencing employees' eating behaviours in the office-based workplace: a systematic review. *Obes Rev*. 2019;20(12):1771–80.
27. Prevalence of Short Sleep Duration by Industry and Occupation in the National Health Interview Survey. | SLEEP | Oxford Academic [Internet]. [cited 2024 Apr 25]. <https://academic.oup.com/sleep/article/33/2/149/2454438>
28. Väisänen D, Kallings LV, Andersson G, Wallin P, Hemmingsson E, Ekblom-Bak E. Lifestyle-associated health risk indicators across a wide range of occupational groups: a cross-sectional analysis in 72,855 workers. *BMC Public Health*. 2020;20:1656.
29. Kirk MA, Rhodes RE. Occupation correlates of adults' participation in leisure-time physical activity: a systematic review. *Am J Prev Med*. 2011;40(4):476–85.
30. Harknett K, Schneider D, Wolfe R. Losing sleep over work scheduling? The relationship between work schedules and sleep quality for service sector workers. *SSM - Popul Health*. 2020;12:100681.
31. Tanaka R, Tsuji M, Tsuchiya T, Kawamoto T. Association between Work-related factors and Diet: a review of the literature. *Workplace Health Saf*. 2019;67(3):137–45.
32. Reed JL, Prince SA, Pipe AL, Attallah S, Adamo KB, Tulloch HE, et al. Influence of the workplace on physical activity and cardiometabolic health: results of the multi-centre cross-sectional Champlain nurses' study. *Int J Nurs Stud*. 2018;81:49–60.
33. Weston G, Zilanawala A, Webb E, Carvalho L, McMunn A. Work hours, weekend working, nonstandard work schedules and sleep quantity and quality: findings from the UK household longitudinal study. *BMC Public Health*. 2024;24(1):309.
34. Åkerstedt T, Nordin M, Alfredsson L, Westerholm P, Kecklund G. Sleep and sleepiness: impact of entering or leaving Shiftwork—a prospective study. *Chronobiol Int*. 2010;27(5):987–96.
35. Watts A, Laska M, Larson N, Neumark-Sztainer D. Millennials at work: workplace environments of young adults and associations with weight-related health. *J Epidemiol Community Health*. 2015;70.
36. University of Essex, Institute for Social and Economic Research. Understanding Society: Waves 1–13, 2009–2022 and Harmonised BHPS: Waves 1–18, 1991–2009. UK Data Service. SN: 6614; 2023.
37. Consent information [Internet]. Understanding Society. [cited 2024 Oct 28]. <https://www.understandingsociety.ac.uk/documentation/mainstage/consents/>
38. HILDA Survey [Internet]. [cited 2024 Oct 28]. <https://melbourneinstitute.unimelb.edu.au/hilda>
39. Pot GK, Richards M, Prynne CJ, Stephen AM. Development of the eating choices Index (ECI): a four-item index to measure healthiness of diet. *Public Health Nutr*. 2014;17(12):2660–6.
40. Shahid A, Wilkinson K, Marcu S, Shapiro CM. Pittsburgh Sleep Quality Index (PSQI). In: Shahid A, Wilkinson K, Marcu S, Shapiro CM, editors. STOP, THAT and One Hundred Other Sleep Scales [Internet]. New York, NY: Springer New York; 2011 [cited 2024 Oct 25]. pp. 279–83. [https://link.springer.com/https://doi.org/10.1007/978-1-4419-9893-4\\_67](https://link.springer.com/https://doi.org/10.1007/978-1-4419-9893-4_67)
41. Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc*. 2003;35(8):1381–95.
42. Committee IR. Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ)-Short and Long Forms. <http://www.ipaq.ki.se/scoring.pdf> [Internet]. 2005 [cited 2024 Jul 5]; <https://cir.nii.ac.jp/crid/1573950400545533440>
43. Robustness of linear mixed-effects models to violations of distributional assumptions - Schielzeth – 2020 - Methods in Ecology and Evolution - Wiley Online Library [Internet]. [cited 2024 Jul 5]. <https://besjournals.onlinelibrary.wiley.com/doi/full/https://doi.org/10.1111/2041-210X.13434>
44. easystats [Internet]. 2019 [cited 2024 Oct 25]. Check your (Mixed) Model for Multicollinearity with 'performance'. [https://easystats.github.io/blog/posts/performance\\_check\\_collinearity/](https://easystats.github.io/blog/posts/performance_check_collinearity/)
45. Popham F, Mitchell R. Relation of employment status to socioeconomic position and physical activity types. *Prev Med*. 2007;45(2):182–8.
46. Gabriel PE, Schmitz S. Gender differences in occupational distributions among workers [Internet]. Bureau of Labor Statistics; 2007. (Monthly Labor Review). <https://www.bls.gov/opub/mlr/2007/06/art2full.pdf>
47. Biswas A, Harbin S, Irvin E, Johnston H, Begum M, Tiong M, et al. Sex and gender differences in Occupational Hazard exposures: a scoping review of the recent literature. *Curr Environ Health Rep*. 2021;8(4):267–80.
48. Cillekens B, Huysmans MA, Holtermann A, van Mechelen W, Straker L, Krause N, et al. Physical activity at work may not be health enhancing. A systematic review with meta-analysis on the association between occupational physical activity and cardiovascular disease mortality covering 23 studies with 655 892 participants. *Scand J Work Environ Health*. 2022;48(2):86–98.
49. Yang L, Panter J, Griffin SJ, Ogilvie D. Associations between active commuting and physical activity in working adults: cross-sectional results from the Commuting and Health in Cambridge study. *Prev Med*. 2012;55(5):453–7.
50. Henriques-Neto D, Peralta M, Garradas S, Pelegriani A, Pinto AA, Sánchez-Miguel PA, et al. Active commuting and physical fitness: a systematic review. *Int J Environ Res Public Health*. 2020;17(8):2721.
51. Grandner MA, Patel NP, Gehrman PR, Xie D, Sha D, Weaver T, et al. Who gets the best sleep? Ethnic and socioeconomic factors related to Sleep complaints. *Sleep Med*. 2010;11(5):470–8.
52. Garde AH, Albertsen K, Persson R, Hansen ÅM, Rugulies R. Bi-directional associations between Psychological Arousal, Cortisol, and Sleep. *Behav Sleep Med*. 2012;10(1):28–40.
53. Akerstedt T, Knutsson A, Westerholm P, Theorell T, Alfredsson L, Kecklund G. Sleep disturbances, work stress and work hours: a cross-sectional study. *J Psychosom Res*. 2002;53(3):741–8.
54. Kecklund G, Akerstedt T. Apprehension of the subsequent working day is associated with a low amount of slow wave sleep. *Biol Psychol*. 2004;66(2):169–76.
55. Winpenny EM, van Sluijs EMF, White M, Klepp KI, Wold B, Lien N. Changes in diet through adolescence and early adulthood: longitudinal trajectories and association with key life transitions. *Int J Behav Nutr Phys Act*. 2018;15(1):86.
56. Fulton SL, McKinley MC, Young IS, Cardwell CR, Woodside JV. The Effect of increasing Fruit and Vegetable Consumption on overall Diet: a systematic review and Meta-analysis. *Crit Rev Food Sci Nutr*. 2016;56(5):802–16.

## Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.