



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/223754/>

Version: Accepted Version

Article:

Kottner, J. and Coleman, S. (2023) The theory and practice of pressure ulcer/injury risk assessment: a critical discussion. *Journal of Wound Care*, 32 (9). pp. 560-569. ISSN: 0969-0700

<https://doi.org/10.12968/jowc.2023.32.9.560>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

The theory and practice of pressure ulcer/injury risk assessment and early detection: a critical discussion

Abstract

Pressure ulcer/injury risk assessment is widely considered as an essential component in clinical practice. It is a complex and broad concept including different approaches such as clinical judgement, using standardized risk assessment instruments, skin assessments or using devices to measure skin or tissues properties. A distinction between pressure ulcer/injury risk assessment and early detection is important. Pressure ulcer/injury risk measures the individual susceptibility to develop a pressure ulcer/injury under a specific exposure (primary prevention) and early detection includes the assessment of early (sub)clinical signs and symptoms to prevent progression and to support healing (secondary prevention). Pressure ulcer/injury risk is measured using prognostic/risk factors or prognostic models. Every risk estimate is a probability statement containing varying degrees of uncertainty. It follows, that every clinical decision based on risk estimates contains uncertainty, too. Indeed, pressure ulcer risk assessment and prevention is a complex intervention, where delivery contains several interacting components. There is a huge body of evidence indicating that risk assessment, outcomes of risk assessment, the selection of preventive interventions, and pressure ulceration are not well connected. Methods for prognostic model development and testing in pressure ulcer/injury risk research must be improved following state-of-the-art methodological standards. Despite these challenges, we do have substantial knowledge about pressure ulcer/injury risk factors that helps to make better clinical decisions. A striking next step in development of pressure ulcer/injury risk prediction might be the combination of clinical and other predictors for more individualized care. Any prognostic test or procedure must lead to better prevention at acceptable costs.

Key words

Pressure injury, pressure ulcer, risk assessment, early diagnosis

INTRODUCTION

Pressure ulcers/injuries (PUs) are localised skin and underlying soft tissue damage, wounds or necrosis due to prolonged pressure, or pressure in combination with shear.¹ They typically occur over bony prominences or due to the prolonged contact with medical devices.² Underlying pathways of PU development include deformation damage leading directly to cell death, ischaemia and reperfusion injury and impaired lymphatic function.²⁻⁴ Like many other health problems and diseases, PUs seem to be as old as mankind itself⁵ and recent systematic reviews indicate high prevalence and incidence across various populations and settings.⁶⁻⁹

Because of the severity of this unwanted condition and the substantial impact on individuals and healthcare systems, PU prevention is critical.² State-of-the-art PU prevention includes risk assessment, skin and tissue assessment, and preventive interventions including but not limited to repositioning and early mobilization, use of special support surfaces, skin care or nutrition.² Because of the importance of effective prevention, various quality and patient safety indicators have been proposed to measure the quality of PU prevention whereby PU incidence is most often used.¹⁰ PU occurrence has also been identified as one core outcome to be measured in clinical PU prevention trials.¹¹ However, despite available clinical practice guidelines^{2 12 13} and high quality evidence summaries,¹⁴⁻¹⁶ there are many areas in PU prevention where the quality of evidence is low, where evidence is missing or difficult to generate leading to ongoing debates and controversies about best clinical practice.^{17 18} One such area is PU risk assessment.

PU risk assessment is a complex and broad concept including different approaches such as clinical judgement, using standardized risk assessment instruments, skin assessments looking at erythema or using devices to measure skin or tissues properties such as temperature or oedema.² The concept of risk describes the probability with which a health outcome will occur.¹⁹ Per definition probabilities range from 0 to 1, but because they are probabilities, they are never 0 or 1. Related to PUs, even the best risk assessment method cannot predict with certainty whether a PU will develop or not. Every risk estimate is a probability statement containing varying degrees of uncertainty. It follows, that every clinical decision based on PU risk estimates (e.g. allocation of special support

surfaces) contains uncertainty, too, which contributes to the ongoing discussion of over- and undersupply of PU preventive measures.

When considering PU prevention and management it is important to make a distinction between PU risk assessment and PU early detection (Table 1). PU risk estimates the individual susceptibility to develop a PU under a specific exposure. The skin and underlying tissues are intact and the diagnosis is a predictive statement. The overall preventive goal is to keep the skin and tissues intact (primary prevention). Early detection includes the assessment of early (sub)clinical signs and symptoms (e.g. oedema, erythema, pain, blood or tissue markers) to prevent progression and to support healing (secondary prevention).

Table 1. Distinction between pressure ulcer/injury risk assessment and ulcer/injury early detection

Both strategies are relevant and there are overlaps, but they follow different measurement principles and have unique challenges. In recent years, PU early detection received increasing attention in PU research²⁰⁻²² and clinical practice^{2,23} and a number of reviews are available.²⁴⁻²⁸ Therefore, the focus of this contribution is a critical discussion of the theory, practice and current challenges of PU risk assessment, and to discuss implications for risk assessment practice and research.

PRESSURE ULCER/INJURY RISK ASSESSMENT

Risk assessment is widely considered as an essential component in clinical practice to identify individuals who are susceptible to PU development. The basic programme theory is that once the individual risk has been identified, tailored individual PU prevention is implemented (Figure 1).²

> Fig 1. Programme theory for pressure ulcer risk assessment <

Although in theory this appears straightforward, in practice risk assessment is a complex intervention, where their delivery contains several interacting components,²⁹ including the assessment itself (and associated challenges regarding the concept of risk and how this is summarised and communicated), the potential outcomes and decisions about care interventions and their implementation set within the

context of complex health care environments that are responsible for the delivery of safe and quality care. There is a huge body of evidence indicating that risk assessment, outcomes of risk assessment, the selection of preventive interventions, and PU incidence are not well connected.³⁰⁻³⁶ However, results of diagnostic or prognostic tests must improve clinical decision making, otherwise any risk measurement makes little sense.³⁷ Establishing evidence linkages between the different steps of the PU prevention (Figure 1) and evidence from adequately designed diagnostic randomized controlled trials are urgently needed.³⁰⁻³⁷ To support this further work is also needed to gain a deeper understanding of ‘what works, how, for whom, in what circumstances and to what extent.’³⁸⁻³⁹ Structured PU risk assessment is a resource to clinicians and its impact on care is dependent on how it is used in practice, which will differ according to context.⁴⁰ This requires an understanding of causality via consideration of context, mechanisms and outcome configurations or realist/programme theories, to clarify how different contexts elicit particular nursing team responses and give rise to different outcomes.⁴¹ This approach is becoming more common in the evaluation of complex interventions⁴² and is considered particularly appropriate for the evaluation of new interventions, to explore how an intervention can be adapted for different contexts and where trials have shown conflicting outcomes, to better understand why these inconsistencies occur.⁴³

Quantifying pressure ulcer/injury risk

In clinical research and practice, risks are measured using prognostic factors, often referred to as risk factors in the PU context. A prognostic factor is any variable that is associated with the probability of a subsequent health outcome among people with a particular health condition.⁴⁴ In the PU field, hundreds of prognostic factors have been proposed and are investigated.²⁻⁴⁵⁻⁴⁶

It is essential to point out that the strength and direction of associations of prognostic factors with subsequent PU development highly depends on the population and setting, because most factors are associated with each other and interact. For example, mobility and other common PU risk factors are associated with health and illness in general. This explains the various reports showing for example associations between PU risk and sarcopenia,⁴⁷ disease severity of patients admitted to intensive care units,⁴⁸ length of stay and outcomes in geriatric rehabilitation or internal medicine,⁴⁹⁻⁵⁰

or coronary artery disease.⁵¹ PU risk may be regarded as a general marker of poor health, severe disease and functional impairments.³¹

Interaction describes a situation, when one prognostic factor modifies the effect of another with regard to the occurrence of an outcome.⁵² In PU research this typically occurs, when the effect of immobility (prognostic factor) on PU development (outcome) differs depending whether the patient is placed on a support surface or not (effect modifier). Therefore, preventive interventions must be taken into account in every prognostic study, otherwise obtained prediction estimates and models are wrong or biased.^{53 54}

Furthermore, the association of prognostic factors with risks of subsequent health outcomes do not automatically imply causality.^{52 55} The majority of known PU risk factors do not cause pressure ulceration, rather increase or decrease the likelihood, but only if direct causal factors (e.g. immobility, poor perfusion) are present.⁵⁶ This conceptual distinction has major consequences for clinical practice and research. For example, only interventions aiming at reducing the magnitude and duration of mechanical loads (including repositioning, mobility promotion, offloading) will have direct and immediate preventive effects.⁵⁷ All other preventive interventions such as improving nutrition, the improvement of a diabetic metabolic state, or managing incontinence are indirect.

Irrespectively from multiple associations and interaction, there is a huge body of evidence supporting the relevance of a number of prognostic factors for PU development including activity and mobility limitations, diabetes mellitus, perfusion and circulation, impaired nutritional status and many more.² Although useful for PU assessment and clinical decision making, there are at least two additional challenges:

(1) Many risk factors describe ‘systemic’ characteristics such as age, diabetes mellitus, or ‘skin maturity’ in children,² but PU development is always a local phenomenon. Most prognostic factors are not skin area specific and the actual PU risk depends on the actual loading condition while sitting, lying, standing or being exposed to medical devices.

(2) There is a huge body of evidence that ‘external’ mechanical loads (e.g. lying on a stiff support surface) lead to highly variable and individual internal local stresses and strains within loaded soft tissues.⁵⁸ For example, the anatomical shapes of the calcaneus or sacrum influence how much the

tissues in close proximity to these bones are deformed leading to different intrinsic PU risks.⁵⁹⁻⁶¹ A specific tolerance to soft tissue deformation damage is also highly likely.⁶² It is currently impossible to determine this highly individual ‘biomechanical risk’^{59 63} externally or using any clinical predictor. Therefore, every currently known prognostic factor contains a large amount of error, because it is impossible to observe the actual degree of tissue deformation under the skin surface.

Standardized pressure ulcer/injury risk assessment instruments and scores

Standardized PU risk assessment instruments, scales or scores play an important role in risk assessment practice for decades.² The first standardized PU risk instrument was developed by Norton et al. in the 1960es.⁶⁴ In fact, this instrument was developed only for research purposes, but “... nurses expressed regret when these forms were removed from the wards on completion of the investigation.”⁶⁴ Since then, these scales entered clinical practice. Today, there are probably more than 100 so-called risk assessment scales and new scales are still constantly being developed and existing instruments ‘validated’ and/or modified.⁶⁵⁻⁶⁹ Their use in clinical practice is well established and is still supported by nurses, who value them as an aide memoir to support clinical judgement and an important mechanism for PU prevention.⁷⁰

PU risk assessment instruments incorporate varying prognostic factors which are traditionally used to create an overall score to estimate PU risk. They are prognostic prediction models. Compared to single prognostic factors, prognostic prediction models estimate the individual probability of a future health outcome (here pressure ulceration) by combining information from multiple prognostic factors from an individual.^{71 72} Criticisms of many risk assessment instruments include inadequate development methods (including lack of statistical modelling) and lack of conceptual framework and limited evidence of target population involvement during development i.e. clinical nurses and patient/carers, all of which have led to inconsistent prognostic/risk factor inclusion raising concern about their content validity.^{54 73}

There is a huge and growing body of evidence about measurement properties of PU risk scores including reliability, sensitivity, specificity, predictive validity, and receiver operating characteristic curves. Evidence summaries indicate that these measurement properties vary widely^{2 13} and

discussions about risk score measurement performance are ongoing. Similar to single prognostic factors, the population and setting substantially affect the discrimination ability to distinguish between individuals developing or not developing the outcome of interest. In general, the more homogenous a population (or study sample is), the less discrimination can be achieved by prediction models⁷¹ also leading to low reliability.⁷⁴ Even if the relative measurement error in terms of reliability is low, the absolute measurement errors in terms of proportions of agreement or limits of agreement are much too high, to support any use of overall risk scores or cut-offs.^{31 69 75} Therefore, similar to nearly every other measurement instrument in general, PU risk assessment scales cannot be ‘validated’ in terms of their predictive validity.⁷⁶ Validity or reliability estimates are not fixed properties of a score.⁷⁷ Under specific conditions, obtained risk scores may show certain measurement properties that will be fundamentally different in (slightly) other situations. If authors claim that a prediction model is clinically valuable, it should at least be studied in another independent study in a different physical location.⁷² Newly developed scores should never be tested using the data set with which the score was developed.⁷⁷ Overall, methods for prognostic model development and testing in PU risk research seem to be flawed and do not follow methodological standards.⁵⁴ Further limitations are that studies often focus on assessing the predictive validity of only one instrument, rather than multiple models in the same study population for comparison.⁷⁸

It is important to emphasize, that PU risk scores are causal indicators⁷⁹ or formative models (Figure 2).⁸⁰ The individual prognostic factors of the model determine the concept of PU risk; they are not effects or consequences of PU risk.⁸¹ The individual factors may or may not be associated with each other but the concept of internal consistency does not apply. Therefore, although widely done,^{82 83} it is inappropriate and meaningless to calculate or to interpret Cronbach’s alpha or similar measures in the context of PU risk scores.

> Fig 2. The conceptual structure of pressure ulcer/injury prediction models <

Another major limitation of PU risk scores is that they consist of a fixed number of prognostic factors. Not every factor of the used score is relevant for everyone and there might be other factors not captured by the score relevant for the individual. The International Clinical Practice Guideline explicitly says, that a risk assessment tool “does not replace a comprehensive assessment conducted by an appropriately qualified health professional.”²

Instead of using overall risk scores, a practical criticism of traditional risk assessment instruments, it has been proposed to focus on single items of these scores to guide clinical decision making.^{67 84} This approach seems to be questionable too, because often the range of risk factors included is limited/variable and far from a comprehensive assessment.²

Finally, evidence about measurement properties is different from clinical effects or effectiveness. Even if single prognostic factors or prognostic models show good measurement properties, evidence about the clinical effectiveness is needed to answer the final question of clinical relevance regarding patient outcomes.^{32 85}

Alternative approaches to measure pressure ulcer/injury risk

To overcome the many shortcomings of prognostic factors and models, there is a strong interest in alternative strategies. One is real-time pressure mapping.² It is a widely applied method, to visualize the amount of pressure at the skin support surface interface indicating areas of high compression. This information can be immediately used to reposition and/or off-load subjects or skin areas to reduce local pressure.⁸⁶ However, while there is a clear relationship between interface pressure, position, support surface and individual characteristics,⁸⁷⁻⁸⁹ the association with internal soft tissue deformation and PU development seems to be weaker.^{90 91} Therefore, possible interface pressure thresholds and durations are also unknown⁹² and it is unclear how the interface pressure readings inform clinical decisions. Similar to standardised risk assessment scales, evidence linking pressure mapping with improved patient outcomes is weak.^{2 93}

While there are a many parameters including skin and underlying soft tissue properties,^{22 23} inflammatory local^{28 94} or systemic blood markers²¹ associated with (early) pressure ulceration, it is

much more complicated to find similar biomarkers measuring intrinsic PU risk when skin and tissues are intact. Systemic blood biomarkers including albumin have been proposed proposed,^{2 95} but it is unlikely that any single chemical, biological or biophysical parameter or reading will have better predictive validity than any of the known clinical prognostic factors, because they face the same limitations as described above.

A more promising way may be measuring the individual resistance against mechanical loading. Compared to other biological systems, the functional capacity is much more meaningful to characterise the likely response of skin and soft tissues compared to snapshot or ‘basal’ measurements.⁹⁶⁻⁹⁸ These tests involve artificial controlled stress, stimulus or irritation applied to the skin or tissue and are widely used in skin research.⁹⁹⁻¹⁰⁵ Based on the reaction, conclusions can be drawn about the response, capacity and likely resistance. One popular approach in PU risk research is to measure blood flow responses to characterise microvascular function. For example in healthy individuals, moderate compression of the skin leads to vasodilation, which is called pressure induced vasodilation and considered to be a compensatory response. Evidence suggests, that the absence of this physiological response is associated with skin fragility and increased PU risk.^{106 107} Recently, artificially induced reactive and heat hyperaemia responses were compared between patients with and without PUs¹⁰⁸ and other ‘microvascular stress tests’ might be possible.

The advantages of these functional parameters are that they capture highly individual characteristics leading to individual risk quantification and individualized prevention. On the other hand, evidence suggests, that there is a high degree of biological variability in skin responses after mechanical loading and irritation.^{94 105 109} Another open question is to what extent the measurement of specific functional markers is meaningful with regard to the overall complex multifactorial PU development process.²⁻⁴

WHAT DOES THIS ALL MEAN FOR CLINICAL PRACTICE?

Optimal methods of PU risk assessment are unknown. Currently, a two step-approach is considered best practice (Figure 3)^{2 73 110}: (1) As soon as possible after admission to care services or institutions, a

screening should be conducted. Although the concept of screening may have various meanings in healthcare, it mainly describes examinations or testing procedures in healthy people to detect ‘something putatively prognostic’¹¹¹ to initiate prompt interventions. Regarding PU risk, screening includes a (fast) standardized assessment of *major risk factors* (such as mobility limitations) that are relevant for the population and setting.² (2) If a PU risk is likely, a comprehensive PU risk assessment needs to be conducted. This includes a detailed evaluation of major and condition specific risk factors.²

> Fig 3. Process of pressure ulcer/injury risk assessment <

Summaries of major and specific prognostic factors can be found in systematic reviews,^{45 46} clinical guidelines² or diagnostic manuals such as the NANDA International.¹¹² Setting and population specific clinical algorithms¹¹³⁻¹¹⁵ or risk assessment pathways⁷³ seem to be helpful to support this process. It is the responsibility of the organisation and management to select factors for screening and comprehensive assessment and to develop and implement a risk assessment pathway that fits to the respective service, setting and supports clinical decision making.^{2 110 116}

There are two practical consequences of this two-step approach; one is that those who are clearly not at risk can be quickly screened out, without the need for a more time consuming comprehensive assessment (allowing more appropriate use of nurses time) and the other is that the screening step is not needed at all in high risk settings. For example in intensive care, in the operating room or in spinal cord injury patients it not helpful to perform regular screenings, because a PU risk can be assumed for everyone.

It should be kept in mind that the communication, perception and understanding of risks in healthcare is challenging in general. Evidence indicates that there seem to be particular differences between healthcare professionals and care receivers who are at PU risk.¹¹⁷ This needs to be taken into account for care planning and implementation.

WHAT DOES THIS ALL MEAN FOR RESEARCH?

After decades of PU risk research it is important to adhere to highest methodological standards for identifying, developing and testing prognostic factors and models in the future. Compared to clinical effectiveness research, diagnostic and prognostic research also follows a stepwise approach.¹¹⁸ (Table 2).

Table 2. Phases of clinical prognostic pressure ulcer/injury risk research

PU risk research should begin with the development of a theoretical model (phase 1). Various sources of evidence might be used including exploratory empirical studies. The next basic question is, whether subjects with PUs differ regarding candidate factors from subjects without PUs (phase 2a). Cross-sectional or case-control studies might be used and are widely applied in this early stage. Next questions should be, whether subjects with certain test results have or will develop PUs. This should be followed by questions, whether the test results in the relevant population and setting discriminate between those who will develop a PU and those who will not. The only acceptable designs to describe this possible prediction are longitudinal (phase 2b). If prognostic factors are known already, phase 2b studies should be done immediately. Establishing any (additional) cross-sectional association is not helpful. These phase 2b studies should also look at measurement properties.^{77 80} Finally, evidence is needed to show, if and how the testing procedure leads to better outcomes (phase 3) at acceptable costs (phase 4).

We firmly believe that the described phases should be followed subsequently. Depending on the prognostic factor and the underlying evidence it may be possible to combine phases 2a and b, but it makes no sense to look for example a health economic evaluation (phase 4) without showing the superiority of one prognostic procedure compare to another (phase 3). Any prognostic factor or model should be applied in clinical practice only, if all of the questions have been answered satisfactorily using appropriate designs^{55 72} and reporting.^{77 119} High certainty evidence needs to be established between prognostic factors and PU development.¹²⁰

Especially phase 3 studies are challenging, because it may be difficult to connect prognostic factors or models directly with PU development (Figure 1).³² These studies must develop/evaluate clearly defined diagnostic pathways or decision algorithms. The results of the prognostic factor or model must inform PU prevention practice.³⁷

There is a huge body of evidence about prognostic factors for PU development. Therefore, there might not be a need for finding new predictors, but to develop better models following state-of-the-art methods.⁵⁴ Neither clinical predictors nor biomarkers alone will substantially significantly improve PU prediction compared to what we already have today. Most likely, it will come to a combination of different clinical, biological, or biophysical predictors in the future.⁹⁴

CONCLUSIONS

PU risk assessment was, is, and will remain a critical first step in PU prevention. After decades of PU risk research we must acknowledge the underlying uncertainty of each prognostic factor or model and that an ‘objective’ or best assessment does not exist. Despite this uncertainty, we do have substantial knowledge about PU risk factors that helps to make better clinical decisions that support PU prevention. The challenge is, how this existing knowledge is translated locally. A striking next step in development of PU risk prediction might be the combination of clinical and other (e.g., tissue markers) predictors for more individualized care and how these can be implemented in healthcare organisations to improve processes of care and outcomes. Any prognostic test or procedure must lead to better prevention at acceptable costs.

References

1. World Health Organization. EH90 Pressure ulceration 2023 [Available from: <https://icd.who.int/browse11/l-m/en#/http://id.who.int/icd/entity/455330172> accessed 7 March 2023].
2. European Pressure Ulcer Advisory Panel, National Pressure Injury Advisory Panel, Pan Pacific Pressure Injury Alliance. Prevention and Treatment of Pressure Ulcers/Injuries: Clinical Practice Guideline.2019.

3. Oomens CW, Bader DL, Loerakker S, et al. Pressure induced deep tissue injury explained. *Ann Biomed Eng* 2015;43(2):297-305. doi: 10.1007/s10439-014-1202-6 [published Online First: 2014/12/07]
4. Gefen A, Brienza DM, Cuddigan J, et al. Our contemporary understanding of the aetiology of pressure ulcers/pressure injuries. *Int Wound J* 2022;19(3):692-704. doi: 10.1111/iwj.13667 [published Online First: 2021/08/13]
5. Rowling JT. Pathological changes in mummies. *Proc R Soc Med* 1961;54(5):409-15. [published Online First: 1961/05/01]
6. Labeau SO, Afonso E, Benbenishty J, et al. Prevalence, associated factors and outcomes of pressure injuries in adult intensive care unit patients: the DecubICUs study. *Intensive Care Med* 2021;47(2):160-69. doi: 10.1007/s00134-020-06234-9 [published Online First: 2020/10/10]
7. Li Z, Lin F, Thalib L, et al. Global prevalence and incidence of pressure injuries in hospitalised adult patients: A systematic review and meta-analysis. *Int J Nurs Stud* 2020;105:103546. doi: 10.1016/j.ijnurstu.2020.103546 [published Online First: 2020/03/01]
8. Triantafyllou C, Chorianopoulou E, Kourkouni E, et al. Prevalence, incidence, length of stay and cost of healthcare-acquired pressure ulcers in pediatric populations: A systematic review and meta-analysis. *Int J Nurs Stud* 2021;115:103843. doi: 10.1016/j.ijnurstu.2020.103843 [published Online First: 2020/12/30]
9. Sardo PMG, Teixeira JPF, Machado A, et al. A systematic review of prevalence and incidence of pressure ulcers/injuries in hospital emergency services. *J Tissue Viability* 2023;32(2):179-87. doi: 10.1016/j.jtv.2023.02.001 [published Online First: 2023/02/03]
10. Kottner J, Hahnel E, Lichterfeld-Kottner A, et al. Measuring the quality of pressure ulcer prevention: A systematic mapping review of quality indicators. *Int Wound J* 2018;15(2):218-24. doi: 10.1111/iwj.12854 [published Online First: 2017/11/28]
11. Lechner A, Coleman S, Balzer K, et al. Core outcomes for pressure ulcer prevention trials: results of an international consensus study: Classification: Outcomes and qualitative research. *Br J Dermatol* 2022;187(5):743-52. doi: 10.1111/bjd.21741 [published Online First: 2022/07/06]
12. Kottner J, El Genedy-Kalyoncu M. The uptake of the international pressure ulcer/injury prevention and treatment guidelines in the scientific literature: A systematic analysis of two major citation databases. *J Tissue Viability* 2022;31(4):763-67. doi: 10.1016/j.jtv.2022.07.011 [published Online First: 2022/08/13]
13. Qaseem A, Mir TP, Starkey M, et al. Risk assessment and prevention of pressure ulcers: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2015;162(5):359-69. doi: 10.7326/M14-1567 [published Online First: 2015/03/04]
14. Greenwood C, Nelson EA, Nixon J, et al. Comparative effectiveness of heel-specific medical devices for the prevention of heel pressure ulcers: A systematic review. *J Tissue Viability* 2022;31(4):579-92. doi: 10.1016/j.jtv.2022.09.009 [published Online First: 2022/10/23]
15. Avsar P, Moore Z, Patton D, et al. Repositioning for preventing pressure ulcers: a systematic review and meta-analysis. *J Wound Care* 2020;29(9):496-508. doi: 10.12968/jowc.2020.29.9.496 [published Online First: 2020/09/15]
16. Shi C, Dumville JC, Cullum N, et al. Beds, overlays and mattresses for preventing and treating pressure ulcers: an overview of Cochrane Reviews and network meta-analysis. *Cochrane Database Syst Rev* 2021;8(8):CD013761. doi: 10.1002/14651858.CD013761.pub2 [published Online First: 2021/08/17]
17. Black J. Pressure ulcer prevention and management: a dire need for good science. *Ann Intern Med* 2015;162(5):387-8. doi: 10.7326/M15-0190 [published Online First: 2015/03/04]
18. Haesler E, Kottner J, Cuddigan J, et al. The 2014 International Pressure Ulcer Guideline: methods and development. *J Adv Nurs* 2017;73(6):1515-30. doi: 10.1111/jan.13241 [published Online First: 2016/12/18]
19. Higgins JPT LT, Deeks JJ. Chapter 6: Choosing effect measures and computing estimates of effect. In: Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, ed. *Cochrane Handbook for Systematic Reviews of Interventions* version 63 (updated February 2022): Cochrane 2022.

20. Kimura N, Nakagami G, Minematsu T, et al. Non-invasive detection of local tissue responses to predict pressure ulcer development in mouse models. *J Tissue Viability* 2020;29(1):51-57. doi: 10.1016/j.jtv.2019.11.001 [published Online First: 2019/11/24]
21. Traa WA, Strijkers GJ, Bader DL, et al. Myoglobin and troponin concentrations are increased in early stage deep tissue injury. *J Mech Behav Biomed Mater* 2019;92:50-57. doi: 10.1016/j.jmbbm.2018.12.026 [published Online First: 2019/01/19]
22. Volzer B, Kottner J. Associations between skin structural and functional changes after loading in healthy aged females at sacral and heel skin: A secondary data analysis. *J Tissue Viability* 2022;31(2):239-44. doi: 10.1016/j.jtv.2022.01.010 [published Online First: 2022/02/08]
23. Martins de Oliveira AL, O'Connor T, Patton D, et al. Sub-epidermal moisture versus traditional and visual skin assessments to assess pressure ulcer risk in surgery patients. *J Wound Care* 2022;31(3):254-64. doi: 10.12968/jowc.2022.31.3.254 [published Online First: 2022/02/25]
24. Chaboyer W, Coyer F, Harbeck E, et al. Oedema as a predictor of the incidence of new pressure injuries in adults in any care setting: A systematic review and meta-analysis. *Int J Nurs Stud* 2022;128:104189. doi: 10.1016/j.ijnurstu.2022.104189 [published Online First: 2022/02/27]
25. Shi C, Bonnett LJ, Dumville JC, et al. Nonblanchable erythema for predicting pressure ulcer development: a systematic review with an individual participant data meta-analysis. *Br J Dermatol* 2020;182(2):278-86. doi: 10.1111/bjd.18154 [published Online First: 2019/05/24]
26. Wang N, Lv L, Yan F, et al. Biomarkers for the early detection of pressure injury: A systematic review and meta-analysis. *J Tissue Viability* 2022;31(2):259-67. doi: 10.1016/j.jtv.2022.02.005 [published Online First: 2022/03/02]
27. Wilson H, Moore Z, Avsar P, et al. Exploring the Role of Pain as an Early Indicator for Individuals at Risk of Pressure Ulcer Development: A Systematic Review. *Worldviews Evid Based Nurs* 2021;18(4):299-307. doi: 10.1111/wvn.12528 [published Online First: 2021/07/25]
28. Bader DL, Worsley PR. Technologies to monitor the health of loaded skin tissues. *Biomed Eng Online* 2018;17(1):40. doi: 10.1186/s12938-018-0470-z [published Online First: 2018/04/14]
29. Skivington K, Matthews L, Simpson SA, et al. A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance. *BMJ* 2021;374:n2061. doi: 10.1136/bmj.n2061 [published Online First: 2021/10/02]
30. Lovegrove J, Ven S, Miles SJ, et al. Comparison of pressure injury risk assessment outcomes using a structured assessment tool versus clinical judgement: A systematic review. *J Clin Nurs* 2021. doi: 10.1111/jocn.16154 [published Online First: 2021/12/03]
31. Kottner J, Balzer K. Do pressure ulcer risk assessment scales improve clinical practice? *Journal of Multidisciplinary Healthcare* 2010;3:103-11.
32. Moore ZE, Patton D. Risk assessment tools for the prevention of pressure ulcers. *Cochrane Database Syst Rev* 2019;1(1):CD006471. doi: 10.1002/14651858.CD006471.pub4 [published Online First: 2019/02/01]
33. Fossum M, Alexander GL, Goransson KE, et al. Registered nurses' thinking strategies on malnutrition and pressure ulcers in nursing homes: a scenario-based think-aloud study. *J Clin Nurs* 2011;20(17-18):2425-35. doi: 10.1111/j.1365-2702.2010.03578.x [published Online First: 2011/06/22]
34. Van Herck P, Sermeus W, Jylha V, et al. Using hospital administrative data to evaluate the knowledge-to-action gap in pressure ulcer preventive care. *J Eval Clin Pract* 2009;15(2):375-82. doi: 10.1111/j.1365-2753.2008.01019.x [published Online First: 2009/04/02]
35. Skytt B, Engstrom M, Martensson G, et al. A longitudinal qualitative study of health care personnel's perceptions of simultaneous implementation of three risk assessment scales on falls, malnutrition and pressure ulcers. *J Clin Nurs* 2016;25(13-14):1912-22. doi: 10.1111/jocn.13207 [published Online First: 2016/04/15]
36. Samuriwo R, Dowding D. Nurses' pressure ulcer related judgements and decisions in clinical practice: a systematic review. *Int J Nurs Stud* 2014;51(12):1667-85. doi: 10.1016/j.ijnurstu.2014.04.009 [published Online First: 2014/05/20]

37. Balzer K, Kopke S, Luhmann D, et al. Designing trials for pressure ulcer risk assessment research: methodological challenges. *Int J Nurs Stud* 2013;50(8):1136-50. doi: 10.1016/j.ijnurstu.2013.02.008 [published Online First: 2013/04/02]
38. Pawson R, Tilley N. Realist Evaluation. In: Office BC, ed. http://www.communitymatters.com.au/RE_chapter.pdf, 2004.
39. Wong G, Westhorp G, Manzano A, et al. RAMESES II reporting standards for realist evaluations. *Bmc Medicine* 2016;14 doi: ARTN 96
10.1186/s12916-016-0643-1
40. Greenhalgh T, Wong G, Jagosh J, et al. Protocol—the RAMESES II study: developing guidance and reporting standards for realist evaluation. *BMJ Open* 2015;5(8) doi: 10.1136/bmjopen-2015-008567
41. Greenhalgh T, Humphrey C, Hughes J, et al. How Do You Modernize a Health Service ? A Realist Evaluation of Whole-Scale Transformation in London. *Milbank Q* 2009;87(2):391-416. doi: 10.1111/j.1468-0009.2009.00562.x
42. Moore GF, Audrey S, Barker M, et al. Process evaluation of complex interventions: Medical Research Council guidance. *BMJ* 2015;350 doi: 10.1136/bmj.h1258
43. Richards DA, Hallberg IR. Complex Interventions in Health: An Overview of Research Methods. London: Routledge Taylor and Francis Group 2015.
44. Riley RD, Moons KGM, Snell KIE, et al. A guide to systematic review and meta-analysis of prognostic factor studies. *BMJ* 2019;364:k4597. doi: 10.1136/bmj.k4597 [published Online First: 2019/02/01]
45. Coleman S, Gorecki C, Nelson EA, et al. Patient risk factors for pressure ulcer development: systematic review. *Int J Nurs Stud* 2013;50(7):974-1003. doi: 10.1016/j.ijnurstu.2012.11.019 [published Online First: 2013/02/05]
46. Alderden J, Rondinelli J, Pepper G, et al. Risk factors for pressure injuries among critical care patients: A systematic review. *Int J Nurs Stud* 2017;71:97-114. doi: 10.1016/j.ijnurstu.2017.03.012 [published Online First: 2017/04/07]
47. Cha YH, Song SY, Park KS, et al. Relationship between pressure ulcer risk and sarcopenia in patients with hip fractures. *J Wound Care* 2022;31(6):532-36. doi: 10.12968/jowc.2022.31.6.532 [published Online First: 2022/06/10]
48. Tang W, Zha ML, Zhang WQ, et al. APACHE scoring system and pressure injury risk for intensive care patients: A systematic review and meta-analysis. *Wound Repair Regen* 2022;30(4):498-508. doi: 10.1111/wrr.13021 [published Online First: 2022/05/20]
49. Justo D, Vislapu N, Shvedov V, et al. Admission Norton scale scores (ANSS) correlate with rehabilitation outcome and length in elderly patients following hip arthroplasty. *Arch Gerontol Geriatr* 2011;53(1):e33-6. doi: 10.1016/j.archger.2010.10.008 [published Online First: 2010/11/27]
50. Leshem-Rubinow E, Vaknin A, Sherman S, et al. Norton scale, hospitalization length, complications, and mortality in elderly patients admitted to internal medicine departments. *Gerontology* 2013;59(6):507-13. doi: 10.1159/000353710 [published Online First: 2013/08/08]
51. Wang Y, Chen R, Ding J, et al. Predictive value of pressure ulcer risk for obstructive coronary artery disease. *Nurs Open* 2021;8(4):1848-55. doi: 10.1002/nop2.835 [published Online First: 2021/03/07]
52. Szklo M, Nieto FJ. Epidemiology Beyond the Basics. 4th ed. Burlington: Jones & Bartlett Learning 2019.
53. Defloor T, Grypdonck MF. Validation of pressure ulcer risk assessment scales: a critique. *J Adv Nurs* 2004;48(6):613-21. doi: 10.1111/j.1365-2648.2004.03250.x [published Online First: 2004/11/19]
54. Shi C, Dumville JC, Cullum N. Evaluating the development and validation of empirically-derived prognostic models for pressure ulcer risk assessment: A systematic review. *Int J Nurs Stud* 2019;89:88-103. doi: 10.1016/j.ijnurstu.2018.08.005 [published Online First: 2018/10/24]

55. Riley RD, Hayden JA, Steyerberg EW, et al. Prognosis Research Strategy (PROGRESS) 2: prognostic factor research. *PLoS Med* 2013;10(2):e1001380. doi: 10.1371/journal.pmed.1001380 [published Online First: 2013/02/09]
56. Coleman S, Nixon J, Keen J, et al. A new pressure ulcer conceptual framework. *J Adv Nurs* 2014;70(10):2222-34. doi: 10.1111/jan.12405 [published Online First: 2014/04/02]
57. Lechner A, Kottner J, Coleman S, et al. Outcomes for Pressure Ulcer Trials (OUTPUTs) project: review and classification of outcomes reported in pressure ulcer prevention research. *Br J Dermatol* 2021;184(4):617-26. doi: 10.1111/bjd.19304 [published Online First: 2020/06/09]
58. Abou Karam M, Mukhina E, Daras N, et al. Reliability of B-mode ultrasound and shear wave elastography in evaluating sacral bone and soft tissue characteristics in young adults with clinical feasibility in elderly. *J Tissue Viability* 2022;31(2):245-54. doi: 10.1016/j.jtv.2022.02.003 [published Online First: 2022/03/04]
59. Sprigle S, Sonenblum S. Visualizing Tissue Strain Under the Sacrum and Coccyx in Different Supine Postures: A Case Series. *Adv Skin Wound Care* 2019;32(6):264-71. doi: 10.1097/01.ASW.0000554445.59743.44 [published Online First: 2019/04/09]
60. Sonenblum SE, Sprigle SH, Cathcart JM, et al. 3D anatomy and deformation of the seated buttocks. *J Tissue Viability* 2015;24(2):51-61. doi: 10.1016/j.jtv.2015.03.003 [published Online First: 2015/05/04]
61. Gefen A. The biomechanics of heel ulcers. *J Tissue Viability* 2010;19(4):124-31. doi: 10.1016/j.jtv.2010.06.003 [published Online First: 2010/06/30]
62. Traa WA, van Turnhout MC, Nelissen JL, et al. There is an individual tolerance to mechanical loading in compression induced deep tissue injury. *Clin Biomech (Bristol, Avon)* 2019;63:153-60. doi: 10.1016/j.clinbiomech.2019.02.015 [published Online First: 2019/03/22]
63. Sonenblum SE, Sprigle SH. Buttock tissue response to loading in men with spinal cord injury. *PLoS One* 2018;13(2):e0191868. doi: 10.1371/journal.pone.0191868 [published Online First: 2018/02/08]
64. Norton D, McLaren R, Exton-Smith AN. An Investigation of Geriatric Nursing Problems in Hospital. London: Churchill Livingstone 1975.
65. Heikkilä A, Kotila J, Junttila K. Validation of the Helsinki University Hospital prevent pressure Injury Risk Assessment Tool: a prospective observational study. *BMC Nurs* 2022;21(1):18. doi: 10.1186/s12912-021-00799-6 [published Online First: 2022/01/19]
66. Shi HY, Chen HL. Optimized Norton Scale for Pressure Injury Risk Assessment: Need for Additional Predictive Validity Analysis. *J Wound Ostomy Continence Nurs* 2021;48(2):174-75. doi: 10.1097/WON.0000000000000743 [published Online First: 2021/03/11]
67. Wahlin I, Ek AC, Lindgren M, et al. Development and validation of an ICU-specific pressure injury risk assessment scale. *Scand J Caring Sci* 2021;35(3):769-78. doi: 10.1111/scs.12891 [published Online First: 2020/07/16]
68. Higgins J, Casey S, Taylor E, et al. Comparing the Braden and Jackson/Cubbin Pressure Injury Risk Scales in Trauma-Surgery ICU Patients. *Crit Care Nurse* 2020;40(6):52-61. doi: 10.4037/ccn2020874 [published Online First: 2020/12/02]
69. Kohta M, Ohura T, Tsukada K, et al. Inter-Rater Reliability of a Pressure Injury Risk Assessment Scale for Home Care: A Multicenter Cross-Sectional Study. *J Multidiscip Healthc* 2020;13:2031-41. doi: 10.2147/JMDH.S291162 [published Online First: 2020/12/31]
70. Coleman S, Greenhalgh J, Schoonhoven L, et al. Use of PURPOSE-T in Clinical Practice. Annual Meeting of the European Pressure Ulcer Advisory Panel. Prague, Czech Republic, 2022.
71. Debray TP, Damen JA, Snell KI, et al. A guide to systematic review and meta-analysis of prediction model performance. *BMJ* 2017;356:i6460. doi: 10.1136/bmj.i6460 [published Online First: 2017/01/07]
72. Steyerberg EW, Moons KG, van der Windt DA, et al. Prognosis Research Strategy (PROGRESS) 3: prognostic model research. *PLoS Med* 2013;10(2):e1001381. doi: 10.1371/journal.pmed.1001381 [published Online First: 2013/02/09]

73. Coleman S, Smith IL, McGinnis E, et al. Clinical evaluation of a new pressure ulcer risk assessment instrument, the Pressure Ulcer Risk Primary or Secondary Evaluation Tool (PURPOSE T). *J Adv Nurs* 2018;74(2):407-24. doi: 10.1111/jan.13444 [published Online First: 2017/08/24]
74. Kottner J, Audige L, Brorson S, et al. Guidelines for Reporting Reliability and Agreement Studies (GRRAS) were proposed. *Int J Nurs Stud* 2011;48(6):661-71. doi: 10.1016/j.ijnurstu.2011.01.016 [published Online First: 2011/04/26]
75. Kottner J, Dassen T. An interrater reliability study of the Braden scale in two nursing homes. *Int J Nurs Stud* 2008;45(10):1501-11. doi: 10.1016/j.ijnurstu.2008.02.007 [published Online First: 2008/04/09]
76. American Educational Research Association, American Psychological Association, National Council on Measurement in Education. Standards for Educational and Psychological Testing. Washington 2014.
77. Streiner DL, Kottner J. Recommendations for reporting the results of studies of instrument and scale development and testing. *J Adv Nurs* 2014;70(9):1970-79. doi: 10.1111/jan.12402 [published Online First: 2014/04/02]
78. Ferrante di Ruffano L, Hyde CJ, McCaffery KJ, et al. Assessing the value of diagnostic tests: a framework for designing and evaluating trials. *BMJ* 2012;344:e686. doi: 10.1136/bmj.e686 [published Online First: 2012/02/23]
79. Bollen K, Lennox R. Conventional Wisdom on Measurement - a Structural Equation Perspective. *Psychol Bull* 1991;110(2):305-14. doi: 10.1037/0033-2909.110.2.305
80. De Vet H, Terwee C, Mokkink L, et al. Measurement in Medicine: Cambridge University Press 2011.
81. Kottner J, Streiner DL. Internal consistency and Cronbach's alpha: A comment on Beeckman et al. (2010). *Int J Nurs Stud* 2010;47(7):926-8. doi: 10.1016/j.ijnurstu.2009.12.018 [published Online First: 2010/03/24]
82. Puspitasari JD, Nurhaeni N, Waluyanti FT. Testing of Braden QD Scale for predicting pressure ulcer risk in the Pediatric Intensive Care Unit. *Pediatr Rep* 2020;12(Suppl 1):8694. doi: 10.4081/pr.2020.8694 [published Online First: 2020/09/10]
83. Lovegrove J, Fulbrook P, Miles S. International consensus on pressure injury preventative interventions by risk level for critically ill patients: A modified Delphi study. *Int Wound J* 2020;17(5):1112-27. doi: 10.1111/iwj.13461 [published Online First: 2021/02/17]
84. Magnan MA, Maklebust J. The nursing process and pressure ulcer prevention: making the connection. *Adv Skin Wound Care* 2009;22(2):83-92; quiz 93-4. doi: 10.1097/01.ASW.0000345279.13156.d0 [published Online First: 2009/01/22]
85. Vach W, Bibiza E, Gerke O, et al. A potential for seamless designs in diagnostic research could be identified. *J Clin Epidemiol* 2021;129:51-59. doi: 10.1016/j.jclinepi.2020.09.019 [published Online First: 2020/09/30]
86. Scott RG, Thurman KM. Visual Feedback of Continuous Bedside Pressure Mapping to Optimize Effective Patient Repositioning. *Adv Wound Care (New Rochelle)* 2014;3(5):376-82. doi: 10.1089/wound.2013.0519 [published Online First: 2014/05/08]
87. Urasaki M, Nakagami G, Sanada H, et al. Interface pressure distribution of elderly Japanese people in the sitting position. *Disabil Rehabil Assist Technol* 2011;6(1):38-46. doi: 10.3109/17483107.2010.512968 [published Online First: 2010/09/08]
88. Kawabata T, Sugama J. Relationship between mattress internal air pressure and interface pressure distribution in the lateral position. *Int Wound J* 2022;19(8):2115-23. doi: 10.1111/iwj.13817 [published Online First: 2022/04/28]
89. Grap MJ, Munro CL, Wetzel PA, et al. Backrest Elevation and Tissue Interface Pressure by Anatomical Location During Mechanical Ventilation. *Am J Crit Care* 2016;25(3):e56-63. doi: 10.4037/ajcc2016317 [published Online First: 2016/05/03]
90. Kawasaki S, Nishimura Y, Kamijo YI, et al. Relationship between ultrasonographically low-echoic lesions under the skin, wheelchair sitting time, and interface pressure on ischial region in

- individuals with chronic spinal cord injury. *J Spinal Cord Med* 2021;44(6):978-84. doi: 10.1080/10790268.2020.1746873 [published Online First: 2020/05/23]
91. Reenalda J, Jannink M, Nederhand M, et al. Clinical use of interface pressure to predict pressure ulcer development: a systematic review. *Assist Technol* 2009;21(2):76-85. doi: 10.1080/10400430903050437 [published Online First: 2009/09/01]
 92. Sving EBM, Gunningberg LAC, Baath CB, et al. Using pressure mapping intraoperatively to prevent pressure ulcers-A quasi-experimental study. *Health Sci Rep* 2023;6(2):e1112. doi: 10.1002/hsr2.1112 [published Online First: 2023/02/24]
 93. Gunningberg L, Sedin IM, Andersson S, et al. Pressure mapping to prevent pressure ulcers in a hospital setting: A pragmatic randomised controlled trial. *Int J Nurs Stud* 2017;72:53-59. doi: 10.1016/j.ijnurstu.2017.04.007 [published Online First: 2017/05/02]
 94. Jayabal H, Abiakam NS, Filingeri D, et al. Inflammatory biomarkers in sebum for identifying skin damage in patients with a Stage I pressure ulcer in the pelvic region: A single centre observational, longitudinal cohort study with elderly patients. *Int Wound J* 2023 doi: 10.1111/iwj.14131 [published Online First: 2023/03/07]
 95. Bogie KM, Schwartz K, Li Y, et al. Exploring adipogenic and myogenic circulatory biomarkers of recurrent pressure injury risk for persons with spinal cord injury. *J Circ Biomark* 2020;9:1-7. doi: 10.33393/jcb.2020.2121 [published Online First: 2021/02/19]
 96. Blume-Peytavi U, Kottner J, Sterry W, et al. Age-Associated Skin Conditions and Diseases: Current Perspectives and Future Options. *Gerontologist* 2016;56 Suppl 2:S230-42. doi: 10.1093/geront/gnw003 [published Online First: 2016/03/20]
 97. Kottner J, Beeckman D, Vogt A, et al. Skin health and integrity. In: Gefen A, ed. *Innovations and Emerging Technologies in Wound Care*. London: Elsevier 2020:183-96.
 98. Ghadially R. Aging and the epidermal permeability barrier: implications for contact dermatitis. *Am J Contact Dermat* 1998;9(3):162-9. [published Online First: 1998/09/24]
 99. El Genedy-Kalyoncu M, Richter C, Surber C, et al. The effect of a basic skin care product on the structural strength of the dermo-epidermal junction: An exploratory, randomised, controlled split-body trial. *Int Wound J* 2022;19(2):426-35. doi: 10.1111/iwj.13643 [published Online First: 2021/06/15]
 100. Fluhr JW, Kuss O, Diepgen T, et al. Testing for irritation with a multifactorial approach: comparison of eight non-invasive measuring techniques on five different irritation types. *Br J Dermatol* 2001;145(5):696-703. doi: 10.1046/j.1365-2133.2001.04431.x [published Online First: 2001/12/12]
 101. Loffler H, Aramaki J, Effendy I. Response to thermal stimuli in skin pretreated with sodium lauryl sulfate. *Acta Derm Venereol* 2001;81(6):395-7. doi: 10.1080/000155501317208309 [published Online First: 2002/02/28]
 102. Rosado C, Pinto P, Rodrigues LM. Assessment of moisturizers and barrier function restoration using dynamic methods. *Skin Res Technol* 2009;15(1):77-83. doi: 10.1111/j.1600-0846.2008.00331.x [published Online First: 2009/01/21]
 103. Roure R, Lanctin M, Nollent V, et al. Methods to Assess the Protective Efficacy of Emollients against Climatic and Chemical Aggressors. *Dermatol Res Pract* 2012;2012:864734. doi: 10.1155/2012/864734 [published Online First: 2012/09/07]
 104. Kanto H, Washizaki K, Ito M, et al. Optimal patch application time in the evaluation of skin irritation. *J Dermatol* 2013;40(5):363-9. doi: 10.1111/1346-8138.12004 [published Online First: 2013/02/19]
 105. Jayabal H, Bates-Jensen BM, Abiakam NS, et al. The identification of biophysical parameters which reflect skin status following mechanical and chemical insults. *Clin Physiol Funct Imaging* 2021;41(4):366-75. doi: 10.1111/cpf.12707 [published Online First: 2021/05/03]
 106. Zwanenburg PR, Backer SFM, Obdeijn MC, et al. A Systematic Review and Meta-Analysis of the Pressure-Induced Vasodilation Phenomenon and Its Role in the Pathophysiology of Ulcers. *Plast Reconstr Surg* 2019;144(4):669e-81e. doi: 10.1097/PRS.00000000000006090 [published Online First: 2019/10/01]

107. Vouillarmet J, Josset-Lamaugarny A, Michon P, et al. Neurovascular Response to Pressure in Patients With Diabetic Foot Ulcer. *Diabetes* 2019;68(4):832-36. doi: 10.2337/db18-0694 [published Online First: 2019/01/27]
108. Tzen YT, Champagne PT, Rao V, et al. Utilisation of skin blood flow as a precursor for pressure injury development in persons with acute spinal cord injury: A proof of concept. *Int Wound J* 2022;19(8):2191-99. doi: 10.1111/iwj.13829 [published Online First: 2022/05/12]
109. Lechner A, Rancan F, Hadam S, et al. Comparing the effects of three different multilayer dressings for pressure ulcer prevention on sacral skin after prolonged loading: An exploratory crossover trial. *Wound Repair Regen* 2021;29(2):270-79. doi: 10.1111/wrr.12883 [published Online First: 2020/12/22]
110. Deutsches Netzwerk für Qualitätsentwicklung in der Pflege. Expertenstandard Dekubitusprophylaxe in der Pflege, 2. Aktualisierung 2017.
111. Speechley M, Kunnilathu A, Aluckal E, et al. Screening in Public Health and Clinical Care: Similarities and Differences in Definitions, Types, and Aims - A Systematic Review. *J Clin Diagn Res* 2017;11(3):LE01-LE04. doi: 10.7860/JCDR/2017/24811.9419 [published Online First: 2017/05/18]
112. NANDA International. Nursing Diagnoses Definitions and Classifications 2021-2023. New York: Thieme 2021.
113. National Institute for Health and Care Excellence. Algorithm for risk assessment, prevention and management in children 2014.
114. National Institute for Health and Care Excellence. Algorithm for risk assessment, prevention and management in adults 2014.
115. van Rijswijk L, Beitz JM. Creating a pressure ulcer prevention algorithm: systematic review and face validation. *Ostomy Wound Manage* 2013;59(11):28-40. [published Online First: 2013/11/10]
116. Healthcare Improvement Scotland. Prevention and Management of Pressure Ulcers, 2020.
117. Ledger LJ, Worsley PR, Hope J, et al. Patient perceptions and understanding of pressure ulcer risk in the community: Empirical Research Qualitative. *J Adv Nurs* 2023 doi: 10.1111/jan.15637 [published Online First: 2023/03/16]
118. Haynes RB, You JJ. The architecture of diagnostic research. In: Knottnerus JA, Buntix F, eds. The Evidence Base of Clinical Diagnosis. 2nd ed. Oxford: Wiley-Blackwell 2009.
119. Collins GS, Reitsma JB, Altman DG, et al. Transparent Reporting of a multivariable prediction model for Individual Prognosis Or Diagnosis (TRIPOD). *Ann Intern Med* 2015;162(10):735-6. doi: 10.7326/L15-5093-2 [published Online First: 2015/05/20]
120. Foroutan F, Guyatt G, Zuk V, et al. GRADE Guidelines 28: Use of GRADE for the assessment of evidence about prognostic factors: rating certainty in identification of groups of patients with different absolute risks. *J Clin Epidemiol* 2020;121:62-70. doi: 10.1016/j.jclinepi.2019.12.023 [published Online First: 2020/01/27]