

This is a repository copy of *Kernel operators with variable intervals of integration in Lebesgue spaces and applications*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/47020/>

Version: Published Version

---

**Article:**

Stepanov, Vladimir D and Ushakova, Elena P (2010) Kernel operators with variable intervals of integration in Lebesgue spaces and applications. *Mathematical Inequalities Applications*. 449–510.

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# KERNEL OPERATORS WITH VARIABLE INTERVALS OF INTEGRATION IN LEBESGUE SPACES AND APPLICATIONS

VLADIMIR D. STEPANOV AND ELENA P. USHAKOVA

*Dedicated to Professor Josip Pečarić  
on the occasion of his 60th birthday*

*(Communicated by L.-E. Persson)*

**Abstract.** New criteria of  $L_p - L_q$  boundedness of Hardy-Steklov type operator (1.1) with both increasing on  $(0, \infty)$  boundary functions  $a(x)$  and  $b(x)$  are obtained for  $1 < p \leq q < \infty$  and  $0 < q < p < \infty$ ,  $p > 1$ . This result is applied for two-weighted  $L_p - L_q$  characterization of the corresponding geometric Steklov operator (1.3) and other related problems.

## 1. Introduction

Let  $0 < p < \infty$ ,  $\|f\|_p := (\int_0^\infty |f(x)|^p dx)^{1/p}$  and  $L_p$  denotes the Lebesgue space of all measurable functions on  $\mathbb{R}^+ := [0, \infty)$  such that  $\|f\|_p < \infty$ .

Assume  $w(x)$  and  $v(y)$  be locally integrable and almost everywhere positive functions (weights). We study the  $L_p - L_q$  boundedness of the Hardy-Steklov operator of the form

$$\mathcal{H}f(x) := w(x) \int_{a(x)}^{b(x)} f(y)v(y)dy, \quad (1.1)$$

where the boundaries  $a(x)$  and  $b(x)$  satisfy the following conditions:

- (i)  $a(x)$  and  $b(x)$  are differentiable and strictly increasing on  $(0, \infty)$ ;
- (ii)  $a(0) = b(0) = 0$ ,  $a(x) < b(x)$  for  $0 < x < \infty$ ,  $a(\infty) = b(\infty) = \infty$ .

In the limiting cases  $a(x) = 0$  or  $b(x) = \infty$  the operator  $\mathcal{H}$  is reduced to the Hardy-type operators with variable upper or lower bound and this relation stands behind of the so-called *block-diagonal method* for investigation of  $\mathcal{H}$  by a suitable decomposition into a sequence of Hardy-type operators with non-overlapping domains.

*Mathematics subject classification* (2010): Primary 26D10; Secondary 26D15, 26D07.

*Keywords and phrases:* Integral operators, Lebesgue spaces, weights, boundedness.

The Swedish Institute is the grant-giving authority for research of the second author (Project 00105/2007 Visby Programme 382). The work was also partially supported by the Russian Foundation for Basic Research (Projects 09-01-00093, 09-01-98516 and 07-01-00054) and by the Far-Eastern Branch of the Russian Academy of Sciences (Project 09-II-CO-01-003).

However, apart from the limiting cases some properties of  $\mathcal{H}$  could be rather different. For instance,  $\mathcal{H}$  is a self-adjoint operator in  $L_2$  for  $w = v$  and  $a(x) = b^{-1}(x)$  and to find the spectrum of  $\mathcal{H}$  in this case is an interesting problem.

We give two alternative pairs of criteria for the  $L_p - L_q$  boundedness of  $\mathcal{H}$  (§ 4). The first pair is a complete analog of Tomaselli-Muckenhoupt-Bradley and Mazya-Rozin conditions for the Hardy-type operators in the cases  $1 < p \leq q < \infty$  and  $0 < q < p < \infty$ ,  $p > 1$ , respectively. The second pair is new and allows to characterize the weighted  $L_p - L_q$  boundedness of the geometric Steklov operator (§ 5)

$$\mathcal{G}f(x) := \exp\left(\frac{1}{b(x) - a(x)} \int_{a(x)}^{b(x)} \log f(y) dy\right), \quad f(y) \geq 0. \quad (1.3)$$

Both pairs of criteria involve a notion of the *fairway* - a curve between the graphs of  $a(x)$  and  $b(x)$  with such an equilibrium property which allows to squeeze up discrete portions produced by the block-diagonal method into one piece.

We demonstrate the block-diagonal method (§ 3) for an even more general operator

$$\mathcal{K}f(x) := w(x) \int_{a(x)}^{b(x)} k(x, y) f(y) v(y) dy, \quad (1.4)$$

where the kernel  $k(x, y) \geq 0$  satisfies the *Oinarov-type condition* of the form

$$k(x, y) \approx k(x, b(z)) + k(z, y), \quad z \leq x, \quad a(x) \leq y \leq b(z). \quad (1.5)$$

The same method works for a formally dual operator

$$\mathcal{K}^*f(x) := w(x) \int_{a(x)}^{b(x)} k(y, x) f(y) v(y) dy \quad (1.6)$$

with the kernel  $k(y, x) \geq 0$  satisfying

$$k(y, x) \approx k(y, z) + k(a(z), x), \quad x \leq z, \quad a(z) \leq y \leq b(x).$$

However, in both cases two-sided estimates of the norms have discrete forms, which are rather inconvenient for further applications. When  $p \leq q$  the forms can be refined up to "continuous" ones, but with a double supremum. As for the case  $q < p$  is concerned the attempts to find out the integral form of criteria, analogous to the Hardy-type case, met some difficulties. Nevertheless, it gives a solution for the  $L_p - L_q$  boundedness of  $\mathcal{H}$  narrowed on the cone of monotone functions (§ 6.2).

Our next observation is that  $L_p - L_q$  boundedness of  $\mathcal{H}$  is equivalent to the validity of the differential inequality

$$\|Fw\|_q \leq C \|F'/v\|_p \quad (1.7)$$

restricted to a non-linear class of absolutely continuous function defined by the border functions  $a(x)$  and  $b(x)$ , which in turn is closely related to the embedding of the weighted Sobolev space to the weighted Lebesgue space to hold (§ 6.1), that is to inequalities of the form

$$\|Fw\|_q \leq C (\|Fu\|_s + \|F'/v\|_p). \quad (1.8)$$

We start the paper with the Preliminaries (§ 2) contained auxiliary results and conclude by the Bibliographical remarks.

Throughout of the paper products of the form  $0 \cdot \infty$  are taken to be equal to 0. Relations  $A \ll B$  mean  $A \leq cB$  with some constants  $c$  depending only on parameters of summations and, possibly, on the constants of equivalence in the inequalities of the type (1.5). We write  $A \approx B$  instead of  $A \ll B \ll A$  or  $A = cB$ .  $\mathbb{Z}$  and  $\mathbb{N}$  denote the sets of all integers and all positive integers, respectively.  $\chi_E$  stands for a characteristic function (indicator) of a subset  $E \subset \mathbb{R}^+$ . Also we make use of marks  $: =$  and  $=:$  for introducing new quantities and denote  $p' := p/(p-1)$  for  $0 < p < \infty$ ,  $p \neq 1$  and  $r := pq/(p-q)$  for  $0 < q < p < \infty$ . In Section 5 we denote  $L_{p,v}$  the weighted Lebesgue space with the norm  $\|f\|_{p,v} := \|fv\|_p$ .

## 2. Preliminaries

### 2.1. Hardy and Hardy type operators

Here we collect some known results for Hardy operator

$$Hf(x) := w(x) \int_c^x f(y)v(y)dy, \quad 0 \leq c \leq x \leq d \leq \infty, \quad (2.1)$$

and Hardy type operator of the form

$$Kf(x) := w(x) \int_c^x k(x,y)f(y)v(y)dy, \quad 0 \leq c \leq x \leq d \leq \infty, \quad (2.2)$$

with a non-negative kernel  $k(x,y)$  from Oinarov's class  $\mathcal{O}$ .

DEFINITION 2.1. Let  $k(x,y) \geq 0$ ,  $k(x,y) \in \mathcal{O}$  if there exists a constant  $D \geq 1$  such that

$$D^{-1}k(x,y) \leq k(x,z) + k(z,y) \leq Dk(x,y), \quad 0 \leq c \leq y \leq z \leq x \leq d \leq \infty. \quad (2.3)$$

THEOREM 2.1. Let the operator  $H : L_p(c,d) \rightarrow L_q(c,d)$  be defined by (2.1).

(a) If  $1 < p \leq q < \infty$ , then  $\|H\|_{L_p(c,d) \rightarrow L_q(c,d)} \approx A_M \approx A_T$ , where

$$A_M := \sup_{c \leq t \leq d} \left( \int_t^d w^q(x)dx \right)^{\frac{1}{q}} \left( \int_c^t v^{p'}(y)dy \right)^{\frac{1}{p'}}, \quad (2.4)$$

$$A_T := \sup_{c \leq t \leq d} \left( \int_c^t \left[ \int_c^x v^{p'}(y)dy \right]^q w^q(x)dx \right)^{\frac{1}{q}} \left( \int_c^t v^{p'}(y)dy \right)^{-\frac{1}{p}}. \quad (2.5)$$

(b) Let  $0 < q < p < \infty$ ,  $p > 1$ . Then  $\|H\|_{L_p(c,d) \rightarrow L_q(c,d)} \approx B_{MR} \approx B_{PS}$ , where

$$B_{MR} := \left( \int_c^d \left[ \int_t^d w^q(x)dx \right]^{\frac{r}{p}} \left[ \int_c^t v^{p'}(y)dy \right]^{\frac{r}{p'}} w^q(t)dt \right)^{\frac{1}{r}}, \quad (2.6)$$

$$B_{PS} := \left( \int_c^d \left[ \int_c^t \left\{ \int_c^x v^{p'}(y) dy \right\}^q w^q(x) dx \right]^{\frac{t}{p}} \left[ \int_c^t v^{p'}(y) dy \right]^{q-\frac{t}{p}} w^q(t) dt \right)^{\frac{1}{r}}. \quad (2.7)$$

REMARK 2.1. Since  $\|H\|_{L_p(c,d) \rightarrow L_q(c,d)} = \|H^*\|_{L_{q'}(c,d) \rightarrow L_{p'}(c,d)}$  for  $1 < p, q < \infty$ , where

$$H^*g(y) := v(y) \int_y^d g(x)w(x)dx, \quad 0 \leq c \leq y \leq d \leq \infty, \quad (2.8)$$

the above equivalences ought to be supplemented by

$$A_T^* := \sup_{c \leq t \leq d} \left( \int_t^d \left[ \int_y^d w^q(x) dx \right]^{p'} v^{p'}(y) dy \right)^{\frac{1}{p'}} \left( \int_t^d w^q(x) dx \right)^{-\frac{1}{q'}}, \quad (2.9)$$

$$B_{PS}^* := \left( \int_c^d \left[ \int_t^d \left\{ \int_y^d w^q(x) dx \right\}^{p'} v^{p'}(y) dy \right]^{\frac{t}{q'}} \left[ \int_t^d w^q(x) dx \right]^{p'-\frac{t}{q'}} v^{p'}(t) dt \right)^{\frac{1}{r}}. \quad (2.10)$$

For the Hardy type operator we have the following.

THEOREM 2.2. Let the operator  $K : L_p(c, d) \rightarrow L_q(c, d)$  be defined by (2.2) with  $k(x, y) \in \mathcal{O}$ .

(a) If  $1 < p \leq q < \infty$ , then  $\|K\|_{L_p(c,d) \rightarrow L_q(c,d)} \approx A_0 + A_1$ , where

$$A_0 := \sup_{c \leq t \leq d} \left( \int_t^d k^q(x, t) w^q(x) dx \right)^{\frac{1}{q}} \left( \int_c^t v^{p'}(y) dy \right)^{\frac{1}{p'}}, \quad (2.11)$$

$$A_1 := \sup_{c \leq t \leq d} \left( \int_t^d w^q(x) dx \right)^{\frac{1}{q}} \left( \int_c^t k^{p'}(t, y) v^{p'}(y) dy \right)^{\frac{1}{p'}}. \quad (2.12)$$

(b) Let  $1 < q < p < \infty$ . Then  $\|K\|_{L_p(c,d) \rightarrow L_q(c,d)} \approx B_0 + B_1$ , where

$$B_0 := \left( \int_c^d \left[ \int_t^d k^q(x, t) w^q(x) dx \right]^{\frac{t}{q}} \left[ \int_c^t v^{p'}(y) dy \right]^{\frac{t}{q'}} v^{p'}(t) dt \right)^{\frac{1}{r}}, \quad (2.13)$$

$$B_1 := \left( \int_c^d \left[ \int_t^d w^q(x) dx \right]^{\frac{t}{p}} \left[ \int_c^t k^{p'}(t, y) v^{p'}(y) dy \right]^{\frac{t}{p'}} w^q(t) dt \right)^{\frac{1}{r}}. \quad (2.14)$$

Let  $b : [c, d] \rightarrow [0, \infty)$  be a strictly increasing differentiable function and let  $K_b : L_p(b(c), b(d)) \rightarrow L_q(c, d)$  be an operator of the form

$$K_b f(x) := w(x) \int_{b(c)}^{b(x)} k(x, y) f(y) v(y) dy, \quad 0 \leq c \leq x \leq d \leq \infty, \quad (2.15)$$

where a non-negative kernel  $k(x, y)$  satisfies the following definition.

DEFINITION 2.2.  $k(x, y) \in \mathcal{O}_b$  if there exists a constant  $D \geq 1$  such that

$$D^{-1}k(x, y) \leq k(x, b(z)) + k(z, y) \leq Dk(x, y), \quad \begin{cases} 0 \leq c \leq z \leq x \leq d \leq \infty, \\ 0 \leq b(c) \leq y \leq b(z). \end{cases} \quad (2.16)$$

COROLLARY 2.1. Let the operator  $K_b$  be an operator given by (2.15) with a strictly increasing differentiable function  $b(x) \geq 0$  and  $k(x, y) \in \mathcal{O}_b$ .

(a) If  $1 < p \leq q < \infty$ , then

$$\|K_b\|_{L_p(b(c), b(d)) \rightarrow L_q(c, d)} \approx A_{b,0} + A_{b,1}, \quad (2.17)$$

where

$$A_{b,0} := \sup_{c \leq t \leq d} \left( \int_t^d k^q(x, b(t)) w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{b(c)}^{b(t)} v^{p'}(y) dy \right)^{\frac{1}{p'}}, \quad (2.18)$$

$$A_{b,1} := \sup_{c \leq t \leq d} \left( \int_t^d w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{b(c)}^{b(t)} k^{p'}(t, y) v^{p'}(y) dy \right)^{\frac{1}{p'}}. \quad (2.19)$$

(b) If  $1 < q < p < \infty$ , then

$$\|K_b\|_{L_p(b(c), b(d)) \rightarrow L_q(c, d)} \approx B_{b,0} + B_{b,1}, \quad (2.20)$$

where

$$B_{b,0} := \left( \int_{b(c)}^{b(d)} \left[ \int_{b^{-1}(t)}^d k^q(x, t) w^q(x) dx \right]^{\frac{t}{q}} \left[ \int_{b(c)}^t v^{p'}(y) dy \right]^{\frac{t}{q'}} v^{p'}(t) dt \right)^{\frac{1}{r}}, \quad (2.21)$$

$$B_{b,1} := \left( \int_c^d \left[ \int_t^d w^q(x) dx \right]^{\frac{t}{p}} \left[ \int_{b(c)}^{b(t)} k^{p'}(t, y) v^{p'}(y) dy \right]^{\frac{t}{p'}} w^q(t) dt \right)^{\frac{1}{r}}. \quad (2.22)$$

*Proof.* By the substitution  $\tau = b^{-1}(y)$  we see that the inequality

$$\left( \int_c^d (K_b f)^q(x) dx \right)^{\frac{1}{q}} \leq C \left( \int_{b(c)}^{b(d)} f^p(y) dy \right)^{\frac{1}{p}} \quad (2.23)$$

is equivalent to the inequality

$$\left( \int_c^d \left[ \int_c^x \tilde{k}(x, \tau) f(\tau) \tilde{v}(\tau) d\tau \right]^q w^q(x) dx \right)^{\frac{1}{q}} \leq C \left( \int_c^d f^p(\tau) d\tau \right)^{\frac{1}{p}} \quad (2.24)$$

with  $\tilde{v}(\tau) = v(b(\tau)) [b'(\tau)]^{1/p'}$ ,  $\tilde{k}(x, \tau) = k(x, b(\tau)) \in \mathcal{O}$ . The result follows from Theorem 2.2.

REMARK 2.1. If  $k(x, y) \equiv 1$  the result of Corollary 2.1(b) is true for  $0 < q < p < \infty$ ,  $p > 1$ .

Similar characterization is valid for the operator  $K_a : L_p(a(c), a(d)) \rightarrow L_q(c, d)$  with a lower variable limit of integration of the form

$$K_a f(x) := w(x) \int_{a(x)}^{a(d)} k(y, x) f(y) v(y) dy, \quad 0 \leq c \leq x \leq d \leq \infty, \quad (2.25)$$

with a non-negative strictly increasing differentiable function  $a(x)$  and a non-negative kernel  $k(y, x)$  from Oinarov's type class  $\mathcal{O}_a$  defined as follows.

DEFINITION 2.3.  $k(y, x) \in \mathcal{O}_a$ , if there exists a constant  $D \geq 1$  such that

$$D^{-1}k(y, x) \leq k(y, z) + k(a(z), x) \leq Dk(y, x), \quad \begin{cases} 0 \leq c \leq x \leq z \leq d \leq \infty, \\ 0 \leq a(z) \leq y \leq a(d). \end{cases} \quad (2.26)$$

COROLLARY 2.2. (a) If  $1 < p \leq q < \infty$ , then

$$\|K_a\|_{L_p(a(c), a(d)) \rightarrow L_q(c, d)} \approx A_{a,0} + A_{a,1}, \quad (2.27)$$

where

$$A_{a,0} := \sup_{c \leq t \leq d} \left( \int_c^t k^q(a(t), x) w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{a(d)} v^{p'}(y) dy \right)^{\frac{1}{p'}}, \quad (2.28)$$

$$A_{a,1} := \sup_{c \leq t \leq d} \left( \int_c^t w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{a(d)} k^{p'}(y, t) v^{p'}(y) dy \right)^{\frac{1}{p'}}. \quad (2.29)$$

(b) If  $1 < q < p < \infty$ , then

$$\|K_a\|_{L_p(a(c), a(d)) \rightarrow L_q(c, d)} \approx B_{a,0} + B_{a,1}, \quad (2.30)$$

where


$$B_{a,0} := \left( \int_{a(c)}^{a(d)} \left[ \int_c^{a^{-1}(t)} k^q(t, x) w^q(x) dx \right]^{\frac{r}{q}} \left[ \int_t^{a(d)} v^{p'}(y) dy \right]^{\frac{r}{q'}} v^{p'}(t) dt \right)^{\frac{1}{r}}, \quad (2.31)$$

$$B_{a,1} := \left( \int_c^d \left[ \int_c^t w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{a(t)}^{a(d)} k^{p'}(y, t) v^{p'}(y) dy \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}}. \quad (2.32)$$

REMARK 2.2. In the case  $k(y, x) \equiv 1$  the part (b) holds for  $0 < q < p < \infty$ ,  $p > 1$ .

In lemmas 2.1–2.4 we state norm estimates for certain Hardy type operators with only one variable limit. Such estimates are given above in a less general form (see corollaries 2.1–2.2 with  $k \equiv 1$ ), but the authors find it difficult to give precise references. Therefore, we state these results here together with sketches of the proof for some of them. We start from the case  $1 < p \leq q < \infty$ .

LEMMA 2.1. *Let  $1 < p \leq q < \infty$  and let  $0 \leq c < d \leq \infty$ ,  $0 \leq a < \infty$ . Suppose that the function  $b(x)$  is differentiable, strictly increasing and such that  $a \leq b(x) < \infty$ ,  $x \in [c, d]$ , and let*



$$Sf(x) := w(x) \int_a^{b(x)} f(y)v(y)dy. \quad (2.33)$$

*Then for the norm of  $S$  we have the following two-sided estimates with coefficients of equivalence depending on  $p$  and  $q$  only:*

$$\|S\|_{L_p(a,b(d)) \rightarrow L_q(c,d)} \approx A_b := \sup_{c \leq t \leq d} \left( \int_t^d w^q(x)dx \right)^{\frac{1}{q}} \left( \int_a^{b(t)} v^{p'}(y)dy \right)^{\frac{1}{p'}}, \quad (2.34)$$

$$\begin{aligned} \|S\|_{L_p(a,b(d)) \rightarrow L_q(c,d)} &\approx \mathbb{A}_b \\ &:= \sup_{c \leq t \leq d} \left( \int_c^t \left[ \int_a^{b(x)} v^{p'}(y)dy \right]^q w^q(x)dx \right)^{\frac{1}{q}} \left( \int_a^{b(t)} v^{p'}(y)dy \right)^{-\frac{1}{p}}. \end{aligned} \quad (2.35)$$

*Proof.* Necessity for both (2.34) and (2.35) follows from applying the test function  $f_t(y) = [v(y)]^{p'-1} \chi_{[a,b(t)]}(y)$  to the inequality

$$\left( \int_c^d (Sf)^q(x)dx \right)^{\frac{1}{q}} \leq C \left( \int_a^{b(d)} f^p(y)dy \right)^{\frac{1}{p}}. \quad (2.36)$$

To prove sufficiency in (2.34) we note that the least possible constant  $C \geq 0$  of the inequality (2.36) coinciding with the operator norm  $C = \|S\|_{L_p(a,b(d)) \rightarrow L_q(c,d)}$  is equivalent to the sum of the least possible constants  $C \approx C_1 + C_2$  of the inequalities

$$\begin{aligned} \int_a^{b(c)} f(y)v(y)dy \left( \int_c^d w^q(x)dx \right)^{\frac{1}{q}} &\leq C_1 \left( \int_a^{b(d)} f^p(y)dy \right)^{\frac{1}{p}}, \\ \left( \int_c^d \left[ \int_{b(c)}^{b(x)} f(y)v(y)dy \right]^q w^q(x)dx \right)^{\frac{1}{q}} &\leq C_2 \left( \int_a^{b(d)} f^p(y)dy \right)^{\frac{1}{p}}, \end{aligned}$$

which, in their turn, are equivalent to the following two inequalities:

$$\int_a^{b(c)} f(y)v(y)dy \left( \int_c^d w^q(x)dx \right)^{\frac{1}{q}} \leq C_3 \left( \int_a^{b(c)} f^p(y)dy \right)^{\frac{1}{p}}, \quad (2.37)$$



$$\left( \int_c^d \left[ \int_{b(c)}^{b(x)} f(y)v(y)dy \right]^q w^q(x)dx \right)^{\frac{1}{q}} \leq C_4 \left( \int_{b(c)}^{b(d)} f^p(y)dy \right)^{\frac{1}{p}}, \quad (2.38)$$

respectively, therefore,  $C \approx C_3 + C_4$ . From (2.37) by the reverse Hölder inequality we find

$$C_3 = \left( \int_c^d w^q(x)dx \right)^{\frac{1}{q}} \left( \int_a^{b(c)} v^{p'}(y)dy \right)^{\frac{1}{p'}} \leq A_b$$

and for (2.38) by Corollary 2.1 (a) for  $k(x,y) = 1$  we have

$$C_4 \approx \sup_{c \leq t \leq d} \left( \int_t^d w^q(x)dx \right)^{\frac{1}{q}} \left( \int_{b(c)}^{b(t)} v^{p'}(y)dy \right)^{\frac{1}{p'}} \leq A_b$$

and the sufficiency of (2.34) follows.

For an upper estimate in (2.35) we write the inequality

$$I^{\frac{1}{p'}} := \left( \int_a^{b(d)} \left[ \int_{b_0^{-1}(y)}^d g(x)w(x)dx \right]^{p'} dV(y) \right)^{\frac{1}{p'}} \leq C \left( \int_c^d g^{q'}(x)dx \right)^{\frac{1}{q}},$$

dual to (2.36), where  $V(y) := \int_a^y v^{p'}(z)dz$  and  $b_0^{-1}(y) = \max \{c, b^{-1}(y)\}$ . Integrating by parts and applying Hölder's inequality we find

$$\begin{aligned} I &= p' \int_{b(c)}^{b(d)} \left( \int_{b^{-1}(y)}^d g(x)w(x)dx \right)^{p'-1} V(y)g(b^{-1}(y))w(b^{-1}(y))db^{-1}(y) \\ &\leq p' \left( \int_{b(c)}^{b(d)} [g(b^{-1}(y))]^{q'} db^{-1}(y) \right)^{\frac{1}{q'}} \\ &\quad \times \left( \int_{b(c)}^{b(d)} \left[ \int_{b^{-1}(y)}^d g(x)w(x)dx \right]^{\frac{q}{p-1}} V^q(y)w^q(b^{-1}(y))db^{-1}(y) \right)^{\frac{1}{q}} \\ &=: p' \left( \int_c^d g^{q'}(x)dx \right)^{\frac{1}{q'}} I_1^{\frac{1}{q}}. \end{aligned}$$

Now

$$\begin{aligned} I_1 &= \int_{b(c)}^{b(d)} \left\{ \int_y^{b(d)} d \left[ - \left( \int_{b^{-1}(t)}^d g(x)w(x)dx \right)^{\frac{q}{p-1}} \right] \right\} V^q(y)w^q(b^{-1}(y))db^{-1}(y) \\ &= \int_{b(c)}^{b(d)} \left( \int_{b(c)}^t V^q(y)w^q(b^{-1}(y))db^{-1}(y) \right) d \left[ - \left( \int_{b^{-1}(t)}^d g(x)w(x)dx \right)^{\frac{q}{p-1}} \right] \\ &= \int_{b(c)}^{b(d)} \left( \int_c^{b^{-1}(t)} [V(b(x))]^q w^q(x)dx \right) d \left[ - \left( \int_{b^{-1}(t)}^d g(x)w(x)dx \right)^{\frac{q}{p-1}} \right] \\ &= \int_c^d \left( \int_c^s [V(b(x))]^q w^q(x)dx \right) d \left[ - \left( \int_s^d g(x)w(x)dx \right)^{\frac{q}{p-1}} \right] \end{aligned}$$

$$\leq \mathbb{A}_b^q \int_c^d \left( \int_a^{b(s)} v^{p'}(y) dy \right)^{\frac{q}{p}} d \left[ - \left( \int_s^d g(x) w(x) dx \right)^{\frac{q}{p-1}} \right].$$

Furthermore, by Minkowskii's inequality

$$\begin{aligned} & \int_c^d \left( \int_a^{b(s)} v^{p'}(y) dy \right)^{\frac{q}{p}} d \left[ - \left( \int_s^d g(x) w(x) dx \right)^{\frac{q}{p-1}} \right] \\ & \ll \left( \int_c^d g(x) w(x) dx \right)^{\frac{q}{p-1}} \left( \int_a^{b(c)} v^{p'}(y) dy \right)^{\frac{q}{p}} \\ & \quad + \left( \int_{b(c)}^{b(d)} \left\{ \int_{b^{-1}(y)}^d d \left[ - \left( \int_s^d g(x) w(x) dx \right)^{\frac{q}{p-1}} \right] \right\}^{\frac{p}{q}} dV(y) \right)^{\frac{q}{p}} \\ & = \left( \int_c^d g(x) w(x) dx \right)^{\frac{q}{p-1}} \left( \int_a^{b(c)} v^{p'}(y) dy \right)^{\frac{q}{p}} \\ & \quad + \left( \int_{b(c)}^{b(d)} \left[ \int_{b^{-1}(y)}^d g(x) w(x) dx \right]^{p'} dV(y) \right)^{\frac{q}{p}} \approx I_p^{\frac{q}{p}}. \end{aligned}$$


Thus,

$$I \ll \mathbb{A}_b \left( \int_c^d g^{q'}(x) dx \right)^{\frac{1}{q'}} I^{\frac{1}{p}}$$

and the proof of the upper bound  $C \ll \mathbb{A}_b$  is completed.

For the Hardy type operator with a lower variable limit of integration the similar norm estimates can be obtained analogously.

LEMMA 2.2. *Let  $1 < p \leq q < \infty$  and let  $0 \leq c < d \leq \infty$ ,  $0 < b \leq \infty$ . Suppose that the function  $a(x)$  is differentiable, strictly increasing and such that  $0 < a(x) \leq b$ ,  $x \in (c, d]$ , and*



$$Tf(x) := w(x) \int_{a(x)}^b f(y) v(y) dy. \quad (2.39)$$

Then

$$\|T\|_{L_p(a(c), b) \rightarrow L_q(c, d)} \approx A_a := \sup_{c \leq t \leq d} \left( \int_c^t w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^b v^{p'}(y) dy \right)^{\frac{1}{p'}}, \quad (2.40)$$

$$\|T\|_{L_p(a(c), b) \rightarrow L_q(c, d)} \approx \mathbb{A}_a := \sup_{c \leq t \leq d} \left( \int_t^d \left[ \int_{a(x)}^b v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^b v^{p'}(y) dy \right)^{-\frac{1}{p}}. \quad (2.41)$$

Next two lemmas are concerned to the case  $0 < q < p < \infty$ ,  $p > 1$ .

LEMMA 2.3. *Let  $0 < q < p < \infty$ ,  $p > 1$ ,  $1/r = 1/q - 1/p$  and let  $0 \leq c < d \leq \infty$ ,  $0 \leq a < \infty$ . Suppose that the function  $b(x)$  is differentiable, strictly increasing and such that  $a \leq b(x) < \infty$ ,  $x \in [c, d)$ , and let the operator  $S$  be defined by (2.33). Then*

$$\|S\|_{L_p(a, b(d)) \rightarrow L_q(c, d)}^r \approx B_b^r := \int_c^d \left( \int_t^d w^q(x) dx \right)^{\frac{r}{p}} \left( \int_a^{b(t)} v^{p'}(y) dy \right)^{\frac{r}{p'}} w^q(t) dt \quad (2.42)$$

$$\begin{aligned} &\approx \left( \int_c^d w^q(x) dx \right)^{\frac{r}{q}} \left( \int_a^{b(c)} v^{p'}(y) dy \right)^{\frac{r}{p'}} \\ &\quad + \int_c^d \left( \int_t^d w^q(x) dx \right)^{\frac{r}{q}} d \left( \int_a^{b(t)} v^{p'}(y) dy \right)^{\frac{r}{p'}} =: \widehat{B}_b^r, \end{aligned} \quad (2.43)$$

$$\begin{aligned} \|S\|_{L_p(a, b(d)) \rightarrow L_q(c, d)}^r &\approx \mathbb{B}_b^r := \int_c^d \left( \int_c^t \left[ \int_a^{b(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{p}} \left( \int_a^{b(t)} v^{p'}(y) dy \right)^{q - \frac{r}{p}} w^q(t) dt \end{aligned} \quad (2.44)$$

$$\begin{aligned} &\approx \left( \int_c^d \left[ \int_a^{b(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{q}} \left( \int_a^{b(d)} v^{p'}(y) dy \right)^{-\frac{r}{p}} \\ &\quad + \int_c^d \left( \int_c^t \left[ \int_a^{b(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{q}} d \left[ - \left( \int_a^{b(t)} v^{p'}(y) dy \right)^{-\frac{r}{p}} \right]. \end{aligned} \quad (2.45)$$

*Proof.* We start as in the proof of Lemma 2.1 and by the reverse Hölder inequality

$$E_0 := \left( \int_c^d w^q(x) dx \right)^{\frac{1}{q}} \left( \int_a^{b(c)} v^{p'}(y) dy \right)^{\frac{1}{p'}} = C_3,$$

and by Corollary 2.1 (b) with taking into account Remark 2.1 we obtain

$$E_1 := \left( \int_c^d \left[ \int_t^d w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{b(c)}^{b(t)} v^{p'}(y) dy \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}} \approx C_4,$$

and we conclude that

$$\begin{aligned} C &\approx E_0 + E_1 \\ &\approx \left( \int_c^d \left\{ \int_t^d w^q(x) dx \right\}^{\frac{r}{p}} \left[ \left( \int_a^{b(c)} v^{p'}(y) dy \right)^{\frac{r}{p'}} + \left( \int_{b(c)}^{b(t)} v^{p'}(y) dy \right)^{\frac{r}{p'}} \right] w^q(t) dt \right)^{\frac{1}{r}} \\ &\approx B_b. \end{aligned}$$

For the proof of the equivalence (2.42) to the sum (2.43) suppose first that  $B_b^r < \infty$ . Then

$$\int_t^d w^q(x) dx < \infty \quad \text{for any } t \in (c, d] \quad (2.46)$$

and, therefore,

$$\begin{aligned} 0 &= \lim_{s \rightarrow d} \int_s^d \left( \int_a^{b(t)} v^{p'}(y) dy \right)^{\frac{r}{p'}} d \left[ - \left( \int_t^d w^q(x) dx \right)^{\frac{r}{q}} \right] \\ &\geq \lim_{s \rightarrow d} \left( \int_a^{b(s)} v^{p'}(y) dy \right)^{\frac{r}{p'}} \int_s^d d \left[ - \left( \int_t^d w^q(x) dx \right)^{\frac{r}{q}} \right] \\ &= \lim_{s \rightarrow d} \left( \int_a^{b(s)} v^{p'}(y) dy \right)^{\frac{r}{p'}} \left( \int_s^d w^q(x) dx \right)^{\frac{r}{q}}. \end{aligned} \quad (2.47)$$

It implies by integration by parts that  $\infty > B_b^r \approx \widehat{B}_b^r$ . In the reverse direction let  $\widehat{B}_b^r =: I_1 + I_2 < \infty$ , then

$$\begin{aligned} \infty > I_1 + I_2 &= I_1 + \left( \int_t^d w^q(x) dx \right)^{\frac{r}{q}} \left( \int_a^{b(t)} v^{p'}(y) dy \right)^{\frac{r}{p'}} \Big|_c^d \\ &\quad + \int_c^d \left( \int_a^{b(t)} v^{p'}(y) dy \right)^{\frac{r}{p'}} d \left[ - \left( \int_t^d w^q(x) dx \right)^{\frac{r}{q}} \right] \\ &\geq \int_c^d \left( \int_a^{b(t)} v^{p'}(y) dy \right)^{\frac{r}{p'}} d \left[ - \left( \int_t^d w^q(x) dx \right)^{\frac{r}{q}} \right] \approx B_b^r. \end{aligned}$$

Hence,  $B_b^r < \infty$ , therefore, by (2.46) – (2.47) we have  $B_b^r \approx \widehat{B}_b^r$ .

To prove the second part of the lemma we put  $V_a(t) := \int_a^{b(t)} v^{p'}(y) dy$  and let

$$\mathbb{B}_{PS}^r := \int_c^d \left( \int_c^t V_a^q(x) w^q(x) dx \right)^{\frac{r}{q}} d \left( -[V_a(t)]^{-r/p} \right).$$

It was proved in [18, Theorem 3] that

$$\mathbb{B}_b \approx \mathbb{B}_{PS}, \quad \text{if } V_a(d) = \infty \quad (2.48)$$

and

$$\mathbb{B}_b^r \approx \left( \int_c^d V_a^q(x) w^q(x) dx \right)^{\frac{r}{q}} [V_a(d)]^{-r/p} + \mathbb{B}_{PS}^r, \quad \text{if } 0 < V_a(d) < \infty. \quad (2.49)$$

For the upper estimate  $C \ll \mathbb{B}_b$  we assume that  $\mathbb{B}_b < \infty$  and suppose first  $V_a(d) = \infty$ . Then by Hölder's inequality with the exponents  $r/q$  and  $p/q$  we have

$$\begin{aligned} J &:= \int_c^d \left( \int_a^{b(x)} f(y) v(y) dy \right)^q w^q(x) dx \\ &= \int_c^d \left( \int_a^{b(x)} f(y) v(y) dy \right)^q w^q(x) V_a^q(x) V_a^{-q}(x) dx \end{aligned}$$

$$\begin{aligned}
&= q \int_c^d \left( \int_a^{b(x)} f(y)v(y)dy \right)^q w^q(x) V_a^q(x) \left( \int_x^d [V_a(s)]^{-q-1} dV_a(s) \right) dx =: J_0 \\
&= q \int_c^d [V_a(s)]^{-q-1} \left[ \int_c^s \left( \int_a^{b(x)} f(y)v(y)dy \right)^q w^q(x) V_a^q(x) dx \right] dV_a(s) \\
&\leq q \int_c^d \left\{ \left( \int_a^{b(s)} f(y)v(y)dy \right)^q V_a^{-q}(s) \right\} \left\{ \left( \int_c^s w^q(x) V_a^q(x) dx \right) V_a^{-1}(s) \right\} dV_a(s) \\
&\ll \left( \int_c^d \left[ \int_a^{b(s)} f(y)v(y)dy \right]^p V_a^{-p}(s) dV_a(s) \right)^{\frac{q}{p}} \mathbb{B}_{PS}^q.
\end{aligned}$$

Hence, by using the estimate (2.34) from Lemma 2.1 we find

$$J \ll \mathbb{B}_{PS}^q \left( \int_a^{b(d)} f^p(y)dy \right)^{\frac{q}{p}}.$$

Therefore, in view of (2.48) the inequality  $C \ll \mathbb{B}_b$  holds. If  $V_a(d) < \infty$ , then

$$V_a^{-q}(x) = V_a^{-q}(d) + q \int_x^d [V_a(s)]^{-q-1} dV_a(s)$$

and analogously we find that

$$J = J_0 + V_a^{-q}(d) \int_c^d \left( \int_a^{b(x)} f(y)v(y)dy \right)^q w^q(x) V_a^q(x) dx =: J_0 + J_1.$$

From the above estimate for  $J_0$  and (2.49) it follows that

$$J_0 \ll \mathbb{B}_b^q \left( \int_a^{b(d)} f^p(y)dy \right)^{\frac{q}{p}}.$$

To estimate  $J_1$  let  $\{x_k\}_{k \in \mathbb{Z}}$  be such a sequence that  $\int_{b(c)}^{b(x_k)} f(y)v(y)dy = 2^k$ ,  $k \leq N$ . Then

$$\begin{aligned}
V_a^q(d)J_1 &= \int_c^d \left[ \int_a^{b(c)} f(y)v(y)dy + \int_{b(c)}^{b(x)} f(y)v(y)dy \right]^q w^q(x) V_a^q(x) dx \\
&\approx \left( \int_a^{b(c)} f(y)v(y)dy \right)^q \int_c^d w^q(x) V_a^q(x) dx \\
&\quad + \sum_{k \leq N} \int_{x_k}^{x_{k+1}} \left( \int_{b(c)}^{b(x)} f(y)v(y)dy \right)^q w^q(x) V_a^q(x) dx =: J_{1,1} + J_{1,2}.
\end{aligned}$$

By applying Hölder's inequality and (2.49) we find that

$$J_{1,1} \leq \left( \int_a^{b(c)} f^p(y)dy \right)^{\frac{q}{p}} [V_a(c)]^{q/p'} \int_c^d w^q(x) V_a^q(x) dx \leq V_a^q(d) \mathbb{B}_b^q \left( \int_a^{b(c)} f^p(y)dy \right)^{\frac{q}{p}}.$$

For the second term  $J_{1,2}$  we have by Hölder's inequality with the exponents  $r/q$  and  $p/q$

$$\begin{aligned}
 J_{1,2} &\leq \sum_{k \leq N} 2^{q(k+1)} \int_{x_k}^{x_{k+1}} w^q(x) V_a^q(x) dx \\
 &\leq 4^q \sum_{k \leq N} \left( \int_{b(x_{k-1})}^{b(x_k)} f(y) v(y) dy \right)^q \int_{x_k}^{x_{k+1}} w^q(x) V_a^q(x) dx \\
 &\leq 4^q \sum_{k \leq N} \left( \int_{b(x_{k-1})}^{b(x_k)} f^p(y) dy \right)^{\frac{q}{p}} \left( \int_{b(x_{k-1})}^{b(x_k)} v^{p'}(y) dy \right)^{\frac{q}{p'}} \int_{x_k}^{x_{k+1}} w^q(x) V_a^q(x) dx \\
 &\ll \left( \int_{b(c)}^{b(d)} f^p(y) dy \right)^{\frac{q}{p}} \left( \sum_{k \leq N} V_a(x_k)^{\frac{r}{p'}} \left( \int_{x_k}^{x_{k+1}} w^q(x) V_a^q(x) dx \right)^{\frac{r}{q}} \right)^{\frac{q}{r}} \\
 &\ll \left( \int_{b(c)}^{b(d)} f^p(y) dy \right)^{\frac{q}{p}} \left( \int_c^d \left( \int_c^t w^q(x) V_a^q(x) dx \right)^{\frac{r}{p}} w^q(t) [V_a(t)]^{q+\frac{r}{p'}} dt \right)^{\frac{q}{r}} \\
 &\ll V_a^q(d) \mathbb{B}_b^q \left( \int_{b(c)}^{b(d)} f^p(y) dy \right)^{\frac{q}{p}}.
 \end{aligned}$$

By combining the above estimates we obtain the upper bound  $C \ll \mathbb{B}_b$ .

For the lower estimate we suppose that  $C < \infty$  and note that in view of (2.42) the inequality  $C \gg B_b$  holds. Under the condition  $V_a(d) = \infty$  let us prove first that  $B_b \gg \mathbb{B}_b$ . We have

$$\begin{aligned}
 \int_c^t w^q(x) V_a^q(x) dx &= \int_c^t V_a^q(x) d \left( - \int_x^t w^q(s) ds \right) \\
 &= V_a^q(c) \int_c^t w^q(x) dx + q \int_c^t \left( \int_x^t w^q(s) ds \right) [V_a(x)]^{q-1} dV_a(x).
 \end{aligned}$$

Then

$$\begin{aligned}
 \mathbb{B}_b^r &\approx \mathbb{B}_{PS}^r \approx V_a^r(c) \int_c^d \left( \int_c^t w^q(s) ds \right)^{\frac{r}{q}} d \left( -[V_a(t)]^{-r/p} \right) \\
 &+ \int_c^d \left\{ \int_c^t \left( \int_x^t w^q(s) ds \right) [V_a(x)]^{q-1} dV_a(x) \right\}^{\frac{r}{q}} d \left( -[V_a(t)]^{-r/p} \right) =: I_1 + I_2.
 \end{aligned}$$

Obviously,

$$I_1 \leq [V_a(c)]^{r/p'} \left( \int_c^d w^q(s) ds \right)^{\frac{r}{q}} \ll B_b^r.$$

To estimate  $I_2$  we write

$$\begin{aligned}
 &\int_c^t \left( \int_x^t w^q(s) ds \right) [V_a(x)]^{q-1} dV_a(x) \\
 &= \int_c^t \left\{ \left( \int_x^t w^q(s) ds \right) [V_a(x)]^{q-1+q/2p} \right\} [V_a(x)]^{-q/2p} dV_a(x)
 \end{aligned}$$

$$\begin{aligned} &\leq \left\{ \int_c^t \left( \int_x^t w^q(s) ds \right)^{\frac{r}{q}} [V_a(x)]^{(q-1+q/2p)r/q} dV_a(x) \right\}^{\frac{q}{r}} \left( \int_c^t [V_a(x)]^{-1/2} dV_a(x) \right)^{\frac{q}{p}} \\ &\ll [V_a(t)]^{q/2p} \left( \int_c^t \left( \int_x^t w^q(s) ds \right)^{\frac{r}{q}} [V_a(x)]^{(q-1+q/2p)r/q} dV_a(x) \right)^{\frac{q}{r}}. \end{aligned}$$

By applying this estimate and the equivalence (2.42) and (2.43) we find that

$$\begin{aligned} I_2 &\ll \int_c^d \left\{ \int_c^t \left( \int_x^t w^q(s) ds \right)^{\frac{r}{q}} [V_a(x)]^{(q-1+q/2p)r/q} dV_a(x) \right\} [V_a(t)]^{r/2p} d \left( -[V_a(t)]^{-r/p} \right) \\ &\ll \int_c^d \left( \int_x^d w^q(s) ds \right)^{\frac{r}{q}} [V_a(x)]^{(q-1+q/2p)r/q} \left( \int_x^d [V_a(t)]^{r/2p-r/p-1} dV_a(t) \right) dV_a(x) \\ &\ll \int_c^d \left( \int_x^d w^q(s) ds \right)^{\frac{r}{q}} [V_a(x)]^{r/q'} dV_a(x) \ll B_b^r. \end{aligned}$$

Thus, since  $V_a(d) = \infty$ , we get that  $\mathbb{B}_b^r \approx I_1 + I_2 \ll B_b^r \ll C^r$ . If  $0 < V_a(d) < \infty$ , it follows from (2.36) with  $f(x) = v^{p'-1}(x)$  that

$$C \geq [V_a(d)]^{-1/p} \left( \int_c^d w^q(x) V_a^q(x) dx \right)^{\frac{1}{q}}.$$

By combining this estimate and the previous one we find in view of (2.49) that  $\mathbb{B}_b \ll C$ . Now (2.44) is proved. The equivalence (2.44) and (2.45) follows from (2.49).

**LEMMA 2.4.** *Let  $0 < q < p < \infty$ ,  $p > 1$ ,  $1/r = 1/q - 1/p$  and let  $0 \leq c < d \leq \infty$ ,  $0 < b \leq \infty$ . Suppose that the function  $a(x)$  is differentiable, strictly increasing and such that  $0 < a(x) \leq b$ ,  $x \in (c, d]$ , and let the operator  $T$  be defined by (2.39). Then*

$$\|T\|_{L_p(a(c),b) \rightarrow L_q(c,d)}^r \approx B_a := \int_c^d \left( \int_c^t w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{a(t)}^b v^{p'}(y) dy \right)^{\frac{r}{p'}} w^q(t) dt \quad (2.50)$$

$$\begin{aligned} &\approx \left( \int_c^d w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{a(d)}^b v^{p'}(y) dy \right)^{\frac{r}{p'}} \\ &\quad + \int_c^d \left( \int_c^t w^q(x) dx \right)^{\frac{r}{q}} d \left[ - \left( \int_{a(t)}^b v^{p'}(y) dy \right)^{\frac{r}{p'}} \right], \end{aligned} \quad (2.51)$$

$$\begin{aligned} \|T\|_{L_p(a(c),b) \rightarrow L_q(c,d)}^r &\approx \mathbb{B}_a := \int_c^d \left( \int_t^d \left[ \int_{a(x)}^b v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{a(t)}^b v^{p'}(y) dy \right)^{q-\frac{r}{p}} w^q(t) dt \end{aligned} \quad (2.52)$$

$$\begin{aligned} &\approx \left( \int_c^d \left[ \int_{a(x)}^b v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{a(c)}^b v^{p'}(y) dy \right)^{-\frac{r}{p}} \\ &\quad + \int_c^d \left( \int_t^d \left[ \int_{a(x)}^b v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{q}} d \left( \int_{a(t)}^b v^{p'}(y) dy \right)^{-\frac{r}{p}}. \end{aligned} \quad (2.53)$$

## 2.2. Fairway-function and technical lemmas.


We start with the following definition.

DEFINITION 2.4. Given boundary functions  $a(x)$  and  $b(x)$ , satisfying the conditions (1.2), a number  $p \in (1, \infty)$  and a weight function  $v(x)$  such that  $0 < v(x) < \infty$  a.e.  $x \in \mathbb{R}^+$  and  $v^{p'}(x)$  is locally integrable on  $\mathbb{R}^+$ , we define the *fairway-function*  $\sigma(x)$  such that  $a(x) < \sigma(x) < b(x)$  and

$$\int_{a(x)}^{\sigma(x)} v^{p'}(y) dy = \int_{\sigma(x)}^{b(x)} v^{p'}(y) dy \quad \text{for all } x \in \mathbb{R}^+. \quad (2.54)$$

It is possible to prove that the fairway-function is differentiable and strictly increasing.

LEMMA 2.5. Let the functions  $a(x)$  and  $b(x)$  satisfy the conditions (1.2) and  $\sigma(x)$  is the fairway-function. For any  $c \in (0, \infty)$  put  $c^- = \sigma^{-1}(a(c))$  and let  $[\mathcal{N}^-]$  be the integral part of the number



$$\mathcal{N}^- := \log_2 \frac{\int_{a(c)}^{b(c)} v^{p'}(y) dy}{\int_{a(c)}^{b(c^-)} v^{p'}(y) dy},$$

that is

$$2^{[\mathcal{N}^-]} \leq \frac{\int_{a(c)}^{b(c)} v^{p'}(y) dy}{\int_{a(c)}^{b(c^-)} v^{p'}(y) dy} < 2^{[\mathcal{N}^-] + 1}.$$

Let the point sequence  $\{x_j\}_{j=0}^{j_b}$ , where  $j_b = \begin{cases} [\mathcal{N}^-], & \text{if } \mathcal{N}^- = [\mathcal{N}^-], \\ [\mathcal{N}^-] + 1, & \text{if } \mathcal{N}^- > [\mathcal{N}^-], \end{cases}$  be defined by:

- (1)  $x_0 = c^-$ ,  $x_{j_b} = c$ ;
- (2) if  $[\mathcal{N}^-] = 0$  or  $\mathcal{N}^- = [\mathcal{N}^-] = 1$ , then  $j_b = 1$ ;
- (3) if  $\mathcal{N}^- > 1$ , then the points  $x_j$  for  $1 \leq j \leq [\mathcal{N}^-]$  are taken so that

$$\int_{a(c)}^{b(x_j)} v^{p'}(y) dy = 2 \int_{a(c)}^{b(x_{j-1})} v^{p'}(y) dy. \quad (2.55)$$



Then for any  $t \in [x_j, x_{j+1}]$ ,  $0 \leq j \leq j_b - 1$ , we have that

$$\int_{a(t)}^{b(t)} v^{p'}(y) dy \approx \int_{a(x_j)}^{b(x_{j+1})} v^{p'}(y) dy, \quad (2.56)$$

and for all  $t \in [c^-, c]$

$$\int_{a(t)}^{b(t)} v^{p'}(y) dy < \int_{a(c^-)}^{b(c)} v^{p'}(y) dy \approx \int_{a(c)}^{b(c)} v^{p'}(y) dy. \quad (2.57)$$

Moreover, if  $j_b \geq 2$  and  $0 \leq j \leq j_b - 2$  it holds that

$$\int_{a(c)}^{b(x_{j+1})} v^{p'}(y) dy = 2^l \int_{a(c)}^{b(x_j)} v^{p'}(y) dy \quad \text{for all } l \in \{1, \dots, j_b - j - 1\}. \quad (2.58)$$

*Proof.* Clearly

$$\int_{a(x_{j+1})}^{b(x_j)} v^{p'}(y) dy < \int_{a(t)}^{b(t)} v^{p'}(y) dy < \int_{a(x_j)}^{b(x_{j+1})} v^{p'}(y) dy \quad (2.59)$$

is trivial for any  $t \in [x_j, x_{j+1}]$ ,  $0 \leq j \leq j_b - 1$ . First we prove (2.56) for  $t = x_j$ . From the definition  $x_{j+1}$  we have

$$\begin{aligned} \int_{a(x_j)}^{b(x_{j+1})} v^{p'}(y) dy &\stackrel{(2.55)}{=} \int_{a(x_j)}^{a(c)} v^{p'}(y) dy \\ &\quad + 2 \int_{a(c)}^{b(x_j)} v^{p'}(y) dy \leq 3 \int_{a(x_j)}^{b(x_j)} v^{p'}(y) dy. \end{aligned} \quad (2.60)$$

Now (2.59) and (2.60) yield (2.56) for  $t = x_j$ . If  $t = x_{j+1}$ , then in view of Lemma's condition

$$\begin{aligned} \int_{a(x_{j+1})}^{b(x_{j+1})} v^{p'}(y) dy &\geq \int_{a(c)}^{b(x_{j+1})} v^{p'}(y) dy = \int_{\sigma(c^-)}^{b(x_{j+1})} v^{p'}(y) dy \\ &\geq \int_{\sigma(x_j)}^{b(x_j)} v^{p'}(y) dy + \int_{b(x_j)}^{b(x_{j+1})} v^{p'}(y) dy = \frac{1}{2} \int_{a(x_j)}^{b(x_j)} v^{p'}(y) dy \\ &\quad + \int_{b(x_j)}^{b(x_{j+1})} v^{p'}(y) dy \geq \frac{1}{2} \int_{a(x_j)}^{b(x_{j+1})} v^{p'}(y) dy. \end{aligned} \quad (2.61)$$

Thus, (2.61) and (2.59) imply (2.56) for  $t = x_{j+1}$ . For  $t \in (x_j, x_{j+1})$  we write

$$\begin{aligned} \int_{a(t)}^{b(t)} v^{p'}(y) dy &\geq \int_{a(c)}^{b(x_j)} v^{p'}(y) dy \stackrel{(2.55)}{=} \frac{1}{2} \int_{a(c)}^{b(x_{j+1})} v^{p'}(y) dy = \frac{1}{2} \int_{\sigma(c^-)}^{b(x_{j+1})} v^{p'}(y) dy \\ &\geq \frac{1}{2} \int_{\sigma(x_{j+1})}^{b(x_{j+1})} v^{p'}(y) dy = \frac{1}{4} \int_{a(x_{j+1})}^{b(x_{j+1})} v^{p'}(y) dy \stackrel{(2.61)}{\geq} \frac{1}{8} \int_{a(x_j)}^{b(x_{j+1})} v^{p'}(y) dy \end{aligned}$$

and in view of (2.59) the proof of (2.56) for  $t \in (x_j, x_{j+1})$  is completed.


The left inequality in (2.57) is obvious. Since  $\int_{a(c)}^{b(c^-)} v^{p'}(y) dy = \int_{a(c^-)}^{a(c)} v^{p'}(y) dy$ , then

$$\int_{a(c)}^{b(c)} v^{p'}(y) dy = \int_{a(c)}^{b(c)} v^{p'}(y) dy + \int_{a(c)}^{b(c^-)} v^{p'}(y) dy \leq 2 \int_{a(c)}^{b(c)} v^{p'}(y) dy \quad (2.62)$$

and (2.57) follows. The relation (2.58) is a simple consequence of (2.55).

The proof of the next lemma is analogous.

**LEMMA 2.6.** *Let the functions  $a(x)$  and  $b(x)$  satisfy the conditions (1.2) and  $\sigma(x)$  is the fairway-function. For any  $c \in (0, \infty)$  put  $c^+ = \sigma^{-1}(b(c))$  and let  $[\mathcal{N}^+]$  be the integral part of the number*



$$\mathcal{N}^+ := \log_2 \frac{\int_{a(c)}^{b(c)} v^{p'}(y) dy}{\int_{a(c^+)}^{b(c)} v^{p'}(y) dy},$$

that is

$$2^{[\mathcal{N}^+]} \leq \frac{\int_{a(c)}^{b(c)} v^{p'}(y) dy}{\int_{a(c^+)}^{b(c)} v^{p'}(y) dy} < 2^{[\mathcal{N}^+] + 1}.$$

Let the point sequence  $\{x_j\}_{j=0}^{j_a}$ , where  $j_a = \begin{cases} [\mathcal{N}^+], & \text{if } \mathcal{N}^+ = [\mathcal{N}^+], \\ [\mathcal{N}^+] + 1, & \text{if } \mathcal{N}^+ > [\mathcal{N}^+], \end{cases}$  be defined by:

- (1)  $x_0 = c, \quad x_{j_a} = c^+;$
- (2) if  $[\mathcal{N}^+] = 0$  or  $\mathcal{N}^+ = [\mathcal{N}^+] = 1$  then  $j_a = 1;$
- (3) if  $\mathcal{N}^+ > 1$  then the points  $x_j$  for  $1 \leq j \leq [\mathcal{N}^+]$  are taken so that

$$\int_{a(x_j)}^{b(c)} v^{p'}(y) dy = \frac{1}{2} \int_{a(x_{j-1})}^{b(c)} v^{p'}(y) dy. \quad (2.63)$$

Then for any  $t \in [x_j, x_{j+1}]$ ,  $0 \leq j \leq j_a - 1$ , we have (2.56), and for all  $t \in [c, c^+]$


$$\int_{a(t)}^{b(t)} v^{p'}(y) dy \leq \int_{a(c)}^{b(c^+)} v^{p'}(y) dy \approx \int_{a(c)}^{b(c)} v^{p'}(y) dy. \quad (2.64)$$

Moreover, if  $j_a \geq 2$  and  $0 \leq j \leq j_a - 2$  it holds that

$$2^l \int_{a(x_{j+l})}^{b(c)} v^{p'}(y) dy = \int_{a(x_j)}^{b(c)} v^{p'}(y) dy \quad \text{for all } l \in \{1, \dots, j_a - j - 1\}. \quad (2.65)$$

In conclusion of the section we provide two lemmas with a different kind of decomposition than in the previous two lemmas.

LEMMA 2.7. Let the functions  $a(x)$  and  $b(x)$  satisfy the conditions (1.2) and



$\sigma(x)$  is the fairway-function. For any  $d \in (0, \infty)$  put  $d^- = \sigma^{-1}(a(d))$ ,  $d^+ = \sigma^{-1}(b(d))$  and let the point sequence  $\{x_j\}_{j=-j_a}^0$  be defined by:

- (1)  $x_{-j_a} = d$ ,  $x_0 = d^+$ ;
- (2) if  $(d^+)^- \leq d$ , then  $j_a = 1$ ;
- (3) if  $(d^+)^- > d$ , then  $j_a > 1$  and  $x_{j-1} = (x_j)^-$ , where  $(x_j)^- > d$  and  $-j_a + 2 \leq j \leq 0$ .

Then for any  $t \in [x_j, x_{j+1}]$ ,  $-j_a \leq j \leq -1$ , we have (2.56). Moreover, if  $d \leq x^- \leq t \leq x \leq d^+$ ,

$$\int_{a(t)}^{b(t)} v^{p'}(y) dy \approx \int_{a(x^-)}^{b(x)} v^{p'}(y) dy \approx \int_{a(x)}^{b(x)} v^{p'}(y) dy. \quad (2.66)$$

*Proof.* We start with the proof of (2.66). Since  $\int_{a(x^-)}^{a(x)} v^{p'}(y) dy = \int_{a(x)}^{b(x^-)} v^{p'}(y) dy$  and  $d \leq x^- \leq t \leq x \leq d^+$ , then


$$\begin{aligned} \int_{a(t)}^{b(t)} v^{p'}(y) dy &\leq \int_{a(x^-)}^{b(x)} v^{p'}(y) dy = \int_{a(x)}^{b(x^-)} v^{p'}(y) dy + \int_{a(x)}^{b(x)} v^{p'}(y) dy \\ &\leq 2 \int_{a(x)}^{b(x)} v^{p'}(y) dy \leq 2 \int_{a(x^-)}^{b(x)} v^{p'}(y) dy, \end{aligned} \quad (2.67)$$

and the second equivalence in (2.66) is proved. On the other hand, in view of  $d \leq x^- \leq t \leq x \leq d^+$  and Lemma's condition we have that

$$\int_{a(t)}^{b(t)} v^{p'}(y) dy \geq \int_{a(x)}^{b(d)} v^{p'}(y) dy = \int_{a(x)}^{\sigma(d^+)} v^{p'}(y) dy \geq \int_{a(x)}^{\sigma(x)} v^{p'}(y) dy$$

$$= \frac{1}{2} \int_{a(x)}^{b(x)} v^{p'}(y) dy \stackrel{(2.67)}{\geq} \frac{1}{4} \int_{a(x^-)}^{b(x)} v^{p'}(y) dy, \quad (2.68)$$

and now (2.66) is proved. Since  $d \leq x_j = (x_{j+1})^- \leq t \leq x_{j+1} \leq d^+$ , (2.56) follows from (2.66).



LEMMA 2.8. Let the functions  $a(x)$  and  $b(x)$  satisfy the conditions (1.2) and

$\sigma(x)$  is the fairway-function.

For any  $d \in (0, \infty)$  put  $d^- = \sigma^{-1}(a(d))$ ,  $d^+ = \sigma^{-1}(b(d))$  and let the point sequence  $\{x_j\}_{j=0}^{j_b}$  be given by:

- (1)  $x_0 = d^-$ ,  $x_{j_b} = d$ ;
- (2) if  $(d^-)^+ \geq d$ , then  $j_b = 1$ ;
- (3) if  $(d^-)^+ < d$ , then  $j_b > 1$  and  $x_{j+1} = (x_j)^+$ , where  $(x_j)^+ < d$  and  $0 \leq j \leq j_b - 2$ .

Then for any  $t \in [x_j, x_{j+1}]$ ,  $0 \leq j \leq j_b - 1$ , we have (2.56). Moreover, if  $d^- \leq x \leq t \leq x^+ \leq d$ ,

$$\int_{a(t)}^{b(t)} v^{p'}(y) dy \approx \int_{a(x)}^{b(x^+)} v^{p'}(y) dy \approx \int_{a(x)}^{b(x)} v^{p'}(y) dy. \quad (2.69)$$

### 3. Block-diagonal method

#### 3.1. Key lemma.

We need the following assertion about a block-diagonal operator.

LEMMA 3.1. Let  $U = \sqcup_k U_k$  and  $V = \sqcup_k V_k$  be unions of non-overlapping measurable sets and  $T = \sum_k T_k$ , where  $T_k: L_p(U_k) \rightarrow L_q(V_k)$ . Then

$$\|T\|_{L_p(U) \rightarrow L_q(V)} = \sup_k \|T_k\|_{L_p(U_k) \rightarrow L_q(V_k)}, \quad 0 < p \leq q < \infty \quad (3.1)$$

and

$$\|T\|_{L_p(U) \rightarrow L_q(V)} \approx \left( \sum_k \|T_k\|_{L_p(U_k) \rightarrow L_q(V_k)}^r \right)^{1/r}, \quad 0 < q < p < \infty, \quad 1/r = 1/q - 1/p. \quad (3.2)$$

*Proof.* Let  $0 < p \leq q < \infty$ . Since  $\|Tf\|_{L_q(V)} \geq \|T_k f\|_{L_q(V_k)}$  for all  $k$ , the lower estimate in (3.1) is trivial. For the upper bound, applying Jensen's inequality, we write

$$\begin{aligned} \|Tf\|_{L_q(V)}^q &= \sum_k \|T_k f\|_{L_q(V_k)}^q \leq \left( \sup_k \|T_k\|_{L_p(U_k) \rightarrow L_q(V_k)} \right)^q \sum_k \|f \chi_k\|_{L_p(U_k)}^q \\ &\leq \left( \sup_k \|T_k\|_{L_p(U_k) \rightarrow L_q(V_k)} \right)^q \|f\|_{L_p(U)}^q. \end{aligned} \quad (3.3)$$

Let  $0 < q < p < \infty$ . The upper bound in (3.2) follows similar to (3.3) by application of Hölder's inequality. For the lower bound let  $0 < \lambda < 1$  and the functions  $f_k \in L_p(U_k)$  are such that

$$\lambda \|T_k\|_{L_p(U_k) \rightarrow L_q(V_k)} \|f_k\|_{L_p(U_k)} \leq \|T_k f_k\|_{L_p(U_k)}.$$

Because of homogeneity we can choose  $f_k$  so that

$$\|f_k\|_{L_p(U_k)} = \|T_k\|_{L_p(U_k) \rightarrow L_q(V_k)}^{r/p}.$$

If we put  $f = \sum_k \chi_{U_k} f_k$ , then in view of  $r/q = r/p + 1$

$$\begin{aligned} \lambda^q \sum_k \|T_k\|_{L_p(U_k) \rightarrow L_q(V_k)}^r &= \lambda^q \sum_k \left( \|T_k\|_{L_p(U_k) \rightarrow L_q(V_k)} \|f_k\|_{L_p(U_k)} \right)^q \\ &\leq \sum_k \|T_k f_k\|_{L_q(V_k)}^q = \|Tf\|_{L_q(V)}^q \leq \|T\|_{L_p(U) \rightarrow L_q(V)}^q \|f\|_{L_p(U)}^q \\ &= \|T\|_{L_p(U) \rightarrow L_q(V)}^q \left( \sum_k \|T_k\|_{L_p(U_k) \rightarrow L_q(V_k)}^r \right)^{q/p} \end{aligned}$$

and the lower bound in (3.2) follows by tending  $\lambda \rightarrow 1$ .

### 3.2. Integral operators with Oinarov type kernels.

We demonstrate the block-diagonal method for the operator (1.6)

$$\mathcal{K}^* f(x) := w(x) \int_{a(x)}^{b(x)} k(y, x) f(y) v(y) dy$$

with a kernel  $k(y, x) \geq 0$  satisfying


$$k(y, x) \approx k(y, z) + k(a(z), x), \quad x \leq z, \quad a(z) \leq y \leq b(x). \quad (3.4)$$

Given functions  $a(x)$  and  $b(x)$  satisfying (1.2) we take a sequence of points  $\{\xi_k\}_{k \in \mathbb{Z}} \subset (0, \infty)$  such that

$$\xi_0 = 1, \quad \xi_k = (a^{-1} \circ b)^k(1), \quad k \in \mathbb{Z}, \quad (3.5)$$

and put

$$\eta_k = a(\xi_k) = b(\xi_{k-1}), \quad \Delta_k = [\xi_k, \xi_{k+1}), \quad \delta_k = [\eta_k, \eta_{k+1}), \quad k \in \mathbb{Z}. \quad (3.6)$$



Breaking the semiaxis  $(0, \infty)$  into intervals by points of the sequence  $\{\xi_k\}_{k \in \mathbb{Z}}$ , we decompose the operator  $\mathcal{K}^*$  into the sum

$$\mathcal{K}^* = \mathcal{T} + \mathcal{S} \quad (3.7)$$

of block-diagonal operators  $\mathcal{T}$  and  $\mathcal{S}$  such that

$$\mathcal{T} = \sum_{k \in \mathbb{Z}} T_k, \quad \mathcal{S} = \sum_{k \in \mathbb{Z}} S_k, \quad (3.8)$$

where

$$T_k f(x) = w(x) \int_{a(x)}^{a(\xi_{k+1})} k(y, x) f(y) v(y) dy, \quad T_k : L_p(\delta_k) \rightarrow L_q(\Delta_k),$$

$$S_k f(x) = w(x) \int_{b(\xi_k)}^{b(x)} k(y, x) f(y) v(y) dy, \quad S_k : L_p(\delta_{k+1}) \rightarrow L_q(\Delta_k).$$

$\mathcal{K}^*$ ,  $\mathcal{T}$  and  $\mathcal{S}$  are integral operators with non-negative kernels, then

$$\|\mathcal{K}^*\|_{L_p \rightarrow L_q} \approx \|\mathcal{T}\|_{L_p \rightarrow L_q} + \|\mathcal{S}\|_{L_p \rightarrow L_q}.$$

Since  $\sqcup \Delta_k = \sqcup \delta_k = (0, \infty)$  by Lemma 3.1 we can estimate the norms of  $\mathcal{T}$  and  $\mathcal{S}$  via the norms of  $T_k$  and  $S_k$ . Moreover, kernels  $k(y, x)$  of the operators  $T_k$  and  $S_k$  satisfy the condition (3.4) for  $x \leq z$ ,  $x \in \Delta_k$  and

$$a(z) \leq y \leq a(\xi_{k+1}), \quad b(\xi_k) \leq y \leq b(x), \quad (3.9)$$

respectively. It allows us to apply preliminary results (see Section 2).

More precisely we have the following theorem.

**THEOREM 3.1.** *Let  $1 < p \leq q < \infty$ . Then*

$$\|\mathcal{K}^*\|_{L_p \rightarrow L_q} \approx \mathcal{A}^* := \mathcal{A}_0^* + \mathcal{A}_1^*, \quad (3.10)$$

where

$$\mathcal{A}_0^* := \sup_{s > 0} \mathcal{A}_0^*(s) = \sup_{s > 0} \sup_{s \leq t \leq a^{-1}(b(s))} \left( \int_s^t k^q(a(t), x) w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} v^{p'}(y) dy \right)^{\frac{1}{p'}},$$

$$\mathcal{A}_1^* := \sup_{s > 0} \mathcal{A}_1^*(s) = \sup_{s > 0} \sup_{s \leq t \leq a^{-1}(b(s))} \left( \int_s^t w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} k^{p'}(y, t) v^{p'}(y) dy \right)^{\frac{1}{p'}}.$$

Moreover,  $\mathcal{K}^*$  is compact if and only if  $\mathcal{A}^* < \infty$  and  $\lim_{s \rightarrow 0} \mathcal{A}_i^*(s) = \lim_{s \rightarrow \infty} \mathcal{A}_i^*(s) = 0$ ,  $i = 0, 1$ .

If  $1 < q < p < \infty$ , then

$$\|\mathcal{K}^*\|_{L_p \rightarrow L_q} \approx \mathcal{B}^* := \left( \sum_k \left[ (\mathcal{B}_{k,1}^*)^r + (\mathcal{B}_{k,2}^*)^r + (\mathcal{B}_{k,3}^*)^r + (\mathcal{B}_{k,4}^*)^r \right] \right)^{\frac{1}{r}}, \quad (3.11)$$

where

$$\mathcal{B}_{k,1}^* := \left( \int_{a(\xi_k)}^{a(\xi_{k+1})} \left[ \int_{\xi_k}^{a^{-1}(t)} k^q(t, x) w^q(x) dx \right]^{\frac{r}{q}} \left[ \int_t^{a(\xi_{k+1})} v^{p'}(y) dy \right]^{\frac{r}{q'}} v^{p'}(t) dt \right)^{\frac{1}{r}},$$

$$\mathcal{B}_{k,2}^* := \left( \int_{\xi_k}^{\xi_{k+1}} \left[ \int_{\xi_k}^t w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{a(t)}^{a(\xi_{k+1})} k^{p'}(y, t) v^{p'}(y) dy \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}},$$

$$\mathcal{B}_{k,3}^* := \left( \int_{b(\xi_k)}^{b(\xi_{k+1})} \left[ \int_{b^{-1}(t)}^{\xi_{k+1}} k^q(a(\xi_{k+1}), x) w^q(x) dx \right]^{\frac{r}{q}} \left[ \int_{b(\xi_k)}^t v^{p'}(y) dy \right]^{\frac{r}{q'}} v^{p'}(t) dt \right)^{\frac{1}{r}},$$

$$\mathcal{B}_{k,4}^* := \left( \int_{\xi_k}^{\xi_{k+1}} \left[ \int_t^{\xi_{k+1}} w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{b(\xi_k)}^{b(t)} k^{p'}(y, \xi_{k+1}) v^{p'}(y) dy \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}},$$

and  $\mathcal{K}^*$  is compact if and only if  $\mathcal{B}^* < \infty$ .

*Proof.* Put

$$\|\mathcal{K}^*\| = \|\mathcal{K}^*\|_{L_p \rightarrow L_q}, \quad \|T_k\| = \|T_k\|_{L_p(\delta_k) \rightarrow L_q(\Delta_k)}, \quad \|S_k\| = \|S_k\|_{L_p(\delta_{k+1}) \rightarrow L_q(\Delta_k)}.$$

Let  $1 < p \leq q < \infty$ . It follows from (3.7) – (3.8) and Lemma 3.1 that

$$\|\mathcal{K}^*\| \approx \sup_k \|T_k\| + \sup_k \|S_k\|. \quad (3.12)$$

The norm of the operator  $T_k$  is characterized by Corollary 2.2 (a):

$$\begin{aligned} \|T_k\| &\approx \sup_{t \in \Delta_k} \left( \int_{\xi_k}^t k^q(a(t), x) w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{a(\xi_{k+1})} v^{p'}(y) dy \right)^{\frac{1}{p'}} \\ &\quad + \sup_{t \in \Delta_k} \left( \int_{\xi_k}^t w^q(y) dy \right)^{\frac{1}{q}} \left( \int_{a(t)}^{a(\xi_{k+1})} k^{p'}(y, t) v^{p'}(y) dy \right)^{\frac{1}{p'}}. \end{aligned}$$

For the kernel of the operator  $S_k$  it follows from (3.4) provided  $z = \xi_{k+1}$ ,  $x \in \Delta_k$  and (3.9) that

$$k(y, x) \approx k(y, \xi_{k+1}) + k(a(\xi_{k+1}), x).$$

Therefore,  $S_k f(x) \approx S_{k,1} f(x) + S_{k,2} f(x)$ , where

$$\begin{aligned} S_{k,1} f(x) &= w(x) \int_{b(\xi_k)}^{b(x)} k(y, \xi_{k+1}) f(y) v(y) dy, & x \in \Delta_k, \\ S_{k,2} f(x) &= w(x) k(a(\xi_{k+1}), x) \int_{b(\xi_k)}^{b(x)} f(y) v(y) dy, & x \in \Delta_k. \end{aligned}$$

The norm estimates for  $S_{k,1}$  and  $S_{k,2}$  follow from Lemma 2.1:

$$\begin{aligned} \|S_k\| &\approx \sup_{t \in \Delta_k} \left( \int_t^{\xi_{k+1}} k^q(a(\xi_{k+1}), x) w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{b(\xi_k)}^{b(t)} v^{p'}(y) dy \right)^{\frac{1}{p'}} \\ &\quad + \sup_{t \in \Delta_k} \left( \int_t^{\xi_{k+1}} w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{b(\xi_k)}^{b(t)} k^{p'}(y, \xi_{k+1}) v^{p'}(y) dy \right)^{\frac{1}{p'}}. \end{aligned}$$

Define the following functions

$$\begin{aligned} \mathcal{A}_0(s, t) &= \left( \int_s^t k^q(a(t), x) w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} v^{p'}(y) dy \right)^{\frac{1}{p'}}, \\ \mathcal{A}_1(s, t) &= \left( \int_s^t w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} k^{p'}(y, t) v^{p'}(y) dy \right)^{\frac{1}{p'}}, \end{aligned}$$

where  $s < t < a^{-1}(b(s))$ . Then

$$\|T_k\| \approx \sup_{t \in \Delta_k} \mathcal{A}_0(\xi_k, t) + \sup_{t \in \Delta_k} \mathcal{A}_1(\xi_k, t), \quad (3.13)$$

$$\|S_k\| \approx \sup_{t \in \Delta_k} \mathcal{A}_0(t, \xi_{k+1}) + \sup_{t \in \Delta_k} \mathcal{A}_1(t, \xi_{k+1}) \quad (3.14)$$

and

$$\begin{aligned} \mathcal{A}_0^* &= \sup_{s>0} \sup_{s \leq t \leq a^{-1}(b(s))} \mathcal{A}_0(s, t), & \mathcal{A}_1^* &= \sup_{s>0} \sup_{s \leq t \leq a^{-1}(b(s))} \mathcal{A}_1(s, t), \\ \sup_{t \in \Delta_k} \mathcal{A}_i(\xi_k, t) &\leq \sup_{\xi_k \leq t \leq a^{-1}(b(\xi_k))} \mathcal{A}_i(\xi_k, t) \leq \mathcal{A}_i^*, & i &= 0, 1, \\ \sup_{t \in \Delta_k} \mathcal{A}_i(t, \xi_{k+1}) &\leq \sup_{t \leq \xi_{k+1} \leq a^{-1}(b(t))} \mathcal{A}_i(t, \xi_{k+1}) \leq \mathcal{A}_i^*, & i &= 0, 1. \end{aligned}$$

Now it follows from (3.12) – (3.14) that

$$\|\mathcal{K}^*\| \ll \mathcal{A}_0^* + \mathcal{A}_1^*.$$



For the lower bound in (3.10) we suppose that  $\|\mathcal{K}^*\| < \infty$  and  $s < x < t < a^{-1}(b(s))$ . If

$$f(y) = \chi_{[a(t), b(s)]}(y) v^{p'-1}(y),$$

then

$$\mathcal{K}^* f(x) \geq w(x) \int_{a(t)}^{b(s)} k(y, x) v^{p'}(y) dy.$$

The condition (3.4) with  $z = t$  implies  $k(y, x) \gg k(a(t), x)$ , therefore

$$\mathcal{K}^* f(x) \gg w(x) k(a(t), x) \int_{a(t)}^{b(s)} v^{p'}(y) dy.$$

Hence,

$$\|\mathcal{K}^*\| \geq \frac{\|\mathcal{K}^* f\|_q}{\|f\|_p} \geq \left( \int_s^t k^q(a(t), x) w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} v^{p'}(y) dy \right)^{\frac{1}{p'}}.$$

Therefore,  $\|\mathcal{K}^*\| \geq \mathcal{A}_0(s, t)$  for all  $s < t < a^{-1}(b(s))$ . Consequently,  $\|\mathcal{K}^*\| \geq \mathcal{A}_0^*$ . The inequality  $\|\mathcal{K}^*\| \geq \mathcal{A}_1^*$  can be proved analogously by applying a test function

$$g(x) = \chi_{[s, t]}(x) w^{q-1}(x)$$

into the inequality for the operator dual to  $\mathcal{K}^*$ .

To prove (3.11) we note that by Lemma 3.1

$$\|\mathcal{K}^*\| \approx \left( \sum_k \|T_k\|^r \right)^{\frac{1}{r}} + \left( \sum_k \|S_k\|^r \right)^{\frac{1}{r}}. \quad (3.15)$$

Norms of the operators  $T_k$  and  $S_k$  are estimated with the help of Corollary 2.2 (b) and Lemma 2.3 by the following way:

$$\|T_k\| \approx \mathcal{B}_{k,1}^* + \mathcal{B}_{k,2}^*, \quad \|S_k\| \approx \mathcal{B}_{k,3}^* + \mathcal{B}_{k,4}^* \quad (3.16)$$

and the required result follows.

The proof of compactness assertion of the theorem for  $1 < p \leq q < \infty$  follows from representation of the operator by the sum of a compact operator and an operator with a small norm. For  $1 < q < p < \infty$  the required result follows by applying Ando's theorem (see [11], [16] and [27]).

Using decomposition (3.5) for the operator  $\mathcal{K}$  defined by (1.4) with the kernel  $k(x, y) \geq 0$  satisfying the condition (1.5) we obtain the analogous result for  $\mathcal{K}$ .

**THEOREM 3.2.** *If  $1 < p \leq q < \infty$ , then*

$$\|\mathcal{K}\|_{L_p \rightarrow L_q} \approx \mathcal{A} := \mathcal{A}_0 + \mathcal{A}_1,$$

where

$$\mathcal{A}_0 := \sup_{t>0} \mathcal{A}_0(t) = \sup_{t>0} \sup_{b^{-1}(a(t)) \leq s \leq t} \left( \int_s^t k^q(x, b(s)) w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} v^{p'}(y) dy \right)^{\frac{1}{p'}},$$

$$\mathcal{A}_1 := \sup_{t>0} \mathcal{A}_1(t) = \sup_{t>0} \sup_{b^{-1}(a(t)) \leq s \leq t} \left( \int_s^t w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} k^{p'}(s, y) v^{p'}(y) dy \right)^{\frac{1}{p'}}.$$

Moreover,  $\mathcal{K}$  is compact if and only if  $\mathcal{A} < \infty$  and  $\lim_{t \rightarrow 0} \mathcal{A}_i(t) = \lim_{t \rightarrow \infty} \mathcal{A}_i(t) = 0, i = 0, 1$ .

If  $1 < q < p < \infty$ , then

$$\|\mathcal{K}\|_{L_p \rightarrow L_q} \approx \mathcal{B} := \left( \sum_k [\mathcal{B}_{k,1}^r + \mathcal{B}_{k,2}^r + \mathcal{B}_{k,3}^r + \mathcal{B}_{k,4}^r] \right)^{\frac{1}{r}},$$

where

$$\mathcal{B}_{k,1} := \left( \int_{a(\xi_k)}^{a(\xi_{k+1})} \left[ \int_{\xi_k}^{a^{-1}(t)} k^q(x, b(\xi_k)) w^q(x) dx \right]^{\frac{r}{q}} \left[ \int_t^{a(\xi_{k+1})} v^{p'}(y) dy \right]^{\frac{r}{q'}} v^{p'}(t) dt \right)^{\frac{1}{r}},$$

$$\mathcal{B}_{k,2} := \left( \int_{\xi_k}^{\xi_{k+1}} \left[ \int_{\xi_k}^t w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{a(t)}^{a(\xi_{k+1})} k^{p'}(\xi_k, y) v^{p'}(y) dy \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}},$$

$$\mathcal{B}_{k,3} := \left( \int_{b(\xi_k)}^{b(\xi_{k+1})} \left[ \int_{b^{-1}(t)}^{\xi_{k+1}} k^q(x, t) w^q(x) dx \right]^{\frac{r}{q}} \left[ \int_{b(\xi_k)}^t v^{p'}(y) dy \right]^{\frac{r}{q'}} v^{p'}(t) dt \right)^{\frac{1}{r}},$$

$$\mathcal{B}_{k,4} := \left( \int_{\xi_k}^{\xi_{k+1}} \left[ \int_t^{\xi_{k+1}} w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{b(\xi_k)}^{b(t)} k^{p'}(t, y) v^{p'}(y) dy \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}},$$


and the operator  $\mathcal{K}$  is compact if and only if  $\mathcal{B} < \infty$ .

REMARK 3.1. It is curious to note, that in spite of duality of Theorems 3.1 and 3.2 the conditions (3.4) and (1.5) are independent in general, that is one of them can hold whereas the other is broken. In practice the operators (1.4) and (1.6) are almost indistinguishable and, therefore, a form of criterion entirely depends on which of conditions (3.4) or (1.5) holds. This phenomenon is caused by the following. The condition (3.4) extends (2.3) for the Volterra operator with variable lower limit, but (1.5) generalizes (2.3) for the operator with variable upper limit. Losing Volterra's form the operator ((1.4) or (1.6)) forgets the origin, but its kernel remembers.

## 4. Hardy-Steklov operator

### 4.1. Muckenhoupt and Mazya-Rosin type criteria

In this section we give complete analogs of the conditions (2.4) and (2.6) for the operator (1.1) basing on the fairway-function conception.



Put

$$\Delta(x) = [a(x), b(x)],$$

$$\delta(x) = [b^{-1}(\sigma(x)), a^{-1}(\sigma(x))],$$

where  $a^{-1}(y)$  and  $b^{-1}(y)$  are the functions converse to  $y = a(x)$  and  $y = b(x)$ , respectively.

**THEOREM 4.1.** *Let the operator  $\mathcal{H}$  of the form (1.1) be given with the boundary functions  $a(x)$  and  $b(x)$  satisfying the conditions (1.2). Then for the norm of  $\mathcal{H}$  and  $1 < p \leq q < \infty$  the estimate*

$$\|\mathcal{H}\|_{L_p \rightarrow L_q} \approx \mathcal{A}_M, \quad (4.1)$$

holds, where

$$\mathcal{A}_M := \sup_{t>0} \mathcal{A}_M(t) = \sup_{t>0} \left( \int_{\delta(t)} w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{\frac{1}{p'}}. \quad (4.2)$$

Moreover,  $\mathcal{H} : L_p \rightarrow L_q$  is compact if and only if  $\mathcal{A}_M < \infty$  and  $\lim_{t \rightarrow 0} \mathcal{A}_M(t) = \lim_{t \rightarrow \infty} \mathcal{A}_M(t) = 0$ .

If  $0 < q < p < \infty$ ,  $p > 1$ ,  $1/r = 1/q - 1/p$ , then

$$\|\mathcal{H}\|_{L_p \rightarrow L_q} \approx \mathcal{B}_{MR}, \quad (4.3)$$

where

$$\mathcal{B}_{MR} := \left( \int_0^\infty \left[ \int_{\delta(t)} w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{\Delta(t)} v^{p'}(y) dy \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}} \quad (4.4)$$

and  $\mathcal{H} : L_p \rightarrow L_q$  is compact if and only if  $\mathcal{B}_{MR} < \infty$ .

*Proof.* First we consider the case  $1 < p \leq q < \infty$ . It follows from Theorem 3.1 with  $k(y, x) = 1$  that  $\mathcal{A}_0^* = \mathcal{A}_1^*$  and by (3.10)

$$\|\mathcal{H}\| := \|\mathcal{H}\|_{L_p \rightarrow L_q} \approx \mathbb{A} := \sup_{s>0} \sup_{s \leq t \leq a^{-1}(b(s))} \mathbb{A}(s, t), \quad (4.5)$$

where

$$\mathbb{A}(s, t) = \left( \int_s^t w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} v^{p'}(y) dy \right)^{\frac{1}{p'}}.$$

Let  $\mathbb{A}^* = \sup_{s>0} \sup_{b^{-1}(a(s)) \leq t \leq s} \mathbb{A}(t, s)$ . Then  $\mathbb{A} = \mathbb{A}^*$  and using (2.54) we find for all  $t > 0$

$$\begin{aligned} \mathcal{A}_M(t) &\approx \left( \int_{b^{-1}(\sigma(t))}^t w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{\sigma(t)} v^{p'}(y) dy \right)^{\frac{1}{p'}} \\ &\quad + \left( \int_t^{a^{-1}(\sigma(t))} w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{\sigma(t)}^{b(t)} v^{p'}(y) dy \right)^{\frac{1}{p'}} \\ &= \mathbb{A}(b^{-1}(\sigma(t)), t) + \mathbb{A}^*(t, a^{-1}(\sigma(t))) \leq 2\mathbb{A}. \end{aligned}$$

It implies  $\mathcal{A}_M \ll \mathbb{A}$ . For the proof of the opposite inequality we put  $\tau = \sigma^{-1}(b(s))$  and write

$$\mathbb{A} \leq \sup_{s \leq t \leq \tau} \mathbb{A}(s, t) + \sup_{\tau \leq t \leq a^{-1}(b(s))} \mathbb{A}(s, t) \ll \mathcal{A}_M.$$


Indeed, if  $s \leq t \leq \tau$ , then  $(s, t) \subset (b^{-1}(\sigma(t)), a^{-1}(\sigma(t)))$ ,  $(a(t), b(s)) \subset (a(t), b(t))$  and  $\sup_{s \leq t \leq \tau} \mathbb{A}(s, t) \leq \mathcal{A}_M$ . If  $s \leq \tau \leq t \leq a^{-1}(b(s))$ , then  $(s, t) \subset (s, a^{-1}(b(s))) = (b^{-1}(\sigma(\tau)), a^{-1}(\sigma(\tau)))$ ,  $(a(t), b(s)) \subset (a(\tau), b(\tau))$  and  $\sup_{\tau < t < a^{-1}(b(s))} \mathbb{A}(s, t) \leq \mathcal{A}_M$ . Therefore,  $\mathcal{A}_M \approx \mathbb{A}$  and (4.1) follows from (4.5). The criterion of compactness of the operator  $\mathcal{H}$  follows from Theorem 3.1.

Now we consider the case  $0 < q < p < \infty$ ,  $p > 1$ . For the proof (4.3) we show first that  $\|\mathcal{H}\| \ll \mathcal{B}_{MR}$ . To this end we introduce some notations:

$$\begin{aligned} \xi_k^- &= \sigma^{-1}(a(\xi_k)), \quad \xi_k^+ = \sigma^{-1}(b(\xi_k)), \quad \Delta_k = [\xi_k^-, \xi_k^+] = \Delta_k^- \cup \Delta_k^+, \\ \Delta_k^- &= [\xi_k^-, \xi_k], \quad \Delta_k^+ = [\xi_k, \xi_k^+], \quad k \in \mathbb{Z}, \end{aligned}$$

where  $\{\xi_k\}_{k \in \mathbb{Z}}$  are given by relation (3.5). The operator  $\mathcal{H}$  with  $x \in \Delta_k$  splits into the sum of four operators

$$\mathcal{H}f(x) = T_{k,1}f(x) + T_{k,2}f(x) + S_{k,1}f(x) + S_{k,2}f(x), \quad x \in \Delta_k,$$



where

$$T_{k,1}f(x) = w(x) \int_{a(x)}^{a(\xi_k)} f(y)v(y)dy, \quad x \in \Delta_k^-,$$

$$T_{k,2}f(x) = w(x) \int_{a(x)}^{b(\xi_k)} f(y)v(y)dy, \quad x \in \Delta_k^+,$$

$$S_{k,1}f(x) = w(x) \int_{b(\xi_k)}^{b(x)} f(y)v(y)dy, \quad x \in \Delta_k^+,$$

$$S_{k,2}f(x) = w(x) \int_{a(\xi_k)}^{b(x)} f(y)v(y)dy, \quad x \in \Delta_k^-.$$

Applying known estimates of these operators (see lemmas 2.1 – 2.4) we receive

$$\begin{aligned} \|T_{k,1}\|^r &\approx \int_{\xi_k^-}^{\xi_k} \left( \int_{\xi_k^-}^t w^q(x)dx \right)^{\frac{r}{p}} \left( \int_{a(t)}^{a(\xi_k)} v^{p'}(y)dy \right)^{\frac{r}{p'}} w^q(t)dt \\ &\leq \frac{q}{r} \left( \int_{a(\xi_k^-)}^{\sigma(\xi_k^-)} v^{p'}(y)dy \right)^{\frac{r}{p'}} \left( \int_{\xi_k^-}^{\xi_k} w^q(x)dx \right)^{\frac{r}{q}} = \\ &\stackrel{(2.54)}{=} \left( \int_{\sigma(\xi_k^-)}^{b(\xi_k^-)} v^{p'}(y)dy \right)^{\frac{r}{p'}} \left( \int_{\xi_k^-}^{\xi_k} \left( \int_t^{\xi_k} w^q(x)dx \right)^{\frac{r}{p}} w^q(t)dt \right). \end{aligned}$$

For  $t \in [\xi_k^-, \xi_k]$  we have  $a(t) \leq \sigma(\xi_k^-)$ ,  $b(\xi_k^-) \leq b(t)$ ,  $\xi_k \leq a^{-1}(\sigma(t))$ , then  $[\sigma(\xi_k^-), b(\xi_k^-)] \subseteq \Delta(t)$ ,  $[t, \xi_k] \subset [t, a^{-1}(\sigma(t))] \subset \delta(t)$ , therefore

$$\|T_{k,1}\|^r \ll \int_{\xi_k^-}^{\xi_k} \left( \int_{\delta(t)} w^q(x)dx \right)^{\frac{r}{p}} \left( \int_{\Delta(t)} v^{p'}(y)dy \right)^{\frac{r}{p'}} w^q(t)dt =: J_{k,1}. \quad (4.6)$$

Analogously, we find

$$\|T_{k,2}\|^r \approx \int_{\xi_k}^{\xi_k^+} \left( \int_{\xi_k}^t w^q(x)dx \right)^{\frac{r}{p}} \left( \int_{a(t)}^{b(\xi_k)} v^{p'}(y)dy \right)^{\frac{r}{p'}} w^q(t)dt.$$

Since for  $t \in [\xi_k, \xi_k^+]$  we have  $\xi_k \geq b^{-1}(\sigma(t))$ ,  $b(\xi_k) \leq b(t)$ , then  $[\xi_k, t] \subset (b^{-1}(\sigma(t)), t) \subset \delta(t)$ ,  $[a(t), b(\xi_k)] \subset \Delta(t)$ , hence

$$\|T_{k,2}\|^r \ll \int_{\xi_k}^{\xi_k^+} \left( \int_{\delta(t)} w^q(x)dx \right)^{\frac{r}{p}} \left( \int_{\Delta(t)} v^{p'}(y)dy \right)^{\frac{r}{p'}} w^q(t)dt =: J_{k,2}. \quad (4.7)$$

Similarly, we prove that

$$\|S_{k,1}\|^r \ll J_{k,2}, \quad \|S_{k,2}\|^r \ll J_{k,1}. \quad (4.8)$$

Using a decomposition of operator  $\mathcal{H}$  and the estimates (4.6) – (4.8) we obtain the required inequality  $\|\mathcal{H}\| \ll \mathcal{B}_{MR}$ .

For the proof of the estimate  $\|\mathcal{H}\| \gg \mathcal{B}_{MR}$  we suppose first that

$$\sigma(x) = x, \quad (4.9)$$

i.e. fairway is the bisectrix of the first quadrant. In this case we denote

$$\begin{aligned} x^- &= a(x), & x^+ &= b(x), & \Delta(x) &= [x^-, x^+] = \Delta^-(x) \cup \Delta^+(x), \\ \Delta^-(x) &= [x^-, x], & \Delta^+(x) &= [x, x^+] \end{aligned}$$

and we name *basic* an interval of the form  $[x^-, x^+]$ . We need the following

**DEFINITION 4.1.** Let  $p \in (1, \infty)$  and the functions  $a(x)$ ,  $b(x)$  and weight function  $v(x)$  satisfy the conditions of Definition 2.4 provided (4.9) is fulfilled. Define  $\mathcal{L}$  as the set of all absolutely continuous functions  $F$  on  $(0, \infty)$  such that  $\|F'/v\|_p < \infty$  and if  $F \in \mathcal{L}$ , then there exist mutually disjoint intervals  $I_k = (\alpha_k, \beta_k) \subset (0, \infty)$  and basic intervals  $J_k = [c_k^-, c_k^+]$  such that  $I_k \subset J_k$ ,  $\text{supp} F \subset \bigcup_k I_k$  and  $F(\alpha_k) = F(\beta_k) = 0$  for all  $k$ .

We consider the inequality

$$\|Fw\|_q \leq C \|F'/v\|_p, \quad F \in \mathcal{L}, \quad (4.10)$$

with a constant  $C$  independent of  $F \in \mathcal{L}$  and chosen as the least possible. We show that

$$C \leq \|\mathcal{H}\|. \quad (4.11)$$

To this end for any function  $F \in \mathcal{L}$  we write

$$\int_0^\infty |F(x)|^q w^q(x) dx = \sum_k \int_{I_k} |F(x)|^q w^q(x) dx.$$

Since  $I_k = (\alpha_k, \beta_k) \subset [c_k^-, c_k^+] = J_k$  the only three variants are possible:

$$(i) \quad \beta_k \leq c_k, \quad (ii) \quad c_k \leq \alpha_k, \quad (iii) \quad c_k \in I_k.$$

In the case (i) we have

$$\int_{I_k} |F(x)|^q w^q(x) dx \leq \int_{I_k} \left( \int_{\alpha_k}^x |F'(y)| dy \right)^q w^q(x) dx \leq \int_{I_k} \left( \int_{\Delta(x)} |F'(y)| dy \right)^q w^q(x) dx,$$

because  $(\alpha_k, x) \subset (a(x), x) \subset \Delta(x)$  on the strength of  $\alpha_k \geq c_k^- = a(c_k) \geq a(\beta_k) \geq a(x)$ . Analogously, for the case (ii) we write

$$\int_{I_k} |F(x)|^q w^q(x) dx \leq \int_{I_k} \left( \int_x^{\beta_k} |F'(y)| dy \right)^q w^q(x) dx \leq \int_{I_k} \left( \int_{\Delta(x)} |F'(y)| dy \right)^q w^q(x) dx$$

since  $(x, \beta_k) \subset (x, b(x)) \subset \Delta(x)$  which follows from the chain of inequalities  $\beta_k \leq c_k^+ = b(c_k) \leq b(\alpha_k) \leq b(x)$ . Thus, using the above arguments, we obtain for the case (iii)

$$\begin{aligned} \int_{I_k} |F(x)|^q w^q(x) dx &\leq \int_{\alpha_k}^{c_k} \left( \int_{\alpha_k}^x |F'(y)| dy \right)^q w^q(x) dx + \int_{c_k}^{\beta_k} \left( \int_x^{\beta_k} |F'(y)| dy \right)^q w^q(x) dx \\ &\ll \int_{I_k} \left( \int_{\Delta(x)} |F'(y)| dy \right)^q w^q(x) dx. \end{aligned}$$

Consequently,

$$\int_0^\infty |F(x)|^q w^q(x) dx \leq \sum_k \int_{I_k} (\mathcal{H}|F'/v|)^q(x) dx \leq \|\mathcal{H}|F'/v|\|_q^q \leq \|\mathcal{H}\|^q \|F'/v\|_p^q$$


and the inequality (4.11) is proved. Thus, Theorem 4.1 will be established under the condition (4.9) if we show that

$$\mathcal{B}_{MR} \ll C. \quad (4.12)$$

For this purpose it is sufficient to prove that  $\mathcal{B}_{MR}^\pm \ll C$ , where

$$\mathcal{B}_{MR}^\pm = \left( \int_0^\infty \left[ \int_{\delta^\pm(t)} w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{\Delta(t)} v^{p'}(y) dy \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}},$$

and  $\delta^-(t) = [b^{-1}(t), t]$ ,  $\delta^+(t) = [t, a^{-1}(t)]$ . We prove the inequality  $\mathcal{B}_{MR}^+ \ll C$ , arguments for  $\mathcal{B}_{MR}^- \ll C$  are similar.



Let  $\eta_0 = 1$  and the sequence  $\{\eta_k\}_{k \in \mathbb{Z}} \subset (0, \infty)$  be defined by

$$\eta_{k+1} = a^{-1}(\eta_k), \quad k \in \mathbb{Z}.$$

Put

$$\eta_k^* = \min(\eta_k^+, \eta_{k+1}).$$

For a fixed  $k \in \mathbb{Z}$  we take five neighboring points  $\eta_{k-1}$ ,  $\eta_k$ ,  $\eta_{k+1}$ ,  $\eta_{k+2}$ ,  $\eta_{k+2}^*$  and let

$$g_k(t) = \chi_{[\eta_{k-1}, \eta_{k+2}]}(t) \int_{\eta_{k-1}}^t \left( \int_s^{\eta_{k+2}} w^q(x) dx \right)^{\frac{r}{pq}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{p'}} v^{p'}(s) ds.$$

On the interval  $[\eta_{k-1}, \eta_{k+2}^*]$  we define the function

$$h_k(t) = \begin{cases} g_k(t), & t \notin [\eta_{k+2}, \eta_{k+2}^*], \\ g_k(\eta_{k+2}) \Omega_{k+2}(t), & t \in [\eta_{k+2}, \eta_{k+2}^*], \end{cases}$$

where

$$\Omega_l(t) = \left( \int_{\eta_l}^{\eta_l^*} v^{p'}(y) dy \right)^{-1} \int_t^{\eta_l^*} v^{p'}(s) ds.$$

Put

$$\lambda_k := \int_{\eta_{k-1}}^{\eta_{k+2}} \left( \int_s^{\eta_{k+2}} w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{q'}} v^{p'}(s) ds. \quad (4.13)$$

and observe, that  $\lambda_k \in (0, \infty)$  for all  $k$ . Now we find decomposition

$$h_k(t) = \sum_{i=1}^3 h_{k,i}(t), \quad (4.14)$$

with  $h_{k,i} \in \mathcal{L}$ ,  $i = 1, 2, 3$ . To this end we determine  $h_{k,1}$  by

$$h_{k,1}(t) = \begin{cases} 0, & t \notin [\eta_{k-1}, \eta_k^*], \\ h_k(t), & t \in [\eta_{k-1}, \eta_k], \\ g_k(\eta_k) \Omega_k(t), & t \in [\eta_k, \eta_k^*]. \end{cases}$$

Since  $\eta_k^- = \eta_{k-1}$ , then  $\text{supp } h_{k,1} \subseteq [\eta_k^-, \eta_k^*] \subseteq [\eta_k^-, \eta_k^+]$ , therefore  $h_{k,1} \in \mathcal{L}$ . Now, let

$$h_k^{(1)}(t) := h_k(t) - h_{k,1}(t).$$

Define

$$h_{k,2}(t) = \begin{cases} 0, & t \notin [\eta_k, \eta_{k+1}^*], \\ h_k^{(1)}(t), & t \in [\eta_k, \eta_{k+1}], \\ h_k^{(1)}(\eta_{k+1}) \Omega_{k+1}(t), & t \in [\eta_{k+1}, \eta_{k+1}^*]. \end{cases}$$

Obviously,  $h_{k,2} \in \mathcal{L}$  and  $h_{k,3} \in \mathcal{L}$  too, where

$$h_{k,3}(t) = h_k^{(1)}(t) - h_{k,2}(t).$$

Now it is clear that (4.14) holds. We also need the following estimates

$$\|h'_{k,i}/v\|_p^p \ll \lambda_k, \quad i = 1, 2, 3. \quad (4.15)$$

Write

$$\kappa_1 := \|h'_{k,1}/v\|_p^p = \int_{\eta_{k-1}}^{\eta_k} |h'_k(s)|^p v^{-p}(s) ds + g_k^p(\eta_k) \int_{\eta_k}^{\eta_k^*} |\Omega'_k(s)|^p v^{-p}(s) ds =: \kappa_{1,1} + \kappa_{1,2}.$$

Evidently,

$$\kappa_{1,1} = \int_{\eta_{k-1}}^{\eta_k} \left( \int_s^{\eta_{k+2}} w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{q'}} v^{p'}(s) ds \leq \lambda_k.$$

Hölder's inequality yields

$$g_k^p(\eta_k) \leq \int_{\eta_{k-1}}^{\eta_k} \left( \int_s^{\eta_{k+2}} w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{q'}} v^{p'}(s) ds \left( \int_{\eta_{k-1}}^{\eta_k} v^{p'}(y) dy \right)^{p-1}.$$



Since

$$\int_{\eta_k}^{\eta_k^*} |\Omega'_k(s)|^p v^{-p}(s) ds = \left( \int_{\eta_k}^{\eta_k^*} v^{p'}(y) dy \right)^{1-p},$$

then

$$\kappa_{1,2} \leq \lambda_k \left( \int_{\eta_k^-}^{\eta_k} v^{p'}(y) dy \right)^{p-1} \left( \int_{\eta_k}^{\eta_k^*} v^{p'}(y) dy \right)^{1-p}.$$

If  $\eta_k^* = \eta_k^+$ , then  $\kappa_{1,2} \leq \lambda_k$  because of (2.54) and (4.9). If  $\eta_k^* = \eta_{k+1}$ , we observe that

$$\int_{\eta_k^-}^{\eta_k} v^{p'}(y) dy = \int_{\eta_k}^{\eta_k^+} v^{p'}(y) dy \leq \int_{\eta_k^-}^{\eta_{k+1}^+} v^{p'}(y) dy = 2 \int_{\eta_k}^{\eta_{k+1}} v^{p'}(y) dy, \quad (4.16)$$

therefore,  $\kappa_{1,2} \leq 2^{p-1} \lambda_k$ . Thus, (4.15) is proved for  $i = 1$ . Since  $h_k^{(1)}(\eta_{k+1}) = h_k(\eta_{k+1})$ , the other cases of (4.15) follow from the first. Now, on the strength of

$$\begin{aligned} g_k(t) &\geq \left( \int_t^{\eta_{k+2}} w^q(x) dx \right)^{\frac{r}{pq}} \int_{\eta_{k-1}}^t \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{pq}} v^{p'}(s) ds \\ &\approx \left( \int_t^{\eta_{k+2}} w^q(x) dx \right)^{\frac{r}{pq}} \left( \int_{\eta_{k-1}}^t v^{p'}(y) dy \right)^{\frac{r}{p'q}}, \quad t \in [\eta_{k-1}, \eta_{k+2}], \end{aligned}$$

the lower bound

$$\begin{aligned} \|h_k w\|_q^q &\geq \int_{\eta_{k-1}}^{\eta_{k+2}} |g_k(t)|^q w^q(t) dt \\ &\gg \int_{\eta_{k-1}}^{\eta_{k+2}} \left( \int_t^{\eta_{k+2}} w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{\eta_{k-1}}^t v^{p'}(y) dy \right)^{\frac{r}{p'}} w^q(t) dt \end{aligned}$$

holds and integration by parts brings

$$\|h_k w\|_q^q \gg \lambda_k. \quad (4.17)$$

Now we construct the functions

$$h_i = \sum_{|k| \leq N} h_{k,i} = \sum_{|k| \leq N} h_{2k,i} + \sum_{|k| \leq N} h_{2k+1,i} =: F_{1,i} + F_{2,i},$$

where  $N \in \mathbb{N}$ . For each  $i = 1, 2, 3$  the supports of  $h_{2k,i}$ ,  $k \in \mathbb{Z}$ , are mutually disjoint. Therefore,  $F_{1,i} \in \mathcal{L}$  and by the same reason  $F_{2,i} \in \mathcal{L}$ . Observe that

$$1 \leq \sum_{|k| \leq N} \chi_{\text{supp } h_k}(x) \leq 4, \quad x \in \bigcup_{|k| \leq N} \text{supp } h_k.$$

Letting

$$\Lambda_N := \sum_{|k| \leq N} \lambda_k \in (0, \infty)$$

and using (4.14) – (4.17) and (4.10) we find

$$\begin{aligned} \Lambda_N^{1/q} &\ll \left\| \sum_{|k| \leq N} h_k w \right\|_q \leq \sum_{i=1}^3 \sum_{j=1}^2 \|F_{j,i} w\|_q \leq C \sum_{i=1}^3 \sum_{j=1}^2 \|F'_{j,i}/v\|_p \\ &\ll C \left( \sum_{|k| \leq N, i} \|h'_{k,i}/v\|_p^p \right)^{\frac{1}{p}} \ll C \Lambda_N^{1/p}. \end{aligned}$$

It implies  $C \gg \Lambda_N^{1/r}$ . Letting  $N \rightarrow \infty$  we obtain  $C \gg \sum_k \lambda_k$  and because of definition (4.13) we have

$$\begin{aligned} \lambda_k &\gg \int_{\eta_k}^{\eta_{k+1}} \left( \int_s^{\eta_{k+2}} w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{p'}} w^q(s) ds \\ &\geq \int_{\eta_k}^{\eta_{k+1}} \left( \int_{\delta^+(s)} w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{\Delta^-(s)} v^{p'}(y) dy \right)^{\frac{r}{p'}} w^q(s) ds. \end{aligned}$$

Then the required estimate  $C \gg \mathcal{B}_{MR}^+$  follows from (2.54) and (4.9). The inequality  $C \gg \mathcal{B}_{MR}^-$  can be proved by a similar construction using the intervals formed by the upper boundary function  $b(x)$ . Thus, the estimate (4.12) is true. Consequently, inequality  $\|\mathcal{H}\| \gg \mathcal{B}_{MR}$  is proved in the case when the fairway  $\sigma(x) = x$ .

Now we set free from this constraint. Let

$$\tilde{f}(t) = f(\sigma(t))[\sigma'(t)]^{1/p}, \quad \tilde{a}(x) = \sigma^{-1}(a(x)), \quad \tilde{b}(x) = \sigma^{-1}(b(x)). \quad (4.18)$$

By change of variables in the left and right hand sides of

$$\|\mathcal{H}f\|_q \leq \|\mathcal{H}\| \|f\|_p \quad (4.19)$$

it follows from the inequality of the form

$$\left( \int_0^\infty w^q(x) \left| \int_{\tilde{a}(x)}^{\tilde{b}(x)} \tilde{f}(y) \tilde{v}(y) dy \right|^q dx \right)^{\frac{1}{q}} \leq \|\mathcal{H}\| \left( \int_0^\infty |\tilde{f}(y)|^p dy \right)^{\frac{1}{p}}, \quad (4.20)$$

where  $\tilde{v}(y) = v(\sigma(y))[\sigma'(y)]^{1/p'}$ . It is easy to see, that

$$\int_{\tilde{a}(x)}^x \tilde{v}^{p'}(y) dy = \int_{a(x)}^{\sigma(x)} v^{p'}(y) dy = \int_{\sigma(x)}^{b(x)} v^{p'}(y) dy = \int_x^{\tilde{b}(x)} \tilde{v}^{p'}(y) dy, \quad (4.21)$$


therefore the fairway for  $\tilde{a}$ ,  $\tilde{b}$  and  $\tilde{v}$  is  $\tilde{\sigma}(x) = x$ . It has already been proved above that  $\tilde{\mathcal{B}}_{MR} \ll \|\mathcal{H}\|$ , where

$$\tilde{\mathcal{B}}_{MR}^r = \int_0^\infty \left( \int_{\tilde{\delta}(t)} w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{\tilde{\Delta}(t)} \tilde{v}^{p'}(y) dy \right)^{\frac{r}{p'}} w^q(t) dt.$$

Because of  $\tilde{\Delta}(t) := [\tilde{a}(t), \tilde{b}(t)]$ ,  $\tilde{\delta}(t) := [\tilde{b}^{-1}(\tilde{\sigma}(t)), \tilde{a}^{-1}(\tilde{\sigma}(t))] = [b^{-1}(\sigma(t)), a^{-1}(\sigma(t))]$  =  $\delta(t)$  and  $\int_{\tilde{\Delta}(t)} \tilde{v}^{p'}(y) dy = \int_{\delta(t)} v^{p'}(y) dy$  the equality  $\tilde{\mathcal{B}}_{MR} = \mathcal{B}_{MR}$  follows. The assertion about compactness for  $q < p$  is a direct corollary of obtained criterion of the boundedness and Ando's theorem.

## 4.2. Tomaselli and Persson-Stepanov type criteria

Now we give complete analogs of the alternative boundedness conditions for the operator (1.1) similar to (2.5) and (2.7).



Let  $\sigma^{-1}(y)$  denote the inverse function to the fairway  $\sigma(x)$  and put

$$\Delta(t) = (a(t), b(t)),$$

$$\theta(t) = (\sigma^{-1}(a(t)), \sigma^{-1}(b(t))).$$

**THEOREM 4.2.** *Let the hypotheses of Theorem 4.1 hold. If  $1 < p \leq q < \infty$ , then*

$$\|\mathcal{H}\|_{L_p \rightarrow L_q} \approx \mathcal{A}_T, \quad (4.22)$$

where

$$\mathcal{A}_T := \sup_{t>0} \left( \int_{\theta(t)} \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{-\frac{1}{p}}.$$

If  $0 < q < p < \infty$ ,  $p > 1$ ,  $1/r = 1/q - 1/p$ , then

$$\|\mathcal{H}\|_{L_p \rightarrow L_q} \approx \mathcal{B}_{PS}, \quad (4.23)$$

where

$$\mathcal{B}_{PS} := \left( \int_0^\infty \left[ \int_{\theta(t)} \left\{ \int_{\Delta(x)} v^{p'}(y) dy \right\}^q w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{\Delta(t)} v^{p'}(y) dy \right]^{q-\frac{r}{p}} w^q(t) dt \right)^{\frac{1}{r}}.$$

*Proof. The upper bounds.* Given boundary functions  $a(x)$  and  $b(x)$  satisfying the conditions (1.2) let a point sequence  $\{\xi_k\}_{k \in \mathbb{Z}} \subset (0, \infty)$  be given by relation (3.5) and put as before

$$\xi_k^- = \sigma^{-1}(a(\xi_k)), \quad \xi_k^+ = \sigma^{-1}(b(\xi_k)), \quad \Delta_k = [\xi_k^-, \xi_k^+] = \Delta_k^- \cup \Delta_k^+,$$

$$\Delta_k^- = [\xi_k^-, \xi_k), \quad \Delta_k^+ = [\xi_k, \xi_k^+), \quad k \in \mathbb{Z}.$$

Note that in view of (2.54)

$$\frac{1}{2} \int_{\Delta(t)} v^{p'}(y) dy \leq \int_{a(\xi_k)}^{b(t)} v^{p'}(y) dy \leq \int_{\Delta(t)} v^{p'}(y) dy \quad \text{for } t \in \Delta_k^- \quad (4.24)$$

and

$$\frac{1}{2} \int_{\Delta(t)} v^{p'}(y) dy \leq \int_{a(t)}^{b(\xi_k)} v^{p'}(y) dy \leq \int_{\Delta(t)} v^{p'}(y) dy \quad \text{for } t \in \Delta_k^+. \quad (4.25)$$

As in Theorem 4.1 we split the operator  $\mathcal{H}$  into the sum of four sequences of operators

$$\mathcal{H}f(x) = T_{k,1}f(x) + T_{k,2}f(x) + S_{k,1}f(x) + S_{k,2}f(x), \quad x \in \Delta_k, \quad (4.26)$$

then

$$\begin{aligned} \|\mathcal{H}f\|_q^q &= \sum_k \|\mathcal{H}f\|_{L_q(\Delta_k)}^q \approx \sum_k \|T_{k,1}f\|_{L_q(\Delta_k^-)}^q + \sum_k \|S_{k,2}f\|_{L_q(\Delta_k^-)}^q \\ &\quad + \sum_k \|T_{k,2}f\|_{L_q(\Delta_k^+)}^q + \sum_k \|S_{k,1}f\|_{L_q(\Delta_k^+)}^q. \end{aligned} \quad (4.27)$$

Let us start with trapeze-shaped operators  $S_{k,2}$  and  $T_{k,2}$ . If  $1 < p \leq q < \infty$ , we find by using the estimate (2.35) of Lemma 2.1

$$\begin{aligned} \|S_{k,2}\|_{L_p(a(\xi_k), b(\xi_k)) \rightarrow L_q(\Delta_k^-)}^q &\approx \sup_{t \in \Delta_k^-} \left( \int_{\xi_k^-}^t \left[ \int_{a(\xi_k)}^{b(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right) \left( \int_{a(\xi_k)}^{b(t)} v^{p'}(y) dy \right)^{-\frac{q}{p}}. \end{aligned}$$

For any  $t \in [\xi_k^-, \xi_k]$  we have  $(\xi_k^-, t) \subset \theta(t)$ , and  $(a(\xi_k), b(x)) \subset \Delta(x)$  for  $x \in (\xi_k^-, t) \subset (\xi_k^-, \xi_k)$ . Therefore, in view of (4.24),

$$\begin{aligned} \|S_{k,2}\|_{L_p(a(\xi_k), b(\xi_k)) \rightarrow L_q(\Delta_k^-)}^q &\ll \sup_{t \in \Delta_k^-} \left( \int_{\theta(t)} \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right) \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{-\frac{q}{p}} \leq \mathcal{A}_T^q. \end{aligned} \quad (4.28)$$

Analogously, by applying (2.41) of Lemma 2.2 and (4.25) we obtain

$$\begin{aligned} \|T_{k,2}\|_{L_p(a(\xi_k), b(\xi_k)) \rightarrow L_q(\Delta_k^+)}^q &\ll \sup_{t \in \Delta_k^+} \left( \int_{\theta(t)} \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right) \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{-\frac{q}{p}} \ll \mathcal{A}_T^q. \end{aligned} \quad (4.29)$$

If  $0 < q < p < \infty$ ,  $p > 1$  we have by applying the estimate (2.44) of Lemma 2.3

$$\begin{aligned} \|S_{k,2}\|_{L_p(a(\xi_k), b(\xi_k)) \rightarrow L_q(\Delta_k^-)}^p &\approx \int_{\xi_k^-}^{\xi_k} \left( \int_{\xi_k^-}^t \left[ \int_{a(\xi_k)}^{b(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{a(\xi_k)}^{b(t)} v^{p'}(y) dy \right)^{q-\frac{r}{p}} w^q(t) dt. \end{aligned}$$

For  $t \in [\xi_k^-, \xi_k]$  it yields that  $(\xi_k^-, t) \subset \theta(t)$ ,  $(a(\xi_k), b(t)) \subset \Delta(t)$  and also  $(a(\xi_k), b(x)) \subset \Delta(x)$  for any  $x \in (\xi_k^-, t) \subset (\xi_k^-, \xi_k)$ . Moreover,  $a(\xi_k) \leq \sigma(t)$  for  $t \in [\xi_k^-, \xi_k]$ . Therefore, in view of (4.24)

$$\begin{aligned} \|S_{k,2}\|_{L_p(a(\xi_k), b(\xi_k)) \rightarrow L_q(\Delta_k^-)}^r &\leq \int_{\xi_k^-}^{\xi_k} \left( \int_{\theta(t)} \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{p}} \\ &\quad \times \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^q \left( \int_{\sigma(t)}^{b(t)} v^{p'}(y) dy \right)^{-\frac{r}{p}} w^q(t) dt \\ &\stackrel{(2.54)}{\leq} 2^{\frac{r}{p}} \int_{\xi_k^-}^{\xi_k} \left( \int_{\theta(t)} \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{p}} \\ &\quad \times \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{q-\frac{r}{p}} w^q(t) dt \\ &=: \left( \mathcal{B}_{\Delta_k^-} \right)^r. \end{aligned} \quad (4.30)$$

Analogously, by applying (2.52) of Lemma 2.4 and (4.25) we can get that

$$\begin{aligned} \|T_{k,2}\|_{L_p(a(\xi_k), b(\xi_k)) \rightarrow L_q(\Delta_k^+)}^r &\ll \int_{\xi_k}^{\xi_k^+} \left( \int_{\theta(t)} \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{q-\frac{r}{p}} w^q(t) dt \\ &=: \left( \mathcal{B}_{\Delta_k^+} \right)^r. \end{aligned} \quad (4.31)$$

To estimate the norms  $\|T_{k,1}\|_{L_p(a(\xi_k^-), a(\xi_k)) \rightarrow L_q(\Delta_k^-)}^r$  and  $\|S_{k,1}\|_{L_p(b(\xi_k), b(\xi_k^+)) \rightarrow L_q(\Delta_k^+)}^r$  of triangle-shaped operators we introduce sequences  $\{t_i\}_{i=0}^{i_b(k)-1}$  and  $\{s_j\}_{j=0}^{j_a(k)-1}$  according to the constructions of Lemmas 2.5 and 2.6 with  $c = \xi_k$ . Then the operators  $T_{k,1}f(x)$ ,  $x \in \Delta_k^-$ , and  $S_{k,1}f(x)$ ,  $x \in \Delta_k^+$ , split into the following two sums:

$$\begin{aligned} T_{k,1}f(x) &= \sum_{i=0}^{i_b(k)-1} T_{k,1}^{(i)}f(x) := \sum_{i=0}^{i_b(k)-1} [T_{k,1}f(x) \chi_{(t_i, t_{i+1})}(x)], \\ S_{k,1}f(x) &= \sum_{j=0}^{j_a(k)-1} S_{k,1}^{(j)}f(x) := \sum_{j=0}^{j_a(k)-1} [S_{k,1}f(x) \chi_{(s_j, s_{j+1})}(x)]. \end{aligned} \quad (4.32)$$

If  $1 < p \leq q < \infty$  we have for  $T_{k,1}^{(i)}$ ,  $0 \leq i \leq i_b - 1$ , by applying the estimate (2.40) of Lemma 2.2

$$\begin{aligned} \|T_{k,1}^{(i)}\|_{L_p(a(t_i), a(\xi_k)) \rightarrow L_q(t_i, t_{i+1})}^q &\approx \sup_{t_i \leq t \leq t_{i+1}} \left( \int_{t_i}^t w^q(x) dx \right) \left( \int_{a(t)}^{a(\xi_k)} v^{p'}(y) dy \right)^{\frac{q}{p'}} \\ &\leq \sup_{t_i \leq t \leq t_{i+1}} \left( \int_{t_i}^t w^q(x) dx \right) \left( \int_{a(\xi_k^-)}^{\sigma(\xi_k^-)} v^{p'}(y) dy \right)^{\frac{q}{p'}} \end{aligned}$$

$$\begin{aligned}
 & \stackrel{(2.54)}{=} \left( \int_{t_i}^{t_{i+1}} w^q(x) dx \right) \left( \int_{a(\xi_k)}^{b(\xi_k^-)} v^{p'}(y) dy \right)^{\frac{q}{p'}} \\
 & \stackrel{(2.58)}{=} \left( \frac{1}{2^i} \right)^{\frac{q}{p'}} \left( \int_{t_i}^{t_{i+1}} w^q(x) dx \right) \left( \int_{a(\xi_k)}^{b(t_i)} v^{p'}(y) dy \right)^{\frac{q}{p'}}.
 \end{aligned}$$

Since  $q/p' = q - q/p$  and  $(t_i, t) \subseteq (\xi_k^-, t) \subset \theta(t)$  for  $t \in [\xi_k^-, \xi_k]$  we get in view of (2.55) that

$$\begin{aligned}
 & \|T_{k,1}^{(i)}\|_{L_p(a(t_i), a(\xi_k)) \rightarrow L_q(t_i, t_{i+1})}^q \\
 & \stackrel{(2.55)}{\ll} 2^{\frac{q}{p}} \left( \frac{1}{2^i} \right)^{\frac{q}{p'}} \left( \int_{t_i}^{t_{i+1}} w^q(x) dx \right) \left( \int_{a(\xi_k)}^{b(t_i)} v^{p'}(y) dy \right)^q \left( \int_{a(\xi_k)}^{b(t_{i+1})} v^{p'}(y) dy \right)^{-\frac{q}{p}} \\
 & \leq 2^{\frac{q}{p}} \left( \frac{1}{2^i} \right)^{\frac{q}{p'}} \sup_{t_i \leq t \leq t_{i+1}} \left( \int_{t_i}^t \left[ \int_{a(\xi_k)}^{b(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right) \left( \int_{a(\xi_k)}^{b(t)} v^{p'}(y) dy \right)^{-\frac{q}{p}} \\
 & \stackrel{(4.24)}{\ll} 2^{\frac{q}{p}} \left( \frac{1}{2^i} \right)^{\frac{q}{p'}} \sup_{t_i \leq t \leq t_{i+1}} \left( \int_{\theta(t)} \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right) \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{-\frac{q}{p}} \\
 & \ll \left( \frac{1}{2^i} \right)^{\frac{q}{p'}} \mathcal{A}_T^q.
 \end{aligned} \tag{4.33}$$

Hence, since  $p \leq q$  it holds that

$$\begin{aligned}
 \|T_{k,1} f\|_{L_q(\Delta_k^-)}^q &= \sum_{i=0}^{i_b-1} \|T_{k,1}^{(i)} f\|_{L_q(t_i, t_{i+1})}^q \leq \sum_{i=0}^{i_b-1} \|T_{k,1}^{(i)}\|_{L_p(a(t_i), a(\xi_k)) \rightarrow L_q(t_i, t_{i+1})}^q \|f\|_{L_p(a(t_i), a(\xi_k))}^q \\
 & \stackrel{(4.33)}{\ll} \mathcal{A}_T^q \|f\|_{L_p(a(\xi_k^-), a(\xi_k))}^q \sum_{i=0}^{i_b-1} \left( \frac{1}{2^i} \right)^{\frac{q}{p'}} \leq \frac{2^{q/p'}}{2^{q/p'} - 1} \mathcal{A}_T^q \|f\|_{L_p(a(\xi_k^-), a(\xi_k))}^q.
 \end{aligned} \tag{4.34}$$

Analogously, from (2.34) of Lemma 2.1 by using (2.65), (2.63) and (4.25) we can get that

$$\|S_{k,1} f\|_{L_q(\Delta_k^+)}^q \ll \frac{2^{q/p'}}{2^{q/p'} - 1} \mathcal{A}_T^q \|f\|_{L_p(b(\xi_k), b(\xi_k^+))}^q. \tag{4.35}$$

If  $0 < q < p < \infty$ ,  $p > 1$  we have for  $T_{k,1}^{(i)}$ ,  $0 \leq i \leq i_b - 1$ , by applying the estimate (2.50) of Lemma 2.4

$$\begin{aligned}
 \|T_{k,1}^{(i)}\|_{L_p(a(t_i), a(\xi_k)) \rightarrow L_q(t_i, t_{i+1})}^r &\approx \int_{t_i}^{t_{i+1}} \left( \int_{t_i}^t w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{a(t)}^{a(\xi_k)} v^{p'}(y) dy \right)^{\frac{r}{p'}} w^q(t) dt \\
 &\leq \left( \int_{a(\xi_k^-)}^{\sigma(\xi_k^-)} v^{p'}(y) dy \right)^{\frac{r}{p'}} \int_{t_i}^{t_{i+1}} \left( \int_{t_i}^t w^q(x) dx \right)^{\frac{r}{p}} w^q(t) dt
 \end{aligned}$$

$$\begin{aligned}
& \stackrel{(2.54)}{=} \left( \int_{a(\xi_k)}^{b(\xi_k^-)} v^{p'}(y) dy \right)^{\frac{r}{p'}} \int_{t_i}^{t_{i+1}} \left( \int_{t_i}^t w^q(x) dx \right)^{\frac{r}{p}} w^q(t) dt \\
& \stackrel{(2.58)}{=} \left( \frac{1}{2^i} \right)^{\frac{r}{p'}} \left( \int_{a(\xi_k)}^{b(t_i)} v^{p'}(y) dy \right)^{\frac{r}{p'}} \int_{t_i}^{t_{i+1}} \left( \int_{t_i}^t w^q(x) dx \right)^{\frac{r}{p}} w^q(t) dt.
\end{aligned}$$

By using the relation  $r/p' = q \cdot r/p + q - r/p$  we write

$$\begin{aligned}
& \|T_{k,1}^{(i)}\|_{L_p(a(t_i), a(\xi_k)) \rightarrow L_q(t_i, t_{i+1})}^r \\
& \ll \left( \frac{1}{2^i} \right)^{\frac{r}{p'}} \left( \int_{a(\xi_k)}^{b(t_i)} v^{p'}(y) dy \right)^{q \cdot \frac{r}{p} + q - \frac{r}{p}} \int_{t_i}^{t_{i+1}} \left( \int_{t_i}^t w^q(x) dx \right)^{\frac{r}{p}} w^q(t) dt \\
& \stackrel{(2.55)}{=} 2^{\frac{r}{p}} \left( \frac{1}{2^i} \right)^{\frac{r}{p'}} \left( \int_{a(\xi_k)}^{b(t_i)} v^{p'}(y) dy \right)^{q \cdot \frac{r}{p} + q} \left( \int_{a(\xi_k)}^{b(t_{i+1})} v^{p'}(y) dy \right)^{-\frac{r}{p}} \\
& \quad \times \int_{t_i}^{t_{i+1}} \left( \int_{t_i}^t w^q(x) dx \right)^{\frac{r}{p}} w^q(t) dt \\
& \leq 2^{\frac{r}{p}} \left( \frac{1}{2^i} \right)^{\frac{r}{p'}} \int_{t_i}^{t_{i+1}} \left( \int_{t_i}^t \left[ \int_{a(\xi_k)}^{b(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{a(\xi_k)}^{b(t)} v^{p'}(y) dy \right)^q \\
& \quad \times \left( \int_{a(\xi_k)}^{b(t)} v^{p'}(y) dy \right)^{-\frac{r}{p}} w^q(t) dt \\
& \stackrel{(4.24)}{\approx} \left( \frac{1}{2^i} \right)^{\frac{r}{p'}} \int_{t_i}^{t_{i+1}} \left( \int_{t_i}^t \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{q - \frac{r}{p}} w^q(t) dt.
\end{aligned}$$

As before  $(t_i, t) \subseteq (\xi_k^-, t) \subset \theta(t)$  for  $t \in [\xi_k^-, \xi_k]$ . Therefore, we have

$$\begin{aligned}
& \|T_{k,1}^{(i)}\|_{L_p(a(t_i), a(\xi_k)) \rightarrow L_q(t_i, t_{i+1})}^r \\
& \ll \left( \frac{1}{2^i} \right)^{\frac{r}{p'}} \int_{t_i}^{t_{i+1}} \left( \int_{\theta(t)} \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{r}{p}} \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{q - \frac{r}{p}} w^q(t) dt \\
& =: \left( \frac{1}{2^i} \right)^{\frac{r}{p'}} \left( \mathcal{B}_{\Delta_k^-}^{(i)} \right)^r.
\end{aligned} \tag{4.36}$$

By applying Hölder's inequality with the powers  $r/q$  and  $p/q$  we get that

$$\begin{aligned}
\|T_{k,1} f\|_{L_q(\Delta_k^-)}^q &= \sum_{i=0}^{i_b-1} \|T_{k,1}^{(i)} f\|_{L_q(t_i, t_{i+1})}^q \\
&\leq \sum_{i=0}^{i_b-1} \|T_{k,1}^{(i)}\|_{L_p(a(t_i), a(\xi_k)) \rightarrow L_q(t_i, t_{i+1})}^q \|f\|_{L_p(a(t_i), a(\xi_k))}^q \\
&\stackrel{(4.36)}{\ll} \|f\|_{L_p(a(\xi_k^-), a(\xi_k))}^q \sum_{i=0}^{i_b-1} \left( \frac{1}{2^i} \right)^{\frac{q}{p'}} \left( \mathcal{B}_{\Delta_k^-}^{(i)} \right)^q
\end{aligned}$$

$$\begin{aligned}
 &\leq \|f\|_{L_p(a(\xi_k^-), a(\xi_k))}^q \left( \sum_{i=0}^{i_b-1} \left( \mathcal{B}_{\Delta_k^-}^{(i)} \right)^r \right)^{\frac{q}{r}} \left( \sum_{i=0}^{i_b-1} \left( \frac{1}{2^i} \right)^{\frac{p}{p'}} \right)^{\frac{q}{p}} \\
 &\leq \left( \frac{2^{p-1}}{2^{p-1}-1} \right)^{\frac{q}{p}} \left( \mathcal{B}_{\Delta_k^-} \right)^q \|f\|_{L_p(a(\xi_k^-), a(\xi_k))}^q. \quad (4.37)
 \end{aligned}$$

Analogously, from (2.42) of Lemma 2.3 by using (4.25), (2.65) and (2.63) we can get that

$$\|S_{k,1}f\|_{L_q(\Delta_k^+)}^q \ll \left( \frac{2^{p-1}}{2^{p-1}-1} \right)^{\frac{q}{p}} \left( \mathcal{B}_{\Delta_k^+} \right)^q \|f\|_{L_p(b(\xi_k), b(\xi_k^+))}^q. \quad (4.38)$$

Since every series  $\{T_{k,1}\}$ ,  $\{T_{k,2}\}$ ,  $\{S_{k,1}\}$  or  $\{S_{k,2}\}$  is block-diagonal, we have the following pair of inequalities coming from Lemma 3.1 for each term in (4.27). Namely, if  $1 < p \leq q < \infty$  then

$$\sum_{k \in \mathbb{Z}} \|T_{k,1}f\|_{L_q(\Delta_k^-)}^q \stackrel{(4.34)}{\ll} \frac{2^{q/p'}}{2^{q/p'}-1} \mathcal{A}_T^q \sum_{k \in \mathbb{Z}} \|f\|_{L_p(a(\xi_k^-), a(\xi_k))}^q \leq \frac{2^{q/p'}}{2^{q/p'}-1} \mathcal{A}_T^q \|f\|_p^q.$$

If  $0 < q < p < \infty$ ,  $p > 1$  then

$$\sum_{k \in \mathbb{Z}} \|T_{k,1}f\|_{L_q(\Delta_k^-)}^q \stackrel{(4.37)}{\ll} \left( \sum_{k \in \mathbb{Z}} \left( \mathcal{B}_{\Delta_k^-} \right)^r \right)^{\frac{1}{r}} \|f\|_p \leq \mathcal{B}_{PS} \|f\|_p.$$

Analogous inequalities hold for  $\{T_{k,2}\}$ ,  $\{S_{k,1}\}$  and  $\{S_{k,2}\}$  on the strength of (4.29), (4.35), (4.28) for  $1 < p \leq q < \infty$  and (4.31), (4.38), (4.30) in the case  $0 < q < p < \infty$ ,  $p > 1$ , respectively. By combining these estimates with (4.27) we obtain the upper bounds in (4.22) and (4.23).

*The lower bound.* Let  $1 < p \leq q < \infty$ .

First we prove that  $\mathcal{A}_T \approx \mathcal{A}_{T,1} + \mathcal{A}_{T,2}$ , where  $\mathcal{A}_{T,1} := \sup_{t>0} \sup_{b^{-1}(\sigma(t)) \leq s \leq t} \mathcal{A}_{T,1}(s, t)$  and  $\mathcal{A}_{T,2} := \sup_{t>0} \sup_{t \leq s \leq a^{-1}(\sigma(t))} \mathcal{A}_{T,2}(t, s)$  with

$$\mathcal{A}_{T,1}(s, t) := \left( \int_s^t \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(s)}^{\sigma(t)} v^{p'}(y) dy \right)^{-\frac{1}{p}}$$

and

$$\mathcal{A}_{T,2}(t, s) := \left( \int_t^s \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{\sigma(t)}^{b(s)} v^{p'}(y) dy \right)^{-\frac{1}{p}}.$$

Indeed, in view of (2.54)

$$\frac{1}{2} \int_{a(s)}^{b(s)} v^{p'}(y) dy \leq \int_{a(s)}^{\sigma(t)} v^{p'}(y) dy \leq \int_{a(s)}^{b(s)} v^{p'}(y) dy, \quad b^{-1}(\sigma(t)) \leq s \leq t. \quad (4.39)$$

Therefore, it holds that

$$\mathcal{A}_{T,1} \approx \sup_{s>0} \left( \int_{a(s)}^{b(s)} v^{p'}(y) dy \right)^{-\frac{1}{p}} \sup_{s \leq t \leq \sigma^{-1}(b(s))} \left( \int_s^t \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}}$$



$$= \sup_{s>0} \left( \int_s^{\sigma^{-1}(b(s))} \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{\Delta(s)} v^{p'}(y) dy \right)^{-\frac{1}{p}}.$$

Analogously, because of (2.54)

$$\frac{1}{2} \int_{a(s)}^{b(s)} v^{p'}(y) dy \leq \int_{\sigma(t)}^{b(s)} v^{p'}(y) dy \leq \int_{a(s)}^{b(s)} v^{p'}(y) dy, \quad t \leq s \leq a^{-1}(\sigma(t)). \quad (4.40)$$

Thus,

$$\mathcal{A}_{T,2} \approx \sup_{s>0} \left( \int_{\sigma^{-1}(a(s))}^s \left[ \int_{\Delta(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{\Delta(s)} v^{p'}(y) dy \right)^{-\frac{1}{p}}.$$

Now suppose that  $\|\mathcal{H}\| < \infty$  and insert the function  $f_{s,t}(y) = v^{p'-1}(y) \chi_{[a(s), \sigma(t)]}(y)$ , where  $b^{-1}(\sigma(t)) \leq s \leq t$ , into the inequality

$$\left( \int_0^\infty (\mathcal{H}f)^q(x) dx \right)^{\frac{1}{q}} \leq \|\mathcal{H}\| \left( \int_0^\infty f^p(y) dy \right)^{\frac{1}{p}}.$$

Since  $a(s) \leq a(x)$  and  $\sigma(x) \leq \sigma(t)$  for  $s \leq x \leq t$ , we have in view of (2.54) that

$$\|\mathcal{H}\| \geq \frac{1}{2} \left( \int_s^t \left[ \int_{a(x)}^{b(x)} v^{p'}(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(s)}^{\sigma(t)} v^{p'}(y) dy \right)^{-\frac{1}{p}} \approx \mathcal{A}_{T,1}(s, t).$$

Therefore,  $\|\mathcal{H}\| \gg \mathcal{A}_{T,1}(s, t)$  for all  $s \leq t$  and, thus,  $\|\mathcal{H}\| \gg \mathcal{A}_{T,1}$ . Analogously, by applying the function  $f_{t,s}(y) = v^{p'-1}(y) \chi_{[\sigma(t), b(s)]}(y)$  with  $t \leq s \leq a^{-1}(\sigma(t))$  we get that  $\|\mathcal{H}\| \gg \mathcal{A}_{T,2}$ .

Let  $0 < q < p < \infty$ ,  $p > 1$ . First, similar to Theorem 4.1 we prove  $\|\mathcal{H}\| \gg \mathcal{B}_{PS}$  under the condition (4.9). Let us remind that in this case we denote

$$x^- = a(x), \quad x^+ = b(x), \quad \Delta(x) = [x^-, x^+]$$


and establish the estimate

$$C \gg \mathcal{B}_{PS}, \quad (4.41)$$

where  $C$  is the least possible constant of the inequality (4.10) which holds for the function class from Definition 4.1. For this purpose it is sufficient to prove that  $C \gg \mathcal{B}_{PS}^\pm$ , where

$$\mathcal{B}_{PS}^\pm := \left( \int_0^\infty \left[ \int_{\Delta^\pm(t)} \left\{ \int_{\Delta(x)} v^{p'}(y) dy \right\}^q w^q(x) dx \right]^{\frac{r}{p}} \left[ \int_{\Delta(t)} v^{p'}(y) dy \right]^{q-\frac{r}{p}} w^q(t) dt \right)^{\frac{1}{r}}$$

and  $\Delta^-(t) = (t^-, t)$ ,  $\Delta^+(t) = (t, t^+)$ .



Further we utilize in part notations from the proof of Theorem 4.1 with sometimes a different meaning. To prove  $C \gg \mathcal{B}_{PS}^-$  we put  $\eta_0 = 1$  and define the sequence  $\{\eta_k\}_{k \in \mathbb{Z}} \subset (0, \infty)$  by

$$\eta_{k+1} = a^{-1}(\eta_k), \quad k \in \mathbb{Z}.$$

Let

$$\eta_k^* = \min(\eta_k^+, \eta_{k+1}).$$

For a fixed  $k \in \mathbb{Z}$  we take four neighboring points  $\eta_{k-1}$ ,  $\eta_k$ ,  $\eta_{k+1}$ ,  $\eta_{k+1}^*$  and let

$$f_k(t) = \chi_{[\eta_{k-1}, \eta_{k+1}]}(t) [g_k(t) + h_k(t)],$$

where

$$g_k(t) = \int_{\eta_{k-1}}^t \left( \int_s^{\eta_{k+1}} w^q(x) dx \right)^{\frac{r}{pq}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{pq}} v^{p'}(s) ds,$$

$$h_k(t) = \left( \int_{\eta_{k-1}}^{\eta_{k+1}} v^{p'}(y) dy \right)^{-\frac{r}{pq}} \left( \int_{\eta_{k-1}}^{\eta_{k+1}} \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{pq}} \left( \int_{\eta_{k-1}}^t v^{p'}(s) ds \right).$$

On the interval  $[\eta_{k-1}, \eta_{k+1}^*]$  we define the function

$$\phi_k(t) = \begin{cases} f_k(t), & t \in [\eta_{k-1}, \eta_{k+1}], \\ f_k(\eta_{k+1}) \Omega_{k+1}(t), & t \in [\eta_{k+1}, \eta_{k+1}^*], \end{cases}$$

where

$$\Omega_l(t) = \left( \int_{\eta_l}^{\eta_l^*} v^{p'}(y) dy \right)^{-1} \int_t^{\eta_l^*} v^{p'}(s) ds.$$

Put

$$v_k := \lambda_k + \mu_k,$$

where

$$\lambda_k := \int_{\eta_{k-1}}^{\eta_{k+1}} \left( \int_s^{\eta_{k+1}} w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{q}} v^{p'}(s) ds,$$

$$\mu_k := \left( \int_{\eta_{k-1}}^{\eta_{k+1}} v^{p'}(y) dy \right)^{-\frac{r}{p}} \left( \int_{\eta_{k-1}}^{\eta_{k+1}} \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{q}}.$$

Now we find a decomposition

$$\phi_k(t) = \sum_{i=1}^2 \phi_{k,i}(t) \quad (4.42)$$

with  $\phi_{k,i} \in \mathcal{L}$ ,  $i = 1, 2$ . To this end we first determine  $\phi_{k,1}(t)$  by

$$\phi_{k,1}(t) = \begin{cases} 0, & t \notin [\eta_{k-1}, \eta_k^*], \\ \phi_k(t), & t \in [\eta_{k-1}, \eta_k], \\ \phi_k(\eta_k)\Omega_k(t), & t \in [\eta_k, \eta_k^*]. \end{cases}$$

Since  $\eta_k^- = \eta_{k-1}$ , it yields that  $\text{supp } \phi_{k,1} \subseteq [\eta_k^-, \eta_k^*] \subseteq [\eta_k^-, \eta_k^+]$  and, hence,  $\phi_{k,1} \in \mathcal{L}$ . Obviously that  $\phi_{k,2}(t) = \phi_k(t) - \phi_{k,1}(t)$  is in  $\mathcal{L}$ . It is clear that (4.42) holds. Our next step is to prove that

$$\|\phi'_{k,i}/v\|_p^p \ll v_k, \quad i = 1, 2. \quad (4.43)$$

If

$$\kappa_1 := \|\phi'_{k,1}/v\|_p^p = \int_{\eta_{k-1}}^{\eta_k} |\phi'_k(s)|^p v^{-p}(s) ds + f_k^p(\eta_k) \int_{\eta_k}^{\eta_k^*} |\Omega'_k(s)|^p v^{-p}(s) ds =: \kappa_{1,1} + \kappa_{1,2},$$

then

$$\begin{aligned} \kappa_{1,1} &= \int_{\eta_{k-1}}^{\eta_k} \left( \int_s^{\eta_{k+1}} w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{q'}} v^{p'}(s) ds \\ &\quad + \left( \int_{\eta_{k-1}}^{\eta_{k+1}} v^{p'}(y) dy \right)^{-\frac{r}{q}} \left( \int_{\eta_{k-1}}^{\eta_{k+1}} \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{q}} \\ &\quad \times \left( \int_{\eta_{k-1}}^{\eta_k} v^{p'}(s) ds \right) \leq \lambda_k + \mu_k = v_k. \end{aligned}$$

By Hölder's inequality,

$$\begin{aligned} f_k^p(\eta_k) &\approx \left( \int_{\eta_{k-1}}^{\eta_k} \left[ \int_s^{\eta_{k+1}} w^q(x) dx \right]^{\frac{r}{pq}} \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^{\frac{r}{pq'}} v^{p'}(s) ds \right)^p \\ &\quad + \left( \int_{\eta_{k-1}}^{\eta_{k+1}} v^{p'}(y) dy \right)^{-\frac{r}{q}} \left( \int_{\eta_{k-1}}^{\eta_{k+1}} \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^{\eta_k} v^{p'}(s) ds \right)^p \\ &\leq \int_{\eta_{k-1}}^{\eta_k} \left( \int_s^{\eta_{k+1}} w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{q'}} v^{p'}(s) ds \left( \int_{\eta_{k-1}}^{\eta_k} v^{p'}(s) ds \right)^{p-1} \\ &\quad + \mu_k \left( \int_{\eta_{k-1}}^{\eta_k} v^{p'}(s) ds \right)^{p-1} \\ &\leq (\lambda_k + \mu_k) \left( \int_{\eta_{k-1}}^{\eta_k} v^{p'}(s) ds \right)^{p-1} = v_k \left( \int_{\eta_{k-1}}^{\eta_k} v^{p'}(s) ds \right)^{p-1}. \end{aligned}$$

Since

$$\int_{\eta_k}^{\eta_k^*} |\Omega'_k(s)|^p v^{-p}(s) ds = \left( \int_{\eta_k}^{\eta_k^*} v^{p'}(s) ds \right)^{1-p},$$

we have that

$$\kappa_{1,2} \ll v_k \left( \int_{\eta_{k-1}}^{\eta_k} v^{p'}(s) ds \right)^{p-1} \left( \int_{\eta_k}^{\eta_k^*} v^{p'}(s) ds \right)^{1-p}$$

and in view of (2.54), (4.9) and (4.16) it holds that  $\kappa_{1,2} \ll v_k$ . Thus, (4.43) is proved for  $i = 1$ . Since  $\phi_{k,2}(\eta_{k+1}) = \phi_k(\eta_{k+1})$ , we can write for the second case of (4.43) that

$$\begin{aligned} \kappa_2 &:= \|\phi'_{k,2}/v\|_p^p = \int_{\eta_k}^{\eta_{k+1}} |\phi'_{k,2}(s)|^p v^{-p}(s) ds + \phi_k^p(\eta_{k+1}) \int_{\eta_{k+1}}^{\eta_{k+1}^*} |\Omega'_{k+1}(s)|^p v^{-p}(s) ds \\ &= \int_{\eta_k}^{\eta_{k+1}} |\phi'_k(s) - \phi'_{k,1}(s)|^p v^{-p}(s) ds + \phi_k^p(\eta_{k+1}) \int_{\eta_{k+1}}^{\eta_{k+1}^*} |\Omega'_{k+1}(s)|^p v^{-p}(s) ds \\ &\ll \int_{\eta_k}^{\eta_{k+1}} |\phi'_k(s)|^p v^{-p}(s) ds + f_k^p(\eta_k) \int_{\eta_k}^{\eta_{k+1}} \chi_{[\eta_k, \eta_k^*]}(s) |\Omega'_k(s)|^p v^{-p}(s) ds \\ &\quad + f_k^p(\eta_{k+1}) \int_{\eta_{k+1}}^{\eta_{k+1}^*} |\Omega'_{k+1}(s)|^p v^{-p}(s) ds =: \kappa_{2,1} + \kappa_{2,2} + \kappa_{2,3}. \end{aligned}$$

Obviously,  $\kappa_{2,2} = \kappa_{1,2} \ll v_k$ . Similar to  $\kappa_{1,1}$  and  $\kappa_{1,2}$  we can prove that  $\kappa_{2,1} + \kappa_{2,3} \ll v_k$ . Therefore, (4.43) is true for  $i = 2$  also.

Now, since

$$\begin{aligned} g_k(t) &\geq \left( \int_t^{\eta_{k+1}} w^q(x) dx \right)^{\frac{r}{pq}} \int_{\eta_{k-1}}^t \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{pq'}} v^{p'}(s) ds \\ &\approx \left( \int_t^{\eta_{k+1}} w^q(x) dx \right)^{\frac{r}{pq}} \left( \int_{\eta_{k-1}}^t v^{p'}(y) dy \right)^{\frac{r}{p'q}}, \quad t \in [\eta_{k-1}, \eta_{k+1}], \end{aligned}$$

the lower bound

$$\|\phi_k w\|_q^q \gg v_k \quad (4.44)$$

follows. Next we construct the functions

$$F_i = \sum_{|k| \leq N} \phi_{k,i} = \sum_{|k| \leq N} \phi_{2k,i} + \sum_{|k| \leq N} \phi_{2k+1,i} =: F_{1,i} + F_{2,i},$$

where  $N \in \mathbb{N}$ . For every  $i = 0, 1$  the supports of  $\phi_{2k,i}$ ,  $k \in \mathbb{Z}$ , are mutually disjoint. Therefore,  $F_{1,i} \in \mathcal{L}$  and by the same reason  $F_{2,i} \in \mathcal{L}$ . Observe that

$$2 \leq \sum_{|k| \leq N} \chi_{\text{supp} \phi_k}(x) \leq 3, \quad x \in \bigcup_{|k| \leq N} \text{supp} \phi_k.$$

On the strength of (4.42), (4.43) and (4.44) we find that

$$\begin{aligned} \left( \sum_{|k| \leq N} v_k \right)^{1/q} &\ll \left\| \sum_{|k| \leq N} \phi_k w \right\|_q \leq \sum_{i=1}^2 \sum_{j=1}^2 \|F_{j,i} w\|_q \leq C \sum_{i=1}^2 \sum_{j=1}^2 \|F'_{j,i}/v\|_p \\ &\ll C \left( \sum_{|k| \leq N, i} \|\phi'_{k,i}/v\|_p^p \right)^{1/p} \ll C \left( \sum_{|k| \leq N} v_k \right)^{1/p}. \end{aligned} \quad (4.45)$$

Therefore, by letting  $N \rightarrow \infty$  we obtain  $C \gg (\sum_{k \in \mathbb{Z}} v_k)^{1/r}$ . Now we put

$$\tilde{\lambda}_k := \int_{\eta_{k-1}}^{\eta_{k+1}} \left( \int_{\eta_{k-1}}^t \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^t v^{p'}(y) dy \right)^{-\frac{r}{q}} v^{p'}(t) dt,$$

$$\lambda_k^* := \int_{\eta_{k-1}}^{\eta_{k+1}} \left( \int_{\eta_{k-1}}^t \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{p}} \left( \int_{\eta_{k-1}}^t v^{p'}(y) dy \right)^{q - \frac{r}{p}} w^q(t) dt.$$

Write

$$\tilde{\lambda}_k = \int_{\eta_{k-1}}^{\eta_{k+1}} \left( \int_{\eta_{k-1}}^t \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^q d \left[ - \int_s^{\eta_{k+1}} w^q(x) dx \right] \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^t v^{p'}(y) dy \right)^{-\frac{r}{q}} v^{p'}(t) dt,$$

and note that

$$\begin{aligned} & \int_{\eta_{k-1}}^t \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^q d \left( - \int_s^{\eta_{k+1}} w^q(x) dx \right) \\ & \leq q \int_{\eta_{k-1}}^t \left( \int_s^{\eta_{k+1}} w^q(x) dx \right) \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{q-1} v^{p'}(s) ds \\ & \approx \int_{\eta_{k-1}}^t \left\{ \left( \int_s^{\eta_{k+1}} w^q(x) dx \right) \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{q-1 + \frac{q}{2p}} \right\} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{-\frac{q}{2p}} v^{p'}(s) ds \end{aligned}$$

[applying Hölder's inequality with the powers  $r/q$  and  $p/q$ ]

$$\begin{aligned} & \leq \left( \int_{\eta_{k-1}}^t \left[ \int_s^{\eta_{k+1}} w^q(x) dx \right]^{\frac{r}{q}} \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^{\frac{r}{q} + \frac{r}{2p}} v^{p'}(s) ds \right)^{\frac{q}{r}} \\ & \quad \times \left( \int_{\eta_{k-1}}^t \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^{-\frac{1}{2}} v^{p'}(s) ds \right)^{\frac{q}{p}}. \end{aligned}$$

This implies

$$\begin{aligned} \tilde{\lambda}_k & \ll \int_{\eta_{k-1}}^{\eta_{k+1}} \int_{\eta_{k-1}}^t \left( \int_s^{\eta_{k+1}} w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{q} + \frac{r}{2p}} v^{p'}(s) ds \\ & \quad \times \left( \int_{\eta_{k-1}}^t v^{p'}(y) dy \right)^{\frac{r}{2p} - \frac{r}{q}} v^{p'}(t) dt \\ & = \int_{\eta_{k-1}}^{\eta_{k+1}} \left( \int_s^{\eta_{k+1}} w^q(x) dx \right)^{\frac{r}{q}} \left( \int_{\eta_{k-1}}^s v^{p'}(y) dy \right)^{\frac{r}{q} + \frac{r}{2p}} v^{p'}(s) ds \\ & \quad \times \int_s^{\eta_{k+1}} \left( \int_{\eta_{k-1}}^t v^{p'}(y) dy \right)^{\frac{r}{2p} - \frac{r}{q}} v^{p'}(t) dt \ll \lambda_k. \end{aligned}$$

Observe that

$$\begin{aligned} \tilde{\lambda}_k & = \frac{p}{r} \int_{\eta_{k-1}}^{\eta_{k+1}} \left( \int_{\eta_{k-1}}^t \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{q}} d \left( - \left[ \int_{\eta_{k-1}}^t v^{p'}(y) dy \right]^{-\frac{r}{p}} \right) \\ & = -\frac{p}{r} \mu_k + \frac{p}{q} \lambda_k^*. \end{aligned}$$

Therefore,

$$\lambda_k^* = \frac{q}{p} \tilde{\lambda}_k + \frac{q}{r} \mu_k,$$

and, hence,

$$\lambda_k^* \ll v_k.$$

This implies  $C \gg (\sum_{k \in \mathbb{Z}} v_k)^{1/r} \gg (\sum_{k \in \mathbb{Z}} \lambda_k^*)^{1/r}$ . Note that

$$\int_{\eta_{k-1}}^t v^{p'}(y) dy \leq \int_{\eta_{k-1}}^{t^+} v^{p'}(y) dy \leq 2 \int_{\eta_k}^{t^+} v^{p'}(y) dy \quad (4.46)$$

and

$$\frac{1}{2} \int_{\Delta(t)} v^{p'}(y) dy \leq \int_{\eta_k}^{t^+} v^{p'}(y) dy \leq \int_{\Delta(t)} v^{p'}(y) dy \quad (4.47)$$

for any  $t \in [\eta_k, \eta_{k+1}]$ . Therefore, in view of (2.54) and (4.9) we obtain that

$$\begin{aligned} \lambda_k^* &\geq \int_{\eta_k}^{\eta_{k+1}} \left( \int_{\eta_{k-1}}^t \left[ \int_{\eta_{k-1}}^s v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{p}} \left( \int_{\eta_{k-1}}^t v^{p'}(y) dy \right)^{q-\frac{r}{p}} w^q(t) dt \\ &\stackrel{(4.46)}{\geq} 2^{-\frac{r}{p}} \int_{\eta_k}^{\eta_{k+1}} \left( \int_{\Delta^-(t)} \left[ \int_{\Delta^-(s)} v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{p}} \\ &\quad \times \left( \int_{\Delta^-(t)} v^{p'}(y) dy \right)^q \left( \int_{\eta_k}^{t^+} v^{p'}(y) dy \right)^{-\frac{r}{p}} w^q(t) dt \\ &\stackrel{(4.47)}{\gg} \int_{\eta_k}^{\eta_{k+1}} \left( \int_{\Delta^-(t)} \left[ \int_{\Delta(s)} v^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{p}} \left( \int_{\Delta(t)} v^{p'}(y) dy \right)^{q-\frac{r}{p}} w^q(t) dt. \end{aligned}$$

Hence, the required estimate  $C \gg \mathcal{B}_{PS}^-$  is proved. The estimate  $C \gg \mathcal{B}_{PS}^+$  can be proved similarly with intervals formed by the fairway  $\sigma(x)$  and the upper boundary function  $b(x)$ . Thus, the estimate (4.41) holds and, therefore, the inequality  $\|\mathcal{H}\| \gg \mathcal{B}_{PS}$  is proved in the case when the fairway  $\sigma(x) = x$ . The general case follows similar to the proof of Theorem 4.1. By changing variables (4.18) in both hand sides of the inequality (4.19) we arrive to the inequality (4.20). Since (4.21) is true the fairway-function for  $\tilde{a}$ ,  $\tilde{b}$  and  $\tilde{v}$  is just  $\tilde{\sigma}(x) = x$ . Therefore, in view of the obtained estimate  $\tilde{\mathcal{B}}_{PS} \ll \|\mathcal{H}\|$  with

$$\tilde{\mathcal{B}}_{PS}^r = \int_0^\infty \left( \int_{\tilde{\Delta}(t)} \left[ \int_{\tilde{\Delta}(s)} \tilde{v}^{p'}(y) dy \right]^q w^q(s) ds \right)^{\frac{r}{p}} \left( \int_{\tilde{\Delta}(t)} \tilde{v}^{p'}(y) dy \right)^{q-\frac{r}{p}} w^q(t) dt$$

and because of  $\tilde{\Delta}(t) = [\tilde{a}(t), \tilde{b}(t)] = \theta(t)$ ,  $\int_{\tilde{\Delta}(t)} \tilde{v}^{p'}(y) dy = \int_{\Delta(t)} v^{p'}(y) dy$  we get  $\tilde{\mathcal{B}}_{PS} = \mathcal{B}_{PS}$ .

## 5. Geometric Steklov operator

In this section we study the geometric Steklov operator

$$\mathcal{G}f(x) := \exp \left( \frac{1}{b(x) - a(x)} \int_{a(x)}^{b(x)} \log f(y) dy \right), \quad f(y) \geq 0,$$

acting in weighted Lebesgue spaces with the border functions  $a(x)$  and  $b(x)$  satisfying the conditions (1.2). This operator is closely related to the Hardy operator (1.1) because of Jensen's inequality

$$[\mathcal{G}(fv)](x)w(x) \leq \frac{\mathcal{H}f(x)}{b(x) - a(x)}, \quad (5.1)$$

from which the upper bound for the weighted  $L_p - L_q$  boundedness of  $\mathcal{G}$  easy follows if  $L_p - L_q$  boundedness of  $\mathcal{H}$  has a suitable characterization.

**THEOREM 5.1.** *Let  $0 < p \leq q < \infty$  and the operator  $\mathcal{G}$  be given by (1.3) with the boundary functions  $a(x)$  and  $b(x)$  satisfying the condition (1.2). Then for the "norm"*

$$\|\mathcal{G}\|_{L_{p,v} \rightarrow L_{q,w}} := \sup_{f \geq 0} \frac{\|(\mathcal{G}f)w\|_q}{\|fv\|_p} \quad \text{the estimate}$$

$$\|\mathcal{G}\|_{L_{p,v} \rightarrow L_{q,w}} \approx \sup_{t > 0} \left( \int_{\sigma_0^{-1}(a(t))}^{\sigma_0^{-1}(b(t))} u^q(x) dx \right)^{\frac{1}{q}} [b(t) - a(t)]^{-\frac{1}{p}}, \quad (5.2)$$

holds.

If  $0 < q < p < \infty$  then

$$\|\mathcal{G}\|_{L_{p,v} \rightarrow L_{q,w}} \approx \left( \int_0^\infty \left[ \int_{\sigma_0^{-1}(a(t))}^{\sigma_0^{-1}(b(t))} u^q(s) ds \right]^{\frac{r}{p}} [b(t) - a(t)]^{-\frac{r}{p}} u^q(t) dt \right)^{\frac{1}{r}} =: \mathcal{B}_{\mathcal{G}}, \quad (5.3)$$

where

$$\sigma_0(t) := \frac{a(t) + b(t)}{2}, \quad u(t) := (\mathcal{G}v^{-1})(t)w(t). \quad (5.4)$$

*Proof.* The proof of the estimate (5.2) is due to L.-E. Persson and D.V. Prokhorov. We prove the estimate (5.3).

*The upper bound.* Let  $0 < q < p < \infty$  and  $\mathcal{B}_{\mathcal{G}} < \infty$ . According to Jensen's inequality (5.1) we can get the upper bound in (5.3) by using Theorem 4.2 with suitable summation parameters and proper weight functions. Indeed, since  $\mathcal{G}(f^s) = (\mathcal{G}f)^s$  and  $\mathcal{G}(f \cdot g) = (\mathcal{G}f) \cdot (\mathcal{G}g)$  the equalities

$$\|\mathcal{G}\|_{L_{p,v} \rightarrow L_{q,w}} = \|\mathcal{G}\|_{L_{p,1} \rightarrow L_{q,u}} = \|\mathcal{G}\|_{L_{\frac{p}{s},1}^{\frac{1}{s}} \rightarrow L_{\frac{q}{s},u^s}^{\frac{1}{s}}} \quad (5.5)$$

hold with  $u$  defined by (5.4). Note that for  $\mathcal{H}: L_{\bar{p},1} \rightarrow L_{\bar{q},\bar{u}}$  with  $\bar{u} = u^s$ ,  $\bar{p} = p/s$  and  $\bar{q} = q/s$  the corresponding parameter  $\bar{r}$  is equal to  $r/s$  and, respectively,  $\bar{r}/\bar{p} = r/p$ . Moreover, the condition  $\bar{p} > 1$  of Theorem 4.2 is satisfied if we put  $0 < s < q < p < \infty$ . Therefore, according to (5.1) and Theorem 4.2 with  $v = 1$  and  $w = u^s$  we have for any  $0 < 1 < q/s < p/s < \infty$  that

$$\|\mathcal{G}\|_{L_{\frac{p}{s},1}^{\frac{1}{s}} \rightarrow L_{\frac{q}{s},u^s}^{\frac{1}{s}}} \leq \|\mathcal{H}\|_{L_{\frac{p}{s}}^{\frac{1}{s}} \rightarrow L_{\frac{q}{s}}^{\frac{1}{s}}} \ll \mathcal{B}_{\mathcal{G}}. \quad (5.6)$$

Now, in view of (5.5) the upper bound in (5.3) is proved.

*The lower bound.* Suppose that  $\|\mathcal{G}\|_{L_{p,v} \rightarrow L_{q,w}} = \|\mathcal{G}\|_{L_{p,1} \rightarrow L_{q,u}} = \|\mathcal{G}\|_{L_{\tilde{p},1} \rightarrow L_{\tilde{q},\tilde{u}}}^{1/s} < \infty$ , where  $\tilde{u} = u^s$  and  $\tilde{p} = p/s$ ,  $\tilde{q} = q/s$  for any  $0 < q < p < s < \infty$ . In other words the inequality

$$\|(\mathcal{G}f)\tilde{u}\|_{\tilde{q}} \leq C\|f\|_{\tilde{p}} \quad (5.7)$$

holds for all  $f \in L_{\tilde{p}}$  and  $0 < \tilde{q} < \tilde{p} < 1$  with  $C < \infty$  independent on  $f$ . Now the fairway-function  $\sigma(x)$  is such that  $b(x) - \sigma(x) = \sigma(x) - a(x)$ . Let us remind that as before we denote

$$x^- = \sigma^{-1}(a(x)), \quad x^+ = \sigma^{-1}(b(x)), \quad \theta(x) = [x^-, x^+].$$


Similarly to the sections 4.1 and 4.2 we prove that

$$\left( \int_0^\infty \left[ \int_{\theta^\pm(x)} \tilde{u}^{\tilde{q}}(s) ds \right]^{\tilde{p}} [b(x) - a(x)]^{-\tilde{p}} \tilde{u}^{\tilde{q}}(x) dx \right)^{\frac{1}{\tilde{r}}} =: (\tilde{\mathcal{B}}_{\mathcal{G}})^\pm \ll \|\mathcal{G}\|_{L_{\tilde{p},1} \rightarrow L_{\tilde{q},\tilde{u}}},$$

where  $\theta^-(x) = (x^-, x)$ ,  $\theta^+(x) = (x, x^+)$  and  $1/\tilde{r} = 1/\tilde{q} - 1/\tilde{p}$ . To prove the inequality with  $(\tilde{\mathcal{B}}_{\mathcal{G}})^-$  we put  $\xi_0 = 1$  and define the sequence  $\{\xi_k\} \subset (0, \infty)$  as before such that

$$\xi_k = (a^{-1} \circ b)^k(1), \quad k \in \mathbb{Z}.$$

Then  $\xi_k < \xi_k^+ = \xi_{k+1}^- < \xi_{k+1}$ . Moreover,  $\bigcup_k [\xi_k, \xi_{k+1}] = (0, \infty)$  and  $\bigcup_k [\sigma(\xi_k), \sigma(\xi_{k+1})] = (0, \infty)$ .



Further, put  $\xi_k^+ = \xi_{k+1}^- = m_k^0$  and construct on every segment  $[\xi_k, \xi_{k+1}]$  the sequence  $\{m_k^j\}$  with  $-j_a(k) \leq j \leq j_b(k)$  by the following way: for  $-j_a(k) \leq j \leq 0$  we use the construction of Lemma 2.7 with  $v = 1$  and  $d = \xi_k$ , while for  $0 \leq j \leq j_b(k)$  we use Lemma 2.5 with  $v = 1$  and  $c = \xi_{k+1}$ . Actually we have the sequence  $\{m_k^j\}$ ,  $-j_a(k) \leq j \leq j_b(k)$ , defined by:

- (1)  $m_k^{-j_a(k)} = \xi_k$ ,  $m_k^{j_b(k)} = \xi_{k+1}$ ;
- (2) if  $(\xi_k^+)^- \leq \xi_k$  then  $j_a(k) = 1$ ;  $j_b(k) = 1$  if  $[\mathcal{N}^-(k)] = 0$  or  $\mathcal{N}^-(k) = [\mathcal{N}^-(k)] = 1$ ;
- (3) if  $(\xi_k^+)^- > \xi_k$ , then  $j_a(k) > 1$  and  $m_k^{j-1} = (m_k^j)^-$ , where  $(m_k^j)^- > \xi_k$  and  $-j_a(k) + 2 \leq j \leq 0$ ;



(4) if  $[\mathcal{N}^-(k)] > 0$ , then the points  $m_k^j$  for  $1 \leq j \leq [\mathcal{N}^-(k)]$  are taken so that

$$b(m_k^j) - a(\xi_{k+1}) = 2 [b(m_k^{j-1}) - a(\xi_{k+1})]. \quad (5.8)$$

Then we have  $\bigcup_k \bigcup_{j=-j_a(k)}^{j_b(k)-1} [m_k^j, m_k^{j+1}) = (0, \infty)$  and the following useful properties of  $b(t) - a(t)$  coming from Lemmas 2.5 and 2.7:

1°) if  $t \in (m_k^j, m_k^{j+1})$ , then

$$b(t) - a(t) \approx b(m_k^{j+1}) - a(m_k^j) \quad (5.9)$$

and

$$b(m_k^{j+1}) - a(m_k^j) \approx b(m_k^j) - a(m_k^j) \approx b(m_k^{j+1}) - a(m_k^{j+1}); \quad (5.10)$$

2°) if  $t \in [x^-, x]$ , then

$$b(t) - a(t) \approx b(x) - a(x^-) \approx b(x) - a(x) \quad \text{for} \quad \xi_k \leq x^- < t < x \leq \xi_k^+, \quad (5.11)$$

and

$$b(t) - a(t) < b(x) - a(x^-) \approx b(x) - a(x); \quad (5.12)$$

3°) if  $0 \leq j \leq j_b(k) - 2$ , then

$$b(m_k^{j+i}) - a(\xi_{k+1}) = 2^i [b(m_k^j) - a(\xi_{k+1})] \quad \text{for some } i \in \{1, \dots, j_b(k) - j - 1\}. \quad (5.13)$$

Now we apply in (5.7) a test function of the form  $f(y) = \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} f_{k,j}(y)$ , where

$$f_{k,j}(y) = \chi_{(a(m_k^j), b(m_k^{j+1}))}(y) l_{k,j}$$

and

$$l_{k,j} := \left( \int_{(m_k^j)^-}^{m_k^{j+1}} \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{pq}} [b(m_k^{j+1}) - a(m_k^j)]^{-\frac{\tilde{r}}{pq}}.$$

Then we have

$$\begin{aligned} \|(\mathcal{G}f)\tilde{u}\|_{\tilde{q}}^{\tilde{q}} &= \int_0^\infty \left( \exp \frac{1}{b(x) - a(x)} \int_{a(x)}^{b(x)} \log f(y) dy \right)^{\tilde{q}} \tilde{u}^{\tilde{q}}(x) dx \\ &= \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} \int_{m_k^j}^{m_k^{j+1}} \left( \exp \frac{1}{b(x) - a(x)} \int_{a(x)}^{b(x)} \log \left[ \sum_{n \in \mathbb{Z}} \sum_{i=-j_a(n)}^{j_b(n)-1} f_{n,i}(y) \right] dy \right)^{\tilde{q}} \\ &\quad \times \tilde{u}^{\tilde{q}}(x) dx \\ &\geq \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} l_{k,j}^{\tilde{q}} \int_{m_k^j}^{m_k^{j+1}} \tilde{u}^{\tilde{q}}(x) dx \end{aligned}$$

$$\begin{aligned}
 & \stackrel{(5.9)}{\gg} \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} \int_{m_k^j}^{m_k^{j+1}} \left( \int_{(m_k^j)^-}^x \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} (b(x) - a(x))^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(x) dx \\
 & =: \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} \gamma_{k,j}.
 \end{aligned} \tag{5.14}$$

On the other hand in view of  $\tilde{p} < 1$  we have that

$$\begin{aligned}
 \|f\|_{\tilde{p}}^{\tilde{p}} &= \sum_{k \in \mathbb{Z}} \int_{\sigma(\xi_k)}^{\sigma(\xi_{k+1})} \left( \sum_{n \in \mathbb{Z}} \sum_{i=-j_a(n)}^{j_b(n)-1} f_{n,i}(y) \right)^{\tilde{p}} dy \leq \sum_{k \in \mathbb{Z}} \int_{\sigma(\xi_k)}^{\sigma(\xi_{k+1})} \left( \sum_{n \in \mathbb{Z}} \sum_{i=-j_a(n)}^{j_b(n)-1} f_{n,i}^{\tilde{p}}(y) \right) dy \\
 &= \sum_{k \in \mathbb{Z}} \int_{\sigma(\xi_k)}^{\sigma(\xi_{k+1})} \left( \sum_{n \in \mathbb{Z}} \sum_{i=-j_a(n)}^{j_b(n)-1} \chi_{(a(m_n^i), b(m_n^{i+1}))}(y) l_{n,i}^{\tilde{p}} \right) dy.
 \end{aligned}$$

Denote

$$M_{n,i}(k) := (\sigma(\xi_k), \sigma(\xi_{k+1})) \cap (a(m_n^i), b(m_n^{i+1}))$$

and observe that  $M_{n,i}(k) = \emptyset$  if  $n \leq k-2$  and  $n \geq k+2$ , while  $M_{n,i}(k) \subseteq (a(m_n^i), b(m_n^{i+1}))$  for the rest  $k-1 \leq n \leq k+1$  and all corresponding them  $-i_a(n) \leq i \leq i_b(n)-1$ . Therefore,

$$\begin{aligned}
 \|f\|_{\tilde{p}}^{\tilde{p}} &\leq \sum_{k \in \mathbb{Z}} \int_{\sigma(\xi_k)}^{\sigma(\xi_{k+1})} \left( \sum_{n=k-1}^{k+1} \sum_{i=-j_a(n)}^{j_b(n)-1} \chi_{M_{n,i}(k)}(y) l_{n,i}^{\tilde{p}} \right) dy \\
 &= \sum_{k \in \mathbb{Z}} \int_{\sigma(\xi_k)}^{\sigma(\xi_{k+1})} \left( \sum_{j=-j_a(k-1)}^{j_b(k-1)-1} \chi_{M_{k-1,j}(k)}(y) l_{k-1,j}^{\tilde{p}} + \sum_{j=-j_a(k)}^{j_b(k)-1} \chi_{M_{k,j}(k)}(y) l_{k,j}^{\tilde{p}} \right. \\
 &\quad \left. + \sum_{j=-j_a(k+1)}^{j_b(k+1)-1} \chi_{M_{k+1,j}(k)}(y) l_{k+1,j}^{\tilde{p}} \right) dy \leq \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k-1)}^{j_b(k-1)-1} l_{k-1,j}^{\tilde{p}} [b(m_{k-1}^{j+1}) - a(m_{k-1}^j)] \\
 &\quad + \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} l_{k,j}^{\tilde{p}} [b(m_k^{j+1}) - a(m_k^j)] + \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k+1)}^{j_b(k+1)-1} l_{k+1,j}^{\tilde{p}} [b(m_{k+1}^{j+1}) - a(m_{k+1}^j)] \\
 &= \sum_{k \in \mathbb{Z}} \left( \sum_{j=-j_a(k-1)}^{j_b(k-1)-1} \tau_{k-1,j} + \sum_{j=-j_a(k)}^{j_b(k)-1} \tau_{k,j} + \sum_{j=-j_a(k+1)}^{j_b(k+1)-1} \tau_{k+1,j} \right) \\
 &\leq 3 \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} \tau_{k,j},
 \end{aligned} \tag{5.15}$$

where

$$\tau_{k,j} := \left( \int_{(m_k^j)^-}^{m_k^{j+1}} \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{q}} [b(m_k^{j+1}) - a(m_k^j)]^{-\frac{\tilde{r}}{p}}$$

and

$$\tau_{k,j} = \frac{q}{r} [b(m_k^{j+1}) - a(m_k^j)]^{-\frac{\tilde{r}}{p}} \int_{(m_k^j)^-}^{m_k^{j+1}} \left( \int_{(m_k^j)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt$$

$$\begin{aligned}
&= \frac{q}{r} [b(m_k^{j+1}) - a(m_k^j)]^{-\frac{\tilde{r}}{p}} \int_{(m_k^j)^-}^{m_k^j} \left( \int_{(m_k^j)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&\quad + \frac{q}{r} [b(m_k^{j+1}) - a(m_k^j)]^{-\frac{\tilde{r}}{p}} \int_{m_k^j}^{m_k^{j+1}} \left( \int_{(m_k^j)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&=: I_{k,j} + II_{k,j}.
\end{aligned} \tag{5.16}$$

On the strength of (5.9)  $II_{k,j} \approx \gamma_{k,j}$  and we can write that

$$\|f\|_{\tilde{p}}^{\tilde{p}} \ll \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} [I_{k,j} + \gamma_{k,j}]. \tag{5.17}$$

Now we need to estimate  $\sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} I_{k,j}$  by  $\sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} \gamma_{k,j}$ . Let  $j = -j_a(k)$ , then  $(m_k^{-j_a(k)})^- = m_{k-1}^0$  and

$$\begin{aligned}
I_{k,-j_a(k)} &= [b(m_k^{-j_a(k)+1}) - a(m_k^{-j_a(k)})]^{-\frac{\tilde{r}}{p}} \int_{\xi_k^-}^{\xi_k} \left( \int_{\xi_k^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&\stackrel{(5.10)}{\approx} [b(\xi_k) - a(\xi_k)]^{-\frac{\tilde{r}}{p}} \int_{\xi_k^-}^{\xi_k} \left( \int_{\xi_k^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&\stackrel{(5.12)}{\ll} \int_{\xi_k^-}^{\xi_k} \left( \int_{\xi_k^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&\leq \sum_{j=0}^{j_b(k-1)-1} \int_{m_{k-1}^j}^{m_{k-1}^{j+1}} \left( \int_{(m_{k-1}^j)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&= \sum_{j=0}^{j_b(k-1)-1} \gamma_{k-1,j}.
\end{aligned} \tag{5.18}$$

Let  $j = 0$ , then we have if  $j_a(k) = 1$  that

$$\begin{aligned}
I_{k,0} &= [b(m_k^1) - a(m_k^0)]^{-\frac{\tilde{r}}{p}} \int_{(m_k^0)^-}^{m_k^0} \left( \int_{(m_k^0)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&\stackrel{(5.10), (5.12)}{\ll} \int_{(m_k^0)^-}^{m_k^0} \left( \int_{(m_k^0)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&\leq \sum_{j=0}^{j_b(k-1)-1} \int_{m_{k-1}^j}^{m_{k-1}^{j+1}} \left( \int_{(m_{k-1}^j)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&\quad + \int_{m_{k-1}^0}^{m_k^0} \left( \int_{(m_{k-1}^0)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt
\end{aligned}$$

$$\leq \sum_{j=0}^{j_b(k-1)-1} \gamma_{k-1,j} + \gamma_{k,-1}. \quad (5.19)$$

If  $j_a(k) > 1$  and  $-j_a(k) + 1 \leq j \leq 0$ , then

$$\sum_{j=-j_a(k)+1}^0 I_{k,j} = I_{k,-j_a(k)+1} + \sum_{j=-j_a(k)+2}^0 I_{k,j}. \quad (5.20)$$

Analogously to (5.19),

$$I_{k,-j_a(k)+1} \ll \sum_{j=0}^{j_b(k-1)-1} \gamma_{k-1,j} + \gamma_{k,-j_a(k)}, \quad (5.21)$$

while

$$\sum_{j=-j_a(k)+2}^0 I_{k,j} \leq \sum_{j=-j_a(k)+2}^0 \gamma_{k,j-1} = \sum_{j=-j_a(k)+1}^{-1} \gamma_{k,j}. \quad (5.22)$$

Finally we get from (5.18) – (5.22) that

$$\sum_{j=-j_a(k)}^0 I_{k,j} \ll 2 \sum_{j=0}^{j_b(k-1)-1} \gamma_{k-1,j} + \sum_{j=-j_a(k)}^{-1} \gamma_{k,j} \leq 2 \left( \sum_{j=0}^{j_b(k-1)-1} \gamma_{k-1,j} + \sum_{j=-j_a(k)}^{j_b(k)-1} \gamma_{k,j} \right). \quad (5.23)$$

Further, if  $j_b(k) = 1$ , the required estimate follows from (5.23). Let  $j_b(k) > 1$  and  $1 \leq j \leq j_b(k) - 1$ . Since  $(m_k^0)^- < (m_k^j)^- < m_k^0 < m_k^j$  we have for  $1 \leq j \leq j_b(k) - 1$  that

$$\begin{aligned} I_{k,j} &\leq \left[ b(m_k^{j+1}) - a(m_k^j) \right]^{-\frac{\tilde{r}}{p}} \int_{(m_k^0)^-}^{m_k^0} \left( \int_{(m_k^0)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\ &+ \left[ b(m_k^{j+1}) - a(m_k^j) \right]^{-\frac{\tilde{r}}{p}} \sum_{i=1}^j \int_{m_k^{i-1}}^{m_k^i} \left( \int_{(m_k^i)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt =: I'_{k,j} + I''_{k,j}. \end{aligned}$$

To estimate  $I'_{k,j}$  observe that  $b(m_k^j) - a(m_k^j) \approx b(m_k^j) - a(\xi_{k+1})$  and in view of (5.10) and (5.13)

$$b(m_k^{j+1}) - a(m_k^j) \approx b(m_k^j) - a(m_k^j) \approx b(m_k^j) - a(\xi_{k+1}) = 2^j [b(m_k^0) - \sigma(m_k^0)].$$

Analogously to (5.19) we have for each  $1 \leq j \leq j_b(k) - 1$  that

$$\begin{aligned} I'_{k,j} &\ll 2^{-\frac{\tilde{r}}{p}} [b(m_k^0) - \sigma(m_k^0)]^{-\frac{\tilde{r}}{p}} \int_{(m_k^0)^-}^{m_k^0} \left( \int_{(m_k^0)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\ &\stackrel{(5.12)}{\ll} 2^{-\frac{\tilde{r}}{p}} \int_{(m_k^0)^-}^{m_k^0} \left( \int_{(m_k^0)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \end{aligned}$$

$$\begin{aligned}
&\leq 2^{-\frac{j\tilde{r}}{p}} \left[ \sum_{i=0}^{j_b(k-1)-1} \int_{m_{k-1}^i}^{m_k^{i+1}} \left( \int_{(m_{k-1}^i)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \right. \\
&\quad \left. + \sum_{n=-j_a(k)}^{-1} \int_{m_k^n}^{m_k^{n+1}} \left( \int_{(m_k^n)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \right] \\
&= 2^{-\frac{j\tilde{r}}{p}} \left[ \sum_{i=0}^{j_b(k-1)-1} \gamma_{k-1,i} + \sum_{n=-j_a(k)}^{-1} \gamma_{k,n} \right]. \tag{5.24}
\end{aligned}$$

Therefore,

$$\begin{aligned}
\sum_{j=1}^{j_b(k)-1} I'_{k,j} &\ll \left[ \sum_{j=1}^{j_b(k)-1} 2^{-\frac{j\tilde{r}}{p}} \right] \left[ \sum_{i=0}^{j_b(k-1)-1} \gamma_{k-1,i} + \sum_{n=-j_a(k)}^{-1} \gamma_{k,n} \right] \\
&\leq \frac{2^{\tilde{r}/\tilde{p}}}{2^{\tilde{r}/\tilde{p}} - 1} \left[ \sum_{i=0}^{j_b(k-1)-1} \gamma_{k-1,i} + \sum_{n=-j_a(k)}^{-1} \gamma_{k,n} \right]. \tag{5.25}
\end{aligned}$$

Further, in view of (5.10),

$$I''_{k,j} \ll [b(m_k^j) - a(m_k^j)]^{-\frac{\tilde{r}}{p}} \sum_{i=1}^j \int_{m_k^{i-1}}^{m_k^i} \left( \int_{(m_k^{i-1})^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt. \tag{5.26}$$

Therefore, on the strength of (5.13) and the equivalence  $b(m_k^j) - a(m_k^j)$  to  $b(m_k^j) - a(\xi_{k+1})$  we get

$$\begin{aligned}
\sum_{j=1}^{j_b(k)-1} I''_{k,j} &\ll \sum_{j=1}^{j_b(k)-1} [b(m_k^j) - a(\xi_{k+1})]^{-\frac{\tilde{r}}{p}} \sum_{i=1}^j \int_{m_k^{i-1}}^{m_k^i} \left( \int_{(m_k^{i-1})^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&= [b(m_k^1) - a(\xi_{k+1})]^{-\frac{\tilde{r}}{p}} \int_{m_k^0}^{m_k^1} \left( \int_{(m_k^1)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&\quad + [b(m_k^2) - a(\xi_{k+1})]^{-\frac{\tilde{r}}{p}} \left( \left[ \int_{m_k^0}^{m_k^1} + \int_{m_k^1}^{m_k^2} \right] \left[ \int_{(m_k^2)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right]^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \right) + \dots \\
&\quad \dots + [b(m_k^{j_b(k)-1}) - a(\xi_{k+1})]^{-\frac{\tilde{r}}{p}} \\
&\quad \times \left( \left[ \int_{m_k^0}^{m_k^1} + \dots + \int_{m_k^{j_b(k)-2}}^{m_k^{j_b(k)-1}} \right] \left[ \int_{(m_k^{j_b(k)-1})^-}^t \tilde{u}^{\tilde{q}}(s) ds \right]^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \right) \\
&\stackrel{(5.13)}{\leq} \left[ \sum_{j=0}^{j_b(k)-1} 2^{-\frac{j\tilde{r}}{p}} \right] [b(m_k^1) - a(\xi_{k+1})]^{-\frac{\tilde{r}}{p}} \int_{m_k^0}^{m_k^1} \left( \int_{(m_k^1)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
&\quad + \left[ \sum_{j=0}^{j_b(k)-1} 2^{-\frac{j\tilde{r}}{p}} \right] [b(m_k^2) - a(\xi_{k+1})]^{-\frac{\tilde{r}}{p}} \int_{m_k^1}^{m_k^2} \left( \int_{(m_k^2)^-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt + \dots
\end{aligned}$$

$$\begin{aligned}
 & \dots + \left[ \sum_{j=0}^{j_b(k)-1} 2^{-\frac{\tilde{r}}{p}} \right] \left[ b(m_k^{j_b(k)-1}) - a(\xi_{k+1}) \right]^{-\frac{\tilde{r}}{p}} \int_{m_k^{j_b(k)-2}}^{m_k^{j_b(k)-1}} \\
 & \times \left( \int_{(m_k^{j_b(k)-1})_-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \\
 (5.9), (5.10), (5.12) \quad & \ll \frac{2^{\tilde{r}/\tilde{p}}}{2^{\tilde{r}/\tilde{p}} - 1} \left[ \int_{m_k^0}^{m_k^1} \left( \int_{(m_k^1)_-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt + \dots \right. \\
 & \left. \dots + \int_{m_k^{j_b(k)-2}}^{m_k^{j_b(k)-1}} \left( \int_{(m_k^{j_b(k)-1})_-}^t \tilde{u}^{\tilde{q}}(s) ds \right)^{\frac{\tilde{r}}{p}} [b(t) - a(t)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(t) dt \right] \\
 & \leq \frac{2^{\tilde{r}/\tilde{p}}}{2^{\tilde{r}/\tilde{p}} - 1} \sum_{j=0}^{j_b(k)-1} \gamma_{k,j}. \tag{5.27}
 \end{aligned}$$

Now we have from (5.25) – (5.27) that

$$\sum_{j=1}^{j_b(k)-1} I_{k,j} \ll \frac{2^{\tilde{r}/\tilde{p}}}{2^{\tilde{r}/\tilde{p}} - 1} \left[ \sum_{j=0}^{j_b(k-1)-1} \gamma_{k-1,j} + \sum_{j=-j_a(k)}^{j_b(k)-1} \gamma_{k,j} \right]. \tag{5.28}$$

By combining (5.23) and (5.28) we conclude that

$$\begin{aligned}
 \sum_{j=-j_a(k)}^{j_b(k)-1} I_{k,j} & \ll \max \left\{ 2, \frac{2^{\tilde{r}/\tilde{p}}}{2^{\tilde{r}/\tilde{p}} - 1} \right\} \left[ \sum_{j=0}^{j_b(k-1)-1} \gamma_{k-1,j} + \sum_{j=-j_a(k)}^{j_b(k)-1} \gamma_{k,j} \right] \\
 & \leq \max \left\{ 2, \frac{2^{\tilde{r}/\tilde{p}}}{2^{\tilde{r}/\tilde{p}} - 1} \right\} \left[ \sum_{j=-j_a(k-1)}^{j_b(k-1)-1} \gamma_{k-1,j} + \sum_{j=-j_a(k)}^{j_b(k)-1} \gamma_{k,j} \right]. \tag{5.29}
 \end{aligned}$$

It follows from (5.17) and (5.29) that

$$\|f\|_{\tilde{p}}^{\tilde{p}} \ll \sum_{k \in \mathbb{Z}} \left[ \sum_{j=-j_a(k-1)}^{j_b(k-1)-1} \gamma_{k-1,j} + \sum_{j=-j_a(k)}^{j_b(k)-1} \gamma_{k,j} \right] \leq 2 \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} \gamma_{k,j}. \tag{5.30}$$

Finally, we have by combining (5.14) and (5.30) that

$$\begin{aligned}
 \|\mathcal{G}\|_{L_{\tilde{p},1} \rightarrow L_{\tilde{q},\tilde{u}}} & \gg \left( \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} \gamma_{k,j} \right)^{\frac{1}{r}} \\
 & \geq \left( \sum_{k \in \mathbb{Z}} \sum_{j=-j_a(k)}^{j_b(k)-1} \int_{m_k^j}^{m_k^{j+1}} \left[ \int_{\theta^-(x)} \tilde{u}^{\tilde{q}}(s) ds \right]^{\frac{\tilde{r}}{p}} [b(x) - a(x)]^{-\frac{\tilde{r}}{p}} \tilde{u}^{\tilde{q}}(x) dx \right)^{\frac{1}{r}} \\
 & = (\tilde{\mathcal{B}}_{\mathcal{G}})^-.
 \end{aligned}$$

Analogously we can prove the inequality

$$\|\mathcal{G}\|_{L_{\tilde{p},1} \rightarrow L_{\tilde{q},\tilde{u}}} \gg (\tilde{\mathcal{B}}_{\mathcal{G}})^+.$$

Hence, in view of the equivalence  $\mathcal{B}_{\mathcal{G}} \approx (\mathcal{B}_{\mathcal{G}})^- + (\mathcal{B}_{\mathcal{G}})^+$  and since  $\tilde{r}/\tilde{p} = r/p$  and  $\tilde{u}^{\tilde{q}} = u^q$  the lower bound for  $\|\mathcal{G}\|_{L_{p,v} \rightarrow L_{q,w}}$  is proved.

## 6. Applications

### 6.1. Embeddings

Let  $u$ ,  $v$  and  $w$  be non-negative measurable weight functions such that for  $0 < p, q, s < \infty$ ,  $p > 1$  the powers  $u^s$ ,  $w^q$ ,  $v^{-p}$  and  $v^{p'}$  are locally integrable on  $(0, \infty)$  and  $0 < u(x), v(x), w(x) < \infty$  for a.e.  $x \in \mathbb{R}^+$ . The Sobolev space  $W_{p,s}^1$  consists of all absolutely continuous functions on  $(0, \infty)$  such that

$$\|F\|_{W_{p,s}^1} := \|Fu\|_s + \|F'/v\|_p < \infty. \quad (6.1)$$

We assume for simplicity that  $W_{p,s}^1$  is a closure of the set of all finitely supported differentiable functions with respect to the norm (6.1), i.e.

$$W_{p,s}^1 = \overset{\circ}{W}_{p,s}^1. \quad (6.2)$$

In [15] necessary and sufficient conditions for  $u$  and  $v$  are pointed out under which (6.2) holds. In this case by arguing as in [15] it is possible to construct the boundary functions  $a(x)$  and  $b(x)$  satisfying (1.2) such that  $a(x) < x < b(x)$  and for all  $x \in (0, \infty)$

$$\int_{\Delta^-(x)} v^{p'}(y) dy = \int_{\Delta^+(x)} v^{p'}(y) dy, \quad (6.3)$$

$$\left( \int_{\Delta(x)} u^s(z) dz \right)^{\frac{1}{s}} \left( \int_{\Delta(x)} v^{p'}(y) dy \right)^{\frac{1}{p'}} = 1, \quad (6.4)$$

where  $\Delta(x) = [a(x), b(x)]$ ,  $\Delta^-(x) = [a(x), x]$ ,  $\Delta^+(x) = [x, b(x)]$ . The equality (6.3) says that the fairway  $\sigma(x) = x$ , and (6.4) guarantees inequalities

$$\left( \int_{\Delta(x)} |F(z)|^s u^s(z) dz \right)^{\frac{1}{s}} \leq 2 \left( \int_{\Delta(x)} |F'(y)|^p v^{-p}(y) dy \right)^{\frac{1}{p}} \quad (6.5)$$

for every absolutely continuous function  $F$  such that  $F(t) = 0$  for some  $t \in \Delta(x)$  and

$$\sup_{t \in \Delta(x)} |F(t)| \leq \left( \int_{\Delta(x)} v^{p'}(y) dy \right)^{\frac{1}{p'}} \left[ \left( \int_{\Delta(x)} |F(z)|^s u^s(z) dz \right)^{\frac{1}{s}} + \left( \int_{\Delta(x)} |F'(y)|^p v^{-p}(y) dy \right)^{\frac{1}{p}} \right],$$

which follow by applying Hölder's inequality.

Let  $\mathcal{L} \subset \overset{\circ}{W}_{p,s}^1$  be the set of functions satisfying Definition 4.1. Using (6.5) it is easy to see that for  $1 < p \leq s < \infty$

$$\|F\|_{W_{p,s}^1} \approx \|F'/v\|_p, \quad F \in \mathcal{L}. \quad (6.6)$$

Applying Theorem 4.1 we obtain the following assertion, where the constants  $\mathcal{A}_M$  and  $\mathcal{B}_{MR}$  are given by (4.2) and (4.4), respectively.

THEOREM 6.1. *Let  $1 < p \leq q < \infty$ ,  $s > 0$ . Then the inequality*

$$\|Fw\|_q \leq C \|F\|_{W_{p,s}^1}, \quad F \in \mathcal{L} \quad (6.7)$$

*is valid if and only if  $\mathcal{A}_M < \infty$ . Moreover, for the least constant  $C$  in (6.7) the equivalence  $C \approx \mathcal{A}_M$  holds. If  $0 < q < p < \infty$ ,  $s \geq p > 1$ , then (6.7) is valid if and only if  $\mathcal{B}_{MR} < \infty$ , moreover  $C \approx \mathcal{B}_{MR}$ .*

*Proof.* The upper bounds  $C \ll \mathcal{A}_M$  and  $C \ll \mathcal{B}_{MR}$  follow from (4.11) and the upper bounds for the norm  $\|\mathcal{H}\|$  established in Theorem 4.1. The estimate  $C \gg \mathcal{B}_{MR}$  follows from the proof of Theorem 4.1 and (6.6). To justify  $C \gg \mathcal{A}_M$  we fix an arbitrary number  $t > 0$  and consider a test function  $F_t := F(x)$  determined by

$$F(x) = \begin{cases} 0, & x \notin [(t^-)^-, t^+], \\ \int_{(t^-)^-}^x v^{p'}(y) dy, & x \in [(t^-)^-, t^-], \\ \int_{(t^-)^-}^{t^-} v^{p'}(y) dy, & x \in [t^-, t], \\ \int_x^{t^+} v^{p'}(y) dy \frac{\int_{(t^-)^-}^{t^-} v^{p'}(y) dy}{\int_t^{t^+} v^{p'}(y) dy}, & x \in [t, t^+]. \end{cases}$$

Obviously,  $F$  is absolutely continuous,  $\text{supp} F \subseteq [(t^-)^-, t^+]$  and

$$\left( \int_{\Delta^-(t)} w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right) \leq \|Fw\|_q. \quad (6.8)$$

Let us show that

$$\|F\|_{W_{p,s}^1} \ll \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{1}{p}}. \quad (6.9)$$

We have

$$\begin{aligned} \|Fu\|_s^s &= \int_{\Delta^-(t^-)} \left( \int_{(t^-)^-}^z v^{p'}(y) dy \right)^s u^s(z) dz + \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^s \int_{\Delta^-(t)} u^s(z) dz \\ &\quad + \left( \frac{\int_{\Delta^-(t^-)} v^{p'}(y) dy}{\int_{\Delta^+(t)} v^{p'}(y) dy} \right)^s \int_{\Delta^+(t)} \left( \int_z^{t^+} v^{p'}(y) dy \right)^s u^s(z) dz =: I_1 + I_2 + I_3, \\ \|F'/v\|_p^p &= \int_{\Delta^-(t^-)} v^{p'}(y) dy + \left( \frac{\int_{\Delta^-(t^-)} v^{p'}(y) dy}{\int_{\Delta^+(t)} v^{p'}(y) dy} \right)^p \int_{\Delta^+(t)} v^{p'}(y) dy =: I_4 + I_5. \end{aligned}$$



Using (6.3) and (6.4) we find

$$\begin{aligned}
 I_1 &\leq \left( \int_{\Delta^-(t^-)} u^s(z) dz \right) \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^s \leq \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{s}{p}}, \\
 I_2 &\leq \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{s}{p}} \left( \int_{\Delta(t^-)} v^{p'}(y) dy \right)^{\frac{s}{p'}} \int_{\Delta(t)} u^s(z) dz \\
 &= 2^{s/p'} \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{s}{p}} \left( \int_{t^-}^{(t^-)^+} v^{p'}(y) dy \right)^{\frac{s}{p'}} \int_{\Delta(t)} u^s(z) dz \\
 &\leq 2^{s/p'} \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{s}{p}}, \\
 I_3 &\leq \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{s}{p}} \left( \int_{t^-}^{(t^-)^+} v^{p'}(y) dy \right)^{\frac{s}{p'}} \left( \int_{\Delta^+(t)} u^s(z) dz \right) \\
 &\leq \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{s}{p}}, \\
 I_5 &\leq 2^{p-1} \int_{\Delta^-(t^-)} v^{p'}(y) dy.
 \end{aligned}$$

Putting these estimates together we get (6.9). Now we represent the function  $F$  in the form

$$F = F_1 + F_2,$$

where

$$F_1(x) = \begin{cases} 0, & x \notin \Delta(t^-), \\ \int_{(t^-)^-}^x v^{p'}(y) dy, & x \in \Delta^-(t^-), \\ \int_x^{(t^-)^+} v^{p'}(y) dy, & x \in \Delta^+(t^-). \end{cases}$$

$F_1 \in \mathcal{L}$  because of (6.3), therefore  $F_2 := F - F_1$  belongs to  $\mathcal{L}$  too. It is easy to see, that

$$\begin{aligned}
 \|F_1\|_{W_{p,s}^1} &\leq \int_{\Delta^-(t^-)} v^{p'}(y) dy \left( \int_{\Delta^-(t^-)} u^s(z) dz \right)^{\frac{1}{s}} + \int_{\Delta^-(t^-)} v^{p'}(y) dy \left( \int_{\Delta^+(t^-)} u^s(z) dz \right)^{\frac{1}{s}} \\
 &\quad + 2 \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{1}{p}} \ll \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{1}{p}}
 \end{aligned} \tag{6.10}$$

and by the inequality (6.9)

$$\|F_2\|_{W_{p,s}^1} \leq \|F\|_{W_{p,s}^1} + \|F_1\|_{W_{p,s}^1} \ll \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{1}{p}}. \tag{6.11}$$

From (6.8), (6.10) and (6.11) applying (6.7) to  $F_1$  and  $F_2$  we find

$$\left( \int_{\delta^+(t^-)} w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right) \leq \|F_1 w\|_q + \|F_2 w\|_q$$

$$\leq C \left[ \|F_1\|_{W_{p,s}^1} + \|F_2\|_{W_{p,s}^1} \right] \ll C \left( \int_{\Delta^-(t^-)} v^{p'}(y) dy \right)^{\frac{1}{p}}.$$

It implies

$$\left( \int_{\delta^+(t^-)} w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{\Delta(t^-)} v^{p'}(y) dy \right)^{\frac{1}{p'}} \ll C,$$

where  $\delta^+(t) = [t, a^{-1}(t)]$ ,  $\delta^-(t) = [b^{-1}(t), t]$ . Similar inequality with  $\delta^-(t^+)$  and  $\Delta(t^+)$  instead of  $\delta^+(t^-)$  and  $\Delta(t^-)$  follows from analogous consideration with test function  $G_t := G$  of the form

$$G(x) = \begin{cases} 0, & x \notin [t^-, (t^+)^+], \\ \int_{t^-}^x v^{p'}(y) dy \frac{\int_{t^+}^{(t^+)^+} v^{p'}(y) dy}{\int_{t^-}^{t^+} v^{p'}(y) dy}, & x \in [t^-, t], \\ \int_{t^+}^{(t^+)^+} v^{p'}(y) dy, & x \in [t, t^+], \\ \int_x^{(t^+)^+} v^{p'}(y) dy, & x \in [t^+, (t^+)^+]. \end{cases}$$

Since  $t > 0$  was arbitrary we obtain  $\mathcal{A}_M \ll C$ .

## 6.2. Inequalities on monotone functions

In this section we study the operator (1.1) from weighted  $L_p$  to  $L_q$  on subclasses of non-increasing ( $f \downarrow$ ) or non-decreasing ( $f \uparrow$ ) non-negative functions. The border functions  $a(x)$  and  $b(x)$  as before satisfy the conditions (1.2).

Applying the Sawyer criterion [21] we reduce the problem to the  $L_p - L_q$  characterization of integral operators with Oinarov's kernels considered in Section 3. Using the results of Theorem 3.1 and 3.2 we characterize the operator  $\mathcal{H}$  on the cones of monotone functions.

Let  $U(x) := \int_0^x u^p(y) dy$ . Put

$$\begin{aligned} \mathcal{A}_0^\downarrow &= \left( \int_0^\infty \left[ \int_{a(x)}^{b(x)} v(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} U^{-1/p}(\infty), \\ \mathcal{A}_{1,0}^\downarrow &:= \sup_{t>0} \left( \int_0^t \left[ \int_{a(x)}^{b(x)} v(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{b(t)}^\infty U^{-p'}(z) u^p(z) dz \right)^{\frac{1}{p'}}, \\ \mathcal{A}_{1,1}^\downarrow &= \sup_{s>0} \sup_{s \leq t \leq a^{-1}(b(s))} \left[ \mathbb{A}_0^\downarrow(s, t) + \mathbb{A}_1^\downarrow(s, t) \right], \end{aligned}$$

where

$$\mathbb{A}_0^\downarrow(s, t) = \left( \int_s^t \left[ \int_{a(x)}^{a(t)} v(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} U^{-p'}(z) u^p(z) dz \right)^{\frac{1}{p'}},$$

$$\mathbb{A}_1^\downarrow(s, t) = \left( \int_s^t w^q(x) dx \right)^{\frac{1}{q}} \left( \int_{a(t)}^{b(s)} \left[ \int_{a(t)}^z v(y) dy \right]^{p'} U^{-p'}(z) u^p(z) dz \right)^{\frac{1}{p'}}.$$

Also we notate

$$\begin{aligned} \mathcal{B}_{1,0}^\downarrow &:= \left( \int_0^\infty \left[ \int_0^t \left\{ \int_{a(x)}^{b(x)} v(y) dy \right\}^q w^q(x) dx \right]^{\frac{r}{p}} \right. \\ &\quad \times \left. \left[ \int_{b(t)}^\infty U^{-p'}(z) u^p(z) dz \right]^{\frac{r}{p'}} \left[ \int_{a(t)}^{b(t)} v(y) dy \right]^q w^q(t) dt \right)^{\frac{1}{r}}, \\ \mathcal{B}_{1,1}^\downarrow &= \left[ \sum_{k \in \mathbb{Z}} \left( \mathbb{B}_{k,1}^\downarrow \right)^r + \left( \mathbb{B}_{k,2}^\downarrow \right)^r + \left( \mathbb{B}_{k,3}^\downarrow \right)^r + \left( \mathbb{B}_{k,4}^\downarrow \right)^r \right]^{\frac{1}{r}}, \end{aligned}$$

where

$$\begin{aligned} \mathbb{B}_{k,1}^\downarrow &= \left( \int_{a(\xi_k)}^{a(\xi_{k+1})} \left[ \int_{\xi_k}^{a^{-1}(t)} \left\{ \int_{a(x)}^t v(y) dy \right\}^q w^q(x) dx \right]^{\frac{r}{q}} \right. \\ &\quad \times \left. \left[ \int_t^{a(\xi_{k+1})} U^{-p'}(z) u^p(z) dz \right]^{\frac{r}{q'}} U^{-p'}(t) u^p(t) dt \right)^{\frac{1}{r}}, \\ \mathbb{B}_{k,2}^\downarrow &= \left( \int_{\xi_k}^{\xi_{k+1}} \left[ \int_{\xi_k}^t w^q(x) dx \right]^{\frac{r}{p}} \right. \\ &\quad \times \left. \left[ \int_{a(t)}^{a(\xi_{k+1})} \left\{ \int_{a(t)}^z v(y) dy \right\}^{p'} U^{-p'}(z) u^p(z) dz \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}}, \\ \mathbb{B}_{k,3}^\downarrow &= \left( \int_{b(\xi_k)}^{b(\xi_{k+1})} \left[ \int_{b^{-1}(t)}^{\xi_{k+1}} \left\{ \int_{a(x)}^{a(\xi_{k+1})} v(y) dy \right\}^q w^q(x) dx \right]^{\frac{r}{q}} \right. \\ &\quad \times \left. \left[ \int_{b(\xi_k)}^t U^{-p'}(z) u^p(z) dz \right]^{\frac{r}{q'}} U^{-p'}(t) u^p(t) dt \right)^{\frac{1}{r}}, \\ \mathbb{B}_{k,4}^\downarrow &= \left( \int_{\xi_k}^{\xi_{k+1}} \left[ \int_t^{\xi_{k+1}} w^q(x) dx \right]^{\frac{r}{p}} \right. \\ &\quad \times \left. \left[ \int_{b(\xi_k)}^{b(t)} \left\{ \int_{a(\xi_{k+1})}^z v(y) dy \right\}^{p'} U^{-p'}(z) u^p(z) dz \right]^{\frac{r}{p'}} w^q(t) dt \right)^{\frac{1}{r}}, \end{aligned}$$

and  $\xi_k = (a^{-1} \circ b)^k(\xi_0)$ ,  $\xi_0 = 1$ .

THEOREM 6.2. *For the least possible constant  $C$  in the inequality*

$$\|\mathcal{H}f\|_q \leq C \|fu\|_p, \quad f \downarrow, \quad (6.12)$$

the estimates

$$C \approx \begin{cases} \mathcal{A}_0^\downarrow + \mathcal{A}_{1,0}^\downarrow + \mathcal{A}_{1,1}^\downarrow, & 1 < p \leq q < \infty, \\ \mathcal{A}_0^\downarrow + \mathcal{B}_{1,0}^\downarrow + \mathcal{B}_{1,1}^\downarrow, & 1 < q < p < \infty, \end{cases}$$

hold.

*Proof.* We start with analysis of the Sawyer criterion for the three-weighted inequality

$$\|Tf\|_q \leq C \|fu\|_p, \quad f \downarrow, \quad (6.13)$$

where

$$Tf(x) = w(x) \int_0^\infty t(x, y) f(y) v(y) dy$$

is an integral operator with a non-negative kernel  $t(x, y)$ . Let

$$T^*g(y) = v(y) \int_0^\infty t(x, y) g(x) w(x) dx$$

be a formally adjoint operator to  $T$ . By Sawyer criterion [21] the inequality (6.13) is equivalent, when  $1 < p, q < \infty$ , to the following two inequalities

$$\left( \int_0^\infty \left[ U^{-1}(z) u^{p-1}(z) \int_0^z T^*g(y) dy \right]^{p'} dx \right)^{\frac{1}{p'}} \leq C_1 \left( \int_0^\infty g^{q'}(x) dx \right)^{\frac{1}{q'}}, \quad g \geq 0, \quad (6.14)$$

and

$$\int_0^\infty T^*g(y) dy \leq C_2 \left( \int_0^\infty g^{q'}(x) dx \right)^{\frac{1}{q'}} U^{1/p}(\infty), \quad g \geq 0. \quad (6.15)$$

We assume the constants  $C$ ,  $C_1$  and  $C_2$  as the least possible. The second inequality (6.15) is easily characterized by the duality in Lebesgue's space and

$$C_2 = \left( \int_0^\infty \left[ \int_0^\infty t(x, y) v(y) dy \right]^q w^q(x) dx \right)^{\frac{1}{q}} U^{-1/p}(\infty).$$

As for the first inequality (6.14) it is more convenient for our purpose to use its dual form

$$\begin{aligned} & \left( \int_0^\infty \left[ w(x) \int_0^\infty t(x, y) v(y) \left\{ \int_y^\infty U^{-1}(z) u^{p-1}(z) g(z) dz \right\} dy \right]^q dx \right)^{\frac{1}{q}} \\ & \leq C_1 \left( \int_0^\infty g^p(z) dz \right)^{\frac{1}{p}}, \quad g \geq 0. \end{aligned} \quad (6.16)$$

Now, applying this scheme for the least possible constant  $C$  in (6.12) we have

$$C \approx \mathcal{A}_0^\downarrow + \mathcal{A}_1^\downarrow,$$

where  $\mathcal{A}_1^\downarrow$  is a characterization constant for the inequality

$$\begin{aligned} & \left( \int_0^\infty \left[ w(x) \int_{a(x)}^{b(x)} v(y) \left\{ \int_y^\infty U^{-1}(z) u^{p-1}(z) g(z) dz \right\} dy \right]^q dx \right)^{\frac{1}{q}} \\ & \leq C_1 \left( \int_0^\infty g^p(z) dz \right)^{\frac{1}{p}}, \quad g \geq 0. \end{aligned} \quad (6.17)$$

We write for the left hand side of (6.17)

$$\begin{aligned} F &:= \int_0^\infty \left[ w(x) \int_{a(x)}^{b(x)} v(y) \left\{ \int_y^\infty U^{-1}(z) u^{p-1}(z) g(z) dz \right\} dy \right]^q dx \\ &\approx \int_0^\infty \left[ w(x) \int_{a(x)}^{b(x)} v(y) dy \left\{ \int_{b(x)}^\infty U^{-1}(z) u^{p-1}(z) g(z) dz \right\} \right]^q dx \\ &\quad + \int_0^\infty \left[ w(x) \int_{a(x)}^{b(x)} U^{-1}(z) u^{p-1}(z) g(z) \left\{ \int_{a(x)}^z v(y) dy \right\} dz \right]^q dx. \end{aligned}$$

Hence,

$$\begin{aligned} F &\approx \int_0^\infty \left[ w(x) \int_{a(x)}^{b(x)} v(y) dy \left\{ \int_{b(x)}^\infty U^{-1}(z) u^{p-1}(z) g(z) dz \right\} \right]^q dx \\ &\quad + \int_0^\infty \left[ w(x) \int_{a(x)}^{b(x)} k(z, x) g(z) U^{-1}(z) u^{p-1}(z) dz \right]^q dx, \end{aligned}$$

where  $k(z, x) := \int_{a(x)}^z v(y) dy$ . Clearly, the kernel  $k(z, x) \geq 0$  satisfies the condition (3.4). Thus, the inequality (6.17) is equivalent to the following two inequalities

$$\begin{aligned} & \left( \int_0^\infty \left[ w(x) \int_{a(x)}^{b(x)} v(y) dy \left\{ \int_{b(x)}^\infty U^{-1}(z) u^{p-1}(z) g(z) dz \right\} \right]^q dx \right)^{\frac{1}{q}} \\ & \leq C_{1,0} \left( \int_0^\infty g^p(z) dz \right)^{\frac{1}{p}}, \quad g \geq 0, \end{aligned} \quad (6.18)$$

and

$$\begin{aligned} & \left( \int_0^\infty \left[ w(x) \int_{a(x)}^{b(x)} k(z, x) g(z) U^{-1}(z) u^{p-1}(z) dz \right]^q dx \right)^{\frac{1}{q}} \\ & \leq C_{1,1} \left( \int_0^\infty g^p(z) dz \right)^{\frac{1}{p}}, \quad g \geq 0, \end{aligned} \quad (6.19)$$

with  $C_1 \approx C_{1,0} + C_{1,1}$ . The constant  $C_{1,0}$  of (6.18) is characterized by Lemma 2.2 and 2.4: for  $1 < p \leq q < \infty$  we have  $C_{1,0} \approx \mathcal{A}_{1,0}^\downarrow$ , and  $C_{1,0} \approx \mathcal{B}_{1,0}^\downarrow$  for  $1 < q < p < \infty$ . Applying Theorem 3.1 for the inequality (6.19) we finish the proof.

REMARK 6.1. The similar result as Theorem 6.2 is also true for non-decreasing functions. We omit details.

### Bibliographical Remarks

*Section 1.* Investigation of the operators (1.1) in a primitive form was started in [2], the case  $a(x) = x$ ,  $b(x) = 2x$  was later completely characterized in PhD Thesis by E.N. Batuev [1]. The regular study of the  $L_p - L_q$  boundedness of (1.1) was initiated in [7] and for (1.4) in [6] in the Banach function spaces setting and continued in [28], [5]. In particular, the important conception of a fairway-function was introduced in [28].

*Section 2.* The standard references for the weighted Hardy inequalities are the monographs [14], [10], [9] and [12] (§1.3) with original papers [31], [13], [4] and [23]. The Hardy type operators were studied in [16], [3], [26] (see also [20]). The characterization constant (2.7) was discovered in [18].

*Section 3.* Lemma 3.1 was stated in [29]. Theorems 3.1, 3.2 improve and correct the related results of [28].

*Section 4.* Less general form of Theorems 4.1 and 4.2 can be found in [28], [30].

*Section 5.* Theorem 5.1 in case  $0 < p \leq q < \infty$  is proved in [17].

*Section 6.* Theorem 6.1 is closely related to the results of [15] and Theorem 6.2 to the results of [24], [25], [7], [22], [19], [8].

### REFERENCES

- [1] BATUEV, E.N., *Weighted inequalities of Hardy type and applications*, PhD Thesis, 1991, Khabarovsk University of Technology.
- [2] BATUEV, E.N.; STEPANOV, V.D., *On weighted inequalities of Hardy type*, Siberian Math. J., **30** (1989), 8–16.
- [3] BLOOM, S.; KERMAN, R., *Weighted norm inequalities for operators of Hardy type*, Proc. Amer. Math. Soc., **113** (1991), 135–141.
- [4] BRADLEY, J.S., *Hardy inequalities with mixed norms*, Canad. Math. Bull., **21** (1978), 405–408.
- [5] CHEN, T.; SINNAMON, G., *Generalized Hardy operators and normalizing measures*, J. Ineq. Appl., **7** (2002), 829–866.
- [6] GOGATISHVILI, A.; LANG, J., *The generalized Hardy operators with kernel and variable integral limits in Banach function spaces*, J. Inequal. Appl., **4** (1999), 1–16.
- [7] HEINIG, H.P.; SINNAMON, G., *Mapping properties of integral averaging operators*, Studia Math., **129** (1998), 157–177.
- [8] JOHANSSON, M.; STEPANOV, V. D.; USHAKOVA, E.P., *Hardy inequality with three measures on monotone functions*, Math. Inequal. Appl., **11**, 3 (2008), 393–413.
- [9] KUFNER, A.; MALIGRANDA, L.; PERSSON, L.-E., *The Hardy inequality. About its history and some related results*, Vydavatelský Servis, Plzeň, 2007.
- [10] KUFNER, A.; PERSSON, L.-E., *Weighted inequalities of Hardy type*, World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
- [11] LOMAKINA, E.N.; STEPANOV, V.D., *On the Hardy-type integral operators in Banach function spaces*, Publ. Mat., **42** (1998), 165–194.
- [12] MAZ'YA V.G., *Sobolev Spaces*, Springer-Verlag, Berlin, 1985.
- [13] MUCKENHOUT, B., *Hardy's inequality with weights*, Studia Math., **44** (1972), 31–38.
- [14] OPIC, B.; KUFNER, A., *Hardy-type inequalities*, Pitman Research Notes in Mathematics Series 219, Longman Scientific & Technical, Harlow, 1990.
- [15] OINAROV, R., *On weighted norm inequalities with three weights*, J. London Math. Soc., **48** (1993), 103–116.

- [16] OINAROV, R., *Two-sided estimates of the norm of some classes of integral operators*, Proc. Steklov Inst. Math., **204**, 3 (1994), 205–214.
- [17] PERSSON, L.-E.; PROKHOROV, D., *Integral inequalities for some weighted geometric mean operator with variable limits*, Arch. Ineq. Appl., **2** (2004), 475–482.
- [18] PERSSON, L.-E.; STEPANOV, V. D., *Weighted integral inequalities with the geometric mean operator*, J. Inequal. Appl., **7** (2002), 727–746.
- [19] PERSSON, L.-E.; STEPANOV, V. D.; USHAKOVA, E.P., *Equivalence of Hardy-type inequalities with general measures on the cones of non-negative respective non-increasing functions*, Proc. Amer. Math. Soc., **134**, 8 (2006), 2363–2372.
- [20] PROKHOROV, D., *Inequalities of Hardy type for a class of integral operators with measures*, Anal. Math., **33** (2007), 199–225.
- [21] SAWYER, E.T., *Boundedness of classical operators on classical Lorentz spaces*, Studia Math., **99** (1990), 135–158.
- [22] SINNAMON, G., *Hardy's inequality and monotonicity*, Function spaces, Differential Operators and Nonlinear Analysis, Prague, 2005, pp. 292–310.
- [23] SINNAMON, G.; STEPANOV, V.D., *The weighted Hardy inequality: new proofs and the case  $p=1$* , J. London Math. Soc., **54** (1996), 89–101.
- [24] STEPANOV, V.D. *Integral operators on the cone of monotone functions*, J. London Math. Soc., 1993, vol. 48, pp. 465–487.
- [25] STEPANOV, V.D., *The weighted Hardy's inequality for nonincreasing functions*, Trans. Amer. Math. Soc., **338** (1993), 173–186.
- [26] STEPANOV, V.D., *Weighted norm inequalities of Hardy type for the class of integral operators*, J. London Math. Soc., **50** (1994), 105–120.
- [27] STEPANOV, V.D., *Weighted norm inequalities for integral operators and related topics*, Nonlinear analysis, function spaces and applications, Prague, 1994, vol. 5, pp. 139–175.
- [28] STEPANOV, V.D.; USHAKOVA, E.P., *On integral operators with variable limits of integration*, Proc. Steklov Inst. Math., **232** (2001), 290–309.
- [29] STEPANOV, V.D.; USHAKOVA, E.P., *Hardy operator with variable limits on monotone functions*, J. Funct. Spaces Appl., **1** (2003), 1–15.
- [30] STEPANOV, V.D.; USHAKOVA, E.P., *On the geometric mean operator with variable limits of integration*, Proc. Steklov Inst. Math., **260** (2008), 264–288.
- [31] TOMASELLI, G., *A class of inequalities*, Boll. Unione Mat. Ital., **2** (1969), 622–631.

(Received August 14, 2008)

Vladimir D. Stepanov  
 Department of Mathematical Analysis and Function Theory  
 Peoples Friendship University  
 117198 Moscow  
 Russia  
 e-mail: vstepanov@sci.pfu.edu.ru

Elena P. Ushakova  
 Computing Centre of Far Eastern Branch of Russian Academy of Sciences  
 680000 Khabarovsk  
 Russia  
 Department of Mathematics  
 Uppsala University, Box 480  
 751 06 Uppsala  
 Sweden  
 e-mail: elena@math.uu.se