



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/74798/>

Proceedings Paper:

Beyersdorff, O (2004) Representable Disjoint NP-Pairs. In: FSTTCS 2004: Foundations of Software Technology and Theoretical Computer Science. FSTITCS 2004, 16-18 Dec 2004, Chennai, India. Springer Verlag, 122 - 134 . ISBN: 3-540-24058-6. ISSN: 0302-9743.

https://doi.org/10.1007/978-3-540-30538-5_11

Reuse

See Attached

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Representable Disjoint NP-Pairs

Olaf Beyersdorff

Institut für Informatik,
Humboldt-Universität zu Berlin,
D-10099 Berlin, Germany
beyersdo@informatik.hu-berlin.de

Abstract. We investigate the class of disjoint NP-pairs under different reductions. The structure of this class is intimately linked to the simulation order of propositional proof systems, and we make use of the relationship between propositional proof systems and theories of bounded arithmetic as the main tool of our analysis. Specifically we exhibit a pair which is complete under strong reductions for all disjoint NP-pairs representable in a theory. We use these pairs to explain the simulation order of NP-pairs under these reductions. As corollaries we also get simplified proofs of results obtained earlier in [3] and [5].

1 Introduction

Disjoint NP-pairs (DNPP) naturally occur in cryptography (cf. [4]). The investigation of disjoint NP-pairs in connection with propositional proof systems was initiated by Razborov [12] and further developed by Pudlák [11] and Köbler et al. [5]. These applications attracted more complexity theoretic research on the structure of the class of disjoint NP-pairs (cf. [2, 3, 5]). Various reductions between NP-pairs were introduced by Grollmann and Selman [4]. For the most usual form of a many-one-reduction between DNPP a polynomial time computable function is required to map the components of the two pairs to each other. We denote this reduction here by \leq_p . Later Köbler et al. defined in [5] a strong reduction (denoted by \leq_s), where additionally to \leq_p the reduction function has to map the complements of the pairs to each other.

One of the most prominent questions regarding disjoint NP-pairs is whether there exist complete pairs for the class of all DNPP under these reductions. These problems remain open and various oracle results from [2] indicate that these are indeed difficult questions. Under the assumption that there is an optimal proof system, however, Razborov showed the existence of a \leq_p -complete pair. This was improved by Köbler et al. in [5] to the existence of a complete pair for \leq_s .

Razborov associates to a proof system a canonical disjoint NP-pair and uses the relationship between theories of bounded arithmetic and propositional proof systems for his investigation. In this paper we define another canonical pair for a proof system which plays the same role for the stronger \leq_s -reduction as Razborov's pair for \leq_p . We show that these canonical pairs are quite typical for the class of all DNPP in the sense that every DNPP is \leq_s -reducible to such a

canonical pair, and if there exists a \leq_s -complete pair then it is equivalent to a canonical pair. As one consequence we obtain that, while \leq_p and \leq_s are generally different, the existence of a \leq_p -complete pair already implies the existence of a \leq_s -complete pair. This was also observed by Glaßer et al. in [3] using direct arguments where no reference to proof systems is made.

In this paper, however, we aim to explain some facts about the structure of disjoint NP-pairs by using the close relationship between NP-pairs, proof systems and bounded arithmetic. This also considerably simplifies proofs of earlier results in [3] and [5] which were originally shown by more involved simulation techniques.

Pursuing the afore mentioned goal we start in Sect. 2 by reviewing relevant facts about the connection between propositional proof systems and bounded arithmetic. We only give a very brief presentation tailored to our applications in later sections and refer the reader to [6] or [10] for a detailed account of this rich relationship.

In Sect. 3 we define and separate the afore mentioned reductions between NP-pairs.

In Sect. 4 we start to explain the relationship between disjoint NP-pairs and propositional proof systems by restricting the class of all DNPP to the DNPP representable in some theory T of bounded arithmetic, where a DNPP is called representable in T if the disjointness of the pair is provable in the theory T . We present a \leq_s -complete pair for all DNPP representable in sufficiently strong theories. To make the paper self contained we also reprove some known results.

In Sect. 5 we show that if \leq_s -complete pairs exist then these are equivalent to a canonical pair from Sect. 4 and derive some consequences on the relationship between the simulation order of proof systems and the class of DNPP.

In Sect. 6 we discuss separators and Turing reductions. We show that our pairs from Sect. 4 are candidates for NP-pairs which can not be separated by sets from P, and that the class of all DNPP representable in some theory T is closed under smart Turing-reductions implying that even the existence of smart Turing-complete pairs suffices for the existence of \leq_s -complete DNPP which is also shown in [3].

2 Preliminaries

Propositional proof systems were defined in a very general way by Cook and Reckhow in [1] as polynomial time functions P which have as its range the set of all tautologies. A string π with $P(\pi) = \varphi$ is called a P -proof of the tautology φ . By $P \vdash_{\leq m} \varphi$ we indicate that there is a P -proof of φ of length $\leq m$. If φ_n is a sequence of propositional formulas we write $P \vdash_* \varphi_n$ if there is a polynomial p such that $P \vdash_{\leq p(|\varphi_n|)} \varphi_n$.

Given two proof systems P and S we say that S simulates P (denoted by $P \leq S$) if there exists a polynomial p such that for all tautologies φ and P -proofs π of φ there is a S -proof π' of φ with $|\pi'| \leq p(|\pi|)$. If such a proof π' can even be computed from π in polynomial time we say that S p-simulates P and denote

this by $P \leq_p S$. A proof system is called (p-)optimal if it (p-)simulates all proof systems.

In this paper we are only concerned with sufficiently strong proof systems simulating the extended Frege proof system EF , where EF is a usual textbook proof system based on axioms and rules and augmented by the possibility to abbreviate complex formulas by propositional variables to reduce the proof size (see e.g. [6]). For simplicity we call proof systems simulating EF strong. A method how to actually construct strong proof systems was recently described in [7].

We now review the relationship between theories of arithmetic and proof systems. Let L be the language of arithmetic (cf. [6]). Bounded L -formulas are formulas in the language of L where only quantifiers of the form $(\forall x \leq t(y))$ and $(\exists x \leq s(y))$ occur with L -terms t and s . In the following we are particularly interested in Π_1^b and Σ_1^b -formulas where only bounded universal and bounded existential quantifiers are allowed, respectively.

To explain the connection to propositional proof systems we have to translate L -formulas into propositional formulas. Let $\varphi(x)$ be a Π_1^b -formula. We can assume that φ is of the form $(\forall y) |y| \leq |x|^k \rightarrow \psi(x, y)$ with some polynomial time computable predicate ψ . Hence we can compute $\psi(x, y)$ by polynomial size boolean circuits C_n for numbers x of length n . From C_n we build a propositional formula $\|\varphi\|^n$ with atoms p_1, \dots, p_n for the bits of x , atoms q_1, \dots, q_n for the bits of y and auxiliary atoms $r_1, \dots, r_{n^{O(1)}}$ for the inner nodes of C_n . The formula $\|\varphi\|^n$ describes that if the values for \bar{r} are correctly computed from \bar{p} and \bar{q} then the output of the computation of C_n is 1. Thus we get a sequence of propositional formulas $\|\varphi\|^n$ of polynomial size in n and $\|\varphi\|^n$ is a tautology iff $\varphi(x)$ holds for all natural numbers of length $\leq n$.

Encoding propositional formulas as numbers in some straightforward way we can in a theory T speak of propositional formulas, assignments and proofs. Let $\text{Prf}_P(\pi, \varphi)$ be a L -formula describing that π is the encoding of a correct P -proof of the propositional formula encoded by φ . Similarly, let $\text{Taut}(\varphi)$ be a L -formula asserting that all assignments satisfy the formula φ . Because P is computable in polynomial time Prf_P is definable by a Σ_1^b -formula whereas Taut is in Π_1^b .

The reflection principle for a propositional proof system P is the L -formula

$$\text{RFN}(P) = (\forall \pi)(\forall \varphi)\text{Prf}_P(\pi, \varphi) \rightarrow \text{Taut}(\varphi)$$

and states a strong form of the consistency of the proof system P . From the last remark it follows that $\text{RFN}(P)$ is a $\forall \Pi_1^b$ -formula.

In [8] a general correspondence between L -theories T and propositional proof systems P is introduced. Pairs (T, P) from this correspondence possess in particular the following two properties:

1. For all $\varphi(x) \in \Pi_1^b$ with $T \vdash (\forall x)\varphi(x)$ we have polynomially long P -proofs of the tautologies $\|\varphi(x)\|^n$.
2. T proves the correctness of P , i.e. $T \vdash \text{RFN}(P)$. Furthermore P is the strongest proof system for which T proves the correctness, i.e. $T \vdash \text{RFN}(S)$ for a proof system S implies $S \leq_p P$.

The most prominent example for this correspondence is the pair (S_2^1, EF) where S_2^1 is a L -theory with induction for Σ_1^b -formulas. This in particular allows the formalization of polynomial time computations and the provability of its basic properties (see e.g. [6] Chapter 6).

To every L -theory $T \supseteq S_2^1$ with a polynomial time set of axioms we can associate a proof system P which is unique up to \leq_p -equivalence by property 2 above. Conversely every strong proof system has a corresponding theory, but here according to property 1 only the $\forall\Pi_1^b$ -consequences of T are determined by P .

As the correspondence only works for sufficiently strong proof systems we will restrict ourselves to proof systems $P \geq EF$ and theories $T \supseteq S_2^1$.

By \mathcal{N} we denote the standard model of arithmetic which is in particular a submodel of all models of theories T considered here.

3 Reductions between NP-Pairs

A pair (A, B) is called a disjoint NP-pair (DNPP), if $A, B \in \text{NP}$ and $A \cap B = \emptyset$. To exclude trivial cases we additionally require $A \neq \emptyset$ and $B \neq \emptyset$. We consider the following reductions between disjoint NP-pairs.

Definition 1. *Let (A, B) and (C, D) be DNPP.*

1. (A, B) is polynomially reducible to (C, D) ($(A, B) \leq_p (C, D)$), if there exists a function $f \in FP$ such that $f(A) \subseteq C$ and $f(B) \subseteq D$.
2. (A, B) is strongly reducible to (C, D) ($(A, B) \leq_s (C, D)$), if there exists a function $f \in FP$ such that $f^{-1}(C) = A$ and $f^{-1}(D) = B$.
3. As usual we write $(A, B) \equiv_p (C, D)$ for $(A, B) \leq_p (C, D)$ and $(C, D) \leq_p (A, B)$. \equiv_s is defined in the same way.

$(A, B) \leq_p (C, D)$ does not in general imply that A and B are reducible to C and D , respectively, but if f realizes a \leq_s -reduction from (A, B) to (C, D) , then f is simultaneously a many-one-reduction between A and C as well as between B and D . Equivalently we can also view \leq_s as a reduction between triples. In addition to the two conditions $f(A) \subseteq C$ and $f(B) \subseteq D$ for \leq_p we also require $f(\overline{A \cup B}) \subseteq \overline{C \cup D}$.

Obviously \leq_s is a refinement of \leq_p . Under the assumption $\text{P} \neq \text{NP}$ this is indeed a proper refinement. The reason for this lies in the following proposition:

Proposition 2. *For every DNPP (A, B) there exists a DNPP (A', B') such that $(A, B) \equiv_p (A', B')$ and A', B' are NP-complete.*

Proof. Choose $A' = A \times \text{SAT}$ and $B' = B \times \text{SAT}$. Then we have $(A, B) \leq_p (A', B')$ via $x \mapsto (x, \varphi_0)$ with a fixed formula $\varphi_0 \in \text{SAT}$, and $(A', B') \leq_p (A, B)$ via the projection $(x, \varphi) \mapsto x$. \square

With this proposition we can easily separate the reductions \leq_p and \leq_s under the assumption $\text{P} \neq \text{NP}$. Namely, let A and B be nonempty sets in P such that

$\overline{A \cup B}$ is also nonempty. Choose A' and B' as in the last proposition. Then $(A, B) \equiv_p (A', B')$ but $(A, B) \not\equiv_s (A', B')$ because $(A', B') \leq_s (A, B)$ would imply in particular $A' \leq_m^p A$ and hence $P=NP$. On the other hand if $P=NP$ then all DNPP (A, B) where all three components $A, B, \overline{A \cup B}$ are nonempty would be \leq_s -equivalent. This equivalence of $P \neq NP$ and the separation of \leq_p from \leq_s for DNPP with all three components nonempty (or equivalently for DNPP with all three components infinite) is also observed in [3].

4 Representable NP-Pairs

In the following we investigate the relationship between disjoint NP-pairs and propositional proof systems. We will use the correspondence between proof systems and arithmetical theories as explained in Sect. 2. For this section let P be a strong proof system and T be a corresponding theory.

Following Razborov we call a Σ_1^b -formula φ a representation of an NP-set A , if for all natural numbers a

$$\mathcal{N} \models \varphi(a) \iff a \in A .$$

A DNPP (A, B) is representable in T , if there are Σ_1^b -formulas φ and ψ representing A and B , respectively, such that

$$T \vdash (\forall x)(\neg\varphi(x) \vee \neg\psi(x)) .$$

For the last line we also use the abbreviation $T \vdash A \cap B = \emptyset$. Since $A \cap B = \emptyset$ is a $\forall\Pi_1^b$ -formula we can also express the disjointness of A and B propositionally by the sequence of tautologies $\|\neg\varphi(x) \vee \neg\psi(x)\|^n$, which we again shortly denote by $\|A \cap B = \emptyset\|^n$.

The DNPP representable in T can also be characterized via the corresponding proof system P in the following way:

Proposition 3. *A DNPP (A, B) is representable in T if and only if*

$$P \vdash_* \|A \cap B = \emptyset\|^n$$

for suitable representations of A and B .

Proof. Let φ and ψ be representations for A and B , respectively, such that

$$T \vdash (\forall x)(\neg\varphi(x) \vee \neg\psi(x)) .$$

Because this is a $\forall\Pi_1^b$ -formula, we have $P \vdash_* \|\neg\varphi(x) \vee \neg\psi(x)\|^n$, which we write by definition as $P \vdash_* \|A \cap B = \emptyset\|^n$.

For the other direction let φ and ψ be representations of A and B , such that for some natural number k we have $P \vdash_{\leq n^k} \|\neg\varphi(x) \vee \neg\psi(x)\|^n$. Consider the formula

$$\psi'(x) = \psi(x) \wedge (\exists\pi)|\pi| \leq |x|^k \wedge \text{Prf}_P(\pi, \|\neg\varphi(y) \vee \neg\psi(y)\|^{|x|}) .$$

We have $\psi' \in \Sigma_1^b$ and furthermore $\mathcal{N} \models (\forall x)\psi'(x) \leftrightarrow \psi(x)$, i.e. ψ' is also a representation of B . From $T \vdash \text{RFN}(P)$ it follows that $T \vdash (\forall x)(\neg\varphi(x) \vee \neg\psi'(x))$, hence (A, B) is representable in T . \square

The next lemma shows that all pairs representable in a theory form a reasonable complexity class.

Lemma 4 (Razborov [12]). *The set of all DNPP representable in T is closed under \leq_p -reductions.*

Proof. Let (A, B) and (C, D) be DNPP such that $f : (A, B) \leq_p (C, D)$ and $T \vdash C \cap D = \emptyset$. Consider the NP-sets

$$\begin{aligned} A' &= \{x \mid x \in A \text{ and } f(x) \in C\} \\ B' &= \{x \mid x \in B \text{ and } f(x) \in D\} . \end{aligned}$$

Obviously $A = A'$ and $B = B'$. From $T \supseteq S_2^1$ and $f \in FP$ we get $T \vdash (\forall x)(\exists! y)f(x) = y$. Hence

$$T \vdash (\forall x)(x \in A' \cap B' \rightarrow f(x) \in C \cap D)$$

and with $T \vdash C \cap D = \emptyset$ we conclude $T \vdash A' \cap B' = \emptyset$. □

Following Razborov [12] we associate a disjoint NP-pair $(\text{Ref}(P), \text{SAT}^*)$ with a proof system P with

$$\begin{aligned} \text{Ref}(P) &= \{(\varphi, 1^m) \mid P \vdash_{\leq m} \varphi\} \\ \text{SAT}^* &= \{(\varphi, 1^m) \mid \neg\varphi \in \text{SAT}\} . \end{aligned}$$

$(\text{Ref}(P), \text{SAT}^*)$ is called the canonical pair of P .

Lemma 5 (Razborov [12]). *The canonical pair of P is representable in T .*

Proof. We argue in T . Let $(\varphi, 1^m) \in \text{Ref}(P)$. Then there is a P -proof π of φ . Since $\text{RFN}(P)$ is available in T we conclude from $\text{Prf}_P(\pi, \varphi)$ the formula $\text{Taut}(\varphi)$, hence $\neg\varphi \notin \text{SAT}$ and therefore $(\varphi, 1^m) \notin \text{SAT}^*$. □

Now we associate a second disjoint NP-pair with a proof system P . For a propositional formula φ let $\text{Var}(\varphi)$ be the set of propositional variables occurring in φ . Let

$$\begin{aligned} U_1(P) &= \{(\varphi, \psi, 1^m) \mid \text{Var}(\varphi) \cap \text{Var}(\psi) = \emptyset, \neg\varphi \in \text{SAT} \text{ and } P \vdash_{\leq m} \varphi \vee \psi\} \\ U_2 &= \{(\varphi, \psi, 1^m) \mid \text{Var}(\varphi) \cap \text{Var}(\psi) = \emptyset \text{ and } \neg\psi \in \text{SAT}\} . \end{aligned}$$

As for the canonical pair we get:

Lemma 6. *The pair $(U_1(P), U_2)$ is representable in T .*

Proof. Let $(\varphi, \psi, 1^m) \in U_1(P)$ and π be a P -proof of $\varphi \vee \psi$ of length $\leq m$. Because $\neg\varphi \in \text{SAT}$ we have an assignment α with $\varphi(\alpha) = 0$. If we substitute the variables of φ by 0 or 1 according to α , we get from the proof π a proof π' of ψ . Hence we have $T \vdash (\exists\pi')\text{Prf}_P(\pi', \psi)$. Because T proves the correctness of P , we get $T \vdash \text{Taut}(\psi)$ and thus $T \vdash (\varphi, \psi, 1^m) \notin U_2$. □

Now we come to the main theorem of this section which states the completeness of $(U_1(P), U_2)$ for all DNPP representable in T under \leq_s -reductions.

Theorem 7. *A DNPP (A, B) is representable in T if and only if $(A, B) \leq_s (U_1(P), U_2)$.*

Proof. Let (A, B) be a DNPP such that $T \vdash A \cap B = \emptyset$. Let the NP-sets A and B be of the form

$$\begin{aligned} A &= \{x \mid (\exists y)|y| \leq |x|^{O(1)} \wedge (x, y) \in C\} \\ B &= \{x \mid (\exists z)|z| \leq |x|^{O(1)} \wedge (x, z) \in D\} \end{aligned}$$

with polynomial time predicates C and D . Because of the correspondence between T and P there is a polynomial p for the $\forall\Pi_1^b$ -formula $A \cap B = \emptyset$ such that

$$P \vdash_{\leq p(n)} \|A \cap B = \emptyset\|^n .$$

Here the formula $\|A \cap B = \emptyset\|^n$ is more explicitly $\|(x, y) \notin C \vee (x, z) \notin D\|^n$ and has propositional variables for x, y and z and auxiliary variables for the computation of boolean circuits for C and D . We can plug into this formula natural numbers a of length n for x by substituting the propositional variables corresponding to x by the bits of a . We indicate this by the suffix (x/a) .

Now we claim that the function

$$f(a) = (\|(x, y) \notin C\|^{|a|}(x/a), \|(x, z) \notin D\|^{|a|}(x/a), 1^{p(|a|)})$$

realizes a \leq_s -reduction from (A, B) to $(U_1(P), U_2)$.

If we choose different auxiliary variables for the computation of C and D and also disjoint variables for y and z , then the formulas $\|(x, y) \notin C\|^{|a|}(x/a)$ and $\|(x, z) \notin D\|^{|a|}(x/a)$ have no common variables. Furthermore for every natural number a the formulas

$$\begin{aligned} \|(x, y) \notin C\|^{|a|}(x/a) \vee \|(x, z) \notin D\|^{|a|}(x/a) &= \\ \|(x, y) \notin C \vee (x, z) \notin D\|^{|a|}(x/a) &= \\ \|A \cap B = \emptyset\|^{|a|}(x/a) & \end{aligned}$$

have P -proofs of length $\leq p(|a|)$, which we get from the P -proofs of $\|A \cap B = \emptyset\|^{|a|}$ by substituting the variables for x by the bits of a .

The last thing to check is that the formula

$$\neg \|(x, y) \notin C\|^{|a|} = \|(x, y) \in C\|^{|a|},$$

expressing, that there is a correct accepting computation of C with input (x, y) , is satisfiable if and only if the variables of x are substituted by the bits of a number $a \in A$.

Similarly, $\neg \|(x, z) \notin D\|^{|a|}(x/a)$ is satisfiable if and only if $a \in B$.

The backward implication follows from Lemma 6 and the fact, that the DNPP representable in T are closed under \leq_p and hence also under \leq_s according to Lemma 4. \square

The pair $(U_1(P), U_2)$ strongly resembles the interpolation pair defined by Pudlák in [11]:

$$\begin{aligned} I_P^0 &= \{(\varphi, \psi, \pi) \mid P(\pi) = \varphi \vee \psi, \text{Var}(\varphi) \cap \text{Var}(\psi) = \emptyset \text{ and } \neg\varphi \in \text{SAT}\} \\ I_P^1 &= \{(\varphi, \psi, \pi) \mid P(\pi) = \varphi \vee \psi, \text{Var}(\varphi) \cap \text{Var}(\psi) = \emptyset \text{ and } \neg\psi \in \text{SAT}\} . \end{aligned}$$

This pair is p-separable, if and only if the proof system P has the efficient interpolation property. For $\|\cdot\|$ -translations of $\forall\Pi_1^b$ -formulas provable in T we can efficiently construct polynomially long P -proofs (i.e. with functions from FP). Hence the proof of the last theorem also shows the \leq_s -completeness of (I_P^0, I_P^1) for all DNPP representable in T .

In [11] Pudlák defined a DNPP (A, B) to be symmetric if $(B, A) \leq (A, B)$. With Lemma 6 also the pair $(U_2, U_1(P))$ is representable in T , hence by the last theorem $(U_1(P), U_2)$ is symmetric even with respect to the stronger \leq_s -reduction.

As a corollary of Theorem 7 we obtain the \leq_p -completeness of the canonical pair for all DNPP representable in T , which was shown by Razborov:

Theorem 8 (Razborov [12]). *A DNPP (A, B) is representable in T if and only if $(A, B) \leq_p (\text{Ref}(P), \text{SAT}^*)$.*

Proof. For the forward implication we reduce $(U_1(P), U_2)$ to $(\text{Ref}(P), \text{SAT}^*)$ via the projection

$$(\varphi, \psi, 1^m) \mapsto (\psi, 1^{m+p(|\varphi|)})$$

with a suitable polynomial p .

Let $(\varphi, \psi, 1^m) \in U_1(P)$. Then there is a P -proof π of length $\leq m$ of $\varphi(\bar{x}) \vee \psi(\bar{y})$. The formula $\neg\varphi(\bar{x})$ is satisfiable, so by substituting a satisfying assignment \bar{a} into the proof π we get a proof π' with $|\pi'| \leq m$ for $\varphi(\bar{a}) \vee \psi(\bar{y})$. Since $\varphi(\bar{a})$ is a false formula without free variables we can evaluate it in polynomially long P -proofs to \perp . Let p be a corresponding polynomial. Thus we get a P -proof of length $\leq m + p(|\varphi|)$ for ψ .

If $(\varphi, \psi, 1^m) \in U_2$, then $\neg\psi$ is satisfiable and hence $(\psi, 1^{m+p(|\varphi|)}) \in \text{SAT}^*$.

This \leq_p -reduction from $(U_1(P), U_2)$ to $(\text{Ref}(P), \text{SAT}^*)$ yields together with the last theorem the \leq_p -completeness of $(\text{Ref}(P), \text{SAT}^*)$ for all DNPP representable in T .

The backward implication follows from Lemma 4 and Lemma 5. \square

Thus the pairs $(\text{Ref}(P), \text{SAT}^*)$ and $(U_1(P), U_2)$ are complete for all DNPP representable in T under \leq_p - and \leq_s -reductions, respectively.

5 NP-Pairs and the Simulation Order of Proof Systems

Now we use the results of the last section to make some observations about the connection between the simulation order of proof systems and disjoint NP-pairs.

In Sect. 3 it was shown that the reductions \leq_p and \leq_s are different under the assumption $P \neq NP$. Still we have:

Proposition 9. *For all strong proof systems P and DNPP (A, B) it holds*

$$(A, B) \leq_p (U_1(P), U_2) \iff (A, B) \leq_s (U_1(P), U_2) .$$

Proof. Let $(A, B) \leq_p (U_1(P), U_2)$. $(U_1(P), U_2)$ is representable in T . Hence with Lemma 4 also (A, B) is representable in T , from which we conclude with Theorem 7 $(A, B) \leq_s (U_1(P), U_2)$.

The opposite implication follows by definition. \square

Corollary 10. *Let P and S be strong proof systems. Then we have:*

$$(\text{Ref}(P), \text{SAT}^*) \leq_p (\text{Ref}(S), \text{SAT}^*) \iff (U_1(P), U_2) \leq_s (U_1(S), U_2) .$$

Proof. For the first direction we get from

$$(U_1(P), U_2) \leq_p (\text{Ref}(P), \text{SAT}^*) \leq_p (\text{Ref}(S), \text{SAT}^*) \leq_p (U_1(S), U_2)$$

together with the last proposition $(U_1(P), U_2) \leq_s (U_1(S), U_2)$.

The other implication follows from

$$(\text{Ref}(P), \text{SAT}^*) \leq_p (U_1(P), U_2) \leq_p (U_1(S), U_2) \leq_p (\text{Ref}(S), \text{SAT}^*) .$$

\square

The following proposition is well known (see e.g. [11]):

Proposition 11. *If P and S are proof systems with $P \leq S$, then we have*

$$(\text{Ref}(P), \text{SAT}^*) \leq_p (\text{Ref}(S), \text{SAT}^*) .$$

Proof. By assumption there is a polynomial p , such that for all formulas φ and P -proofs π of φ there is a S -proof π' of length $\leq p(|\pi|)$. Therefore the mapping

$$(\varphi, 1^m) \mapsto (\varphi, 1^{p(m)})$$

is a \leq_p -reduction from $(\text{Ref}(P), \text{SAT}^*)$ to $(\text{Ref}(S), \text{SAT}^*)$. \square

Proposition 11 and Corollary 10 yield:

Corollary 12. *If P and S are strong proof systems with $P \leq S$, then we have*

$$(U_1(P), U_2) \leq_s (U_1(S), U_2) .$$

Köbler, Messner and Torán proved in [5] that the existence of an optimal proof system implies the existence of \leq_s -complete NP-pairs. This result also follows from the last corollary. Additionally we can exhibit a complete DNPP in this case:

Corollary 13. *If P is an optimal proof system, then $(U_1(P), U_2)$ is \leq_s -complete for the class of all DNPP.*

Proof. Let P be an optimal proof system and (A, B) a DNPP. The sequence of tautologies $\|A \cap B = \emptyset\|^n$ can be constructed in polynomial time. Hence there is a proof system S with polynomially long proofs of these tautologies (for example just add these tautologies as axioms to the extended Frege system). Using Proposition 3 and Theorem 7 we get $(A, B) \leq_s (U_1(S), U_2)$. By assumption we have $S \leq P$. Together with the previous corollary this yields $(U_1(S), U_2) \leq_s (U_1(P), U_2)$, and hence $(A, B) \leq_s (U_1(P), U_2)$.

Therefore the pair $(U_1(P), U_2)$ is \leq_s -complete for all DNPP. \square

Proposition 14. *Let (A, B) be \leq_p -complete for the class of all DNPP. Then we have $(A, B) \equiv_p (\text{Ref}(P), \text{SAT}^*)$ for some proof system P .*

Proof. As in the last proof let P be a proof system with $P \vdash_* \|A \cap B = \emptyset\|^n$. Then $(A, B) \leq_p (\text{Ref}(P), \text{SAT}^*)$ and by assumption $(\text{Ref}(P), \text{SAT}^*) \leq_p (A, B)$. \square

In the same way we get:

Proposition 15. *Let (A, B) be \leq_s -complete for the class of all DNPP. Then we have $(A, B) \equiv_s (U_1(P), U_2)$ for some proof system P .*

The following proposition is also observed in [3]:

Proposition 16. *The class of all DNPP contains a \leq_p -complete DNPP if and only if it contains a \leq_s -complete DNPP.*

Proof. For the first direction we can assume with Proposition 14 that the \leq_p -complete DNPP has the form $(\text{Ref}(P), \text{SAT}^*)$ for some proof system P . Then by Theorem 7 and Theorem 8 all DNPP are \leq_s -reducible to $(U_1(P), U_2)$.

The other direction holds by definition. \square

6 Separators and Turing Reductions

For disjoint NP-pairs we can also study Turing reductions as defined by Grollmann and Selman in [4]. For this we need the notion of a separator.

Definition 17. *A set S is a separator for the DNPP (A, B) if $A \subseteq S$ and $B \subseteq \overline{S}$.*

Of central interest is the case where a given DNPP has a separator belonging to P . Such a pair is called p -separable. The set of all p -separable DNPP form the lowest degree with respect to the \leq_p -reduction. For the stronger \leq_s -reduction this minimal degree shrinks to the set of all p -separable pairs with empty complement, i.e. sets of the form (A, \overline{A}) with $A \in P$. But also the set of all p -separable pairs with nonempty complement splits into different \equiv_s -degrees. Namely, if (A, B) is a p -separable DNPP then the pair $(A \times \text{SAT}, B \times \text{SAT})$ is also p -separable and both of its components are NP-complete, hence we have:

Proposition 18. *$P \neq NP$ iff there exist p -separable pairs (A, B) and (C, D) , such that $\overline{A \cup B}$ and $\overline{C \cup D}$ are nonempty and $(A, B) \not\equiv_s (C, D)$.*

The question whether p-inseparable pairs exist is open. Candidates for p-inseparable pairs come from cryptography (cf. [4]) and proof systems. Namely, Krajíček and Pudlák demonstrate in [9] that a pair (A_0, A_1) associated with the RSA-cryptosystem is representable in the theory S_2^1 corresponding to EF . By the results from Sect. 4 this means that $(A_0, A_1) \leq_p (\text{Ref}(EF), \text{SAT}^*)$ and $(A_0, A_1) \leq_s (U_1(EF), U_2)$. Assuming the security of RSA the pair (A_0, A_1) is not p-separable, hence under this assumption neither $(\text{Ref}(P), \text{SAT}^*)$ nor $(U_1(P), U_2)$ is p-separable for any $P \geq EF$.

If we look at the property of symmetry of pairs under \leq_s it is clear that a DNPP (A, B) can not be symmetric if we choose A from P and B NP-complete. In other words:

Proposition 19. *$P \neq NP$ iff there exist non-symmetric pairs with respect to \leq_s .*

A similar result for \leq_p is not known as \leq_p -non-symmetric pairs are p-inseparable and it is not clear how to derive the existence of p-inseparable pairs from the assumption $P \neq NP$.

We now come to the definition of Turing-reductions between DNPP from [4]:

Definition 20. *Let (A, B) and (C, D) be DNPP. (A, B) is Turing reducible to (C, D) ($(A, B) \leq_T (C, D)$), if there exists a polynomial time oracle Turing machine M such that for every separator T of (C, D) $L(M^T)$ separates (A, B) .*

If for inputs from $A \cup B$ the machine M makes only queries to $C \cup D$ we call the reduction performed by M a smart Turing reduction.

In [3] Glaßer et al. prove that the existence of a complete DNPP under smart Turing reductions already implies the existence of a \leq_p -complete DNPP (and hence by Proposition 16 also of a \leq_s -complete pair). We can easily reprove their result in our framework by noticing:

Lemma 21. *The set of all DNPP representable in a theory T is closed under smart Turing reductions.*

Proof. Let the pair (A, B) be smartly Turing reducible to (C, D) via the deterministic oracle Turing machine M , and let (C, D) be representable in T . Consider the NP-sets

$$\begin{aligned} A' &= \{x \mid x \in A \text{ and } M(x) \text{ accepts}\} \\ B' &= \{x \mid x \in B \text{ and } M(x) \text{ rejects}\} . \end{aligned}$$

By " $M(x)$ accepts" we mean that M accepts the input x by a computation where all oracle queries that are positively answered are verified by a computation of a nondeterministic machine for C and all negative answers are verified by D . Since the reduction is smart we have $A = A'$ and $B = B'$. For $T \vdash A' \cap B' = \emptyset$ it suffices to show in T the uniqueness of the computation of M at inputs x from $A \cup B$. T can prove the uniqueness of computations of the deterministic machine M , and the possibility to answer an oracle query both positively and negatively is excluded by $T \vdash C \cap D = \emptyset$. \square

From this we conclude:

Proposition 22. *Suppose (A, B) is a smart \leq_T -complete pair. Then for any theory T such that $T \vdash A \cap B = \emptyset$ the pair $(U_1(P), U_2)$ is \leq_s -complete for all DNPP where P is the proof system corresponding to T .*

Proof. Choose a theory T with $T \vdash A \cap B = \emptyset$. Then by the last lemma all DNPP are representable in T and hence by Theorem 7 the pair $(U_1(P), U_2)$ is \leq_s -complete. \square

It is not clear whether the class of pairs representable in some theory T is also closed under \leq_T -reductions. This corresponds to the open problem from [3] whether the existence of a \leq_T -complete pair implies the existence of a \leq_p -complete DNPP.

Acknowledgments. For helpful conversations and suggestions on this work I am very grateful to Johannes Köbler, Jan Krajíček, and Pavel Pudlák.

References

1. S. Cook and R. Reckhow. The relative efficiency of propositional proof systems. *Journal of Symbolic Logic* **44**:36-50, 1979.
2. C. Glaßer, A. Selman, S. Sengupta, and L. Zhang. Disjoint NP-pairs. In *Proceedings 18th Annual IEEE Conference on Computational Complexity*, pages 313-332, 2003.
3. C. Glaßer, A. Selman, and S. Sengupta. Reductions between disjoint NP-pairs. In *Proceedings 19th Annual IEEE Conference on Computational Complexity*, 2004.
4. J. Grollmann and A. Selman. Complexity measures for public-key cryptosystems. *SIAM Journal on Computing* **17**(2):309-335, 1988.
5. J. Köbler, J. Messner, and J. Torán. Optimal proof systems imply complete sets for promise classes. *Information and Computation* **184**:71-92, 2003.
6. J. Krajíček. *Bounded Arithmetic, Propositional Logic, and Complexity Theory*. Encyclopedia of Mathematics and Its Applications, volume **60**, Cambridge University Press, Cambridge, 1995.
7. J. Krajíček. Implicit proofs. *Journal of Symbolic Logic* **69**(2):387-397, 2004.
8. J. Krajíček and P. Pudlák. Quantified propositional calculi and fragments of bounded arithmetic. *Zeitschr. f. math. Logik und Grundlagen d. Math.* **36**:29-46, 1990.
9. J. Krajíček and P. Pudlák. Some consequences of cryptographical conjectures for S_2^1 and EF . *Information and Computation* **140**(1):82-94, 1998.
10. P. Pudlák. The lengths of proofs. In *Handbook of Proof Theory*, S. R. Buss ed., pages 547-637, Elsevier, Amsterdam, 1998.
11. P. Pudlák. On reducibility and symmetry of disjoint NP-pairs. In *Proceedings 26th International Symposium on Mathematical Foundations of Computer Science*, volume 2136 of *Lecture Notes in Computer Science*, pages 621-632. Springer-Verlag, Berlin, 2001.
12. A. A. Razborov. On provably disjoint NP-pairs. Technical Report TR94-006, Electronic Colloquium on Computational Complexity, 1994.