

This is a repository copy of *Functional genomics reveals that Clostridium difficile Spo0A* coordinates sporulation, virulence and metabolism.

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/78494/

Version: Supplemental Material

Article:

Pettit, L.J., Browne, H.P., Yu, L. et al. (10 more authors) (2014) Functional genomics reveals that Clostridium difficile Spo0A coordinates sporulation, virulence and metabolism. BMC Genomics, 15. 160 - 160. ISSN 1471-2164

https://doi.org/10.1186/1471-2164-15-160

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.





Additional file 1: Identification of differentially expressed genes in *C. difficile* 630 *spo0A* mutant by transcriptional profiling. Scatter plot of the log2 fold changes against the normalised mean read abundance per gene (calculated at the base level). Red dots represent genes considered to be significantly differentially expressed ($P = \le 0.01$). Black dots signify genes not deemed to be significantly differentially expressed according to these criteria.