



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/80337/>

Version: Submitted Version

Article:

Perkins, J, Reed, M, Akanyang, L et al. (2013) Making land management more sustainable: experiences implementing a new methodological framework in Botswana. *Land Degradation & Development*, 24 (5). 463 - 477. ISSN: 1085-3278

<https://doi.org/10.1002/ldr.1142>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



**Making land management more sustainable: experience
implementing a new methodological framework in
Botswana**

Journal:	<i>Land Degradation & Development</i>
Manuscript ID:	LDD-10-0221.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	27-Jun-2011
Complete List of Authors:	Perkins, J; University of Botswana, Dept. of Environmental Sciences Reed, M.S; Aberdeen Centre for Environmental Sustainability and Centre for Planning and Environmental Management, School of Geosciences, University of Aberdeen Akanyang, L; University of Botswana, Dept. of Environmental Sciences Atlhopheng, J; University of Botswana, Dept. of Environmental Sciences Chanda, R; University of Botswana, Dept. of Environmental Sciences Magole, L; University of Botswana, Dept. of Environmental Sciences Mphinyane, W; University of Botswana, Dept. of Environmental Sciences Mulale, K; University of Botswana, Dept. of Environmental Sciences Sebego, R; University of Botswana, Dept. of Environmental Sciences Fleskens, L; University of Leeds, Sustainability Research Institute Irvine, B; University of Leeds, Sustainability Research Institute Kirkby, Mike; University of Leeds, School of Geography
Keywords:	knowledge management, sustainable land management, drylands, Botswana

SCHOLARONE™
Manuscripts

Making land management more sustainable: experience implementing a new methodological framework in Botswana

Perkins JS¹, Reed MS^{2*}, Akanyang L³, Athlopheng JR¹, Chanda R¹, Magole L⁴, Mphinyane W¹, Mulale K¹, Sebegu RJ¹, Fleskens L⁵, Irvine B⁶, Kirkby MJ⁶

¹Department of Environmental Science, University of Botswana, Private Bag 00704, Gaborone, Botswana

² Aberdeen Centre for Environmental Sustainability, Centre for Planning and Environmental Management and Centre for Sustainable International Development, School of Geosciences, University of Aberdeen, St Mary's, Aberdeen AB24 3UF, UK

³ Botswana College of Agriculture, Department of Animal Science and Production, Botswana College of Agriculture P/Bag 0027, Gaborone, Botswana.

⁴ Okavango Research Institute, University of Botswana, Private Bag 285, Maun, Botswana

⁵ Sustainability Research Institute, School of Earth & Environment, University of Leeds, Leeds, West Yorkshire LS2 9JT, UK

⁶ School of Geography, University of Leeds, Leeds, West Yorkshire LS2 9JT, UK

*Corresponding author, e-mail: m.reed@abdn.ac.uk.

Abstract

It is increasingly recognized that tackling land degradation through more sustainable land management (SLM) depends on incorporating multiple perspectives using a variety of methods at multiple scales, including the perspectives of those who manage and/or use the land. This paper reports experience implementing a previously proposed methodological framework that is designed to facilitate knowledge sharing between researchers and stakeholders about land degradation severity and extent, and SLM options. Empirical findings are presented from the Botswana site of the EU-funded Desertification and Remediation of Land (DESIRE) project. The paper reflects upon the challenges and benefits of the proposed framework, and identifies a number of benefits, notably related to insights arising from the integration of local and scientific knowledge, and the ownership of the SLM strategies that emerged from the process. However, implementing the framework was not without challenges, and levels of poverty and formal education may limit the implementation of the framework in some developing world contexts.

1 Introduction

Global environmental problems demand interdisciplinary assessment. Although this is now widely recognised by academics (e.g. Kates et al., 2002; Lélé and Norgaard, 2005) and policy-makers (UNCBD, 1992; UNCCD, 1994), there is little consensus about how this can best be achieved. The challenge of marrying results about environmental change from different disciplines is further compounded by the difficulty of reconciling potentially conflicting perceptions among and between researchers and different stakeholders. This article will ask whether we can disentangle the complex arguments that characterise these complex relationships, to arrive at an assessment of environmental change that can effectively learn from these diverse forms of knowledge.

This challenge is particularly acute in the assessment of land degradation¹. Land degradation is an anthropocentric concept that can only be defined in relation to the objectives of those who use and manage the land: “land degradation is contextual” (Warren, 2001:449). For land degradation assessment to be accurate and reliable, it must therefore incorporate multiple perspectives using a variety of methods at multiple scales, including the perspectives of those who manage and/or use the land (Reed *et al.*, in press).

To address this challenge, Reed *et al.* (2011) present a methodological framework for land degradation and sustainable land management (SLM) monitoring and assessment (Figure 1). The framework integrates approaches used by the Desertification Mitigation and Remediation of Land (DESIRE) project, the World Overview of Conservation Approaches and Technologies (WOCAT), the Dryland Development Paradigm (DDP), and the UN Food & Agriculture Organisation’s UNEP/GEF-funded Land Degradation Assessment in Drylands (LADA) project. It groups methodological steps under four broad themes:

¹ Land degradation: i) is a human-induced phenomenon that cannot be caused by natural processes alone; ii) decreases the capacity of the land system as managed to meet its user demands; and iii) threatens the long-term biological and/or economic resilience and adaptive capacity of the ecosystem. This definition is based on a synthesis of: Holling, 1986; Abel and Blaikie, 1989; UNEP, 1992; Turner & Benjamin, 1993; UNCCD, 1994; Dean et al., 1995; Kasperson et al., 1995; UNEP, 1997; Holling, 2001; and IPCC, 2001.

- i) Establishing land degradation and SLM context and sustainability goals;
- ii) Identifying, evaluating and selecting SLM strategies;
- iii) Selecting land degradation and SLM indicators; and
- iv) Applying SLM options and monitoring land degradation and progress towards SLM goals.

This methodological framework is now being applied and evaluated through the DESIRE project in 16 of the most degraded drylands of the world, representing a wide range of land degradation processes and environmental, socio-cultural, economic and policy contexts. One of these sites is Botswana, which has been described as “one of the most desertified countries in sub-Saharan Africa” (Barrow, 1991: 191).

This paper for the first time reports experience implementing the DESIRE/WOCAT/LADA methodological framework proposed by Reed *et al.* (2011), using the DESIRE study site in Botswana to critically evaluate its application. Elements of the framework have been tested elsewhere (e.g. Reed *et al.*, 2007, 2008; Schwilch *et al.*, 2009, 2011), but this is the first time that all the elements brought together by Reed *et al.* (2011) have been applied and evaluated. It starts by exploring the case study context, including an assessment of key constraints to sustainable land management. It goes on to describe the SLM strategies selected by local stakeholders for field trials, and the indicators chosen to monitor progress in their efforts to tackle land degradation. The paper ends by reflecting upon the practical challenges of applying the framework, in particular reconciling multiple perceptions of environmental change.

[Figure 1 here]

2 Case study context

The Boteti study site is situated in the Central District of Botswana. Within Boteti, the focus is on the villages of Mopipi, Mokoboxane (Figure 2) and Rakops (not in Figure 2) with an estimated combined area of 3,000 sq km. This study site is regarded as a

1
2
3 'desertification' hotspot and was the focus of Botswana's 1993 case study for the
4 Intergovernmental Convention to Combat Desertification (INCD), and is identified on the
5 GLASOD map as an area of extreme human induced wind erosion (Government of
6 Botswana, 1994). The Boteti area has consequently been a foci of efforts to "combat
7 desertification" and from 2002 was one of the sites for the Indigenous Vegetation Project
8 (IVP), a five-year Botswana Government-GEF-funded pilot project for "community-
9 driven rehabilitation of degraded rangelands"². DESIRE builds on these past and on-
10 going efforts, which are all in line with the recently adopted National Action Programme
11 to Combat Desertification (Department of Environmental Affairs, 2006).
12
13
14
15
16
17
18
19
20

21 [Figure 2 here]
22
23
24

25 The climate is semi-arid and the rains are concentrated in the summer season
26 (October to April). The average rainfall is about 350 mm/yr and has a variability of about
27 38%. The rate of evapotranspiration for the area is taken to fall within the Botswana
28 average of 2000 mm/yr. The drought cycle in the area is estimated at 9-15 years
29 recurrence interval (Ministry of Agriculture, 1994).
30
31
32
33
34

35 The Boteti River spans two major ecosystems, the Makgadikgadi to the north east
36 and the Kalahari System to the south-west and has undergone pronounced change over
37 the last thirty years. Water from rains in the Angolan highlands pass through the
38 Okavango Delta and reach the lower Boteti in the dry season, providing essential surface
39 water and access to the surrounding forage (plains grasslands and riparian zones) for wild
40 ungulates and domestic animals. As part of wildlife seasonal migration, zebra move from
41 the Pans in the wet season to the Boteti River in the dry. About 80-90% mortality of the
42 Makgadikgadi zebra population in the 1982-86 resulted from drought, partly exacerbated
43 by competition with livestock (over 80,000 animals died). Other wildlife (e.g. wildebeest
44 and hartebeest) and livestock suffered a similar fate.
45
46
47
48
49
50
51
52

53 A major grievance of the communities in the area is that they do not benefit from
54 wildlife related tourism, but endure crop and livestock losses from wild predators.
55
56

57
58 ² <http://www.indigenousvegetation.net>
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Consequently, an electrified game-proof fence was erected along the Boteti in 2004 to try and reduce predation. Seasonally intense grazing by mobile populations of wildebeest and zebra on the western plains area, including Lake Xau, has today been replaced by permanent grazing by livestock facilitated by intensified borehole and shallow well drilling.

The study site is located on tribal land and falls under the responsibility of the Ngwato Land Board. Tribal land in Botswana is communally owned; hence its use is also communal except where individuals or groups of people have been granted exclusive rights to use a particular piece of land. Most livestock are kept at cattle posts, located away from villages to avoid conflict with other land uses such as crop production. Indeed, in the 1970s and part of the 1980s, Mopipi was a fishing and “molapo” farming village, according to the villagers. With the loss of flow along the Boteti River since the late 1980s, the people converted the productive flood recession “molapo” agricultural system along the fluvisols of the Boteti river to the inherently more risky rain-fed production. Since then arable agriculture has also expanded into grazing land to the west of Rakops and north of Mokoboxane slightly reducing land available for livestock and wildlife grazing.

Provision of water is by drilling boreholes, and farmers do not have exclusive rights to either water or grazing resources. However, farmers can apply for exclusive rights, and a block of leasehold farms are located south of Mopipi. These farms were allocated under the Tribal Grazing Land Policy of (1975) and the revised/new Policy on Agricultural Development of 1991 whose aim was to primarily reduce grazing pressure from the communal areas, improve grazing management and increase livestock production. According to the Botswana Poverty Map, the Boteti District population falls within the 40-50% headcount poverty level making it the third poorest region after Ngamiland west and Kgalagadi South Sub-Districts (CSO, 2008). Livelihood sources in the area include livestock rearing, arable farming, casual employment, government support, formal employment, and gathering of veld products. Although hunting and gathering was an important source of protein and food for many households in the recent past, due to the drastic decline of wildlife populations at the end of the last century, pastoralism is now by far the most important source of livelihoods in the study area. The

1
2
3 agropastoral system and practice in Botswana separates crop fields and livestock
4 completely during cropping season (October-June) and at night for almost the whole year.
5 Arable farmers who use cow dung to improve soil fertility let the cows graze in the fields
6 for about six months after harvest. Even these however would still kraal their cattle at
7 night to be able to get milk and protect them from stock theft and predators. Cow dung
8 for bio-gas will be mainly collected from watering points and kraals. For these reasons
9 there is no competition for cow dung.
10
11
12
13
14

15
16 Fishing, as indicated earlier, ceased to be an important source of livelihood in the
17 late 1980s when the Boteti River stopped receiving water from the Okavango delta
18 system. Although the floods have returned to the system today, fishing has been slow to
19 re-establish. Livelihoods are supplemented by harvesting a variety of rangeland products,
20 including building poles, thatching grass, firewood, wild fruits and vegetables, medicine
21 and Mopane worms³. Although these products are used mainly for subsistence, Reed
22 (2005) found that some families derived significant income, for example from the sale of
23 traditional beer brewed from wild fruits. Reed (2005) found that thatching grass was
24 increasingly hard to find in the study area and that firewood was a problem for families
25 without access to transport. It was found in some workshops conducted under the
26 DESIRE project that all other rangeland products have been commercialized to some
27 extent.
28
29
30
31
32
33
34
35
36

37
38 Until the 1930s, the study area was sparsely populated and Basarwa (San) were
39 the predominant ethnic group. The Bakalanga, who migrated into the area in the 1930s,
40 are now the largest ethnic group in the area. Population density is still low (estimated at
41 about 1 person per square kilometer). However, the population in the study sites grew
42 significantly during the 1991-2001 decade. The population of Rakops grew from 3122
43 inhabitants in 1991 to 4555 in 2001. Mopipi village population grew from 2264 to 3066
44 inhabitants during the same period. Mokoboxane registered the highest growth rate from
45 614 inhabitants in 1991 to 1290 by 2001. Around 40% of the population of the study area
46 is under the age of 15. Furthermore the area has a high proportion of adults above the age
47 of 20 who have never been to school, with significantly more illiterate women than men.
48
49
50
51
52
53
54
55
56

57
58 ³ The caterpillar of the Mopane Moth, *Imbrasia belina*.
59
60

1
2
3 For instance, in Mopipi 41% of women compared to 34% of men in this age bracket had
4 never been to school (Republic of Botswana, 2004).
5
6
7
8
9

10 **3 Methods**

11
12 The following text will describe the methods used to apply the framework described in
13 Figure 1, structured by each of the steps in the framework.
14
15
16

17 *3.1 Establishing land degradation and SLM context and sustainability goals*

18
19 Identifying system boundaries, stakeholders and their goals (Figure 1, step 1): A detailed
20 account of the environmental, socio-economic and policy context in which land
21 degradation and SLM occur in this area are provided in section 2 (see also Atlhopheng et
22 al., 2009 on DESIRE website: <http://www.desire-his.eu>). System boundaries,
23 stakeholders and their goals were determined through Workshop 1 with land users,
24 conducted following the WOCAT approach described in Bachman et al. (2007a&b). The
25 participatory exercises established not only sustainability goals, but also the influence of
26 various stakeholders and institutions on local land management (cf. Magole et al. 2008).
27
28
29

30
31 Describing the socio-cultural, economic, technological, political and environmental
32 context and identifying key drivers of change (Figure 1, step 2): This was also done
33 during Workshop 1 through an exercise that elicited from land users land degradation
34 cause-effect linkages or impact chains (cf. Magole et al., 2008). This information was
35 supplemented through literature (cf. Atlhopheng et al., 2009 on DESIRE website:
36 <http://www.desire-his.eu>) and policy reviews (cf. Chanda et al., 2009a; Atlhopheng, et al.
37 2009a; Mulale and Chanda, 2009).
38
39

40
41 Determining current land degradation status, future land degradation risk and existing
42 SLM measures using existing indicators (Figure 1, step 3): This was done in two ways.
43 First, participatory methods were used to identify indicators that could measure progress
44 towards sustainability goals, and that could be used easily by communities themselves.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 This was done through Workshop 1, reflecting on land degradation indicators participants
4 had observed during a transect walk, which took place prior to the workshop. This built
5 on previous participatory indicator development by Reed et al., (2008). Second, a more
6 standardised procedure adopted across the DESIRE project (which had previously been
7 used to determine land degradation indicators in the Mediterranean region by the
8 DESERTLINKS project) was used to survey indicators in the most important land use
9 types in Boteti (agriculture, pasture, woodland/forest, settlement, pan/dam). A structured
10 indicator questionnaire was completed for each process deemed significant in a land use
11 type using appropriate weighing indices (cf. Chanda et al., 2009b). The weighing indices
12 provided the quantitative data for subsequent statistical analysis and determination of
13 significant degradation indicators in land use types.
14
15
16
17
18
19
20
21
22
23

24 *3.2 Identifying, evaluating and selecting SLM strategies*

25 Identifying, assessing and prioritising possible SLM options (Figure 1, steps 4 and 5):

26 Potential SLM options were listed by participants during Workshop 1, and evaluated
27 using WOCAT standard questionnaires and then during structured small group
28 discussions about the strengths and weaknesses of each option (cf. Magole et al., 2008).
29 SLM options deemed appropriate by participants in Workshop 1 were further investigated
30 by the research team (e.g. cost, current use in Botswana, etc), and then further evaluated
31 during Workshop 2 using Multi-Criteria Evaluation. After reviewing the top-ranked
32 remediation options from Workshop 1, participants discussed and agreed on evaluation
33 criteria and scored each SLM option using 'facilitator software' (see Magole et al., 2009).
34
35
36
37
38
39
40
41
42
43
44

45 Trial SLM options at field scale (Figure 1, step 6): Although the most preferred SLM 46 strategy was community game ranching, on account of affordability the land users settled 47 for biogas production and use, with local cow dung as the main input to the gas 48 production process (see section 4 for more details). The plan for implementing SLM 49 options in Boteti therefore centres on piloting the impact of household biogas use on 50 household fuelwood consumption and woody biomass conservation. This entails 51 monitoring the rate of fuel wood consumption by a family utilizing biogas as the primary 52 53 54 55 56 57 58 59 60

1
2
3 energy source compared to a control family of similar characteristics still dependent on
4 firewood as the primary energy source. It will also be necessary to establish baseline
5 information on current (i.e. pre-biogas) household energy options, fuel wood
6 consumption rates, estimates of current (pre-biomass) standing woody biomass, as well as
7 estimates of available feedstock (cow dung) for the biogas plant. Current household
8 energy options were captured through 42 household surveys using a structured
9 questionnaire, conducted during winter (July to September) of 2009. Fuelwood
10 consumption rates were estimated from a 2-week monitoring and measurement of
11 firewood supply and use using a spring scale for a subsample of surveyed families in
12 Mopipi village. With regard to estimating feedstock for biogas production, two kraals
13 (cattle-posts) were chosen to estimate dung produced by cattle over-night. Each morning,
14 farmers usually allow their cattle to go out to graze leaving their calves behind. In the
15 evening, when they return to the water point they are kraaled – although in good rainfall
16 seasons many animals do not return to the water point but stay out to graze. With the
17 consent of two kraal owners ten ‘standard’ fresh dung pats were weighed after night time
18 kraaling over a period of three days. Assessment of woody biomass was made within a
19 radius of 12 km from Mopipi using stepwise sampling points at distances of 2, 4, 6, 8, 10
20 and 12 km from the village. At each sampling point, a cluster of three quadrats of 50m x
21 10m each were located with the aid of tapes and ranging poles. Coordinates of each
22 quadrat were recorded. A boundary of 100m between quadrats was allowed. Individual
23 rooted woody species within the quadrat were identified at species level; their stem
24 diameters (at ankle height) and woody biomass estimated via the related regression
25 equations developed by Tietema (1993).
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 Up-scale/aggregate biophysical and socio-economic effects of SLM from field to regional
47 and national scales to further prioritise SLM options (Figure 1, step 7): Experimental
48 work as done in the previous step can give valuable information to evaluate a technology,
49 but cannot easily be generalized to a larger scale, nor can scenario analyses be made to
50 answer ‘what-if’ questions. In DESIRE, an integrated environmental modelling approach
51 was adopted to address these issues (for details see Fleskens et al., 2009). For the case of
52 biogas, modelling is focussing on the economic and environmental effects of widespread
53
54
55
56
57
58
59
60

1
2
3 adoption of biogas in the Mopipi and Mokoboxane villages. In this paper, a simplified
4 preliminary economic evaluation is presented. Investment costs for a biogas installation
5 were estimated by participants in Workshop 2 (above). Potential collection sites for cow
6 dung were assessed in relation to the demand for dung if all households would switch to
7 biogas. The collection sites are assumed to be adjacent to boreholes, of which there are
8 378 in the study area. The price (transport and labour costs) for dung collection was
9 explicitly taken into account. Moreover, a bi-annual maintenance of the biogas
10 installation was assumed and estimated to be 5% of the investment costs. The economic
11 life of the biogas installation was set at 20 years, and the discount rate for cost-benefit
12 analysis was set at 10%, reflecting the local opportunity cost of capital. Importantly,
13 biogas releases time (and perhaps financial resources) that would otherwise be spent
14 collecting firewood. The cost of firewood is in the current analyses a model output, i.e.
15 the daily average household firewood requirements are priced at break-even point (using
16 10% rate of internal return).

17
18 The biophysical effects will be simulated with the PESERA model (Kirkby et al.,
19 2008). Biophysical models offer an opportunity to merge physical and social constraints
20 on the production and availability of wood and grazing biomass. Once established, the
21 model can be used to explore the spatial interaction between fuel demand, fuel production
22 (biomass and dung) and land degradation under current and SLM practice. This
23 modelling approach allows an assessment of the scale at which biogas offers resilience to
24 the natural system with uncertainty in cattle numbers and rain fed biomass production.
25 Monitoring and observation of current rates and patterns of biomass removal (fuel and
26 grazing) provide essential data when considering modelling the bio-mass balance for the
27 local area or region (Figure 3). Such data, combined with land use, vegetation data and
28 climate are the primary input into bio-physical models. The biophysical component of the
29 Pan-European Soil Erosion Assessment (PESERA) model. PESERA is a process-based
30 model that is designed to estimate long term average erosion rates at 1 km resolution and
31 has been applied in locations across Africa, Latin America and Asia through the DESIRE
32 project. The model is built around a partition of precipitation into components for
33 overland flow (infiltration excess, saturation excess and snowmelt), evapo-transpiration
34 and changes in soil moisture storage. Transpiration is used to drive a generic plant

1
2
3 growth model for biomass, constrained as necessary by land use decisions, primarily on a
4 monthly time step. Leaf fall, with corrections for cropping, grazing etc, also drives a
5 simple model for soil organic matter. The runoff threshold for infiltration excess
6 overland flow depends dynamically on vegetation cover, organic matter and soil
7 properties, varying over the year. The distribution of daily rainfall totals has been fitted
8 to a Gamma distribution for each month, and drives overland flow and sediment transport
9 (proportional to the sum of overland flow squared) by summing over this distribution.
10 Total erosion is driven by erodibility, derived from soil properties, squared overland flow
11 discharge and gradient; it is assessed at the slope base to estimate total loss from the land,
12 and delivered to stream channels. The combined biophysical and biomass offer the
13 potential to consider resource management options under future scenarios and inform
14 further discussions between the study site researchers and stakeholders.
15
16
17
18
19
20
21
22
23
24
25

26 [Figures 3 & 4 here]
27
28
29
30
31

32 *3.3 Selecting land degradation and SLM indicators*

33
34

35 Finalise selection of indicators (in collaboration with likely users) to represent relevant
36 system components for ongoing monitoring by land managers (Figure 1, step 8):
37

38 Indicators identified during Workshop 1 (see section step 3, section 3.1) were evaluated
39 and selected on the basis of their ease of use (i.e. the ability of land managers to measure
40 them) and capacity to represent land degradation processes or the long-term sustainability
41 of land management (cf. Magole et al., 2008 and Bachman et al., 2007a&b). This built on
42 previous work by the team where indicators that had been identified in interviews were
43 evaluated and shortlisted in workshops using Multi-Criteria Evaluation and further
44 evaluated using field methods (Reed et al., 2008).
45
46
47
48
49
50
51

52 *3.4 Promoting the application of SLM options and monitoring land degradation and* 53 *progress towards SLM goals* 54 55 56 57 58 59 60

1
2
3
4 Disseminate strategies and indicators for extension and national and international policy
5 (Figure 1, step 9): Although literacy levels are relatively high in Botswana (average 81%
6 according to Central Statistics Office (2004)), Boteti land users have no easy access to
7 computers and the internet. Consequently, the most appropriate means of disseminating
8 information to them would be through workshops, translated leaflets, manuals and
9 posters. Participatory workshops have already proven to be an effective means of local
10 stakeholder engagement. However, for national policy development and implementation,
11 web-based dissemination would be possible. SLM options from this project and previous
12 work (Reed and Dougill, 2010) were integrated, and linked to relevant indicators of land
13 degradation and SLM in a manual, targeted at land managers and extension workers. In
14 addition to biogas and game ranching, the manual includes a range of other SLM options
15 deemed relevant to the study area through previous work (Reed and Dougill, 2010), such
16 as changing livestock breeds, shifting grazing, managing bushes and various soil
17 management techniques. These materials and other more detailed results from the project
18 are also available via the DESIRE project's "Harmonised Information System", available
19 at: <http://www.desire-his.eu/>.
20
21
22
23
24
25
26
27
28
29
30
31

32
33 Apply SLM strategies, monitor degradation and progress towards SLM goals, upscaling
34 or aggregating to district and national levels (Figure 1, step 10): The results of the biogas
35 project with respect to the environmental and socio-economic objectives for which it was
36 adopted will be reviewed with land managers as soon as results become available.
37 Depending on its impact on woody vegetation conservation and socio-economic welfare,
38 the strategy could be implemented more broadly within the study village and in areas
39 within the region and beyond with similar circumstances. Ongoing monitoring using
40 indicators by land managers is facilitated through the dissemination of manuals linking
41 indicators to SLM options (see previous step). These manuals are being used as part of
42 the wider Government of Botswana's implementation of its UNCCD National Action
43 Plan, under which it is hoped that these activities can be up-scaled to district and national
44 scales, and progress towards SLM goals can be monitored nationally.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Adjust strategies to ensure goals are met (Figure 1, step 11): Finally, it should be noted
4 that as goals are met and contexts change, it may be necessary to develop or prioritise
5 new SLM strategies and indicators in future. Consequently, this framework is iterative,
6 represented by the dashed arrow between steps 11 and 4.
7
8
9

10 11 12 13 14 **4 Results**

15 16 17 *4.1 Establishing land degradation and SLM context and sustainability goals*

18
19
20
21 Figure 5 shows land degradation drivers and desertification risk according to structured
22 indicator questionnaires that were completed for each process deemed significant in each
23 of the land use types found in the study area (Figure 1, step 3, section 3.1 above). Boteti
24 stakeholders identified sustainability goals that addressed ecological (environmental)
25 integrity (as suggested by the objective of reducing the depletion of trees), economic
26 (livelihood) security (as suggested by the objective of minimizing the impact of drought
27 on arable production) and social equity (as indicated by the objective of reducing
28 poverty). These sustainability goals were a response to problems that are well
29 documented in the literature in this study area. For instance, the Boteti area has had the
30 highest proportion of permanent destitutes among the 5 sub-regions of the Central
31 District of Botswana (Central District Council, 2003). However, despite the many studies
32 that have identified rangeland degradation as a problem in this area (e.g. Ministry of
33 Agriculture, 1993; Ringrose et al., 1996; Perkins, 2007; Chanda, et al., 2007),
34 environmental sustainability goals focussed on the arable system and the availability of
35 water and fuel-wood.
36
37
38
39
40
41
42
43
44
45
46
47
48

49 [Table 1 here]
50
51
52

53 This may reflect a number of potential problems with the research that initially
54 identified rangeland degradation. In particular, Ringrose et al. (1996) inferred land
55 degradation from the presence of three indicators: (1) increased wind erosion and dust
56
57
58
59
60

1
2
3 storms; (2) *localised* permanent damage to vegetation around villages and along the river;
4
5 (3) die-back of riverine forest (it is assumed due to lowering of water table or salinisation
6
7 due to unsustainable water extraction). It is clear that the riverine system has been
8
9 degraded in Boteti, due mainly to the loss of the Boteti flows, with losses in fishing and
10
11 flood plain agriculture. However there is less evidence to support the claim that grassland
12
13 resources in this area are degraded, and not simply responding to drought. Although
14
15 increased wind erosion may be a degradation indicator, it is also an indicator of drought.
16
17 Evidence of wind erosion features from remotely sensed data may simply indicate
18
19 drought conditions. Ringrose *et al.* (1996) detected wind erosion features from remote
20
21 sensing data collected in 1984, 1989 and 1993. Two of the three years were drought
22
23 years.

24
25 Permanent changes are evident around settlements, however these changes are
26
27 highly localised in extent and do not suppress the productivity of the livestock system.
28
29 Reed (2005) found that grass cover was high 2-4 km from villages in Boteti. The
30
31 dominant grass species, *Cynodon dactylon* was classified as “high grazing value” by van
32
33 Oudlshoorn’s (1999). In common with Ringrose *et al.* (1996), community members cited
34
35 a range of environmental problems, which they blamed predominantly on drought (Reed,
36
37 2005). When probed about the capacity of the land to recover after rain, the majority of
38
39 respondents interviewed by Reed (2005) emphasised the resilience of the grassland zone,
40
41 noting that rainfall produced sufficient fodder to maintain herds for at least two years. As
42
43 such few land users were constrained by their natural capital. However, community
44
45 members agreed that the mopane veld zone, which is less important for livestock
46
47 production, had experienced a decline in productivity. This was due to increased
48
49 dominance of *C. mopane* in response to grazing, leading to the suppression of grass.
50
51 Elsewhere in the mopane region of Botswana, the species is highly valued for forage.
52
53 However, elsewhere in Botswana it is usually valued in relation to available alternatives,
54
55 which tend to comprise less palatable annual grasses (Ringrose, pers. comm.), and it is
56
57 inferior to forage in the grassland zone in Boteti.
58
59

60
61
62 In conclusion, the riverine system in the study area is clearly degraded due to
63
64 declining river water levels and heavy stocking pressure within the riparian woodland.
65
66 The absence of flows along the Boteti from the late 1980s until recently has significantly

1
2
3 constrained livelihood options for the local community, who once relied on fishing and
4 flood plain agriculture. Localised rangeland degradation is occurring around villages and
5 in the mopane veld zone. At present levels, this does not appear to threaten the
6 sustainability of livestock production in Boteti, but there are major concerns over fuel-
7 wood availability and water shortages. These concerns are reflected in the identification
8 and selection of SLM options that emerged from the two workshops.
9
10
11
12
13

14 15 16 *4.2 Identifying, evaluating and selecting SLM strategies*

17
18
19 Altogether, land users identified 13 SLM strategies during Workshop 1, out of which they
20 prioritized 4, viz: biogas as an alternative energy source, community wildlife (game)
21 farming, irrigation and dam-building (water harvesting). The DESIRE team investigated
22 three of these (biogas, water harvesting and game ranching) and another technology not
23 mentioned by the land users (solar power via solar cooker). In order to help them make
24 informed decisions about the technologies, the findings of the investigations were shared
25 with land users during Workshop 2. The latter four technologies were evaluated as
26 outlined under section 3.2 above. As Figures 6-8 below indicate, the technologies with
27 the highest beneficial environmental (ecological), economic and socio-cultural impacts
28 according to land users were community game ranching and biogas (cf. Magole et al,
29 2009). Figure 6 shows that there was considerable disagreement over the ecological
30 benefits of game ranching (indicated by the length of the horizontal line) compared to
31 biogas. Figures 7 and 8 show that there was considerable agreement however, over the
32 economic and socio-cultural benefits of game ranching, and that these benefits were
33 considered greater for game ranching than for other technologies.
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 The land users' prioritization of game ranching was based on a number of social
49 and environmental considerations. Game ranching was perceived to have the potential to
50 bring economic returns for poverty alleviation (Table 1) and also to be a more viable
51 alternative to livestock farming which causes overgrazing. They reasoned that wildlife
52 species are better adapted to the semi-arid environment which they utilize more optimally
53 as grazers and browsers. On the other hand, biogas production and use was expected to
54
55
56
57
58
59
60

1
2
3 reduce the need to collect firewood which is the main source of energy in the study area.
4
5 The stakeholders reasoned that by using biogas one can save time, use less labour and
6
7 save trees. Although it was marginally less popular than game ranching, the land users
8
9 ultimately settled for the biogas production strategy because the initial set-up costs were
10
11 lower, and hence could be implemented within the funding and time available within the
12
13 DESIRE project.

14
15
16 [Figure 6-8 here]

17 18 19 *4.3 Selecting land degradation and SLM indicators*

20
21
22 During Workshop 1, land users identified a wide array of indicators of both land
23
24 degradation and SLM (Table 2). Participants considered the positive indicators as
25
26 depicting past environmental conditions, confirming findings from earlier studies that the
27
28 people of Boteti perceive a long-term deterioration in their environment which, as
29
30 observed earlier, they readily link to climate desiccation and failure of Boteti floods (cf.
31
32 Chanda, 1996; Penning de Vries, 2007; Chanda and Darkoh, 2007).

33
34
35 [Table 2 here]

36 37 38 39 40 *4.4 Applying SLM options and monitoring land degradation and progress towards SLM* 41 42 *goals*

43
44
45 Monitoring of environmental and socio-economic impacts of adopting biogas as an
46
47 alternative domestic energy source is ongoing. However, the following baseline data has
48
49 been generated using the methods described under section 3.2 above:

- 50
51 ● There is indeed a very high dependency on firewood as a source of energy for
52
53 cooking (100% - Mopipi; 98.4% Mokoboxane), space heating (77% - Mopipi; 96% -
54
55 Mokoboxane), warming bath water (98.7% - Mopipi; 98.4% - Mokoboxane) and
56
57 various family events or ceremonies (78.5% - Mopipi; 83% - Mokoboxane);
58
59
60

- There has been mounting scarcity of firewood within more accessible areas (86% of Mopipi and 90% of Mokoboxane respondents). This is supported by the increasing use of donkey cart and motor vehicles in firewood collection, especially in the larger village – Mopipi (88.9% of respondents);
- On average, families use 10kg of firewood per day;
- Most firewood is collected from communal land predominantly (not solely) lying in the easterly direction;
- There is enough cattle dung in the Mopipi area to support domestic biogas production on a sustainable basis without compromising its use as organic fertiliser in arable agriculture. The total dung output per animal per night was estimated at 5.4 kg wet weight; and
- Biomass of both live and dead fuel wood increases linearly with distance from the village. Biomass of live trees (25 989 kg per ha average) was far greater than that of dead wood (919 kg per ha average), suggesting depletion of the latter stock as people currently depend on dead, rather than live, wood for energy. *Colophospermum mopane* (the most preferred firewood species) contributed the most biomass of live tree species and, expectedly, the least biomass to dead tree species – underscoring its popularity as a firewood resource.

These results portray a situation of a decreasing stock of dead tree biomass for firewood within collection zones. Thus the land users' choice of biogas production as an alternative energy source could be viewed as a proactive move intended to pre-empt cutting of the now more accessible live trees for firewood.

Regional assessment of introducing biomass was elaborated based on the above indicators. Figure 9 shows the location of boreholes as potential dung collection sites. To simplify the analysis, we classified boreholes based on distance into 5 classes: less than 1 km (1.6%); 1-2km (0.5%); 2-5 km (4.0%); 5-11 km (18.8%); 11-20 km (41.5%); and more than 20 km (33.6%). Importantly, dung collection from these sites is assumed to be undertaken on foot (in the case of the first class), by cart (next two classes), or by van (the remaining classes).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The results from scenarios of introducing biogas are shown in Table 3. If we assume a uniform distribution of cattle over boreholes (potential collection sites), all dung required for operating biogas installations for the 872 households inhabiting Mopipi and Mokoboxane will be collected from within a range of 11 km from the villages; note that in this scenario (A) the six boreholes at less than a kilometre from the villages are disregarded as dung collection on foot is uneconomical. Biogas is financially attractive if firewood collection is about twice as costly as collecting dung. In view of less attractive rangelands close to the villages, it is more likely that cattle is kept away from the village.

If we assume all cattle is held beyond 11 km (scenario B), dung collection costs for operating all biogas installations will rise by about 24%. The cost of firewood collection at which biogas becomes a viable investment rises by about 12% relative to Scenario A. Similarly, if cattle is kept at more than 20 km (Scenario C), dung collection costs almost double (+81%) but firewood opportunity costs would need to rise 40% in order to make the investment viable. Neither of Scenarios A-C is likely to be correct; however these analyses show that cattle herding dynamics (e.g. in response to drought) do not have to form an obstacle to adopting biogas.

Scenario D addresses what happens if the biogas installations do not perform (or are not managed) efficiently and can only fulfil 75% of energy demand, with the remaining 25% continuing to be provided by firewood. This obviously renders the investment in biogas less attractive (i.e. a higher firewood opportunity cost is needed). Importantly, the assessment method employed requires the share of firewood energy substituted by biogas to be larger than 50% - any share lower than that means the technology will never be viable from an individual decision-making perspective.

Scenario E addresses the fact that firewood collection costs may not yet be as high as to push people to other alternatives, but may be on a rising trend. If we assume an annual increase of 10%, reflecting both increased scarcity (and thus time) and distance (transport and time) of firewood collection, adoption of biogas can anticipate these future changes and be viable even if the current cost of firewood collection is lower than dung collection costs.

1
2
3 Scenario F represents Scenario A but assuming cattle dung from collection areas
4 close to the village would not involve opportunity costs of labour. This drops the cost of
5 dung collection and the minimum required opportunity cost of firewood collection to
6 67% and 60% of Scenario A respectively. In fact, taking the perspective of a pioneering
7 adopter of biogas who would exclusively collect dung from a nearby collection area
8 (from his/her own kraal for instance), the tipping point for opportunity cost of firewood
9 collection would go down further to P6.98 (36%).
10
11

12
13
14
15
16 Finally, scenario G reflects the extremely high cost and tipping point when dung
17 collection needs to be done on foot – taking into account labour opportunity costs.
18 Although this situation may be highly theoretical (a biogas adopter who is not in a
19 position to source dung using more economical means of transport would probably not
20 invest in the installation in the first place), it does show that there is considerable scope
21 for some entrepreneurs to create a local dung market and organize supply of dung to
22 biogas installations, or to develop a biogas market. The latter does however require
23 additional investment for pressure filling equipment and suitable cylinders to transport
24 biogas.
25
26
27
28
29
30
31
32

33 [Table 3 here]
34
35
36
37

38 **5 Discussion**

39
40 This paper builds upon a number of previous studies conducted in this area, and through
41 the integration of local and scientific knowledge, has provided a number of novel insights
42 that contrast with these previous studies. Most notably, the present study challenges the
43 nature and causes of land degradation identified previously, and hence suggests quite
44 different options for tackling land degradation in the study area. The study started by
45 identifying types and causes of land degradation with local communities, rather than
46 inferring this via natural science assessments of land degradation indicators. In contrast to
47 previous research based primarily on remote sensing (e.g. Ministry of Agriculture, 1993;
48 Ringrose et al., 1996), and studies by Reed et al. (2007, 2008) which focussed on
49 rangeland degradation, participants in this study emphasised problems associated with
50
51
52
53
54
55
56
57
58
59
60

1
2
3 poverty, and degradation processes related to the depletion of fuel wood and water
4 shortages. In contrast to previous studies, which had a strong environmental and
5 rangeland focus, the present study considered land degradation and SLM in the broadest
6 possible context. In this way, although localised rangeland degradation issues were
7 identified (with associated indicators) they were not given undue importance, and other
8 issues were prioritised, leading to the selection of SLM options for trial that were
9 focussed on tackling poverty and fuel wood shortages.

10
11
12 In particular, the scenario analyses demonstrate considerable scope for biogas
13 production. The currently felt scarcity of firewood seems to indicate that the resource is
14 being overexploited. With 10 kg of firewood needed for each household each day, and
15 using the average of 919 kg/ha of dead wood biomass observed, annually at least an area
16 the size of 3460 ha is needed to satisfy the demand. This involves a 3.3 km search radius
17 from the villages. However, the current level of availability of firewood is already much
18 lower than average close to the village, and the harvesting of all dead wood is unlikely to
19 be sustainable. Therefore, it seems safe to assume the real search radius for firewood is
20 already much larger. Unlike dung, which is collected from kraals, firewood needs to be
21 collected from highly disperse areas, so that time and transport requirements are likely
22 much higher than for dung.

23
24
25 Simulations with the biophysical PESERA model will provide valuable insight in
26 the environmental effects of both firewood and dung collection dynamics. However,
27 ultimately an important social aspect is involved. As firewood is collected by women and
28 girls, perhaps at a perceived opportunity cost of zero, whether biogas will be taken up
29 depends on community recognition of the importance of freeing them of an onerous task.
30 Even when this is recognized, investment in biogas is expensive and beyond the
31 individual capacity to sustain; institutional arrangements are therefore needed, which may
32 be most effective if embedded in national level policy. Apart from financial viability and
33 environmental benefits, the rationale for supporting biogas development can be
34 considerably broadened, including potential industry development when scaling up,
35 energy saving, energy security, and health benefits by reducing smoke-related respiratory
36 diseases.

1
2
3 The use of Multi-Criteria Evaluation makes the reasons why participants selected
4 particular SLM options transparent, and it is interesting to note that despite considerable
5 disagreement over the ecological benefits of game ranching, it was prioritised by
6 participants primarily for socio-cultural and economic reasons. Again, this emphasises
7 the need to consider the socio-cultural and economic aspects of land degradation and
8 sustainable land management alongside environmental dimensions. The contrast between
9 the results of this research and previous research in the area may therefore reflect a
10 difference in the relative priority that researchers and local stakeholders give to these
11 different dimensions of sustainability.
12
13
14
15
16
17
18

19 This further emphasises the need for methodological approaches such as those
20 proposed by Reed et al. (2011) and used in this research, which can integrate local and
21 scientific knowledge of land degradation and SLM. Rather than pitching local knowledge
22 (as collected in this research) against scientific knowledge (as per previous publications
23 in this study area), this research combines local and scientific knowledge of land
24 degradation problems and SLM options. Scientific knowledge contributed to the
25 understanding of these issues via land degradation indicators that were developed by
26 researchers and evaluating and applying SLM options (such as biogas) suggested by
27 researchers. Local knowledge then ensured that relevant SLM options were considered,
28 based on a local appreciation of land degradation problems (which differed to the
29 perceptions of most researchers), and only those that could best meet local needs and
30 priorities were implemented.
31
32
33
34
35
36
37
38
39

40 Although for some these different priorities may represent a conflict of interests
41 between researchers and local stakeholders, the participatory approach facilitated a close
42 working relationship between the research team and local communities, which translated
43 into strong local ownership of the adopted SLM strategies. The adopted SLM strategies
44 addressed an environmental problem with a strong link to priority livelihood issues.
45 Thus, environmental conservation has a concrete rather than theoretical or abstract
46 meaning to the lives of the land users. There was a very high land user appreciation of
47 what they referred to as the “DESIRE project process” of solving land use and
48 management problems. Land users unequivocally observed that DESIRE made them
49 realize that they possessed the ability to analyze their socio-economic circumstances and
50
51
52
53
54
55
56
57
58
59
60

1
2
3 ultimately identify expert validated strategies to address locally genuine livelihood and
4 environmental problems. They further applauded the process for promoting cooperation
5 among various stakeholders and for affording them the opportunity to learn about the
6 various remediation strategies from DESIRE experts (see online supplementary
7 material). However, the Boteti experience also exposed a number of challenges to the
8 operationalization of the approach, which may be relevant to other developing country
9 contexts:

- 10
11
12
13
14
15
16
17 i) The high illiteracy rate among land users constrained progress with some of the
18 more technical exercises in the WOCAT process (e.g. impact chain analysis and
19 scoring of the strategies);
20
21
22 ii) High poverty levels among land users meant some remediation strategies were
23 beyond most people's reach, requiring capital outlays (e.g. biogas production),
24 which delayed the piloting of agreed strategies. Indeed, it is on this account that
25 the most preferred (and arguably the most effective) remediation strategy
26 (community-based game ranching) could not be adopted for piloting. The
27 approach assumes that there will be a number of SLM options that are not already
28 being used by land users, that will not require significant capital investment or be
29 associated with significant opportunity costs, and this assumption may not be
30 valid in some of the poorest communities of the world
31
32 iii) WOCAT workshop 2 requires access to electricity to run the 'facilitator
33 software'. While this was not a problem in the Boteti case, it would surely be an
34 obstacle in more remote areas with no or unreliable power supply
35
36
37
38
39
40
41
42
43
44
45
46

47 **6. Conclusion**

48
49
50 The dynamic, context-specific and value-laden nature of land degradation makes it hard
51 to address mechanistically. There can be no simple, universal system for assessing land
52 degradation or identifying relevant SLM options to prevent or tackle the problems it
53 causes. Instead, land degradation assessment must recognise a multiplicity of
54 perspectives, and cannot be judged in isolation from those who face its consequences
55
56
57
58
59
60

1
2
3 (Warren, 2002; Reed et al., 2011) propose a methodological framework for land
4 degradation and SLM monitoring and assessment (Figure 1) that attempts to marry
5 information about environmental change from different research disciplines and
6 stakeholders. This paper shows how this framework has been applied in the Boteti
7 District of Botswana, one of the study sites for the DESIRE project.
8
9

10
11
12 The paper illustrates methods that may be used to operationalise Reed et al.'s
13 (2011) methodological framework, which could be applied in a variety of contexts
14 internationally. It identifies a number of benefits associated with the proposed
15 framework, notably related to insights arising from the integration of local and scientific
16 knowledge, and the ownership of the SLM strategies that emerged from the process.
17 However, implementing the framework was not without challenges. In some developing
18 country situations, the operationalization of the methodology might be restrained by
19 various challenges related to general underdevelopment or, as Penning de Vries (2007)
20 puts it, "capability" problems. As the framework is applied in different contexts through
21 the DESIRE project and elsewhere internationally (e.g. Ravera et al., in press), it will be
22 possible to learn more about how researchers and stakeholders can work more effectively
23 together to monitor and respond to land degradation.
24
25
26
27
28
29
30
31
32
33
34

35 **Acknowledgements**

36
37
38 This research was funded under EU Framework 6 Desertification Mitigation & Remediation of Land – a
39 Global Approach for Local Solutions (DESIRE) project contract no. 037046. It was conducted under
40 Government of Botswana research permit EWT3/3/8 XXXIX (21).and was supported by the University of
41 Botswana. Mark Reed is also funded by a British Academy Research Development Award. Special thanks
42 to the communities of Rakops, Mopipi and Mokoboxane.
43
44
45
46
47

48 **References**

49
50
51 Abel NOJ, Blaikie PM 1989. Land degradation, stocking rates and conservation policies in the communal
52 rangelands of Botswana and Zimbabwe. *Land Degradation & Rehabilitation* 1:101-123.
53
54 Athlhopeng JR, Chanda R, Mphinyane W, Sebego RJ 2009. *Study site description, site 13 – Boteti*.
55 DESIRE research project report. <http://www.desire-project.eu>
56
57
58
59
60

- 1
2
3 Bachman F, Schwilch G, Gabathuler E, Liniger H 2007a. *Guidelines for WB3 Part I: stakeholder workshop*
4 *1. Identification of existing and potential prevention and mitigation strategies*. Centre for Development
5 and Environment (CDE), University of Bern, Switzerland. Report Number 6 of Manuals and Training
6 Guides. In collaboration with DESIRE Project.
7
8
9 Barrow CJ 1991. *Land Degradation: Development and Breakdown of Terrestrial Environments*.
10 Cambridge University Press, Cambridge. Central District Council 2003. Central District Development
11 Plan 6: 2003-2009. Central District Development Committee, Ministry of Local Government, Republic
12 of Botswana.
13
14 Central Statistics Office 2004. *Botswana Literacy Survey 2003*.
15 http://www.cso.gov.bw/html/liter_survey.htm
16
17 Chanda R 1996. Human perceptions of environmental degradation in a part of the Kalahari ecosystem.
18 *Geojournal* **39**: 65-71.
19
20 Chanda R, Darkoh MBK 2007. *Feasibility of community-based rangeland management in Boteti,*
21 *Botswana: the role of tradition and policy*. Component 2 Report for the UB-IVP-CAR collaborative
22 research project.
23
24 Chanda R, Atlhopheng JR, Magole L, Mphinyane W, Mulale K, Perkins JS, Sebego RJ 2009a. *Drivers and*
25 *policy context of land degradation in the Boteti area, Botswana*. DESIRE research report
26
27 Chanda R, Atlhopheng JR, Magole L, Mphinyane W, Mulale K, Perkins JS, Sebego RJ 2009b. *Land*
28 *degradation indicators in Boteti – a triangulated approach*. DESIRE research report
29
30 CSO (Central Statistics Office) 2008. *Botswana Census-Based Poverty Map: District and Sub-district level*
31 *Results*. Ministry of Finance and Development, Gaborone, Botswana.
32
33 Dean WRJ, Hoffman MT, Meadows ME, Milton SJ 1995. Desertification in the semiarid Karoo, South
34 Africa - review and reassessment *Journal of Arid Environments* **30**: 247-264.
35
36 Department of Environmental Affairs 2006. *Botswana National Action Programme to Combat*
37 *Desertification*. Gaborone, Botswana.
38
39 de Vries Penning, L 2007. *Combating desertification: bridging the gap – compatibility analysis of*
40 *Botswana's national strategy to combat desertification with local land user's perceptions and action*
41 *strategies*. Dissertation paper; DESIRE, Wageningen University, Wageningen.
42
43 Fleskens L, Irvine B, Kirkby MK, Nainggolan D, Reed MS, Termansen M 2009. A model that integrates
44 the main bio-physical and socio-economic processes interacting within an agro-ecosystem. DESIRE
45 Report 63 D5.1.1 Aug09. Government of Botswana 1994. *Desertification and possible solutions in the*
46 *Mid-Boteti River area: a Botswana case study for the Inter-governmental Convention to Combat*
47 *Desertification (INCD)*. Final Report, Volume 1 - The desertification process and its impacts. Ministry
48 of Agriculture, Gaborone, Botswana.
49
50 Holling CS 1986. The resilience of terrestrial ecosystems, local surprise and global change. In *Sustainable*
51 *development of the biosphere*, Clark WC, Mtmn RE (eds), Cambridge University Press, Cambridge;
52 292-317.
53
54
55
56
57
58
59
60

- 1
2
3 Holling CS 2001. Understanding the Complexity of Economic, Ecological, and Social Systems. *Ecosystems*
4 **4**: 390-405.
- 5
6 IPCC (International Panel of Climate Change) 2001. *Climate Change 2001: Impacts, Adaptation, and*
7 *Vulnerability*, Cambridge University Press, Cambridge.
- 8
9 Kasperson J, Kasperson R, Turner B 1995. *Regions at Risk*. United Nations University Press.
10 <http://www.unu.edu/unupress/unupbooks/uu14re/uu14re00.htm>
- 11
12 Kates RW, Clark WC, Corell R, Hall JM, Jaeger CC, Lowe I, McCarthy JJ, Schnellhuber HJ, Bolin B,
13 Huntley B, Jäger J, Jodha NS, Kasperson RE, Mabogunje A, Matson P, Mooney H, Moore B,
14 O’Riordan T, Svedin U 2001. Sustainability Science. *Science* **292**: 641-642.
- 15
16 Kirkby MJ, Irvine BJ, Jones RJA, Govers G, the PESERA team 2008. The PESERA coarse scale erosion
17 model for Europe: I – Model rationale and implementation. *European Journal of Soil Science* **59**: 1293-
18 1306.
- 19
20
21 Lele SR, Norgaard RB 2005. Practicing interdisciplinarity. *BioScience* **55**: 967–975.
- 22
23 Magole L, Atlhopheng JR, Chanda R 2008. *Stakeholder workshop 1: Land degradation and desertification*
24 *– existing and potential prevention and conservation strategies*. Boteti First DESIRE WOCAT
25 workshop report.
- 26
27 Magole, L, Atlhopheng, JR and Chanda, R 2009. *Stakeholder workshop 2: Selection and decision on*
28 *technologies / approaches to be implemented*. Boteti First DESIRE WOCAT workshop report.
- 29
30 Ministry of Agriculture 1994. Desertification and possible solutions in the mid-Boteti River area: a
31 Botswana case study for the intergovernmental convention to combat desertification (INCD). Final
32 report: Volume 1 – the desertification process and its impacts. A Consultancy report prepared by a team
33 from the Department of Environmental Science, University of Botswana, April 1994 (164 pages).
- 34
35 Mulale, K, Chanda, R 2009. *Drivers of desertification in the Boteti area of Botswana: law and policy*.
36 DESIRE research report.
- 37
38 Perkins JS 2007. *Ecological impact evaluation of current rangeland management regimes in Botswana,*
39 *with special reference to the eastern Kalahari*. Component 1 Report for the UB-IVP-CAR collaborative
40 research project.
- 41
42
43 Reed MS 2005. *Participatory Rangeland Monitoring and Management in the Kalahari, Botswana*. PhD
44 Thesis, School of Earth & Environment, University of Leeds.
- 45
46 Reed MS, Dougill AJ, Taylor MJ 2007. Integrating local and scientific knowledge for adaptation to land
47 degradation: Kalahari rangeland management options. *Land Degradation & Development* **18**: 249-268.
- 48
49 Reed MS, Dougill AJ, Baker T 2008. Participatory indicator development: what can ecologists and local
50 communities learn from each other? *Ecological Applications* **18**: 1253–1269.
- 51
52 Reed MS, Dougill AJ 2010. Linking Degradation Assessment to Sustainable Land Management: a decision
53 support system for Kalahari pastoralists. *Journal of Arid Environments* **74**: 149–155.
- 54
55 Reed MS, Buenemann, M, Atlhopheng J, Akhtar-Schuster M, Bachmann F, Bastin G, Bigas H, Chanda R,
56 Dougill AJ, Essahli W, Evely AC, Fleskens L, Geeson N, Glass JH, Hessel R, Holden J, Ioris A, Kruger
57
58
59
60

- 1
2
3 B, Liniger HP, Mphinyane W, Nainggolan D, Perkins J, Raymond CM, Ritsema CJ, Schwilch G,
4 Sebege R, Seely M, Stringer LC, Thomas R, Twomlow S, Verzandvoort S 2011. Cross-scale
5 monitoring and assessment of land degradation and sustainable land management: a methodological
6 framework for knowledge management. *Land Degradation & Development*.
7
8
9 Reed MS, Fazey I, Stringer LC, Raymond CM, Akhtar-Schuster M, Begni G, Bigas H, Brehm S, Briggs J,
10 Bryce R, Buckmaster S, Chanda R, Davies J, Diez E, Essahli W, Evely A, Geeson N, Hartmann I,
11 Holden J, Hubacek K, Ioris I, Kruger B, Laureano P, Phillipson J, Prell C, Quinn CH, Reeves AD, Seely
12 M, Thomas R, van der Werff Ten Bosch MJ, Vergunst P, Wagner L (in press) Knowledge management
13 for land degradation monitoring and assessment: an analysis of contemporary thinking. *Land*
14 *Degradation & Development*.
15
16
17
18 Republic of Botswana (2004) *Summary statistics on small areas - Volume VII. 2001 Population and*
19 *Housing Census*. Central Statistics Office, Government Printer, Gaborone, Botswana.
20
21 Ringrose S, Chanda R, Nkambwe M, Sepe F 1996. Environmental change in the mid-boteti area of North-
22 central Botswana: biophysical processes and human perceptions. *Environmental Management* **20**: 397-
23 410
24
25
26
27 Schwilch G, Liniger HP, Bachmann F 2007b. *Guidelines for WB3 Part II: Assessment of Conservation*
28 *Strategies. Assessment and documentation of existing and potential prevention and mitigation*
29 *strategies*. Centre for Development and Environment (CDE), University of Bern, Switzerland. Report
30 Number 7 of Manuals and Training Guides. In collaboration with DESIRE Project.
31
32 Schwilch G, Bachmann F, Liniger H 2008. *Guidelines for WB 3 Part III: stakeholder workshop 2. Selection*
33 *and decision on prevention and mitigation strategies to be implemented*. Centre for Development and
34 Environment (CDE), University of Bern, Switzerland. Report Number 17 of Manuals and Training
35 Guides. Schwilch, G., Bachmann, F., Liniger H.P. 2009. Appraising and selecting conservation
36 measures to mitigate desertification and land degradation based on stakeholder participation and global
37 best practices. *Land Degradation & Development* **20**: 308-326.
38
39
40 Schwilch G., Bestelmeyer B, Bunning S., Critchley W., Herrick J., Kellner K., Liniger H.P., Nachtergaele
41 F., Ritsema C.J., Schuster B., Tabo R., van Lynden G., Winslow M. 2011. Experiences in Monitoring
42 and Assessment of Sustainable Land Management. *Land Degradation & Development* (in press).
43
44 Doi10.1002/ldr.1040
45
46 Tietema T 1993. Biomass determination of fuelwood trees and bushes of Botswana, Southern Africa,
47 *Forest Ecology and Management* **60**: 257-269.
48
49 Turpie JK, Crowe TM 1994. Patterns of distribution, diversity and endemism of larger African mammals
50 *South African Journal of Zoology* **29**: 19-32 .
51
52 Turner BL II, Bejamin P, 1993. Fragile lands and their management. In *Agriculture, environment and*
53 *health: Towards sustainable development into 21st century*, Ruttan VW (ed). University of Minnesota
54 Press, Minneapolis.
55
56 UNCBD 1992. *United Nations convention on biological diversity*. UNCBD Secretariat: Montreal.
57
58
59
60

1
2
3 UNCCD (1994) *United Nations Convention to Combat Desertification in Those Countries Experiencing*
4 *Serious Drought and/or Desertification Particularly in Africa: Text with Annexes*, UNEP, Nairobi.

5
6 Warren A. 2002. Land degradation is contextual. *Land Degradation and Development* **13**: 449–459.

7
8 Werger MJA, Coetzee BJ 1978. The Sudano-Zambezian Region. In *Biogeography and ecology of southern*
9 *Africa*, Werger MJA (ed). Junk, The Hague; 301–46.

10
11 White F 1983. *The vegetation of Africa, a descriptive memoir to accompany the UNESCO/AETFAT/UNSO*
12 *Vegetation Map of Africa* (3 Plates, Northwestern Africa, Northeastern Africa, and Southern Africa,
13 1:5,000,000), UNESCO, Paris.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 1: Boteti land users' objectives for sustainable land management (source: Workshop 1)

Objective	Appropriate technology	Adequate approaches	Responsible stakeholders
To address the issue of shortage of water	Building dams	- Submit application for land. - Submit application for financial assistance.	- Land Board - Farmers - Agricultural extension workers - Financial assistance agencies
To reduce the impact of drought on harvests	Irrigation farming	- Submit application for land. -Produce a management plan. - Submit application for financial assistance -Produce a training plan for farmers.	- Village leaders (Village Development Committee). - Land Board - Agricultural extension workers and soil specialists - Arable farmers
To reduce depletion of trees	Use bio-gas as energy for cooking	-Submit application for land. - Submit application for financial assistance -Produce a training plan for implementers and or users.	- Village leaders (Village Development Committee). - Land Board - Local community - Rural Industry Innovation Centre (RIIC)
To reduce poverty	Divert to wildlife ranching	- Submit application for land. -Produce a management plan. - Submit application for financial assistance -Produce a training plan on wildlife ranching.	-Department of Wildlife and national parks (DWNP) and wildlife ranching experts - Village leaders (Village Development Committee). - Land Board - Local community

Source: Magole et al. (2008), p.10.

Table 2: Land user identified indicators

Negative or degradation indicators – Degradation	The positive attributes or indicators – Conservation
Lack of vegetation, no regeneration and germination	The river starts flowing again
Uncontrollable winds	Good rains
High temperatures	Good harvests (high crop yields)
Wild fires	Acceptable levels of soil and water salinity
Depletion of underground water	Soil fertility increases
No dew (due to dry atmospheric conditions)	Improved (raised) underground water table
High mortality rate of livestock	Dams, ponds, pans and lakes fill up with water.
Low weight and weak livestock	More food for both animals and people
Trees and other vegetation (e.g. weeds) die	Good vegetation cover
Browning (as opposed to greening) of the land	Animals recover and start to reproduce
Pans, dams and lakes dry up	Increased and improved quality of wild fruits
Extinction of certain plants	Wild animals re-appear
Poor soil conditions, crusting	Food becomes abundant
Some grasses disappear	Greening of the land
Low yields	Birds and other wildlife begin to sing
Shortage of water for both animals and people	
Some of the soils become disturbed (loose, poor)	
High water salinity	
High soil salinity	
Wildlife disappears e.g. animals, butterflies, etc	
Drying of the river (Boteti R.)	
Too much dust	
Poverty and hunger	
Cattle tracks expose soil leading to erosion and sand mounds	
Increasing and spreading of certain vegetation species	

Source: Magole et al. (2008), p.7

Table 3. Financial performance of biogas expressed in firewood collection opportunity costs under various scenarios.

	Daily household dung collection costs		Daily household firewood collection opportunity costs	
	Pula	% ^a	Pula	% ^a
Cattle distribution				
Scenario A: Cattle uniformly distributed ^a	6.84	100	13.82	100
Scenario B: Cattle herds from 11 km	8.51	124	15.46	112
Scenario C: Cattle herds beyond 20km	12.37	181	19.35	140
Efficiency and socio-economic dynamics				
Scenario D: Biogas covers 75% of demand	11.09	162	18.06	131
Scenario E: Firewood price rise trend 10%	6.84	100	6.45-18.47	47-134
Scenario F: No cost dung collection <1km	4.59	67	11.58	60
Scenario G: On foot dung collection only	95.93	1402	102.92	745

^a Scenario A is used as baseline; percentages are relative to this scenario.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review




Figure 1: Integrated methodological framework for land degradation and SLM monitoring and assessment, building on the DESIRE, WOCAT, LADA and DDP approaches, providing examples in italics around the outside of the figure that show how each step may be operationalised (drawing mainly on experience from the DESIRE project). Dashed arrows represent potential links that may not always be realised (adapted from Reed *et al.*, 2011).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2: The Botswana DESIRE study site (Source: Sebego, RJ, Botswana DESIRE team)

Review




Figure 3: Modelling alternative uses of biomass resource

Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4: Schematic diagram showing how the Pan-European Erosion Assessment (PESERA) model partitions rainfall into components for overland flow, evapotranspiration and changes in soil moisture storage .




Figure 5: Land degradation drivers and desertification risk in Boteti, showing the number of land use types within the study site where different drivers of land degradation were deemed to be occurring, according to an expert assessment of indicators by researchers and land managers (Source: Chanda et al., 2009a)




Figure 6: Perceived ecological impact of solar cooker, rainwater harvesting, game ranching and biogas, where 1.0 is a beneficial ecological impact an 0.0 is a negative ecological impact, and width of bar represents the breadth of responses received and hence the level of agreement or disagreement between stakeholders (narrow bar represents high levels of agreement and wide bar represents a wide range of different answers) (Source: Workshop 2 report; Magole et al., 2009, p. 11)

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 7: Perceived economic impact of solar cooker, rainwater harvesting, game ranching and biogas, where 1.0 is a beneficial economic impact an 0.0 is a negative economic impact, and width of bar represents the breadth of responses received and hence the level of agreement or disagreement between stakeholders (narrow bar represents high levels of agreement and wide bar represents a wide range of different answers) (Source: Workshop 2 report; Magole et al., 2009, p. 11)




Figure 8: Perceived socio-cultural impact of solar cooker, rainwater harvesting, game ranching and biogas, where 1.0 is a beneficial socio-cultural impact and 0.0 is a negative socio-cultural impact, and width of bar represents the breadth of responses received and hence the level of agreement or disagreement between stakeholders (narrow bar represents high levels of agreement and wide bar represents a wide range of different answers) (Source: Workshop 2 report; Magole et al., 2009, p. 12)

For Peer Review




Figure 9: Location of the 378 boreholes (potential dung collection sites) in the study site relative to the villages Mopipi (centre) and Mokoboxane (south-east of Mopipi)

Online supplementary material

Video of local residents from the villages of Rakops, Mopipi and Mokoboxane talking about the environmental problems they are currently facing and their hopes for how the DESIRE project will help them:

http://www.desire-his.eu/index.php?option=com_content&view=article&id=388:interviews-with-local-people&catid=228:boteti-botswana&Itemid=155&lang=en

For Peer Review