



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/81908/>

Version: Published Version

Article:

Tsigaridis, K, Daskalakis, N, Kanakidou, M et al. (2014) The AeroCom evaluation and intercomparison of organic aerosol in global models. *Atmospheric Chemistry and Physics*, 14 (19). pp. 10845-10895. ISSN: 1680-7316

<https://doi.org/10.5194/acp-14-10845-2014>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



The AeroCom evaluation and intercomparison of organic aerosol in global models

K. Tsigaridis^{1,2}, N. Daskalakis^{3,4}, M. Kanakidou³, P. J. Adams^{5,6}, P. Artaxo⁷, R. Bahadur⁸, Y. Balkanski⁹, S. E. Bauer^{1,2}, N. Bellouin^{10,a}, A. Benedetti¹¹, T. Bergman¹², T. K. Berntsen^{13,14}, J. P. Beukes¹⁵, H. Bian¹⁶, K. S. Carslaw¹⁷, M. Chin¹⁸, G. Curci¹⁹, T. Diehl^{18,20}, R. C. Easter²¹, S. J. Ghan²¹, S. L. Gong²², A. Hodzic²³, C. R. Hoyle^{24,25}, T. Iversen^{11,26,13}, S. Jathar⁵, J. L. Jimenez²⁷, J. W. Kaiser^{28,11,29}, A. Kirkevåg²⁶, D. Koch^{1,2,b}, H. Kokkola¹², Y. H. Lee^{5,c}, G. Lin³⁰, X. Liu^{21,d}, G. Luo³¹, X. Ma^{32,e}, G. W. Mann^{33,34}, N. Mihalopoulos³, J.-J. Morcrette¹¹, J.-F. Müller³⁵, G. Myhre¹⁴, S. Myriokefalitakis^{3,4}, N. L. Ng³⁶, D. O'Donnell^{37,f}, J. E. Penner³⁰, L. Pozzoli³⁸, K. J. Pringle^{39,29}, L. M. Russell⁹, M. Schulz²⁶, J. Sciare⁹, Ø. Seland²⁶, D. T. Shindell^{2,1,g}, S. Sillman³⁰, R. B. Skeie¹⁴, D. Spracklen¹⁷, T. Stavrakou³⁵, S. D. Steenrod²⁰, T. Takemura⁴⁰, P. Tiitta^{15,41}, S. Tilmes²³, H. Tost⁴², T. van Noije⁴³, P. G. van Zyl¹⁵, K. von Salzen³², F. Yu³¹, Z. Wang⁴⁴, Z. Wang⁴⁵, R. A. Zaveri²¹, H. Zhang⁴⁴, K. Zhang^{21,37}, Q. Zhang⁴⁶, and X. Zhang⁴⁵

¹Center for Climate Systems Research, Columbia University, New York, NY, USA

²NASA Goddard Institute for Space Studies, New York, NY, USA

³Environmental Chemical Processes Laboratory, Department of Chemistry, University of Crete, Heraklion, Greece

⁴Institute of Chemical Engineering, Foundation for Research and Technology Hellas (ICE-HT FORTH), Patras, Greece

⁵Department of Civil and Environmental Engineering, Carnegie Mellon University, Pittsburgh, PA, USA

⁶Department of Engineering and Public Policy, Carnegie Mellon University, Pittsburgh, PA, USA

⁷University of São Paulo, Department of Applied Physics, Brazil

⁸Scripps Institution of Oceanography, University of California San Diego, CA, USA

⁹Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France

¹⁰Met Office Hadley Centre, Exeter, UK

¹¹ECMWF, Reading, UK

¹²Finnish Meteorological Institute, Kuopio, Finland

¹³University of Oslo, Department of Geosciences, Oslo, Norway

¹⁴Center for International Climate and Environmental Research – Oslo (CICERO), Oslo, Norway

¹⁵Environmental Sciences and Management, North-West University, Potchefstroom, South Africa

¹⁶University of Maryland, Joint Center for Environmental Technology, Baltimore County, MD, USA

¹⁷School of Earth and Environment, University of Leeds, Leeds, UK

¹⁸NASA Goddard Space Flight Center, Greenbelt, MD, USA

¹⁹Department of Physics CETEMPS, University of L'Aquila, Italy

²⁰Universities Space Research Association, Greenbelt, MD, USA

²¹Pacific Northwest National Laboratory; Richland, WA, USA

²²Air Quality Research Branch, Meteorological Service of Canada, Toronto, Ontario, Canada

²³National Center for Atmospheric Research, Boulder, CO, USA

²⁴Paul Scherrer Institute, Villigen, Switzerland

²⁵Swiss Federal Institute for Forest Snow and Landscape Research (WSL) – Institute for Snow and Avalanche Research (SLF), Davos, Switzerland

²⁶Norwegian Meteorological Institute, Oslo, Norway

²⁷University of Colorado, Department of Chemistry & Biochemistry, Boulder, CO, USA

²⁸King's College London, Department of Geography, London, UK

²⁹Department of Atmospheric Chemistry, Max Planck Institute for Chemistry, Mainz, Germany

³⁰Department of Atmospheric, Oceanic, and Space Sciences, University of Michigan, Ann Arbor, MI, USA

³¹State University of New York, Albany, NY, USA

³²Environment Canada, Victoria, Canada

³³National Centre for Atmospheric Science, University of Leeds, Leeds, UK

³⁴School of Earth and Environment, University of Leeds, Leeds, UK

³⁵Belgian Institute for Space Aeronomy, Brussels, Belgium

³⁶School of Chemical and Biomolecular Engineering and School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

³⁷Max Planck Institute for Meteorology, Hamburg, Germany

³⁸Eurasia Institute of Earth Sciences, Istanbul Technical University, Turkey

³⁹Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, UK

⁴⁰Research Institute for Applied Mechanics, Kyushu University, Fukuoka, Japan

⁴¹Fine Particle and Aerosol Technology Laboratory, Department of Environmental Science, University of Eastern Finland, Kuopio, Finland

⁴²Institute for Atmospheric Physics, Johannes Gutenberg University, Mainz, Germany

⁴³Royal Netherlands Meteorological Institute (KNMI), De Bilt, the Netherlands

⁴⁴Laboratory for Climate Studies, Climate Center, China Meteorological Administration, Beijing, China

⁴⁵Chinese Academy of Meteorological Sciences, Beijing, China

⁴⁶Department of Environmental Toxicology, University of California, Davis, CA, USA

^anow at: Department of Meteorology, University of Reading, Reading, UK

^bnow at: Department of Energy, Office of Biological and Environmental Research, Washington, DC, USA

^cnow at: Center for Climate Systems Research, Columbia University, New York, NY, USA and NASA Goddard Institute for Space Studies, New York, NY, USA

^dnow at: University of Wyoming, Department of Atmospheric Science, Laramie, WY, USA

^enow at: State University of New York, Department of Atmospheric Science, Albany, NY, USA

^fnow at: Finnish Meteorological Institute, Helsinki, Finland

^gnow at: Nicholas School of the Environment, Duke University, Durham, NC, USA

Correspondence to: K. Tsigaridis (kostas.tsigaridis@columbia.edu) and M. Kanakidou (mariak@chemistry.uoc.gr)

Received: 11 February 2014 – Published in Atmos. Chem. Phys. Discuss.: 7 March 2014

Revised: 3 September 2014 – Accepted: 3 September 2014 – Published: 15 October 2014

Abstract. This paper evaluates the current status of global modeling of the organic aerosol (OA) in the troposphere and analyzes the differences between models as well as between models and observations. Thirty-one global chemistry transport models (CTMs) and general circulation models (GCMs) have participated in this intercomparison, in the framework of AeroCom phase II. The simulation of OA varies greatly between models in terms of the magnitude of primary emissions, secondary OA (SOA) formation, the number of OA species used (2 to 62), the complexity of OA parameterizations (gas-particle partitioning, chemical aging, multi-phase chemistry, aerosol microphysics), and the OA physical, chemical and optical properties. The diversity of the global OA simulation results has increased since earlier AeroCom experiments, mainly due to the increasing complexity of the SOA parameterization in models, and the implementation of new, highly uncertain, OA sources. Diversity of over one order of magnitude exists in the modeled vertical distribution of OA concentrations that deserves a dedicated future study. Furthermore, although the OA/OC ratio depends on OA sources and atmospheric processing, and is important

for model evaluation against OA and OC observations, it is resolved only by a few global models.

The median global primary OA (POA) source strength is 56 Tg a^{-1} (range $34\text{--}144 \text{ Tg a}^{-1}$) and the median SOA source strength (natural and anthropogenic) is 19 Tg a^{-1} (range $13\text{--}121 \text{ Tg a}^{-1}$). Among the models that take into account the semi-volatile SOA nature, the median source is calculated to be 51 Tg a^{-1} (range $16\text{--}121 \text{ Tg a}^{-1}$), much larger than the median value of the models that calculate SOA in a more simplistic way (19 Tg a^{-1} ; range $13\text{--}20 \text{ Tg a}^{-1}$, with one model at 37 Tg a^{-1}). The median atmospheric burden of OA is 1.4 Tg (24 models in the range of $0.6\text{--}2.0 \text{ Tg}$ and 4 between 2.0 and 3.8 Tg), with a median OA lifetime of 5.4 days (range $3.8\text{--}9.6$ days). In models that reported both OA and sulfate burdens, the median value of the OA/sulfate burden ratio is calculated to be 0.77; 13 models calculate a ratio lower than 1, and 9 models higher than 1. For 26 models that reported OA deposition fluxes, the median wet removal is 70 Tg a^{-1} (range $28\text{--}209 \text{ Tg a}^{-1}$), which is on average 85 % of the total OA deposition.

Fine aerosol organic carbon (OC) and OA observations from continuous monitoring networks and individual field

campaigns have been used for model evaluation. At urban locations, the model–observation comparison indicates missing knowledge on anthropogenic OA sources, both strength and seasonality. The combined model–measurements analysis suggests the existence of increased OA levels during summer due to biogenic SOA formation over large areas of the USA that can be of the same order of magnitude as the POA, even at urban locations, and contribute to the measured urban seasonal pattern.

Global models are able to simulate the high secondary character of OA observed in the atmosphere as a result of SOA formation and POA aging, although the amount of OA present in the atmosphere remains largely underestimated, with a mean normalized bias (MNB) equal to -0.62 (-0.51) based on the comparison against OC (OA) urban data of all models at the surface, -0.15 ($+0.51$) when compared with remote measurements, and -0.30 for marine locations with OC data. The mean temporal correlations across all stations are low when compared with OC (OA) measurements: 0.47 (0.52) for urban stations, 0.39 (0.37) for remote stations, and 0.25 for marine stations with OC data. The combination of high (negative) MNB and higher correlation at urban stations when compared with the low MNB and lower correlation at remote sites suggests that knowledge about the processes that govern aerosol processing, transport and removal, on top of their sources, is important at the remote stations. There is no clear change in model skill with increasing model complexity with regard to OC or OA mass concentration. However, the complexity is needed in models in order to distinguish between anthropogenic and natural OA as needed for climate mitigation, and to calculate the impact of OA on climate accurately.

1 Introduction

Atmospheric aerosols are important drivers of air quality and climate. The organic component of aerosols can contribute 30–70 % of the total submicron dry aerosol mass, depending on location and atmospheric conditions (Kanakidou et al., 2005; Murphy et al., 2006). The majority of fine aerosol mass (PM_{10} : particulate matter of dry diameter smaller than $1 \mu\text{m}$) consists of non-refractory material, and has been found to contain large amounts of organic matter (Zhang et al., 2007; Jimenez et al., 2009), as measured by the Aerosol Mass Spectrometer (AMS).

Global model estimates of the dry organic aerosol (OA) direct radiative forcing at the top of the atmosphere are $-0.14 \pm 0.05 \text{ W m}^{-2}$ based on AeroCom phase I experiments (Schulz et al., 2006), which was decomposed during AeroCom phase II to $-0.03 \pm 0.01 \text{ W m}^{-2}$ for primary organic aerosol (POA) from fossil fuel and biofuel, $-0.02 \pm 0.09 \text{ W m}^{-2}$ for secondary organic aerosol (SOA) and $0.00 \pm 0.05 \text{ W m}^{-2}$ for the combined OA and black

carbon from biomass burning (Myhre et al., 2013). IPCC (2013) assessed the contribution of anthropogenic primary and secondary organic aerosols to the radiative forcing from aerosol–radiation interactions (RFari) to be -0.12 (-0.4 to $+0.1$) W m^{-2} . Spracklen et al. (2011) estimated the climate forcing of the anthropogenically driven natural SOA alone (including the presence of water on hydrophilic OA) at $-0.26 \pm 0.15 \text{ W m}^{-2}$ (direct effect) and $-0.6_{-0.14}^{+0.24} \text{ W m}^{-2}$ (indirect effect). These amounts largely depend on the atmospheric loadings of OA simulated by the models under past, present and future climate conditions, and on the properties they attribute to them. Indeed, Myhre et al. (2013) calculated a SOA load of $0.33 \pm 0.32 \text{ Tg}$, while Spracklen et al. (2011) estimated a SOA load of 1.84 Tg , which resulted in an order of magnitude higher radiative forcing. There is therefore an urgent need for a consensus between models and agreement with observations, in order to constrain the large variability between models and, consequently, the OA impact on climate.

1.1 Definitions

OA can be emitted directly as POA or formed via gas-phase reactions and subsequent condensation of semi-volatile vapors, resulting in SOA. In addition, multiphase and heterogeneous processes can also contribute to SOA formation. Emissions of volatile organic compounds (VOCs) from terrestrial vegetation are 10 times larger than from anthropogenic sources (Guenther et al., 1995; Kanakidou et al., 2005, and references therein). In addition, the mass of organic carbon emitted in the gas phase exceeds by more than a factor of 10 that emitted directly as primary particulate matter (Goldstein and Galbally, 2007; Kanakidou et al., 2012). VOCs therefore have a large potential to contribute to SOA formation. However, the exact formation processes and composition of OA are poorly understood. Fuzzi et al. (2006) and Hallquist et al. (2009) provided a number of marker compounds and observations that could be used to distinguish the various OA sources. Most OA observational techniques measure the particulate organic carbon content of OA mass, either total (OC) or the water soluble component (WSOC), while some of the variability of OA is accounted for by oxygen, nitrogen and other elements in the organic compounds. Significant discrepancies in OC concentrations determined by different techniques have been identified (Kanakidou et al., 2005), and have been addressed by protocols of the definition of OC/EC (elemental carbon) measurements (Cavalli et al., 2010). The use of OC historically corresponded to its easier measurement. Recently, Aerosol Mass Spectrometer (AMS) observations started providing very high temporal resolution information on the OA mass of the non-refractory PM_{10} . It has to be emphasized that it is the OA mass, not the OC, which determines aerosol properties such as chemical composition, size, hygroscopicity and hygroscopic growth, each of which is an important factor affecting aerosol scattering,

absorption and the ability to act as cloud condensation nuclei (CCN). Therefore, the ratio of OA to OC mass (Turpin and Lim, 2001; Aiken et al., 2008) requires careful investigation. Furthermore, OA compounds differ in their volatility, solubility, hygroscopicity, chemical reactivity and their physical and optical properties. Due to the chemical complexity of the organic component of aerosols (Goldstein and Galbally, 2007), only simplified representations are introduced in global chemistry climate models (Kanakidou et al., 2005; Hallquist et al., 2009). As a compromise between simplicity and accuracy, the net effect of the complex mixture of OA is described by only a limited number of representative compounds or surrogates.

1.2 Sources

Kanakidou et al. (2005) reviewed how organic aerosols were incorporated into global chemistry transport models (CTMs) and general circulation models (GCMs), and identified gaps in knowledge that deserved further investigation. The POA sources include fossil fuel, biofuel and biomass burning, as well as the less understood sources of marine OA, biological particles and soil organic matter on dust (Kanakidou et al., 2012, and references therein). Biogenic VOCs (BVOCs) greatly contribute to OA formation (e.g., Griffin et al., 1999b; Kanakidou et al., 2012), implying that significant feedbacks exist between the biosphere, the atmosphere and climate that affect the OA levels in the atmosphere, which was also demonstrated by more recent studies (Tsigaridis et al., 2005; Arneth et al., 2010; Carslaw et al., 2010; Paasonen et al., 2013). In addition, oxidant and pollutant enhancement by human-induced emissions is expected to increase OA levels, even those chemically formed by BVOC (Hoyle et al., 2011, and references therein); it is therefore conceivable that some portion of the ambient biogenic SOA, which would have been absent under preindustrial conditions, can be removed by controlling emissions of anthropogenic pollutants (Carlton et al., 2010). Goldstein and Galbally (2007) estimated that SOA formation could be as high as 910 Tg C a^{-1} , which is at least an order of magnitude higher than any SOA formation modeling study, as shown here. Spracklen et al. (2011) were able to reconcile AMS observations (mostly from the Northern Hemisphere mid-latitudes during summer) with global CTM simulations by estimating a large SOA source (140 Tg a^{-1}). 100 Tg a^{-1} was characterized as anthropogenically controlled, 90% of which was possibly linked to anthropogenically enhanced SOA formation from BVOC oxidation. Similar conclusions were reached by Heald et al. (2011) by comparing aircraft AMS observations of submicron OA with the results of another global model, and by Heald et al. (2010) by accounting for the satellite-measured aerosol optical depth that could possibly be due to OA. Recently, Carlton and Turpin (2013) showed that anthropogenically enhanced aerosol water in the eastern USA could lead to an increase in WSOC from

BVOC. Although large uncertainties still exist in SOA modeling, there is a need for models to document and improve treatments of solubility, hygroscopicity, volatility and optical properties of the OA from different sources. The SOA formation from anthropogenic VOCs, despite a recent estimate of 13.5 Tg a^{-1} that makes it a non-negligible SOA source in polluted regions (De Gouw and Jimenez, 2009), is frequently neglected by global models.

1.3 Atmospheric processing

Improvement in our understanding and quantification of the emissions of POA and SOA precursors demonstrated from earlier review studies (Kanakidou et al., 2005; Fuzzi et al., 2006) motivated a number of experimental, chamber and field studies that have also significantly enhanced our knowledge on the OA atmospheric cycle. Aging, both physical (e.g., condensation and coagulation) and chemical (in any phase), has been suggested as a significant contributor to the observed OA levels (Kanakidou et al., 2005; Fuzzi et al., 2006; Hallquist et al., 2009), which influences the amount and properties of organic material in the aerosol phase, and occurs at different rates and via different mechanisms in the various atmospheric compartments (e.g., urban/rural/marine boundary layers, low/middle/upper troposphere) (e.g., Molina et al., 2004; Ervens et al., 2011). Despite these advances in understanding, such OA processing remains to date either missing or very poorly parameterized in global models, since advances in OA parameterizations are limited by weak observational constraints. Zhang et al. (2007) and Jimenez et al. (2009) compiled experimental evidence showing that most of the OA in the atmosphere has undergone chemical aging, most likely via SOA formation, and is significantly oxygenated, with lower volatility and higher hygroscopicity than its precursors. To explain these large amounts of oxygenated OA, several chemical pathways have been suggested (Hallquist et al., 2009; Jimenez et al., 2009), which differ in the O/C atomic ratio and in the volatility changes they induce in the parent compounds. Donahue et al. (2006) suggested lumping organic compounds according to their volatility and developed the volatility-basis set (VBS) algorithm to parameterize the many organic compounds present in the atmosphere into several lumped OA species of different volatilities. Chemical aging via gas-phase reactions in the parameterization resulted in changes in the volatility of the species; this has been implemented for SOA from VOCs (e.g., Tsimpidi et al., 2010) and also for SOA from semi-volatile and intermediate volatility species (Robinson et al., 2007). However, the implementation of VBS into global models is hindered both by the large number of tracers required, and the underlying uncertainties and free parameters involved. The VBS method was recently expanded to account for the degree of oxidation of OA, by tracking the O/C content of the organics per volatility class; the method is called 2-D VBS (Donahue et al., 2011) and

has been successfully used to simulate the evolution of OA in field campaigns (Murphy et al., 2011, 2012). Unfortunately, this new approach needs an even larger number of tracers, which makes it extremely difficult to implement in a global climate model without a large performance penalty. Still, it certainly adds value to our OA understanding, since the ratio of organic aerosol mass (OA) to organic carbon (OA/OC), an alternative way to describe the degree of oxidation of OA, does greatly vary in time and space (Turpin and Lim, 2001). This variability is either neglected or taken into account in a very simplistic way in models.

Yu (2011) extended the two-product SOA formation scheme in the GEOS-Chem model by taking into account the volatility changes of secondary organic gases arising from the oxidative aging process (Jimenez et al., 2009) as well as the kinetic condensation of low-volatility secondary organic gases. It was shown that, over many parts of the continents, low-volatility secondary organic gas concentrations are generally a factor of ~ 2 – 20 higher than those of sulfuric acid gas, and the kinetic condensation of low-volatility secondary organic gases significantly enhances particle growth rates. Based on this computationally efficient new SOA formation scheme, annual mean SOA mass concentrations in many parts of the boundary layer increase by a factor of 2–10, in better agreement with Aerosol Mass Spectrometer (AMS) SOA measurements (Yu, 2011).

Hallquist et al. (2009) also summarized new laboratory data that provided insight into the chemical reaction pathways for the formation of oligomers and other higher molecular weight products observed in SOA. They determined higher production rates of SOA from their precursors' oxidation than earlier measurement studies and linked the dependence of SOA yield from VOC oxidation to the oxidant levels. In chamber experiments, Volkamer et al. (2009) have shown that even small (C_2) molecules undergoing aqueous-phase reactions can produce low-volatility material and contribute to SOA formation in the atmosphere, a process that was reviewed by Ervens et al. (2011) and Lim et al. (2013). The global modeling study of Myriokefalitakis et al. (2011) has shown that multiphase reactions of organics significantly increase the OA mass (5–9% when expressed as OC) and its oxygen content, while Murphy et al. (2012) suggested that these reactions are not enough to explain the observed O/C content of OA.

1.4 Losses

Hallquist et al. (2009) used the VBS concept and estimated the atmospheric deposition of OA to be 150 Tg a^{-1} , higher than earlier estimates and similar to the total particulate OC deposition of 147 Tg a^{-1} (109 Tg a^{-1} of WSOC) calculated by Kanakidou et al. (2012). Dry and wet removal of organic vapors that are in thermodynamic equilibrium with SOA becomes increasingly important with atmospheric processing (Hodzic et al., 2013) and was found to lead to 10–30% (up

to 50%) removal of anthropogenic (biogenic) SOA (Hodzic et al., 2014). Volatilization of OA upon heterogeneous oxidation has been observed for laboratory and ambient particles (George and Abbatt, 2010) and might be a significant OA sink (Heald et al., 2011).

1.5 Motivation and aim

During the AeroCom phase I modeling experiments (Textor et al., 2006), although most of the models considered both primary and secondary OA sources, OA was simulated in a very simplified way in which both primary and secondary OA were treated as non-volatile. OA was only allowed to age via hydrophobic-to-hydrophilic conversion, and was removed from the atmosphere by particle deposition. Comparisons of individual models with OA observations have shown a large underestimation of the organic aerosol component by models, especially in polluted areas (Volkamer et al., 2006, and references therein). They showed that the underestimation of SOA by models increases with photochemical age, which can be partially correlated with long-range transport, with the largest discrepancies in the free troposphere, suggesting missing sources or underestimated atmospheric processing of organics in models.

Several global models now treat SOA as semi-volatile, as detailed below, which enables potentially more accurate model calculations. Some models also account for intermediate-volatility organics, multiphase chemistry and semi-volatile POA (e.g., Pye and Seinfeld, 2010; Jathar et al., 2011; Myriokefalitakis et al., 2011; Lin et al., 2012), with encouraging results in reducing the difference between models and observations. Indeed, the modeled SOA concentrations in Mexico City were much closer to observations when intermediate-volatility organics were taken into account in a regional model, although it was unclear if the model–observation gap was reduced for the right reasons (Hodzic et al., 2010). However, OA simulations have many degrees of freedom due to incomplete knowledge of the behavior and fate of OA in the troposphere. Thus, several assumptions made are translated to model tuning parameters that vary greatly between models.

This organic aerosol AeroCom intercomparison aims to update the evaluation of OA modeling by documenting the current status of global modeling of OA in the troposphere, identifying weaknesses that still exist in models, as well as explaining the similarities and differences that exist between models and observations. It quantifies the uncertainties in surface OA concentrations and attributes them to major contributors. It also attempts to identify and analyze potential model systematic biases. The ensemble of the simulations is used to build an integrated and robust view of our understanding of organic aerosol sources and sinks in the troposphere. The target year of simulations was selected to be 2006, with a free choice for each modeling group on the meteorological conditions and emission inventories to be used.

1.6 Terminology

In atmospheric OA research, several naming conventions and abbreviations are used, often ambiguously and inconsistently between authors. To avoid confusion, we clarify here the conventions adopted in this paper, which we use throughout. Note that some aspects of our terminology are different from the very recent VBS-centered attempt by Murphy et al. (2014) to clarify this ambiguity systematically; new model development is required from modelers to adopt the new naming convention in future model simulations.

- Organic aerosol (OA) and the main OA components, i.e., primary and secondary OA (POA and SOA, respectively): we use these terms to refer to the total mass that organic compounds have in the aerosol phase, including H and O, and potentially other elements like N, S and P. Other authors have used the term organic matter (OM), which is synonymous with our OA definition. The units used are $\mu\text{g m}^{-3}$ for surface mass concentrations at ambient conditions and Tg for burden and budget calculations. OA amounts exclude the water associated with it (assuming that OA is hygroscopic), an important additional component that affects particle size, refractive index and light scattering efficiency.
- Organic carbon (OC), together with other OC components, like, e.g., primary and secondary OC (POC and SOC, respectively): these terms refer to the mass of carbon present in OA, instead of to the total OA mass. The units used here are $\mu\text{gC m}^{-3}$ for surface mass concentrations. This is typically the terminology that is used when comparing model results with filter measurements analyzed by thermal–optical methods.

OA mass can increase for constant OC, due to oxidative aging; this is something that very few models calculate, and should be improved in the future. The OA/OC ratio is discussed in more detail in Sect. ???. Care should also be taken for the case of methane sulfonic acid (MSA), since the letter A stands for “acid”, not “aerosol”, as in OA. When reporting MSA results, we refer to the total methane sulfonic acid mass present in OA and not its carbon mass only, unless clearly stated otherwise.

1.7 OA/OC and O/C ratios

To calculate the total organic aerosol mass concentration for each model, we apply the following equation:

$$\text{OA}_i = \text{OC}_i * (\text{OA}/\text{OC})_i \quad (1)$$

where $(\text{OA}/\text{OC})_i$ is the organic aerosol to organic carbon ratio for aerosol tracer i (Table 1). OA/OC, frequently termed as OM/OC in the literature (OM: organic matter), was found to correlate extremely well with the O/C ratio in Mexico City and chamber data (Aiken et al., 2008), because of low

N/C ratios. A low OA/OC ratio is also indicative of “fresh” OA as deduced from observations (Turpin and Lim, 2001; Philip et al., 2014). The OA/OC ratio varies greatly between models, with many of them setting OA/OC = 1.4 as a constant for all OA sources. Some models use different OA/OC ratios for every OA tracer: IMAGES, IMPACT, and the two TM4-ECPL models calculate the specific OA/OC ratio for each of their aerosol tracers, depending on their sources and chemical identity. CAM4-Oslo uses 1.4 for fossil fuel and biofuel, OsloCTM2 and SPRINTARS use 1.6, while all three models use 2.6 for biomass burning. In the case of CAM4-Oslo and SPRINTARS, it is not possible to calculate the OC concentration from the model fields accurately, since they only track one tracer. For this, we used a single value, that of the fossil fuel each model is using, which will lead to an underestimation of their OC concentration (but not of OA) close to biomass burning sources. The remaining models use a constant OA/OC ratio: GEOS-Chem and GEOS-Chem-APM use a specified value of 2.1, GISS-CMU-VBS and GISS-CMU-TOMAS use 1.8, and all other models use 1.4. Observations (Turpin and Lim, 2001; Aiken et al., 2008) suggest that OA/OC values of 1.6 ± 0.2 and 2.1 ± 0.2 are good approximations for urban and non-urban aerosols, respectively, indicating that most models might use OA/OC values that are low. The study of both the OA/OC and O/C ratios is extremely important and warrants a dedicated investigation; although this will be mentioned in the present work, it will be studied in detail in the future.

1.8 Organic aerosol speciation

In the present work, we have separated organic aerosols into five categories, as described below and summarized in Table 1. The models are then grouped based on their OA parameterizations in Table 2.

1. tPOA, for terrestrial primary organic aerosol, which includes primary emissions from fossil fuel, biofuel and biomass burning. All models participating in this intercomparison include these three tPOA sources. Several models also consider a biogenic secondary organic aerosol source that is included in tPOA (BCC, CAM4-Oslo, CanAM-PAM, ECHAM5-HAMMOZ, ECHAM5-SALSA, ECMWF-GEMS, EMAC, GISS-CMU-TOMAS, GISS-MATRIX, GISS-TOMAS, GMI, GOCART, LMDz-INCA, SPRINTARS and TM5), as discussed earlier. This is considered to be linked with monoterpene emissions (Guenther et al., 1995), producing non-volatile aerosol mass with a fixed yield as discussed in Sect. ???. Some models have a simplified chemistry that produces non-volatile SOA, also included in tPOA: in GISS-CMU-TOMAS and GISS-TOMAS a generic SOA precursor is emitted in the gas phase representing all SOA precursor gases (Dentener et al., 2006; 15% of the monoterpene

emissions, emitted in the gas phase) with a chemical lifetime of 12 h that forms a non-volatile SOA tracer (which is included in tPOA). In GISS-TOMAS the SOA precursor emissions are based on terpenes, with a 10 % yield, while *a*-pinene oxidation by all major oxidants (OH, O₃, NO₃) produces non-volatile SOA (included in tPOA) with a 13 % yield in GLOMAPbin and GLOMAPmode. SPRINTARS has a 9.2 % yield of non-volatile SOA (Griffin et al., 1999a, b) from monoterpene emissions, and considers this tracer as inert and tracks it separately, in contrast to the other models that produce non-volatile SOA and track it together with tPOA. SOA from anthropogenic VOCs is included in only a few models, and is not included in tPOA.

2. mPOA, for primary organic aerosol from marine sources. CAM4-Oslo has a primary marine organic source of 8 Tg a⁻¹ (Spracklen et al., 2008) with the same emissions distribution as sea salt (provided by Dentener et al., 2006) included in tPOA. IMPACT includes a mPOA source of 35 Tg a⁻¹ (Gantt et al., 2009a), which scales with chlorophyll *a* and sea salt as a proxy of marine biological activity (O'Dowd et al., 2004), while GISS-modelE-G/I and TM4-ECPL-F/FNP include a similar source of submicron mPOA based on Vignati et al. (2010). The GISS-modelE-G/I source is described in Tsigaridis et al. (2013) and the TM4-ECPL-F/FNP mPOA source in Myriokefalitakis et al. (2010). It has to be noted that these two studies have a factor of 10 difference in submicron mPOA source strength, despite having very similar source function parameterizations. This results from differences in sea-spray size distribution assumptions, as discussed in Tsigaridis et al. (2013). In addition to the fine-mode mPOA source, TM4-ECPL-F/FNP accounts for about 30 Tg C a⁻¹ of coarse-mode mPOA (Kanakidou et al., 2012), but that was not taken into account in the present study, since all measurements used here are for fine aerosols.
3. trSOA, for “traditional” secondary organic aerosol, which is produced by gas to particle mass transfer of secondary organic material, either assuming the material has a finite vapor pressure (a gas-particle partitioning process) or that it has zero vapor pressure (a condensation process). The most common precursors of SOA used across models are isoprene and terpenes, although few models have other precursors as well, as presented in Sect. 2. All models have some form of trSOA, either included in tPOA (as explained above), or via an explicit treatment of the semi-volatile oxidation products of the precursor VOCs. For the models other than the ones presented in (a) above that treat SOA as part of tPOA, the approach used and species taken into account differ. CAM5-MAM3 prescribes mass yields from 5

trSOA precursor categories (isoprene, terpenes, aromatics, higher molecular weight alkanes and alkenes, with yields of 6.0, 37.5, 22.5, 7.5, and 7.5 %, respectively), which then reversibly and kinetically partition to the aerosol phase. GISS-CMU-VBS uses the volatility-basis set, but without aging for the biogenic trSOA. The rest of the models use the two-product model approach to calculate trSOA; see the references column in Table 3 for more details. GEOS-Chem-APM considers the volatility changes of the gaseous semi-volatile compounds arising from the oxidation aging process, as well as the kinetic condensation of low-volatility gases (Yu, 2011). HadGEM2-ES does not calculate trSOA online; instead, it uses an offline 3-D monthly mean trSOA climatology obtained from the STOCHEM CTM (Derwent et al., 2003). The two-product model implemented in IMAGES was modified to account for the effect of water uptake on the partitioning of semi-volatile organics, through activity coefficients parameterized using a detailed model for *α*-pinene SOA (Ceulemans et al., 2012). IMPACT predicts semi-volatile SOA from organic nitrates and peroxides using the gas-particle partitioning parameterization with an explicit gas-phase organic chemistry. These condensed semi-volatile compounds are assumed to undergo further aerosol-phase reactions to form non-evaporative SOA with a fixed 1-day e-folding time (Lin et al., 2012). The two TM4-ECPL models account for SOA aging by gas-phase oxidation by OH with a rate of 10⁻¹² cm³ molec⁻¹ s⁻¹, while the conversion of insoluble POA to soluble is parameterized as described by Tsigaridis and Kanakidou (2003) with a decay rate that depends on O₃ concentration and water vapor availability, which corresponds to an approximately 1-day global mean turnover time, with strong spatial variability.

4. ntrSOA, for non-traditional secondary organic aerosol, which comes from a variety of sources, as explained below. GISS-CMU-VBS includes the VBS (Robinson et al., 2007), which allows tPOA to evaporate and age (via oxidation) in the gas phase, producing less volatile gas-phase products, which can again partition between the gas and aerosol phases. This model, which is the only one in the present study that takes into account the intermediate-volatility species as additional sources of OA, enables the application of the partitioning theory to POA and its associated vapors as well, not only SOA. The aerosol phase of these oxidized products is termed ntrSOA. The impact of this process strongly affects the chemical composition of SOA and will be discussed later (Sect. 4.3.3). Other models, namely IMAGES, IMPACT, and TM4-ECPL-F/FNP, include an aqueous-phase oxidation pathway of small organic molecules like glyoxal and methylglyoxal that produces low-volatility compounds and oligomers in cloud and

aerosol water (Fu et al., 2008, 2009; Stavrou et al., 2009; Myriokefalitakis et al., 2011), with the two TM4-ECPL models having a primary glyoxal source from the oceans of 4.1 TgC a^{-1} , which is not present in the other two models. Glyoxal and methylglyoxal are highly reactive species in the aqueous phase. The aqueous-phase reactions can occur both in aerosol water and cloud droplets; after droplet evaporation, the residual organic compounds remain in the aerosol phase in the form of OA. By applying a reactive uptake (γ) of glyoxal and methylglyoxal on aqueous particles and cloud drops (Liggio et al., 2005), IMAGES and IMPACT parameterized the irreversible surface-controlled uptake of these soluble gas-phase species. On the other hand, Myriokefalitakis et al. (2011) applied a much more detailed aqueous-phase chemical scheme in cloud droplets in order to produce oxalate. For IMPACT, 52 % of the total SOA comes from glyoxal and methylglyoxal multiphase chemistry (Lin et al., 2012). IMPACT also includes ntrSOA formation from the uptake of gas-phase epoxides onto aqueous sulfate aerosol (Paulot et al., 2009), which contributes by 25.1 Tg a^{-1} (21 %) to the total SOA formation (Lin et al., 2012).

5. MSA, an oxidation product of DMS, is also a SOA component. Although a minor organic aerosol component on the global scale, MSA can be very important in remote oceanic regions, especially when mPOA is relatively low: observations indicate that MSA can be at least 10 % of the total WSOC mass (Sciare et al., 2001; Facchini et al., 2008) at marine locations. Only CAM4-Oslo, GEOS-Chem-APM, GISS-modelE-G/I, IMPACT, LMDz-INCA, TM4-ECPL-F/FNP and TM5 have this tracer, which has been typically neglected from the organic aerosol budget in modeling studies. In CAM4-Oslo, MSA is included in tPOA, in IMPACT it is included in mPOA (which is in turn included in tPOA), whereas in the other models, it is individually tracked.

A summary of the OA processes included in the models is presented in Table 2. The total organic aerosol mass is calculated as follows:

$$\text{OA} = \text{tPOA} + \text{mPOA} + \text{trSOA} + \text{ntrSOA} + \text{MSA} \quad (2)$$

The models that have mPOA, SOA and/or MSA included in tPOA do not track them separately, so there is no risk of double-counting any OA species. In addition to this categorization, in order to compare with AMS data (see Sect. ??) we separate the modeled OA into HOA (hydrocarbon-like OA) and OOA (oxygenated OA) as defined by Zhang et al. (2005), when sufficient information on hydrophobic/hydrophilic speciation from the models is available. We use the terminology HOA/OOA instead of water soluble/insoluble OC (WSOC/WIOC), and compare only with AMS organic aerosol data, in order to

contrast with the OC measurements that refer to organic carbon. The separation into HOA and OOA has been provided by only a few models: ECHAM5-HAM2, ECMWF-GEMS, EMAC, GISS-modelE-G, GISS-modelE-I, GISS-TOMAS, GLOMAPbin, GLOMAPmode, IMAGES, LMDz-INCA, TM4-ECPL-F, TM4-ECPL-FNP and TM5. From the AMS perspective, the total OA is calculated as follows:

$$\text{OA} = \text{HOA} + \text{OOA} \quad (3)$$

Further subdivisions into other categories of OOA (Jimenez et al., 2009) are neglected in this study. In addition, the term POA used in Zhang et al. (2011) as a surrogate for different HOA categories is also not taken into account here.

2 Description of models

The models participating in the present study differ in (a) the spatial resolution, both horizontal and vertical, (b) the underlying model with which the aerosol calculations are coupled, which can be either a CTM or a GCM, and will be named “host model” from now on, (c) the emissions used, both for POA and SOA precursors, as well as for other gaseous and aerosol tracers, (d) the inclusion or not of aerosol microphysics, which are implemented in multiple ways (Mann et al., 2014), and (e) the OA processes simulated, i.e., the chemical and physical processes that change existing OA (such as oxidative aging), and the representation of SOA formation.

The complexity of the OA calculations varies greatly between models (Table 3). There are differences in OA emission source strength, both for primary particles (Table 4) and precursors of secondary OA (Table 5), as well as in the total number of OA tracers used (2 to 62; Table 1) and their properties, especially with regard to the temperature dependence of their vapor pressure (Sect. 6). Although a classification is difficult, one can categorize the models in various groups when considering OA modeling from different perspectives. The classification used here will be presented later (Sect. 1.8).

Some models using the same host model have very specific (and not necessarily a few) differences. ECHAM5-HAM2, ECHAM5-HAMMOZ, ECHAM5-SALSA and EMAC use the same host model (ECHAM5) but different aerosol parameterizations: the first two use M7 (modal), ECHAM5-SALSA uses SALSA (sectional) and EMAC uses a modified version of M7. ECHAM5-HAMMOZ uses the previous version of the HAM aerosol module, which does not take into account the detailed SOA formation introduced in ECHAM5-HAM2 (O’Donnell et al., 2011). GEOS-Chem and GEOS-Chem-APM use the same host model (GEOS-Chem) but different aerosol representations: the first uses the default bulk aerosol scheme, while the latter uses a size-resolved (bin) advanced particle microphysics (APM) module (Yu and Luo, 2009). GISS-CMU-VBS and GISS-CMU-TOMAS use the same host GCM (GISS-II’), with the only

difference being in the calculation of OA: the first one uses a bulk aerosol scheme with the VBS approach (Donahue et al., 2006; Jathar et al., 2011), and the second one the aerosol microphysics scheme TOMAS (Adams and Seinfeld, 2002; Lee and Adams, 2010, 2012). Similarly, GISS-MATRIX, the two GISS-modelE models and GISS-TOMAS use the same host GCM (GISS-E2), but they have different aerosol representations: GISS-MATRIX uses the aerosol microphysics module MATRIX (Bauer et al., 2008), the two modelE versions have a bulk aerosol scheme (Koch et al., 2006, 2007; Miller et al., 2006; Tsigaridis and Kanakidou, 2007; Tsigaridis et al., 2013) and GISS-TOMAS uses the same aerosol microphysics scheme as GISS-CMU-TOMAS (Lee and Adams, 2012; Lee et al., 2014). GISS-modelE-G and GISS-modelE-I only differ in the emissions used; they both have CMIP5 anthropogenic emissions for all tracers (Lamarque et al., 2010), but GISS-modelE-G uses GFED3 (van der Werf et al., 2010) for biomass burning. GLOMAPbin and GLOMAPmode use the same host CTM (TOMCAT; Chipperfield, 2006), with the only difference being the sectional and modal aerosol microphysics calculations (Mann et al., 2012). TM4-ECPL-FNP is almost identical to TM4-ECPL-F, but also takes into account the contribution to OA from primary biological particles and soil dust in the fine and coarse modes (Kanakidou et al., 2012). These two models also use different biogenic and anthropogenic VOC emission inventories (Tables 4 and 5).

All model results presented here come from monthly mean data, while measurements are averaged in monthly mean values, prior to any comparison with model data.

2.1 Meteorology

One major difference between the configurations of the models is the meteorology and meteorological year used. This affects aerosol transport, removal, chemistry (e.g., temperature dependence of reaction rates) and gas-particle partitioning of semi-volatile species. In some models, meteorology also directly affects natural aerosol emissions, like wind-driven sea salt, marine organic aerosol, dust and VOC emissions from the vegetation and oceans. Indirectly, meteorology affects MSA sources, since MSA is produced via dimethyl sulfide (DMS) oxidation, whose source is affected by wind speed and its oxidation depends on chemical rates.

Several climate models that participated in this intercomparison calculate the meteorology online. These are BCC, CAM4-Oslo, CAM5-MAM3, CanAM-PAM, GISS-CMU-VBS and GISS-CMU-TOMAS. In addition, climate models GISS-MATRIX, GISS-modelE-G, GISS-modelE-I and SPRINTARS are nudged to the NCEP reanalysis (Kalnay et al., 1996), GISS-TOMAS is nudged to MERRA meteorology (Rienecker et al., 2011), HadGEM2-ES and ECHAM5-HAMMOZ are nudged to the operational ECMWF meteorology (<http://www.ecmwf.int/products/data/archive/descriptions/od>), and LMDz-INCA is nudged to ECMWF reanalysis from the Integrated Forecast System.

The remaining models use a variety of prescribed meteorology data sets for the year 2006 (Table 3), except that GISS-CMU-VBS uses 2008, IMPACT uses 1997, and TM4-ECPL-FNP uses 2005.

2.2 Emissions

All participating models include POA in their simulations. The sources are both anthropogenic and biogenic, and can be classified as follows:

1. Fuel emissions. These exclusively anthropogenic sources include fossil fuel and biofuel burning. All models include these sources, but the emission inventories used are not always the same (Table 4). A number of models used emissions for the year 2000; others used emissions for the year 2006, and one for the year 2005 (TM4-ECPL-FNP). Cooking emissions, which can contribute up to 50 % of the POA in many urban areas (Mohr et al., 2012) are not included in any model.
2. Biomass burning. As in the case of fuel emissions, not all models use the same sources or representative years. Only about half of the models use biomass burning emissions from the year 2006 (Table 4), which is the reference year in the present study. Biomass burning is the largest POA source; it has significant interannual and strong seasonal variability and is the most uncertain POA source on a global scale (Andreae and Merlet, 2001), making it extremely important for comparison with measurements, especially at remote sites, to properly represent this source. Comparisons of several model simulations with the smoke aerosol optical depth (AOD) observed by MODIS have indicated a systematic underestimation when emissions from bottom-up inventories like GFED, used by several models here, are used. The underestimation may be as high as a factor of 3 on the global scale (Kaiser et al., 2012, and references therein), and strongly varies by region (Petrenko et al., 2012).
3. Marine sources. Few models take into account marine sources of organic aerosols (see Sect. 1.8); these depend on sea spray emissions. The GISS-modelE-G and GISS-modelE-I source depends on SeaWiFS chlorophyll *a* measurements from the year 2000 (Tsigaridis et al., 2013), while IMPACT and TM4-ECPL-F/FNP calculations use the MODIS chlorophyll *a* data from the corresponding simulated year. However, recent observations indicate the presence of marine organic aerosol over oceanic oligotrophic areas (Long et al., 2011); this can be either due to long-range transport, or a missing source not accounted for with the current source parameterizations, or both. CAM4-Oslo also has marine organic emissions, with a global flux based on Spracklen et al. (2008), and a spatial distribution given by the

prescribed AeroCom phase I fine-mode sea salt emissions (Dentener et al., 2006).

4. Other primary sources. TM4-ECPL-FNP (Kanakidou et al., 2012) includes some fine-mode POA sources that do not exist in any other global model in this intercomparison. These consist of primary biological particle emissions from plants (25 Tg a^{-1}) and soil organic matter on dust (0.2 Tg a^{-1}).
5. “Pseudo” primary non-volatile SOA fluxes. A number of models parameterize SOA chemical production in the atmosphere as a source of non-volatile aerosol emitted directly from vegetation. SOA is then modified similarly to POA by processes like transport, chemical aging, growth, coagulation and condensation, among others, depending on the model. BCC, CanAM-PAM, ECHAM5-HAMMOZ, ECHAM5-SALSA, ECMWF-GEMS, EMAC, GISS-CMU-TOMAS, LMDz-INCA and TM5 use a global source of 19.1 Tg a^{-1} (Dentener et al., 2006). This source is equivalent to a 15 % yield from the year 1990 monoterpene emissions (Guenther et al., 1995) and is identical to the source used during the AeroCom phase I experiments. GISS-CMU-TOMAS, GISS-TOMAS, GLOMAPbin and GLOMAPmode also use the same approach (based on the Guenther et al. (1995) emissions, except GISS-TOMAS, which is based on Lathière et al., 2005), but with SOA produced according to an assumed molar yield following oxidation (see Sect. 1.8 and Table 1), which results in a calculated SOA source of 19.1, 17.1, 23.1, and 23.0 Tg a^{-1} , respectively. GISS-MATRIX and GISS-TOMAS use a 10 % yield (17.1 Tg a^{-1}) from monoterpene emissions for the year 1990 from Lathière et al. (2005), while GMI and GOCART assume a 10 % yield (12.7 Tg a^{-1}) from the Guenther et al. (1995) monoterpene emissions. In the case of CAM4-Oslo, the strength of the secondary source suggested by Dentener et al. (2006) has been scaled up to 37.5 Tg a^{-1} , based on Hoyle et al. (2007).

In addition to the primary aerosol emissions, the inventories used for the precursors of secondary organic aerosols are also both very diverse and of great importance. These are presented in Table 5.

3 Measurements

The compilations of $\text{PM}_{2.5}$ OC measurements by Bahadur et al. (2009) and PM_1 OA measurements by Zhang et al. (2007) form the basis for the present study. Additional OC and OA observations from continuous monitoring networks and individual case studies reported in the literature have been used to increase the spatial and temporal coverage of the observational database for model evaluation.

The OC measurements reported by Bahadur et al. (2009) include data from

- The Interagency Monitoring of Protected Visual Environments (IMPROVE; <http://vista.cira.colostate.edu/IMPROVE>), which is the American monitoring network for national parks and wilderness areas, for 1988–2006.
- The Speciated Trends Network (STN) administered by the Environmental Protection Agency (Air Quality System Environmental Protection Agency (AQSEPA); <http://www.epa.gov/ttn/airs/airsaqs>), which mainly consists of urban monitoring stations within the USA, for 2000–2007.
- The North American Research Strategy for Tropospheric Ozone (NARSTO; <http://www.narsto.org>), which consists of measurements in Mexico, the USA and Canada, for 1999–2005.
- The New England Air Quality Studies (NEAQS; <http://www.esrl.noaa.gov/csd/projects/neaqs>), which contains measurements from the New England region, as a part of NOAA field studies, for 2002.
- The Southeastern Aerosol Research and Characterization Study (SEARCH; Hansen et al., 2003), which is a monitoring network for the southeastern United States, for 1998–2007.
- The European Monitoring and Evaluation Programme (EMEP; <http://www.emep.int>). EMEP is a European monitoring network with a few hundred monitoring stations all over Europe; only a few measure OC, which are used here, for 2002–2006.
- The Construction, Use and Delivery of a European Aerosol Database (CREATE; <http://www.nilu.no/projects/ccc/create>). CREATE is a database that compiles aerosol data from eight European countries, for 2000–2006.
- The Hong Kong Environmental Protection Agency (<http://www.epd.gov.hk/epd/eindex.html>), with measurements from the extended area of Hong Kong, for 2000–2002.

These data sets have been extended by numerous new measurements from published studies (Chow et al., 1993; Smith et al., 1996; Zappoli et al., 1999; Kim et al., 2000; Eatough et al., 2001, 2003; Krivacsy et al., 2001; Artaxo et al., 2002; Balasubramanian et al., 2003; Gatari and Boman, 2003; Graham et al., 2003; Long et al., 2003; Fan et al., 2004; He et al., 2004; Ho et al., 2004; Jeong et al., 2004; Salma et al., 2004; Sawant et al., 2004; Ward et al., 2004; Xu et al., 2004; Hueglin et al., 2005; Kim et al., 2005; Fuzzi et al., 2006; Koulouri et al., 2008; Pindado et al., 2009; Sciare et al., 2009; Li et al., 2010; Shakya et al., 2010; X. Y. Zhang et al., 2012) enhancing primarily the spatial, but also the temporal availability of comparison points.

The IMPROVE and AQSEPA networks cover most of the United States more than adequately. The EMEP monitoring network together with the European Integrated project on Aerosol, Cloud, Climate, and Air Quality Interactions (EU-CAARI) and CREATE data sets and other studies found in the literature provide good coverage of a large part of Europe, with stations in 17 countries. Although the spatial and temporal coverage is not as extensive as in the USA, it provides a comprehensive representation of different sources and chemical environments over Europe. There are limited measurements from Asia, with many of them being at urban or urban-influenced locations in India and China. South America, Africa and Oceania have very poor spatial and temporal coverage, despite the importance of the tropical forests of the former two on the global OA budget. Marine areas are almost exclusively covered by short-term measurement campaigns, with the exception of Amsterdam Island in the southern Indian Ocean (Sciare et al., 2009). All OC measurements are $PM_{2.5}$ or smaller sizes, e.g., $PM_{1.8}$ (Koulouri et al., 2008).

A rapidly increasing number of AMS OA measurements has been reported in the literature since the work of Zhang et al. (2007). Most of these AMS measurements are available online, in a web page created and maintained by Q. Zhang and J.-L. Jimenez (<http://tinyurl.com/ams-database>). We include in this analysis most of the ground-based data available as of January 2013. These data include the only AMS measurements so far available for a whole year (using the ACSM instrument, which is a monitoring version of the AMS; Ng et al., 2011), from Welgegund, South Africa (Tiitta et al., 2014); all other stations were measuring for about a month or less. The geographical coverage of the AMS stations is far less dense than the OC measurement locations, but the number of stations is rapidly increasing. Longer records are also starting to appear in the literature (Tiitta et al., 2014), and are expected to increase in the near future. It is important to note that the OA values provided by the AMS-type instruments have uncertainties (30 %) inherent in quantifying the detection efficiency for the wide range of organic molecules that make up complex SOA material (Canagaratna et al., 2007; Middlebrook et al., 2012). Care should be taken when using AMS-type OA data in models that estimate organic aerosol content.

All station data have been classified in three main categories: urban, remote and marine. Urban sites are defined as those that are either in cities or highly influenced by them. AMS stations characterized as “urban downwind” fall in this category. Remote sites are defined as those not influenced by local anthropogenic activities, and include forested regions, mountains, rural areas, etc. Marine sites are all measurements from ships or from coastal stations that are highly influenced by the marine atmosphere. Only two AMS stations fall into this category (Okinawa, Japan, and Mace Head, Ireland), and for simplicity, they were classified in the “remote” category.

The two databases (OC and OA measurements) have been kept separate because of the added complexity related to the

OA/OC ratio (Sect. ??). Almost all models calculate OA mass concentration, integrated across the fine-mode size distribution where appropriate, which can be compared with AMS measurements without any unit conversion. To compare with filter measurements of OC, we used the models' assumptions about the OA/OC ratio to convert the modeled OA to OC. As mentioned earlier, the importance of the OA/OC ratio will be explored in the future. The cutoff diameter of aerosols can also be an issue (Koulouri et al., 2008), but it is not expected to be significant in the present study, given the assumptions that the models adopt for the primary OA sources. No model adds fine OA mass from coarse-mode sources, and no model allows partitioning of semi-volatile gases to the coarse mode; thus, the difference between the $PM_{2.5}$ filter measurements and PM_1 AMS data is not expected to be properly resolved by models, even if they include aerosol microphysics calculations.

4 Results

4.1 Global budgets

Many global models have evolved significantly since the AeroCom phase I intercomparison studies. During phase I, the first experiment, AeroCom A (ExpA), was designed in a very similar way to the AeroCom phase II model simulations described here (Schulz et al., 2009). For the second, AeroCom B (ExpB), all models used the same emission inventories. The outcomes of these studies have been summarized by Textor et al. (2006) for ExpA and Textor et al. (2007) for ExpB and is compared with the present study in detail here (Fig. 1). The two AeroCom phase I studies focused on the total aerosol budget, but the individual aerosol components were also studied. Sixteen models participated in ExpA and twelve in ExpB, most of which are earlier versions of the models that participated in the present intercomparison.

The large number of models used in this study adds a significant level of complexity to the interpretation of results, due to the large diversity of inputs and configurations used by the different modeling groups. Despite the large differences between model formulations, on the global scale, several interesting similarities and patterns appear, which are frequently associated with the parameterizations and emission inventories used.

4.1.1 Emissions

Global mean model POA emissions used in the models are in the range of 34–144 $Tg a^{-1}$. The emissions in most models lie below 80 $Tg a^{-1}$ (Fig. 2), with a median value of 56 $Tg a^{-1}$. Notable exceptions are the two GISS-modelE models (G and I), in which about two thirds of the POA emissions come from marine sources (Tsigaridis et al., 2013); without this source, these two models have the same emissions as GISS-MATRIX (39.5 $Tg a^{-1}$), which falls below the




Figure 1. Box and whisker plot for all POA, SOA and OA global budgets and comparison with AeroCom phase I (Textor et al., 2006, 2007) results. The boxes represent the first and third quartile range (50 % of the data), the line is the median value, the star is the mean, and the error bars represent the 9/91 % of the data. Outliers are presented with x-symbols, with the corresponding color of the model, and the numbers of models participating in each bars statistics are presented with a grey number at the top. The AeroCom phase I outliers are presented with black color, since there is no direct correspondence with the models that participate in the present study. Bar colors are POA (brown), SOA (green), OA (blue), AeroCom A (red; Textor et al., 2006), and AeroCom B (orange; Textor et al., 2007).

25 % quantile. CAM4-Oslo also has the highest terrestrial sources of all models (144 Tg a^{-1}), followed by IMPACT (98 Tg a^{-1}) and EMAC (92 Tg a^{-1}). All models appear to have similar seasonality in POA emissions that are driven by tPOA, with increased emissions during Northern Hemisphere summer due to the enhanced contribution of Northern Hemisphere biomass burning emissions from temperate and boreal forests to the total POA fluxes. In addition, several models include SOA sources in tPOA as explained earlier, scaled by BVOC emissions, which also peak during Northern Hemisphere summer (Guenther et al., 1995, 2006); this contributes to a seasonal cycle of tPOA that is caused by the trSOA treatment as part of tPOA, and should not be interpreted as a tPOA seasonality. Also note that contrary to biomass burning, anthropogenic tPOA sources have no seasonality in their emission inventories. The IMPACT model appears to have the opposite seasonality, with maximum POA emissions during winter and minimum from late spring to early summer, due to the fossil fuel emissions scaling to fit observations (Wang et al., 2009). The minimum of the emissions for all models except IMPACT is during Northern Hemisphere spring, when neither biomass burning nor the photochemical trSOA sources (included in tPOA by many models) are high.

The POA emissions variability from phase II is roughly the same as that of the OA variability from ExpA, which indicates that the significant uncertainties in the POA emissions in global models since AeroCom phase I have not been reduced. However, some models have very high POA emissions, due to the recently developed parameterizations of mPOA sources in global models. These highly uncertain sources were absent in AeroCom phase I.

4.1.2 Chemical production

The chemical production of SOA is much more complex compared to the POA emissions. Firstly, many models include SOA sources as primary emissions, which are included in tPOA (see Sect. 1.8 and Table 1). This type of source was used during AeroCom phase I experiments (Dentener et al., 2006). The direct consequence of this assumption is that any uncertainties resulting from the OA sources in ExpA are only related to the POA emissions, since the SOA sources were identical across models. For AeroCom phase II, 13 out of 31 models still use this source parameterization (Table 2), while 5 models use a simple SOA production rate based on gas-phase oxidation, which then forms non-volatile SOA. These 18 models have a median SOA source strength of 19.1 Tg a^{-1} (mean 20 Tg a^{-1}) and a standard deviation of 4.9 Tg a^{-1} (Fig. 2). Very few models that include this source have provided budget information on the seasonal variability of its SOA source, since it is implicitly included in the tPOA sources and is not tracked separately. However, it has a virtually identical seasonality to that of the monoterpene emissions adopted in each model.

From the other models that include a more complex calculation of SOA chemical production, there is a large inter-model variability in the source flux, with median 51 Tg a^{-1} (mean 59 Tg a^{-1}) and 38 Tg a^{-1} standard deviation, based on 12 out of 14 models that include such parameterizations and have submitted budget information. This is more than twice as high as the models that use the AeroCom phase I parameterization, and with much larger model diversity. The seasonality of OA emissions in all these models peaks during Northern Hemisphere summer (Fig. 2), when VOC fluxes from temperate and boreal forests are at a maximum, while emissions from tropical forests are high year-round. Six




Figure 2. Top row: POA emissions included in models (before POA evaporation in the case of GISS-CMU-VBS); middle row: SOA chemical production (including the pseudo-primary SOA source, where applicable); bottom row: total OA sources (sum of top and middle rows) for the annual mean (left column; short dashes: mean; long dashes: median; dotted lines: 25/75 % of the data) and seasonal variability (right column). Note that not all models have submitted annual budget data, and fewer have submitted seasonal information; thus, their corresponding columns/lines are not shown. The models are grouped based on their complexity, as separated by vertical solid lines in the annual mean budgets. Groups from left to right: SOA is directly emitted as a non-volatile tracer; SOA is chemically formed in the atmosphere, but is considered non-volatile; SOA is semi-volatile; SOA is semi-volatile and also has VBS (GISS-CMU-VBS) or multiphase chemistry sources.

models (IMAGES, IMPACT, GISS-CMU-VBS, HadGEM2-ES, OsloCTM2 and TM4-ECPL-F) include very strong SOA sources of 120, 119, 79, 64, 53 and 49 Tg a^{-1} , respectively, followed by CCSM4-Chem (33 Tg a^{-1}) and GEOS-Chem (31 Tg a^{-1}). About 42 % (50 Tg a^{-1}) in IMAGES are due to non-traditional sources (glyoxal and methylglyoxal). The traditional SOA source in IMAGES accounts for water uptake, which is found to increase the partitioning of semi-volatile intermediates (Müller, 2009). Monoterpenes alone account for about 40 Tg a^{-1} . This large contribution is due to the very high SOA yields (~ 0.4) in the oxidation of monoterpenes by OH in low- NO_x conditions, which are justified by the formation of low-volatility compounds like hydroxy di-hydroperoxides (Surratt et al., 2010). IMPACT has sev-

eral non-traditional SOA sources from aqueous chemistry, which locally can contribute as much as 80 % of the total OA mass. CAM5-MAM3 and IMPACT also include anthropogenic precursors. CAM5-MAM3 also uses a factor of 1.5 SOA yield increase in order to reduce anthropogenic aerosol indirect forcing, by elevating the importance of SOA during the preindustrial period (Liu et al., 2012). As mentioned before, HadGEM2-ES does not calculate SOA production explicitly; instead, it uses the Derwent et al. (2003) climatology from STOCHEM, which calculates an SOA formation of 64 Tg a^{-1} . For comparison, satellite-constrained studies estimate that the total OA formation (primary and secondary) can be as high as 150 Tg a^{-1} , with 80 % uncertainty (Heald et al., 2010); AMS-constrained estimates put the total SOA

formation rate between 50 and 380 Tg a⁻¹, with 140 Tg a⁻¹ being the best estimate (Spracklen et al., 2011), while Halquist et al. (2009) estimated, using a top-down approach, that the best estimate for the total biogenic SOA formation is 88 TgC a⁻¹, out of a total 150 TgC a⁻¹ of OC.

The case of GISS-CMU-VBS deserves focus. This model calculates SOA production based on the VBS approach. Its secondary source of 79 Tg a⁻¹ includes not only newly formed SOA both from POA and intermediate-volatility organics, but also gas-phase chemical conversion of organic mass that has evaporated from emitted POA, to produce less volatile organics, i.e., mass that has undergone aging in the atmosphere. The traditional SOA sources from biogenic VOC are included in this model like in other models that use the two-product model, but also the chemical conversion of intermediate-volatility organics to less volatile OA is taken into account, again with the use of the VBS. Overall, GISS-CMU-VBS presents a similar seasonal pattern of SOA chemical production as other models, but shifted by one month, i.e., peaking in August, when biomass burning is at its maximum in the Northern Hemisphere, instead of maximizing in July, when photochemical activity and biogenic VOC emissions are higher globally. This might be due to the inclusion of the intermediate-volatility compounds as SOA precursors, which also have large biomass burning sources. CCSM4-Chem and GEOS-Chem also have a shift in the seasonal maximum. For CCSM4-Chem this is due to strong production from biomass burning sources, while in the case of GEOS-Chem the seasonal cycle seems to be driven by production from Amazonia, which is related with both biogenic and biomass burning emissions.

The total OA sources during ExpA were very similar to the total sources from the phase II experiments (median 97 Tg a⁻¹ both in ExpA and here), while ExpB had much lower total OA sources, 67 Tg a⁻¹ (Fig. 1). All of these sources include SOA, either as pseudo-emissions (phase I) or from a variety of parameterizations (phase II). The models from phase II present a much higher variability in their total OA sources, which is primarily attributed to the SOA chemical production variability that was not present in ExpA.

4.1.3 Burden

From the models that have submitted POA burden data (also termed load; the mean total mass in the atmosphere), both its seasonality and amplitude largely follow those of the corresponding POA emissions (Fig. 3), with two notable differences. The two GISS-modelE models have much lower POA burdens (but similar seasonality) than their emissions would imply. The reason is that the mPOA fraction of POA has a very short lifetime of ~ 1.5 days, since mPOA is assumed to be internally mixed with fine-mode sea salt, which is removed efficiently due to wet scavenging (Tsigaridis et al., 2013). This keeps the overall load of POA fairly low, and comparable with the models that do not have mPOA. The

other difference is GISS-CMU-VBS, which also has a much lower POA load than their emissions would suggest. This is due to the POA aging parameterization, which converts POA into SOA, drastically reducing the POA burden. The other models appear to have the expected POA load, given their emissions, including IMPACT, whose different seasonal variability of the emissions is also reflected on its OA load.

For the computed SOA load (Fig. 3), all models assume that SOA is very soluble, with 80–100 % of its total mass considered soluble, which results in similar globally averaged removal rates across the models. This means that the differences in the SOA loads are expected to be driven primarily by the SOA chemical production, similar to how the POA load is driven by emissions. This is indeed the case for almost all models, with GISS-CMU-VBS, IMAGES, IMPACT, CCSM4-Chem and CAM5-MAM3 having the highest loads, exceeding 1 Tg, with the first two models being as high as 2.3 and 2.2 Tg, respectively, and GEOS-Chem being just below 1 Tg. Spracklen et al. (2011) estimated a global SOA burden of 1.84 Tg, similar to the high-end models that participate in the current intercomparison, but for a SOA formation rate of 140 Tg a⁻¹, which is about 20 % higher than IMPACT and IMAGES (the models with the strongest SOA formation here), and about 3 times higher than the median SOA formation rate of the models that have a complex SOA parameterization. ECHAM5-HAM2 calculates an increasing load over the course of 1 year, which is related to the short spin-up time of 3 months, which is not sufficient for the upper tropospheric SOA to reach equilibrium. GEOS-Chem simulates an inverse seasonality when compared with other models, with the maximum load calculated during Northern Hemisphere winter and the minimum during Northern Hemisphere summer. The cycle seems to be dominated by the SOA load over the Southern Ocean; probably the removal processes are slower than other models there, thus SOA may form a uniform band between 30 and 50° S during the whole austral summer.

With regard to the total OA load, a median of 1.4 Tg (mean 1.6 Tg) and standard deviation of 0.8 Tg is calculated; half the models lie within the range of 1–1.6 Tg (Fig. 3). CAM4-Oslo calculates a global burden of 3.8 Tg, reflecting the very high POA emissions, while IMAGES, IMPACT, GISS-CMU-VBS and CCSM4-Chem calculate a burden of 3.7, 2.6, 2.4 and 2 Tg, respectively, as a result of their high SOA production. Overall, the models calculate very similar total OA load seasonality, which peaks during the Northern Hemisphere summer season, when both primary (biomass burning) and secondary (chemical production) OA sources are high, and minimizes during Northern Hemisphere spring, when neither biomass burning nor SOA chemical production is significant in the Northern Hemisphere. The tropical biomass burning and SOA production around December and January both contribute to the secondary maximum that all models calculate during that time. The relative importance of SOA over POA will be discussed in Sect. 4.3.3.




Figure 3. Same as in Fig. 2, for the POA/SOA/OA load.

The total OA load is calculated to be mostly lower than the sulfate load in the models that reported budget values for both aerosol components, with a median value of the $\text{OA}/\text{SO}_4^{2-}$ mass load ratio of 0.77 (mean 0.95). The ratio lies in the range 0.26–2.0; CAM4-Oslo, CAM5-MAM3, GEOS-Chem, GISS-modelE-G/I, IMAGES, IMPACT, and TM4-ECPL-F/FNP calculate values above 1, which means that annually on the global scale OA dominates over sulfate aerosols. That was the case for 5 out of 16 models during AeroCom phase I (Textor et al., 2006). Note however that AeroCom phase I models were simulating the year 2000, while here we simulate the year 2006; interactive chemistry, new sources (isoprene, mPOA and ntrSOA) and different emission inventories also contribute to significant differences between the two studies. One has to be reminded that even in AeroCom phase II, many models used some emission inventories from a year other than 2006 (Tables 4 and 5).

4.1.4 Deposition

Dry deposition is a minor removal pathway for OA, accounting for a median of 13 Tg a^{-1} (range $2\text{--}36 \text{ Tg a}^{-1}$) and a mean of 15 Tg a^{-1} (standard deviation of 10 Tg a^{-1} ; Fig. 4). On average, dry deposition is responsible for 15 % of the total OA removal across models. The two TOMAS models and TM5 calculate by far the lowest dry deposition flux of all, followed by three of the ECHAM5 models, excluding EMAC. The two TOMAS models use essentially the same aerosol microphysics parameterization in two different host models, GISS-II' for GISS-CMU-TOMAS and GISS-E2 for GISS-TOMAS. GISS-modelE-G/I and GISS-MATRIX use the same host model and identical emissions as GISS-TOMAS, a fact that suggests the TOMAS aerosol module (Adams and Seinfeld, 2002) either is less efficient in scavenging OA via dry deposition, or is more efficient in removing OA from the system via wet deposition, or both. The latter, though, would mean that the OA load (Fig. 4) would be

much smaller in GISS-TOMAS in order to have low enough dry deposition fluxes, which does not appear to be the case.

Other than the two TOMAS models, of the remaining models that have submitted dry deposition flux data, three models calculate very low fluxes: ECHAM5-HAM2, ECHAM5-HAMMOZ, and TM5, with the latter already mentioned earlier. The first two models use ECHAM5 as the host model, and all three use the M7 aerosol microphysics module (Vignati et al., 2004). As for the TOMAS case, this is strong evidence that the M7 module does not allow OA to deposit as fast as in most other models; ECHAM5-SALSA, which uses the same host model as ECHAM5-HAM2 and ECHAM5-HAMMOZ, calculates higher dry deposition fluxes than the two ECHAM5 models with M7. The largest difference in dry deposition between the two aerosol microphysics schemes comes from the treatment of external mixing of OA in the accumulation sized particles. ECHAM5-SALSA includes soluble and insoluble OA in the accumulation mode, while ECHAM5-HAMMOZ and ECHAM5-HAM2 include only soluble OA. In addition, EMAC, which uses a sectional version of M7 called GMXe, does not calculate as low a dry deposition as the models that use the modal version of M7. The fact that there are other models with aerosol microphysics parameterizations in this intercomparison, both modal and sectional, that do not calculate such low dry deposition fluxes, suggests that it is not a general aerosol microphysics calculation issue.

Comparisons of phase I models results for ExpA and ExpB strengthen this conclusion, since the model with the lowest OA dry deposition flux of ExpA (MPI_HAM; 5 Tg a^{-1}) and that of ExpB (TM5; 1.7 Tg a^{-1}) both use the aerosol microphysics module (M7). This scheme appears to be responsible for the lowest dry deposition fluxes calculated by the models that participate in the present intercomparison: the updated versions of these two phase I models, ECHAM5-HAM2, ECHAM5-HAMMOZ and TM5, participate in the phase II experiment and simulated the lowest dry deposition fluxes among all phase II models, together with the GISS-CMU-TOMAS and GISS-TOMAS models that did not participate in phase I. Whether the above explanation suffices to explain the low dry deposition, or other processes are involved as well, like very strong wet removal that does not allow time to dry deposition to become effective, the calculated aerosol size distribution, the aerosol properties that impact dry deposition rates, or something else, remains to be explored by dedicated deposition flux model–data comparisons. Also note that we have not assessed this feature of the models against observations, so we do not know which models are closer to observations.

CAM4-Oslo has the highest dry deposition flux of 36 Tg a^{-1} , which is due to the high OA load. BCC follows with 33 Tg a^{-1} , which is then followed by the two GISS-modelE models and IMAGES with $\sim 28 \text{ Tg a}^{-1}$. In the case of the two GISS-modelE models, this is due to the strong removal of mPOA, which is internally mixed with sea salt (as

explained earlier), while for IMAGES, it is due to the high OA load, as a result of strong trSOA formation. BCC uses a smaller mass mean diameter than the size distribution of POA emissions, which can explain the high dry deposition flux (Zhang et al., 2012). Despite these large differences between models, the calculated dry deposition fluxes follow the same seasonal pattern as the aerosol load presented earlier (Sect. ?? and Fig. 4).

The effective dry deposition rate coefficient, defined as the ratio of the dry deposition flux over the aerosol burden that is being deposited (Textor et al., 2006), ranges from 0.005 to 0.13 days^{-1} , with a median value of 0.025 days^{-1} , a mean value of 0.029 days^{-1} and a standard deviation of 0.046 days^{-1} . The diversity (defined as the standard deviation over the mean) has increased since AeroCom phase I, from 0.62 to 0.87. BCC has the largest effective dry deposition rate coefficient, 0.13 days^{-1} , more than double that of any other model. The models with very low dry deposition fluxes are the ones that have the lowest effective dry deposition rate coefficients, all below 0.014 days^{-1} , supporting the hypothesis that their dry deposition flux is probably too low.

By far the most important removal mechanism across all models is wet deposition (Fig. 4). Due to similar OA solubility assumptions across all models, the wet deposition flux largely follows the OA load, both in the annual budget and the seasonality. IMPACT has the highest wet deposition flux of all models (209 Tg a^{-1}), followed by IMAGES (163 Tg a^{-1}), CAM4-Oslo (146 Tg a^{-1}), CAM5-MAM3 (134 Tg a^{-1}), OsloCTM2 (128 Tg a^{-1}) and GISS-modelE-G/I ($120/125 \text{ Tg a}^{-1}$, respectively). These are the models with the highest OA sources (Fig. 2), thus also with the highest sinks. Wet removal of OA is simulated to range from 28 to 209 Tg a^{-1} for the 26 models that reported fluxes, with mean (median) standard deviation values of 86 (70) 43 Tg a^{-1} , which is on average 85 % of the total OA deposition.

The effective wet deposition rate coefficient ranges from 0.09 to 0.24 days^{-1} , with a median value of 0.15 days^{-1} , a mean value of 0.16 days^{-1} and a standard deviation of 0.04 days^{-1} . The diversity since AeroCom phase I has virtually not changed, with a slight increase from 0.27 to 0.28. OsloCTM2 has the highest effective wet deposition rate coefficient, and LMDz-INCA the lowest.

Wet removal, which together with aerosol sources is a major driver of the calculated aerosol lifetime and load, presents a much higher variability in the phase II models (Fig. 1). This is largely due to the consideration of SOA formation, which is responsible for the large variability in OA sources and burden in the models, as well as to differences in the assumptions on SOA solubility and aging.

4.1.5 Lifetime

The combination of all sources and sinks affects the load and lifetime of OA, either directly or indirectly. The lifetime of




Figure 4. Same as in Fig. 2, for the dry/wet OA deposition.

a species is calculated as the ratio of the species burden over its total removal; in the case of aerosols, the removal is dry and wet deposition. Unfortunately, while most model groups have submitted total OA diagnostics to calculate the OA lifetime, few have submitted the diagnostics required to calculate the global mean POA and SOA lifetimes.

The calculated median POA lifetime from the 13 models that reported relevant data is 4.8 days (mean 4.8 ± 1.4 days). The modeled lifetime ranges from 2.7 days for the two GISS-modelE models to 7.6 days for IMAGES (Fig. 5). The GISS-modelE models have the lowest lifetime, which is consistent with roughly two-thirds of POA being removed rapidly with sea salt (as mPOA). There is no clear seasonal signal on the calculated POA lifetime.

The SOA lifetime calculated by 12 out of 31 models also lacks a clear seasonal signal (Fig. 5). The GISS-modelE-G/I models, CCSM4-Chem, ECHAM5-HAM2 and GISS-CMU-VBS have the highest SOA lifetime of 15/14, 14, 13 and 10 days, respectively, which is related to large amounts of SOA in the upper troposphere, where there is virtually no removal mechanism and therefore SOA lifetime is enhanced, until atmospheric circulation or sedimentation brings it to lower layers where it becomes susceptible to removal. For the remaining models that provide information, the calculated SOA lifetime ranges from 2.4 to 6.8 days. The median SOA lifetime from all models that provide budget information is calculated to be 6.1 days (range 2.4–14.8 days), higher than the median POA lifetime. Anthropogenic POA, which in general is more hydrophobic than SOA, is almost exclusively emitted close to surface and below clouds, making

it more susceptible to dry and wet removal; biomass burning POA can be emitted at higher altitudes (Dentener et al., 2006), while a significant amount of SOA is formed above clouds in the models, where temperatures are low. For instance, in TM4-ECPL-FNP, about 42 % of the total SOA mass is formed in the free troposphere, while 98 % of POA mass is emitted in the boundary layer. Furthermore, although one might expect that SOA is more soluble, thus more susceptible to removal, this does not appear to be reflected in the model results; the reason is that SOA can be formed above clouds and avoid removal for long periods of time.

Twenty-four models provide sufficient information to calculate the total OA lifetime, which lies in the range of 3.8–9.6 days, with a median of 5.4 days and a mean of 5.7 ± 1.6 days (Fig. 5). GISS-CMU-TOMAS has a very strong seasonality in the calculated OA lifetime, with a maximum during late Northern Hemisphere spring and a minimum during late Northern Hemisphere fall, and GISS-CMU-VBS has the highest OA lifetime of all the models. As in the case of POA and SOA, there is no clear seasonality in the OA lifetime across models.

The high wet removal variability across all AeroCom phase II models is also reflected in the total OA load and lifetime (Fig. 1), where SOA presents a very high variability between models, especially in the case of SOA lifetime. This slightly increases the calculated variability of the total OA by the phase II models compared to phase I. This change is not so pronounced in the OA burden, due to the relatively low contribution of SOA to the OA load calculated by the models. This might change in the future, though, since SOA is

computed very high OA load, the other models have organic AOD spanning almost an order of magnitude, from 0.004 to 0.023, with a median value of 0.014. This is 8 % of the total AOD calculated by the same models.

4.2 Median model annual mean

4.2.1 Surface distribution

The composite annual mean OC and OA surface air concentrations, defined as the median of the regridded model fields to a $5^\circ \times 5^\circ$ horizontal resolution, exceed $0.5 \mu\text{g C m}^{-3}$ (or $\mu\text{g m}^{-3}$) across most continental regions, with maximum concentrations primarily over biomass burning regions and secondarily over industrialized areas (Fig. 7). The model diversity, defined as the ratio of the standard deviation of all models over their corresponding mean value calculated on the same grid, is smallest over and downwind continental regions, with ratios below 1 over most continental areas, and above 1 over the remote oceans (Fig. 7).

Diversity that exceeds 2 is evident over most of the oceanic regions south of 30°S and Antarctica, which is a result of the marine OA sources being present in only a few models. Ratios approaching 2 are also found over the northern Pacific and Atlantic oceans, and are also related to the marine OA sources. However, these local maxima are not as pronounced as in the Southern Hemisphere, due to (a) the much stronger seasonality, and (b) the stronger influence of continental aerosol sources in the Northern Hemisphere.

Over and close to the continents, the model diversity is low, except in three areas that present striking differences. Two are located over biomass burning regions, Indonesia and the Pacific borders of the USA and Canada, where the different emissions used by the models produce a large local diversity in concentrations. The third case is off the Pacific coast of Mexico; although this might also be related to biomass burning, the exact reason for the high model diversity is not clear, since this is not over an aerosol source area. Marine sources or different precipitation patterns in the models can also be part of the explanation; however, there are very few measurements (Shank et al., 2012) over that region, which hinders a definite conclusion.

Overall, it appears that the model diversity is low over and downwind of continental source regions, since the primary sources of aerosols are constrained by the availability of only a few different emissions inventories to be selected by the models. In addition, less constrained parameters like SOA and mPOA formation, long-range transport and OA removal (which affects OA lifetime) increase the model diversity over remote areas.

4.2.2 Vertical distribution

The vertical distribution of the mean OC simulated by all except three models (GOCART has only submitted surface

data, and GISS-CMU-TOMAS and GISS-CMU-VBS have not submitted all necessary fields for unit conversions) shows concentrations increasing with height up to a mean pressure level of about 800–900 hPa, and then decreasing with altitude (Fig. 8). The increase in concentration is due to (a) a maximum OC concentration over the tropics, where strong convection raises OC from the surface sources to the lower troposphere, (b) the SOA formation that largely takes place above the surface, (c) the biomass burning emissions that some models distribute to more layers than just the surface one, and (d) the absence of dry deposition above the surface (Fig. 9). A local maximum also exists at low altitudes over the industrialized northern mid-latitudes, although less pronounced than the tropical one. From the middle to the upper troposphere, the OC concentrations simulated by most models decline steeply with altitude. Some models show a secondary maximum at around 100–200 hPa, with concentrations much lower than the maximum near the surface, above which the concentrations decline even faster with height: CCSM4-Chem, ECHAM5-HAM2, ECHAM5-HAMMOZ, GISS-modelE-G/I, IMAGES, LMDZ-INCA, OsloCTM2 and SPRINTARS present a local minimum in concentrations around 400 hPa, which then increase, before dropping again above 100 hPa. The increase around the tropopause is due to the low temperatures that allow condensation of the semi-volatile SOA precursors that had not condensed at lower layers, or OA accumulation above clouds, where wet deposition does not happen, or both. The models that explicitly calculate SOA seem to have slower removal of SOA from these altitudes than the other models. In addition, uplift in strong convective regions of OA (both primary and secondary) can also explain this local maximum, due to transport of aerosols to layers of the atmosphere with very slow removal. The modeled vertical distribution of OA presents a diversity that spans over one order of magnitude.

The model diversity is relatively low in the lower troposphere (below 600 hPa) between 30°S and 60°N , but very high over the poles and near the tropopause (Fig. 9). A similar pattern was found for BC, sulfate aerosol and particles larger than $100 \mu\text{m}$ in dry diameter in another AeroCom phase II intercomparison study that focused on aerosol microphysics (Mann et al., 2014). This points out three important features: (a) the areas directly affected by strong primary and secondary sources around the tropics and northern mid-latitudes do not present a large diversity, due to the fairly similar emission inventories in the different models; (b) the primary marine sources of OA however are both highly uncertain and not present in many models, resulting in the high model diversity close to the surface over the Southern Ocean; and (c) the processes that involve low temperatures (which favor condensation of semi-volatile compounds) are not well constrained either, and they are also absent in many models, leading to very high model diversity over the poles and above 200 hPa. The vertical distribution of OA is thus very




Figure 6. Same as in Fig. 2, for OA all-sky aerosol optical depth at 550 nm.




Figure 7. Annual mean of the median model surface air concentration (left) and model diversity (right), defined as the standard deviation of the models over their mean, for OC (top) and OA (bottom) on a $5^\circ \times 5^\circ$ degree grid.

poorly understood, much less than its surface concentration, and deserves a dedicated study with thorough analysis.

4.3 Comparison with measurements

Many model–measurement comparisons can be performed with the extensive data set used here. The focus of the comparisons in the present study is to identify model strengths and weaknesses, and try to explain where and why the models are failing to simulate the measured concentrations. This will provide insight to directions for future model improvements. In parallel, we are also interested in understanding where and why the models successfully reproduce the observations, and focus on these areas in order to understand the role of the different model complexities on simulations

with comparable skill. It is not within the scope of this work to identify which model is the “winner” in simulating OA concentrations, especially since one model is unlikely to outperform the others on all metrics, but to provide information on the robustness of the model results. The present study focuses on the surface OC and OA concentrations. The sources and amount of OA in the upper layers of the atmosphere are not explicitly studied here, although accounted for in the OA budget terms discussed above. The detailed analysis of the vertical distribution of OA will be the topic of a future study.

Due to the very inhomogeneous spatial variability of measurements (supplementary material), only a general global model performance benchmark is performed here. Most data have been collected in the USA, followed by Europe and




Figure 8. Annual mean vertical profiles of OC (at ambient conditions) interpolated at 50 hPa steps from the surface to 50 hPa for OC: (a) Global, (b) northern mid-latitudes, (c) South America and (d) tropics.

China. The rest of the world, including some very important regions with regard to OA, are severely under-represented, or not represented at all. Such regions include all tropical forest areas (the Amazon basin, Africa and Southeast Asia) and the boreal forests of Canada and Russia. Long-term measurements in these areas are extremely scarce, with the only notable exception being Alta Floresta in the Amazon, where OC measurements for more than ten years are available.

4.3.1 Model skill

One of the major challenges when comparing global models with observations is whether the measurement locations are representative of the regional levels of the measured quantity in question. For most urban measurements, this is not the case, since the aerosol concentrations at urban centers are usually much higher than the regional background concentrations. Even a model with a very high horizontal resolution for a global model (like SPRINTARS) is not expected to cap-

ture the measurements at urban locations, since its grid cells are of the order of 100×100 km, which is still too coarse to accurately resolve urban pollution. Many of the “urban downwind” AMS data are also expected to fall into this category; thus we included them in the “urban” category.

For all stations, there are several instances where more than one measurement locations are present in a given grid box for a certain model. When this is the case, we use the arithmetic mean of the measurements for that specific grid box, in order to compare the single aerosol concentration the model is providing with a single measurement value.

When discussing the model ensemble results we use the median of all models, while we also analyze the mean normalized bias (MNB) of the models against measurements. The perfect comparison should have a $MNB = 0$ and correlation $r = 1$. The normalized bias (NB) at a given grid box is




Figure 9. Annual zonal mean of the median model results for OC interpolated at 50 hPa steps (left) and model diversity (standard deviation over mean; right).

calculated as follows:

$$NB_i = \frac{C_{\text{model},i} - C_{\text{meas},i}}{C_{\text{meas},i}} \quad (4)$$

where $C_{\text{model},i}$ is the modeled concentration in grid box i , and $C_{\text{meas},i}$ is the measured concentration in the same grid box. If more than one station exists in the same grid box, $C_{\text{meas},i}$ is the arithmetic mean of the individual stations. The model's MNB is derived as the arithmetic mean of all NB_i values.

Urban locations

The models perform poorly at urban locations, as expected. Most models strongly underestimate the measurements, having a median MNB of -0.64 (mean -0.62 , range -0.04 to -0.86) for OC (Fig. 10) and -0.51 (mean -0.48 , range -0.1 to -0.85) for OA (Fig. 11). CAM5-MAM3 appears in both OC and OA as an outlier, with a slightly negative MNB for OC and $+0.24$ for OA. As mentioned earlier, CAM5-MAM3 has an enhancement factor of 1.5 for the SOA formation, which might be part of the reason for the generally higher OA concentrations, which result in less bias, compared to the other models. Interestingly, the correlation of model results with measurements is slightly higher for the OA data; a median value of 0.54 is calculated for OA (mean 0.52, range 0.11 to 0.77), compared to 0.47 for OC (mean 0.43, range -0.09 to 0.70). Note though that the locations and temporal resolution of OC and OA measurements differ greatly, making a conclusive comparison between them impossible. In addition, these results are not representative of the overall performance of the models on the global scale; they only represent the models' ability to capture the available measurements, which are very inhomogeneously distributed in space and time (Supplement).

Remote locations

The models show a completely different behavior when compared with measurements of OC (Fig. 12) and OA (Fig. 13) at remote locations. Compared with the models' performance at urban stations, more models have more negative than positive MNB in the case of OC at remote locations, with the range spanning from -0.61 to 1.29 (median -0.15 , mean -0.02), while most models have a positive MNB in the case of OA, with a range from -0.38 to 2.17 (median 0.51, mean 0.70). It has to be noted, though, that the locations and times of OC and OA measurements are not the same, which means the model performance for OC and OA data are not directly comparable, due to the different spatial and temporal coverage of the stations. Only four models present relatively high positive MNB values when compared with the OC data: CAM5-MAM3 (1.3), EMAC (0.9), ECHAM5-SALSA (0.7) and ECMWF-GEMS (0.6). CAM5-MAM3 has the third highest SOA source of all models, but none of the other three models with strong positive MNB has exceptionally high POA or SOA sources (Fig. 2) and sinks (Fig. 4). All of EMAC, ECHAM5-SALSA and ECMWF-GEMS present a very strong maximum in the OC concentrations at the western border of the USA with Canada; monthly mean concentrations exceeding $200 \mu\text{gC m}^{-3}$ in EMAC (Fig. S3 in the Supplement) might be the reason for the positive MNB. Also note that EMAC emits all biomass burning aerosols at the surface, while most other models distribute them to a number of layers above the surface, typically within the boundary layer. The other models that present a positive correlation are all linked with either strong POA sources (CAM4-Oslo) or strong SOA sources (HadGEM2-ES and IMPACT), as presented in Fig. 2, but that is not the case for IMAGES, which has the highest SOA source, but presents a MNB of only $+0.1$, and TM4-ECPL-FNP, which has the 7th strongest SOA




Figure 10. Mean normalized bias (top) and correlation (bottom) of all models with urban OC measurements. When more than one measurement data point exists in a model grid, the corresponding mean of the measurements was used (see text). The mean across all models is shown with the short-dashed line, and the median with the long-dashed line. The perfect model has $MNB = 0$ and correlation = 1. The models are grouped like in Fig. 2.




Figure 11. Same as Fig. 10, for urban OA measurements.

source from the models that submitted their SOA chemical production, but presents the second strongest negative MNB of all the models.

Many models have a lower correlation with remote OC and OA measurements than with urban OC and OA. Al-

though this might appear unexpected, a possible explanation might be that urban pollution probably adds a large offset in the comparison, which does not affect correlation. In remote sites on the other hand, long-range transport adds one additional level of uncertainty in the model calculations,




Figure 12. Same as Fig. 10, for remote OC measurements.




Figure 13. Same as Fig. 10, for remote OA measurements.

which can result in lower correlation of the model results with measurements. The correlation coefficient against OC remote measurements rarely exceeds 0.5, with the correlation for about half of the models lying below 0.4 (median 0.39, mean 0.40, range 0.11–0.67), while when compared against the remote OA measurements the correlations are slightly lower, with a median and mean value of 0.37 (range

0.07–0.55). It is possible that either a remote source is missing or treated in a too simplistic way, or that the transport and lifetime (which largely depend on solubility, representation of precipitation from clouds, and poorly represented ageing processes) of organic aerosols in the regional and remote atmosphere are not properly calculated in models, or that the seasonality of sources is not represented accurately,




Figure 15. Annual mean OA/OC at surface as calculated by IMAGES (a), IMPACT (b), OsloCTM2 (c) and TM4-ECPL-F (d).

able to simulate aqueous-phase OA formation and comparison with targeted observations are needed to consolidate the importance of this process for the OA budget. The primary marine source also improves the comparison over the oceans (Fig. 23), but further work is needed to constrain this source. Overall, the median and mean correlations are very close (0.25 and 0.24, respectively), and the correlation range is from -0.03 to $+0.41$.

Importance of model complexity

In the comparisons of model results with urban station data, the correlations with OA observations were higher than those with OC. Urban aerosols are mostly fresh, compared to the more aged ones at remote locations. All models simulate OA, and then the OA/OC ratio is used to convert from OA to OC, in order to compare with OC data. Emission inventories however are frequently in units of carbon, not organic matter, adding an additional conversion, thus uncertainty, in the models. Using the same OA/OC ratio to convert emissions and then the simulated concentrations implies that the OA/OC ratio has not changed with atmospheric processing. This is clearly a weak assumption, since OA/OC is different at emission time and after atmospheric processing. Since all models have some aging parameterization in their calculations, this strongly suggests that the OA/OC ratio in models has to be revisited. As a general rule, models are expected to

underestimate OA/OC, since several of them use a constant value of 1.4 throughout the entire troposphere. Three models (CAM4-Oslo, OsloCTM2 and SPRINTARS) use OA/OC ratio of 2.6 for biomass burning aerosol, a value that came from measurements (Formenti et al., 2003), which is above the high-end value recently suggested in the literature for ambient aerosol (2.5; Aiken et al., 2008). Four models account for temporally and spatially variable OA/OC ratios dependent on the OA speciation in the atmosphere, but their results are completely different (Fig. 15). Measurements of OA and OC at the same location have a different seasonality, as presented later (Sect. 4.3.3) for Finokalia, Greece, which is not evident in the model results. This shows that the OA/OC ratio changes with atmospheric processing, and as applied in the model simulations (in most cases by a spatially and temporally fixed ratio), is not appropriate. A dedicated study aiming to tackle the OA/OC ratio is clearly needed.

Overall, the increased model complexity does not improve the comparison with measurements. The MNB of the urban OA comparison appears to be lower in the models that take into account the semi-volatile nature of SOA, but the correlation degrades to values as low as 0.1. The correlation of model results with remote OC data is higher for models that include semi-volatile SOA, but the difference is really small. In all other cases, no change in model skill is observed. However, the complexity is needed in models in order to distinguish between anthropogenic and natural OA and accurately

calculate the OA physical, chemical and optical properties, and their impact on climate.

4.3.2 Seasonality

Most measurements, especially at locations with at least a full year of data, are located in the USA, although recently observations have been made available from the EU-SAAR/ACTRIS observational network in Europe. Throughout the USA, where data availability is the highest, the general finding is that all models have a pronounced seasonal cycle, with minimum concentrations during winter and maximum concentrations during summer, except for the western coast, where agricultural and biomass fuel burning invert the picture, in line with previous results (Bahadur et al., 2009). This seasonal cycle is primarily caused by the presence of SOA, whose chemical production maximizes during summer, due to both elevated precursor emissions and enhanced photochemistry. Biomass burning also contributes to this summertime increase, although some models simulate excessively high monthly mean OA concentrations that can exceed $200 \mu\text{g m}^{-3}$, due to biomass burning emissions.

Although a global model is not the best tool to study urban aerosol levels, useful results can be extracted by collective comparison of OC measurements with model results. In the western states of the USA, as well as in Alaska and Florida, the typical observed urban OC seasonality presents maximum concentrations during winter and minimum during summer. This would have been expected for primary anthropogenic material due to, e.g., enhanced residential emissions from heating during winter, as well as due to enhanced agricultural and biofuel burning during winter on the west coast of the USA, seasonal patterns currently absent from most emission inventories. However, the observed seasonality is opposite of what the models calculate, which compute an OA maximum during summer, following biogenic SOA formation (Fig. 16a). In the southeast, the typical urban measured pattern does not present a pronounced seasonal cycle, with most urban locations showing a fairly flat or noisy seasonality in observed OA with no unique pattern (Fig. 16b). In most other urban cases in USA, either there is no clear seasonal pattern, or the two cases described earlier are repeated, with one unique characteristic: a peak during summer, which distorts the seasonality described above (Fig. 16c, d). Thus, the combined model–measurements analysis, given the limitations global models have when compared against urban data, suggests the existence of increased OA levels during summer due to biogenic SOA formation over large areas of the USA. This summertime OA can be of the same order of magnitude as the anthropogenic OA, even inside cities. The absolute OC values are generally still underestimated, especially during winter.

The reason why this is not the case in the western states, Alaska and Florida, might be that these areas have a strong marine influence, with air masses that do not have very aged




Figure 16. Typical seasonal distribution of OC measurements and comparison with model results for urban stations. Stars show the monthly mean of all measurements from all years that data are available, error bars present the standard deviation of the averaged measurements per month, and lines show model results, colored as in the previous figures. The grey bars show the number of measurements per month. The stations used are Arizona (a; 112.1° W, 33.5° N, years 2000–2007); Georgia (b; 83.64° W, 32.78° N, years 2001–2008); Colorado (c; 104.83° W, 38.83° N, years 2002–2006); Ohio (d; 81.68° W, 41.49° N, years 2001–2003 and 2005–2007).

SOA. For Alaska, due to its location at very high latitudes, even during summer photochemistry is less intense than at mid-latitudes, resulting in lower SOA formation rates. On the other hand, it is not clear why the OA observations in the southeastern USA do not show a peak during summer; this area is well known for its strong SOA formation potential (Carlton et al., 2010), due to both vicinity of sources and abundance in solar radiation, especially during summer. One explanation might be that wintertime emissions are much stronger there than in other areas in USA, enhancing the wintertime OA levels and masking the summertime SOA contribution. Additionally, enhanced anthropogenic aerosols like sulfate might increase aerosol water content substantially in the southeast USA (Dick et al., 2000), counterbalancing the photochemical production of SOA, an effect currently absent from all models participating in this study that do not take into account aqueous SOA formation. All these hypotheses need to be investigated in the future by both field and modeling studies in more detail.

The absence of seasonality measured at several urban locations might be due to a combination of stronger anthropogenic primary sources and reduced dispersion during winter and enhanced SOA formation during summer, as well as missing processes from the models, flattening the seasonal cycle. The missing processes include the intermediate-volatility organic compounds, which are expected to condense more during winter, and the assumption of semi-volatile POA, which will favor POA evaporation during summer. The combination of these two processes will lead to




Figure 17. Same as in Fig. 16, for remote stations. Arizona (**a**): 114.07° W, 36.02° N, years 2000–2006); Georgia (**b**): 82.13° W, 30.74° N, years 1993–2006); Colorado (**c**): 107.80° W, 37.66° N, years 2000–2006); Ohio (**d**): 81.34° W, 39.94° N, years 1998–2004).

higher OA concentrations during winter and lower during summer when compared with the current OA parameterizations. This is expected to vary spatially, depending on the availability of these species and that of preexisting aerosols, and assuming no seasonality in their sources. Whether SOA dominates over anthropogenic POA, appears to be the decisive factor for the seasonal pattern. However, this is only a hypothesis that is driven by the model results, that needs to be explored in the field. The fact that the models appear to be (a) missing an urban source, and (b) underestimating the pollution levels in cities, is also supported by the comparison of the model results with remote stations close to the urban ones presented in Fig. 16, where the models are able to capture both the magnitude and seasonality of measurements much better (Fig. 17). An important thing to note is that the measurements are roughly a factor of 5 lower in these remote stations compared to their urban counterparts, except the case of Ohio, where the remote station appears to be influenced by urban pollution: its levels are only half that of the Ohio urban station, while its seasonality resembles the seasonality present in several urban stations discussed earlier.

4.3.3 Chemical composition

Unfortunately, it is impractical to present and analyze every individual station used in the present study. Instead, a number of stations have been selected, based on a number of criteria: they must be far enough away from each other geographically, have enough data to capture both their seasonality and, where present, their interannual variability, and/or be potentially interesting for any other reason if none of the other criteria are met. Only one station has a full year of AMS data (Welgegund, South Africa, using an ACSM for real-time aerosol composition data), and only one station has both OC

and more than a couple of months of AMS data (Finokalia, Greece).

The stations that are analyzed here are the remote stations Finokalia (Greece), Welgegund (South Africa), Alaska (USA), and Manaus (Brazil), as well as the marine station Amsterdam Island (southern Indian Ocean). For clarity, only a few models are presented in the following discussion and in the figures. The remaining models (which have at least both tPOC and trSOC tracers submitted) are presented in the Supplementary Material. In addition, a number of other interesting stations are discussed in the Supplement: the urban and remote Colorado US stations discussed in Sect. 4.3.2, the remote stations LinAn (China), Alta Floresta (Brazil), Melpitz (Germany) and Mace Head (Ireland), and the marine station Okinawa (Japan).

The remote station Finokalia, Greece, has both OC and OA (AMS) measurements. The OC data (Fig. 18) do not exhibit any seasonality, in contrast to all models that underestimate the wintertime measurements by simulating a wintertime minimum and a summertime maximum. The measured OA concentrations (Fig. 19), although from only four out of twelve months, appear to be higher during summer, a feature that is captured both in shape and magnitude by a small number of models. The air masses that arrive at Finokalia are aged, since there are no significant sources upwind for at least 300 km (Mihalopoulos et al., 1997). This is also evident from the GISS-CMU-VBS results, where virtually all POA is calculated to be ntrSOA (aged primary), which means that photochemistry, which is expected to be higher during summer, has already contributed to the aging of the air masses arriving at the station. If this is indeed the case, it means that the OA/OC ratio during summer is higher than the winter value, a fact that is implied by the measurements. Note however that it is not trivial to compare the $PM_{1.8}$ OC data with the PM_1 AMS data and calculate an OA/OC ratio (Koulouri et al., 2008); it is also not straightforward to calculate OA/OC from O/C that the AMS provides, without introducing an additional level of uncertainty, due to the small, but not negligible, contribution of other heteroatoms like N, S, and P in OA. In any case, the fact that OA/OC appears to be changing with seasons is something that has to be taken into account by models that use a constant OA/OC ratio in their calculations. The evaluation of OA/OC will be studied in detail in the future; as a first estimation, since many models calculate high SOA during summer at that station, it is anticipated that the modeled OA/OC ratio will also be higher during summer. Two of the models that include multiphase chemistry of organics (IMAGES and IMPACT) calculate a significant contribution of ntrSOA to the total OC over Finokalia.

Welgegund, South Africa, is the only station for which we have been able to obtain a full year of AMS data from Fig. 20; unfortunately, no OC measurements in our database are in the same area to perform the same analysis as in Finokalia. Welgegund is a station that is strongly affected by seasonal biomass burning, and occasional anthropogenic




Figure 18. OC seasonality as calculated by all models (a) and chemical composition in GEOS-Chem-APM (b), GISS-CMU-VBS (c), GISS-modelE-I (d), IMAGES (e) and TM4-ECPL-FNP (f) for Finokalia, Greece (remote, years 2004–2007). The coordinates in panel (a) show the location of the station, while those in (b–f) show the center of the grid box of the corresponding model. (a) is similar with those presented in Figs. 16 and 17; for the chemical composition in (b–f), brown is tPOC, green is trSOC, cyan is mPOC, blue is ntrSOC (aged tPOC for GISS-CMU-VBS, OA formed via multiphase chemistry for all other cases), and orange is MSA. The chemical composition of the remaining models that have submitted at least both tPOC and trSOC data are presented in the Supplement. Note the different scales on the y axes.




Figure 19. Same as in Fig. 18, for OA (years 2008 and 2009). The chemical composition in (b–f) (where available) is presented as defined by the AMS: HOA (grey) and OOA (purple).

pollution (Tiitta et al., 2014). Besides EMAC, which overpredicts the biomass burning seasonal maximum by a factor of more than 3, most models appear to capture both the seasonal variability and levels at that station. EMAC uses the GFED inventory, the same as ECHAM5-SALSA (which lies at the high end of the models but does not stand out) and BCC, which strongly underestimates the biomass burning peak. The reason why the OC load calculated by EMAC is so high, which is evident in comparisons with several stations that are strongly affected by biomass burning, might be the fact that EMAC puts all biomass burning emissions at the first model layer, in contrast to the other models that distribute them between many layers close to the surface. Several mod-

els simulate peak OC values during September, in line with a September–October maximum in the measurements, which can be attributed to biomass burning. Caution has to be taken for the exact interpretation of the absolute values or even the peaks in the data set, since the measurements are from the year 2011, and no model has used emissions or meteorology from that year. Since biomass burning has a strong interannual variability, either multi-year data are needed in order to construct a climatology and then compare with a model year that is not exactly the same as that of the data, or the simulations should use emission inventories and meteorology for the specific year that the measurements have been performed. There is agreement between the models that the September




Figure 20. Same as in Fig. 18, for OA at Welgegund, South Africa (remote, years 2010–2011). The chemical composition in (b–f) (where available) is presented as defined by the AMS: HOA (grey) and OOA (purple).

maximum is due to POA, while SOA is fairly constant year-round; aqueous chemistry also contributes a small amount to the total OA, which is enhanced during the wet season. GISS-CMU-VBS calculates that most of the POA is already aged, although during the biomass burning season, there is a non-negligible amount that is still fresh.

In Alaska, USA (Fig. 21), many models simulate a summer maximum, in agreement with the measurements; this is due to biomass burning sources. TM4-ECPL-FNP calculates a very strong contribution from primary biological particles to the total OC, resulting in a slight overestimation of measurements throughout the year. The four models that have provided mPOA concentrations (two GISS-modelE and two TM4-ECPL models) suggest that marine organics are present in significant quantities. Multiphase chemistry is also calculated to contribute during the summer months. ECMWF-GEMS shows a very wide peak in OC during summer, in contrast with the other models, resulting in higher concentrations than the measured ones for half of the year. This might be caused by the averaging of biomass burning emissions over six fire seasons that this model uses, which exhibit a large interannual variability and which broaden the biomass burning contribution over many months. The remaining models generally underestimate the measurements, although they capture the observed seasonality rather well; more than half of the models have a correlation coefficient against measurements greater than 0.8. An interesting pattern is that of the two GISS-modelE models, which simulate a significant contribution of trSOA to the total OC, especially during winter. These two models are the only models that include semi-volatile SOA, and use the Lathière et al. (2005) VOC emissions, in which strong summer emissions in southern Alaska are present (Tsigaridis et al., 2005). It is very likely that the distribution of VOC sources (which differs from that of the other models), when combined with the low temperatures

in Alaska during winter (which favors partitioning to the aerosol phase), leads to the enhanced trSOA formation.

As expected, only the models that include a marine source of mPOA are able to capture the OA concentrations at remote marine stations. This is particularly true for the two versions of GISS-modelE (Tsigaridis et al., 2013), which have the strongest source of mPOA of all models that participate in the intercomparison. Although most of the remote marine data we have are single measurements and their seasonality cannot be studied, it is important to note that their chemical composition is dominated by mPOA. Fortunately, there is one station with five years of data in a remote marine environment: Amsterdam Island, in the southern Indian Ocean (Fig. 23). As at Mace Head, the models that include mPOA sources are closer to the measurements, while the rest of the models simulate extremely low OC concentrations. There are three notable exceptions: one is the two GISS-modelE models, which strongly overestimate the measurements, as discussed by Tsigaridis et al. (2013). Second, the ECMWF-GEMS model, which, although it does not have a marine OA source, simulates higher-than-expected OC concentrations there. Third, the IMAGES model, which is able to capture some of the measured data due to high ntrSOA amounts calculated there. Multiphase chemistry appears to contribute significantly to the OC mass calculated at Amsterdam Island in other models as well, which reproduce the long-range transport of biomass burning aerosol from southern Africa from August to October (Fig. 23), which is also seen in the observations (Sciare et al., 2009). The meteorology used appears to affect ntrSOA production in the two TM4-ECPL models significantly, due to differences in the availability of water in aerosols and the distribution of clouds between the years simulated: 2005 for TM4-ECPL-F and 2006 for TM4-ECPL-FNP.




Figure 21. Same as in Fig. 18, for Alaska, USA (remote, years 2002–2006). For the chemical composition in (b–f), brown is tPOC, green is trSOC, cyan is mPOC, blue is ntrSOC, and orange is MSA.




Figure 22. Same as in Fig. 18, for Manaus, Brazil (remote, years 2008–2010). For the chemical composition in (b–f), brown is tPOC, green is trSOC, cyan is mPOC, blue is ntrSOC, and orange is MSA.

5 Conclusions

This study shows that the diversity of the global OA modeling results has increased since AeroCom phase I, mainly due to both the increased complexity, as well as the increased diversity of the OA parameterizations and sources in the models, which is evident in the different chemical compositions simulated by the models at the various stations analyzed here. Increased number of tracers, however, does not necessarily mean increased complexity of OA parameterizations; models with aerosol microphysics must have a large number of organic aerosol tracers, even when they may simulate OA production in a very simplistic way. At present, about half of the thirty-one participating models include explicit treatment of semi-volatile SOA formation in the atmosphere. Four models also account for multiphase chemistry and six models for natural sources of POA, in particular the marine source, with

one model including the emissions of primary biological particles.

The POA sources in the thirty-one AeroCom models range from 34 to 144 Tg a⁻¹ with a median value of 56 Tg a⁻¹. Secondary OA sources show larger model diversity spanning from 12.7 to 121 Tg a⁻¹, with a median value for the 12 out of 14 models that parameterize SOA chemical production of 51 Tg a⁻¹ (mean 59 Tg a⁻¹ with standard deviation of 38 Tg a⁻¹). In the four models that account for multiphase chemistry of organics, its contribution to SOA levels is calculated to be significant (up to 50 % of total SOA formation), at least regionally.

The wet removal of OA is simulated to range from 28 to 209 Tg a⁻¹ for 26 of the models, with median 70 Tg a⁻¹, which is on average 85 % of the total OA deposition. The high wet removal variability, together with the large variability of OA sources, is attributed primarily to the diversity of




Figure 23. Same as in Fig. 18, for Amsterdam Island, Indian Ocean (marine, years 2003–2007). For the chemical composition in (b–f), brown is tPOC, green is trSOC, cyan is mPOC, blue is ntrSOC, and orange is MSA.

SOA formation, which affects the total OA load and lifetime. The very high variability of SOA budgets between models is especially evident in the SOA lifetime (2.4 days to 15 days). This slightly increases the calculated variability of the total OA by the phase II models compared to phase I, where the SOA model diversity was essentially zero.

The treatment of aerosol microphysics in the models appears to have a significant impact on the calculated OA load and dry deposition. The range in dry deposition flux for OA ($2\text{--}36\text{ Tg a}^{-1}$ in the present study) has been greatly increased since both AeroCom ExpA and ExpB, by a factor of 2 or more, while the M7 and TOMAS aerosol microphysics parameterizations, used by three and two models, respectively, simulate very low dry deposition rates when compared to the other models and thus contribute a lot to this change in diversity.

The annual median atmospheric burden of OA is calculated to be 1.4 Tg by the AeroCom phase II models, with values that vary mostly between 0.6 Tg and 1.8 Tg . Four models simulate loadings higher than 2.0 Tg , up to 3.8 Tg . The models calculate very similar OA load seasonality, which maximizes during Northern Hemisphere summer, when both primary (biomass burning) and secondary (chemical production) OA are high and minimize during Northern Hemisphere spring. A median OA lifetime of about 5.4 days (ranging from 3.8 to 9.6 days) is derived from the present study. The median POA lifetime of 4.8 days (ranging from 2.7 to 7.6 days) from this study is slightly shorter than the median SOA lifetime of 6.1 days (range from 2.4 to 14.8 days).

For many models that reported both OA and SO_4^{2-} loads, the OA load is calculated to be lower than that of SO_4^{2-} , with a median value of the $\text{OA}/\text{SO}_4^{2-}$ mass load ratio of 0.77. Simulated values of this ratio span from 0.25 to 2.0, with 9 models having a value greater than 1, indicating that there is a low level of understanding of the relative importance of

OA and SO_4^{2-} aerosol components between models, although modeling studies indicate that this ratio will increase in the future due to sulfur emission controls. This ratio is also affected by multiphase chemistry of organics and deserves further attention in the future.

A significant (up to 45 %) but highly variable contribution of multiphase chemistry to global SOA formation is calculated by models that account for this process. The comparison with observations indicates that the lower estimate of this source might be closer to reality, but this has to be revisited when more models will include multiphase SOA formation. In addition, a gas-phase source of SOA, either new or an enhanced pre-existing one, has the potential to improve the comparison with measurements in the same way multiphase chemistry does; OA chemical composition measurements can help identify which one of the two, or both, is the case. Further investigation of the importance of multiphase chemistry on the global scale and evaluation against targeted observations and field campaigns is needed.

The models show a large diversity (about two orders of magnitude) in the free troposphere, pointing to uncertainties in the temperature-dependent partitioning of SOA, uncertainties in free tropospheric sources, and the impact of meteorology and transport. A systematic comparison of model results with the limited available free tropospheric observations would give important insights into the large model differences in the middle and upper troposphere.

Despite the increasing diversity between models since AeroCom phase I experiments, the models are now able to simulate the secondary nature of OA observed in the atmosphere as a result of SOA formation and POA aging, although the absolute amount of OA present in the atmosphere remains underestimated. The median MNB of all models against urban measurements at the surface is calculated to be -0.62 for OC and -0.51 for OA and with correlations 0.47 and 0.54, respectively, while for remote surface measurements

the MNB is -0.15 for OC and $+0.51$ for OA with correlations of 0.39 and 0.37, respectively. Due to the different locations, number of stations and measurement times where OA and OC data are available, a direct comparison between the OC and OA statistics results is not straightforward, and should be avoided.

Comparison of model results with OA and OC, where available, shows that the models capture the submicron OA mass better than the $PM_{2.5}$ OC mass near the surface. Although this indicates a possible overestimate of the OA/OC ratio by the models, this is not necessarily the case, since virtually all OC and OA measurements were taken at different locations and different times. Most models use a constant value of 1.4, and only four models in this study calculate it prognostically. The limited number of observations that can be used to derive the OA/OC ratio indicate dependence on sources, atmospheric conditions and season; this will be revisited in a future study.

The flat seasonality measured at several urban locations is not reproduced by the models. The comparisons indicate a missing or underestimated source of OA in the models, either anthropogenic primary (for instance domestic wood burning), or secondary, primarily during winter. Improvements in the seasonality and strength of the anthropogenic POA sources in models can reduce the differences between model results and observations, but not eliminate them, since most global models cannot resolve urban pollution due to their large grid size.

6 Future directions

Available OC and OA observations and thus model evaluations are concentrated in the USA and Europe, but additional long-term observations from tropical, boreal, Southern Hemisphere and remote marine regions also from the free troposphere are needed to complement the global OA observational database.

Natural POA sources are important components of the OA global budget; however, among the thirty-one models participating in this intercomparison, only six account for the marine source of OA and one for the primary biogenic particles. Comparison of model results to observations over remote marine locations can provide constraints on our understanding of the marine POA source. The statistics on model performance calculated here are not able to quantify the importance or the understanding of this source because seasonal data from remote marine locations are limited. The magnitude of the marine source and the properties of marine OA remain highly uncertain and are an active area of research.

Primary biogenic particles can also be significant contributors to OA, particularly over land, but are taken into account only in one model. While the parameterization of the primary biogenic source of OA is extremely uncertain, model comparison with measurements is improved when accounting for

this source in that model, by reducing the MNB. The correlation of the model results with observations does not change significantly when including or excluding this source. However, station-by-station comparison indicates a low level of understanding of the spatial and seasonal variability of this natural source, which deserves further investigation and improvement.

Both the model diversity that increased with increasing model complexity over the past decade, as well as the comparison of model results with station data, reveal important gaps in our understanding of OA concentrations, sources and sinks in the atmosphere, and point towards the need for better understanding of sources and chemical aging of OA. Although the increasing complexity did not significantly improve the model performance, model complexity is imposed by the need to provide information for future developments that will help quantify the anthropogenic impact to climate via the aerosol direct and indirect effects. The existence of significant secondary sources of OA that are enhanced by interactions of natural with anthropogenic emissions remains an open question that cannot be answered by a simple OA parameterization. Furthermore, the OA impact on climate depends on the OA physical, chemical and optical properties, as well as the OA distribution in the atmosphere, which is affected by continuous evaporation/condensation processes of semi-volatile organic material and consequent change of hygroscopicity.

In this respect, new information from dedicated field campaigns that either occurred over the past few years or are planned to take place soon, is expected to shed light on the OA formation processes and how these are altered in the presence of anthropogenic pollution. The model development related to OA is expected to accelerate in the near future and must be performed in parallel with extensive model evaluation. Important processes currently not included in many models that need to receive high priority from modeling groups include the semi-volatile nature of OA, the temperature-dependent OA formation and aging, which affects their volatility, and an improved parameterization of the OA/OC ratio. Improved laboratory measurements of SOA formation are also crucial for the model improvements (Zhang et al., 2014).

Table 1. Organic aerosol representation in the models.

Model	OA types ¹	No. of OA tracers	trSOA precursors	trSOA calculations	OA/OC	Comments
BCC	tPOA	12	Monoterpenes	15 % yield from terpene emissions (Dentener et al., 2006); included in tPOA.	1.4	
CAM4-Oslo	tPOA ^{2,3}	3	Monoterpenes	37.5 Tg a ⁻¹ from terpene emissions based on the Dentener et al. (2006) distribution; included in tPOA.	1.4 for fossil and biofuel burning and 2.6 for biomass burning	
CAM5-MAM3	tPOA, trSOA	3 ⁴	Isoprene, terpenes, aromatics, higher molecular weight alkanes and alkenes	Prescribed mass yields for the 5 trSOA precursor categories (6.0, 37.5, 22.5, 7.5, and 7.5 %, respectively) that form a single semi-volatile species that then kinetically but reversibly partitions to the OA phase.	1.4	Precursor VOCs are lumped species from MOZART. Yields listed include a 1.5 times increase to reduce anthropogenic aerosol indirect forcing. The single semi-volatile gas has a saturation mixing ratio of 0.1 ppbv at 298 K. Includes aerosol microphysics (MAM3; modal).
CanAM-PAM	tPOA ³	3	Monoterpenes	15 % yield from terpene emissions (Dentener et al., 2006); included in tPOA.	1.4	1 tracer in 3 size classes, internally mixed with BC and ammonium sulfate (2 moments).
CCSM4-Chem	tPOA, trSOA	7	Isoprene, monoterpenes, toluene, benzene, xylene	Two-product model	1.4	Isoprene+OH uses the high-NO _x pathway.
ECHAM5-HAM2	tPOA, trSOA	24	Isoprene, monoterpenes, benzene, toluene, xylene	Two-product model	1.4	Includes aerosol microphysics (M7; modal).
ECHAM5-HAMMOZ	tPOA ³	4	Monoterpenes	15 % yield from terpene emissions (Dentener et al., 2006); included in tPOA.	1.4	Includes aerosol microphysics (M7; modal).
ECHAM5-SALSA	tPOA ³	11	Monoterpenes	15 % yield from terpene emissions (Dentener et al., 2006); included in tPOA.	1.4	Includes aerosol microphysics (SALSA; sectional).
ECMWF-GEMS	tPOA ³	2	Monoterpenes	15 % yield from terpene emissions (Dentener et al., 2006); included in tPOA.	1.4	
EMAC	tPOA ³	2	Monoterpenes	15 % yield from terpene emissions (Dentener et al., 2006); included in tPOA.	1.4	Includes aerosol microphysics (GMXe, based on M7; sectional).
GEOS-Chem	tPOA, trSOA	5	Isoprene, monoterpenes, sesquiterpenes	Two-product model	2.1	
GEOS-Chem-APM	tPOA, trSOA, MSA	24	Isoprene, monoterpenes, limonene, sesquiterpenes, alcohols, benzene, toluene, xylene	Two-product model + aging/condensation	2.1	Considers the volatility changes of the gaseous semi-volatile compounds arising from the oxidation aging process, as well as the kinetic condensation of low-volatility gases; includes aerosol microphysics (bins).
GISS-CMU-TOMAS	tPOA ³	24	Terpenes	A generic SOA precursor (Dentener et al., 2006) representing all SOA precursor gases is emitted and forms non-volatile SOA (included in tPOA) with a chemical lifetime of 12 h.	1.8	Includes aerosol microphysics (sectional).
GISS-CMU-VBS	tPOA, trSOA, ntr-SOM	26	Isoprene, monoterpenes, sesquiterpenes, alkanes, alkenes and aromatics (VOCs with C* < 10 ⁶ μg m ⁻³)	Volatility-basis set	1.8	tPOA is treated as semi-volatile and reactive. ntrSOA is formed from the gas-phase oxidation of tPOA.
GISS-MATRIX	tPOA ³	3	Monoterpenes	10 % yield from monoterpene emissions (Lathière et al., 2005); included in tPOA.	1.4	Includes aerosol microphysics (moments).

Table 1. Continued.

Model	OA types ¹	No. of OA tracers	trSOA precursors	trSOA calculations	OA/OC	Comments
GISS-modelE-G	tPOA, mPOA, trSOA, MSA	9	Isoprene, monoterpenes, sesquiterpenes	Two-product model	1.4	
GISS-modelE-I	tPOA, mPOA, trSOA, MSA	9	Isoprene, monoterpenes, sesquiterpenes	Two-product model	1.4	
GISS-TOMAS	tPOA ³	24	Monoterpenes	A generic SOA precursor (Lathi�re et al., 2005) representing all SOA precursor gases is emitted and forms non-volatile SOA (included in tPOA) with a chemical lifetime of 12 h.	1.4	Includes aerosol microphysics (sectional).
GLMAPbin	tPOA ³	40	Monoterpenes	α -pinene + all oxidants \rightarrow 13 % non-volatile SOA (included in tPOA).	1.4	Includes aerosol microphysics (bin).
GLMAPmode	tPOA ³	5	Monoterpenes	α -pinene + all oxidants \rightarrow 13 % non-volatile SOA (included in tPOA).	1.4	Includes aerosol microphysics (modal).
GMI	tPOA ³	3	Monoterpenes	10 % yield from monoterpene emissions (GEIA); included in tPOA.	1.4	
GOCART	tPOA ³	2	Monoterpenes	10 % yield from monoterpene emissions (GEIA); included in tPOA.	1.4	50 % of anthropogenic and biomass burning OC is emitted as hydrophobic and 50 % as hydrophilic (Cooke et al., 1999); hydrophobic OC becomes hydrophilic in an e-folding time of 2.5 days.
HadGEM2-ES	tPOA, trSOA	3	Terpenes	Fixed 3-D monthly climatology obtained from STOCHEM (Derwent et al., 2003)	1.4	3 tracers for fossil fuel organic carbon aerosols (fresh, aged, dissolved in cloud water).
IMAGES	tPOA, trSOA, ntrSOA	26	Isoprene, α -pinene, sesquiterpenes, benzene, toluene, xylene	Two-product model	Varying	trSOA includes the effect of water uptake on partitioning. ntrSOA is glyoxal and methylglyoxal from cloud chemistry and aqueous aerosol processing.
IMPACT	tPOA ⁵ , trSOA, ntrSOA	33	Isoprene, monoterpenes, aromatics	SOA comes from organic nitrates and peroxides using the traditional gas-particle partitioning with an explicit full chemistry. The condensed SOA is further assumed to form oligomers with a 1 day e-folding time.	Varying	ntrSOA from the uptake of gas-phase glyoxal and methylglyoxal onto clouds and aqueous sulfate aerosol (Fu et al., 2008, 2009) and uptake of gas-phase epoxides onto aqueous sulfate aerosol (Paulot et al., 2009).
LMDz-INCA	tPOA ³ , MSA	3	Monoterpenes	15 % yield from terpene emissions (Dentener et al., 2006); included in tPOA.	1.4	
OsloCTM2	tPOA, trSOA	62	Isoprene, 5 classes of terpenoid compounds (Griffin et al., 1999b), 2 classes of aromatics	Two-product model	1.6 for fossil and biofuel burning and 2.6 for biomass burning	
SPRINTARS	tPOA, trSOA	2	Monoterpenes ⁶	9.2 % yield of non-volatile trSOM (Griffin et al., 1999a; Griffin et al., 1999b) from monoterpene emissions (GEIA) ⁶	1.6 for fossil fuel and biofuel, 2.6 for other	
TM4-ECPL-F	tPOA, mPOA, trSOA, ntrSOA, MSA	22	Isoprene, monoterpenes, sesquiterpenes, aromatics	Two-product model	Varying	ntrSOA is oxalic acid, glyoxilic acid and glyoxal oligomers from cloud chemistry.
TM4-ECPL-FNP	tPOA, mPOA, trSOA, ntrSOA, MSA	24	Isoprene, monoterpenes, sesquiterpenes, aromatics	Two-product model	Varying	tPOA includes primary biogenic particles and organics associated with soil dust; ntrSOA is oxalic acid, glyoxilic acid and glyoxal oligomers from cloud chemistry.
TM5	tPOA ³ , MSA	4	Monoterpenes	15 % yield from terpene emissions (Dentener et al., 2006); included in tPOA.	1.4	Includes aerosol microphysics (M7; modal).

¹: tPOA: terrestrial primary organic aerosol mass; mPOA: marine primary organic aerosol mass; trSOA: traditional secondary organic aerosol mass; ntrSOA: non-traditional secondary organic aerosol mass; MSA: methane sulfonic acid.

²: tPOA also includes mPOA and MSA.

³: tPOA also includes trSOA.

⁴: tPOA in accumulation mode; trSOA in accumulation and Aitken modes. Aitken mode mass is minor.

⁵: tPOA also includes mPOA.

⁶: SPRINTARS also has a two-product model configuration (not presented here), with trSOA coming from isoprene and α -pinene.

Table 2. Summary of organic aerosol processes taken into account by the models.

Model	SOA like tPOA	mPOA	Simple SOA, irreversible partitioning	Reversible partitioning (equilibrium)	Includes ntrSOA	MSA	Microphys. aging
BCC	X						
CAM4-Oslo	X	In tPOA				In tPOA	X
CAM5-MAM3				Kinetically			X
CanAM-PAM	X						
CCSM4-Chem				X			
ECHAM5-HAM2				X			X
ECHAM5-HAMMOZ	X						X
ECHAM5-SALSA	X						X
ECMWF-GEMS	X						
EMAC	X						X
GEOS-Chem				X			
GEOS-Chem-APM				X		X	X
GISS-CMU-TOMAS			X				X
GISS-CMU-VBS				X	VBS		
GISS-MATRIX	X						X
GISS-modelE-G		X		X		X	
GISS-modelE-I		X		X		X	
GISS-TOMAS			X				X
GLOMAPbin			X				X
GLOMAPmode			X				X
GMI	X						
GOCART	X						
HadGEM2-ES				Offline			
IMAGES				X	Aqueous		
IMPACT		In tPOA		X	Aqueous	In tPOA	
LMDz-INCA	X					X	
OsloCTM2				X			
SPRINTARS	X						
TM4-ECPL-F		X		X	Aqueous	X	Chemical
TM4-ECPL-FNP		X		X	Aqueous	X	Chemical
TM5	X					X	X

Table 3. Host model description and year of simulation. If multiple years were simulated, the data used for the present study are those from the year 2006, if available; otherwise, the year closest to 2006 was selected.

Model (AeroCom experiment name)	Simulated year(s)	Horizontal resolution (lat./lon.)	Vertical resolution	Meteorology	Model references
BCC (BCC_AGCM2.0.1_CAM.A2.HCA-FIX)	2000 ¹	2.8125 × 2.8125	26 (hybrid sigma) to 2.9 hPa	Online	Zhang et al. (2012a)
CAM4-Oslo (CAM4-Oslo-Vemip5.A2.CTRL)	2006 ²	1.875 × 2.5	26 (hybrid sigma) to 2.19 hPa	Online ²	Kirkevåg et al. (2013)
CAM5-MAM3 (CAM5.1-MAM3-PNNL.A2.CTRL)	2000 ¹	1.875 × 2.5	30 (hybrid sigma) to 2.3 hPa	Online	Liu et al. (2012)
CanAM-PAM (CCCma.A2.CTRL)	2006	3.75 × 3.71	35 (hybrid sigma) to 1 hPa	Online	von Salzen et al. (2005); von Salzen (2006)
CCSM4-Chem (CCSM4)	2006	1.9 × 2.5	26 (sigma) to 3.5 hPa	GEOS 5	Heald et al. (2008); Lamarque et al. (2012)
ECHAM5-HAM2 (MPIHAM_V2_KZ.A2.CTRL)	2006–2008	1.875 × 1.875	31 (hybrid sigma) to 10 hPa	ERA	Stier et al. (2005); Kazil et al. (2010); O'Donnell et al. (2011); K. Zhang et al. (2012)
ECHAM5-HAMMOZ (ECHAM5-HAMMOZ.A2.HCA-0)	2000–2005	2.8125 × 2.8125	31 (hybrid sigma) to 10 hPa	ECMWF ERA40 and operational	Stier et al. (2005); Pozzoli et al. (2008, 2011)
ECHAM5-SALSA (SALSA_V1_TB.A2.CTRL)	2006	1.875 × 1.875	31 (hybrid sigma) to 10 hPa	ECMWF operational	Stier et al. (2005); Kokkola et al. (2008); Bergman et al. (2012)
ECMWF-GEMS (n/a)	2003–2008	1.5 × 1.5 ³	60 (hybrid sigma) to 0.1 hPa	ECMWF-GEMS operational	Benedetti et al. (2009); Morcrette et al. (2009)
EMAC (ECHAM-MESSy-GMXe.A2.CTRL)	2006	2.8125 × 2.8125	19 (hybrid) to 10 hPa	ECMWF reanalysis	Jockel et al. (2005); Pringle et al. (2010)
GEOS-Chem (GEOSCHEM-v822.AEROCOM_A2.CTRL)	2006	2 × 2.5	47 (hybrid sigma) to 0.01 hPa	GMAO version 5 (GEOS-5)	Bey et al. (2001); Park et al. (2003); Henze and Seinfeld (2006); Liao et al. (2007)
GEOS-Chem-APM (GEOS-Chem-APM.A2.CTRL)	2006	2 × 2.5	47 (hybrid sigma) to 0.01 hPa	GEOS-5.2.0	Bey et al. (2001); Park et al. (2003); Yu and Luo (2009); Yu (2011)
GISS-CMU-TOMAS (GISS-TOMAS.A2.CTRL)	2006 ¹	4 × 5	9 (hybrid sigma) to 10 hPa	Online	Hansen et al. (1983); Adams and Seinfeld (2002); Lee and Adams (2010, 2012)
GISS-CMU-VBS (GISS-CMU.A2.CTRL)	2008 ¹	4 × 5	7 (hybrid sigma) to 10 hPa	Online	Hansen et al. (1983); Farina et al. (2010); Jathar et al. (2011)
GISS-MATRIX (GISS-MATRIX.A2.CTRL)	2006–2008 ⁴	2 × 2.5	40 (sigma) to 0.1 hPa	NCEP reanalysis and online	Bauer et al. (2008)
GISS-modelE-G (GISS-modelE.A2.CTRL)	2000–2008 ⁴	2 × 2.5	40 (sigma) to 0.1 hPa	NCEP reanalysis and online	Koch et al. (2007); Tsigaridis and Kanakidou (2007); Tsigaridis et al. (2013)
GISS-modelE-I (GISS-modelE.A2.HCA-IPCC)	2000–2008 ⁴	2 × 2.5	40 (sigma) to 0.1 hPa	NCEP reanalysis and online	Koch et al. (2007); Tsigaridis and Kanakidou (2007); Tsigaridis et al. (2013)
GISS-TOMAS (n/a)	2006 ⁵	2 × 2.5	40 (sigma) to 0.1 hPa	MERRA reanalysis and online	Lee and Adams (2010); Lee et al. (2014)
GLOMAPbin (GLOMAPbin1pt1.A2.CTRL)	2006	2.8125 × 2.8125	31 (hybrid sigma) to 10 hPa	ECMWF operational	Mann et al. (2012)
GLOMAPmode (GLOMAPmodev6R.A2.CTRL)	2006	2.8125 × 2.8125	31 (hybrid sigma) to 10 hPa	ECMWF operational	Mann et al. (2012)

Table 3. Continued.

Model (AeroCom experiment name)	Simulated year(s)	Horizontal resolution (lat/lon)	Vertical resolution	Meteorology	Model references
GMI (GMI-v3.A2.CTRL)	2006	2 × 2.5	42 (hybrid sigma) to 0.01 hPa	NASA GMAO GEOS4	Liu et al. (2007); Bian et al. (2009)
GOCART (GOCART-v4.A2.HCA-0)	2000–2007	2 × 2.5	30 (hybrid sigma) to 0.01 hPa	NASA GMAO GEOS4	Chin et al. (2000); Ginoux et al. (2001); Chin et al. (2002)
HadGEM2-ES (HadGEM2-ES.A2.CTRL)	2006–2008	1.25 × 1.875	38 (hybrid height) to 39 km	ECMWF operational and online	Bellouin et al. (2011), and references therein
IMAGES (n/a)	2006	2 × 2.5	40 (hybrid) to 44 hPa	ECMWF ERA-Interim	Müller (2009); Stavrakou et al. (2009); Ceulemans et al. (2012)
IMPACT (IMPACT-C.A2)	1997	4 × 5	46 (hybrid sigma) to 0.147 hPa	NASA DAO GEOS-STRAT	Lin et al. (2012)
LMDz-INCA (LSCE-vRV.A2.CTRL)	2006	1.875 × 3.75	19 (sigma) to 3 hPa	ECMWF IMF and online	Schulz (2007); Balkanski (2011); Szopa et al. (2013)
OsloCTM2 (OsloCTM2-v2.A2.CTRL)	2006	2.8125 × 2.8125	60 (hybrid sigma) to 2 hPa	ECMWF IFS	Hoyle et al. (2007, 2009); Myhre et al. (2009)
SPRINTARS (SPRINTARS-v384.A2.CTRL)	2006	1.125 × 1.125	56 (sigma) to ~ 1 hPa	NCEP reanalysis and online	Takemura et al. (2000, 2002, 2005, 2009)
TM4-ECPL-F (TM4-ECPL-F.A2.CTRL)	2006	2 × 3	34 (hybrid sigma) to 0.1 hPa	ECMWF ERA-Interim	Myriokefalitakis et al. (2008); Myriokefalitakis et al. (2010); Myriokefalitakis et al. (2011)
TM4-ECPL-FNP (TM4-ECPL-FNP.A2.CTRL)	2005	2 × 3	34 (hybrid sigma) to 0.1 hPa	ECMWF ERA-Interim	Myriokefalitakis et al. (2008, 2010, 2011); Kanakidou et al. (2012)
TM5 (TM5-V3.A2.HCA-IPCC)	2000–2009	2 × 3	34 (hybrid sigma) to 0.5 hPa	ECMWF ERA-Interim	Huijnen et al. (2010); Aan de Brugh et al. (2011); van Noije et al. (2014)

¹ Meteorology calculated by the model's climate.² 5 year mean of model's calculated meteorology, driven by offline CAM4 aerosols and cloud droplet number concentration.³ The model is run at a TL159 L60 resolution, meaning a reduced physical grid of 1.125 × 1.125. The data extraction was carried out on a 1.5 × 1.5 regular grid.⁴ Horizontal winds are nudged to NCEP reanalysis (Kalnay et al., 1996), with the rest of the climate parameters being calculated online.⁵ Horizontal winds are nudged to MERRA reanalysis (Rienecker et al., 2011), with the rest of the climate parameters being calculated online.

Table 4. Primary organic aerosol emissions adopted by the models.

Model	Fossil fuel and biofuel	Year	Biomass burning	Year	Other sources/comments
BCC	Bond et al. (2004)	2000	GFED	2000	
CAM4-Oslo	AeroCom; mPOA based on Spracklen et al. (2008).	2006	AeroCom	2006	Emitted particle sizes (with some adjustments) and fire emission injection heights from Dentener et al. (2006).
CAM5-MAM3	CMIP5	2000	CMIP5	2000	Emitted particle sizes (with some adjustments) and fire emission injection heights from Dentener et al. (2006).
CanAM-PAM	AeroCom	2006	AeroCom	2006	
CCSM4-Chem	POET, REAS over Asia	2006	GFED2	2006	
ECHAM5-HAM2	AeroCom	2006	AeroCom	2006	
ECHAM5-HAMMOZ	AeroCom	2000–2005	GFED2	2000–2005	For 1980–2000 (not studied here), the model uses RETRO emissions.
ECHAM5-SALSA	Dentener et al. (2006)	2000	GFED	2000	
ECMWF-GEMS	Dentener et al. (2006)	2000	GFED2	2003–2008	Biofuel emissions have a prescribed diurnal cycle.
EMAC	Dentener et al. (2006)	2000	GFED	2000	
GEOS-Chem	Bond et al. (2007)	2000	GFED2	2006	
GEOS-Chem-APM	Bond et al. (2007)	2000	GFED2	2006	
GISS-CMU-TOMAS	AeroCom	2006	AeroCom	2006	
GISS-CMU-VBS	Bond et al. (2004)*	2000	GFED2	2005	
GISS-MATRIX	CMIP5 RCP4.5	2000	GFED3	2006–2008	
GISS-modelE-G	CMIP5 RCP4.5	2000–2008	GFED3	2000–2008	mPOA calculated online. (Tsigaridis et al., 2013)
GISS-modelE-I	CMIP5 RCP4.5	2000–2008	CMIP5 RCP4.5	2000–2008	mPOA calculated online. (Tsigaridis et al., 2013)
GISS-TOMAS	CMIP5 RCP4.5	2006	CMIP5 RCP4.5	2006	
GLOMAPbin	AeroCom	2000	GFED2	climatology (Dentener et al., 2006)	
GLOMAPmode	AeroCom	2000	GFED2	climatology (Dentener et al., 2006)	
GMI	CMIP5	2000	CMIP5	2000	
GOCART	AeroCom	2006	AeroCom	2006	Details about emissions from different sources are in Chin et al. (2009).
HadGEM2-ES	CMIP5	2000	CMIP5	2000	
IMAGES	Bond et al. (2004)	2000	GFED2	2000	
IMPACT	Ito and Penner (2005)	2000	Ito and Penner (2005)	2000	mPOA calculated online, based on Gantt et al. (2009b). Fossil fuel adjusted as in Wang et al. (2009)
LMDz-INCA	CMIP5	2000	CMIP5	2000	
OsloCTM2	CMIP5	2000	CMIP5	2000	
SPRINTARS	AeroCom	2006	AeroCom	2006	
TM4-ECPL-F	CIRCE	2006	GFED2	2006	mPOA calculated online (Myriokefalitakis et al., 2010)
TM4-ECPL-FNP	CMIP5	2005	CMIP5	2005	mPOA calculated online (Myriokefalitakis et al., 2010); POA from primary biogenic sources are scaled on leaf area index and dust distribution (Kanakidou et al., 2012).
TM5	CMIP5 RCP4.5	2000–2009	CMIP5 RCP4.5	2000–2009	

North America emissions come from Park et al. (2003).

Table 5. SOA precursor emissions adopted by the models. Models that do not calculate semi-volatile SOA have been omitted.

Model	Isoprene	Year	Terpenes	Year	Aromatics	Year	Other	Year
BCC			GEIA	1990				
CAM5-MAM3	POET	2000	POET	2000	POET	2000	POET	2000
CCSM4-Chem	MEGAN 2.1	2006	MEGAN 2.1	2006	POET, with REAS over Asia	2006		
ECHAM5-HAM2	MEGAN	2006	MEGAN	2006	EDGAR v3.2 (Olivier et al., 2001)	2000		
GEOS-Chem	MEGAN 2.04	2006	MEGAN 2.04	2006				
GEOS-Chem-APM	MEGAN 2	2006	MEGAN 2	2006	EDGAR v2	1985	Online (MEGAN 2)	2006
GISS-CMU-VBS	GEIA	1990	GEIA	1990	(Farina et al., 2010)	1999	(Farina et al., 2010)	1999
GISS-modelE-G	Online (Guenther et al., 1995)	2000–2008	Lathièrè et al. (2005)	1990				
GISS-modelE-I	Online (Guenther et al., 1995)	2000–2008	Lathièrè et al. (2005)	1990				
HadGEM2-ES			GEIA	1990				
IMAGES	MEGAN	2006	GEIA	1990	RETRO	2000		
IMPACT	Online (Guenther et al., 1995)	1997	Online (Guenther et al., 1995)	1997	(Piccot et al., 1992)	1985		
OsloCTM2	POET, scaled to 220 Tg a ⁻¹	2000	GEIA	1990	CMIP5	2000		
TM4-ECPL-F	POET	2000	POET/GEIA	2000	CIRCE	2006		
TM4-ECPL-FNP	MEGAN	2005	MEGAN	2005	CMIP5	2005		

Table 6. Enthalpies of vaporization used by the models that include semi-volatile OA.

Model	ΔH (kJ mol ⁻¹)	References
CAM5-MAM3	156	Strader et al. (1999)
CCSM4-Chem	42	Heald et al. (2008)
ECHAM5-HAM2	Isoprene: 42 Monoterpenes: 59 Aromatics: 0	Henze and Seinfeld (2006) Saathoff et al. (2009)
GEOS-Chem	42	Chung and Seinfeld (2002)
GEOS-Chem-APM	47–64	Yu (2011)
GISS-CMU-VBS	30	Farina et al. (2010)
GISS-modelE-G	Isoprene: 42 Terpenes: 72.9	Henze and Seinfeld (2006) Tsigaridis et al. (2006)
GISS-modelE-I	Isoprene: 42 Terpenes: 72.9	Henze and Seinfeld (2006) Tsigaridis et al. (2006)
HadGEM2-ES	Not reported	Derwent et al. (2003)
IMAGES	Isoprene: 42 Aromatics: 18 <i>a</i> -pinene: 25.8–153.7 Sesquiterpenes: 42	Henze and Seinfeld (2006) Offenberg et al. (2006) Capouet et al. (2008); Ceulemans et al. (2012) Offenberg et al. (2006)
IMPACT	42	Heald et al. (2008)
OsloCTM2	42	Hoyle et al. (2007)
TM4-ECPL-F	Isoprene: 42 <i>a</i> -pinene: 38 β -pinene: 40 Aromatics: 40	Henze and Seinfeld (2006) Svendby et al. (2008) Svendby et al. (2008) Svendby et al. (2008)
TM4-ECPL-FNP	Isoprene: 42 <i>a</i> -pinene: 38 β -pinene: 40 Aromatics: 40	Henze and Seinfeld (2006) Svendby et al. (2008) Svendby et al. (2008) Svendby et al. (2008)

Appendix A

Table A1. List of acronyms.

AeroCom	Aerosol Comparisons between Observations and Models. For hindcast emissions, see Diehl et al. (2012).
ACSM	Aerosol Chemical Specification Monitor, a mini-AMS (Ng et al., 2011).
AMS	Aerosol Mass Spectrometer (Jayne et al., 2000).
BVOC	Biogenic VOC.
CIRCE	Climate Change and Impact Research: the Mediterranean Environment (http://www.circeproject.eu ; Doering et al., 2009).
CMIP5	Coupled Model Intercomparison Project phase 5 (http://cmip-pcmdi.llnl.gov/cmip5). For historical emissions, see (Lamarque et al., 2010).
DMS	DiMethyl Sulfide, CH ₃ SCH ₃ .
ECMWF	European Center for Medium-Range Weather Forecasts.
GEIA	Global Emissions Inventory Activity (http://geiacenter.org). For BVOC emissions, see Guenther et al. (1995).
GFED	Global Fire Emissions Database (van der Werf et al., 2003).
GFED2	Global Fire Emissions Database, version 2 (van der Werf et al., 2006).
GFED3	Global Fire Emissions Database, version 3 (van der Werf et al., 2010).
HOA	Hydrocarbon-like OA.
IFS	Integrated Forecast System.
IMF	Isobaric mapping functions.
LAD	Least absolute deviation technique.
MEGAN	Model of Emissions of Gases and Aerosols from Nature (Guenther et al., 2006).
mPOA/mPOC	marine POA/POC.
MERRA	Modern-Era Retrospective analysis for Research and Applications.
MNB	Mean normalized bias.
MSA	Methane sulfonic acid, CH ₃ SO ₃ H.
NCEP	National Centers for Environmental Prediction.
ntrSOA/ntrSOC	non-traditional SOA/SOC. For IMAGES, IMPACT and TM4-ECPL-F/FNP this is OA produced from multiphase chemistry, while for GISS-CMU-VBS it is OA formed from the VBS gas-phase chemistry.
OA	Organic aerosol and organic aerosol mass (as appropriate)
OC	Organic carbon.
OOA:	Oxygenated OA.
ntrSOA/ntrSOC	non-traditional secondary organic aerosol mass/carbon.
POA/POC	Primary OA / OC.
POET	Present and future surface emissions of atmospheric compounds (http://accent.aero.jussieu.fr/POET.php ; Granier et al., 2003)
RETRO	REanalysis of the TROpospheric chemical composition over the past 40 years (http://retro.enes.org ; Schultz et al., 2007)
SOA	Secondary organic aerosol.
tPOA/tPOC	terrestrial POA/POC.
trSOA/trSOC	traditional SOA/SOC.
VBS	Volatility-basis set.
VOC	Volatile organic compounds.
WSOC	Water soluble organic compounds.
WIOC	Water insoluble organic compounds.

The Supplement related to this article is available online at doi:10.5194/acp-14-10845-2014-supplement.

Acknowledgements. K. Tsigaridis and S. E. Bauer were supported by NASA-MAP (NASA Award Number: NNX09AK32G); N. Daskalakis, T. Iversen, A. Kirkevåg, Ø. Seland, K. S. Carslaw, G. W. Mann and L. Pozzoli by PEGASOS (FP7-ENV-2010-265148); M. Kanakidou and S. Myriokefalitakis by ECLIPSE (FP7-ENV-2011, 282688); T. Iversen, A. Kirkevåg and Ø. Seland also by the Research Council of Norway through the EarthClim (207711/E10) and NOTUR/NorStore projects, and through the EU project ACCESS; A. Kirkevåg also received funding from the Norwegian Space Center through the PM-VRAE project; A. Benedetti, J. W. Kaiser, and J.-J. Morcrette were funded through the MACC and MACC-II projects funded by the European Commission under the EU Seventh Research Framework Programme, contract numbers 218793 and 283576, respectively; J.-F. Müller and T. Stavrakou by the Belspo projects PRODEX-A3C and BIOSOA (SD/CS/05A); T. Takemura by the Funding Program for Next Generation World-Leading Researchers of the Cabinet Office, Government of Japan (GR079); Y. H. Lee by the NASA MAP and ACPMAP programs; N. Bellouin by the Joint DECC/Defra Met Office Hadley Centre Climate Programme (Ga01101); G. Luo and F. Yu by NASA-ACMAP (NNX13AK20G) and NSF (AGS-0942106); G. Curci by the Italian Space Agency in the frame of QUITSAT and PRIMES projects; R. C. Easter, S. J. Ghan, X. Liu, R. A. Zaveri and K. Zhang by the US Department of Energy, Office of Science, Scientific Discovery through Advanced Computing (SciDAC) Program and by the Office of Science Earth System Modeling Program; J. L. Jimenez and Q. Zhang through DOE (BER/ASR) DE-FG02-11ER65293; J. L. Jimenez also through DOE (BER/ASR) DE-SC0006035 and DE-SC0006711, NOAA NA13OAR4310063, and NSF AGS-1243354. The Pacific Northwest National Laboratory is operated for DOE by Battelle Memorial Institute under Contract DE-AC06-76RLO 1830. G. Lin, J. E. Penner and S. Sillman are funded by EPA STAR program (grant no. 83337701). A. Hodzic and S. Tilmes were supported by the National Center for Atmospheric Research, which is operated by the University Corporation for Atmospheric Research on behalf of the National Science Foundation. Resources supporting this work were provided by the NASA High-End Computing (HEC) Program through the NASA Center for Climate Simulation (NCCS) at Goddard Space Flight Center. C. R. Hoyle was funded through Swiss National Science Foundation (SNSF) (grant number 200021 140663). ECHAM5-HAMMOZ simulations were supported by the Deutsches Klimarechenzentrum (DKRZ) and the Forschungszentrum Juelich. Model simulations with OsloCTM2 have received support from the Research Council of Norway through the SLAC project. The AeroCom database benefits greatly from caretaking by Jan Griesfeller and the infrastructure support from IT at the Norwegian Meteorological Institute. Q. Zhang acknowledges DE-SC0001673. We also acknowledge D. Plummer for helpful comments.

Edited by: J. H. Seinfeld

References

- Aan de Brugh, J. M. J., Schaap, M., Vignati, E., Dentener, F., Kahnert, M., Sofiev, M., Huijnen, V., and Krol, M. C.: The European aerosol budget in 2006, *Atmos. Chem. Phys.*, 11, 1117–1139, doi:10.5194/acp-11-1117-2011, 2011.
- Adams, P. J. and Seinfeld, J. H.: Predicting global aerosol size distributions in general circulation models, *J. Geophys. Res.-Atmos.*, 107, 4370, doi:10.1029/2001jd001010, 2002.
- Aiken, A. C., Decarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ. Sci. Technol.*, 42, 4478–4485, doi:10.1021/es703009q, 2008.
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 955–966, doi:10.1029/2000gb001382, 2001.
- Arneth, A., Harrison, S. P., Zechle, S., Tsigaridis, K., Menon, S., Bartlein, P. J., Feichter, J., Korhola, A., Kulmala, M., O'Donnell, D., Schurgers, G., Sorvari, S., and Vesala, T.: Terrestrial biogeochemical feedbacks in the climate system, *Nat. Geosci.*, 3, 525–532, doi:10.1038/ngeo905, 2010.
- Artaxo, P., Martins, J. V., Yamasoe, M. A., Procopio, A. S., Pauliquevis, T. M., Andreae, M. O., Guyon, P., Gatti, L. V., and Leal, A. M. C.: Physical and chemical properties of aerosols in the wet and dry seasons in Rondonia, Amazonia, *J. Geophys. Res.-Atmos.*, 107, 8081, doi:10.1029/2001JD000666, 2002.
- Bahadur, R., Habib, G., and Russell, L. M.: Climatology of PM_{2.5} organic carbon concentrations from a review of ground-based atmospheric measurements by evolved gas analysis, *Atmos. Environ.*, 43, 1591–1602, 2009.
- Balasubramanian, R., Qian, W. B., Decesari, S., Facchini, M. C., and Fuzzi, S.: Comprehensive characterization of PM_{2.5} aerosols in Singapore, *J. Geophys. Res.-Atmos.*, 108, 4523, doi:10.1029/2002JD002517, 2003.
- Balkanski, Y.: L'influence des aérosols sur le climat, Thèse d'habilitation à diriger des recherches, Université Versailles Saint Quentin, St Quentin en Yvelines, 2011.
- Bauer, S. E., Wright, D. L., Koch, D., Lewis, E. R., McGraw, R., Chang, L.-S., Schwartz, S. E., and Ruedy, R.: MATRIX (Multiconfiguration Aerosol TRacker of mIXing state): an aerosol microphysical module for global atmospheric models, *Atmos. Chem. Phys.*, 8, 6003–6035, doi:10.5194/acp-8-6003-2008, 2008.
- Bellouin, N., Rae, J., Jones, A., Johnson, C., Haywood, J., and Boucher, O.: Aerosol forcing in the Climate Model Intercomparison Project (CMIP5) simulations with HadGEM2-ES and the role of ammonium nitrate, *J. Geophys. Res.-Atmos.*, 116, D20206, doi:10.1029/2011jd016074, 2011.
- Benedetti, A., Morcrette, J. J., Boucher, O., Dethof, A., Engelen, R. J., Fisher, M., Flentje, H., Huneus, N., Jones, L., Kaiser, J. W., Kinne, S., Mangold, A., Razingger, M., Simmons, A. J., and Suttie, M.: Aerosol analysis and forecast in the European Centre for Medium-Range Weather Forecasts Integrated Forecast System: 2. Data assimilation, *J. Geophys. Res.-Atmos.*, 114, D13205, doi:10.1029/2008jd011115, 2009.

- Bergman, T., Kerminen, V.-M., Korhonen, H., Lehtinen, K. J., Makkonen, R., Arola, A., Mielonen, T., Romakkaniemi, S., Kulmala, M., and Kokkola, H.: Evaluation of the sectional aerosol microphysics module SALSA implementation in ECHAM5-HAM aerosol-climate model, *Geosci. Model Dev.*, 5, 845–868, doi:10.5194/gmd-5-845-2012, 2012.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q. B., Liu, H. G. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.-Atmos.*, 106, 23073–23095, doi:10.1029/2001jd000807, 2001.
- Bian, H., Chin, M., Rodriguez, J. M., Yu, H., Penner, J. E., and Strahan, S.: Sensitivity of aerosol optical thickness and aerosol direct radiative effect to relative humidity, *Atmos. Chem. Phys.*, 9, 2375–2386, doi:10.5194/acp-9-2375-2009, 2009.
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: An investigative review, *Aerosol Sci. Tech.*, 40, 27–67, doi:10.1080/02786820500421521, 2006.
- Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J. H., and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.-Atmos.*, 109, D14203, doi:10.1029/2003jd003697, 2004.
- Bond, T. C., Bhardwaj, E., Dong, R., Jogani, R., Jung, S. K., Roden, C., Streets, D. G., and Trautmann, N. M.: Historical emissions of black and organic carbon aerosol from energy-related combustion, 1850–2000, *Global Biogeochem. Cy.*, 21, GB2018, doi:10.1029/2006gb002840, 2007.
- Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, *Mass Spectrom. Rev.*, 26, 185–222, doi:10.1002/mas.20115, 2007.
- Capouet, M., Mueller, J. F., Ceulemans, K., Compernelle, S., Vereecken, L., and Peeters, J.: Modeling aerosol formation in alpha-pinene photo-oxidation experiments, *J. Geophys. Res.-Atmos.*, 113, D02308, doi:10.1029/2007jd008995, 2008.
- Carlton, A. G. and Turpin, B. J.: Particle partitioning potential of organic compounds is highest in the Eastern US and driven by anthropogenic water, *Atmos. Chem. Phys.*, 13, 10203–10214, doi:10.5194/acp-13-10203-2013, 2013.
- Carlton, A. G., Pinder, R. W., Bhave, P. V., and Pouliot, G. A.: To What Extent Can Biogenic SOA be Controlled?, *Environ. Sci. Technol.*, 44, 3376–3380, doi:10.1021/es903506b, 2010.
- Carslaw, K. S., Boucher, O., Spracklen, D. V., Mann, G. W., Rae, J. G. L., Woodward, S., and Kulmala, M.: A review of natural aerosol interactions and feedbacks within the Earth system, *Atmos. Chem. Phys.*, 10, 1701–1737, doi:10.5194/acp-10-1701-2010, 2010.
- Cavalli, F., Viana, M., Yttri, K. E., Genberg, J., and Putaud, J.-P.: Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: the EUSAAR protocol, *Atmos. Meas. Tech.*, 3, 79–89, doi:10.5194/amt-3-79-2010, 2010.
- Ceulemans, K., Compernelle, S., and Müller, J.-F.: Parameterising secondary organic aerosol from α -pinene using a detailed oxidation and aerosol formation model, *Atmos. Chem. Phys.*, 12, 5343–5366, doi:10.5194/acp-12-5343-2012, 2012.
- Chin, M., Rood, R. B., Lin, S. J., Müller, J. F., and Thompson, A. M.: Atmospheric sulfur cycle simulated in the global model GOCART: Model description and global properties, *J. Geophys. Res.-Atmos.*, 105, 24671–24687, doi:10.1029/2000jd900384, 2000.
- Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: Tropospheric aerosol optical thickness from the GOCART model and comparisons with satellite and Sun photometer measurements, *J. Atmos. Sci.*, 59, 461–483, doi:10.1175/1520-0469(2002)059<0461:taotft>2.0.co;2, 2002.
- Chin, M., Diehl, T., Dubovik, O., Eck, T. F., Holben, B. N., Sinyuk, A., and Streets, D. G.: Light absorption by pollution, dust, and biomass burning aerosols: a global model study and evaluation with AERONET measurements, *Ann. Geophys.*, 27, 3439–3464, doi:10.5194/angeo-27-3439-2009, 2009.
- Chipperfield, M. P.: New version of the TOMCAT/SLIMCAT offline chemical transport model: Intercomparison of stratospheric tracer experiments, *Q. J. Roy. Meteor. Soc.*, 132, 1179–1203, doi:10.1256/qj.05.51, 2006.
- Chow, J. C., Watson, J. G., Lowenthal, D. H., Solomon, P. A., Magliano, K. L., Ziman, S. D., and Richards, L. W.: PM(10) and PM(2.5) Compositions in California San Joaquin Valley, *Aerosol Sci. Tech.*, 18, 105–128, doi:10.1080/02786829308959588, 1993.
- Chung, S. H. and Seinfeld, J. H.: Global distribution and climate forcing of carbonaceous aerosols, *J. Geophys. Res.-Atmos.*, 107, 4407, doi:10.1029/2001JD001397, 2002.
- Cooke, W. F., Liousse, C., Cachier, H., and Feichter, J.: Construction of a 1 degrees x 1 degrees fossil fuel emission data set for carbonaceous aerosol and implementation and radiative impact in the ECHAM4 model, *J. Geophys. Res.-Atmos.*, 104, 22137–22162, doi:10.1029/1999jd900187, 1999.
- De Gouw, J. and Jimenez, J. L.: Organic Aerosols in the Earth's Atmosphere, *Environ. Sci. Technol.*, 43, 7614–7618, doi:10.1021/es9006004, 2009.
- Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J. J., Ito, A., Marelli, L., Penner, J. E., Putaud, J.-P., Textor, C., Schulz, M., van der Werf, G. R., and Wilson, J.: Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom, *Atmos. Chem. Phys.*, 6, 4321–4344, doi:10.5194/acp-6-4321-2006, 2006.
- Derwent, R. G., Jenkin, M. E., Johnson, C. E., and Stevenson, D. S.: The global distribution of secondary particulate matter in a 3-D Lagrangian chemistry transport model, *J. Atmos. Chem.*, 44, 57–95, 2003.
- Dick, W. D., Saxena, P., and McMurry, P. H.: Estimation of water uptake by organic compounds in submicron aerosols measured during the Southeastern Aerosol and Visibility Study, *J. Geophys. Res.-Atmos.*, 105, 1471–1479, doi:10.1029/1999jd901001, 2000.
- Diehl, T., Heil, A., Chin, M., Pan, X., Streets, D., Schultz, M., and Kinne, S.: Anthropogenic, biomass burning, and volcanic emissions of black carbon, organic carbon, and SO₂ from 1980 to 2010 for hindcast model experiments, *Atmos. Chem. Phys.*

- Discuss., 12, 24895–24954, doi:10.5194/acpd-12-24895-2012, 2012.
- Doering, U., van Aardenne, J., Monni, S., Pagliari, V., Orlandini, L., and San Martin, F.: CIRCE report D8.1.3 Update of gridded emission inventories, addition of period 1990–2005 and the years 2010, 2015, 2050, 2009.
- Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environ. Sci. Technol.*, 40, 2635–2643, doi:10.1021/es052297c, 2006.
- Donahue, N. M., Epstein, S. A., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility basis set: 1. organic-aerosol mixing thermodynamics, *Atmos. Chem. Phys.*, 11, 3303–3318, doi:10.5194/acp-11-3303-2011, 2011.
- Eatough, D. J., Eatough, N. L., Obeidi, F., Pang, Y. B., Modey, W., and Long, R.: Continuous determination of PM_{2.5} mass, including semi-volatile species, *Aerosol Sci. Tech.*, 34, 1–8, 2001.
- Eatough, D. J., Long, R. W., Modey, W. K., and Eatough, N. L.: Semi-volatile secondary organic aerosol in urban atmospheres: meeting a measurement challenge, *Atmos. Environ.*, 37, 1277–1292, 2003.
- Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmos. Chem. Phys.*, 11, 11069–11102, doi:10.5194/acp-11-11069-2011, 2011.
- Facchini, M. C., Decesari, S., Rinaldi, M., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S., Moretti, F., Tagliavini, E., Ceburnis, D., and O’Dowd, C. D.: Important Source of Marine Secondary Organic Aerosol from Biogenic Amines, *Environ. Sci. Technol.*, 42, 9116–9121, doi:10.1021/es0818385, 2008.
- Fan, X. H., Lee, P. K. H., Brook, J. R., and Mabury, S. A.: Improved measurement of seasonal and diurnal differences in the carbonaceous components of urban particulate matter using a denuder-based air sampler, *Aerosol Sci. Tec.*, 38, 63–69, 2004.
- Farina, S. C., Adams, P. J., and Pandis, S. N.: Modeling global secondary organic aerosol formation and processing with the volatility basis set: Implications for anthropogenic secondary organic aerosol, *J. Geophys. Res.-Atmos.*, 115, D09202, doi:10.1029/2009jd013046, 2010.
- Formenti, P., Elbert, W., Maenhaut, W., Haywood, J., Osborne, S., and Andreae, M. O.: Inorganic and carbonaceous aerosols during the Southern African Regional Science Initiative (SAFARI 2000) experiment: Chemical characteristics, physical properties, and emission data for smoke from African biomass burning, *J. Geophys. Res.-Atmos.*, 108, 8488, doi:10.1029/2002jd002408, 2003.
- Fu, Q., Thorsen, T. J., Su, J., Ge, J. M., and Huang, J. P.: Test of Mie-based single-scattering properties of non-spherical dust aerosols in radiative flux calculations, *J. Quant. Spectrosc. Ra.*, 110, 1640–1653, doi:10.1016/j.jqsrt.2009.03.010, 2009.
- Fu, T. M., Jacob, D. J., Wittrock, F., Burrows, J. P., Vrekousis, M., and Henze, D. K.: Global budgets of atmospheric glyoxal and methylglyoxal, and implications for formation of secondary organic aerosols, *J. Geophys. Res.-Atmos.*, 113, D15303, doi:10.1029/2007jd009505, 2008.
- Fuzzi, S., Andreae, M. O., Huebert, B. J., Kulmala, M., Bond, T. C., Boy, M., Doherty, S. J., Guenther, A., Kanakidou, M., Kawamura, K., Kerminen, V.-M., Lohmann, U., Russell, L. M., and Pöschl, U.: Critical assessment of the current state of scientific knowledge, terminology, and research needs concerning the role of organic aerosols in the atmosphere, climate, and global change, *Atmos. Chem. Phys.*, 6, 2017–2038, doi:10.5194/acp-6-2017-2006, 2006.
- Gantt, B., Meskhidze, N., and Kamykowski, D.: A new physically-based quantification of marine isoprene and primary organic aerosol emissions, *Atmos. Chem. Phys.*, 9, 4915–4927, doi:10.5194/acp-9-4915-2009, 2009a.
- Gantt, B., Meskhidze, N., and Kamykowski, D.: A new physically-based quantification of marine isoprene and primary organic aerosol emissions, *Atmos. Chem. Phys.*, 9, 4915–4927, doi:10.5194/acp-9-4915-2009, 2009b.
- Gatari, M. J. and Boman, J.: Black carbon and total carbon measurements at urban and rural sites in Kenya, East Africa, *Atmos. Environ.*, 37, 1149–1154, 2003.
- George, I. J. and Abbatt, J. P. D.: Heterogeneous oxidation of atmospheric aerosol particles by gas-phase radicals, *Nat. Chem.*, 2, 713–722, doi:10.1038/nchem.806, 2010.
- Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S. J.: Sources and distributions of dust aerosols simulated with the GOCART model, *J. Geophys. Res.-Atmos.*, 106, 20255–20273, doi:10.1029/2000jd000053, 2001.
- Goldstein, A. H. and Galbally, I. E.: Known and unexplored organic constituents in the earth’s atmosphere, *Environ. Sci. Technol.*, 41, 1514–1521, doi:10.1021/es072476p, 2007.
- Graham, B., Guyon, P., Maenhaut, W., Taylor, P. E., Ebert, M., Matthias-Maser, S., Mayol-Bracero, O. L., Godoi, R. H. M., Artaxo, P., Meixner, F. X., Moura, M. A. L., Rocha, C., Van Grieken, R., Glovsky, M. M., Flagan, R. C., and Andreae, M. O.: Composition and diurnal variability of the natural Amazonian aerosol, *J. Geophys. Res.-Atmos.*, 108, 4765, doi:10.1029/2003JD004049, 2003.
- Granier, C., Lamarque, J. F., Mieville, A., Muller, J. F., Olivier, J., Orlando, J., Peters, J., Petron, G., Tyndall, G., and Wallens, S.: POET, a database of surface emissions of ozone precursors, available on internet at <http://www.aero.jussieu.fr/projet/ACCENT/POET.php>, 2005.
- Griffin, R. J., Cocker, D. R., Flagan, R. C., and Seinfeld, J. H.: Organic aerosol formation from the oxidation of biogenic hydrocarbons, *J. Geophys. Res.-Atmos.*, 104, 3555–3567, doi:10.1029/1998jd100049, 1999a.
- Griffin, R. J., Cocker, D. R., Seinfeld, J. H., and Dabdub, D.: Estimate of global atmospheric organic aerosol from oxidation of biogenic hydrocarbons, *Geophys. Res. Lett.*, 26, 2721–2724, doi:10.1029/1999gl900476, 1999b.
- Guenther, A., Hewitt, C. N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W. A., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global model of natural volatile organic compound emissions, *J. Geophys. Res.-Atmos.*, 100, 8873–8892, 10.1029/94jd02950, 1995.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George,

- C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.
- Hansen, D. A., Edgerton, E. S., Hartsell, B. E., Jansen, J. J., Kandasamy, N., Hidy, G. M., and Blanchard, C. L.: The southeastern aerosol research and characterization study: Part 1-overview, *J. Air Waste Manage.*, 53, 1460–1471, 2003.
- Hansen, J., Russell, G., Rind, D., Stone, P., Lacis, A., Lebedeff, S., Ruedy, R., and Travis, L.: Efficient Three-Dimensional Global Models for Climate Studies: Models I and II, *Mon. Weather Rev.*, 111, 609–662, 1983.
- He, Z., Kim, Y. J., Ogunjobi, K. O., Kim, J. E., and Ryu, S. Y.: Carbonaceous aerosol characteristics of PM_{2.5} particles in north-eastern Asia in summer 2002, *Atmos. Environ.*, 38, 1795–1800, 2004.
- Heald, C. L., Henze, D. K., Horowitz, L. W., Feddema, J., Lamarque, J. F., Guenther, A., Hess, P. G., Vitt, F., Seinfeld, J. H., Goldstein, A. H., and Fung, I.: Predicted change in global secondary organic aerosol concentrations in response to future climate, emissions, and land use change, *J. Geophys. Res.-Atmos.*, 113, D05211, doi:10.1029/2007JD009092, 2008.
- Heald, C. L., Ridley, D. A., Kreidenweis, S. M., and Drury, E. E.: Satellite observations cap the atmospheric organic aerosol budget, *Geophys. Res. Lett.*, 37, L24808, doi:10.1029/2010gl045095, 2010.
- Heald, C. L., Coe, H., Jimenez, J. L., Weber, R. J., Bahreini, R., Middlebrook, A. M., Russell, L. M., Jolleys, M., Fu, T.-M., Allan, J. D., Bower, K. N., Capes, G., Crosier, J., Morgan, W. T., Robinson, N. H., Williams, P. I., Cubison, M. J., DeCarlo, P. F., and Dunlea, E. J.: Exploring the vertical profile of atmospheric organic aerosol: comparing 17 aircraft field campaigns with a global model, *Atmos. Chem. Phys.*, 11, 12673–12696, doi:10.5194/acp-11-12673-2011, 2011.
- Henze, D. K. and Seinfeld, J. H.: Global secondary organic aerosol from isoprene oxidation, *Geophys. Res. Lett.*, 33, L09812, doi:10.1029/2006gl025976, 2006.
- Ho, K. F., Cao, J. J., Harrison, R. M., Lee, S. C., and Bau, K. K.: Indoor/outdoor relationships of organic carbon (OC) and elemental carbon (EC) in PM_{2.5} in roadside environment of Hong Kong, *Atmos. Environ.*, 38, 6327–6335, 2004.
- Hodzic, A., Jimenez, J. L., Madronich, S., Canagaratna, M. R., DeCarlo, P. F., Kleinman, L., and Fast, J.: Modeling organic aerosols in a megacity: potential contribution of semi-volatile and intermediate volatility primary organic compounds to secondary organic aerosol formation, *Atmos. Chem. Phys.*, 10, 5491–5514, doi:10.5194/acp-10-5491-2010, 2010.
- Hodzic, A., Madronich, S., Aumont, B., Lee-Taylor, J., Karl, T., Camredon, M., and Mouchel-Vallon, C.: Limited influence of dry deposition of semivolatile organic vapors on secondary organic aerosol formation in the urban plume, *Geophys. Res. Lett.*, 40, 3302–3307, doi:10.1002/grl.50611, 2013.
- Hodzic, A., Aumont, B., Knute, C., Lee-Taylor, J., Madronich, S., and Tyndall, G.: Volatility Dependence of Henry's Law Constants Of Condensable Organics: Application to Estimate Depositional Loss of Secondary Organic Aerosols, *Geophys. Res. Lett.*, 41, 4795–4804, doi:10.1002/2014GL060649, 2014.
- Hoyle, C. R., Berntsen, T., Myhre, G., and Isaksen, I. S. A.: Secondary organic aerosol in the global aerosol – chemical transport model Oslo CTM2, *Atmos. Chem. Phys.*, 7, 5675–5694, doi:10.5194/acp-7-5675-2007, 2007.
- Hoyle, C. R., Myhre, G., Berntsen, T. K., and Isaksen, I. S. A.: Anthropogenic influence on SOA and the resulting radiative forcing, *Atmos. Chem. Phys.*, 9, 2715–2728, doi:10.5194/acp-9-2715-2009, 2009.
- Hoyle, C. R., Boy, M., Donahue, N. M., Fry, J. L., Glasius, M., Guenther, A., Hallar, A. G., Huff Hartz, K., Petters, M. D., Petäjä, T., Rosenoern, T., and Sullivan, A. P.: A review of the anthropogenic influence on biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 11, 321–343, doi:10.5194/acp-11-321-2011, 2011.
- Hueglin, C., Gehrig, R., Baltensperger, U., Gysel, M., Monn, C., and Vonmont, H.: Chemical characterisation of PM_{2.5}, PM₁₀ and coarse particles at urban, near-city and rural sites in Switzerland, *Atmos. Environ.*, 39, 637–651, 2005.
- Huijnen, V., Williams, J., van Weele, M., van Noije, T., Krol, M., Dentener, F., Segers, A., Houweling, S., Peters, W., de Laat, J., Boersma, F., Bergamaschi, P., van Velthoven, P., Le Sager, P., Eskes, H., Alkemade, F., Scheele, R., Nédélec, P., and Pätz, H.-W.: The global chemistry transport model TM5: description and evaluation of the tropospheric chemistry version 3.0, *Geosci. Model Dev.*, 3, 445–473, doi:10.5194/gmd-3-445-2010, 2010.
- IPCC: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- Ito, A. and Penner, J. E.: Historical emissions of carbonaceous aerosols from biomass and fossil fuel burning for the period 1870–2000, *Global Biogeochem. Cy.*, 19, GB2028, doi:10.1029/2004gb002374, 2005.
- Jathar, S. H., Farina, S. C., Robinson, A. L., and Adams, P. J.: The influence of semi-volatile and reactive primary emissions on the abundance and properties of global organic aerosol, *Atmos. Chem. Phys.*, 11, 7727–7746, doi:10.5194/acp-11-7727-2011, 2011.
- Jayne, J. T., Leard, D. C., Zhang, X. F., Davidovits, P., Smith, K. A., Kolb, C. E., and Worsnop, D. R.: Development of an aerosol mass spectrometer for size and composition analysis of submicron particles, *Aerosol Sci. Tech.*, 33, 49–70, doi:10.1080/027868200410840, 2000.
- Jeong, C. H., Lee, D. W., Kim, E., and Hopke, P. K.: Measurement of real-time PM_{2.5} mass, sulfate, and carbonaceous aerosols at the multiple monitoring sites, *Atmos. Environ.*, 38, 5247–5256, 2004.
- Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R.,

- Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525–1529, doi:10.1126/science.1180353, 2009.
- Jöckel, P., Sander, R., Kerkweg, A., Tost, H., and Lelieveld, J.: Technical Note: The Modular Earth Submodel System (MESSy) – a new approach towards Earth System Modeling, *Atmos. Chem. Phys.*, 5, 433–444, doi:10.5194/acp-5-433-2005, 2005.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, doi:10.5194/bg-9-527-2012, 2012.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–471, doi:10.1175/1520-0477(1996)077<0437:tnyrp>2.0.co;2, 1996.
- Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.
- Kanakidou, M., Duce, R. A., Prospero, J. M., Baker, A. R., Benitez-Nelson, C., Dentener, F. J., Hunter, K. A., Liss, P. S., Mahowald, N., Okin, G. S., Sarin, M., Tsigaridis, K., Uematsu, M., Zamora, L. M., and Zhu, T.: Atmospheric fluxes of organic N and P to the global ocean, *Global Biogeochem. Cy.*, 26, Gb3026, doi:10.1029/2011gb004277, 2012.
- Kazil, J., Stier, P., Zhang, K., Quaas, J., Kinne, S., O'Donnell, D., Rast, S., Esch, M., Ferrachat, S., Lohmann, U., and Feichter, J.: Aerosol nucleation and its role for clouds and Earth's radiative forcing in the aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 10, 10733–10752, doi:10.5194/acp-10-10733-2010, 2010.
- Kim, B. M., Teffera, S., and Zeldin, M. D.: Characterization of PM_{2.5} and PM₁₀ in the South Coast Air Basin of southern California: Part 1 – Spatial variations, *J. Air Waste Manage.*, 50, 2034–2044, 2000.
- Kim, J., Yoon, S. C., Jefferson, A., Zahorowski, W., and Kang, C. H.: Air mass characterization and source region analysis for the Gosan super-site, Korea, during the ACE-Asia 2001 field campaign, *Atmos. Environ.*, 39, 6513–6523, 2005.
- Kinne, S., Schulz, M., Textor, C., Guibert, S., Balkanski, Y., Bauer, S. E., Bernsten, T., Berglen, T. F., Boucher, O., Chin, M., Collins, W., Dentener, F., Diehl, T., Easter, R., Feichter, J., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Herzog, M., Horowitz, L., Isaksen, I., Iversen, T., Kirkevåg, A., Kloster, S., Koch, D., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Lesins, G., Liu, X., Lohmann, U., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T., and Tie, X.: An AeroCom initial assessment – optical properties in aerosol component modules of global models, *Atmos. Chem. Phys.*, 6, 1815–1834, doi:10.5194/acp-6-1815-2006, 2006.
- Kirkevåg, A., Iversen, T., Seland, Ø., Hoose, C., Kristjansson, J. E., Struthers, H., Ekman, A. M. L., Ghan, S., Griesfeller, J., Nilsson, E. D., and Schulz, M.: Aerosol–climate interactions in the Norwegian Earth System Model – NorESM1-M, *Geosci. Model Dev.*, 6, 207–244, doi:10.5194/gmd-6-207-2013, 2013.
- Koch, D., Schmidt, G. A., and Field, C. V.: Sulfur, sea salt, and radionuclide aerosols in GISS ModelE, *J. Geophys. Res.-Atmos.*, 111, D06206, doi:10.1029/2004jd005550, 2006.
- Koch, D., Bond, T. C., Streets, D., Unger, N., and van der Werf, G. R.: Global impacts of aerosols from particular source regions and sectors, *J. Geophys. Res.-Atmos.*, 112, D02205, doi:10.1029/2005jd007024, 2007.
- Kokkola, H., Korhonen, H., Lehtinen, K. E. J., Makkonen, R., Asmi, A., Järvenoja, S., Anttila, T., Partanen, A.-I., Kulmala, M., Järvinen, H., Laaksonen, A., and Kerminen, V.-M.: SALSA – a Sectional Aerosol module for Large Scale Applications, *Atmos. Chem. Phys.*, 8, 2469–2483, doi:10.5194/acp-8-2469-2008, 2008.
- Koulouri, E., Saarikoski, S., Theodosi, C., Markaki, Z., Gerasopoulos, E., Kouvarakis, G., Makela, T., Hillamo, R., and Mihalopoulos, N.: Chemical composition and sources of fine and coarse aerosol particles in the Eastern Mediterranean, *Atmos. Environ.*, 42, 6542–6550, doi:10.1016/j.atmosenv.2008.04.010, 2008.
- Krivacsy, Z., Gelencser, A., Kiss, G., Meszaros, E., Molnar, A., Hoffer, A., Meszaros, T., Sarvari, Z., Temesi, D., Varga, B., Baltensperger, U., Nyeki, S., and Weingartner, E.: Study on the chemical character of water soluble organic compounds in fine atmospheric aerosol at the Jungfraujoch, *J. Atmos. Chem.*, 39, 235–259, 2001.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schulz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*, 5, 369–411, doi:10.5194/gmd-5-369-2012, 2012.
- Lathièrre, J., Hauglustaine, D. A., De Noblet-Ducoudre, N., Krinner, G., and Folberth, G. A.: Past and future changes in biogenic volatile organic compound emissions simulated with a global dynamic vegetation model, *Geophys. Res. Lett.*, 32, L20818, doi:10.1029/2005GL024164, 2005.
- Lee, Y. H. and Adams, P. J.: Evaluation of aerosol distributions in the GISS-TOMAS global aerosol microphysics model with re-

- mote sensing observations, *Atmos. Chem. Phys.*, 10, 2129–2144, doi:10.5194/acp-10-2129-2010, 2010.
- Lee, Y. H. and Adams, P. J.: A Fast and Efficient Version of the TwO-Moment Aerosol Sectional (TOMAS) Global Aerosol Microphysics Model, *Aerosol Sci. Tech.*, 46, 678–689, doi:10.1080/02786826.2011.643259, 2012.
- Lee, Y. H., Adams, P. J., and Shindell, D. T.: ModelE2-TOMAS development and evaluation using aerosol optical depths, mass and number concentrations, *Geosci. Model Dev. Discuss.*, 7, 5831–5918, doi:10.5194/gmdd-7-5831-2014, 2014.
- Li, L., Wang, W., Feng, J., Zhang, D., Li, H., Gu, Z., Wang, B., Sheng, G., and Fu, J.: Composition, source, mass closure of PM_{2.5} aerosols for four forests in eastern China, *J. Environ. Sci.*, 22, 405–412, 2010.
- Liao, H., Henze, D. K., Seinfeld, J. H., Wu, S. L., and Mickley, L. J.: Biogenic secondary organic aerosol over the United States: Comparison of climatological simulations with observations, *J. Geophys. Res.-Atmos.*, 112, D06201, doi:10.1029/2006JD007813, 2007.
- Liggio, J., Li, S. M., and McLaren, R.: Reactive uptake of glyoxal by particulate matter, *J. Geophys. Res.-Atmos.*, 110, D10304, doi:10.1029/2004jd005113, 2005.
- Lim, Y. B., Tan, Y., and Turpin, B. J.: Chemical insights, explicit chemistry, and yields of secondary organic aerosol from OH radical oxidation of methylglyoxal and glyoxal in the aqueous phase, *Atmos. Chem. Phys.*, 13, 8651–8667, doi:10.5194/acp-13-8651-2013, 2013.
- Lin, G., Penner, J. E., Sillman, S., Taraborrelli, D., and Lelieveld, J.: Global modeling of SOA formation from dicarbonyls, epoxides, organic nitrates and peroxides, *Atmos. Chem. Phys.*, 12, 4743–4774, doi:10.5194/acp-12-4743-2012, 2012.
- Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J.-F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and Mitchell, D.: Toward a minimal representation of aerosols in climate models: description and evaluation in the Community Atmosphere Model CAM5, *Geosci. Model Dev.*, 5, 709–739, doi:10.5194/gmd-5-709-2012, 2012.
- Liu, X. H., Penner, J. E., Das, B. Y., Bergmann, D., Rodriguez, J. M., Strahan, S., Wang, M. H., and Feng, Y.: Uncertainties in global aerosol simulations: Assessment using three meteorological data sets, *J. Geophys. Res.-Atmos.*, 112, D11212, doi:10.1029/2006jd008216, 2007.
- Long, M. S., Keene, W. C., Kieber, D. J., Erickson, D. J., and Maring, H.: A sea-state based source function for size- and composition-resolved marine aerosol production, *Atmos. Chem. Phys.*, 11, 1203–1216, doi:10.5194/acp-11-1203-2011, 2011.
- Long, R. W., Eatough, N. L., Mangelson, N. F., Thompson, W., Fiet, K., Smith, S., Smith, R., Eatough, D. J., Pope, C. A., and Wilson, W. E.: The measurement of PM_{2.5}, including semi-volatile components, in the EMPACT program: results from the Salt Lake City Study, *Atmos. Environ.*, 37, 4407–4417, 2003.
- Mann, G. W., Carslaw, K. S., Ridley, D. A., Spracklen, D. V., Pringle, K. J., Merikanto, J., Korhonen, H., Schwarz, J. P., Lee, L. A., Manktelow, P. T., Woodhouse, M. T., Schmidt, A., Breider, T. J., Emmerson, K. M., Reddington, C. L., Chipperfield, M. P., and Pickering, S. J.: Intercomparison of modal and sectional aerosol microphysics representations within the same 3-D global chemical transport model, *Atmos. Chem. Phys.*, 12, 4449–4476, doi:10.5194/acp-12-4449-2012, 2012.
- Mann, G. W., Carslaw, K. S., Reddington, C. L., Pringle, K. J., Schulz, M., Asmi, A., Spracklen, D. V., Ridley, D. A., Woodhouse, M. T., Lee, L. A., Zhang, K., Ghan, S. J., Easter, R. C., Liu, X., Stier, P., Lee, Y. H., Adams, P. J., Tost, H., Lelieveld, J., Bauer, S. E., Tsigaridis, K., van Noije, T. P. C., Strunk, A., Vignati, E., Bellouin, N., Dalvi, M., Johnson, C. E., Bergman, T., Kokkola, H., von Salzen, K., Yu, F., Luo, G., Petzold, A., Heintzenberg, J., Clarke, A., Ogren, J. A., Gras, J., Baltensperger, U., Kaminski, U., Jennings, S. G., O'Dowd, C. D., Harrison, R. M., Beddows, D. C. S., Kulmala, M., Viisanen, Y., Ulevicius, V., Mihalopoulos, N., Zdimal, V., Fiebig, M., Hansson, H.-C., Swietlicki, E., and Henzing, J. S.: Intercomparison and evaluation of global aerosol microphysical properties among AeroCom models of a range of complexity, *Atmos. Chem. Phys.*, 14, 4679–4713, doi:10.5194/acp-14-4679-2014, 2014.
- Middlebrook, A. M., Bahreini, R., Jimenez, J. L., and Canagaratna, M. R.: Evaluation of Composition-Dependent Collection Efficiencies for the Aerodyne Aerosol Mass Spectrometer using Field Data, *Aerosol Sci. Tech.*, 46, 258–271, doi:10.1080/02786826.2011.620041, 2012.
- Mihalopoulos, N., Stephanou, E., Kanakidou, M., Pilitsidis, S., and Bousquet, P.: Tropospheric aerosol ionic composition in the Eastern Mediterranean region, *Tellus B*, 49, 314–326, doi:10.1034/j.1600-0889.49.issue3.7.x, 1997.
- Miller, R. L., Cakmur, R. V., Perlwitz, J., Geogdzhayev, I. V., Ginoux, P., Koch, D., Kohfeld, K. E., Prigent, C., Ruedy, R., Schmidt, G. A., and Tegen, I.: Mineral dust aerosols in the NASA goddard institute for Space Sciences ModelE atmospheric general circulation model, *J. Geophys. Res.-Atmos.*, 111, D06208, doi:10.1029/2005jd005796, 2006.
- Mohr, C., DeCarlo, P. F., Heringa, M. F., Chirico, R., Slowik, J. G., Richter, R., Reche, C., Alastuey, A., Querol, X., Seco, R., Peñuelas, J., Jiménez, J. L., Crippa, M., Zimmermann, R., Baltensperger, U., and Prévôt, A. S. H.: Identification and quantification of organic aerosol from cooking and other sources in Barcelona using aerosol mass spectrometer data, *Atmos. Chem. Phys.*, 12, 1649–1665, doi:10.5194/acp-12-1649-2012, 2012.
- Molina, M. J., Ivanov, A. V., Trakhtenberg, S., and Molina, L. T.: Atmospheric evolution of organic aerosol, *Geophys. Res. Lett.*, 31, L22104, doi:10.1029/2004gl020910, 2004.
- Morcrette, J. J., Boucher, O., Jones, L., Salmond, D., Bechtold, P., Beljaars, A., Benedetti, A., Bonet, A., Kaiser, J. W., Razinger, M., Schulz, M., Serrar, S., Simmons, A. J., Sofiev, M., Suttie, M., Tompkins, A. M., and Untch, A.: Aerosol analysis and forecast in the European Centre for Medium-Range Weather Forecasts Integrated Forecast System: Forward modeling, *J. Geophys. Res.-Atmos.*, 114, D06206, doi:10.1029/2008jd011235, 2009.
- Müller, J.-F., Ceulemans, K., Compennolle, S., Stavrou, T., Peeters, J., Nguyen, T. L., Vereecken, L., Winterhalter, R., Moortgat, G., Kanawati, B., and Sadezky, A.: Impact of Biogenic emissions on Organic aerosols and Oxidants in the Troposphere (IBOOT), Belgian Science Policy, Brussels, Final report SD/AT/03, 94 pp., 2009.
- Murphy, B. N., Donahue, N. M., Fountoukis, C., and Pandis, S. N.: Simulating the oxygen content of ambient organic aerosol with the 2D volatility basis set, *Atmos. Chem. Phys.*, 11, 7859–7873, doi:10.5194/acp-11-7859-2011, 2011.

- Murphy, B. N., Donahue, N. M., Fountoukis, C., Dall'Osto, M., O'Dowd, C., Kiendler-Scharr, A., and Pandis, S. N.: Functionalization and fragmentation during ambient organic aerosol aging: application of the 2-D volatility basis set to field studies, *Atmos. Chem. Phys.*, 12, 10797–10816, doi:10.5194/acp-12-10797-2012, 2012.
- Murphy, B. N., Donahue, N. M., Robinson, A. L., and Pandis, S. N.: A naming convention for atmospheric organic aerosol, *Atmos. Chem. Phys.*, 14, 5825–5839, doi:10.5194/acp-14-5825-2014, 2014.
- Murphy, D. M., Cziczo, D. J., Froyd, K. D., Hudson, P. K., Matthew, B. M., Middlebrook, A. M., Peltier, R. E., Sullivan, A., Thomson, D. S., and Weber, R. J.: Single-particle mass spectrometry of tropospheric aerosol particles, *J. Geophys. Res.-Atmos.*, 111, D23S32, doi:10.1029/2006JD007340, 2006.
- Myhre, G., Berglen, T. F., Johnsrud, M., Hoyle, C. R., Berntsen, T. K., Christopher, S. A., Fahey, D. W., Isaksen, I. S. A., Jones, T. A., Kahn, R. A., Loebe, N., Quinn, P., Remer, L., Schwarz, J. P., and Yttri, K. E.: Modelled radiative forcing of the direct aerosol effect with multi-observation evaluation, *Atmos. Chem. Phys.*, 9, 1365–1392, doi:10.5194/acp-9-1365-2009, 2009.
- Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Berntsen, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J.-H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmos. Chem. Phys.*, 13, 1853–1877, doi:10.5194/acp-13-1853-2013, 2013.
- Myriokefalitakis, S., Vrekoussis, M., Tsigaridis, K., Wittrock, F., Richter, A., Brühl, C., Volkamer, R., Burrows, J. P., and Kanakidou, M.: The influence of natural and anthropogenic secondary sources on the glyoxal global distribution, *Atmos. Chem. Phys.*, 8, 4965–4981, doi:10.5194/acp-8-4965-2008, 2008.
- Myriokefalitakis, S., Vignati, E., Tsigaridis, K., Papadimas, C., Sciare, J., Mihalopoulos, N., Facchini, M. C., Rinaldi, M., Dentener, F. J., Ceburnis, D., Hatzianastasiou, N., Dowd, C. D., van Weele, M., and Kanakidou, M.: Global Modeling of the Oceanic Source of Organic Aerosols, *Advances in Meteorology*, 2010, 939171, doi:10.1155/2010/939171, 2010.
- Myriokefalitakis, S., Tsigaridis, K., Mihalopoulos, N., Sciare, J., Nenes, A., Kawamura, K., Segers, A., and Kanakidou, M.: In-cloud oxalate formation in the global troposphere: a 3-D modeling study, *Atmos. Chem. Phys.*, 11, 5761–5782, doi:10.5194/acp-11-5761-2011, 2011.
- Ng, N. L., Herndon, S. C., Trimborn, A., Canagaratna, M. R., Croteau, P. L., Onasch, T. B., Sueper, D., Worsnop, D. R., Zhang, Q., Sun, Y. L., and Jayne, J. T.: An Aerosol Chemical Speciation Monitor (ACSM) for Routine Monitoring of the Composition and Mass Concentrations of Ambient Aerosol, *Aerosol Sci. Tech.*, 45, 780–794, doi:10.1080/02786826.2011.560211, 2011.
- O'Donnell, D., Tsigaridis, K., and Feichter, J.: Estimating the direct and indirect effects of secondary organic aerosols using ECHAM5-HAM, *Atmos. Chem. Phys.*, 11, 8635–8659, doi:10.5194/acp-11-8635-2011, 2011.
- O'Dowd, C. D., Facchini, M. C., Cavalli, F., Ceburnis, D., Mircea, M., Decesari, S., Fuzzi, S., Yoon, Y. J., and Putaud, J. P.: Biogenically driven organic contribution to marine aerosol, *Nature*, 431, 676–680, doi:10.1038/nature02959, 2004.
- Offenberg, J. H., Kleindienst, T. E., Jaoui, M., Lewandowski, M., and Edney, E. O.: Thermal properties of secondary organic aerosols, *Geophys. Res. Lett.*, 33, L03816, doi:10.1029/2005gl024623, 2006.
- Olivier, J. G. J., Berdowski, J. J. M., Peters, J. A. H. W., Bakker, J., Visschedijk, A. J. H., and Bloos, J. P. J.: Applications of EDGAR. Including a description of EDGAR 3.0: reference database with trend data for 1970–1995, RIVM, Bilthoven, RIVM report no. 773301 001/NOP report no. 410200 051, 2001.
- Paasonen, P., Asmi, A., Petaja, T., Kajos, M. K., Aijala, M., Junninen, H., Holst, T., Abbatt, J. P. D., Arneth, A., Birmili, W., van der Gon, H. D., Hamed, A., Hoffer, A., Laakso, L., Laaksonen, A., Leaitch, W. R., Plass-Dulmer, C., Pryor, S. C., Raisanen, P., Swietlicki, E., Wiedensohler, A., Worsnop, D. R., Kerminen, V. M., and Kulmala, M.: Warming-induced increase in aerosol number concentration likely to moderate climate change, *Nat. Geosci.*, 6, 438–442, doi:10.1038/ngeo1800, 2013.
- Park, R. J., Jacob, D. J., Chin, M., and Martin, R. V.: Sources of carbonaceous aerosols over the United States and implications for natural visibility, *J. Geophys. Res.-Atmos.*, 108, 4355, doi:10.1029/2002jd003190, 2003.
- Paulot, F., Crouse, J. D., Kjaergaard, H. G., Kurten, A., St Clair, J. M., Seinfeld, J. H., and Wennberg, P. O.: Unexpected Epoxide Formation in the Gas-Phase Photooxidation of Isoprene, *Science*, 325, 730–733, doi:10.1126/science.1172910, 2009.
- Petrenko, M., Kahn, R., Chin, M., Soja, A., Kucsera, T., and Harshvardhan: The use of satellite-measured aerosol optical depth to constrain biomass burning emissions source strength in the global model GOCART, *J. Geophys. Res.-Atmos.*, 117, D18212, doi:10.1029/2012jd017870, 2012.
- Philip, S., Martin, R. V., Pierce, J. R., Jimenez, J. L., Zhang, Q., Canagaratna, M. R., Spracklen, D. V., Nowlan, C. R., Lamsal, L. N., Cooper, M. J., and Krotkov, N. A.: Spatially and seasonally resolved estimate of the ratio of organic mass to organic carbon, *Atmos. Environ.*, 87, 34–40, doi:10.1016/j.atmosenv.2013.11.065, 2014.
- Piccot, S. D., Watson, J. J., and Jones, J. W.: A global inventory of volatile organic compound emissions from anthropogenic sources, *J. Geophys. Res.-Atmos.*, 97, 9897–9912, 1992.
- Pindado, O., Ma Perez, R., Garcia, S., Sanchez, M., Galan, P., and Fernandez, M.: Characterization and sources assignment of PM_{2.5} organic aerosol in a rural area of Spain, *Atmos. Environ.*, 43, 2796–2803, 2009.
- Pozzoli, L., Bey, I., Rast, S., Schultz, M. G., Stier, P., and Feichter, J.: Trace gas and aerosol interactions in the fully coupled model of aerosol-chemistry-climate ECHAM5-HAMMOZ: 1. Model description and insights from the spring 2001 TRACE-P experiment, *J. Geophys. Res.-Atmos.*, 113, D07308, doi:10.1029/2007jd009007, 2008.
- Pozzoli, L., Janssens-Maenhout, G., Diehl, T., Bey, I., Schultz, M. G., Feichter, J., Vignati, E., and Dentener, F.: Re-analysis of tropospheric sulfate aerosol and ozone for the period 1980–2005 using the aerosol-chemistry-climate model ECHAM5-HAMMOZ, *Atmos. Chem. Phys.*, 11, 9563–9594, doi:10.5194/acp-11-9563-2011, 2011.

- Pringle, K. J., Tost, H., Message, S., Steil, B., Giannadaki, D., Nenes, A., Fountoukis, C., Stier, P., Vignati, E., and Lelieveld, J.: Description and evaluation of GMXe: a new aerosol submodel for global simulations (v1), *Geosci. Model Dev.*, 3, 391–412, doi:10.5194/gmd-3-391-2010, 2010.
- Pye, H. O. T. and Seinfeld, J. H.: A global perspective on aerosol from low-volatility organic compounds, *Atmos. Chem. Phys.*, 10, 4377–4401, doi:10.5194/acp-10-4377-2010, 2010.
- Quaas, J., Ming, Y., Menon, S., Takemura, T., Wang, M., Penner, J. E., Gettelman, A., Lohmann, U., Bellouin, N., Boucher, O., Sayer, A. M., Thomas, G. E., McComiskey, A., Feingold, G., Hoose, C., Kristjánsson, J. E., Liu, X., Balkanski, Y., Donner, L. J., Ginoux, P. A., Stier, P., Grandey, B., Feichter, J., Sednev, I., Bauer, S. E., Koch, D., Grainger, R. G., Kirkevåg, A., Iversen, T., Seland, Ø., Easter, R., Ghan, S. J., Rasch, P. J., Morrison, H., Lamarque, J.-F., Iacono, M. J., Kinne, S., and Schulz, M.: Aerosol indirect effects – general circulation model intercomparison and evaluation with satellite data, *Atmos. Chem. Phys.*, 9, 8697–8717, doi:10.5194/acp-9-8697-2009, 2009.
- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications, *J. Climate*, 24, 3624–3648, doi:10.1175/JCLI-D-11-00015.1, 2011.
- Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J. R., and Pandis, S. N.: Rethinking organic aerosols: Semivolatile emissions and photochemical aging, *Science*, 315, 1259–1262, doi:10.1126/science.1133061, 2007.
- Saathoff, H., Naumann, K.-H., Möhler, O., Jonsson, Å. M., Halqu Coast, M., Kiendler-Scharr, A., Mentel, Th. F., Tillmann, R., and Schurath, U.: Temperature dependence of yields of secondary organic aerosols from the ozonolysis of α -pinene and limonene, *Atmos. Chem. Phys.*, 9, 1551–1577, doi:10.5194/acp-9-1551-2009, 2009.
- Salma, I., Chi, X. G., and Maenhaut, W.: Elemental and organic carbon in urban canyon and background environments in Budapest, Hungary, *Atmos. Environ.*, 38, 27–36, 2004.
- Samset, B. H., Myhre, G., Schulz, M., Balkanski, Y., Bauer, S., Bernsten, T. K., Bian, H., Bellouin, N., Diehl, T., Easter, R. C., Ghan, S. J., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Penner, J. E., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Black carbon vertical profiles strongly affect its radiative forcing uncertainty, *Atmos. Chem. Phys.*, 13, 2423–2434, doi:10.5194/acp-13-2423-2013, 2013.
- Sawant, A. A., Na, K., Zhu, X. N., and Cocker, D. R.: Chemical characterization of outdoor PM_{2.5} and gas-phase compounds in Mira Loma, California, *Atmos. Environ.*, 38, 5517–5528, 2004.
- Schultz, M. G., Backman, L., Balkanski, Y., Bjoerndalsaeter, S., Brand, R., Burrows, J. P., Dalsoeren, S., de Vasconcelos, M., Grodtmann, B., Haughustaine, D. A., Heil, A., Hoelzemann, J. J., Isaksen, I. S. A., Kaurola, J., Knorr, W., Ladstaetter-Weißenmayer, A., Mota, B., Oom, D., Pacyna, J., Panasiuk, D., Pereira, J. M. C., Pulles, T., Pyle, J., Rast, S., Richter, A., Savage, N., Schnadt, C., Schulz, M., Spessa, A., Staehelin, J., Sundet, J. K., Szopa, S., Thonicke, K., van het Bolscher, M., van Noije, T., van Velthoven, P., Vik, A. F., and Wittrock, F.: REanalysis of the TROpospheric chemical composition over the past 40 years (RETRO) – A long-term global modeling study of tropospheric chemistry, Final Report, Jülich/Hamburg, Germany, 2007 (Published as report no. 48/2007 in the series “Reports on Earth System Science” of the Max Planck Institute for Meteorology, Hamburg, ISSN 1614-1199), 2007.
- Schulz, M.: Constraining model estimates of the aerosol radiative forcing, Thèse d’habilitation à diriger des recherches, Université Pierre et Marie Curie, Paris VI, 2007.
- Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S., Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, S., Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-industrial simulations, *Atmos. Chem. Phys.*, 6, 5225–5246, doi:10.5194/acp-6-5225-2006, 2006.
- Schulz, M., Chin, M., and Kinne, S.: The aerosol model comparison project, AeroCom, phase II: clearing up diversity, IGAC newsletter, No. 41, 2–11, 2009.
- Sciare, J., Baboukas, E., and Mihalopoulos, N.: Short-term variability of atmospheric DMS and its oxidation products at Amsterdam Island during summer time, *J. Atmos. Chem.*, 39, 281–302, doi:10.1023/a:1010631305307, 2001.
- Sciare, J., Favez, O., Sarda-Estève, R., Oikonomou, K., Cachier, H., and Kazan, V.: Long-term observations of carbonaceous aerosols in the Austral Ocean atmosphere: Evidence of a biogenic marine organic source, *J. Geophys. Res.-Atmos.*, 114, D15302, doi:10.1029/2009jd011998, 2009.
- Shakya, K. M., Ziemba, L. D., and Griffin, R. J.: Characteristics and Sources of Carbonaceous, Ionic, and Isotopic Species of Wintertime Atmospheric Aerosols in Kathmandu Valley, Nepal, *Aerosol Air Qual. Res.*, 10, 219–230, doi:10.4209/aaqr.2009.10.0068, 2010.
- Shank, L. M., Howell, S., Clarke, A. D., Freitag, S., Brekhovskikh, V., Kapustin, V., McNaughton, C., Campos, T., and Wood, R.: Organic matter and non-refractory aerosol over the remote Southeast Pacific: oceanic and combustion sources, *Atmos. Chem. Phys.*, 12, 557–576, doi:10.5194/acp-12-557-2012, 2012.
- Smith, D. J. T., Harrison, R. M., Luhana, L., Pio, C. A., Castro, L. M., Tariq, M. N., Hayat, S., and Quraishi, T.: Concentrations of particulate airborne polycyclic aromatic hydrocarbons and metals collected in Lahore, Pakistan, *Atmos. Environ.*, 30, 4031–4040, doi:10.1016/1352-2310(96)00107-0, 1996.
- Spracklen, D. V., Arnold, S. R., Sciare, J., Carslaw, K. S., and Pio, C.: Globally significant oceanic source of organic carbon aerosol, *Geophys. Res. Lett.*, 35, L12811, doi:10.1029/2008gl033359, 2008.
- Spracklen, D. V., Jimenez, J. L., Carslaw, K. S., Worsnop, D. R., Evans, M. J., Mann, G. W., Zhang, Q., Canagaratna, M. R., Allan, J., Coe, H., McFiggans, G., Rap, A., and Forster, P.: Aerosol mass spectrometer constraint on the global secondary organic aerosol budget, *Atmos. Chem. Phys.*, 11, 12109–12136, doi:10.5194/acp-11-12109-2011, 2011.
- Stavrakou, T., Müller, J.-F., De Smedt, I., Van Roozendael, M., Kanakidou, M., Vrekoussis, M., Wittrock, F., Richter, A., and

- Burrows, J. P.: The continental source of glyoxal estimated by the synergistic use of spaceborne measurements and inverse modelling, *Atmos. Chem. Phys.*, 9, 8431–8446, doi:10.5194/acp-9-8431-2009, 2009.
- Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M., Balkanski, Y., Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 5, 1125–1156, doi:10.5194/acp-5-1125-2005, 2005.
- Strader, R., Lurmann, F., and Pandis, S. N.: Evaluation of secondary organic aerosol formation in winter, *Atmos. Environ.*, 33, 4849–4863, doi:10.1016/s1352-2310(99)00310-6, 1999.
- Surratt, J. D., Chan, A. W. H., Eddingsaas, N. C., Chan, M., Loza, C. L., Kwan, A. J., Hersey, S. P., Flagan, R. C., Wennberg, P. O., and Seinfeld, J. H.: Reactive intermediates revealed in secondary organic aerosol formation from isoprene, *P. Natl. Acad. Sci. USA*, 107, 6640–6645, doi:10.1073/pnas.091114107, 2010.
- Svendby, T. M., Lazaridis, M., and Torseth, K.: Temperature dependent secondary organic aerosol formation from terpenes and aromatics, *J. Atmos. Chem.*, 59, 25–46, doi:10.1007/s10874-007-9093-7, 2008.
- Szopa, S., Balkanski, Y., Schulz, M., Bekki, S., Cugnet, D., Fortems-Cheiney, A., Turquety, S., Cozic, A., Deandreis, C., Hauglustaine, D., Idelkadi, A., Lathiere, J., Lefevre, F., Marchand, M., Vuolo, R., Yan, N., and Dufresne, J. L.: Aerosol and ozone changes as forcing for climate evolution between 1850 and 2100, *Clim. Dynam.*, 40, 2223–2250, doi:10.1007/s00382-012-1408-y, 2013.
- Takemura, T., Okamoto, H., Maruyama, Y., Numaguti, A., Higurashi, A., and Nakajima, T.: Global three-dimensional simulation of aerosol optical thickness distribution of various origins, *J. Geophys. Res.-Atmos.*, 105, 17853–17873, 2000.
- Takemura, T., Uno, I., Nakajima, T., Higurashi, A., and Sano, I.: Modeling study of long-range transport of Asian dust and anthropogenic aerosols from East Asia, *Geophys. Res. Lett.*, 29, 2158–2158, 2002.
- Takemura, T., Nozawa, T., Emori, S., Nakajima, T. Y., and Nakajima, T.: Simulation of climate response to aerosol direct and indirect effects with aerosol transport-radiation model, *J. Geophys. Res.-Atmos.*, 110, D02202, doi:10.1029/2004jd005029, 2005.
- Takemura, T., Egashira, M., Matsuzawa, K., Ichijo, H., O'ishi, R., and Abe-Ouchi, A.: A simulation of the global distribution and radiative forcing of soil dust aerosols at the Last Glacial Maximum, *Atmos. Chem. Phys.*, 9, 3061–3073, doi:10.5194/acp-9-3061-2009, 2009.
- Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, I., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles within AeroCom, *Atmos. Chem. Phys.*, 6, 1777–1813, doi:10.5194/acp-6-1777-2006, 2006.
- Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Feichter, J., Fillmore, D., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, T., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: The effect of harmonized emissions on aerosol properties in global models – an AeroCom experiment, *Atmos. Chem. Phys.*, 7, 4489–4501, doi:10.5194/acp-7-4489-2007, 2007.
- Tiitta, P., Vakkari, V., Croteau, P., Beukes, J. P., van Zyl, P. G., Josipovic, M., Venter, A. D., Jaars, K., Pienaar, J. J., Ng, N. L., Canagaratna, M. R., Jayne, J. T., Kerminen, V.-M., Kokkola, H., Kulmala, M., Laaksonen, A., Worsnop, D. R., and Laakso, L.: Chemical composition, main sources and temporal variability of PM₁ aerosols in southern African grassland, *Atmos. Chem. Phys.*, 14, 1909–1927, doi:10.5194/acp-14-1909-2014, 2014.
- Tsigaridis, K. and Kanakidou, M.: Global modelling of secondary organic aerosol in the troposphere: a sensitivity analysis, *Atmos. Chem. Phys.*, 3, 1849–1869, doi:10.5194/acp-3-1849-2003, 2003.
- Tsigaridis, K. and Kanakidou, M.: Secondary organic aerosol importance in the future atmosphere, *Atmos. Environ.*, 41, 4682–4692, doi:10.1016/j.atmosenv.2007.03.045, 2007.
- Tsigaridis, K., Lathière, J., Kanakidou, M., and Hauglustaine, D. A.: Naturally driven variability in the global secondary organic aerosol over a decade, *Atmos. Chem. Phys.*, 5, 1891–1904, doi:10.5194/acp-5-1891-2005, 2005.
- Tsigaridis, K., Krol, M., Dentener, F. J., Balkanski, Y., Lathière, J., Metzger, S., Hauglustaine, D. A., and Kanakidou, M.: Change in global aerosol composition since preindustrial times, *Atmos. Chem. Phys.*, 6, 5143–5162, doi:10.5194/acp-6-5143-2006, 2006.
- Tsigaridis, K., Koch, D., and Menon, S.: Uncertainties and importance of sea spray composition on aerosol direct and indirect effects, *J. Geophys. Res.-Atmos.*, 118, 220–235, doi:10.1029/2012jd018165, 2013.
- Tsimpidi, A. P., Karydis, V. A., Zavala, M., Lei, W., Molina, L., Ulbrich, I. M., Jimenez, J. L., and Pandis, S. N.: Evaluation of the volatility basis-set approach for the simulation of organic aerosol formation in the Mexico City metropolitan area, *Atmos. Chem. Phys.*, 10, 525–546, doi:10.5194/acp-10-525-2010, 2010.
- Turpin, B. J. and Lim, H. J.: Species contributions to PM_{2.5} mass concentrations: Revisiting common assumptions for estimating organic mass, *Aerosol Sci. Tech.*, 35, 602–610, doi:10.1080/02786820152051454, 2001.
- van der Werf, G. R., Randerson, J. T., Collatz, G. J., and Giglio, L.: Carbon emissions from fires in tropical and subtropical ecosystems, *Glob. Change Biol.*, 9, 547–562, doi:10.1046/j.1365-2486.2003.00604.x, 2003.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Kasibhatla, P. S., and Arellano Jr., A. F.: Interannual variability in global biomass burning emissions from 1997 to 2004, *Atmos. Chem. Phys.*, 6, 3423–3441, doi:10.5194/acp-6-3423-2006, 2006.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

- van Noije, T. P. C., Le Sager, P., Segers, A. J., van Velthoven, P. F. J., Krol, M. C., and Hazeleger, W.: Simulation of tropospheric chemistry and aerosols with the climate model EC-Earth, *Geosci. Model Dev. Discuss.*, 7, 1933–2006, doi:10.5194/gmdd-7-1933-2014, 2014.
- Vignati, E., Wilson, J., and Stier, P.: M7: An efficient size-resolved aerosol microphysics module for large-scale aerosol transport models, *J. Geophys. Res.-Atmos.*, 109, D22202, doi:10.1029/2003jd004485, 2004.
- Vignati, E., Facchini, M. C., Rinaldi, M., Scannell, C., Ceburnis, D., Sciare, J., Kanakidou, M., Myriokefalitakis, S., Dentener, F., and O'Dowd, C. D.: Global scale emission and distribution of sea-spray aerosol: Sea-salt and organic enrichment, *Atmos. Environ.*, 44, 670–677, doi:10.1016/j.atmosenv.2009.11.013, 2010.
- Volkamer, R., Jimenez, J. L., San Martini, F., Dzepina, K., Zhang, Q., Salcedo, D., Molina, L. T., Worsnop, D. R., and Molina, M. J.: Secondary organic aerosol formation from anthropogenic air pollution: Rapid and higher than expected, *Geophys. Res. Lett.*, 33, L17811, doi:10.1029/2006gl026899, 2006.
- Volkamer, R., Ziemann, P. J., and Molina, M. J.: Secondary Organic Aerosol Formation from Acetylene (C_2H_2): seed effect on SOA yields due to organic photochemistry in the aerosol aqueous phase, *Atmos. Chem. Phys.*, 9, 1907–1928, doi:10.5194/acp-9-1907-2009, 2009.
- von Salzen, K.: Piecewise log-normal approximation of size distributions for aerosol modelling, *Atmos. Chem. Phys.*, 6, 1351–1372, doi:10.5194/acp-6-1351-2006, 2006.
- von Salzen, K., McFarlane, N. A., and Lazare, M.: The role of shallow convection in the water and energy cycles of the atmosphere, *Clim. Dynam.*, 25, 671–688, doi:10.1007/s00382-005-0051-2, 2005.
- Wang, M. H., Penner, J. E., and Liu, X. H.: Coupled IMPACT aerosol and NCAR CAM3 model: Evaluation of predicted aerosol number and size distribution, *J. Geophys. Res.-Atmos.*, 114, D06302, doi:10.1029/2008jd010459, 2009.
- Ward, T. J., Hamilton, R. F., and Smith, G. C.: The Missoula, Montana $PM_{2.5}$ speciation study – seasonal average concentrations, *Atmos. Environ.*, 38, 6371–6379, 2004.
- Xu, J., Bergin, M. H., Greenwald, R., Schauer, J. J., Shafer, M. M., Jaffrezo, J. L., and Aymoz, G.: Aerosol chemical, physical, and radiative characteristics near a desert source region of northwest China during ACE-Asia, *J. Geophys. Res.-Atmos.*, 109, D19S03, doi:10.1029/2003JD004239, 2004.
- Yu, F. and Luo, G.: Simulation of particle size distribution with a global aerosol model: contribution of nucleation to aerosol and CCN number concentrations, *Atmos. Chem. Phys.*, 9, 7691–7710, doi:10.5194/acp-9-7691-2009, 2009.
- Yu, F.: A secondary organic aerosol formation model considering successive oxidation aging and kinetic condensation of organic compounds: global scale implications, *Atmos. Chem. Phys.*, 11, 1083–1099, doi:10.5194/acp-11-1083-2011, 2011.
- Zappoli, S., Andracchio, A., Fuzzi, S., Facchini, M. C., Gelencser, A., Kiss, G., Krivacsy, Z., Molnar, A., Meszaros, E., Hansson, H. C., Rosman, K., and Zebuhr, Y.: Inorganic, organic and macromolecular components of fine aerosol in different areas of Europe in relation to their water solubility, *Atmos. Environ.*, 33, 2733–2743, 1999.
- Zhang, H., Wang, Z., Wang, Z., Liu, Q., Gong, S., Zhang, X., Shen, Z., Lu, P., Wei, X., Che, H., and Li, L.: Simulation of direct radiative forcing of aerosols and their effects on East Asian climate using an interactive AGCM-aerosol coupled system, *Clim. Dynam.*, 38, 1675–1693, doi:10.1007/s00382-011-1131-0, 2012.
- Zhang, K., O'Donnell, D., Kazil, J., Stier, P., Kinne, S., Lohmann, U., Ferrachat, S., Croft, B., Quaas, J., Wan, H., Rast, S., and Feichter, J.: The global aerosol-climate model ECHAM-HAM, version 2: sensitivity to improvements in process representations, *Atmos. Chem. Phys.*, 12, 8911–8949, doi:10.5194/acp-12-8911-2012, 2012.
- Zhang, Q., Alfarra, M. R., Worsnop, D. R., Allan, J. D., Coe, H., Canagaratna, M. R., and Jimenez, J. L.: Deconvolution and quantification of hydrocarbon-like and oxygenated organic aerosols based on aerosol mass spectrometry, *Environ. Sci. Technol.*, 39, 4938–4952, doi:10.1021/es0485681, 2005.
- Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., DeCarlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimojo, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, *Geophys. Res. Lett.*, 34, 6, L13801, doi:10.1029/2007g1029979, 2007.
- Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Ulbrich, I. M., Ng, N. L., Worsnop, D. R., and Sun, Y. L.: Understanding atmospheric organic aerosols via factor analysis of aerosol mass spectrometry: a review, *Anal. Bioanal. Chem.*, 401, 3045–3067, doi:10.1007/s00216-011-5355-y, 2011.
- Zhang, X., Cappa, C. D., Jathar, S. H., McVay, R. C., Ensberg, J. J., Kleeman, M. J., and Seinfeld, J. H.: Influence of vapor wall loss in laboratory chambers on yields of secondary organic aerosol, *P. Natl. Acad. Sci. USA*, 111, 5802–5807, doi:10.1073/pnas.1404727111, 2014.
- Zhang, X. Y., Wang, Y. Q., Niu, T., Zhang, X. C., Gong, S. L., Zhang, Y. M., and Sun, J. Y.: Atmospheric aerosol compositions in China: spatial/temporal variability, chemical signature, regional haze distribution and comparisons with global aerosols, *Atmos. Chem. Phys.*, 12, 779–799, doi:10.5194/acp-12-779-2012, 2012.