



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/84198/>

Version: Published Version

Article:

Kirby, A and Herbert, A (2013) Correlations between Income inequality and antimicrobial resistance. PLoS One, 8 (8). e73115. ISSN: 1932-6203

<https://doi.org/10.1371/journal.pone.0073115>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Correlations between Income Inequality and Antimicrobial Resistance

Andrew Kirby^{1*}, Annie Herbert²

1 Department of Microbiology, Leeds Teaching Hospitals National Health Service Trust, Leeds, West Yorkshire, United Kingdom, **2** Department of Research and Development, Pennine Acute National Health Service Trust, Manchester, Greater Manchester, United Kingdom

Abstract

Objectives: The aim of this study is to investigate if correlations exist between income inequality and antimicrobial resistance. This study's hypothesis is that income inequality at the national level is positively correlated with antimicrobial resistance within developed countries.

Data collection and analysis: Income inequality data were obtained from the Standardized World Income Inequality Database. Antimicrobial resistance data were obtained from the European antimicrobial Resistance Surveillance Network and outpatient antimicrobial consumption data, measured by Defined daily Doses per 1000 inhabitants per day, from the European Surveillance of antimicrobial Consumption group. Spearman's correlation coefficient (r) defined strengths of correlations of: > 0.8 as strong, > 0.5 as moderate and > 0.2 as weak. Confidence intervals and p values were defined for all r values. Correlations were calculated for the time period 2003-10, for 15 European countries.

Results: Income inequality and antimicrobial resistance correlations which were moderate or strong, with 95% confidence intervals > 0 , included the following. *Enterococcus faecalis* resistance to aminopenicillins, vancomycin and high level gentamicin was moderately associated with income inequality ($r = \geq 0.54$ for all three antimicrobials). *Escherichia coli* resistance to aminoglycosides, aminopenicillins, third generation cephalosporins and fluoroquinolones was moderately-strongly associated with income inequality ($r = \geq 0.7$ for all four antimicrobials). *Klebsiella pneumoniae* resistance to third generation cephalosporins, aminoglycosides and fluoroquinolones was moderately associated with income inequality ($r = \geq 0.5$ for all three antimicrobials). *Staphylococcus aureus* methicillin resistance and income inequality were strongly associated ($r = 0.87$).

Conclusion: As income inequality increases in European countries so do the rates of antimicrobial resistance for bacteria including *E. faecalis*, *E. coli*, *K. pneumoniae* and *S. aureus*. Further studies are needed to confirm these findings outside Europe and investigate the processes that could causally link income inequality and antimicrobial resistance.

Citation: Kirby A, Herbert A (2013) Correlations between Income Inequality and Antimicrobial Resistance. PLoS ONE 8(8): e73115. doi:10.1371/journal.pone.0073115

Editor: D William Cameron, University of Ottawa, Canada

Received: February 18, 2012; **Accepted:** July 23, 2013; **Published:** August 29, 2013

Copyright: © 2013 Kirby et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors have no support or funding to report.

Competing interests: The authors have declared that no competing interests exist.

* E-mail: kirbyandrew@doctors.org.uk

Introduction

Antimicrobial therapy has contributed greatly to improvements in human health care through the treatment of infectious diseases. Unfortunately, an increase in antimicrobially resistant microorganisms threatens to limit the effectiveness of antimicrobial therapy, and therefore the benefits derived from these drugs [1]. For example, the European Centre for Disease Control recently estimated that 25,000 deaths per year within Europe are directly related to antibiotic resistance [2].

Rates of antimicrobial resistance vary significantly between countries. These variations have been well documented and have been principally related to antimicrobial consumption, i.e. the more antibiotics a country consumes the more resistance is likely to result [3]. However there is limited research to fully explain why resistance rates vary between countries. One area that has yet to be researched is the impact of a nation's economic policy on antimicrobial consumption and the development of antimicrobial resistance [4]. A nation's economic policy determines many outcomes within the societies they impact on, including that country's income inequality. Income inequality is the difference in incomes

between those with the highest and lowest incomes in society. Income inequality at the national level has been proposed as being the underlying cause of numerous undesired outcomes in developed countries, e.g. homicide rates, and particularly undesired health care outcomes [5]. Wilkinson and Pickett have shown that higher rates of diabetes and mental illness (including drug and alcohol addiction) are associated with higher levels of income inequality at the national level [5], within developed countries (data from developing countries are unavailable to make similar comparisons). This raises the question of whether income inequality is associated with infection-related health care outcomes, such as antimicrobial resistance. A causative mechanism might be, for example, that with increased levels of disease, e.g. diabetes, there may be a corresponding increase in infections e.g. diabetic foot infections. This could result in more antimicrobial prescribing, in turn associated with higher rates of antimicrobial resistance [3]. This study therefore investigated the associations between income inequality, antimicrobial resistance and antimicrobial consumption within developed countries, via calculation of correlation coefficients between these variables.

Methods

Income inequality, antimicrobial resistance and antimicrobial consumption were correlated with each other. The analyses were completed on European countries over an eight year time period, 2003-10.

Income inequality data

Data were taken from the Standardized World Income Inequality Database (SWIID) [6]. The SWIID provides estimates of income inequality within a country which is suitable for broad cross-national analyses. The higher the SWIID score the greater a country's income inequality.

Antimicrobial resistance

Data were taken from the European Antimicrobial Resistance Surveillance Network (EARS-Net) [7]. Countries report on the susceptibility of isolates derived from blood cultures and cerebrospinal fluids only. They report data for *Enterococcus faecalis*, *Enterococcus faecium*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Streptococcus pneumoniae*.

Antimicrobial consumption

Data were taken from the European Surveillance of Antimicrobial Consumption (ESAC) group [8]. ESAC provide data on the defined daily doses (DDD) of antibiotic consumption per inhabitant per 1000 days (DID). DDD is the assumed average maintenance dose per day for a drug used for its main indication in adults, as defined by the World Health Organization [9]. Total antimicrobial consumption data (all antimicrobial classes combined) from outpatients were extracted for the period 2003-2009; consumption data for 2010 data were unavailable. It was not possible to compare a specific antibiotic's consumption to its correlated resistance as the ESAC data do not correspond directly with the EARSnet

data e.g. ESAC reports on penicillin consumption but not on aminopenicillins consumption specifically [10]. Insufficient hospital consumption data from these periods were available for analysis.

Country and year selection

Only European countries were included in the study as only these countries had comparable antimicrobial resistance data. Countries were selected from the richest 50 (by Gross National Income per capita at purchasing power parities) in 2002. Countries with a population of less than 3 million were excluded to avoid inclusion of tax havens [5]. The following 15 countries were therefore included in the study: Austria, Belgium, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Netherlands, Norway, Portugal, Spain, Sweden and the United Kingdom. Data were analysed from 2003–2010 as EARSnet data are relatively limited before 2003.

Statistical analysis

Within each antibiotic group, Spearman's rank order correlation coefficient (r) was calculated per year between income inequality (SWIID) and percentage resistance. Pooled correlations for each antibiotic group were calculated from individual year correlations within this group via the Schmidt-Hunter method [11], as well as their confidence intervals (CIs) and an associated p -value. The Schmidt-Hunter method underestimates the correlation when it is greater than 0.5 and therefore provides a conservative estimate. Where r values are > 0.8 , they are classed as strong, > 0.5 moderate, > 0.2 weak and 0 as no association [12]. The CI is given as a range of values which is likely to contain the probable correlation in the general population; a CI not containing 0 indicates sufficient evidence that there is a real correlation in the data. The p -value states how much weight of evidence there is for the observed data, when the true correlation in the population is 0; the smaller the p -value, the more likely a real correlation. These methods were repeated to provide pooled correlations between income inequality and antimicrobial consumption and antimicrobial consumption and antimicrobial resistance. EARS-Net data was incomplete, not every country reported resistance data for all pathogen-resistance combinations every year of the study. Therefore pooled correlations of antimicrobial consumption and antimicrobial resistance were calculated for the subset in which these data were available.

Scatter plots of resistance vs. income inequality by country and year were created for a selection of the antimicrobial resistance and income inequality correlations, with lines of best fit given, per year. Data were initially compiled using Microsoft Excel and were analysed using Stata/IC version 11 and StatsDirect version 3. Graphs were created in Stata/IC version 11.

Results

Correlations between income inequality, antimicrobial resistance and antimicrobial consumption within European countries are given below, and in Tables 1 and 2.

Table 1. Antimicrobial resistance associated with income inequality.

Bacterial species	Antibiotic group	No. years with data available	No. of countries for annual correlations (minimum-maximum)	Pooled correlation	95% Confidence	
					Interval	p-value
<i>E. faecalis</i>	Aminopenicillins	8	13-15	0.54	0.49 to 0.60	<0.0001
	High level gentamicin	8	13-15	0.62	0.55 to 0.69	<0.0001
	Vancomycin	8	13-15	0.73	0.68 to 0.79	<0.0001
<i>E. faecium</i>	Aminopenicillins	8	13-15	0.26	0.06 to 0.44	0.012
	High level gentamicin	8	13-15	0.38	0.17 to 0.58	<0.0001
	Vancomycin	8	13-15	0.72	0.64 to 0.80	<0.0001
<i>E. coli</i>	Aminoglycosides	8	14-15	0.84	0.80 to 0.88	<0.0001
	Aminopenicillins	8	14-15	0.73	0.70 to 0.77	<0.0001
	Carbapenems	8	9-14	0.34	0.18 to 0.49	<0.0001
	Third generation cephalosporins	8	14-15	0.70	0.68 to 0.73	<0.0001
	Fluoroquinolones	8	14-15	0.71	0.60 to 0.82	<0.0001
<i>K. pneumoniae</i>	Aminoglycosides	6	13-15	0.50	0.42 to 0.57	<0.0001
	Carbapenems	6	11-15	0.33	0.29 to 0.37	<0.0001
	Third generation cephalosporins	6	12-15	0.57	0.54 to 0.60	<0.0001
	Fluoroquinolones	6	13-15	0.50	0.42 to 0.57	<0.0001
<i>P. aeruginosa</i>	Amikacin	6	9-11	0.21	0.15 to 0.26	<0.0001
	Aminoglycosides	6	11-15	0.51	0.47 to 0.56	<0.0001
	Carbapenems	6	11-15	0.50	0.45 to 0.55	<0.0001
	Ceftazidime	6	11-15	0.51	0.47 to 0.55	<0.0001
	Fluoroquinolones	6	11-14	0.53	0.48 to 0.58	<0.0001
	Piperacillin-tazobactam	6	11-15	0.46	0.37 to 0.55	<0.0001
<i>S. aureus</i>	Methicillin	8	15	0.86	0.83 to 0.89	<0.0001
	Rifampicin	8	10-13	0.56	0.52 to 0.60	<0.0001
<i>S. pneumoniae</i>	Penicillin	8	12-14	0.34	0.25 to 0.43	<0.0001
	Macrolides	8	12-14	0.28	0.08 to 0.47	<0.0001

doi: 10.1371/journal.pone.0073115.t001

Income inequality and Enterococcal resistance

E. faecalis resistance to aminopenicillins was moderately associated with income inequality, $r=0.54$, 95% confidence interval (CI) 0.49 to 0.6. It was also moderately associated with vancomycin resistance, 0.73, 0.68 to 0.79 (Figure 1) and high level gentamicin resistance, 0.62, 0.55 to 0.69. *E. faecium* vancomycin resistance was moderately associated with income inequality, 0.72, 0.64 to 0.80. *E. faecium* resistance to high level gentamicin and aminopenicillins was weakly associated with income inequality, 0.38, 0.17 to 0.58 and 0.26, 0.06 to 0.44 respectively.

Income inequality and *E. coli* resistance

E. coli resistance to aminoglycosides was strongly associated with income inequality, $r=0.84$, 95% CI: 0.80 to 0.88. Resistance to aminopenicillins, fluoroquinolones and 3rd generation cephalosporins (Figure 2) was moderately associated with income inequality, 0.73, 0.70 to 0.77; 0.71, 0.6 to 0.82 and 0.7, 0.68 to 0.73 respectively. Resistance to carbapenems was weakly associated with income inequality, 0.34, 0.18 to 0.49.

Income inequality and *K. pneumoniae* resistance

K. pneumoniae resistance to aminoglycosides, third generation cephalosporins and fluoroquinolones (Figure 3) was moderately associated with income inequality, $r=0.50$, 95% CI: 0.42 to 0.57; 0.57, 0.54 to 0.60 and 0.50, 0.42 to 0.57 respectively. Carbapenem resistance was weakly associated with income inequality, 0.33, 0.29 to 0.37.

Income inequality and *P. aeruginosa* resistance

P. aeruginosa aminoglycoside, carbapenem, ceftazidime and fluoroquinolone resistance was moderately associated with income inequality, $r=0.51$, 95% CI: 0.47 to 0.56; 0.50, 0.45 to 0.55; 0.51, 0.47 to 0.55 and 0.53, 0.48 to 0.58 respectively. Piperacillin-tazobactam and amikacin resistance were weakly associated with income inequality, 0.46, 0.37 to 0.55 and 0.21, 0.15 to 0.26 respectively.

Income inequality and *S. aureus* resistance data

S. aureus methicillin resistance and income inequality were strongly associated, $r=0.86$, 95% CI: 0.83 to 0.89 (Figure 4). *S.*

Table 2. Antimicrobial resistance associated with antimicrobial consumption.

Bacterial species	Antibiotic group	No. years with data available	No. of countries for annual correlations		95% Confidence	
			(minimum-maximum)	Pooled correlation	Interval	p-value
<i>E. faecalis</i>	Aminopenicillins	7	13-15	0.32	0.24 to 0.39	<0.0001
	High level gentamicin	7	13-15	0.30	0.20 to 0.39	<0.0001
	Vancomycin	7	13-15	0.66	0.62 to 0.70	<0.0001
<i>E. faecium</i>	Aminopenicillins	7	13-15	-0.06	-0.21 to 0.10	0.47
	High level gentamicin	7	13-15	0.13	0.03 to 0.22	0.013
	Vancomycin	7	13-15	0.51	0.45 to 0.57	<0.0001
<i>E. coli</i>	Aminoglycosides	7	14-15	0.49	0.38 to 0.59	<0.0001
	Aminopenicillins	7	14-15	0.27	0.22 to 0.33	<0.0001
	Carbapenems	7	9-14	0.30	0.06 to 0.55	0.015
	Third generation cephalosporins	7	14-15	0.43	0.32 to 0.53	<0.0001
	Fluoroquinolones	7	14-15	0.29	0.19 to 0.38	<0.0001
<i>K. pneumoniae</i>	Aminoglycosides	5	13-15	0.84	0.75 to 0.93	<0.0001
	Carbapenems	5	11-15	0.77	0.72 to 0.81	<0.0001
	Third generation cephalosporins	5	12-15	0.85	0.80 to 0.90	<0.0001
	Fluoroquinolones	5	13-15	0.83	0.73 to 0.92	<0.0001
<i>P. aeruginosa</i>	Amikacin	5	9-11	0.83	0.73 to 0.92	<0.0001
	Aminoglycosides	5	11-15	0.89	0.87 to 0.92	<0.0001
	Carbapenems	5	11-15	0.83	0.76 to 0.90	<0.0001
	Ceftazidime	5	11-15	0.82	0.79 to 0.85	<0.0001
	Fluoroquinolones	5	11-14	0.84	0.83 to 0.86	<0.0001
<i>S. aureus</i>	Piperacillin-tazobactam	5	11-15	0.84	0.78 to 0.90	<0.0001
	Methicillin	7	15	0.67	0.65 to 0.69	<0.0001
	Rifampicin	7	10-13	0.63	0.56 to 0.69	<0.0001
<i>S. pneumoniae</i>	Macrolides	7	12-14	0.86	0.83 to 0.90	<0.0001
	Penicillin	7	12-14	0.45	0.37 to 0.52	<0.0001

doi: 10.1371/journal.pone.0073115.t002

aureus rifampicin resistance was moderately associated with income inequality, 0.56, 0.52 to 0.60.

Income inequality and *S. pneumoniae*:

S. pneumoniae resistance to penicillin and macrolides was weakly associated with income inequality, $r=0.34$, 95% CI: 0.25 to 0.43 and 0.28, 0.08 to 0.47.

Income inequality and antimicrobial consumption

Antimicrobial consumption in outpatients measured by Defined Daily Doses per 1000 inhabitants per day (DID) was weakly associated with income inequality, $r=0.44$, 95% CI: 0.41 to 0.48.

Antimicrobial consumption and antimicrobial resistance

Total antimicrobial consumption in outpatients, measured by Defined Daily Doses per 1000 inhabitants per day (DID), was strongly associated with resistance in the following species-resistance pairings: *K. pneumoniae*-aminoglycosides, $r=0.84$, CI: 0.75 to 0.93, *K. pneumoniae*-third generation

cephalosporins, 0.85, 0.80 to 0.90, *K. pneumoniae*-fluoroquinolones, 0.83, 0.73 to 0.92, *P. aeruginosa*-amikacin, 0.83, 0.73 to 0.92, *P. aeruginosa*-aminoglycosides, 0.89, 0.87 to 0.92, *P. aeruginosa*-carbapenems, 0.83, 0.76 to 0.90, *P. aeruginosa*-ceftazidime, 0.82, 0.79 to 0.85, *P. aeruginosa*-piperacillin-tazobactam, 0.84, 0.78 to 0.90 and *S. pneumoniae*-macrolides, 0.86, 0.83 to 0.90. All other bacterial-resistance pairings analysed were weakly or moderately associated with antimicrobial consumption except for *E. faecium*-aminopenicillins which was not associated with antimicrobial consumption, -0.06, -0.21 to 0.10 (Table 2).

Discussion

This study showed income inequality to be positively correlated with antimicrobial resistance in all seven species analysed: *E. faecalis*, *E. faecium*, *E. coli*, *K. pneumoniae*, *P. aeruginosa*, *S. aureus* and *S. pneumoniae*. In addition, inequality was correlated with antimicrobial consumption, and antimicrobial consumption was correlated with antimicrobial resistance.




Figure 1. Income inequality correlated with vancomycin resistance in *Enterococcus faecalis* (2003-10).

doi: 10.1371/journal.pone.0073115.g001

Strengths of the data analysed in this study include resistance data, antimicrobial consumption data and inequality data being collected by single sources: EARS-Net, ESAC and SWIID, and there being no selection bias in this study as all EARS-net data were analysed. Limitations of the data are: the data is only from developed countries within Europe and data are estimates only. The antimicrobial consumption data used consisted of total out-patient antibiotic data (all classes combined), and this study looks at in-hospital bacterial resistance to specific antibacterials. It has though been shown that out-patient antibiotic consumption is correlated with in-patient antibiotic consumption [13] and antimicrobial consumption for a specific antimicrobial has been related to resistance to that antimicrobial in previous studies [3]. The correlations between income inequality and antimicrobial resistance, seen across all pathogens studied ($n=7$) and all antimicrobials ($n=14$), are not independent of each other in all cases. This is principally an issue in *E. coli* and *K. pneumoniae* due to plasmid-mediated resistance, with plasmids carrying resistance to more than one antimicrobial class [14].

Only correlations are reported in this study and therefore no causality can be proven. Investigating if causality is present between inequality and resistance is not possible given the

data only allows us to calculate correlations, though these correlations reported are consistent with our hypothesis. It is important to consider if biologically plausible causal mechanisms support the correlations. Antimicrobial resistance rates can be increased in two ways. Firstly, exposure to antimicrobials can create resistance de novo and select out resistant strains. Secondly, transmission of antimicrobially resistant bacteria can occur. The associations between income inequality and antimicrobial consumption, and between antimicrobial consumption and antimicrobial resistance, support the former mechanism. In the case of *E. faecium*-aminopenicillin resistance, where there is no correlation between consumption and resistance, as would be expected, there was still a correlation, albeit weak, between income inequality and antimicrobial resistance. This would support the latter mechanism, i.e. that income inequality is involved in resistance through transmission of resistant bacteria, although no mediating factor analogous to antimicrobial consumption has been identified.

To support causality between inequality and consumption, mediating factors are needed, of which there are many possibilities. For example, increased income inequality is associated with higher Body Mass Index (BMI) [5]. Higher BMIs



Figure 2. Income inequality correlated with third generation cephalosporin resistance in *Escherichia coli* (2003-10).

doi: 10.1371/journal.pone.0073115.g002

are associated with comorbidities such as diabetes, resulting in higher rates of skin infections, caused by *S. aureus*, and urinary tract infections, caused by *E. coli* [15,16]. Comorbidities, e.g. obesity, could theoretically result in more severe infections. However, there is evidence that antimicrobial consumption varies independently of infection severity [17]. Given this, there could be factors not related to severity that income inequality impacts upon which affect antibiotic consumption. For example, increased income inequality is associated with lower levels of education, and lower educational levels have been associated with higher rates of antimicrobial consumption [5,18]. A defined dose of an antibiotic may also result in varying risks of developing antibiotic resistance in different countries. For example, patients with higher BMIs have lower antibiotic concentrations in the body. These lower concentrations have been implicated in the development of antibiotic resistance, sub-therapeutic drug concentrations failing to prevent the emergence of resistant bacteria [19,20]. Income inequality is also correlated with factors likely to result in an increased transmission of drug resistance e.g., imprisonment is associated with community associated-MRSA and Multi-Drug Resistant *Mycobacterium tuberculosis* [5,21,22].

Assuming there is causality between income inequality and antimicrobial resistance, the mediating factor which seems likely to be most important is considered to be antimicrobial consumption. A simple triangular model between inequality-consumption-resistance may not though reflect reality. As already demonstrated, *E. faecium* resistance is correlated with inequality even though consumption and resistance and not correlated for this antibiotic. Consideration of the relative strengths of the correlations also challenges this simple model. The correlations between income inequality and antimicrobial resistance are strongest for *S. aureus* and *E. coli*. But the strongest relationships between antimicrobial consumption and antimicrobial resistance are strongest in *K. pneumoniae* and *P. aeruginosa*. Inequality is likely to act differently on bacteria depending if they are more commonly community or hospital pathogens, for example, *S. aureus* and *E. coli* are considered common community pathogens when compared to *K. pneumoniae* and *P. aeruginosa*. Also, consumption affects different bacteria differently. For example, *Klebsiella spp* more easily develop cephalosporin resistance compared to *E. coli* due to their chromosomal carriage of *B*-lactamases [23].

Weak correlations between inequality and resistance were identified for some pathogen-antimicrobial pairings. The weak




Figure 3. Income inequality correlated with quinolone resistance in *Klebsiella pneumoniae* (2005-10).

doi: 10.1371/journal.pone.0073115.g003

association between aminopenicillin antimicrobial resistance in *E. faecium* with income inequality could be expected, as *E. faecium* is commonly resistant to aminopenicillins [24], whereas all other species are typically susceptible to the antibiotics they are tested against. *S. pneumoniae* resistance has been affected by the introduction of *S. pneumoniae* vaccination, causing a replacement of antimicrobial resistance serotypes of *S. pneumoniae* [25]. This vaccine effect may have attenuated links between *S. pneumoniae* resistance and income inequality, though no data support this suggestion. For carbapenems there may be temporal effects present, as these are the most recently introduced class of antibiotics studied.

The associations described here do not tell us how increased levels of resistance are distributed within populations as they are based on population statistics. That is, to suggest these data allow us to say if resistance is more common in the rich or poor is not possible. To do so would be to make a statistical inference error known as the "ecological fallacy". The ecological fallacy occurs when statistical data collected about a group is incorrectly used to make inferences about the nature of individuals e.g. rich or poor. There is some evidence available on who may be affected within the population. It may be that the relatively poor within a society suffer most from

increased antimicrobial resistance as their increased rates of disease e.g. diabetes, make them prone to more infections [26]. However, within less equal societies the rich suffer increased rates of disease compared to the rich within more equal societies [27]. Therefore, as income inequality increases, the rich may have relatively high rates of antibiotic resistance. This association has in fact been shown in a number of studies. In the USA, households with higher median incomes had higher rates of infection with penicillin non-susceptible pneumococci (PNSP) [28]. In Sweden, PNSP have been positively correlated with antibiotic prescribing, which has been positively correlated with per capita income [29]. In the USA study most disease was seen in the relatively poor, but resistance was higher in the relatively rich. It is likely that access to healthcare as well as the presence of an infection is important in the risk of antimicrobial resistance; a study in Ireland showed patients treated privately were prescribed antibiotics more often than those treated by the public health system [30].

The associations between income inequality and antimicrobial resistance should be considered in the context of other factors which impact on antimicrobial resistance, e.g., culture and practices of prescribers and the public, infection



Figure 4. Income inequality correlated with methicillin resistance in *Staphylococcus aureus* (2003-10).

doi: 10.1371/journal.pone.0073115.g004

control practices, intensity of migration and practices of antibiotic use in the agricultural sector [31–33]. These factors may be independently associated with antimicrobial resistance and potentially income inequality; therefore possible confounding by income inequality should always be taken into account in further antibiotic resistance research. It is likely there is a contribution from numerous factors to a country's overall level of antimicrobial resistance and that income inequality, if causality exists, is one of these factors.

Before undertaking this study, correlations between income inequality and antimicrobial resistance had not been quantified. This evidence for a positive association of income inequality and antimicrobial resistance provides the best information available at present with which to consider if increasing income inequality in developed countries does increase antimicrobial resistance; this should be considered by those researching why some countries are better at preventing and controlling the emergence of antimicrobial resistance. To investigate these associations further requires more data and these may come from two areas. Firstly, it may be possible to encourage agencies outside Europe to collect data comparable to that of the EARS-net, e.g., Centers for Disease Control (U.S.A.) could complete state-based surveillance. Secondly, it may also be

possible to monitor how the correlation between income inequality and antimicrobial resistance changes temporally within a country.

It is important for a country to know what to consider a normal level of resistance, and using income inequality correlations may allow this to be achieved. Knowing expected resistance rates is important as organisational reforms may be carried out based on crude rates, without consideration of expected rates. In the UK for example, reforms to the National Health Service (NHS) have been introduced, one of the justifications being that the UK had a higher incidence of MRSA than the European average [34]. If resistance and inequality are causally related then the UK had MRSA rates consistent with its level of income inequality and it would be incorrect to cite past UK MRSA rates in support of reforms to the UK's NHS [34].

In summary, income inequality in developed European countries is associated with antimicrobial resistance. These associations are consistent with a belief that modifying income inequality within a nation may be an effective public health intervention to reduce antimicrobial resistance rates.

Author Contributions

Conceived and designed the experiments: AK. Performed the experiments: AK AH. Analyzed the data: AK AH. Contributed

reagents/materials/analysis tools: Ak AH. Wrote the manuscript: AK AH.

References

- Rice LB (2009) The clinical consequences of antimicrobial resistance. *Curr Opin Microbiol* 12: 476-481. doi:10.1016/j.mib.2009.08.001. PubMed: 19716760.
- ECDC/EMA (2009) The bacterial challenge: time to react, Joint Technical Report from ECDC and EMA. Stockholm, Sweden: European Centre for Disease Prevention and Control.
- Goossens H, Ferech M, Vander Stichele R, Elseviers M (2005) Outpatient antibiotic use in Europe and association with resistance: a cross-national database study. *Lancet* 365: 579-587. doi:10.1016/S0140-6736(05)70799-6. PubMed: 15708101.
- Farmer P (1996) Social inequalities and emerging infectious diseases. *Emerg Infect Dis* 2: 259-269. doi:10.3201/eid0204.960402. PubMed: 8969243.
- Wilkinson RG, Pickett KE (2007) The problems of relative deprivation: why some societies do better than others. *Soc Sci Med* 65: 1965-1978. doi:10.1016/j.socscimed.2007.05.041. PubMed: 17618718.
- Solt F (2009) Standardizing the World Income Inequality Database. *Soc Sci Q* 90: 231-242. doi:10.1111/j.1540-6237.2009.00614.x.
- The European Antimicrobial Resistance Surveillance System (2009) EARSS Annual Report 2008, Bilthoven, The Netherlands
- Adriaenssens N, Coenen S, Versporten A, Muller A, Vankerckhoven V et al. (2011) European Surveillance of Antimicrobial Consumption (ESAC): quality appraisal of antibiotic use in Europe. *J Antimicrob Chemother* 66 Suppl 6: 71-77. doi:10.1093/jac/dkr129. PubMed: 22096068.
- WHO (2004) The anatomical therapeutic chemical classification system with defined daily doses (ATC/DDD). Norway.
- Adriaenssens N, Coenen S, Versporten A, Muller A, Minalu G et al. (2011) European Surveillance of Antimicrobial Consumption (ESAC): outpatient antibiotic use in Europe (1997-2009). *J Antimicrob Chemother* 66 Suppl 6: 3-12. doi:10.1093/jac/dkr119. PubMed: 22096064.
- Hunter JE, Schmidt FL (1990) *Methods of Meta-analysis: correcting error and bias in research findings*. Newbury Park, California; London; New Delhi: SAGE.
- Zou KH, Tuncali K, Silverman SG (2003) Correlation and simple linear regression. *Radiology* 227: 617-622. doi:10.1148/radiol.2273011499. PubMed: 12773666.
- Vander Stichele RH, Elseviers MM, Ferech M, Blot S, Goossens H et al. (2006) Hospital consumption of antibiotics in 15 European countries: results of the ESAC Retrospective Data Collection (1997-2002). *J Antimicrob Chemother* 58: 159-167. doi:10.1093/jac/dkl147. PubMed: 16698845.
- Paterson DL (2006) Resistance in gram-negative bacteria: Enterobacteriaceae. *Am J Infect Control* 34 Suppl 1: 20-28. doi:10.1016/j.ajic.2006.05.175.
- Roberts AD, Simon GL (2012) Diabetic foot infections: the role of microbiology and antibiotic treatment. *Semin Vasc Surg* 25: 75-81. doi:10.1053/j.semvascsurg.2012.04.010. PubMed: 22817856.
- Semins MJ, Shore AD, Makary MA, Weiner J, Matlaga BR (2012) The impact of obesity on urinary tract infection risk. *Urology* 79: 266-269. doi:10.1016/j.urol.2011.09.040. PubMed: 22130358.
- Butler CC, Hood K, Verheij T, Little P, Melbye H et al. (2009) Variation in antibiotic prescribing and its impact on recovery in patients with acute cough in primary care: prospective study in 13 countries. *BMJ* 338: b2242. doi:10.1136/bmj.b2242. PubMed: 19549995.
- Filippini M, Masiero G, Moschetti K (2006) Socioeconomic determinants of regional differences in outpatient antibiotic consumption: evidence from Switzerland. *Health Policy* 78: 77-92. doi:10.1016/j.healthpol.2005.09.009. PubMed: 16290129.
- Singh R, Tam VH (2011) Optimizing dosage to prevent emergence of resistance - lessons from in vitro models. *Curr Opin Pharmacol* 11: 453-456. doi:10.1016/j.coph.2011.07.009. PubMed: 21868287.
- Rea RS, Capitano B, Bies R, Bigos KL, Smith R et al. (2008) Suboptimal aminoglycoside dosing in critically ill patients. *Ther Drug Monit* 30: 674-681. doi:10.1097/FTD.0b013e31818b6b2f. PubMed: 19057371.
- David MZ, Daum RS (2010) Community-associated methicillin-resistant *Staphylococcus aureus*: epidemiology and clinical consequences of an emerging epidemic. *Clin Microbiol Rev* 23: 616-687. doi:10.1128/CMR.00081-09. PubMed: 20610826.
- Ruddy M, Balabanova Y, Graham C, Fedorin I, Malomanova N et al. (2005) Rates of drug resistance and risk factor analysis in civilian and prison patients with tuberculosis in Samara Region, Russia. *Thorax* 60: 130-135. doi:10.1136/thx.2004.026922. PubMed: 15681501.
- Livermore DM (1995) β -Lactamases in laboratory and clinical resistance. *Clin Microbiol Rev* 8: 557-584. PubMed: 8665470.
- Marothi YA, Agnihotri H, Dubey D (2005) Enterococcal resistance--an overview. *Indian J Med Microbiol* 23: 214-219. PubMed: 16327115.
- Liñares J, Ardanuy C, Pallares R, Fenoll A (2010) Changes in antimicrobial resistance, serotypes and genotypes in *Streptococcus pneumoniae* over a 30-year period. *Clin Microbiol Infect* 16: 402-410. doi:10.1111/j.1469-0691.2010.03182.x. PubMed: 20132251.
- Raphael D (2011) Poverty in childhood and adverse health outcomes in adulthood. *Maturitas* 69: 22-26. doi:10.1016/j.maturitas.2011.02.011. PubMed: 21398059.
- Subramanian SV, Kawachi I (2006) Whose health is affected by income inequality? A multilevel interaction analysis of contemporaneous and lagged effects of state income inequality on individual self-rated health in the United States. *Health Place* 12: 141-156. doi:10.1016/j.healthplace.2004.11.001. PubMed: 16338630.
- Chen FM, Breiman RF, Farley M, Plikaytis B, Deaver K et al. (1998) Geocoding and linking data from population-based surveillance and the US Census to evaluate the impact of median household income on the epidemiology of invasive *Streptococcus pneumoniae* infections. *Am J Epidemiol* 148: 1212-1218. doi:10.1093/oxfordjournals.aje.a009611. PubMed: 9867268.
- Nilsson P, Laurell MH (2005) Impact of socioeconomic factors and antibiotic prescribing on penicillin- non-susceptible *Streptococcus pneumoniae* in the city of Malmö. *Scand J Infect Dis* 37: 436-441. doi:10.1080/00365540510037795. PubMed: 16012003.
- Murphy M, Byrne S, Bradley CP (2011) Influence of patient payment on antibiotic prescribing in Irish general practice: a cohort study. *Br J Gen Pract* 61: e549-e555. doi:10.3399/bjgp11X593785. PubMed: 22152734.
- Rogers BA, Aminzadeh Z, Hayashi Y, Paterson DL (2011) Country-to-country transfer of patients and the risk of multi-resistant bacterial infection. *Clin Infect Dis* 53: 49-56. doi:10.1093/cid/cir273. PubMed: 21653302.
- Hulscher ME, Grol RP, van der Meer JW (2010) Antibiotic prescribing in hospitals: a social and behavioural scientific approach. *Lancet Infect Dis* 10: 167-175. doi:10.1016/S1473-3099(10)70027-X. PubMed: 20185095.
- Harada K, Asai T (2010) Role of antimicrobial selective pressure and secondary factors on antimicrobial resistance prevalence in *Escherichia coli* from food-producing animals in Japan. *J Biomed Biotechnol*, 2010: 2010: 180682. PubMed: 20589071
- UK Department of Health (2010) *Liberating the NHS: Legislative framework and next steps*. UK: London.