



UNIVERSITY OF LEEDS

This is a repository copy of *Using theory to develop and test interventions to promote changes in health behaviour: Evidence, issues, and recommendations*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/85120/>

Version: Accepted Version

Article:

Prestwich, A, Webb, TL and Conner, M (2015) Using theory to develop and test interventions to promote changes in health behaviour: Evidence, issues, and recommendations. *Current Opinion in Psychology*, 5. 1 - 5. ISSN 2352-250X

<https://doi.org/10.1016/j.copsyc.2015.02.011>

© 2014, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

This manuscript has been accepted for publication in Current Opinion in Psychology. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all disclaimers that apply to the journal apply to this manuscript. A definitive version was subsequently published in PUBLICATION, [VOL#, ISSUE#, (DATE)] DOI#"

Using theory to develop and test interventions to promote changes in health behaviour:

Evidence, issues, and recommendations

Andrew Prestwich^a

Thomas L. Webb^b

Mark Conner^a

^aSchool of Psychology, University of Leeds, Leeds, LS2 9JT, UK

^bDepartment of Psychology, University of Sheffield, Sheffield, S10 2TP, UK

Correspondence to:

Dr Andrew Prestwich

School of Psychology,

University of Leeds,

Leeds, LS2 9JT

UK

Tel: +44(0)113 343 8559

Email: a.j.prestwich@leeds.ac.uk

Abstract

Using theory to develop health behaviour interventions provides a useful framework to accumulate evidence and has been associated with larger changes in health behaviour. The present review suggests, however, that the evidence that using theory in this way produces greater health behaviour change is mixed. Furthermore, evaluating such evidence is difficult because: (i) a significant proportion of interventions are not based on theory; (ii) those that are, tend not to apply the theory extensively; (iii) theory-based and comparison groups can be confounded; and (iv) interventions may be based on multiple theories. Such difficulties delay theory evolution and the development of interventions that are capable of achieving significant and sustained changes in health behaviours. Recommendations are made to address these issues.

Introduction

Numerous theories have been applied to health behaviours^[1-3]. Theory provides a framework within which to test hypotheses and accumulate evidence, identifies constructs that influence behaviour, and suggests which techniques should be incorporated in behavioural interventions. Theory, therefore, can inform the development of interventions designed to change health-related behaviours like physical activity and healthy eating. In return, the interventions can test and refine the underlying theory, illustrating the reciprocal relationship between theory and interventions. The present review will consider the evidence regarding the effect of using theory on the efficacy of interventions designed to change health behaviour, as well as the associated issues, and recommend possible solutions.

Main text

Several reviews suggest that using theory as a basis for developing interventions generates larger changes in health behaviour than interventions that do not use theory^[4-6]. However, other reviews have provided equivocal support^[7-13], although some of these reviews were based on relatively few studies. More worryingly, some reviews suggest that theory-based interventions are less effective than interventions that are not reliant on theory^[14-16]. Several issues (see Figure 1) could explain the mixed effects of theory on health behaviour change. These issues can, however, be addressed through various solutions that can influence intervention effects directly or indirectly via the development of more appropriate theories to inform health behaviour change interventions.

Issue 1: Little use of theory or poor reporting of theory

Reviews^[13,17] have implied that a relatively high proportion of interventions appear not to use theory at all. Formal tests of this idea suggest that between 36%^[18] and 89%^[19] of health

interventions are not explicitly based on theory. Moreover, even when interventions are explicitly based on theory they often do not apply it extensively^[4,9,13,18,20,21]. For example, in a review of studies focusing on physical activity and diet, Prestwich et al.^[13] found that 90% of studies reporting a ‘theory-based’ intervention did not identify the theoretical construct(s) that was targeted by each of the behaviour change techniques (BCTs) incorporated within the intervention. It was not always clear, therefore, why some BCTs were incorporated within the intervention (see also Mama et al.^[18]). Indeed, others have noted that ‘theory-based’ interventions often employ techniques that are not consistent with the underlying theory^[22]. Many studies in Prestwich et al.’s^[13] review also failed to explicitly target all of the relevant theoretical constructs specified within a theory with a specific BCT. Therefore, only parts of the theory may have been tested rather than the theory as a whole (see also Avery et al.^[4]; Gourlan et al.^[23]). These issues could reflect little actual use of theory and/or poor reporting^[24,25] (see also Abraham, Johnson, de Bruin, & Luszczynska^[26], for associated criticisms), but either way serve to delay the refinement of theory (see issue 6).

Guidelines for the transparent reporting of the use (or non-use) of theory are needed so that it is easier to understand the implications of the effects (or lack thereof) of interventions for theory. Such guidelines also have the potential to increase the extent to which theory is used, because they draw attention to the importance of theory. Michie and Prestwich’s^[27] 19-item Theory Coding Scheme (TCS) may provide a useful basis for such an initiative. The TCS measures various features of theory use (e.g., which theoretical constructs were targeted by which specific BCTs; whether tailoring was delivered on the basis of theory; whether the intervention effects were mediated by theoretical constructs). Because reporting of the materials delivered to control groups is often sparse^[26,28], and differences between the

intervention and control groups should influence efficacy, the use of theory in the development of the materials delivered to the control group should also be clearly reported.

Issue 2: Blunt comparisons

Several reviews considering the effect of the use of theory on changes in health behaviour fail to consider how theory has informed the intervention. Instead, these reviews simply compare effect sizes between studies that explicitly report the theoretical basis for an intervention versus studies that do not^[5,29]. However, it is possible that studies purporting to base an intervention on a specific theory fail to do so fully (see issue 1) and thus conclusions regarding the impact of theory on the effectiveness of interventions are limited. Studies are needed that directly compare the effectiveness of interventions that differ in their use of theory (either in the nature of the theoretical basis, use or non-use of theory, and so on). Experimental tests comparing theory-based interventions (e.g., based on theory X vs. based on theory X + an extra construct vs. based only on the extra construct vs. control) could provide strong tests of theory and a firmer basis for their refinement^[30]. More novel methodological approaches may be required too. For example, evidence attests to non-linear, dynamic shifts in health behaviours that reflect sudden, unexpected changes in the uptake or stopping of health behaviours^[31]. These dynamic shifts might be captured by simulation approaches^[32], and/or frequent measurement of constructs and behaviour (beyond simple pre-post designs).

Issue 3: Potential confounds

Researchers who report using theory to develop and test their interventions may be more inclined to adopt stringent methodological procedures than those using atheoretical interventions, thereby confounding the comparison. For example, Diep et al.^[10] found that,

after controlling for study quality, a weak effect of interventions on vegetable intake remained (initially theory-based interventions $g_+ = 0.18$ vs. non theory-based $g_+ = 0.14$) while effects on fruit only and combined fruit and vegetable intake disappeared. Primary and secondary research should, therefore, employ strict methodological and statistical controls to account for potential confounds.

Issue 4: Combining theories

Given that using theory can render interventions more effective, it might be expected that drawing on multiple theories would lead to larger effects than drawing on a single theory. However, Diep et al.^[10] reported that the number of theories applied by a given intervention was unrelated to the effects of interventions on fruit and vegetable intake. Furthermore, there is some evidence that combining theories may actually reduce effect sizes relative to interventions based on single theories^[13,23]. As noted by Gourlan et al.^[23], many authors using interventions based on combinations of theories do not provide a rationale for their integration. Piecing theories together in this way could render the intervention atheoretical and without an evidence-base^[33]. More careful integration of theories is, therefore, required if new theories are to emerge that are better than the sum of their parts^[34-36].

Issue 5: Selecting appropriate theories for interventions

Certain theories have received extensive correlational support illustrating their ability to predict behaviour but experimental support has been more limited^[37]. Several reviews suggest that interventions based on Social Cognitive Theory (SCT^[38]), the Health Belief Model^[39], Theory of Planned Behaviour^[40] or Protection Motivation Theory^[41] can achieve small to moderate effects on health behaviours, but there is limited evidence that these changes are explained by changes in the relevant theoretical constructs^[42-45]. Other reviews

of experimental evidence^[13,46] show limited support for SCT or for the Transtheoretical Model^[47]. In addition, most popular theories (SCT, being a notable exception) do not specify how to change constructs to facilitate behaviour change^[48]. Consequently, theories that are often used to inform interventions may not be particularly suitable for this purpose.

Several steps could be taken to develop more effective theories for promoting health behaviour change. First, theories should identify evidence-based means to change the putative determinants of behaviour. Systematic reviews of techniques designed to change key determinants including self-efficacy^[49] and perceived susceptibility and worry^[16] can help to address this issue. Second, more rigorous testing and reporting of theories is required (see also Issue 2). Reviews testing the efficacy of techniques designed to change behaviour, particularly combinations of techniques, could examine the effects of techniques aligned with a specific theory. For example, Michie, Abraham, Whittington, McAteer and Gupta^[50] examined the efficacy of techniques derived from Control Theory^[51] (Carver & Scheier, 1982); namely goal-setting, monitoring and feedback (see also Dusseldorp et al. ^[52]). Third, recent research has identified more novel, but promising determinants of health behaviour and its maintenance including implicit processes^[53,54], affect^[55-57], habit^[58,59], justifications or self-licensing^[60,61] and compensatory beliefs^[62]. Further research is required to integrate these ideas within existing theories or to establish these as important theories in their own right. Fourth, theories should consider influences on behaviour beyond those at the level of the individual including dyadic or interpersonal factors such as social support^[63]. Fifth, while there is often tension between the generalisability and utility of a theory, a focus on utility should lead to larger changes in health behaviour. To achieve this, a movement towards theories that focus on specific behaviours or clusters of behaviour^[64] may be useful^[65,66].

Issue 6: Reluctance to change theories

Many theories have remained popular despite little experimental support^[36,37,48,67]. One reason is that when studies provide evidence against a theory, authors are often reluctant to suggest refining the theory^[4,13] (but see Kok & Ruiter^[68]). Such reluctance is likely to slow down the development of interventions capable of achieving significant, sustained health behaviour change. Paying attention to successful and unsuccessful replication attempts^[69] and the adoption of a system to monitor and act on the accumulated evidence is needed^[65]. This system would not discard a theory in the light of a disconfirming test but would be open to changing the theory when the evidence against it is overwhelming. As there is evidence that interventions often do not apply theory extensively^[13,18], such a system should also carefully monitor how theory was applied.

Conclusions

The evidence regarding the association between the use of theory to develop interventions and the resultant change in health behaviour is mixed. Complications include a significant proportion of interventions not extensively applying theory, risk of confounds, and theories being integrated ineffectively. Well designed experimental tests of theory, aided by widely adopted guidelines for reporting the use (and non-use) of theory in developing interventions, are needed. Careful monitoring of the evidence generated by these tests should help to refine existing theory in order to achieve more significant and sustained changes in health behaviours.

References

- ^{[1]*}Conner M, Norman P (Eds.): Predicting and changing health behaviour: Research and practice with social cognition models (3rd Edn.). Open University Press; 2015.
- This book provides a comprehensive overview of a range of theories and how they have been applied to health behaviour change interventions.
- ^[2]Davis R, Campbell R, Hildon Z, Hobbs L: **Theories of behaviour and behaviour change across the social and behavioural sciences: a scoping review**. Health Psychol Rev 2014, **8**:1-22.
- ^[3]Michie S, West R, Campbell R, Brown J, Gainforth H (Eds.): ABC of behaviour change theories. Silverback Publishing; 2014.
- ^[4]Avery KN, Donovan JL, Horwood J, Lane JA: **Behavior theory for dietary interventions for cancer prevention: A systematic review of utilization and effectiveness in creating behavior change**. Cancer Causes Control 2013, **24**:409-420.
- ^[5]Protogerou C, Johnson BT: **Factors underlying the success of behavioral HIV-prevention interventions for adolescents: A meta-review**. AIDS Behav 2014, **18**:1847-1863.
- ^[6]Webb T, Joseph J, Yardley L, Michie S: **Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy**. J Med Internet Res 2010, **12**.
- ^[7]Angus K, Cairns G, Purves R, Bryce S: **Systematic literature review to examine the evidence for the effectiveness of interventions that use theories and models of behaviour change: towards the prevention and control of communicable diseases**. Institute for Social Marketing. Insights into health communication. European Centre for Disease Prevention and Control. 2013.

- ^[8]Ayling K, Brierley S, Johnson B, Heller S, Eiser C: **Efficacy of theory-based interventions for young people with Type 1 diabetes: A systematic review and meta-analysis.** Brit J Health Psych in press.
- ^[9]Bhattarai N, Prevost AT, Wright AJ, Charlton J, Rudisill C, Gulliford MC: **Effectiveness of interventions to promote healthy diet in primary care: systematic review and meta-analysis of randomised controlled trials.** BMC Public Health 2013, **13**:1203.
- ^[10]Diep CS, Chen TA, Davies VF, Baranowski JC, Baranowski T: **Influence of behavioral theory on fruit and vegetable intervention effectiveness among children: a meta-analysis.** J Nutri Educ Behav 2014, **46**: 506-546.
- ^[11]Hill B, Skouteris H, Fuller-Tyszkiewicz M: **Interventions designed to limit gestational weight gain: a systematic review of theory and meta-analysis of intervention components.** Obes Rev 2013, **14**:435–450.
- ^[12]Lara J, Evans EH, O'Brien N, Moynihan PJ, Meyer TD, Adamson AJ, Errington L, Sniehotta, FF, White M, Mathers JC: **Association of behaviour change techniques with effectiveness of dietary interventions among adults of retirement age: a systematic review and meta-analysis of randomised controlled trials: BMC Med** 2014, **12**: 177.
- ^{[13]*}Prestwich A, Sniehotta FF, Whittington C, Dombrowski SU, Rogers L, Michie S: **Does theory influence the effectiveness of health behavior interventions? Meta-analysis.** Health Psychol 2014, **33**:465-474.

Provides an in-depth assessment of how, and the extent to which, theory may influence changes in physical activity and dietary behaviours. Relative to other reviews, it benefits from a large number of studies.

- ^[14]Gardner B, Wardle J, Poston L, Croker H: **Changing diet and physical activity to reduce gestational weight gain: a meta-analysis**. *Obes Rev* 2011, **12**:e602-620.
- ^[15]Mehtälä MAK, Sääkslahti AK, Inkinen ME: **A socio-ecological approach to physical activity interventions in childcare: a systematic review**. *IJBNPA* 2014, **11**: 22.
- ^[16]Portnoy DB, Ferrer RA, Bergman HE, Klein WMP: **Changing deliberative and affective responses to health risk: A meta-analysis**. *Health Psychol* 2013, **8**:296-318.
- ^[17]Ayers S, Olander EK: **What are we measuring and why? Using theory to guide perinatal research and measurement**. *J Reprod Infant Psych* 2013, **31**:439-448.
- ^[18]Mama SK, McNeill LH, McCurdy SA, Evans AE, Diamond PM, Adamus-Leach HJ, Lee RE: **Psychosocial Factors and Theory in Physical Activity Studies in Minorities**. *Am J Health Behav*, 2015, **39**:68-76.
- ^[19]Quirk H, Blake H, Tennyson R, Randell TL: **Physical activity interventions in children and young people with Type 1 diabetes mellitus: a systematic review with meta-analysis**. *Diabetic Med* 2014, **31**: 1163-1173.
- ^[20]Booth HP, Prevost TA, Wright AJ, Gulliford MC: **Effectiveness of behavioural weight loss interventions delivered in a primary care setting: a systematic review and meta-analysis**. *Fam Pract* 2014, **31**: 643-653.
- ^[21]West JH, Hall PC, Arredondo V, Berrett B, Guerra B, Farrell, J: **Health behavior theories in diet apps**. *J Consum Health Internet* 2013, **17**: 10-24.

^[22]van Vugt M, de Wit M, Cleijne WHJJ, Snoek FJ: **Use of behavioral change techniques in web-based self-management programs for type 2 diabetes patients: Systematic Review.** J Med Internet Res 2013, **15**:18-32.

^[23]Gourlan M, Bernard P, Bortholon C, Romain AJ, Lareyre O, Carayol M, Ninot G, Boiche J: **Efficacy of theory-based interventions to promote physical activity. A meta-analysis of randomised controlled trials.** Health Psychol Rev, in press.

^[24]Joseph RP, Daniel CL, Thind H, Benitez TJ: **Applying Psychological Theories to Promote Long-Term Maintenance of Health Behaviors** 2014, Am. J. Lifestyle Med in press.

^[25]Meader N, Semaan S, Halton M, Bhatti H, Chan M, Llewellyn A, Des Jarlais DC: **An international systematic review and meta-analysis of multisession psychosocial interventions compared with educational or minimal interventions on the HIV sex risk behaviors of people of use drugs.** AIDS Behav 2013:1963-1978.

^[26]* Abraham C, Johnson BT, de Bruin M, Luszcznska, A: **Enhancing reporting of behavior change intervention evaluations.** J Acquir Immune Defic Syndr 2014, **66**: S293-9.

The authors suggest ways in which HIV intervention studies can be better reported.

^[27]**Michie S, Prestwich A: **Are interventions theory-based? Development of a theory coding scheme.** Health Psychol 2010, **29**: 1-8.

Presents a 19-item measure that can be used reliably to indicate the extent to which theory has been applied in the development and testing of behavioural interventions.

- [28] Ayling K, Brierley S, Johnson B, Heller S, Eiser C: **How standard is standard care? Exploring control group outcomes in behaviour change interventions for young people with type 1 diabetes.** Psychol Health 2015, **30**: 85-103.
- [29] Ammerman AS, Lindquist CH, Lohr KN, Hersey J: **The efficacy of behavioral interventions to modify dietary fat and fruit and vegetable intake: a review of the evidence.** Prev Med 2002 **35**: 25–41.
- [30] Rhodes RE. **Will the new theories (and theoreticians!) please stand up? A commentary on Sniehotta, Presseau and Araújo-Soares.** Health Psychol Rev in press
- [31] Resnicow K, Page SE: **Embracing chaos and complexity: a quantum change for public health.** Am J Public Health 2008, **98**:1382-1389.
- [32] Orr MG, Plaut DC: **Complex systems and health behavior change: Insights from cognitive science.** Am J Health Behav 2014, **38**: 404-413.
- [33] Bandura A: **Health promotion from the perspective of social cognitive theory.** Psychol Health 1998, **13**:623-649.
- [34] Hagger M, Chatzisarantis NLD: **An Integrated Behavior-Change Model for physical activity.** Exerc Sport Sci Rev in press.
- [35] Pound P, Campbell R: **Exploring the feasibility of theory synthesis: A worked example in the field of health related risk-taking.** Soc Sci Med 2014, **124**:57-65.
- [36] Rhodes RE: **Improving translational research in building theory: A commentary on Head and Noar.** Health Psychol Rev 2014a, **8**: 57-60.
- [37] Schwarzer R: **Life and death of health behaviour theories.** Health Psychol Rev 2014 **8**:53-56.

- [38] Bandura A. (Ed.): **Social Foundations of Thought and Action: A Social Cognitive Theory**. Prentice-Hall; 1986.
- [39] Rosenstock IM: **Why people use health services**. *Milbank Q* 1966, **44**:94-127.
- [40] Ajzen, I: **The theory of planned behavior**. *Organ Behav Hum Dec* 1991, **50**:179-211.
- [41] Rogers RW: **Cognitive and physiological processes in fear appeals and attitude change: A revised theory of protection motivation**. In *Social psychophysiology: A sourcebook*. Edited by Cacioppo JT, Petty RE. Guilford Press 1983: 153-176.
- [42] Stacey FG, James EL, Chapman K: **A systematic review and meta-analysis of social cognitive theory-based physical activity and/or nutrition behavior change interventions for cancer survivors**. *J Cancer Surviv* in press
- [43] Jones CJ, Smith H, Llewellyn C: **Evaluating the effectiveness of health belief model interventions in improving adherence: a systematic review**. *Health Psychol Rev* 2014, **8**:253-269.
- [44] Tyson M, Covey J, Rosenthal HES: **Theory of Planned Behavior Interventions for Reducing Heterosexual Risk Behaviors: A Meta-Analysis**. *Health Psychol* 2014, **33**:1454-1467.
- [45] Bui L, Mullan B, McCaffery K: **Protection motivation theory and physical activity in the general Population: A systematic literature review**, *Psychol, Health Med* 2013, **18**: 522-42.
- [46] Conn VS, Hafdahl AR, Mehr DR: **Interventions to increase physical activity among healthy adults: Meta-analysis of outcomes**. *Am J Public Health* 2011, **101**:751–758.
- [47] Prochaska JO, Velicer WF: **The Transtheoretical Model of Health Behavior Change**. *Am J Health Promot* 1997, **12**, 38–48.

- [48] Sniehotta FF, Pesseau J, Araujo-Soares V: **Time to retire the theory of planned behaviour**. Health Psych Rev 2014, **8**:1-7.
- [49] Prestwich A, Kellar I, Parker R, MacRae S, Learmonth M, Sykes B, Taylor N, Castle H: **How can self-efficacy be increased? Meta-analysis of dietary interventions**. Health Psych Rev 2014, **8**:270-285.
- [50] Michie S, Abraham C, Whittington C, McAteer J, Gupta S: **Effective Techniques in Healthy Eating and Physical Activity Interventions: A Meta-Regression**. Health Psychol, **28**: 690-701.
- [51] Carver CS, Scheier MF: **Control theory: A useful conceptual framework for personality, social, clinical and health psychology**. Psychol Bull 1982, **92**:111-135.
- [52]**Dusseldorp E, van Genugten L, van Buuren S, Verheijden MW, van Empelen P: **Combinations of Techniques That Effectively Change Health Behavior: Evidence From Meta-CART Analysis**. Health Psychol. 2013, **33**: 1530-1540.
- This study uses an innovative statistical method to test the effect of combinations of behaviour change techniques on physical activity and dietary outcomes. Considering combinations of techniques could inform more effective interventions and provide novel tests of theory.
- [53] Keatley D, Clarke DD, Hagger MS: **The predictive validity of implicit measures of self-determined motivation across health-related behaviours**: Brit J Health Psych 2013, **18**:2-17.
- [54]*Sheeran P, Gollwitzer PM, Bargh JA. (2013). **Nonconscious processes and health**. Health Psychol, **32**, 460-473.
- The authors provide an extensive narrative overview of studies examining nonconscious processes for health behaviours. Such processes could inform novel, possibly more efficacious, health behaviour change interventions.
- [55] Allen Catellier JR, Yang ZJ: **The role of affect in the decision to exercise: Does being happy lead to a more active lifestyle?** Psychol Sport Exerc 2013, **14**:275-282.

- ^[56]Conner M, Godin G, Sheeran P, Germain M: **Some feelings are more important: Cognitive attitudes, affective attitudes, anticipated affect, and blood donation.** Health Psychol 2013, **32**:264-272.
- ^[57]Janssen E, Waters EA, van Osch L, Lechner L, de Vries H: **The importance of affectively-laden beliefs about health risks: The case of tobacco use and sun protection.** J Behav Med, **37**:11-21.
- ^[58]Gardner B: **A review and analysis of the use of 'habit' in understanding, predicting and influencing health-related behaviour.** Health Psychol Rev, in press.
- ^[59]Gardner B, Sheals, K, Wardle J: **Putting habit into practice, and practice into habit: a process evaluation and exploration of the acceptability of a habit-based dietary behaviour change intervention.** Int J Behav Nutr Phys Act 2014, **30**: 135.
- ^[60]De Witt, HJC, Evers C, De Ridder DTD: **“Because I am worth it”: A theoretical framework and empirical review of a justification-based account of self-regulation failure.** Pers Soc Psychol Rev 2014, **18**: 119-138.
- ^[61]Taylor C, Webb TL, Sheeran P: **"I deserve a treat!": Justifications for indulgence undermine the translation of intentions into action.** British J Soc Psychol 2014, **53**:501-520.
- ^[62]Berli C, Loretini P, Radtke T, Hornung R, Scholz U: **Predicting physical activity in adolescents: The role of compensatory health beliefs within the Health Action Process Approach.** Psychol Health 2014, **29**:458-474.

[63] Kaufman MR, Cornish F, Zimmerman RS: **Health behavior change models for HIV prevention and AIDS care: practical recommendations for a multi-level approach** J Acquir Immune Defic Syndr 2014, **66**:S250-8.

[64] McEachan RRC, Lawton RJ, Conner M: **Classifying health-related behaviours: Exploring similarities and differences amongst behaviours.** Brit J Health Psych 2010, **15**:347-366.

[65]**Head KJ, Noar SM: **Facilitating progress in health behaviour theory development and modification: the reasoned action approach as a case study.** Health Psychol Rev 2014, **8**: 34-52.

The authors overview a range of barriers and solutions to refining theory. Refining theory could lead to more effective interventions to change health behaviours.

[66] Noar SM, Head KJ: **Mind the gap: Bringing our theories in line with the empirical data-A response to commentaries.** Health Psychol Rev 2014, **8**:65-69.

[67] Rhodes RE: **Adding Depth to the Next Generation of Physical Activity Models.** Exercise Sport Sci R 2014, **42**:43-44.

[68] Kok G, Ruiter RAC: **Who has the authority to change a theory? Everyone! A commentary on Head and Noar.** Health Psychol Rev 2014, **8**:61-64.

[69] O'Carroll RE: **Health psychology interventions.** British J Health Psych 2014, **19**:235-239.