

This is a repository copy of *A Clifford Algebraic Framework for Coxeter Group Theoretic Computations*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/85594/>

Version: Accepted Version

Proceedings Paper:

Dechant, Pierre-Philippe orcid.org/0000-0002-4694-4010 (2014) A Clifford Algebraic Framework for Coxeter Group Theoretic Computations. In: Conference proceedings: Applied Geometric Algebras in Computer Science and Engineering 2012 (AGACSE 2012), July 2-4, La Rochelle, France. Advances in Applied Clifford Algebras . Springer , pp. 89-108.

<https://doi.org/10.1007/s00006-013-0422-4>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

A Clifford algebraic framework for Coxeter group theoretic computations

Pierre-Philippe Dechant

Abstract Real physical systems with reflective and rotational symmetries such as viruses, fullerenes and quasicrystals have recently been modeled successfully in terms of three-dimensional (affine) Coxeter groups. Motivated by this progress, we explore here the benefits of performing the relevant computations in a Geometric Algebra framework, which is particularly suited to describing reflections. Starting from the Coxeter generators of the reflections, we describe how the relevant chiral (rotational), full (Coxeter) and binary polyhedral groups can be easily generated and treated in a unified way in a versor formalism. In particular, this yields a simple construction of the binary polyhedral groups as discrete spinor groups. These in turn are known to generate Lie and Coxeter groups in dimension four, notably the exceptional groups D_4 , F_4 and H_4 . A Clifford algebra approach thus reveals an unexpected connection between Coxeter groups of ranks 3 and 4. We discuss how to extend these considerations and computations to the Conformal Geometric Algebra setup, in particular for the non-crystallographic groups, and construct root systems and quasicrystalline point arrays. We finally show how a Clifford versor framework sheds light on the geometry of the Coxeter element and the Coxeter plane for the examples of the two-dimensional non-crystallographic Coxeter groups $I_2(n)$ and the three-dimensional groups A_3 , B_3 , as well as the icosahedral group H_3 .
IPPP/12/49, DCPT/12/98

Pierre-Philippe Dechant
Institute for Particle Physics Phenomenology
Ogden Centre for Fundamental Physics
Department of Physics
University of Durham
South Road
Durham, DH1 3LE
United Kingdom
e-mail: pierre-philippe.dechant@durham.ac.uk

1 Introduction

Physical systems have to obey the mathematical laws of geometry; in particular, if they possess symmetry – such as invariance under reflections and rotations – this symmetry is heavily constrained by purely geometric considerations. For instance, many physical systems in biology (viruses), chemistry (fullerenes) and physics (crystals and quasicrystals) have polyhedral symmetries. These polyhedral symmetry groups are generated by reflections; via the Cartan-Dieudonné theorem an even number of reflections amounts to a rotation (see e.g. [17] for an exposition in a Clifford algebra context), and physical systems may be invariant only under this rotational (chiral) part, or the full reflection group.

Coxeter group theory [5, 24] axiomatises reflections from an abstract mathematical point of view. Coxeter groups thus encompass the finite Euclidean reflection groups, which include the symmetry groups of the Platonic solids – A_3 for the tetrahedron, B_3 for the dual pair octahedron and cube, and H_3 for the dual pair icosahedron and dodecahedron – as well as the Weyl groups of the simple Lie algebras. A subset of these groups are non-crystallographic, i.e. they describe symmetries that are not compatible with lattices in dimensions equal to their rank. They include the two-dimensional family of symmetry groups $I_2(n)$ of the regular polygons, as well as H_2 (the symmetry group of the decagon), H_3 (the symmetry group of the icosahedron) and the largest non-crystallographic group H_4 (the symmetry group of the hypericosahedron or 600-cell in four dimensions), which are the only Coxeter groups generating rotational symmetries of order 5. The full icosahedral group H_3 and its (chiral) rotational subgroup I are of particular practical importance, as H_3 is the largest discrete symmetry group of physical space. Thus, many 3-dimensional systems with ‘maximal symmetry’, like viruses in biology [49, 4, 51, 26, 52], fullerenes in chemistry [37, 36, 50, 38], quasicrystals in physics [27, 46, 43, 41, 48] as well as polytopes in mathematics [34, 35, 31], can be modeled using Coxeter groups.

Clifford’s Geometric Algebra [18, 15] is a complementary framework that focuses on the geometry of the physical space(-time) that we live in and its given Euclidean/Lorentzian metric. This exposes more clearly the geometric nature of many problems in mathematics and physics. In particular, Clifford’s Geometric Algebra has a uniquely simple formula for performing reflections. Previous research appears to have made exclusive use of one framework at the expense of the other. Here, we combine both paradigms, which results in geometric insights from Geometric Algebra that apparently have been overlooked in Coxeter theory thus far. This approach also has computational and conceptual advantages over standard techniques, in particular through a spinorial or conformal point of view. Hestenes [20] has given a thorough treatment of point and space groups in Geometric Algebra, and Hestenes and Holt [21] have discussed the crystallographic point and space groups from a conformal point of view. Here, we are interested in applying Geometric Algebra in the Coxeter framework, in particular in the context of root systems, the Coxeter element, non-crystallographic groups and quasicrystals, which to our knowledge have not yet been treated at all.

This paper is organised as follows. Section 2 introduces how systems are currently modeled in terms of Coxeter groups, and what kind of computations arise in this context. In Section 3, we present a versor formalism in which the full, chiral and binary polyhedral groups can all be easily generated and treated within the same framework. In particular, this yields a construction of the binary polyhedral groups (discrete subgroups of $SU(2)$ that are the double covers of the chiral (rotation) groups), which we will discuss further in Section 4. In Section 5, we briefly outline how to extend this treatment to the conformal setup, in particular for the non-crystallographic groups, and we demonstrate how to construct root systems and quasicrystalline point sets in this framework. In Section 6, we discuss the two-dimensional non-crystallographic Coxeter groups $I_2(n)$ as well as the icosahedral group H_3 and the other two three-dimensional groups A_3 (tetrahedral) and B_3 (octahedral) in a versor formalism, which elucidates the relation with the Coxeter element and the Coxeter plane. We conclude with a summary and possible further work in Section 7.

2 Coxeter formulation

Coxeter groups are abstract groups describable in terms of mirror symmetries [5]. The elements of finite Coxeter groups can be visualised as reflections at planes through the origin in a Euclidean vector space V . In particular, for $v, \alpha \in V$, then

$$v \rightarrow r_\alpha v = v' = v - \frac{2\alpha \cdot v}{\alpha \cdot \alpha} \alpha \quad (1)$$

corresponds to a Euclidean reflection r_α of the vector v at a hyperplane perpendicular to the so-called root vector α . The structure of the Coxeter group is thus encoded in the collection of all such roots, which form a root system. A subset of the root system, called the simple roots, is sufficient to express every root via a \mathbb{Z} -linear combination with coefficients of the same sign. The root system is therefore completely characterised by this basis of simple roots, which in turn completely characterises the Coxeter group. The number of simple roots is called the rank of the root system, which essentially gives the dimension and therefore indexes the corresponding Coxeter group and root system (e.g. H_3 for the largest discrete symmetry group in three dimensions).

Finite Coxeter groups describe the properties of physical structures, e.g. of a viral protein container or a carbon onion, at a given radial level, as the symmetry only relates features at the same radial distance from the origin. In order to obtain information on how structural properties at different radial levels could collectively be constrained by symmetry, affine extensions of these groups need to be considered. Affine extensions are constructed in the Coxeter framework by adding affine reflection planes not containing the origin [42]. A detailed account of this construction is presented elsewhere [44, 10, 11], but essentially the affine extension amounts to making the reflection group G topologically non-compact by adding a translation

operator T . The structures of viruses follow several different extensions of the (chiral) icosahedral group I by translation operators [28, 12, 29]. Thus, a wide range of empirical observations in virology can be explained by affine Coxeter groups. We now discuss 2D counterparts to the 3D point arrays that predict the architecture of viruses and fullerenes, and explain in what sense the translation operators are distinguished.




Fig. 1 The action of an affine Coxeter group on a pentagon. The translation operator T generates extended point arrays, whilst the compact part G makes the resulting point set rotationally symmetric. Blueprints with degeneracies due to coinciding points correspond to non-trivial group structures and can be used in the modeling of viruses.

For illustration purposes, let us consider a similar construction for a pentagon of unit size, as shown in Fig. 1. The non-compact translation operator T , here chosen to also be of unit length, creates a displaced version of the pentagon. The action of the symmetry group G of the pentagon then generates further copies in such a way that the final point array displays the same rotational symmetries.

The translation operator we have chosen for this example is distinguished because several of the generated points lie on more than one pentagon, for instance the innermost points, or the midpoints of the edges of the large outer pentagon. Certain distinguished translations lead to such point sets with degenerate points, which therefore have lower cardinality than those obtained by a random translation (here 15 points as opposed to 25). This degeneracy yields a non-trivial mathematical structure at the group level, and the corresponding blueprints in three dimensions can be used to model icosahedral viruses.

Fig. 2 shows a similar example for a translation of length of the golden ratio $\tau = \frac{1}{2}(1 + \sqrt{5}) \approx 1.618$. The resulting point set also has degenerate cardinality (now 20 points), and consists of an inner decagon and an outer pentagon. Affine symmetry here means that the relative sizes of the decagon and pentagon are fixed by the group structure. This is a powerful geometric tool for constraining real systems.

The computations necessary in this context are therefore translations, reflections and rotations; one also needs to be able to check degeneracy of points. In the usual vector space approach, these operations are implemented via matrices. We instead




Fig. 2 Translation by the golden ratio results in a point set whose constituent polygons are simultaneously constrained by the affine symmetry.

develop here a versor implementation. This has some computational advantages, as well as offering surprising geometric insights, as we shall see later. Whilst the computational complexity for 3-dimensional applications is limited, equivalent computations in four dimensions, where H_4 – the four-dimensional analogue of the icosahedral group and symmetry group of the hypericosahedron (600-cell) – has order 14,400 and H_4 -symmetric polytopes have upwards of 120 and 600 vertices, are rather more complex.

In the Coxeter setting, therefore, the reflections are fundamental; Geometric Algebra is very efficient at encoding reflections algebraically, and at performing computations with clear geometric content. However, the two frameworks do not appear to have been combined previously. We therefore explore which benefits a Clifford algebraic description might offer for Coxeter group theoretic considerations.

3 Versor framework

The geometric product $xy = x \cdot y + x \wedge y$ of two vectors x and y (with $x \cdot y$ denoting the scalar product and $x \wedge y$ the exterior product) of Geometric Algebra [18, 22, 19, 15] provides a very compact and efficient way of handling reflections in any number of dimensions, and thus by the Cartan-Dieudonné theorem also rotations [17]. For a unit vector α , the two terms in the formula for a reflection of a vector v in the hyperplane orthogonal to α from Eq. (1) simplify to the double-sided action of α via the geometric product

$$v \rightarrow r_\alpha v = v' = -\alpha v \alpha. \quad (2)$$

This prescription for reflecting vectors in hyperplanes is remarkably compact, and applies more generally to all multivectors. Even more importantly, from the Cartan-Dieudonné theorem, rotations are the product of an even number of successive reflections. For instance, compounding the reflections in the hyperplanes defined by the unit vectors α_i and α_j results in a rotation in the plane defined by $\alpha_i \wedge \alpha_j$

$$v'' = \alpha_j \alpha_i v \alpha_i \alpha_j =: \tilde{R} v R, \quad (3)$$

where we have defined the rotor $R = \alpha_i \alpha_j$ and the tilde denotes the reversal of the order of the constituent vectors of a versor, e.g. here $\tilde{R} = \alpha_j \alpha_i$. Rotors satisfy $\tilde{R}R = R\tilde{R} = 1$ and themselves transform single-sidedly under further rotations. They thus form a multiplicative group under the geometric product, called the rotor group, which is essentially the Spin group, and thus a double-cover of the special orthogonal group [18, 15, 45]. Objects in Geometric Algebra that transform single-sidedly are called spinors, so that rotors are normalised spinors.

In fact, the above two cases are examples of a more general theorem on the Geometric Algebra representation of orthogonal transformations. In analogy to the vectors and rotors above, a versor is a multivector $A = a_1 a_2 \dots a_k$ which is the product of k non-null vectors a_i ($a_i^2 \neq 0$). These versors also form a multiplicative group under the geometric product, called the versor group. The Versor Theorem [19] then states that every orthogonal transformation \underline{A} of a vector v can be expressed via unit versors in the canonical form

$$\underline{A} : v \rightarrow v' = \underline{A}(v) = \pm \tilde{A} v A, \quad (4)$$

where the \pm -sign defines the parity of the versor. Since both the versors A and $-A$ encode the same orthogonal transformation \underline{A} , unit versors are double-valued representations of the respective orthogonal transformation, giving a construction of the Pin group $\text{Pin}(p, q)$ [45], the double cover of the orthogonal group $O(p, q)$. Even versors form a double covering of the special orthogonal group $SO(p, q)$, called the Spin group $\text{Spin}(p, q)$. The versor realisation of the orthogonal group is much simpler than conventional matrix approaches. This is particularly useful in the Conformal Geometric Algebra setup, where one uses the fact that the conformal group $C(p, q)$ is homomorphic to $O(p+1, q+1)$ to treat translations as well as rotations in a unified versor framework (see Section 5), making it possible to use all of GA's versor machinery for the analysis of the conformal group.

We now consider which benefits such a versor approach can offer for Coxeter computations, in particular in the context of applications to physical phenomena in three dimensions. The isometry group of three-dimensional space is the orthogonal group $O(3)$, of which the full polyhedral (Coxeter) groups are discrete subgroups. However, $O(3)$ is globally $SO(3) \times \mathbb{Z}_2$, where the special orthogonal group $SO(3)$ is the subgroup of pure rotations (or the chiral part). $SO(3)$ is still not simply-connected, but is doubly covered by the Spin group $\text{Spin}(3) \simeq SU(2)$ (in fact, it is $SO(3) \times \mathbb{Z}_2$ locally, i.e. a fibre bundle). Thus, the chiral polyhedral groups are discrete subgroups of $SO(3)$, the full polyhedral groups (Coxeter) are their preimage

in $O(3)$, and the binary polyhedral groups are their preimage under the universal covering in $\text{Spin}(3)$.

Table 1 Versor framework for a unified treatment of the chiral, full and binary polyhedral groups.

Group	Discrete subgroup	Action Mechanism
$SO(3)$	rotational (chiral)	$x \rightarrow \tilde{R}xR$
$O(3)$	reflection (full)	$x \rightarrow \pm \tilde{A}xA$
$\text{Spin}(3)$	binary	spinors R under spinor multiplication $(R_1, R_2) \rightarrow R_1R_2$

We begin with the simple roots (vertex vectors) which completely characterise a given Coxeter group, and consider their closure under mutual reflections (the root system). We then compute the rotors derivable from all these root vectors/reflections, which encode the rotational part of the respective polyhedral group via the double-sided action in Eq. (3). The rotor group defined by single-sided action can in fact be shown to realise the respective binary polyhedral group, which is the double cover of the chiral polyhedral group under the universal covering homomorphism between $SO(3)$ and $\text{Spin}(3)$. Finally, including the versors of the form $\alpha_i\alpha_j\alpha_k$ via double-sided action gives a realisation of the full polyhedral group (the Coxeter group). The proofs are straightforward calculations in the Geometric Algebra of three dimensions and more details are contained in [8, 9].

Theorem 3.1 (Reflections/Coxeter groups and polyhedra/root systems) *Take the three simple roots for the Coxeter groups $A_1 \times A_1 \times A_1$ (respectively $A_3/B_3/H_3$). Geometric Algebra reflections in the hyperplanes orthogonal to these vectors via Eq. (2) generate further vectors pointing to the 6 (resp. 12/18/30) vertices of an octahedron (resp. cuboctahedron/cuboctahedron with an octahedron/icosidodecahedron), giving the full root system of the group.*

For instance, the simple roots for $A_1 \times A_1 \times A_1$ are $\alpha_1 = e_1$, $\alpha_2 = e_2$ and $\alpha_3 = e_3$ for orthonormal basis vectors e_i . Reflections amongst those then also generate $-e_1$, $-e_2$ and $-e_3$, which all together point to the vertices of an octahedron.

By the Cartan-Dieudonné theorem, combining two reflections yields a rotation, and Eq. (3) gives a rotor realisation of these rotations in Geometric Algebra.

Theorem 3.2 (Spinors from reflections) *The 6 (resp. 12/18/30) reflections in the Coxeter group $A_1 \times A_1 \times A_1$ (resp. $A_3/B_3/H_3$) generate 8 (resp. 24/48/120) rotors.*

For the $A_1 \times A_1 \times A_1$ example above, the spinors thus generated are ± 1 , $\pm e_1e_2$, $\pm e_2e_3$ and $\pm e_3e_1$. In fact, these groups of discrete spinors yield a novel construction of the binary polyhedral groups.

Theorem 3.3 (Spinor groups and binary polyhedral groups) *The discrete spinor group in Theorem 3.2 is isomorphic to the quaternion group Q (resp. binary tetrahedral group $2T$ /binary octahedral group $2O$ /binary icosahedral group $2I$).*

Through the versor theorem, we can therefore describe all three types of groups in the same framework. Vectors are grade 1 versors, and rotors are grade 2 versors. For instance, the 60 rotations of the chiral icosahedral group I are given by 120 rotors acting as $\alpha_i \alpha_j \vee \alpha_j \alpha_i$. 60 operations of odd parity are defined by 120 grade 1 and grade 3 versors (with vector and trivector parts) acting as $-\alpha_i \alpha_j \alpha_k \vee \alpha_k \alpha_j \alpha_i$. However, 30 of them are just the 15 true reflections given by pure vectors, leaving another 45 rotoinversions. Thus, the Coxeter group (the full icosahedral group $H_3 \subset O(3)$) is expressed in accordance with the versor theorem. Alternatively, one can think of 60 rotations and 60 rotoinversions, making $H_3 = I_h = I \times \mathbb{Z}_2$ manifest. However, the rotations operate double-sidedly on a vector, such that the versor formalism actually provides a 2-valued representation of the rotation group $SO(3)$, since the rotors R and $-R$ encode the same rotation. Since $\text{Spin}(3)$ is the universal 2-cover of $SO(3)$, the rotors form a realisation of the preimage of the chiral icosahedral group I , i.e. the binary icosahedral group $2I$. Thus, in the versor approach, we can treat all these different groups in a unified framework, whilst maintaining a clear conceptual separation. In Table 1, we summarise how the three different types of polyhedral groups are realised in the versor framework.

4 Construction of the binary polyhedral groups

In this section, we consider further the implications of our construction of the binary polyhedral groups. Since Clifford algebra is well known to provide a simple construction of the Spin groups, it is perhaps not surprising – from a Clifford algebra point of view – to find that the discrete rotor groups realise the binary polyhedral groups. However, this construction does not seem to be known, and from a Coxeter group point of view, it leads to rather surprising consequences.

rank-3 group	diagram	binary	rank-4 group	diagram
$A_1 \times A_1 \times A_1$		Q	$A_1 \times A_1 \times A_1 \times A_1$	
A_3		$2T$	D_4	
B_3		$2O$	F_4	
H_3		$2I$	H_4	

Table 2 Correspondence between the rank-3 and rank-4 Coxeter groups. The spinors generated from the reflections contained in the respective rank-3 Coxeter group via the geometric product are realisations of the binary polyhedral groups Q , $2T$, $2O$ and $2I$, which in turn generate (mostly exceptional) rank-4 groups.

The Geometric Algebra construction of the binary polyhedral groups is via rotors with (single-sided) rotor multiplication. It is then straightforward to check the group axioms, multiplication table, conjugacy classes and the representation theory.

However, it is also known that the binary polyhedral groups generate some Coxeter groups of rank 4, for instance via quaternionic root systems [8]. In particular Q , $2T$, $2O$ and $2I$ generate $A_1 \times A_1 \times A_1 \times A_1$, D_4 , F_4 and H_4 , respectively, as summarised in Table 2. From a Coxeter perspective, this is surprising. However, in Geometric Algebra, spinors ψ have a natural 4-dimensional Euclidean structure given by $\psi\tilde{\psi}$, and can thus also be interpreted as vectors in a 4D Euclidean space. In fact, one can show that these vertex vectors are again root systems [7, 9, 24], which generate the respective rank-4 Coxeter groups. This demonstrates how in fact the rank-4 groups can be derived from the rank-3 groups via the geometric product of Clifford's Geometric Algebra. This connection has so far been overlooked in Coxeter theory. This 'induction' of higher-dimensional root systems via spinors of lower-dimensional root systems is complementary to the well-known top-down approaches of projection (for instance from E_8 to H_4 [47, 43, 33, 32, 11]), or of taking subgroups by deleting nodes in Coxeter-Dynkin diagrams. It is particularly interesting that this inductive construction relates the exceptional low-dimensional Coxeter groups H_3 , D_4 , F_4 and H_4 to each other as well as to the series A_n , B_n and D_n in novel ways. In particular, it is remarkable that the exceptional dimension-four phenomena D_4 (triality), F_4 (the largest crystallographic Coxeter group in 4D) and H_4 (the largest non-crystallographic Coxeter group) are seen to arise from three-dimensional geometric considerations alone, and it is possible that their existence is due to the 'accidentalness' of the spinor construction. This spinorial view could thus open up novel applications in Coxeter and Lie group theory, as well as in polytopes (e.g. A_4), string theory and triality (D_4), lattice theory (F_4) and quasicrystals (H_4). In particular, this spinorial construction explains the symmetries of these root systems, which otherwise appear rather mysterious [9]. The $I_2(n)$ are self-dual under the corresponding two-dimensional spinor construction [7].

5 Conformal Geometric Algebra and Coxeter groups

The versor formalism is particularly powerful in the Conformal Geometric Algebra approach [22, 15, 6]. The conformal group $C(p, q)$ is 1 – 2-homomorphic to $O(p + 1, q + 1)$ [1, 2], for which one can easily construct the Clifford algebra and find rotor implementations of the conformal group action, including rotations and translations. Thus, translations can also be handled multiplicatively as rotors, for flat, spherical and hyperbolic space-times, making available the 'sandwiching machinery' of GA and simplifying considerably more traditional approaches and allowing novel geometric insight. Hestenes [20, 21] has applied this framework to point and space groups, which is fruitful for the crystallographic groups, as lattice translations can be treated on the same footing as the rotations and reflections, and this approach has helped visualise these space groups [23].

However, the non-crystallographic groups and the root system/Coxeter framework have thus far been neglected in the conformal setup. We have argued earlier and in the papers [44, 10, 11, 28, 12, 29] that in the affine extension frame-

work translations are interesting even for the non-crystallographic groups, leading to quasicrystal-like point arrays that give blueprints for viruses and other three-dimensional physical phenomena. An extension of the conformal framework to translations in the case of non-crystallographic Coxeter groups could therefore have interesting consequences, including for quasilattice theory [27, 46], in particular when quasicrystals are induced via projection from higher dimensions (e.g. via the cut-and-project method) [43, 11, 25]. We therefore briefly outline the basics of such a construction.

Let us consider the conformal space of signature $(+, +, +, +, -)$ achieved by adjoining two additional orthogonal unit vectors e and \bar{e} to the algebra of space [22]. It is therefore spanned by the unit vectors

$$e_1, e_2, e_3, e, \bar{e}, \text{ with } e_i^2 = 1, e^2 = 1, \bar{e}^2 = -1. \quad (5)$$

From these two unit vectors we can define the two null vectors

$$n \equiv e + \bar{e}, \quad \bar{n} \equiv e - \bar{e}. \quad (6)$$

One can then map a 3D vector x into the space of null vectors in the conformal space by defining

$$X \equiv F(x) := x^2 n + 2\lambda x - \lambda^2 \bar{n}. \quad (7)$$

X being null allows for a homogeneous (projective) representation of points, i.e. they are represented by a ray in the conformal space, which tends to be more numerically robust in applications, as for instance the origin is represented by \bar{n} rather than the number 0, which is sensitive to the accumulation of numerical errors. Here, λ is a fundamental length scale that is needed in order to make this expression dimensionally homogeneous, as we think of the position vector x as a dimensional quantity [39]. The equivalent notation in terms of the Amsterdam protocol would be $e = e_+$, $\bar{e} = e_-$, $n = n_\infty$ and $\bar{n} = n_0$. This notation is also consistent with the notion that the above mapping is essentially an embedding into the projective null cone of the embedding space. Originally due to Dirac [14], the idea is that the projective null cone inherits the $SO(4, 1)$ invariance of the ambient space in which $SO(4, 1)$ acts linearly, thereby endowing the projective null cone with a non-linearly realised conformal structure.

The vectors e and \bar{e} and therefore also n and \bar{n} are orthogonal to x and hence anti-commute with it, i.e. $-x^{-1}nx = n$ and $-x^{-1}\bar{n}x = \bar{n}$. Thus, the CGA implementation of a reflection $y' = -x^{-1}yx$ is given by

$$-x^{-1}F(y)x = F(y') = F(-x^{-1}yx). \quad (8)$$

Given the simple roots, one can again generate the whole root system via successive reflections as shown in Fig. 3 (left). We firstly notice that the conformal representation of a root vector $F(\alpha)$ is now different from the implementation of the reflection encoded by it via the versor α . These two roles were treated on an equal footing in 3D, as there α represents both the root vector and the versor encoding a reflection in

the hyperplane perpendicular to the root, and it is debatable whether the conceptual advantages of CGA outweigh this disadvantage.

Secondly, it is often argued that the implementation of rotations in CGA is given by $F(x') = RF(x)\tilde{R}$, since R only contains even blades and thus commutes with the vectors n and \tilde{n} such that $Rn\tilde{R} = n$ and $R\tilde{n}\tilde{R} = \tilde{n}$. However, via the Cartan-Dieudonné theorem, every rotation is given by an even number of successive reflections. Thus, it can be seen that the rotor transformation law actually follows from the more fundamental reflection law in Eq. (8). From the previous sections, we know that the spinors generated by the root vectors are important for the construction of the binary polyhedral groups and 4D polytopes. However, the 3D geometric product does not straightforwardly extend to CGA, such that the spinors and other multivectors are not treated in the same way as vectors. The operators encoding the conformal rotations, however, are still given by the 3D rotors, so that little seems to be gained by going to the conformal setup from the spinorial point of view.




Fig. 3 In the conformal setup, reflections generated by the simple roots (here e.g. α_1 and α_2 for a simple two-dimensional example, H_2) according to Eq. (8) again generate, for instance, the H_2 non-crystallographic root system, the decagon (left). CGA rotor translations via Eq. (9) act multiplicatively, but yield quasicrystalline point sets consistent with the 3D approach; for instance, on the right we show the effect of a translation with length the inverse of the golden ratio acting on a pentagon, in analogy to Figs 1 and 2.

A very salient feature of Conformal Geometric Algebra is that a translation $x \rightarrow x + a$ by a vector a is given by a rotor

$$T_a = \exp\left(\frac{na}{2\lambda}\right) = 1 + \frac{na}{2\lambda}. \quad (9)$$

It is easily checked that this has the desired effect of $T_a F(x) \tilde{T}_a = F(x + a)$, and therefore does indeed represent a 3D translation as a rotor in Conformal Geometric Algebra. One can thus treat reflections, rotations and translations multiplicatively in a unified framework. This allows for a unified construction of the type of point arrays considered earlier, and indeed the construction is entirely equivalent to the lower-dimensional construction (as it must), and can be straightforwardly verified,

for instance, for the non-crystallographic groups $I_2(n)$, H_3 and H_4 . In Fig. 3, we show an example consisting of both one such root system and one quasicrystal-like point array derived entirely in the conformal setup, as a proof of principle. The root system shown is that of H_2 , and the point array is obtained via the action of a translation of length the inverse of the golden ratio on a pentagon.

The CGA approach is naturally more computationally intensive than the 3D approach; however, this could be compensated for by increased numerical stability, as the origin is simply represented by scalar multiples of \bar{n} , as opposed to the number 0, where numerical errors can create artefacts near the origin. Treating both rotations and translations on the same footing as multiplicative rotors is also a nice conceptual shift. However, there are also drawbacks to the conformal approach. Firstly, the conformal representation of the root vectors $F(\alpha)$ is different from their action as generators of reflections α . The relationship between these two functions was more transparent in the conventional approach in 3D, where α represented both. Secondly, the rotors encoding rotations are also the 3D spinors, rather than a conformal representation of those. Thus, CGA affords a nice representation of GA vectors, but not necessarily of the whole GA multivector structure.

Following [40], an interesting approach might be to work in a curved space, for which only one extra dimension is necessary (e or \bar{e}), which should simplify the computations somewhat. One may then finally take the zero curvature limit in order to recover the Euclidean space results. For instance, for Minkowski space-time, the conformal group $C(1,3)$ is 15-dimensional. It has certain well-known ten-dimensional groups as stabiliser subgroups, i.e. groups of transformations that leave a given point (ray) y invariant. If y is spacelike, one gets an $SO(2,3)$ stabiliser subgroup, i.e. the Anti de Sitter group, corresponding to the homogeneous space-time that is the solution of Einstein's field equations with a negative cosmological constant, $\Lambda < 0$. Likewise, for timelike y one obtains the de Sitter ($\Lambda > 0$) group $SO(1,4)$ as the stabiliser (here in the CGA setup, e and \bar{e} are distinguished choices for such spacelike and timelike y). Lastly, when one chooses a null y (e.g. n), one gets an $ISO(1,3)$ subgroup, which is just the Poincaré group [45, 3]. Thus, taking the zero curvature limit essentially corresponds at the group level to the Wigner-Inönü contraction that yields the Poincaré group from the de Sitter and Anti de Sitter groups (see e.g. [16]) and a flat space (which needs two vectors e and \bar{e}) limit from a curved space (for which only one of e or \bar{e} is necessary).

6 $I_2(n)$ and H_3 – the Coxeter element and spinors

In this section, we further analyse the two-dimensional family of non-crystallographic Coxeter groups $I_2(n)$ (the symmetry groups of the regular polygons), as well as the three-dimensional groups A_3 , B_3 and the icosahedral group H_3 , describing the symmetries of the Platonic solids. A versor framework (not necessarily conformal) allows a deeper understanding of the geometry, relating spinors to the Coxeter element

and the Coxeter plane in a novel way, in particular highlighting what the complex structure involved is.

A Coxeter element $w = s_1 \dots s_n$ is the product of the reflections encoded by all the simple roots α_i of a finite Coxeter group W . The Coxeter number h is the order (i.e. $w^h = 1$) of such a Coxeter element. The sequence in which the simple reflections are performed does matter, but all such elements are conjugate, and thus the Coxeter number h is the same (for instance for $I_2(n)$ one has $h = n$). For a given Coxeter element w , there is a unique plane called the Coxeter plane on which w acts as a rotation by $2\pi/h$. At this point in the standard theory, there is a convoluted argument about the need to complexify the situation and taking real sections of the complexification in order to find the complex eigenvalues $\exp(2\pi i/h)$ and $\exp(2\pi i(h-1)/h)$ [24]. It will come as no surprise that in Geometric Algebra the complex structure arises naturally, giving a geometric interpretation for the ‘complex eigenvalues’.

Projection of a root system onto the Coxeter plane is a way of visualising any finite Coxeter group, for instance the well-known representation of E_8 is such a projection of the 240 vertices of the eight-dimensional Gosset root polytope onto the Coxeter plane. Fig. 4 (a) shows such a projection of the root polytope of H_3 (the icosidodecahedron) onto the Coxeter plane.




Fig. 4 Illustration of the geometry of H_3 . (a) shows the projection of the root polytope (the icosidodecahedron with 30 vertices) onto the Coxeter plane. Panel (b) illustrates the action of the Coxeter element on a vector $v = e_1$ denoted by the open circle in the Coxeter plane. w acts by 10-fold rotation generating a decagon clockwise, whereas on a vector n normal to the Coxeter plane it acts by reversal $-n$. Panel (c) displays both sets of vectors, which in turn happen to form the root system of $A_1 \times H_2$.

Without loss of generality, in Geometric Algebra the simple roots for $I_2(n)$ can be taken as $\alpha_1 = e_1$ and $\alpha_2 = -\cos \frac{\pi}{n} e_1 + \sin \frac{\pi}{n} e_2$ (see the H_2 root system in Fig. 3 for $n = 5$). The Cartan matrix $A_{ij} = 2\alpha_i \cdot \alpha_j / \alpha_i^2$ is then correctly given by

$$A(I_2(n)) = \begin{pmatrix} 2 & -2\cos \frac{\pi}{n} \\ -2\cos \frac{\pi}{n} & 2 \end{pmatrix}. \quad (10)$$

The Coxeter versor W describing the rotation encoded by the $I_2(n)$ Coxeter element via the typical GA half-angle formula

$$v \rightarrow wv = \tilde{W}vW \quad (11)$$

is therefore

$$W = \alpha_1 \alpha_2 = -\cos \frac{\pi}{n} + \sin \frac{\pi}{n} e_1 e_2 = -\cos \frac{\pi}{n} + \sin \frac{\pi}{n} I = -\exp\left(-\frac{\pi I}{n}\right) \quad (12)$$

for $I = e_1 e_2$. In GA it is therefore immediately obvious that the action of the $I_2(n)$ Coxeter element is described by a versor (here a rotor/spinor) that encodes rotations in the $e_1 e_2$ -Coxeter-plane and yields $h = n$ since trivially

$$W^n = (-1)^{n+1} \Rightarrow w^n = 1. \quad (13)$$

More generally, the versors belonging to conjugate Coxeter elements could be $W = \pm \exp\left(\pm \frac{\pi I}{n}\right)$ and one immediately finds that $W^n = \pm 1$ as required for w to be of order $h = n$.

Since $I = e_1 e_2$ is the bivector defining the plane of e_1 and e_2 , it anticommutes with both e_1 and e_2 . Thus, in the half-angle formula Eq. (11), one can take W through to the left to write the complex eigenvector equation

$$v \rightarrow wv = \tilde{W}vW = \tilde{W}^2 v = \exp\left(\pm \frac{2\pi I}{n}\right)v, \quad (14)$$

immediately yielding the standard result for the complex eigenvalues. However, in GA it is now obvious that the complex structure is in fact given by the bivector describing the Coxeter plane (trivial for $I_2(n)$), and that the standard complexification is both unmotivated and unnecessary. The ‘complex eigenvalues’ are simply left and right going spinors in the Coxeter rotation plane.

The Pin group/eigenblade description in GA therefore yields a wealth of novel geometric insight and the general case will be the subject of a future publication. However, for instance for the icosahedral group H_3 , standard theory has $h = 10$ and complex eigenvalues $\exp(2\pi mi/h)$ with the exponents $m = \{1, 5, 9\}$. For simple roots $\alpha_1 = e_2$, $-2\alpha_2 = (\tau - 1)e_1 + e_2 + \tau e_3$ and $\alpha_3 = e_3$, one finds the Coxeter plane bivector $B_C = e_1 e_2 + \tau e_3 e_1$. Under the action of the Coxeter element versor $2W = -\tau e_2 - e_3 + (\tau - 1)I$ (here $I = e_1 e_2 e_3$) it gets reversed $-\tilde{W}B_C W = -B_C$ as is expected for an invariant plane under an odd operation. For an ‘eigenvector’ in the Coxeter plane, the two-dimensional argument from Eq. (14) applies and one again finds eigenvalues $\exp\left(\pm \frac{2\pi B_C}{h}\right)$, which corresponds to $m = 1$ and $m = 9$. In fact, this holds true for a general Coxeter group: 1 and $h - 1$ are always exponents and in Geometric Algebra they correspond to ‘eigenvectors’ being rotated in the Coxeter plane via left and right going spinors. However, in Geometric Algebra it is also obvious that in general more complicated geometry is at work, with different complex structures corresponding to different eigenspaces. Going back to our H_3

example, for the vector $b_C = B_C I = -\tau e_2 - e_3$ orthogonal to the Coxeter plane, one has $-\tilde{W}b_C W = -b_C = \exp\left(\pm \frac{5 \cdot 2\pi B_C}{h}\right)b_C$, as is expected for the normal vector for a plane that gets reversed. Thus, in GA this case straightforwardly corresponds to $m = 5$, accounting for the remaining case.

Fig. 4 illustrates this H_3 geometry. Panel (a) shows the projection of the root system, the icosidodecahedron, onto the Coxeter plane. The vector $v = e_1$ lies in the Coxeter plane and the Coxeter element w acts on it by 10-fold rotation via the Coxeter versor W . This is depicted in Panel (b), where v is denoted by the open circle, and rotation via W occurs in the clockwise direction creating a decagon. The Coxeter versor acts on the vector b_C normal to the Coxeter plane simply by reversal, as discussed above. Both sets of vectors (the decagon and \pm the normal) are depicted in Panel (c). Curiously, these vectors form the root system of $A_1 \times H_2 = A_1 \times I_2(5)$.

The geometry for A_3 and B_3 is very similar. They have Coxeter numbers $h = 4$ and $h = 6$, respectively, and exponents $m = \{1, 2, 3\}$ and $m = \{1, 3, 5\}$. The Coxeter versor again inverts the Coxeter bivector, and the exponents 1 and $h - 1$ correspond to left and right going rotations in the Coxeter plane on which the Coxeter element acts by h -fold rotation, whilst the normal to the Coxeter plane gets simply inverted as expected, corresponding to the cases $h/2$ ($m = 2$ and $m = 3$ for A_3 and B_3 , respectively). Again, the combinations of the vectors in the plane and orthogonal to it form the root systems of $A_1 \times A_1 \times A_1 = A_1 \times I_2(2)$ and $A_1 \times A_2 = A_1 \times I_2(3)$.

7 Conclusions

We have investigated what insight a Geometric Algebra description, which lends itself to applications of reflections, can offer when applied to the Coxeter (reflection) group framework. The corresponding computations are conceptually revealing, both for applications to real systems and for purely mathematical considerations. The implementation of orthogonal transformations as versors rather than matrices offers some computational and conceptual advantages, in both the conventional and the conformal approaches. The main benefit in a versor description of the applications, for instance in virology, lies in the simple construction and implementation of the chiral and full polyhedral groups. The Clifford approach then also yields a simple construction of the binary polyhedral groups, and in fact all three groups can be straightforwardly treated in the same framework. This seemingly unknown construction of the binary polyhedral groups also sheds light on the fact why they generate Coxeter groups of rank 4. The natural 4D Euclidean structure of the spinors allows for an alternative interpretation as vectors (in fact, a root system) in a 4D space, which generate Coxeter groups in four dimensions. Thus, one can construct many four-dimensional (exceptional) Lie and Coxeter groups from three-dimensional considerations alone. We have constructed non-crystallographic root systems and groups, as well as quasicrystalline point arrays in the conformal framework. This could be interesting for the latter quasicrystals, as translations (e.g. arising from affine extensions of the Coxeter groups) are treated multiplicatively by

versors in the same way as rotations and reflections. We have discussed the versor framework for the groups $I_2(n)$, A_3 , B_3 and H_3 , in particular in relation to the Coxeter element, the Coxeter plane and complex eigenvalues/exponents. The Geometric Algebra approach gives novel geometric insight, as the complex structure is seen to arise from the Coxeter plane bivector, and the Coxeter element acts as a spinor generating rotations in this Coxeter plane.

We are currently applying the more formal considerations of our recent work to extending the existing paradigm for modeling virus and fullerene structure [12] and to packing problems [30]. The chiral and binary polyhedral groups are attractive as discrete symmetry groups for flavour and neutrino model building in particle physics, and we are currently working on an anomaly analysis (breaking of classical symmetries by quantum effects) for these groups [13]. The two-dimensional groups $I_2(n)$ generate the symmetries of protein oligomers, which we are currently investigating.

Acknowledgements I would like to thank my family and friends for their support, my former PhD supervisor Anthony Lasenby for getting me interested in and teaching me GA (amongst many other things), as well as David Hestenes, Eckhard Hitzer, Joan Lasenby, Mike Hobson, Reidun Twarock, Céline Böhm, Christoph Luhn and Silvia Pascoli for helpful discussions. Many thanks also to the referees, who have led to many valuable improvements in the manuscript.

References

1. Pierre Anglès. Construction de revêtements du groupe conforme d'un espace vectoriel muni d'une métrique de type (p, q) . *Annales de l'institut Henri Poincaré (A) Physique théorique*, 33(1):33, 1980.
2. Pierre Anglès. *Conformal Groups In Geometry And Spin Structures*. Progress in Mathematical Physics. Birkhäuser, 2008.
3. James Emory Baugh. *Regular Quantum Dynamics*. PhD thesis, Georgia Institute of Technology, 2004.
4. D. L. D. Caspar and A. Klug. Physical principles in the construction of regular viruses. *Cold Spring Harbor Symp. Quant. Biol.*, 27:1–24, 1962.
5. H. S. M. Coxeter. Discrete groups generated by reflections. *Ann. of Math*, 35:588–621, 1934.
6. Pierre-Philippe Dechant. *Models of the Early Universe*. PhD thesis, University of Cambridge, UK, 2011.
7. Pierre-Philippe Dechant. Rank-3 root systems induce root systems of rank 4 via a new Clifford spinor construction. *ArXiv e-print 1207.7339*, 2012.
8. Pierre-Philippe Dechant. Clifford algebra unveils a surprising geometric significance of quaternionic root systems of Coxeter groups. *Advances in Applied Clifford Algebras*, 23:301–321, June 2013. 10.1007/s00006-012-0371-3.
9. Pierre-Philippe Dechant. Platonic solids generate their four-dimensional analogues. *to appear in Acta Crystallographica A*, 2013.
10. Pierre-Philippe Dechant, Céline Böhm, and Reidun Twarock. Novel Kac-Moody-type affine extensions of non-crystallographic Coxeter groups. *Journal of Physics A: Mathematical and Theoretical*, 45(28):285202, 2012.
11. Pierre-Philippe Dechant, Céline Böhm, and Reidun Twarock. Affine extensions of non-crystallographic Coxeter groups induced by projection. *accepted by Journal of Mathematical Physics*, 2013.

12. Pierre-Philippe Dechant, Céline Böhm, and Reidun Twarock. Applications of affine extensions of non-crystallographic Coxeter groups in carbon chemistry and virology. *in preparation*, 2013.
13. Pierre-Philippe Dechant, Christoph Luhn, Céline Böhm, and Silvia Pascoli. Discrete anomalies of chiral and binary polyhedral groups and their implications for neutrino and flavour model building. *in preparation*, 2013.
14. P. A. M. Dirac. Wave equations in conformal space. *The Annals of Mathematics*, 37(2):pp. 429–442, 1936.
15. Chris Doran and Anthony N. Lasenby. *Geometric Algebra for Physicists*. Cambridge University Press, Cambridge, 2003.
16. P. G. O. Freund. *Introduction to Supersymmetry*. Cambridge University Press, Cambridge, April 1988.
17. D. J. H. Garling. *Clifford Algebras: An Introduction*. London Mathematical Society Student Texts. Cambridge University Press, 2011.
18. David Hestenes. *Space-Time Algebra*. Gordon and Breach, New York, 1966.
19. David Hestenes. *New foundations for classical mechanics; 2nd ed.* Fundamental theories of physics. Kluwer, Dordrecht, 1999.
20. David Hestenes. *Point Groups and Space Groups in Geometric Algebra*, pages 3–34. Birkhäuser, Boston, 2002.
21. David Hestenes and Jeremy W. Holt. The Crystallographic Space Groups in Geometric Algebra. *Journal of Mathematical Physics*, 48:023514, 2007.
22. David Hestenes and Garret Sobczyk. *Clifford algebra to geometric calculus: a unified language for mathematics and physics*. Fundamental theories of physics. Reidel, Dordrecht, 1984.
23. Eckhard Hitzer and Christian Perwass. Interactive 3D space group visualization with CLUCalc and the Clifford Geometric Algebra description of space groups. *Advances in Applied Clifford Algebras*, 20:631–658, 2010. 10.1007/s00006-010-0214-z.
24. J. E. Humphreys. *Reflection groups and Coxeter groups*. Cambridge University Press, Cambridge, 1990.
25. Giuliana Indelicato, Paolo Cermelli, David Salthouse, Simone Racca, Giovanni Zanzotto, and Reidun Twarock. A crystallographic approach to structural transitions in icosahedral viruses. *Journal of Mathematical Biology*, pages 1–29, 2011. 10.1007/s00285-011-0425-5.
26. A. Janner. Towards a classification of icosahedral viruses in terms of indexed polyhedra. *Acta Crystallographica Section A*, 62(5):319–330, 2006.
27. A. Katz. *Some local properties of the 3-dimensional Penrose tilings, an introduction to the mathematics of quasicrystals*. Academic Press, 1989.
28. T. Keef and R. Twarock. Affine extensions of the icosahedral group with applications to the three-dimensional organisation of simple viruses. *J Math Biol*, 59(3):287–313, 2009.
29. T. Keef, J. Wardman, N.A. Ranson, P. G. Stockley, and R. Twarock. Structural constraints on the 3d geometry of simple viruses. *Acta Crystallographica Section A*, 69:140–150, 2013.
30. Tom Keef, Pierre-Philippe Dechant, and Reidun Twarock. Packings of solids with non-crystallographic symmetry. *in preparation*, 2013.
31. M. Koca, M. Al-Ajmi, and S. Al-Shidhani. Quasi-regular polyhedra and their duals with Coxeter symmetries represented by quaternions ii. *The African Review of Physics*, 6(0), 2011.
32. M. Koca, R. Koc, and M. Al-Barwani. Noncrystallographic Coxeter group H_4 in E_8 . *Journal of Physics A: Mathematical and General*, 34:11201–11213, dec 2001.
33. M. Koca, N. O. Koca, and R. Koç. Quaternionic roots of E_8 related Coxeter graphs and quasicrystals. *Turkish Journal of Physics*, 22:421–436, May 1998.
34. Mehmet Koca, Mudhahir Al-Ajmi, and Ramazan Koç. Polyhedra obtained from Coxeter groups and quaternions. *Journal of Mathematical Physics*, 48(11):113514, 2007.
35. Mehmet Koca, Nazife Ozdes Koca, and Ramazan Koç. Catalan solids derived from three-dimensional root systems and quaternions. *Journal of Mathematical Physics*, 51(4):043501, 2010.
36. H. Kroto. Carbon onions introduce new flavour to fullerene studies. *Nature*, 359:670–671, 1992.

37. H. Kroto, J. R. Heath, S. C. O'Brien, R. F. Curl, and R. E. Smalley. C60: Buckminsterfullerene. *Nature*, 318:162–163, 1985.
38. E. F. Kustov, V. I. Nefedov, A. V. Kalinin, and G. S. Chernova. Classification system for fullerenes. *Russian Journal of Inorganic Chemistry*, 53(9):1384–1395, 2008.
39. A N Lasenby, Joan Lasenby, and Richard Wareham. A covariant approach to geometry using Geometric Algebra. *Technical Report. University of Cambridge Department of Engineering, Cambridge, UK*, 2004.
40. Anthony N. Lasenby. Recent applications of Conformal Geometric Algebra. In Hongbo Li, Peter J. Olver, and Gerald Sommer, editors, *Computer Algebra and Geometric Algebra with Applications: 6th International Workshop, IWMM 2004, Shanghai, China, May 19-21, 2004*, volume 3519 of *Lecture Notes in Computer Science*, pages 298–328. Springer Berlin / Heidelberg, Secaucus, NJ, USA, 2005.
41. L.S. Levitov and J. Rhyner. Crystallography of quasicrystals; application to icosahedral symmetry. *J. Phys. France*, 49(49):1835–1849, 1988.
42. Jon McCammond and T. Petersen. Bounding reflection length in an affine Coxeter group. *Journal of Algebraic Combinatorics*, pages 1–9. 10.1007/s10801-011-0289-1.
43. R. V. Moody and J. Patera. Quasicrystals and icosians. *Journal of Physics A: Mathematical and General*, 26(12):2829, 1993.
44. J. Patera and R. Twarock. Affine extensions of noncrystallographic Coxeter groups and quasicrystals. *Journal of Physics A: Mathematical and General*, 35:1551–1574, 2002.
45. Ian R. Porteous. *Clifford Algebras and the Classical Groups*. Cambridge University Press, Cambridge, 1995.
46. M. Senechal. *Quasicrystals and Geometry*. Cambridge University Press, 1996.
47. O. P. Shcherbak. Wavefronts and reflection groups. *Russian Mathematical Surveys*, 43(3):149, 1988.
48. D. Shechtman, I. Blech, D. Gratias, and J.W. Cahn. Metallic phase with long-range order and no translational symmetry. *Phys. Rev. Lett.*, 53:1951–1953, 1984.
49. P. G. Stockley and R. Twarock. *Emerging Topics in Physical Virology*. Imperial College Press, 2010.
50. R. Twarock. New group structures for carbon onions and carbon nanotubes via affine extensions of noncrystallographic Coxeter groups. *Phys. Lett. A*, 300:437–444, 2002.
51. R. Twarock. Mathematical virology: a novel approach to the structure and assembly of viruses. *Phil. Trans. R. Soc.*, (364):3357–3373, 2006.
52. R. Zandi, D. Reguera, R. F. Bruinsma, W. M. Gelbart, and J. Rudnick. Origin of icosahedral symmetry in viruses. *Proc. Natl. Acad. Sci.*, 101(44):15556–15560, 2004.