



This is a repository copy of *Loss of plakoglobin promotes decreased cell-cell contact, increased invasion, and breast cancer cell dissemination in vivo*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/86806/>

Version: Supplemental Material

Article:

Holen, I., Whitworth, J., Nutter, F. et al. (6 more authors) (2012) Loss of plakoglobin promotes decreased cell-cell contact, increased invasion, and breast cancer cell dissemination in vivo. *Breast Cancer Research*, 14. R86 . ISSN 1465-5411

<https://doi.org/10.1186/bcr3201>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

ERRATUM

Open Access



Erratum to: Loss of *plakoglobin* promotes cell-cell contact, increased invasion and breast cancer cell dissemination in vivo

Ingunn Holen¹, Jacob Whitworth¹, Faith Nutter¹, Alyson Evans¹, Hannah K. Brown¹, Diane V. Lefley¹, Ivana Barbaric², Mark Jones² and Penelope D. Ottewell^{1,3*}

Erratum

After the publication of this study [1] an error was detected in Fig. 2e. The same image was accidentally used for beta-catenin staining of MCF7 2A-1 and T47D 2A-4. This error does not affect the findings or conclusions of the article. The corrected figure is shown below and we apologise for this mistake.

Author details

¹Academic Unit of Clinical Oncology, Beech Hill Road, Sheffield, UK. ²Centre for Stem Cell Research, Biomedical Sciences, Western Bank, University of Sheffield, Sheffield, UK. ³Academic Unit of Clinical Oncology, CR-UK/YCR Sheffield Cancer Research Centre, University of Sheffield, Sheffield S10 2RX, UK.

Received: 6 March 2017 Accepted: 6 March 2017

Published online: 28 March 2017

Reference

1. Holen I, Whitworth J, Nutter F, Evans A, Brown HK, Lefley DV, Barbaric I, Jones M, Ottewell PD. Loss of plakoglobin promotes decreased cell-cell contact, increased invasion and breast cancer cell dissemination in vivo. *Breast Cancer Res.* 2012;14(3):R86.

* Correspondence: P.D.Ottewell@shef.ac.uk

¹Academic Unit of Clinical Oncology, Beech Hill Road, Sheffield, UK

³Academic Unit of Clinical Oncology, CR-UK/YCR Sheffield Cancer Research Centre, University of Sheffield, Sheffield S10 2RX, UK






Fig. 2 Shows relative expression of *plakoglobin* and *e-cadherin* compared with GAPDH \pm SEM before and after siRNA knockdown with (a) scramble sequence or miRNA cassette 2 in MCF7 cells, (b) scramble sequence or miRNA cassette 3 in MCF7 cells, (c) scramble sequence or miRNA cassette 2 in T47D cells. **d** Are Western blots showing gamma catenin and E-cadherin expression following transfection with scramble sequence or miRNA cassettes 2 and 3. **e** Shows immunohistochemical staining for γ -catenin, e-cadherin and beta-catenin (green). In the control cells γ -catenin, e-cadherin and β -catenin are expressed on the cell surface clearly demarcating the cell-cell junctions. In the knock down lines, γ -catenin staining is reduced and e-cadherin and β -catenin is detected in the nucleus and the cytoplasm and β -catenin