



This is a repository copy of *Advances in proteomics for production strain analysis*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/87279/>

Version: Supplemental Material

Article:

Landels, A., Evans, C., Noirel, J. et al. (1 more author) (2015) *Advances in proteomics for production strain analysis*. *Current Opinion in Biotechnology*, 35. 111 - 117. ISSN 0958-1669

<https://doi.org/10.1016/j.copbio.2015.05.001>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Figure 1

A selection from a rank-plot of the 200 most frequently used words in abstracts of production-strain proteomics publications, ranked by frequency. Words in blue are higher-ranked in production-strain proteomics than in proteomics in general; words in red are lower ranked, words in black have not changed relative position. Faded words have changed rank by 5 places or fewer and words in bold-face are only present in the production-strain list. The solid line indicates the change in rank. This figure presents a snapshot of the full list, which is available in the supplementary material.