



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/88243/>

Version: Accepted Version

---

**Conference or Workshop Item:**

Valavanis, A, Dean, P, Burnett, AD et al. Terahertz frequency imaging for explosives and weapons detection. In: Innovative Research Call in Explosives and Weapons Detection (2007) Showcase, 17 Nov 2011, London, United Kingdom.

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# Terahertz frequency imaging for explosives and weapons detection

A. Valavanis, P. Dean, A. D. Burnett, S. Chowdhury, A. Marshall, A. G. Davies and E. H. Linfield

School of Electronic and Electrical Engineering, University of Leeds, Leeds LS2 9JT  
email: a.valavanis@leeds.ac.uk

The terahertz (THz) region of the electromagnetic spectrum lies between microwave and infrared frequencies, and has many potential benefits for security imaging. In particular, terahertz radiation can provide spectral signatures of explosive compounds, illicit drugs and metallic objects, whilst passing through many common packaging and clothing materials with little attenuation. The relatively short wavelength also allows high-resolution imaging of features smaller than  $100\ \mu\text{m}$ . We have developed high power ( $> 50\ \text{mW}$ ) and frequency-tunable sources operating in the 1–5 THz spectral range, and used these in two demonstrations of terahertz imaging:

**Spectroscopic diffuse-reflectance imaging:** We have shown that a common explosive (PETN) can be distinguished from a simple sugar (lactose monohydrate) at THz frequencies (Fig. 1(a)).[1] We are also able to detect signatures of powdered solids at a 1.5 m distance by measuring the back-scattered radiation (Fig. 1(b)). This technique offers several advantages over the state of the art. Firstly, thick and/or highly-absorbing samples can be probed, unlike the case of transmission imaging. Secondly, there is no need for precise detector alignment, unlike specular-reflection imaging. Furthermore, we have undertaken measurements through smooth containers, with no deterioration in signal.

**Self-mixing imaging:** We have removed the need for any external radiation detector by coupling reflected radiation from an object back into the laser cavity.[2] This interference-sensing technique has allowed us to demonstrate high-resolution imaging and sensing at distances in excess of 7 m, and to obtain height-sensing of concealed objects (Fig. 1(c)).

Our two demonstrations illustrate the potential of THz imaging technology as a potential ‘fieldable’ compact system, and we are now seeking industrial partners to develop our research into demonstrator systems for targeted applications.

This project is funded under the Innovative Research Call in Explosives and Weapons Detection (2007), a cross-government programme sponsored by a number of government departments and agencies under the CONTEST strategy.

[1] P. Dean et al., *Opt. Exp.* **17**, 20631 (2009)

[2] P. Dean et al., *Opt. Lett.* **36**, 2587 (2011)




Figure 1: (a) Difference-frequency images at 3.05 and 3.24 THz illustrating contrast between PETN high-explosive and lactose monohydrate [left and right, respectively]. (b) Diffuse-reflectance image at 2.8 THz revealing a bag of powdered PTFE concealed within a visually-opaque envelope. (c) A 2.6 THz self-mixing image of a concealed scalpel blade. Contours in the image reveal the surface-height.