



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/88245/>

Version: Accepted Version

Conference or Workshop Item:

Valavanis, A, Dean, P, Burnett, AD et al. (2011) Spectroscopic diffuse-reflection imaging at a 1.5 m stand-off distance, using a frequency-switchable terahertz quantum cascade laser. In: ITQW 2011: The 11th International Conference on Intersubband Transitions in Quantum Wells, 11-17 Sep 2011, Badesi, Sardinia, Italy.

This is an author produced version of a paper presented in ITQW 2011: The 11th International Conference on Intersubband Transitions in Quantum Wells.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Spectroscopic diffuse-reflection imaging at 1.5 m stand-off, using a frequency-switchable terahertz quantum cascade laser

A. Valavanis, P. Dean, A. D. Burnett, S. Chowdhury, A. G. Davies and E. H. Linfield

Institute of Microwaves and Photonics, School of Electronic and Electrical Engineering,
University of Leeds, Leeds LS2 9JT, United Kingdom
email: a.valavanis@leeds.ac.uk

Numerous potential applications exist for imaging at terahertz (THz) frequencies, within fields including medicine, security, manufacturing quality inspection and chemical sensing. Imaging diffuse reflections from rough or powdered objects offers several advantages over specular-reflection or transmission imaging configurations. Objects of arbitrary thickness may be used, and precise alignment of collection optics is not required, as diffuse reflections spread over a large solid-angle. Furthermore, commonly-used smooth packaging materials have little effect upon diffuse reflection images. Many substances exhibit strong, material-specific spectral responses to THz radiation and spectroscopic THz imaging has been demonstrated in transmission and specular-reflection geometries using both time-domain and frequency-domain techniques. However, spectroscopic imaging of the relatively-weak diffuse reflections is challenging, as a high-intensity multi-frequency THz radiation source is required, along with a sensitive THz detector.

In this work, we demonstrate four-frequency diffuse-reflection imaging using a frequency-switchable THz quantum cascade laser (QCL) as a radiation source. This QCL contains a heterogeneous active region, which may be switched between single-mode emissions at 3.05, 3.21, 3.28 and 3.35 THz by adjusting the bias voltage.