

This is a repository copy of *Khovanov homotopy types and the Dold-Thom functor*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/90031/>

Version: Submitted Version

Article:

Everitt, Brent orcid.org/0000-0002-0395-338X, Lipshitz, Robert, Sarkar, Sucharit et al. (1 more author) (2016) Khovanov homotopy types and the Dold-Thom functor. *Homology, Homotopy and Applications*. pp. 177-181. ISSN: 1532-0081

<https://doi.org/10.4310/HHA.2016.v18.n2.a9>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Khovanov homotopy types and the Dold-Thom functor

Brent Everitt, Robert Lipshitz, Sucharit Sarkar and Paul Turner^{*}

Abstract. We show that the spectrum constructed by Everitt and Turner as a possible Khovanov homotopy type is a product of Eilenberg-MacLane spaces and is thus determined by Khovanov homology. By using the Dold-Thom functor it can therefore be obtained from the Khovanov homotopy type constructed by Lipshitz and Sarkar.

A *Khovanov homotopy type* is a way of associating a (stable) space to each link L so that the classical invariants of the space yield the Khovanov homology of L . There are two recent constructions of Khovanov homotopy types, using different techniques and giving different results [3, 6]. In [3] homotopy limits were employed to build an Ω -spectrum $\mathbf{X}_\bullet L = \{X_k(L)\}$ with the following properties:

- (i). the homotopy type is a link invariant, and
- (ii). the homotopy groups are Khovanov homology:

$$\pi_i(\mathbf{X}_\bullet(L)) = Kh^{-i}(L).$$

The main goal of this note is to prove the following result.

Theorem 1. *Each of the spaces $X_k(L)$ is homotopy equivalent to a product of Eilenberg-MacLane spaces.*

In [6] the programme of Cohen, Jones and Segal [2] was generalized to produce a suspension spectrum $\mathcal{X}_{Kh}(L)$ with the following properties:

- (i). the homotopy type is a link invariant, and
- (ii). the reduced cohomology is Khovanov homology:

$$\tilde{H}^i(\mathcal{X}_{Kh}(L)) = Kh^i(L).$$

As a corollary we obtain that $\mathbf{X}_\bullet(L)$ is homotopy equivalent to the infinite symmetric product of $\mathcal{X}_{Kh}(L)$.

To prove Theorem 1 we use the explicit model, due to McCord [8], of the Eilenberg-MacLane spaces. Given a monoid G and a based topological space X , let $B(G, X)$ denote the set of maps

BRENT EVERITT: Department of Mathematics, University of York, York YO10 5DD, United Kingdom. e-mail: brent.everitt@york.ac.uk. ROBERT LIPSHITZ, SUCHARIT SARKAR: Department of Mathematics, Columbia University, New York NY 10027, USA. e-mail: lipshitz@math.columbia.edu, e-mail: sucharit@math.columbia.edu. PAUL TURNER: Section de mathématiques, Université de Genève, 2-4 rue du Lièvre, CH-1211, Geneva and Département de mathématiques, Université de Fribourg, Chemin du musée, CH-1700 Fribourg, Switzerland. e-mail: pri.maths@gmail.com.

^{*} RL was supported by an NSF grant number DMS-0905796 and a Sloan Research Fellowship. SS was supported by a Clay Foundation Postdoctoral Fellowship. PT was partially supported by Swiss National Science Foundation grant 200021-131967.

$u: X \rightarrow G$ such that $u(x) = 0$ for all but finitely many $x \in X$. Then $B(G, X)$ is a monoid, and if G is a group (the case of interest) then $B(G, X)$ is a group. Moreover, when G is an abelian topological group the set $B(G, X)$ can be topologized in a natural way so that the group operation is continuous. This construction has nice functoriality: letting \mathbf{Ab} , \mathbf{Top}_* and \mathbf{AbTop} denote respectively the categories of abelian groups, based topological spaces and topological abelian groups, one has the following result.

Proposition 1. [8, Proposition 6.7] *McCord's construction is a bifunctor*

$$B(-, -): \mathbf{Ab} \times \mathbf{Top}_* \rightarrow \mathbf{AbTop}.$$

Furthermore, as special case of [8, Theorem 11.4], for an abelian group G the space $B(G, S^n)$ is the Eilenberg-MacLane space $K(G, n)$. Thus we may take as *the* Eilenberg-MacLane space functor:

$$B(-, S^n): \mathbf{Ab} \rightarrow \mathbf{AbTop}.$$

Conversely, the following is [4, Corollary 4K.7, p. 483] (apparently originally due to Moore; cf. [8, p. 295]):

Proposition 2. *A path-connected, commutative topological monoid is a product of Eilenberg-MacLane spaces.*

The spaces $X_k(L)$ are built as homotopy limits of diagrams of spaces. Recall that given a small category \mathbf{C} and a (covariant) functor $D: \mathbf{C} \rightarrow \mathbf{Top}_*$ (a diagram), that $\text{holim}_{\mathbf{C}} D$ is constructed as follows (see, e.g., [1, Section 11.5] or the concise notes [9, Section 3.7]). Consider the product

$$\prod_{\sigma \in N(\mathbf{C})} \text{Hom}(\Delta^n, D(c_n)) = \prod_{n \geq 0} \prod_{\substack{\alpha_1 \rightarrow \dots \rightarrow \alpha_n \\ \alpha_i \neq \text{Id}}} \text{Hom}(\Delta^n, D(c_n)) \quad (1)$$

where $N(\mathbf{C})$ is the subset of the nerve of \mathbf{C} consisting of all sequences of composable morphisms $\sigma = (c_0 \xrightarrow{\alpha_1} c_1 \xrightarrow{\alpha_2} \dots \xrightarrow{\alpha_n} c_n)$ in which none of the morphisms are identity maps, and Hom denotes the space of continuous maps from the standard n -simplex. The homotopy limit $\text{holim}_{\mathbf{C}} D$ is the subspace of this product consisting of those tuples $(f_\sigma)_{\sigma \in N(\mathbf{C})}$ such that the following diagrams commute:

$$\begin{array}{ccc} \Delta^{n-1} & \xrightarrow{f_{d_i \sigma}} & D(c_n) \\ d^i \downarrow & & \downarrow \text{Id} \\ \Delta^n & \xrightarrow{f_\sigma} & D(c_n) \end{array} \quad (2)$$

for each $0 < i < n$, and

$$\begin{array}{ccc} \Delta^{n-1} & \xrightarrow{f_{d_0 \sigma}} & D(c_n) \\ d^0 \downarrow & & \downarrow \text{Id} \\ \Delta^n & \xrightarrow{f_\sigma} & D(c_n) \end{array} \quad \text{and} \quad \begin{array}{ccc} \Delta^{n-1} & \xrightarrow{f_{d_n \sigma}} & D(c_{n-1}) \\ d^n \downarrow & & \downarrow D(\alpha_n) \\ \Delta^n & \xrightarrow{f_\sigma} & D(c_n) \end{array} \quad (3)$$

corresponding to the cases $i = 0$ and $i = n$, respectively. Here the map d^i denotes the i^{th} face inclusion, $d_i \sigma = (c_0 \xrightarrow{\alpha_1} \dots \xrightarrow{\alpha_{i-1}} c_{i-1} \xrightarrow{\alpha_{i+1} \alpha_i} c_{i+1} \dots \xrightarrow{\alpha_n} c_n)$ when $0 < i < n$, and d_0, d_n similarly.

The following is well-known, but for completeness we give its (short) proof.

Proposition 3. *Let $D: \mathcal{C} \rightarrow \text{Top}_*$ be a diagram of topological abelian groups and continuous group homomorphisms. Then the homotopy limit of D is a topological abelian group.*

Proof. Pointwise addition makes the set $\text{Hom}(\Delta^n, D(c_n))$ into an abelian group, and the product in formula (1) is the product (topological abelian) group. It remains to see that the diagrams (2) and (3) describe a subgroup of this product. Suppose that tuples (f_σ) and (g_σ) make these diagrams commute. Then the first two diagrams automatically commute for the pointwise sum $(f_\sigma + g_\sigma)$. The third diagram for the pointwise sum becomes,

$$\begin{array}{ccccccc} \Delta^{n-1} & \longrightarrow & \Delta^{n-1} \times \Delta^{n-1} & \xrightarrow{f_{d_n\sigma} \times g_{d_n\sigma}} & D(c_{n-1}) \times D(c_{n-1}) & \xrightarrow{+} & D(c_{n-1}) \\ d^n \downarrow & & \downarrow d^n \times d^n & & \downarrow D(\alpha_n) \times D(\alpha_n) & & \downarrow D(\alpha_n) \\ \Delta^n & \longrightarrow & \Delta^n \times \Delta^n & \xrightarrow{f_\sigma \times g_\sigma} & D(c_n) \times D(c_n) & \xrightarrow{+} & D(c_n) \end{array}$$

for which the first square obviously commutes, the second commutes since f and g are in the prescribed subspace and the third commutes from the fact that $D(\alpha_n)$ is a group homomorphism. The inverse operation is similarly seen to be closed, hence the subspace defined above is a subgroup. \square

Proof (Proof of Theorem 1). Let L be an oriented link diagram with c negative crossings. The space $X_k(L)$ is constructed as follows. Let I denote the category with objects $\{0, 1\}$ and a single morphism from 0 to 1, and I^n the product of I with itself n times. Let $\bar{0}$ be the initial object in I^n , and let \mathbf{P} be the result of adjoining one more object to I^n and a single morphism from the new object to every object except $\bar{0}$.

In [3] it is shown that there is a functor $F: \mathbf{P} \rightarrow \text{Ab}$ such that the i^{th} derived functor of the inverse limit, $\varprojlim_{\mathbf{P}}^i F$, is isomorphic to the i^{th} unreduced Khovanov homology of L . The space $X_k(L)$ is constructed by composing this functor with the Eilenberg-MacLane space functor $K(-, k+c)$ and taking the homotopy limit of the resulting diagram of spaces.

We may now use the explicit model for Eilenberg-MacLane spaces given by McCord. By applying Proposition 1 we define a diagram $D: \mathbf{P} \rightarrow \text{AbTop}$ as the composition

$$\mathbf{P} \xrightarrow{F} \text{Ab} \xrightarrow{B(-, S^{k+c})} \text{AbTop}.$$

By the homotopy invariance property of the homotopy limit construction we have

$$X_k(L) \simeq \text{holim}_{\mathbf{P}} D.$$

By Proposition 3, the homotopy limit on the right is itself a topological abelian group, and hence, by Proposition 2, a product of Eilenberg-MacLane spaces. \square

Corollary 1. *The homotopy type of $\mathbf{X}_\bullet(L)$ is determined by $Kh(L)$.*

The spectrum $\mathcal{X}_{Kh}(L) = \{\mathcal{X}_{Kh}^{(k)}(L)\}$ constructed in [6] has the additional property that the cellular cochain complex of the space $\mathcal{X}_{Kh}^{(k)}(L)$ is isomorphic to the Khovanov complex of L (up to shift). It follows from the description of the Khovanov homology of the mirror image (see [5]) that

$$\tilde{H}_i(\mathcal{X}_{Kh}(L)) = Kh^{-i}(-L)$$

where $-L$ denotes the mirror of L . The infinite symmetric product $\mathrm{Sym}^\infty \mathcal{X}_{Kh}^{(k)}(L)$ is seen from the Dold-Thom theorem to be

$$\mathrm{Sym}^\infty \mathcal{X}_{Kh}^{(k)}(L) = \prod_n K(\tilde{H}_n(\mathcal{X}_{Kh}^{(k)}(L)), n)$$

from which we have the following.

Corollary 2. *For large enough k , the space $X_k(-L)$ is homotopy equivalent to the infinite symmetric product $\mathrm{Sym}^\infty \mathcal{X}_{Kh}^{(k)}(L)$.*

We end by noting that the analogue of Theorem 1 for the spectra $\mathcal{X}_{Kh}(L)$ is not true. For all alternating knots $\mathcal{X}_{Kh}(L)$ is a wedge of Moore spaces [6], however there are examples of non-alternating knots for which $\mathcal{X}_{Kh}(L)$ is not a wedge of Moore spaces (see [7]).

Acknowledgements

We thank Tyler Lawson for several helpful suggestions, including communicating Proposition 3 to us.

References

- [1] A. K. Bousfield and D. M. Kan, *Homotopy limits, completions and localizations*, Lecture Notes in Mathematics, Vol. 304, Springer-Verlag, Berlin, 1972. MR0365573 (51 #1825)
- [2] R. L. Cohen, J. D. S. Jones, and G. B. Segal, *Floer's infinite-dimensional Morse theory and homotopy theory*, The Floer memorial volume, 1995, pp. 297–325. MR1362832 (96i:55012)
- [3] Brent Everitt and Paul Turner, *The homotopy theory of Khovanov homology*. arXiv:1112.3460.
- [4] Allen Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002. MR1867354 (2002k:55001)
- [5] Mikhail Khovanov, *A categorification of the Jones polynomial*, Duke Math. J. **101** (2000), no. 3, 359–426. MR1740682 (2002j:57025)
- [6] Robert Lipshitz and Sucharit Sarkar, *A Khovanov homotopy type or two*. arXiv:1112.3932.
- [7] ———, *Some Steenrod squares on Khovanov homology*. In preparation.
- [8] M. C. McCord, *Classifying spaces and infinite symmetric products*, Trans. Amer. Math. Soc. **146** (1969), 273–298. MR0251719 (40 #4946)
- [9] Rubén Sánchez-García, *Homotopy limits and colimits*. www.math.uni-duesseldorf.de/~sanchez/.