

This is a repository copy of *Decaying and non-decaying badly approximable numbers*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/90994/>

Version: Submitted Version

Article:

Broderick, Ryan, Fishman, Lior and Simmons, David orcid.org/0000-0002-9136-6635
(2016) Decaying and non-decaying badly approximable numbers. *Acta Arithmetica*. pp. 143-152. ISSN: 1730-6264

<https://doi.org/10.4064/aa8281-10-2016>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

DECAYING AND NON-DECAYING BADLY APPROXIMABLE NUMBERS

RYAN BRODERICK, LIOR FISHMAN, AND DAVID SIMMONS

ABSTRACT. We call a badly approximable number *decaying* if, roughly, the Lagrange constants of integer multiples of that number decay as fast as possible. In this terminology, a question of Y. Bugeaud ('15) asks to find the Hausdorff dimension of the set of decaying badly approximable numbers, and also of the set of badly approximable numbers which are not decaying. We answer both questions, showing that the Hausdorff dimensions of both sets are equal to one. Part of our proof utilizes a game which combines the Banach–Mazur game and Schmidt’s game, first introduced in Fishman, Reams, and Simmons (preprint '15).

1. INTRODUCTION

Fix $d \in \mathbb{N}$. For each $\mathbf{x} \in \mathbb{R}^d$, write

$$\begin{aligned} L(\mathbf{x}) &= \liminf_{q \rightarrow \infty} q^{1/d} d(q\mathbf{x}, \mathbb{Z}^d) \\ &= \sup\{c \geq 0: \|\mathbf{x} - \mathbf{p}/q\| \geq c/q^{1+1/d} \text{ for all but finitely many } (\mathbf{p}, q) \in \mathbb{Z}^d \times \mathbb{N}\}, \end{aligned}$$

where $\|\cdot\|$ denotes some norm on \mathbb{R}^d (e.g. the max norm). The point \mathbf{x} is called *well approximable* if $L(\mathbf{x}) = 0$, and *badly approximable* if $L(\mathbf{x}) > 0$. If \mathbf{x} is badly approximable and $i/j \in \mathbb{Q}$, we can get a trivial bound on $L(\frac{i}{j}\mathbf{x})$ in terms of $L(\mathbf{x})$: if $L(\mathbf{x}) > c$, then for all but finitely many $(\mathbf{p}, q) \in \mathbb{Z}^d \times \mathbb{N}$

$$\left\| \frac{i}{j}\mathbf{x} - \frac{\mathbf{p}}{q} \right\| = \frac{i}{j} \left\| \mathbf{x} - \frac{j\mathbf{p}}{iq} \right\| \geq \frac{i}{j} \frac{c}{(iq)^{1+1/d}} = \frac{c}{i^{1/d}j} \frac{1}{q^2}$$

and thus

$$(1.1) \quad L\left(\frac{i}{j}\mathbf{x}\right) \geq \frac{1}{i^{1/d}j} L(\mathbf{x}).$$

The special case $j = 1$ gives the bound

$$(1.2) \quad L(i\mathbf{x}) \geq L(\mathbf{x})/i^{1/d}.$$

Recently, Y. Bugeaud [3] has investigated the question of whether the bound (1.2) is “optimal”. Precisely, call a badly approximable point \mathbf{x} *decaying* if there exists $C > 0$ such that for all $i \in \mathbb{N}$,

$$L(i\mathbf{x}) \leq C/i^{1/d}.$$

In this terminology, the main result of [3] states that certain algebraic badly approximable points are decaying. Bugeaud also asks the following question:

Question 1.1 ([3, Problem 4.4]). What is the Hausdorff dimension of the set of decaying badly approximable¹ points? of the set of badly approximable points which are not decaying?

The latter question has been recently answered in dimension 1 by a preprint of D. Badziahin and S. Harrap [1, Theorem 15].² In this paper, we give a different proof of Badziahin–Harrap’s result, which is valid in higher dimensions as well. Our proof shows that the set of badly approximable points which are

¹The condition “badly approximable” has been omitted in the statement of [3, Problem 4.4], but the question obviously makes no sense without it.

²In the current version of their paper they do not mention this connection; we informed them of it by private communication while writing this paper.

not decaying is hyperplane winning, which shows that its intersection with certain “nice” fractals has full Hausdorff dimension; see §2 for more details.

We also answer Bugeaud’s first question in dimension 1, showing that the set of decaying badly approximable numbers has full dimension. Our proof uses the so-called Banach–Mazur–Schmidt game recently introduced in [4].

Acknowledgements. The second-named author was supported in part by the Simons Foundation grant #245708. The third-named author was supported in part by the EPSRC Programme Grant EP/J018260/1. We wish to thank Jayadev Athreya for introducing us to a special case of the theorems proved in this paper which motivated this work.

1.1. Main results. In our first main theorem, we find a full dimension set of badly approximable points which are not only non-decaying, but non-decaying when multiplied by any chosen sequence of rational numbers, with as little decay in the Lagrange values as desired. (The full dimension set depends on the chosen sequence and decay rate.) Precisely:

Theorem 1.2. *Let $\frac{i_1}{j_1}, \frac{i_2}{j_2}, \dots \in \mathbb{Q}$ be a sequence of distinct rational numbers, and let $g : \mathbb{N} \rightarrow (0, \infty)$ satisfy $g(k) \rightarrow \infty$ as $k \rightarrow \infty$. Then the set*

$$\text{BA}_d \setminus \left\{ \mathbf{x} \in \text{BA}_d : \limsup_{k \rightarrow \infty} g(k) L\left(\frac{i_k}{j_k} \mathbf{x}\right) = \infty \right\}$$

is hyperplane winning.

This theorem is quite similar to [1, Theorem 15], with three main differences:

- We allow any sequence of rationals $(\frac{i_k}{j_k})_k$, rather than just the sequence of all natural numbers.
- We prove the theorem in any dimension, rather than just dimension 1.
- We prove that the set in question is hyperplane winning, whereas Badziahin and Harrap show that it is Cantor-winning. These concepts share many similar properties and both imply full Hausdorff dimension, but the former is stronger [1, Theorem 12] and implies strong C^1 incompressibility on sufficiently regular fractals (see [2]).

The special case $\frac{i_k}{j_k} = k$, $g(k) = k^{1/d}$ yields an answer to the latter part of Question 1.1, namely it shows that the set of badly approximable points which are not decaying is hyperplane winning, and therefore of full Hausdorff dimension.

We remark that this theorem also answers a related question. Namely, fix an irrational $i/j \in \mathbb{Q}$ and consider the sequence $i_k/j_k = (i/j)^k$. By (1.1) we have $L((i/j)^k \mathbf{x}) \geq \frac{1}{(i^{1/d}j)^k} L(\mathbf{x})$, so it is natural to ask about the size of the set of \mathbf{x} for which

$$(1.3) \quad \lim_{k \rightarrow \infty} (i^{1/d}j)^k L((i/j)^k \mathbf{x}) = \infty.$$

Taking $i_k/j_k = (i/j)^k$ and any $g(k) \in o((i^{1/d}j)^k)$, Theorem 1.2 implies that for a full-dimension set of $\mathbf{x} \in \text{BA}_d$ there exists a sequence $(k_n)_n$ such that $L((i/j)^{k_n} \mathbf{x}) \geq \frac{1}{g(k_n)}$ for all n . Fix such an \mathbf{x} ; applying (1.1) one then obtains $L((i/j)^k \mathbf{x}) \geq (g(k_n)(i^{1/d}j)^{k-k_n})^{-1}$ for all $k_n \leq k < k_{n+1}$. It follows easily from this that (1.3) holds for this \mathbf{x} . Thus, the set of \mathbf{x} satisfying (1.3) has full Hausdorff dimension in \mathbb{R}^d . As a concrete example, there is a full-dimension set of $x \in \mathbb{R}$ such that $L(2^n x) \in \omega(2^{-n})$.

On the other hand, in dimension one we show that the set of x for which $L(\frac{i}{j}x)$ decays quickly also has full dimension, providing a complement to both of the above results. Namely, we obtain full dimension for the set of x for which $\lim_{k \rightarrow \infty} (ij)^k L((\frac{i}{j})^k x) < \infty$, as well as full dimension of the set of decaying badly approximable numbers, via the following theorem:

Theorem 1.3. *The set $\left\{ x \in \text{BA}_1 : L\left(\frac{i}{j}x\right) \leq \frac{1}{ij} \text{ for all reduced } \frac{i}{j} \in \mathbb{Q} \right\}$ has full Hausdorff dimension in \mathbb{R} .*

Since the set in question is contained in the set of decaying badly approximable numbers, Theorem 1.3 answers the first part of Question 1.1 by showing that the set of decaying badly approximable points has full dimension in \mathbb{R} .

2. GAMES

In this section we describe two variants of Schmidt's game which will be used in the following sections. Both games are played by two players, whom we will call Alice and Bob. The *hyperplane absolute game* (or just *hyperplane game*) was introduced in [2] and involves a parameter $0 < \beta < 1/3$ and a target set $S \subseteq \mathbb{R}^d$. Bob begins by choosing a ball $B_0 = B(\mathbf{x}_0, \rho_0) \in \mathbb{R}^d$. Inductively, if $B_i = B(\mathbf{x}_i, \rho_i)$ have been chosen for $i = 1, \dots, k$, Alice chooses a hyperplane \mathcal{L}_{k+1} and a number $0 < \varepsilon_{k+1} \leq \beta$, determining a set $A_{k+1} = \mathcal{N}(\mathcal{L}, \varepsilon_{k+1}\rho_k) = \{\mathbf{x} \in \mathbb{R}^d : d(\mathbf{x}, \mathcal{L}) \leq \varepsilon_{k+1}\rho_k\}$. Bob then must choose a ball $B_{k+1} = B(\mathbf{x}_{k+1}, \rho_{k+1}) \subseteq B_k \setminus A_{k+1}$ satisfying $\rho_{k+1} = \beta\rho_k$.³ The sets $B_0 \supseteq B_1 \supseteq B_2 \supseteq \dots$ are closed and nested with radii tending to zero, so the intersection $B_\infty = \bigcap_{k=0}^\infty B_k$ is a singleton. The unique point \mathbf{x} such that $B_\infty = \{\mathbf{x}\}$ is called the *result* of the game. If $\mathbf{x} \in S$, Alice is declared the winner; otherwise, Bob wins. If Alice has a strategy to win this game regardless of how Bob plays for each $0 < \beta < 1/3$, then S is called *hyperplane absolute winning*, or just *hyperplane winning*. Hyperplane winning sets intersect any sufficiently regular fractal in a set of full dimension, and the class of hyperplane winning sets is closed under diffeomorphisms and countable intersections. Combining these three properties one obtains *strong C^1 -incompressibility* of hyperplane winning sets on “nice” fractals (see [2] for precise definition and discussion):

Theorem 2.1 ([2, Corollary 5.4]). *Let S be a hyperplane winning subset of \mathbb{R}^d , let K be the support of an absolutely decaying and Ahlfors regular measure, and let $f_i : U \rightarrow \mathbb{R}^d$ ($i \in \mathbb{N}$) be nonsingular C^1 maps. Then*

$$\dim \left(\bigcap_{i=1}^\infty f^{-i}(S) \cap K \right) = \dim(K).$$

In particular, $\dim(S) = d$.

Many self-similar fractals are supports of absolutely decaying and Ahlfors regular measures so we obtain for example that when S is a hyperplane winning subset of \mathbb{R} and C is the Cantor ternary set, there is a set of dimension $\dim(C)$ consisting of real numbers $x \in C$ such that x, x^2, x^3, \dots are all in S .

The *Banach–Mazur–Schmidt game*, or *BMS game*, was introduced in [4]. It is played on a complete metric space (X, d) with a target set S and parameter $0 < \beta < 1$ being given. Bob again begins by choosing an arbitrary ball $B_0 = B(x_0, \rho_0)$. Now suppose B_0, B_1, \dots, B_k have been chosen and write $B_i = B(x_i, \rho_i)$. Alice then chooses a ball $A_{k+1} = B(y_{k+1}, \beta\rho_k)$ satisfying $d(x_k, y_{k+1}) + \beta\rho_k \leq \rho_k$. (Note that this implies $A_{k+1} \subseteq B_k$.) Bob then chooses $B_{k+1} = B(x_{k+1}, \rho_{k+1})$ satisfying $d(x_{k+1}, y_{k+1}) + \rho_{k+1} \leq \beta\rho_k$. Here Bob is allowed to choose $0 < \rho_{k+1} \leq \rho_k$ arbitrarily. Again, the balls B_i ($i \in \mathbb{N}$) are closed and nested with radii tending to zero, so the intersection $B_\infty = \bigcap_k B_k$ is a singleton. Again, the *result* x is the unique point such that $B_\infty = \{x\}$, and we call Alice the winner if and only if $x \in S$. If Alice has a strategy to win this game regardless of how Bob plays, we say that S is β -BMS winning. If S is β -BMS winning for some $0 < \beta < 1$ then we say that S is *BMS winning*. Informally, the BMS game is a two-player game in which the first player (Bob) chooses balls according to the relatively lax rules of the Banach–Mazur game, while the second player (Alice) chooses according to the stricter rules of Schmidt's game. Accordingly, the class of BMS winning sets is very restrictive and has strong geometric properties. To make this precise, we need the following definition:

Definition 2.2. Given $\beta > 0$, a set $E \subseteq X$ is said to be *uniformly β -porous* if there exists $r_0 > 0$ such that for every ball $B(x, r) \subseteq X$ with $r \leq r_0$, there exists an open ball $B^\circ(y, \beta r) \subseteq B(x, r)$ such that $B^\circ(y, \beta r) \cap E = \emptyset$.

Theorem 2.3 ([4, Theorem 2.2]). *Let (X, d) be a complete, separable metric space and fix $0 < \beta < 1$. Then a Borel set $S \subseteq X$ is β -BMS winning if and only if $X \setminus \bigcap_{i=1}^\infty S_i$ is a countable union of uniformly β -porous sets.*

³The original hyperplane game only requires Bob's radius to satisfy $\rho_{k+1} \geq \beta\rho_k$ rather than equality, making the game discussed here the “modified hyperplane game”. However, the hyperplane game is equivalent to the modified hyperplane game (this follows from appropriately modifying the proof of [5, Proposition 4.5]), so this distinction is not important.

When (X, d) is given additional structure, this theorem yields an estimate on the Hausdorff measure of a BMS winning set. Specifically, we consider metric spaces (X, d) which support a measure of the following type:

Definition 2.4. A Borel measure on a metric space (X, d) is called *Ahlfors s -regular* if there exist $C > 1$ and $\rho_0 > 0$ such that for any $x \in X$ and $0 < \rho \leq \rho_0$ we have

$$(2.1) \quad \frac{1}{C} \rho^s \leq \mu(B(x, \rho)) \leq C \rho^s.$$

We say μ is Ahlfors regular if it is Ahlfors s -regular for some $s > 0$.

If (X, d) supports some Ahlfors s -regular Borel measure, then the s -dimensional Hausdorff measure on X is Ahlfors s -regular as well [8, Theorem 8.5]. It follows that $\dim X = s$. On a metric space (X, d) supporting such a measure, BMS winning sets are very large in the sense that their complement has positive codimension:

Proposition 2.5 ([4, Corollary 2.5]). *If X is Ahlfors s -regular and $S \subseteq X$ is BMS winning, then $\dim(X \setminus S) < s$.*

It follows from this that any BMS winning set on a metric space supporting an Ahlfors s -regular measure must be conull with respect to the s -dimensional Hausdorff measure.

3. PROOF OF THEOREM 1.2

We will need the following lemma due to Davenport (see [7, Lemma 4]) for a version which implies the one stated here):

Lemma 3.1 (Simplex Lemma). *Let $d \in \mathbb{N}$, $r > 0$, and $\mathbf{x} \in \mathbb{R}^d$. Then the rational points in $B(\mathbf{x}, r)$ with denominator at most $(2d!r)^{-\frac{d}{d+1}}$ all lie in a single hyperplane.*

Proof of Theorem 1.2. Fix $0 < \beta < 1/3$. Let $\rho_0 > 0$ be the radius of B_0 . Since $g(k) \rightarrow \infty$, after extracting a subsequence we can assume without loss of generality that $\beta^{2^k+1}/(4d!) \geq g(k)^{-1/2}$ for all k . Write $\varepsilon_k = g(k)^{-1/2}$. Alice will play the game in such a way that she guarantees that if \mathbf{x} denotes the result of the game, i.e. the unique point so that $\bigcap_{j \geq 0} B_j = \{\mathbf{x}\}$, then for each $k \in \mathbb{N}$, $\left\| \frac{i_k}{j_k} \mathbf{x} - \frac{\mathbf{p}}{q} \right\| \geq \varepsilon_k q^{-1+1/d}$ for all sufficiently large q . She will do this by partitioning \mathbb{N} into infinitely many arithmetic progressions $P_k = 2^{k-1} + 2^k \mathbb{N}_0$ ($k \in \mathbb{N}$) and using her turns corresponding to indices in P_k to ensure that the condition holds for k . (This argument mimics the original proof of the countable intersection stability of Schmidt's winning property.) This ensures that $L\left(\frac{i_k}{j_k} \mathbf{x}\right)$ is larger than ε_k .

Fix $k \in \mathbb{N}$ and $\ell \in P_k$, and we will describe Alice's strategy for the turn corresponding to the index ℓ . Let

$$Q_m = \left(4d! \frac{i_k}{j_k} \beta^\ell \rho_0 \right)^{-\frac{d}{d+1}}$$

and let

$$\mathcal{Q}_m = \left\{ \frac{j_k \mathbf{p}}{i_k q} : Q_{m-1} \leq q < Q_m \right\}.$$

Note that every (reduced) $\frac{j_k \mathbf{p}}{i_k q}$ with q sufficiently large is in some \mathcal{Q}_m (where ‘‘sufficiently large’’ depends on k). Since $\ell \in P_k$, we have $\ell = 2^{k-1} + 2^k m$ for some $m \geq 0$. Given $B_\ell = B_{2^{k-1}+2^k m}$, we consider $\mathcal{Q}_m \cap 2B_\ell$, where $2B_\ell$ denotes the ball with the same center as B_ℓ and twice the radius. Applying Lemma 3.1 with $r = 2(i_k/j_k)\beta^\ell \rho_0$, we see that all rationals in a given ball of radius r of the form $\frac{\mathbf{p}}{q}$ with $Q_{m-1} \leq q < Q_m$ lie on a single hyperplane. Now, the map $\mathbf{x} \mapsto \frac{j_k}{i_k} \mathbf{x}$ sends balls of radius $2(i_k/j_k)\beta^\ell \rho_0$ to balls of radius $2\beta^\ell \rho_0$ and also sends hyperplanes to hyperplanes, so it follows that $\mathcal{Q}_m \cap 2B_\ell$ is contained in a single hyperplane

\mathcal{L} . Alice will choose A_ℓ to be $\mathcal{N}(\mathcal{L}, \beta^{\ell+1}\rho_0)$. Then for any $\mathbf{x} \in A_\ell$ and any $\frac{j_k \mathbf{p}}{i_k q} \in \mathcal{Q}_m \cap 2B_\ell$, we have

$$\begin{aligned} \left\| \mathbf{x} - \frac{j_k \mathbf{p}}{i_k q} \right\| &\geq \beta^{\ell+1} \rho_0 = \frac{\beta^{2^k+1}}{i_k/j_k} (i_k/j_k) \beta^{2^{k-1}+2^k(m-1)} \rho_0 \\ &= \frac{\beta^{2^k+1}}{i_k/j_k} \frac{Q_{m-1}^{-(1+1/d)}}{4d!} \geq \frac{\beta^{2^k+1}}{4d!} \frac{1}{i_k/j_k} q^{-(1+1/d)} \\ &\geq \frac{\varepsilon_k}{i_k/j_k} q^{-(1+1/d)}. \end{aligned}$$

On the other hand, for any $\frac{j_k \mathbf{p}}{i_k q} \in \mathcal{Q}_m \setminus 2B_\ell$ we have

$$\left\| \mathbf{x} - \frac{j_k \mathbf{p}}{i_k q} \right\| \geq \beta^\ell \rho_0 \geq \beta^{\ell+1} \rho_0 \geq \frac{\varepsilon_k}{i_k/j_k} q^{-(1+1/d)}.$$

Thus, if Alice plays according to this strategy and \mathbf{x} denotes the result of the game, then for each $k \in \mathbb{N}$, $\left\| \frac{j_k \mathbf{x}}{j_k} - \frac{\mathbf{p}}{q} \right\| \geq \varepsilon_k q^{-(1+1/d)}$ for all $(\mathbf{p}, q) \in \mathbb{Z}^d \times \mathbb{N}$. \square

4. PROOF OF THEOREM 1.3

Let $F_M \subseteq \mathbb{R}$ be the set of real numbers with partial quotients at most M . Recall that the set of badly approximable numbers is $\bigcup_{M \in \mathbb{N}} F_M$. We will prove that our set has full dimension in \mathbb{R} by showing that it has full dimension within each F_M . To this end we prove the following:

Proposition 4.1. *For each integer $M \geq 2$ and each reduced $\frac{i}{j} \in \mathbb{Q}$, the set $\{x \in F_M : L\left(\frac{i}{j}x\right) \leq \frac{1}{ij}\}$ is a BMS winning set on F_M .*

Before proving this, we show how it can be used to deduce Theorem 1.3.

Proof of Theorem 1.3. Fix $M \in \mathbb{N}$. One can easily check that $F_M \cap [0, 1]$ is the limit set of the contracting family of self-conformal maps $\{f_i\}_{i=1}^M$ on $[a, b]$, where $f_i(x) = \frac{1}{i+x}$, $a = [0; M, 1, M, 1, \dots]$ and $b = [0; 1, M, 1, M, \dots]$, and that this system satisfies the open set condition. It follows (see [9, Lemma 3.14]) that the δ_M -dimensional Hausdorff measure \mathcal{H}^{δ_M} is Ahlfors regular, where $\delta_M = \dim(F_M)$. By Proposition 4.1 the set

$$\left\{ x \in F_M : L\left(\frac{i}{j}x\right) \leq \frac{1}{ij} \text{ for all } i, j \in \mathbb{N} \right\}$$

is a countable intersection of BMS winning sets, hence conull with respect to \mathcal{H}^{δ_M} by Proposition 2.5. In particular,

$$\dim \left\{ x \in \mathbb{R} : L\left(\frac{i}{j}x\right) \leq \frac{1}{ij} \text{ for all } i, j \in \mathbb{N} \right\} \geq \delta_M.$$

Since $\dim(\bigcup_M F_M) = \dim(\text{BA}_1) = 1$, we have $\lim_{M \rightarrow \infty} \delta_M = 1$, so the theorem follows. \square

The main idea of the proof of Proposition 4.1 is as follows: Call an approximation p/q of x “good” if $|x - p/q| < 1/q^2$. If $i/j \in \mathbb{Q}$ is fixed and $(jp)/(iq)$ is a good approximation of x , then $|x - (jp)/(iq)| < 1/(iq)^2$, and rearranging gives $|(i/j)x - p/q| < 1/(ijq^2)$. Thus, if x has infinitely many good approximations whose denominator is a multiple of i and whose numerator is a multiple of j , then $L(\frac{i}{j}x) \leq \frac{1}{ij}$. Alice will direct the play of the game to numbers x with this property by using her turns to choose digits of the continued fraction expansion in such a way that infinitely many truncated expansions (i.e. convergents), which are good approximations to x , have numerator and denominator in the required sets. To show that this is possible we first prove the following lemma:

Lemma 4.2. *For any $i, j \in \mathbb{N}$ and $M \geq 2$, there exists $T = T(i, j, M) \in \mathbb{N}$ such that for any $n \in \mathbb{N}$ and any $a_1, a_2, \dots, a_n \in \mathbb{N}$, there exist $1 \leq t \leq T$ and $a_{n+1}, \dots, a_{n+t} \in \{1, \dots, M\}$ such that $[0; a_1, \dots, a_{n+t}]$ is a rational with numerator in $j\mathbb{N}$ and denominator in $i\mathbb{N}$.*

Proof. Recall [6, Theorem 1] that if $x = [0; a_1, a_2, \dots]$, then the approximations obtained by truncating the continued fraction expansion of x , $\frac{p_n}{q_n} = [0; a_1, a_2, \dots, a_n]$, satisfy the recurrence relation

$$(4.1) \quad \frac{p_n}{q_n} = \frac{a_n p_{n-1} + p_{n-2}}{a_n q_{n-1} + q_{n-2}}.$$

In other words,

$$\begin{bmatrix} p_{n-1} & q_{n-1} \\ p_n & q_n \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & a_n \end{bmatrix} \begin{bmatrix} p_{n-2} & q_{n-2} \\ p_{n-1} & q_{n-1} \end{bmatrix}.$$

Now it is well-known (e.g. [6, Theorem 2]) that

$$(4.2) \quad \det \begin{bmatrix} p_{n-1} & q_{n-1} \\ p_n & q_n \end{bmatrix} = (-1)^n.$$

Fix i, j , and M , and let $a_1, \dots, a_n \in \{1, \dots, M\}$. We may assume i and j are coprime (i.e. that the fraction i/j is reduced). We want to choose $a_{n+1}, a_{n+2}, \dots, a_{n+t}$ in such a way that

$$\begin{bmatrix} p_{n+t-1} & q_{n+t-1} \\ p_{n+t} & q_{n+t} \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & a_{n+t} \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & a_{n+t-1} \end{bmatrix} \cdots \begin{bmatrix} 0 & 1 \\ 1 & a_{n+1} \end{bmatrix} \begin{bmatrix} p_{n-1} & q_{n-1} \\ p_n & q_n \end{bmatrix} \equiv \begin{bmatrix} m_1 & m_2 \\ j & i \end{bmatrix} \pmod{ij},$$

where $m_1, m_2 \in \mathbb{Z}/ij\mathbb{Z}$ satisfy $m_1 j + m_2 i \equiv (-1)^n \pmod{ij}$ (such numbers exist since $\gcd(i, j) = 1$). It will suffice to show that the semigroup H generated by elements of the form $g_a = \begin{bmatrix} 0 & 1 \\ 1 & a \end{bmatrix}$ ($1 \leq a \leq M$) is equal to $G = \mathrm{SL}^\pm(2, \mathbb{Z}/ij\mathbb{Z})$. (Since G is finite, if $H = G$ then there exists T such that every element of G is the product of at most T elements of the form g_a .) Since G is finite, for any $g \in H$ there exist $1 \leq k_1 < k_2$ such that $g^{k_1} = g^{k_2}$ and therefore $g^{-1} = g^{k_2 - k_1 - 1} \in H$, so H is equal to the group generated by the elements g_a . In particular, we have

$$\begin{bmatrix} -a & 1 \\ 1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & a \end{bmatrix}^{-1} = g_a^{-1} \in H$$

for all $1 \leq a \leq M$.

Recall (e.g. [11, p.139]) that $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ generate $\mathrm{SL}(2, \mathbb{Z})$. Since

$$\begin{aligned} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} &= \begin{bmatrix} -1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 2 \end{bmatrix} \in H \\ \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} &= \begin{bmatrix} 0 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} -1 & 1 \\ 1 & 0 \end{bmatrix} \in H \\ \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} &= \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \in H, \end{aligned}$$

it follows that $\mathrm{SL}(2, \mathbb{Z}/ij\mathbb{Z}) \subseteq H$. Since $\det(g_a) = -1$ for all a , we have $G = \mathrm{SL}^\pm(2, \mathbb{Z}/ij\mathbb{Z}) = H$, which completes the proof. \square

Proof of Proposition 4.1. A modification of the argument of [10, Theorem 1.2] shows that the BMS game is invariant under quasisymmetric homeomorphisms, and in particular under the coding map $\pi : \{1, \dots, M\}^\mathbb{N} \rightarrow F_M$ defined by the formula $\pi(a_1, a_2, \dots) = [0; a_1, a_2, \dots]$. When the BMS game is played on the space $\{1, \dots, M\}^\mathbb{N}$, the gameplay is equivalent to the following: on each of Bob's moves, he chooses a word in the alphabet $E = \{1, \dots, M\}$ which is an extension of Alice's most recent word, and on each of Alice's moves, she chooses a word in the alphabet E which is an extension of Bob's most recent word by T letters, where $T \in \mathbb{N}$ is fixed at the start of the game. The result of the game is the unique infinite word ω which extends all of these words, so Alice's goal is to make the number $x = \pi(\omega)$ satisfy $L(\frac{i}{j}x) \leq \frac{1}{ij}$. Now let T be given by Lemma 4.2, and let Alice's strategy be given as follows: if Bob's most recent word is the move a_1, \dots, a_n , then Alice's next move will be the word a_1, \dots, a_{n+T} , where a_{n+1}, \dots, a_{n+t} are given by Lemma 4.2, and $a_{n+t+1}, \dots, a_{n+T}$ are arbitrary. When Alice uses this strategy, each of her moves determines a

new convergent of the continued fraction expansion of the final word whose numerator is in $j\mathbb{N}$ and whose denominator is in $i\mathbb{N}$.

Now, suppose $\frac{p_k}{q_k}$ is a convergent with $p_k \in j\mathbb{N}$ and $q_k \in i\mathbb{N}$, say $p_k = jp$ and $q_k = iq$. Then

$$\left| x - \frac{jp}{iq} \right| = \left| x - \frac{p_k}{q_k} \right| \leq \left| \frac{p_{k+1}}{q_{k+1}} - \frac{p_k}{q_k} \right| = \frac{1}{q_{k+1}q_k} \leq \frac{1}{i^2q^2}$$

and hence

$$\left| \frac{i}{j}x - \frac{p}{q} \right| = \frac{i}{j} \left| x - \frac{jp}{iq} \right| \leq \frac{i}{ji^2q^2} = \frac{1}{ij} \frac{1}{q^2}.$$

Since this holds for infinitely many fractions $\frac{p}{q}$, it follows that $L\left(\frac{i}{j}x\right) \leq \frac{1}{ij}$. \square

REFERENCES

1. D. Badziahin and S. Harrap, *Cantor-winning sets and their applications*, <http://arxiv.org/abs/1503.04738>, preprint 2015.
2. R. Broderick, L. Fishman, D. Y. Kleinbock, A. Reich, and B. Weiss, *The set of badly approximable vectors is strongly C^1 incompressible*, Math. Proc. Cambridge Philos. Soc. **153** (2012), no. 02, 319–339.
3. Y. Bugeaud, *On the multiples of a badly approximable vector*, Acta Arith. **168** (2015), no. 1, 71–81.
4. L. Fishman, V. Reams, and D. S. Simmons, *The Banach–Mazur–Schmidt and Banach–Mazur–McMullen games*, <http://arxiv.org/abs/1503.03428>, preprint 2015.
5. L. Fishman, D. S. Simmons, and M. Urbański, *Diophantine approximation and the geometry of limit sets in Gromov hyperbolic metric spaces*, <http://arxiv.org/abs/1301.5630>, preprint 2013.
6. A. Y. Khinchin, *Continued fractions*, The University of Chicago Press, Chicago, Ill.-London, 1964.
7. S. Kristensen, R. Thorn, and S. L. Velani, *Diophantine approximation and badly approximable sets*, Advances in Math. **203** (2006), 132–169.
8. R. D. Mauldin, T. Szarek, and M. Urbański, *Graph directed Markov systems on Hilbert spaces*, Math. Proc. Cambridge Philos. Soc. **147** (2009), 455–488.
9. R. D. Mauldin and M. Urbański, *Dimensions and measures in infinite iterated function systems*, Proc. London Math. Soc. (3) **73** (1996), no. 1, 105–154.
10. C. T. McMullen, *Winning sets, quasiconformal maps and Diophantine approximation*, Geom. Funct. Anal. **20** (2010), no. 3, 726–740.
11. M. A. P. Newman, *Integral matrices*, Pure and Applied Mathematics, Vol. 45, Academic Press, New York-London, 1972.

340 ROWLAND HALL (BLDG.# 400), UNIVERSITY OF CALIFORNIA, IRVINE, IRVINE, CA 92697-3875, USA
E-mail address: broderir@uci.edu

UNIVERSITY OF NORTH TEXAS, DEPARTMENT OF MATHEMATICS, 1155 UNION CIRCLE #311430, DENTON, TX 76203-5017, USA

E-mail address: lior.fishman@unt.edu

UNIVERSITY OF YORK, DEPARTMENT OF MATHEMATICS, HESLINGTON, YORK YO10 5DD, UK
E-mail address: David.Simmons@york.ac.uk

URL: <https://sites.google.com/site/davidsimmonsmath/>