

ON A THEOREM OF SERRET ON CONTINUED FRACTIONS

PALOMA BENGOCHEA

ABSTRACT. A classical theorem in continued fractions due to Serret shows that for any two irrational numbers x and y related by a transformation γ in $\mathrm{PGL}(2, \mathbb{Z})$ there exist s and t for which the complete quotients x_s and y_t coincide. In this paper we give an upper bound in terms of γ for the smallest indices s and t .

A classical theorem of Serret [Ser66] states that the standard continued fraction expansions of two irrational numbers $x = [n_0, n_1, \dots]$ and $y = [m_0, m_1, \dots]$ are the same after a finite number of steps if and only if x and y are $\mathrm{PGL}(2, \mathbb{Z})$ -equivalent. Serret's theorem does not quantify how large the smallest indices s and t have to be such that the complete quotients $x_s = [n_s, n_{s+1}, \dots]$ and $y_t = [m_t, m_{t+1}, \dots]$ coincide. The more complicated the matrix γ relating x and y is, the larger s or t becomes. A quantification of this fact, namely bounding s and t in terms of γ , is the goal of this paper. The main theorem, in Section 2, gives an upper bound for the smallest s and t such that $x_s = y_t$. The bound we give is the best possible in the sense that for each $\gamma \in \mathrm{PGL}(2, \mathbb{Z})$ one can construct many real numbers x for which s or t almost attain the bound, and for many matrices γ the bound is actually attained. In order to prove the theorem, in the first section we introduce the set $\Gamma(x)$ of transformations of $\mathrm{PGL}(2, \mathbb{Z})$ related to the continued fraction expansion of the real number x . We characterize it by certain linear inequalities in Proposition 1.2. Sets of transformations related to other continued fraction algorithms have been characterized in [KU10] in a similar way.

1. LINEAR INEQUALITIES CHARACTERIZING THE CONTINUED FRACTION TRANSFORMATIONS

We denote by Γ the group $\mathrm{PGL}(2, \mathbb{Z})$ and by ε and T the transformations

$$\varepsilon = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

Key words and phrases. Continued fractions, $\mathrm{PGL}(2, \mathbb{Z})$ -equivalent numbers.
Mathematics Subject Classification: 11A55.

that correspond to the inversion and the translation for the usual action on the projective line

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} x := \frac{ax + b}{cx + d}.$$

The positive continued fraction of a real number x ,

$$x = n_0 + \frac{1}{n_1 + \frac{1}{n_2 + \frac{1}{\ddots}}} \quad (n_0 \in \mathbb{Z}, n_i \geq 1 \ \forall i \geq 1),$$

also denoted by $x = [n_0, n_1, \dots]$, is given by the algorithm

$$\begin{aligned} x_0 &= x, \\ n_i &= \lfloor x_i \rfloor, \quad x_{i+1} = \frac{1}{x_i - n_i} = \varepsilon T^{-n_i}(x_i) \quad \text{if } x_i \notin \mathbb{Z} \quad (i \geq 0). \end{aligned}$$

If $x_i \in \mathbb{Z}$, the algorithm stops after $n_i = \lfloor x_i \rfloor$. Clearly each complete quotient $x_i = [n_i, n_{i+1}, \dots]$ is the image of x by a matrix $\gamma_i = \gamma_{i,x} \in \Gamma$, given explicitly by

$$\gamma_0 := \text{Id}, \quad \gamma_i := \begin{pmatrix} 0 & 1 \\ 1 & -n_{i-1} \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ 1 & -n_0 \end{pmatrix} \quad (i \geq 1)$$

and recursively by

$$(1) \quad \gamma_0 = \text{Id}, \quad \gamma_{i+1} = \varepsilon T^{-n_i} \gamma_i \quad (i \geq 0).$$

A key idea is to replace the sequence $(\gamma_1, \gamma_2, \gamma_3, \dots)$ of elements in Γ by the unordered set

$$\Gamma(x) = \{\gamma_1, \gamma_2, \gamma_3, \dots\} \subset \Gamma.$$

When $x \in \mathbb{R} \setminus \mathbb{Q}$, the set $\Gamma(x)$ is infinite, whereas it is finite if $x \in \mathbb{Q}$. If $x = [n_0, \dots, n_t]$, then

$$\Gamma(x) = \{\gamma_1, \dots, \gamma_{t+1}\}.$$

The i -th convergent of x is denoted by $\frac{p_i}{q_i} = [n_0, \dots, n_i]$. The integers p_i and q_i satisfy the recurrence

$$\begin{aligned} p_{-2} &= 0, & p_{-1} &= 1, & p_i &= n_i p_{i-1} + p_{i-2} & (i \geq 0), \\ q_{-2} &= 1, & q_{-1} &= 0, & q_i &= n_i q_{i-1} + q_{i-2} & (i \geq 0), \end{aligned}$$

the equation

$$p_{i+1} q_i - p_i q_{i+1} = (-1)^i$$

and the inequalities

$$(2) \quad q_i \geq q_{i-1} \geq 0 \quad (i \geq 0), \quad q_i > q_{i-1} > 0 \quad (i \geq 2),$$

$$(3) \quad |p_i| \geq |p_{i-1}| \quad (i \geq 2), \quad |p_i| > |p_{i-1}| \quad (i \geq 4).$$

It is well known that any rational number p/q satisfying

$$\left| \frac{p}{q} - x \right| < \frac{1}{2q^2}$$

is a convergent of x .

The numbers δ_i ($i \geq -1$) defined by

$$\delta_i = (-1)^i (p_{i-1} - q_{i-1}x)$$

satisfy the recurrence

$$\delta_{-1} = x, \quad \delta_0 = 1, \quad \delta_{i+1} = -n_i \delta_i + \delta_{i-1},$$

and the inequalities $1 = \delta_0 > \delta_1 > \dots \geq 0$. If x is rational, then $x_i = p_i/q_i$ for some i and the recurrence stops with $\delta_{i+1} = 0$; if x is irrational, the δ_i are all positive and converge to 0 with exponential rapidity. One has

$$n_i = \left\lfloor \frac{\delta_{i-1}}{\delta_i} \right\rfloor \quad \text{if } \delta_i \neq 0,$$

and

$$\gamma_i^{-1} = \begin{pmatrix} p_{i-1} & p_{i-2} \\ q_{i-1} & q_{i-2} \end{pmatrix}, \quad \gamma_i \begin{pmatrix} x \\ 1 \end{pmatrix} = \begin{pmatrix} \delta_{i-1} \\ \delta_i \end{pmatrix}.$$

Lemma 1.1. *If r/t and s/u are two rational numbers such that $t, u > 0$, $\frac{r}{t} \leq x \leq \frac{s}{u}$ and $ru - st = \pm 1$, then r/t or s/u is a convergent of x .*

Proof. Let x be an element in $\left[\frac{r}{t}, \frac{s}{u}\right]$. If one of the inequalities

$$\left| \frac{r}{t} - x \right| < \frac{1}{2t^2} \quad \text{or} \quad \left| \frac{s}{u} - x \right| < \frac{1}{2u^2}$$

is satisfied, then r/t or s/u is a convergent of x . If not, then

$$\frac{1}{tu} = \left| \frac{r}{t} - \frac{s}{u} \right| = \left| \frac{r}{t} - x \right| + \left| \frac{s}{u} - x \right| \geq \frac{1}{2t^2} + \frac{1}{2u^2}.$$

The inequality above can only hold if $t = u = 1$, in which case $s = r + 1$, so $\left[\frac{r}{t}, \frac{s}{u}\right] = [r, r + 1]$, and then either $\frac{s}{u} = x = \frac{p_0}{q_0}$ or $\frac{r}{t} = \lfloor x \rfloor = \frac{p_0}{q_0}$. \square

Throughout, for any pair of real numbers a, b , we denote by $|a, b|$ the interval $[a, b]$ if $a \leq b$ or $[b, a]$ if $b < a$.

Proposition 1.2. *For any real number x , the set $\Gamma(x)$ equals $W - (W_1 \cup W_2)$, where*

$$W = \{\gamma \in \Gamma \mid -1 \leq \gamma(\infty) \leq 0, \gamma(x) > 1\}$$

and

$$W_1 = \{\gamma \in W \mid \gamma(\infty) = 0, \det(\gamma) = 1\},$$

$$W_2 = \{\gamma \in W \mid \gamma(\infty) = -1, \det(\gamma) = -1\}.$$

The sets W_1 and W_2 have respectively exactly one and at most one element. (The inequality $\gamma(x) > 1$ in the definition of W includes $\gamma(x) = \infty$.)

Proof. One easily checks that

$$W_1 = \left\{ \begin{pmatrix} 0 & -1 \\ 1 & -1 - n_0 \end{pmatrix} \right\},$$

$$W_2 = \begin{cases} \left\{ \begin{pmatrix} -1 & 1 + n_0 \\ 1 & -n_0 \end{pmatrix} \right\} & \text{if } x \in \mathbb{Z} \text{ or } n_1 \geq 2, \\ \emptyset & \text{if } n_1 = 1. \end{cases}$$

This proves the second statement. It is also easy to see that $\Gamma(x) \subseteq W - (W_1 \cup W_2)$. Indeed, $\gamma_i(x) = \frac{\delta_{i-1}}{\delta_i} > 1$ and $\gamma_i(\infty) = -\frac{q_{i-2}}{q_{i-1}}$ for $i \geq 1$ together with the inequalities (2) imply $\gamma_i \in W - (W_1 \cup W_2)$.

Therefore we only have to show that $W \subseteq \Gamma(x) \cup W_1 \cup W_2$. Let $\gamma \in \Gamma$ satisfy

$$-1 \leq \gamma(\infty) < 0, \quad \gamma(x) > 1.$$

Let

$$\gamma^{-1} = \begin{pmatrix} r & s \\ t & u \end{pmatrix}, \quad \gamma = \begin{pmatrix} u & -s \\ -t & r \end{pmatrix}.$$

The conditions $\gamma(\infty) < 0$ and $\gamma(x) > 0$ imply that u and t have the same sign, as well as $ux - s$ and $-tx + r$, so $x \in \left[\frac{r}{t}, \frac{s}{u} \right]$. By Lemma 1.1, r/t or s/u is a convergent of x .

If r/t is a convergent of x , then γ and γ^{-1} are of the form

$$(4) \quad \gamma^{-1} = \begin{pmatrix} p_i & p_{i-1} + kp_i \\ q_i & q_{i-1} + kq_i \end{pmatrix}, \quad \gamma = \begin{pmatrix} q_{i-1} + kq_i & -p_{i-1} - kp_i \\ -q_i & p_i \end{pmatrix}$$

or

$$(5) \quad \gamma^{-1} = \begin{pmatrix} p_i & -p_{i-1} - kp_i \\ q_i & -q_{i-1} - kq_i \end{pmatrix}, \quad \gamma = \begin{pmatrix} -q_{i-1} - kq_i & p_{i-1} + kp_i \\ -q_i & p_i \end{pmatrix}$$

with $k \in \mathbb{Z}$ and $i \geq 0$. In the case (4), since $q_{i-1} \leq q_i$, we have that $-1 \leq \gamma(\infty) = -\frac{q_{i-1}}{q_i} - k < 0$ if and only if $k = 0$ and $i \geq 1$, or $k = 1$

and $i = 0$. Moreover, if $x \notin \mathbb{Z}$, then $\gamma(x) = \frac{\delta_i}{\delta_{i+1}} - k > 1$ if and only

if $k \leq n_{i+1} - 1$ because $\left\lfloor \frac{\delta_i}{\delta_{i+1}} \right\rfloor = n_{i+1}$. Hence $-1 \leq \gamma(\infty) < 0$ and

$\gamma(x) > 1$ if and only if $k = 0$ and $i \geq 1$ or $x \in \mathbb{Z}$, $k = 1$ and $i = 0$, or $n_1 \geq 2$, $k = 1$ and $i = 0$. In the first case

$$\gamma = \begin{pmatrix} q_{i-1} & -p_{i-1} \\ -q_i & p_i \end{pmatrix} = \gamma_{i+1} \quad (i \geq 1);$$

in the other two cases

$$\gamma = \begin{pmatrix} -1 & 1 + n_0 \\ 1 & -n_0 \end{pmatrix} \in W_2.$$

In the case (5), if $-1 \leq \gamma(\infty) = \frac{q_{i-1}}{q_i} + k < 0$, then $k < 0$ and $\gamma(x) = -\frac{\delta_i}{\delta_{i+1}} + k < 0$, so $\gamma \notin W$.

If s/u is a convergent of x , then γ and γ^{-1} are of the form

$$(6) \quad \gamma^{-1} = \begin{pmatrix} p_{i-1} + kp_i & p_i \\ q_{i-1} + kq_i & q_i \end{pmatrix}, \quad \gamma = \begin{pmatrix} q_i & -p_i \\ -q_{i-1} - kq_i & p_{i-1} + kp_i \end{pmatrix}$$

or

$$(7) \quad \gamma^{-1} = \begin{pmatrix} -p_{i-1} - kp_i & p_i \\ -q_{i-1} - kq_i & q_i \end{pmatrix}, \quad \gamma = \begin{pmatrix} q_i & -p_i \\ q_{i-1} + kq_i & -p_{i-1} - kp_i \end{pmatrix}$$

with $k \in \mathbb{Z}$ and $i \geq 0$. In the case (6), $\gamma(x) = \frac{\delta_{i+1}}{\delta_i - k\delta_{i+1}} > 1$ if and only if $k = n_{i+1}$. If $k = n_{i+1}$, then $-1 \leq \gamma(\infty) = -\frac{q_i}{q_{i+1}} < 0$. Hence

$$\gamma = \begin{pmatrix} q_i & -p_i \\ -q_{i+1} & p_{i+1} \end{pmatrix} = \gamma_{i+2} \quad (i \geq 0).$$

In the case (7), if $-1 \leq \gamma(\infty) = \frac{q_i}{q_{i-1} + kq_i} < 0$, then $k < 0$ and $\gamma(x) = \frac{\delta_{i+1}}{-\delta_i + k\delta_{i+1}} \leq 0$, so $\gamma \notin W$.

Finally, let $\gamma = \begin{pmatrix} 0 & 1 \\ 1 & u \end{pmatrix}$ be the unique element of W satisfying $\gamma(\infty) = 0$ and $\det(\gamma) = -1$. We have $1 < \gamma(x) = \frac{1}{x+u}$ if and only if $0 < x+u < 1$, namely $u = -\lfloor x \rfloor$, and $\gamma = \gamma_1 \in \Gamma(x)$.

□

2. MAIN THEOREM

We recall the classic theorem of Serret about continued fraction expansions of equivalent numbers (see [Ser66]):

Theorem 2.1 (Serret). *Two irrational numbers x and y are Γ -equivalent if and only if there exist $s, t \geq 0$ such that $x_s = y_t$.*

Theorem 2.1 (or its proof) does not give any bound independent of x and y for the indexes s and t . The result we state below gives the best possible bound for the index s in terms of the matrix relating x and y (see Remark 2.3).

We introduce the convention that ∞ has zero partial quotients in the rest of the paper.

Theorem 2.2. *Let $\gamma \in \Gamma$ and let r be the number of partial quotients of $\gamma^{-1}(\infty)$. For every real number x there exists an index $s \leq r + 3$ such that the complete quotient x_s is also a complete quotient of $\gamma(x)$.*

Of course the bound for t is obtained in the same way in terms of the number of partial quotients of $\gamma(\infty)$.

Proof. Let $y = \gamma(x)$. Since

$$x_s = \gamma_{s,x}(x) \quad \text{and} \quad y_t = \gamma_{t,y}(y) = (\gamma_{t,y}\gamma)(x),$$

we must prove that there exists $s \leq r + 3$ such that $\gamma_{s,x} = \gamma_{t,y}\gamma$ for some $t \geq 1$. In other words, we prove next that, if $\gamma_{i,x} \neq \gamma_{t,y}\gamma$, then $i \leq r + 2$. Let $i \geq 0$ be such that $\gamma_{i,x}\gamma^{-1} \notin \Gamma(y)$. Proposition 1.2 implies one of the following conditions:

$$(8) \quad \gamma_{i,x}\gamma^{-1}(\infty) \geq 0 \quad \text{or} \quad \gamma_{i,x}\gamma^{-1}(\infty) \leq -1.$$

Suppose $\gamma^{-1}(\infty) \neq \infty$ and set $p/q = \gamma^{-1}(\infty)$ with $(p, q) = 1$. If the first inequality in (8) is satisfied, we have

$$\gamma_{i,x}\left(\frac{p}{q}\right) = \frac{q_{i-2}p - p_{i-2}q}{-q_{i-1}p + p_{i-1}q} \geq 0,$$

which implies $\frac{p}{q} \in \left[\frac{p_{i-2}}{q_{i-2}}, \frac{p_{i-1}}{q_{i-1}}\right]$. Then by Lemma 1.1 we have that either p_{i-1}/q_{i-1} or p_{i-2}/q_{i-2} is a convergent of p/q . Similarly, if the second inequality in (8) is satisfied, we have

$$\gamma_{i,x}\left(\frac{p}{q}\right) = \frac{q_{i-2}p - p_{i-2}q}{-q_{i-1}p + p_{i-1}q} \leq -1,$$

which implies $\frac{p}{q} \in \left[\frac{p_{i-1} - p_{i-2}}{q_{i-1} - q_{i-2}}, \frac{p_{i-1}}{q_{i-1}}\right]$. It is easy to see that $\frac{p_{i-1} - p_{i-2}}{q_{i-1} - q_{i-2}} \in \left[\frac{p_{i-3}}{q_{i-3}}, \frac{p_{i-1}}{q_{i-1}}\right]$. Hence by Lemma 1.1 we have that either p_{i-1}/q_{i-1} or p_{i-3}/q_{i-3} is a convergent of p/q . Thus, if we denote by r the number of partial quotients (or convergents) of p/q , we have that $i - 3 \leq r - 1$.

Now suppose $\gamma^{-1}(\infty) = \infty$. Then we have that, for all $i \geq 1$,

$$\gamma_{i,x}\gamma^{-1}(y) = \gamma_{i,x}(x) \quad \text{and} \quad \gamma_{i,x}\gamma^{-1}(\infty) = \gamma_{i,x}(\infty),$$

so

$$\gamma_{i,x}\gamma^{-1}(y) > 1 \quad \text{and} \quad -1 \leq \gamma_{i,x}\gamma^{-1}(\infty) \leq 0.$$

If, moreover, $\gamma_{i,x}\gamma^{-1}(\infty) \notin \{-1, 0\}$, then, by Proposition 1.2, we have

that $\gamma_{i,x}\gamma^{-1} \in \Gamma(y)$. Now, if $\gamma_{i,x}(\infty) = 0$, then $\gamma_{i,x} = \gamma_{1,x} = \begin{pmatrix} 0 & 1 \\ 1 & -n_0 \end{pmatrix}$.

If $\gamma_{i,x}(\infty) = -1$, then $n_1 = 1$ and $\gamma_{i,x} = \gamma_{2,x} = \begin{pmatrix} 1 & -n_0 \\ -1 & 1 + n_0 \end{pmatrix}$.

Therefore $i \leq 2$.

□

Remark 2.3. Let $\gamma \in \Gamma$ and set $\gamma^{-1}(\infty) = [n_0, \dots, n_{r-1}]$. One can construct many real numbers x for which x_{r+1} is not a complete quotient of $\gamma(x)$ by letting $x = [n_0, \dots, n_{r-2}, n_{r-1} - 1, 1, n_{r+1}, \dots]$ where n_{r+i} are arbitrary for all $i \geq 1$. Indeed, for such an x , we have that $\gamma^{-1}(\infty) = p_r/q_r$, so $\gamma_{r+1,x}\gamma^{-1}(\infty) = \infty$, and then $\gamma_{r+1,x} \notin \Gamma(y)\gamma$. Therefore x_{r+1} is not a complete quotient of $\gamma(x)$. Moreover, $\gamma_{r+2,x}\gamma^{-1}(\infty) = 0$, and if $\det(\gamma) = \det(\gamma_{r+2,x})$, then $\gamma_{r+2,x} \notin \Gamma(y)\gamma$ and x_{r+2} is not a complete quotient of $\gamma(x)$ either.

REFERENCES

- [KU10] Katok, S.; Ugarcovici, I.: *Structure of attractors for (a, b)-continued fraction transformations*, Journal of Modern Dynamics, 4 (2010), 637 - 691.
- [Ser66] Serret, J.A.: *Cours d'algèbre supérieure*, 3ème ed., volume premier (1866).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK, YO10 5DD,
UNITED KINGDOM

E-mail address: paloma.bengoechea@york.ac.uk