



UNIVERSITY OF LEEDS

This is a repository copy of *A three dimensional view of stereopsis in dentistry*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/93456/>

Version: Accepted Version

Article:

Mon-Williams, MA, Mushtaq, F, Wilkie, RM et al. (3 more authors) (2015) A three dimensional view of stereopsis in dentistry. *British Dental Journal*, 219 (10). pp. 479-480. ISSN 0007-0610

<https://doi.org/10.1038/sj.bdj.2015.881>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

A three dimensional view of stereopsis in dentistry

Mark A. Mon-Williams¹, Faisal Mushtaq^{1*}, Richard M. Wilkie¹, Balvinder Khambay², Andrew Keeling² & Michael Manogue²

¹School of Psychology, Faculty of Medicine and Health, University of Leeds, Leeds, West Yorkshire, United Kingdom

²School of Dentistry, Faculty of Medicine and Health, University of Leeds, Leeds, West Yorkshire, United Kingdom

*Correspondence can be addressed to Faisal Mushtaq at the School of Psychology, University of Leeds, Leeds, LS2 9JT, UK. Email: f.mushtaq@leeds.ac.uk. Tel: +44 (0) 113 343 6640

Short Title: Stereopsis and Dentistry

Article Type: Opinion Article

Number of words: 1,007

Does dentistry require the three-dimensional information afforded by binocular vision? Stereopsis and its role in dental practice has been a topic of debate in recent editions of British Dental Journal (BDJ) ^{1,2}. These discussions are particularly timely as they come at a point when virtual reality simulators (often relying on 3D stereo-projected images to create “realistic” dental environments) are becoming increasingly popular in the education of tomorrow’s dentists ³⁻⁵. Such devices have raised questions about whether stereo-projected images are a necessary feature of simulators ⁶⁻⁸. This relates to the larger issue of whether an ability to perceive the information that arises specifically from a binocular view should be a prerequisite for surgical training ^{9,10}.

Syrimi and Ali’s recent review of the literature¹ concluded that “stereopsis should not be considered a requirement for dental training”. This conclusion was supported by a subsequent personal account from an experienced practitioner with a ‘lazy left eye’ and a history of dismal abilities in hitting moving targets, including “high velocity tennis, cricket and squash balls” ². Duffy drew attention to the fact that he had “made a reasonable success of [his] career” and thus “anyone with similar issues reading the article should take some comfort from this”. Nevertheless, we argue that Syrimi and Ali’s conclusion is potentially indefensible on the basis of the (absence of) evidence provided in their review. We further suggest that the use of anecdotal reports to encourage trainees with known perceptual deficits to enter the profession of dentistry is not compatible with a health service that has patient safety as its primary objective – especially when ‘career success’ is considered an outcome measure. We will briefly outline the grounds for our arguments below.

It is first necessary to define what is meant by the term stereopsis (an omission in the review of Syrimi and Ali who appear to regard the term as

synonymous with 'depth perception'). Stereopsis can be defined as the information regarding three dimensional object structure which is made available through retinal image differences (differences that arise because the eyes are horizontally separated by approximately 6cm in humans). Many studies have shown that: (i) horizontal disparities can provide information about the slant, curvature and depth of proximally fixated objects ¹¹⁻¹³; (ii) humans use this information ¹⁴⁻¹⁶; (iii) the use of stereopsis (and other 'cues') is task dependent ^{17,18} (with the binocular information regarding the time to contact of an approaching ball ^{19,20} being used in a different manner to the information used - for example - in gauging object curvature ²¹). Thus, a fundamental and important question arises from the existing literature - is stereopsis a necessary cue in dental tasks?

The implicit reasoning within Syrimi and Ali's review centres on the conjecture that if individuals with stereo-deficits can "function effectively as dental students and practitioners" then it follows that stereopsis is not needed within dentistry. This conjecture appears reasonable and seems to rest on the valid propositional logic of modus tollens. We suggest, however, that Syrimi and Ali's position is actually based on an absence of evidence- which is not the same as 'evidence of absence'. One major difficulty with Syrimi and Ali's proposition is that they have not defined what constitutes a stereo-deficit or 'effective function'. Moreover, there is a lack of acknowledgement that stereo-deficits can arise through a number of aetiological routes ('lazy eye' has a similar level of diagnostic precision as 'tooth rot'). Unfortunately, satisfactory definitions of effective function cannot be achieved without understanding how stereopsis is used (or not) in specific dental tasks (e.g. drilling teeth) and measuring the minimum thresholds of stereo-perception if

stereopsis is required. In our opinion, an absence of studies on this topic cannot logically be used to support the notion that stereopsis is not important.

The discipline of psychology has a long history of empirically investigating the task-specific use of perceptual cues²²⁻²⁵. There is no doubt that stereopsis creates a strong phenomenological sense of three-dimensional space (a fact that is often used to mask the absence of interesting storylines in films such as *Avatar*). But the critical issue is whether the cognitively penetrable sense of an object's stereoscopic dimensions is necessary in dental decision-making (e.g. judging the extent of dental caries). There is also no doubt that stereopsis supports a number of skilled human behaviours – i.e. it can provide essential information for feedforward and feedback motor control mechanisms²⁶⁻²⁸. The crucial issue in this instance is whether specific dental tasks require this perceptual information (or whether other available cues will suffice). As Syrimi and Ali's review demonstrates, there is a notable and disappointing lack of investigation into this topic within the dental literature.

From a patient safety perspective, we would argue that any known perceptual or motor deficit should be an exclusion criterion for dental training unless there is unequivocal evidence that such deficits do not impact on dental task performance. The argument against such an approach relates to the possibility of discriminating against an individual with a physical impairment that is not relevant to the core skills needed within dentistry. We would suggest that such concerns are legitimate but from a safety perspective the dental profession is subject to *onus probandi* – it is reasonable for the public to expect that entrants to the dental profession have no known perceptual-motor deficit unless it has been unequivocally demonstrated that a particular deficit does not impact negatively on any aspect of patient care (with absence of evidence not constituting an adequate demonstration).

In conclusion, we observe that screening for stereo-deficits is easily implemented (with well-defined thresholds of abnormality established) and suggest that it is hard to justify 'turning a blind eye' to a potential impediment to dental proficiency. This reasoning raises the more general (and difficult to address) issue of testing eye-hand coordination competency. We submit that it is incumbent on the dental profession to identify the key perceptual-motor abilities underpinning dental competency so that individuals with known perceptual and motor deficits do not embark on a career that may impact negatively on patient care.

References

- 1 Syrimi M, Ali N. The role of stereopsis (three-dimensional vision) in dentistry: review of the current literature. *Br Dent J* 2015; **218**: 597–598.
- 2 Duffy S. Stereopsis in dentistry: Dismal sporting skills. *Br Dent J* 2015; **219**: 3–3.
- 3 San Diego JP, Cox MJ, Quinn BFA, Newton JT, Banerjee A, Woolford M. Researching haptics in higher education: The complexity of developing haptics virtual learning systems and evaluating its impact on students' learning. *Comput Educ* 2012; **59**: 156–166.
- 4 Tse B, Harwin W, Barrow A, Quinn B, Diego JS, Cox M. Design and development of a haptic dental training system: hapTEL. 2010; : 101–108.
- 5 Bakker D, Lagerweij M, Wesselink P, Vervoorn M. Transfer of manual dexterity skills acquired in the Simodont, a dental haptic trainer with a virtual environment, to reality: a pilot study. *Bio-Algorithms and Med-Systems* 2010; **6**: 21–24.
- 6 McIntire JP, Havig PR, Geiselman EE. Stereoscopic 3D displays and human performance: A comprehensive review. *Displays* 2014; **35**: 18–26.
- 7 Stanney KM, Mourant RR, Kennedy RS. Human Factors Issues in Virtual Environments: A Review of the Literature. *Presence Teleoperators Virtual Environ* 1998; **7**: 327–351.
- 8 Mon-Williams M, Wann JP, Rushton S. Binocular vision in a virtual world: visual deficits following the wearing of a head-mounted display. *Ophthalmic Physiol Opt* 1993; **13**: 387–391.
- 9 Bloch E, Uddin N, Gannon L, Rantell K, Jain S. The effects of absence of stereopsis on performance of a simulated surgical task in two-dimensional and three-dimensional viewing conditions. *Br J Ophthalmol* 2014; **99**: 240–245.
- 10 Dimitrijevic T, Kahler B, Evans G, Collins M, Moule a. Depth and Distance Perception of Dentists and Dental Students. *Oper Dent* 2011; **36**: 467–477.
- 11 Stevens KA, Brookes A. Integrating stereopsis with monocular interpretations of planar surfaces. *Vision Res* 1988; **28**: 371–386.
- 12 Banks MS, Hooge ITC, Backus BT. Perceiving slant about a horizontal axis from stereopsis. *J Vis* 2001; **1**: 1.
- 13 Wickens CD, Merwin DH, Lin EL. Implications of Graphics Enhancements for the Visualization of Scientific Data: Dimensional Integrality, Stereopsis, Motion, and Mesh. *Hum Factors J Hum Factors Ergon Soc* 1994; **36** : 44–61.
- 14 Fielder AR, Moseley MJ. Does stereopsis matter in humans? *Eye* 1996; **10** (Pt 2): 233–238.

- 15 Saladin JJ. Stereopsis from a performance perspective. *Optom Vis Sci* 2005; **82**: 186–205.
- 16 Schreiber K, Crawford JD, Fetter M, Tweed D. The motor side of depth vision. *Nature* 2001; **410**: 819–822.
- 17 Read JCA, Begum SF, McDonald A, Trowbridge J. The binocular advantage in visuomotor tasks involving tools. *Iperception* 2013; **4**: 101–110.
- 18 Greenwald HS, Knill DC. A comparison of visuomotor cue integration strategies for object placement and prehension. *Vis Neurosci* 2008; **26**: 63–72.
- 19 Gray R, Regan D. Chapter 13 The use of binocular time-to-contact information. In: Psychology HH and GJPSBT-A in (ed). *Time-to-Contact*. North-Holland, 2004, pp 303–325.
- 20 Tresilian JR. Visually timed action: time-out for ‘tau’? *Trends Cogn Sci* 1999; **3**: 301–310.
- 21 Pierce RS, Bian Z, Braunstein ML, Andersen GJ. Detection of 3D curved trajectories: the role of binocular disparity. *Front Behav Neurosci* 2013; **7**: 12.
- 22 Bingham GP, Bradley A, Bailey M, Vinner R. Accommodation, occlusion, and disparity matching are used to guide reaching: A comparison of actual versus virtual environments. *J Exp Psychol Percept Perform* 2001; **27**: 1314–1334.
- 23 Loftus A, Servos P, Goodale MA, Mendarozqueta N, Mon-Williams M. When two eyes are better than one in prehension: monocular viewing and end-point variance. *Exp Brain Res* 2004; **158**: 317–327.
- 24 Mon-Williams M, Bingham GP. Ontological issues in distance perception: cue use under full cue conditions cannot be inferred from use under controlled conditions. *Percept Psychophys* 2008; **70**: 551–561.
- 25 Tresilian JR, Mon-Williams M. A curious illusion suggests complex cue interactions in distance perception. *J Exp Psychol Percept Perform* 1999; **25**: 677–687.
- 26 Niechwiej-Szwedo E, Goltz HC, Chandrakumar M, Hirji Z, Crawford JD, Wong a. MF. Effects of Anisometropic Amblyopia on Visuomotor Behavior, Part 2: Visually Guided Reaching. *Invest Ophthalmol Vis Sci* 2010; **52**: 795–803.
- 27 Bradshaw MF, Elliott KM. The role of binocular information in the ‘on-line’ control of prehension. *Spat Vis* 2003; **16**: 295–309.
- 28 Brenner E, Smeets JJ. Two eyes in action. *Exp Brain Res* 2006; **170**: 302–311.