



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/94552/>

Version: Accepted Version

Article:

Collinson, K, Murray, MJ, Orsi, NM et al. (2014) Age-related biological features of germ cell tumors. *Genes, Chromosomes and Cancer*, 53 (3). pp. 215-227. ISSN: 1045-2257

<https://doi.org/10.1002/gcc.22131>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Age-related biological features of germ cell tumours

Kate Collinson¹, Matthew J. Murray², Nicolas M. Orsi³, Michele Cummings³, Janet Shipley⁴, Jonathan K. Joffe⁵, Nicholas Coleman², Daniel Stark⁶

Author details

1. Department of Medical Oncology, St James University Hospital, Leeds, LS9 7TF, UK
2. Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge, CB2 1QP, UK
3. Women's Health Research Group, Section of Cancer & Pathology, Leeds Institute of Cancer Studies and Pathology, Wellcome Trust Brenner Building, St James's University Hospital, Beckett Street, Leeds LS9 7TF, UK
4. Divisions of Molecular Pathology and Cancer Therapeutics, The Institute of Cancer Research, Surrey, UK
5. Macmillan Consultant in Medical Oncology, Departments of Oncology, Calderdale & Huddersfield NHS Foundation Trust, Leeds Teaching Hospitals NHS Trust
6. Leeds Institute of Cancer Studies and Pathology, Leeds Institute of Oncology and St James's University Hospital, Leeds, LS9 7TF, UK

Corresponding author: Dr Daniel Stark, mailing address: St James's University Hospital, Level 4 Bexley Wing, St James's University Hospital, Beckett Street, Leeds, LS9 7TF, Telephone: 0113 2067533, Fax number: 0113 2067871, email: D.P.Stark@leeds.ac.uk

Keywords: biology; germ cell tumour; gonadal; primordial germ cell; teenagers and young adults

Abstract

Germ cell tumours (GCTs) are rare but clinically and pathologically diverse tumours that occur in an extensive range of age groups, from children to older adults and which include both seminomatous and nonseminomatous tumours. Current clinical management for both male and female teenagers and young adults (TYAs) with GCTs remains inconsistent, alternating between paediatric and adult multidisciplinary oncology teams, based on locally defined age cut-offs. We therefore reviewed available literature to determine the biological similarities and differences between GCTs in young children [0-12 years (y)], TYAs (13-24y), and older adults (>24y). GCTs arising in paediatric and adult populations in general showed marked molecular biological differences within identical histological subtypes, whereas there was a distinct paucity of available data for GCTs in the TYA population. These findings highlight that clinical management based simply on chronological age may be inappropriate for TYA and suggests that the optimal future management of GCTs should consider specific molecular biological factors in addition to clinical parameters in the context of patient specific age group specialty.

Aims

The aim of this review was to analyse molecular biological studies of germ cell tumours (GCTs) arising in the paediatric (≤ 12 y) and older adult (> 24 y) age groups for key similarities and differences in an attempt to identify where the biology of GCTs from teenagers and young adult (TYA) patients lies within this framework. The review discusses current knowledge in this area and also highlights gaps that still need to be addressed. In the UK, the TYA age group includes male and females aged 13-24y; accordingly we have used this definition for the purposes of this review). The aims were achieved through a synthesis and analysis of available biological data for GCTs in children, TYAs and older adults. We anticipated that the goal would be to use a combination of both clinical and biological risk factors to improve clinical management decisions for TYA patients with GCTs (Murray et al., 2009).

Background

Regardless of patient age, tumour site and histological subtype, all GCTs are believed to originate from primordial germ cells (PGCs) (Teilum, 1965). The histological appearance of these tumours varies depending on the degree of germ cell differentiation when carcinogenesis takes place (Ross et al., 1999). GCTs can be separated into two major histological subgroups, seminomatous and nonseminomatous tumours. Seminomas (or germinomas, collectively referring to testicular seminomas, ovarian dysgerminomas and extragonadal germinomas) are composed of uniform sheets of tumour cells that resemble undifferentiated PGCs. By contrast, nonseminomas exhibit varying degrees of differentiation that resemble particular cell lineages or stages of embryonic development and include teratomas, embryonal carcinomas (ECs), yolk sac tumours (YSTs) and choriocarcinomas (CHCs) (Palmer et al., 2008). In addition to these, mixed malignant GCTs also exist which are composed of more than one histological subtype.

GCTs are rare, accounting for 4% of all childhood malignancies (Surti et al., 1990). Within this age range, they exhibit a characteristic bimodal age distribution, with a relatively high incidence in the first few years of life, which then declines to very low levels before increasing again in adolescence. As a result, their incidence in the paediatric (4 per million) and older adult (14 per million) populations remains markedly lower than in TYAs (60 per million in males and 8 per million in females) (Figure 1) (Murray et al., 2009).

GCTs can arise as a painless mass in both gonadal and extragonadal sites, including the sacrococcygeum, vagina, mediastinum and central nervous system, typically in the pituitary and pineal region in the case of the latter. It has been proposed that the location of these midline sites may be accounted for by an aberrant path taken by PGCs during their migration from the yolk sac to the gonadal ridge during embryogenesis (Looijenga and Oosterhuis, 1999; Gilbert et al., 2011). In childhood, only 50% of GCTs are gonadal, but this figure increases to 95% in adulthood (Nicholson and Palmer, 2010); where TYA GCTs lie within this spectrum remains unclear.

Historically, malignant GCTs (MGCTs) were rapidly fatal (Nicholson and Palmer, 2010). However, with the advent of effective platinum-based chemotherapy, MGCTs became highly curable, even when diagnosed at advanced clinical stages. The use of bleomycin, etoposide and cisplatin (BEP) in adults with advanced stage GCTs now results in overall survival rates in excess of 90% (Horwich et al., 2006; Krege et al., 2008). Although paediatric GCTs were initially treated with an adult cisplatin-based regime, the neurological and ototoxic sequelae of treatment in young children had a substantial impact upon quality of life, driving the UK to adopt a carboplatin, etoposide and bleomycin (JEB) based approach instead (Mann et al., 1989; Hale et al., 1999). In children, JEB showed comparable survival for all tumour stages (Mann et al., 2000; Stern and Bunin, 2002). Recently, a joint Malignant Germ Cell International Collaboration ('MaGIC') initiative, established between the USA and UK, pooled data from more than 1,100 paediatric patients with GCTs who

had received platinum-based chemotherapy (Hale et al., 2012). The prognostic features of tumours in children and TYAs with GCTs were determined, comparing the outcomes from the US cisplatin-based (PEB) and the UK high dose carboplatin-based approaches (JEB). The results showed that after adjustment for other prognostic factors, the risks of failure for JEB and PEB were not significantly different for patients <11y of age (Hale et al., 2012). In adults, carboplatin therapy has been shown to be inferior to cisplatin, although lower dose carboplatin regimes are typically used in this age group due to the increased risk of myelosuppression (Bokemeyer et al., 1996; Horwich et al., 1997). In younger children, however, higher dose carboplatin resulted in survival rates equivalent to those achieved with cisplatin. These differences may be due to a more aggressive tumour phenotype in older patients, reflect the poor deliverability and tolerance of higher dosing schedules in adults, or a combination of these two factors. It therefore remains unclear as to whether the optimal treatment for TYA GCTs should be as for paediatric cases with relatively high dose carboplatin, or as for adult tumours with cisplatin. If the differing responses to carboplatin therapy are due to tumour biology rather than to treatment dose, then treating TYAs with higher dose carboplatin is likely to be ineffective. In order to know if higher carboplatin doses should be trialled in TYAs, it is necessary to determine whether their GCT biology is most akin to that of paediatric or adult GCTs. This calls for TYA patient-focused clinical studies aimed at establishing their carboplatin dose tolerance, as has been described for acute lymphoblastic leukaemia (Boissel et al., 2003). Ultimately, this may allow clinicians to combine excellent overall survival rates with the potential for reducing the known late neurotoxicity, nephrotoxicity and cardiovascular toxicity associated with cisplatin use. There is ongoing discussion between paediatric and adult (testicular and ovarian) GCT specialists and trialists to consider the application of this in a prospective trial.

The relationship between age, epidemiology and histology in GCTs has been highlighted by Oosterhuis and Looijenga in the design of their classification system (Oosterhuis and Looijenga, 2005). This system divides GCTs into five distinct types. Type I tumours occur in neonates and

infants <5y of age and predominantly comprise nonseminomatous teratomas and YSTs. Type II tumours include both nonseminomatous and seminomatous GCTs and affect the TYA population (Oosterhuis and Looijenga, 2005). As Type III tumours (which comprise spermatocytic seminomas occurring in the elderly male population) are not thought to arise from PGCs (Looijenga et al., 2007), they will not be considered further in this review. Finally, benign ovarian teratomas are categorised as Type IV tumours, while Type V tumours consist of hydatidiform moles (a benign form of gestational trophoblastic disease). According to this classification, GCTs arising in the TYA population should be labelled as Type II tumours. Although this classification is simple, it remains unclear how widely applicable it is to the broad spectrum of paediatric and adult GCTs observed in clinical practice, and whether it can be applied to the TYA GCT population. This review of biological GCT studies in paediatric and adult patients therefore attempts to discern whether a TYA tumour is likely to represent a mature onset childhood-type tumour (Type I), the early onset of a tumour more typical of an older adult (Type II), or in fact represents a distinct group. Importantly, the specific age at which GCTs should be managed as childhood or adult disease remains undefined, resulting in the inconsistent application of treatment regimes and likely poorer outcome for TYAs with GCTs (Murray et al., 2009). We propose that specific features in the biology of these tumours may shed light upon the transition from paediatric to TYA GCTs, and from TYA to adult GCTs, thus enabling rational clinical management decisions to be made for these age groups.

GCT biology

Recent years have witnessed improvements in our overall biological understanding of MGCTs. The similarities and differences in the biology of GCTs within the paediatric and adult age range are discussed, with particular reference to pre-invasive disease, genomic imprinting, chromosomal aberrations, susceptibility loci and both protein-coding and non-protein-coding gene (microRNA) expression.

Pre-invasive disease

All adult testicular GCTs (TGCTs) are thought to arise from a common precursor lesion, originally termed ‘carcinoma *in situ*’ (Skakkebaek, 1972). The currently accepted terminology is ‘intratubular germ cell neoplasia unclassified (IGCNU)’ due to the absence of epithelial differentiation within these lesions. IGCNU appears to develop from undifferentiated PGCs based on studies demonstrating their similar morphology, imprinting patterns and gene expression (van Gurp et al., 1994; Almstrup et al., 2004; Rajpert-De Meyts, 2006). IGCNU does not spontaneously regress and eventually progresses to GCT (Oosterhuis and Looijenga, 2005) in approximately 50% of patients within 5 years (Skakkebaek, 1978; Maase et al., 1986; Giwercman and Skakkebaek, 1993). This progression is highlighted by the fact that IGCNU and TGCT occur at similar frequencies and that IGCNU is found in testicular tissue adjacent to GCTs in 90% of adults (Jacobsen et al., 1981; Giwercman et al., 1991; Dieckmann and Skakkebaek, 1999; Huyghe et al., 2005). This is in marked contrast to paediatric GCTs, where multiple studies have shown IGCNU to be absent (Manivel et al., 1988; Manivel et al., 1989; Jorgensen et al., 1995; Hawkins et al., 1997; Fan and Ulbright, 2012). Indeed, IGCNU in paediatric GCTs has only been described in a small number of isolated cases (Hu et al., 1992; Stamp et al., 1993; Renedo and Trainer, 1994). However, others have questioned whether these paediatric cases truly represent IGCNU (Hawkins et al., 1997). These observations suggest that the presence or absence of IGCNU is likely to prove clinically useful in distinguishing adult from paediatric TGCTs, and, furthermore, that TGCTs in infants and children are likely to arise through a biologically distinct mechanism to TGCTs in adulthood (Hawkins et al., 1997). By contrast, the implications for TYA TGCTs remain unclear given that the exact frequency of IGCNU is unknown in this age group.

Genomic imprinting

Imprinting is an epigenetic phenomenon by which certain genes are selectively differentially expressed in a parental allele-specific manner such that 'silenced' (imprinted) genes are not expressed. This mechanism forms an integral part of the regulation of embryonic cell development (including both in the fetus and placenta). Regulated expression of the imprinted genes *H19* and *IGF-2* (*insulin-like growth factor-2*) is necessary for normal human embryo development (van Gurp et al., 1994). As GCTs are thought to originate from early germ cells, the patterns of allelic expression of the *H19* and *IGF-2* genes were obvious candidates for investigating whether aberrant genomic imprinting is involved in GCT pathogenesis. During normal development, PGCs establish a new sex-specific imprinting pattern. Assuming that GCTs preserve the imprinting status of their cell of origin, they would not be expected to exhibit any imprinting pattern if they have arisen from premeiotic germ cells prior to their arrival at the gonadal ridge (i.e. they would display biallelic expression of both *H19* and *IGF-2*). However, if GCTs develop from PGCs that have entered meiosis instead, they will display a sex-specific imprinting pattern (Schneider et al., 2001). In this regard, the pioneering work of van Gurp and colleagues in adult GCTs reported biallelic expression of *IGF-2* in 55% of TGCTs and biallelic *H19* expression in 70% of TGCTs (van Gurp et al., 1994). The consistent expression of both parental alleles of the *H19* and *IGF2* genes implies that TGCTs may develop from precursor cells that have either erased their imprint or been subjected to a relaxation of the normal monoallelic expression of these genes ensuing from demethylation at the imprinted allele (van Gurp et al., 1994). This is consistent with the finding of Ross and colleagues who also demonstrated loss of imprinting (LOI) of *H19* and *IGF-2* genes in both paediatric and TYA GCTs (Ross et al., 1999). Their results suggested that while LOI of *H19* and *IGF-2* may be common in paediatric and TYA TGCTs, there was a variable pattern of LOI in ovarian GCTs in these age groups, which was proposed to reflect the stage of imprinting during germ cell development at which tumorigenesis occurred.

Schneider and colleagues reported a multipoint imprinting analysis of the *H19*, *IGF-2* and *SNRPN* (*small nuclear ribonucleoprotein polypeptide N*) genes in 42 gonadal and extragonadal GCTs in children and TYAs (Schneider et al., 2001). Forty-six percent of informative tumours showed biallelic *H19* expression, while 47% showed biallelic *IGF-2* expression. Eighty-four percent of gonadal GCTs and 81% of extragonadal GCTs showed absence of *SNRPN* methylation consistent with LOI. Crucially, the frequency of LOI between gonadal and extragonadal GCTs was comparable, and did not correlate with patient age. These results suggest that both gonadal and extragonadal GCTs are derived from PGCs, and have consistent LOI of *SNRPN* and partial LOI of *H19* and *IGF-2*. When compared with adult TGCTs, the frequency of LOI was lower in paediatric testicular and extragonadal GCTs, and in TYA ovarian GCTs. As such, LOI of *H19* and *IGF-2* has been found in the paediatric, TYA and older adult populations although LOI patterns varied across the three populations (van Gurp et al., 1994; Nonomura et al., 1997; Ross et al., 1999; Schneider et al., 2001). Although this suggests that these tumours all arise from PGCs, this may nevertheless occur at different stages of germ cell development. In particular, the partial LOI in paediatric GCTs indicates that these tumours may likely arise from a more immature germ cell, and that TYA GCTs are more similar to GCTs in older adults (Schneider et al., 2001).

Although many chromosomal aberrations in GCTs have been established, the major tumour suppressor genes (TSGs) associated with such areas of physical or functional genomic losses remain to be identified. It has been demonstrated that in adult TGCTs, methylation of the TSGs *APC* (*adenomatous polyposis coli*), *RASSF1A* (*Ras association domain family 1 isoform A*) and *MGMT* (*O-6-methylguanine-DNA methyltransferase*) occurs more frequently in nonseminomas compared to seminomas (Honorio et al., 2002). Methylation of the TSGs *RASSF1A* and *HIC1* (*hypermethylated in cancer 1*) have been shown to be associated with treatment resistance, whilst methylation of *MGMT* is associated with platinum sensitivity (Koul et al., 2004). In paediatric GCTs *APC*, *RASSF1A* and *HIC1* have all been shown to be significantly more methylated in YSTs

than in germinomas (Jeyapalan et al., 2011). In paediatric YSTs, methylation of the TSG *RUNX3* (*Runt-related transcription factor 3*) has been reported, an imprint which has not been reported in adult YSTs, and which may therefore contribute to the pathogenesis of this childhood GCT subtype (Kato et al., 2003; Furukawa et al., 2009). In line with this, a recent study examined the methylation status of 15 selected TSGs in 28 paediatric GCT samples and found deletion of the 1p36-p35 regions in 19 (68%) cases (Ichikawa et al., 2013). *RUNX3*, located at 1p36, was methylated in 16 of the 19 (84%) tumours, demonstrating that hypermethylation appears to be the principal mechanism of inactivation of this gene in paediatric GCTs.

Chromosomal aberrations

The ploidy status of GCTs was reviewed by Veltman and colleagues, who found that TGCTs in TYA and older adult populations were usually aneuploid, with possible overall differences between histological subtypes (Veltman et al., 2003). Seminomas were found to be hypertriploid, whilst nonseminomas were hypotriploid (Looijenga, 2001). By contrast, the pattern of genomic gains and losses between the histological subtypes was broadly similar, except for gains of chromosome 15 in seminomas compared to nonseminomas (Looijenga, 2001). In the paediatric population, DNA ploidy was found to be variable (diploid, triploid and tetraploid) (Looijenga and Oosterhuis, 1999; Bussey et al., 2001; Schneider et al., 2001) while ovarian GCTs are mostly diploid, independent of age (Surti et al., 1990).

Over three decades ago, Atkin and Baker detected an isochromosome of the short arm of chromosome 12 [i(12p)] in TGCTs based on the karyotypic analysis of metaphase chromosomes (Atkin and Baker, 1982). It was subsequently established that i(12p) occurs in 80% of adult TGCTs, while gain of 12p genomic material is invariably present in the other 20% of cases (Looijenga, 2001). Indeed, i(12p) appears to be a characteristic cytogenetic event in both gonadal and extragonadal adult GCTs (Oosterhuis et al., 1997; Bussey et al., 1999; Kraggerud et al., 2000;

Schneider et al., 2002; McKenney et al., 2007; Palmer et al., 2007). By contrast, a gain of 12p was until recently thought to occur only rarely in paediatric cases. In early studies, Perlman and colleagues used comparative genomic hybridization (CGH) to screen for genetic abnormalities in 16 paediatric (<4y) YSTs (10 testicular site and six extragonadal) (Perlman et al., 2000). Gain of 12p was seen in only one of their 16 tumours (a testicular tumour), highlighting the dissimilarity in genomic aberration profile between paediatric and adult GCTs. Bussey and colleagues assessed 12p status in 18 male GCTs, five of which had a 12p gain (Bussey et al., 2001). More specifically, no 12p gain was seen in tumours from patients <9y of age. This led the authors to postulate that gain of 12p was significantly associated with patient age and, accordingly, may relate to the physiological changes brought about by the onset of puberty. By contrast, a more recent paediatric (0-16y) study (Palmer et al., 2007) analysed 11 seminomas, 22 YSTs and one metastatic EC from a range of anatomical sites from 12 male and 22 female patients, including a large proportion of ovarian GCTs. Chromosome 12p gain was identified in 44% of cases, with an incidence increasing with age (29% of those <5y, compared with 53% in those 5-16y). Again, these findings suggest that genomic copy number imbalances can distinguish GCT subgroups primarily by age, rather than by tumour site or histology.

Given that the only consistent structural chromosomal abnormality in TGCTs is 12p gain [predominantly due to i(12p)], the implication is that this genomic aberration is associated with MGCT development. In support of this theory, studies of IGCNU demonstrate a similar pattern of overall gains and losses to those observed in invasive TGCTs, except for 12p gain (Rosenberg et al., 2000; Summersgill et al., 2001; Ottesen et al., 2003).

With regard to paediatric tumours, Schneider and colleagues identified some of the key genomic changes found in mediastinal GCTs in this age range (Schneider et al., 2002). They also reported

that mediastinal teratomas exhibited no genomic gains or losses. By contrast, in young children (<8y of age) mediastinal MGCTs displayed the same pattern of gains and losses as those found in age-matched sacrococcygeal and testicular GCTs, including gain of 1q, 3, and 20q, and loss of 1p, 4q and 6q. Similarly, for children ≥ 8 y of age, mediastinal MGCTs showed identical profiles to gonadal GCTs, particularly with respect to 12p gain. The authors stated that mediastinal MGCTs arising in TYAs carried a poorer prognosis compared to those affecting young children (<8y), likely due to biological age-dependent differences in these tumours (Schneider et al., 2002). Indeed, studies have identified differences in the clinical behaviour of tumours with the same histological subtype, suggesting that these may reflect inherent differences in tumour biology amongst different age groups (Schneider et al., 2002; Marina et al., 2006).

Cytogenetic data on ovarian MGCTs are limited, and the frequency of 12p gain in female GCTs in different age groups is not well established. In paediatric female GCTs, the presence of i(12p) is infrequent, with few, individual cases being reported (Speleman et al., 1990; Speleman et al., 1992; Bussey et al., 1999; Bussey et al., 2001). By contrast, i(12p) is more frequent in adult female GCTs, particularly in dysgerminomas (Atkin and Baker, 1987; Jenkyn and McCartney, 1987; Rodriguez et al., 1995; Dal Cin et al., 1998; Riopel et al., 1998). Using CGH, Kraggerud and colleagues confirmed that the most frequent change in dysgerminomas was gain of 12p, most commonly as i(12p) (Kraggerud et al., 2000). The frequency of 12p gain was distributed largely evenly amongst the age groups (12-32y). More recently, Cossu-Rocca and co-workers used fluorescence *in situ* hybridisation (FISH) to demonstrate chromosome 12p abnormalities in 17/21 (81%) dysgerminomas (Cossu-Rocca et al., 2006). The chromosome abnormalities observed included over-representation of 12p in 5/21 (24%) of cases, and i(12p) in 16/21 (76%) of cases. Patient ages ranged from 10 to 50y, although potential age-dependent variations were not considered. In conclusion, these results suggest that, similarly to males, 12p copy number gains may be associated

with increasing age in females, although more studies are required to confirm this, particularly in the TYA age group.

Other genomic copy number imbalances in MGCTs have also been described in adults. Commonly observed imbalances include gains on chromosomes 1, 7, 8, 12, 21, and X, as well as losses on 4, 5, 11, 13 and 18. However, none of these reported abnormalities are as consistent as gain of 12p (van Echten et al., 1995; Ottesen et al., 1997; Summersgill et al., 1998; Looijenga et al., 2000; Kraggerud et al., 2002; Ichikawa et al., 2013). A wide range of copy number imbalances has also been described in paediatric MGCTs, including gains on chromosomes 1q, 2p, 3, 7, 8, 13, 14, 20q, 21, and X, as well as losses on 1p36, 4q, 6q, 11, 13 and 18 (Perlman et al., 2000; Schneider et al., 2002; Oosterhuis and Looijenga, 2005; Palmer et al., 2007).

Susceptibility loci

Recent genome wide association studies (GWAS) have identified 6 susceptibility gene loci in the development of TGCTs in adults; *activating transcription factor 7 interacting protein (ATF7IP)*, *BCL-2 antagonist/killer 1 (BAKI)*, *doublesex and mab-3 related transcription factor 1 (DMRT1)*, *KIT ligand (KITLG)*, *sprouty homolog 4 (SPRY4)* and *telomerase reverse transcriptase (TERT)* (Kanetsky et al., 2009; Rapley et al., 2009; Turnbull et al., 2010). However the specific mechanisms through which these genes result in tumorigenesis remain unknown. All six genes play a central role in normal PGC development (Gilbert et al., 2011) and, more specifically, may also be important in susceptibility to MGCTs in the TYA (*SPRY4*, *BAKI*) and paediatric (*BAKI*) populations (Poynter et al., 2012).

Protein-coding gene expression

One group studied the global messenger RNA (mRNA) protein-coding gene expression patterns of paediatric MGCTs, and compared them with published data on adult TGCTs (Palmer et al., 2008).

The results showed that paediatric MGCTs differed at the gene expression level according to their histology (Figure 2), irrespective of the anatomical site of the primary disease or the age of the patient (in the paediatric group). Similarly, adult TGCTs were also differentiated by their histology although, importantly, their gene expression profiles differed from their respective paediatric GCT subtype counterparts (Figure 3) (Palmer et al., 2008). These results demonstrated that paediatric and adult tumours with comparable histology can differ in their gene expression profiles, lending weight to the suggestion that the clinical management of these entities should be tailored accordingly. It is possible that differences in gene expression between paediatric and adult germinomas was attributable to contributions from tumour site in addition to patient age, given that the paediatric germinomas studied included both ovarian and extragonadal cases, which were compared to adult testicular seminomas. However, the fact that the YSTs were predominantly testicular in paediatric cases and that clear expression differences were still noted when compared to their adult testicular counterparts, makes a contribution from tumour site unlikely. This notion is supported by other studies, which suggest that site-related gene expression differences in GCTs are likely to be minimal (Korkola et al., 2009).

Korkola and co-workers also reported an mRNA gene expression signature that was predictive of TGCT clinical outcome in adult patients (Korkola et al., 2009). Genome-wide expression profiling was performed on 108 nonseminomatous GCTs from patients treated with cisplatin-based chemotherapy, 33 of which had poor clinical outcomes. More specifically, the results indicated that signatures reflecting enhanced immune function and repression of differentiation were associated with good outcome, while those representing active differentiation were associated with poor outcome, such that the gene expression signature derived added independent prognostic accuracy to existing clinically-used systems. Although patient age ranged from 15 to 65y (median age 29.4y) in the training set and 16 to 45y (median age 28.1y) in the validation set, the majority of cases were from older adults (>24y). Consequently, it is unclear whether this model could be applied directly to the paediatric and TYA setting. Large, future studies aiming to identify how such gene

expression profiles correlate with clinical outcome in paediatric and TYA populations (including direct comparisons with the above prognostic signatures for adult GCTs) are therefore required and should help to determine whether the use of transcriptomic signatures as adjuncts to clinical risk factors may inform treatment choice.

Non-protein-coding gene (microRNA) expression

MicroRNAs are short, non-protein-coding RNAs that modulate the expression of protein-coding genes through interactions with the 3' untranslated region (3'UTR) binding sites of target mRNAs, and whose expression profiles are dysregulated in cancer (Shenouda and Alahari, 2009). One group (Voorhoeve et al., 2006) demonstrated that the miR-371~373 cluster was highly expressed in adult TGCTs, suggesting that this may act as a potential novel oncogene system in TGCTs via inhibition of the tumour suppressor gene *LATS2*. Consistent with this observation, a subsequent study determined the levels of a restricted range of 156 microRNAs in adult MGCTs, and confirmed miR-371~373 over-expression compared with normal testis controls (Gillis et al., 2007).

Palmer and colleagues determined the global microRNA profiles of 48 paediatric and TYA samples (Palmer et al., 2010) and compared these with data from adult GCTs. The results showed that the majority of differentially expressed microRNAs in paediatric GCTs were down-regulated, consistent with observations in other malignancies. However, the most significant finding was that the miR-371~373 and miR-302 clusters were over-expressed in all MGCTs, independent of patient age (paediatric, TYA or adult), tumour histological subtype (YST/ seminoma/EC) or anatomical site (gonadal/extragenital). Furthermore, the expression levels of eight main microRNA members from the miR-371~373 and miR-302 clusters accurately separated MGCTs from non-malignant samples, suggesting potential clinical use as highly sensitive and specific universal biomarkers of MGCTs (Figure 4) (Palmer et al., 2010).

Mechanisms for associations between age and GCT biology

Schneider and colleagues' study suggests a possible transition in biology and prognosis in younger versus older children, and proposes that this coincides with the increase in GCT incidence from 8y of age (Figure 1), although the aetiological factors responsible for this potential transition remain unknown (Schneider et al., 2002). One reasonable presumption is that the transition may follow endocrine changes linked to the onset of puberty. Increasing levels of oestrogen and testosterone associated with this event trigger the completion of the oocyte's first meiotic division and stimulate spermatocytes to enter meiosis, respectively. It is tempting to speculate that deregulations in this tightly orchestrated physiological process may in some way contribute to germ cell tumorigenesis. Thus, Schneider's study provides some genomic evidence pointing to a possible biological transition which may occur between GCTs arising in younger children ('paediatric-like') and those in older children ('TYA-like') and that such changes may, at least in part, be caused by the hormonal *milieu* surrounding the onset of puberty (Schneider et al., 2002).

Conclusion

We have reviewed the current knowledge regarding the biological similarities and differences in GCTs in paediatric and adult populations, and where GCTs in patients aged 13-24y may fit within this framework (summarised in Table 1). A persisting conundrum regards the mechanism and the timing of the 'tumour biological shift' between these populations. Should the inherent differences between paediatric and adult GCTs be accounted for by the onset of puberty, the majority of TYA GCTs may in fact be biologically more similar to adult tumours than previously thought (Schneider et al., 2002). Accordingly, we propose that future studies should correlate GCT biology with documented pubertal status (based on markers including steroid hormone levels and contralateral testicular size in males, as measured by ultrasound at the time of TGCT diagnosis) to investigate this theory. By contrast, there is no analogous physiological/developmental milestone that could

demarcate the presumptive transition between TYA and adult GCTs and their categorisation as distinct entities.

The equivocal boundaries between these groups are further complicated by age group-related differences in GCT clinical presentation and prognostic outlook, in turn reflecting variations in deliverability, tolerance and response to platinum-based therapies. In these cases, identifying tumour-specific biological characteristics (rather than simple patient age-based stratification) may be of greater value in prospectively identifying prognostic risk groups and support the development of novel, more targeted therapeutic regimes that have increased efficacy in poor prognosis tumours and/or cause less long-term toxicity in good prognosis patients.

As a result of these challenges, the authors have established a UK-based GCT consortium, which has generated a clinically annotated database supporting a tumour bank of over 400 MGCTs in patients within the TYA age range. The consortium proposes to utilise recent advances in technology to perform DNA, RNA and immunohistochemical-based studies on the associated paraffin-embedded MGCT tissues. Biological changes will be related to patient age at diagnosis across different histological subtypes and prognostic groups (including platinum-resistant cases), and compared with published results for paediatric and adult tumours. Both these and other future studies should aim to define age-specific GCT biological characteristics, including for TYA GCTs. Such an approach will help identify whether TYA GCTs represent a distinct clinical, physiological and biological entity, assist their optimal future management and may identify biological targets suitable for the development of novel therapeutic agents which may improve clinical outcomes for this under-investigated patient group.

Author Disclosure Statement

No competing financial interests exist.

Table and Figure Legends

Table 1: The biological characteristics of GCTs in paediatric, TYA and older adult age populations

Figure 1: GCT incidence rates by anatomical site, per million person years at risk, for patients aged 0-79y in England, 1995-2003 (from Murray et al., 2009).

Figure 2: Heatmap analysis of paediatric malignant GCT protein-coding gene expression data (from Palmer et al., 2008). The heatmap depicts all 657 differentially expressed probes in the 27 malignant GCTs analysed. In the heatmap, red represents relative mRNA over-expression and blue represents under-expression.

Figure 3: Heatmap comparison of protein-coding gene expression in paediatric malignant GCTs compared with pure (i.e. not mixed) adult malignant TGCTs (from Palmer et al., 2008). Twenty-seven paediatric tumours are compared with 20 adult malignant GCTs for (A) yolk sac tumours and (B) seminomas. In the heatmap, red represents relative mRNA over-expression and blue represents under-expression.

Figure 4: Differential expression of the miR-371~373 and miR-302 clusters in malignant GCTs (from Palmer et al., 2010). Hierarchical clustering analysis based on the eight main microRNAs from the miR-371~373 and miR-302 clusters segregates (A) paediatric and (B) adult malignant GCT samples from non-malignant controls (comprising benign teratomas and normal gonadal controls).

Table 1:

GCTs ^a	Age group	Biological Characteristics
Testicular	Paediatric	<ul style="list-style-type: none"> • Predominantly teratoma/YST^b histology (Oosterhuis and Looijenga 2005) • Diploid (teratoma) or aneuploid (YST) (Oosterhuis and Looijenga 2005; Veltman et al., 2003) • i(12p)^c is seen but to a lesser extent than in the TYA^d population (Palmer et al., 2007) • Other chromosomal abnormalities include deletions of 1p, 4, 6q and gain of 1q, 12p13 and 20q (Oosterhuis and Looijenga 2005) • Overexpression of miR^e-371~373 and miR-302 clusters (Palmer et al., 2010)
	TYA	<ul style="list-style-type: none"> • Predominantly nonseminomatous histology, having a median age of occurrence of 25y^f (Oosterhuis and Looijenga 2005) • Aneuploid (Oosterhuis and Looijenga 2005; Veltman et al., 2003) • i(12p) is present in 80% of these tumours (Looijenga 2001) • Other chromosomal abnormalities include gain of 7, 8, 21 and loss of 1p, 11, 13 and 18 (Oosterhuis and Looijenga 2005) • IGCNU^g seen in 90% (Dieckmann and Skakkebaek 1999; Jacobsen et al., 1981) • Overexpression of miR-371~373 and miR-302 clusters (Palmer et al., 2010)
	Older Adults	<ul style="list-style-type: none"> • Predominantly seminomatous histology, having a median age of occurrence of 35y (Oosterhuis and Looijenga 2005) • Aneuploid (Oosterhuis and Looijenga 2005) • i(12p) is present in 80% of these tumours (Looijenga 2001) • Other chromosomal abnormalities include gain of 7, 8, 21 and loss of 1p, 11, 13 and 18 (Oosterhuis and Looijenga 2005) • IGCNU seen in 90% (Dieckmann and Skakkebaek 1999; Jacobsen et al., 1981) • Overexpression of miR-371~373 and miR-302 clusters (Palmer et al., 2010)
Ovarian	Paediatric	<ul style="list-style-type: none"> • Incidence is very low before age 5y, then increases gradually to peak in early teens (Nicholson and Palmer 2010) • Predominantly teratoma/YST (Oosterhuis and Looijenga 2005) • Presence of i(12p) is infrequent, with only 4 cases reported (Bussey et al., 2001; Bussey et al., 1999; Speleman et al., 1990; Speleman et al., 1992) • Overexpression of miR-371~373 and miR-302 clusters (Palmer et al., 2010)
	TYA	<ul style="list-style-type: none"> • Incidence peaks in early teens (Nicholson and Palmer 2010) • Most commonly teratomas and dysgerminomas (Nicholson and Palmer 2010) • Diploid (Surti et al., 1990) • Overexpression of miR-371~373 and miR-302 clusters (Palmer et al., 2010) • i(12p) is present in a similar proportion to their testicular counterparts (Kraggerud et al., 2000)
	Older Adults	<ul style="list-style-type: none"> • Incidence decreases as age increases (Jha and Karki 2008)
Abbreviations	^a Germ cell tumour, ^b Yolk sac tumour, ^c isochromosome 12p, ^d teenager and young adult, ^e microRNA, ^f years, ^g intratubular germ cell neoplasia unclassified.	

Figure 1:




Figure 2:




Figure 3:





Figure 4:



References

- Almstrup K, Høi-Hansen CE, Wirkner U, Blake J, Schwager C, Ansorge W, Nielsen JE, Skakkebaek NE, Rajpert-De Meyts E, Leffers H. 2004. Embryonic stem cell-like features of testicular carcinoma in situ revealed by genome-wide gene expression profiling. *Cancer Res* 64:4736-4743.
- Atkin NB, Baker MC. 1982. Specific chromosome change, i(12p), in testicular tumours? *Lancet* 2:1349.
- Atkin NB, Baker MC. 1987. Abnormal chromosomes including small metacentrics in 14 ovarian cancers. *Cancer Genet Cytogenet* 26:355-361.
- Boissel N, Auclerc M, Lhéritier V, Perel Y, Thomas X, Leblanc T, Rousselot P, Cayuela J, Gabert J, Fegueux N, Piguet C, Huguet-Rigal F, Berthou C, Boiron J, Pautas C, Michel G, Fièrè D, Leverger G, Dombret H, Baruchel A. 2003. Should adolescents with acute lymphoblastic leukemia be treated as old children or young adults? Comparison of the French FRALLE-93 and LALA-94 trials. *J Clin Oncol.* 1; 21:774-780.
- Bokemeyer C, Köhrmann O, Tischler J, Weißbach L, Ràth U, Haupt A, Schöffski P, Harstrick A, Schmoll H-J. 1996. A randomised trial of cisplatin, etoposide and bleomycin (PEB) versus carboplatin, etoposide and bleomycin (CEB) for patients with 'good risk' metastatic non-seminomatous germ cell tumours. *Ann Oncol* 7:1015-1021.
- Bussey KJ, Lawce HJ, Himoe E, Shu XO, Suijkerbuijk RF, Olson SB, Magenis RE. 2001. Chromosomes 1 and 12 abnormalities in pediatric germ cell tumors by interphase fluorescence in situ hybridization. *Cancer Genet Cytogenet* 125:112-118.
- Bussey KJ, Lawce HJ, Olson SB, Arthur DC, Kalousek DK, Krailo M, Giller R, Heifetz S, Womer R, Magenis RE. 1999. Chromosome abnormalities of eighty-one pediatric germ cell tumors: Sex, age, site, and histopathology related differences a Children's Cancer Group study. *Genes Chromosomes Cancer* 25:134-146.

- Cossu-Rocca P, Zhang S, Roth LM, Eble JN, Zheng W, Karim FW, Michael H, Emerson RE, Jones TD, Hattab EM, Cheng L. 2006. Chromosome 12p abnormalities in dysgerminoma of the ovary: a FISH analysis. *Mod Pathol* 19:611-615.
- Dal Cin P, Dei Tos AP, Qi H, Giannini C, Furlanetto A, Longatti PL, Marynen P, Van den Berghe H. 1998. Immature teratoma of the pineal gland with isochromosome 12p. *Acta Neuropathol (Berl)* 95:107-110.
- Dieckmann KP, Skakkebaek NE. 1999. Carcinoma In situ of the testis: Review of biological and clinical features. *Int J Cancer* 83:815-822.
- Fan R, Ulbright T. 2012. Does intratubular germ cell neoplasia, unclassified type exist in prepubertal, cryptorchid testes? *Fetal Pediatr Pathol* 31:21-24.
- Gilbert D, Rapley E, Shipley J. 2011. Testicular germ cell tumours: predisposition genes and the male germ cell niche. *Nat Rev Cancer* 11:278-288.
- Gillis AJM, Stoop HJ, Hersmus R, Oosterhuis JW, Sun Y, Chen C, Guenther S, Sherlock J, Veltman I, Baeten J, van der Spek PJ, de Alarcon P, Looijenga LHJ. 2007. High-throughput microRNAome analysis in human germ cell tumours. *J Pathol* 213:319-328.
- Giwercman A, Müller J, Skakkebaek NE. 1991. Prevalence of carcinoma in situ and other histopathological abnormalities in testes from 399 men who died suddenly and unexpectedly. *J Urol* 145:77-80.
- Giwercman A, Skakkebaek N. 1993. Carcinoma in situ of the testis: biology, screening and management. *Eur Urol* 23:19-21.
- Hale GA, Marina NM, Jones-Wallace D, Greenwald CA, Jenkins JJ, Rao BN, Luo X, Hudson MM. 1999. Late Effects of Treatment for Germ Cell Tumors During Childhood and Adolescence. *J Pediatr Hematol Oncol* 21.
- Hale J, Olson T, Nicholson J, Dang H, Krailo M, Billmire D, Donachie P, Thornton C, Rodriguez-Galindo C, Frazier A. 2012. Carboplatin (CBP) versus cisplatin (CP) within prognostic

- groups in pediatric extracranial malignant germ cell tumors (MGCTs). ASCO Annual Meeting. Chicago, Illinois
- Hawkins E, Heifetz SA, Giller R, Cushing B. 1997. The prepubertal testis (prenatal and postnatal): Its relationship to intratubular germ cell neoplasia: A combined pediatric oncology group and children's cancer study group. *Hum Pathol* 28:404-409.
- Horwich A, Shipley J, Huddart R. 2006. Testicular germ-cell cancer. *Lancet* 367:754-765.
- Horwich A, Sleijfer DT, Fossa SD, Kaye SB, Oliver RT, Cullen MH, Mead GM, de Wit R, de Mulder PH, Dearnaley DP, Cook PA, Sylvester RJ, Stenning SP. 1997. Randomized trial of bleomycin, etoposide, and cisplatin compared with bleomycin, etoposide, and carboplatin in good-prognosis metastatic nonseminomatous germ cell cancer: a Multiinstitutional Medical Research Council European Organization for Research and Treatment of Cancer Trial. *J Clin Oncol* 15:1844-1852.
- Hu LM, Phillipson J, Barsky SH. 1992. Intratubular Germ Cell Neoplasia in Infantile Yolk Sac Tumor Verification by Tandem Repeat Sequence In Situ Hybridization. *Diagn Mol Pathol* 1.
- Huyghe E, Soulie M, Escourrou G, Mieusset R, Plante P, Thonneau P. 2005. Conservative management of small testicular tumour relative to carcinoma in situ prevalence. *J Urol* 173:820-823.
- Ichikawa M, Arai Y, Haruta M, Furukawa S, Ariga T, Kajii T, Kaneko Y. 2013. Meiosis error and subsequent genetic and epigenetic alterations invoke the malignant transformation of germ cell tumor. *Genes Chromosomes Cancer* 52:274-286.
- Jacobsen GK, Henriksen OB, von der Maase H. 1981. Carcinoma in situ of testicular tissue adjacent to malignant germ-cell tumors: a study of 105 cases. *Cancer Genet Cytogenet* 47:2660-2662.
- Jenkyn DJ, McCartney AJ. 1987. A chromosome study of three ovarian tumors. *Cancer Genet Cytogenet* 26:327-337.

Jorgensen N, Muller J, Giwercman A, Visfeldt J, Møller H, Skakkebaek N. 1995. DNA content and expression of tumour markers in germ cells adjacent to germ cell tumours in childhood: probably a different origin for infantile and adolescent germ cell tumours. *J Pathol* 176:269-278.

Kanetsky PA, Mitra N, Vardhanabhuti S, Li M, Vaughn DJ, Letrero R, Ciosek SL, Doody DR, Smith LM, Weaver J, Albano A, Chen C, Starr JR, Rader DJ, Godwin AK, Reilly MP, Hakonarson H, Schwartz SM, Nathanson KL. 2009. Common variation in KITLG and at 5q31.3 proximate to SPRY4 predispose to testicular germ cell cancer. *Nat Genet* 41:811-815.

Korkola JE, Houldsworth J, Feldman DR, Olshen AB, Qin LX, Patil S, Reuter VE, Bosl GJ, Chaganti RS. 2009. Identification and validation of a gene expression signature that predicts outcome in adult men with germ cell tumors. *J Clin Oncol* 27:5240-5247.

Kraggerud SM, Skotheim RI, Szymanska J, Eknæs M, Fosså SD, Stenwig AE, Peltomäki P, Lothe RA. 2002. Genome profiles of familial/bilateral and sporadic testicular germ cell tumors. *Genes Chromosomes Cancer* 34:168-174.

Kraggerud SM, Szymanska J, Abeler VM, Kaern J, Eknæs M, Heim S, Teixeira MR, Trope CG, Peltomaki P, Lothe RA. 2000. DNA copy number changes in malignant ovarian germ cell tumors. *Cancer Res* 60:3025-3030.

Krege S, Beyer J, Souchon R, Albers P, Albrecht W, Algaba F, Bamberg M, Bodrogi I, Bokemeyer C, Cavallin-Ståhl E, Classen J, Clemm C, Cohn-Cedermark G, Culine S, Daugaard G, De Mulder PH, De Santis M, de Wit M, de Wit R, Derigs HG, Dieckmann KP, Dieing A, Droz JP, Fenner M, Fizazi K, Flechon A, Fosså SD, del Muro XG, Gauler T, Geczi L, Gerl A, Germa-Lluch JR, Gillessen S, Hartmann JT, Hartmann M, Heidenreich A, Hoeltl W, Horwich A, Huddart R, Jewett M, Joffe J, Jones WG, Kisbenedek L, Klepp O, Kliesch S, Koehrmann KU, Kollmannsberger C, Kuczyk M, Laguna P, Galvis OL, Loy V, Mason MD, Mead GM, Mueller R, Nichols C, Nicolai N, Oliver T, Ondrus D, Oosterhof GO, Ares LP,

- Pizzocaro G, Pont J, Pottek T, Powles T, Rick O, Rosti G, Salvioni R, Scheiderbauer J, Schmelz HU, Schmidberger H, Schmoll HJ, Schrader M, Sedlmayer F, Skakkebaek NE, Sohaib A, Tjulandin S, Warde P, Weinknecht S, Weissbach L, Wittekind C, Winter E, Wood L, H vdM. 2008. European Consensus Conference on Diagnosis and Treatment of Germ Cell Cancer: A Report of the Second Meeting of the European Germ Cell Cancer Consensus group (EGCCCG): Part I. *Eur Urol* 53:478-496.
- Looijenga L, Stoop H, Hersmus R, Gillis A, Wolter Oosterhuis J. 2007. Genomic and expression profiling of human spermatocytic seminomas: pathogenetic implications. *Int J Androl* 30:328-336.
- Looijenga LH. 2001. Testis: Germ Cell tumours. *Atlas Genetic Cytogenetic Oncology Haematology*.
- Looijenga LH, Oosterhuis JW. 1999. Pathogenesis of testicular germ cell tumours. *Rev Reprod* 4:90-100.
- Looijenga LH, Rosenberg C, van Gurp R, Geelen E, van Echten-Arends J, de Jong B, Mostert M, Wolter Oosterhuis J. 2000. Comparative genomic hybridization of microdissected samples from different stages in the development of a seminoma and a non-seminoma. *J Pathol* 191:187-192.
- Maase Hvd, Rørth M, Walbom-Jørgensen S, Sørensen B, Christophersen I, Hald T, Jacobsen G, Berthelsen J, Skakkebaek N. 1986. Carcinoma in situ of contralateral testis in patients with testicular germ cell cancer: study of 27 cases in 500 patients. *Br Med J* 293:1398-1401.
- Manivel JC, Reinberg Y, Niehans GA, Fraley EE. 1989. Intratubular germ cell neoplasia in testicular teratomas and epidermoid cysts. Correlation with prognosis and possible biologic significance. *Cancer* 64:715-720.
- Manivel JC, Simonton S, Wold LE, Dehner LP. 1988. Absence of intratubular germ cell neoplasia in testicular yolk sac tumors in children. A histochemical and immunohistochemical study. *Arch Pathol Lab Med* 112:641-645.

- Mann J, Pearson D, Barrett A, Raafat F, Barnes J, Wallendszus K. 1989. Results of the United Kingdom Children's Cancer Study Group's malignant germ cell tumor studies. *Cancer* 63:1657-1667.
- Mann JR, Raafat F, Robinson K, Imeson J, Gornall P, Sokal M, Gray E, McKeever P, Hale J, Bailey S, Oakhill A. 2000. The United Kingdom Children's Cancer Study Group's second germ cell tumor study: carboplatin, etoposide, and bleomycin are effective treatment for children with malignant extracranial germ cell tumors, with acceptable toxicity. *J Clin Oncol* 18:3809-3818.
- Marina N, London WB, Frazier AL, Lauer S, Rescorla F, Cushing B, Malogolowkin MH, Castleberry RP, Womer RB, Olson T. 2006. Prognostic Factors in Children With Extragonadal Malignant Germ Cell Tumors: A Pediatric Intergroup Study. *J Clin Oncol* 24:2544-2548.
- McKenney JK, Heerema-McKenney A, Rouse RV. 2007. Extragonadal germ cell tumors: a review with emphasis on pathologic features, clinical prognostic variables, and differential diagnostic considerations. *Adv Anat Pathol* 14:69-92.
- Murray M, Fern L, Stark D, Eden T, Nicholson J. 2009. Breaking down barriers: improving outcomes for teenagers and young adults with germ cell tumours. *Oncol Rev* 3:201-206.
- Nicholson J, Palmer R. 2010. *Germ Cell Tumours. Pediatric hematology and Oncology: Scientific Principles and Clinical Principles.* Blackwell Publishing Ltd.
- Nonomura N, Miki T, Nishimura K, Kanno N, Kojima Y, Okuyama A. 1997. Altered Imprinting of the H19 and Insulin-Like Growth Factor II Genes in Testicular Tumors. *J Urol* 157:1977-1979.
- Oosterhuis JW, Looijenga LH. 2005. Testicular germ-cell tumours in a broader perspective. *Nat Rev Cancer* 5:210-222.

- Oosterhuis JW, Looijenga LHJ, van Echten J, de Jong B. 1997. Chromosomal constitution and developmental potential of human germ cell tumors and teratomas. *Cancer Genet Cytogenet* 95:96-102.
- Ottesen AM, Kirchhoff M, De-Meyts ER, Maahr J, Gerdes T, Rose H, Lundsteen C, Petersen PM, Philip J, Skakkebaek NE. 1997. Detection of chromosomal aberrations in seminomatous germ cell tumours using comparative genomic hybridization. *Genes Chromosomes Cancer* 20:412-418.
- Ottesen AM, Skakkebaek NE, Lundsteen C, Leffers H, Larsen J, Rajpert-De Meyts E. 2003. High-resolution comparative genomic hybridization detects extra chromosome arm 12p material in most cases of carcinoma in situ adjacent to overt germ cell tumors, but not before the invasive tumor development. *Genes Chromosomes Cancer* 38:117-125.
- Palmer RD, Barbosa-Morais NL, Gooding EL, Muralidhar B, Thornton CM, Pett MR, Roberts I, Schneider DT, Thorne N, Tavaré S, Nicholson JC, Coleman N. 2008. Pediatric malignant germ cell tumors show characteristic transcriptome profiles. *Cancer Res* 68:4239-4247.
- Palmer RD, Foster NA, Vowler SL, Roberts I, Thornton CM, Hale JP, Schneider DT, Nicholson JC, Coleman N. 2007. Malignant germ cell tumours of childhood: new associations of genomic imbalance. *Br J Cancer* 96:667-676.
- Palmer RD, Murray MJ, Saini HK, van Dongen S, Abreu-Goodger C, Muralidhar B, Pett MR, Thornton CM, Nicholson JC, Enright AJ, Coleman N. 2010. Malignant germ cell tumors display common microRNA profiles resulting in global changes in expression of messenger RNA targets. *Cancer Res* 70:2911-2923.
- Perlman E, Hu J, Ho D, Cushing B, Lauer S, Castleberry R. 2000. Genetic analysis of childhood endodermal sinus tumors by comparative genomic hybridization. *J Pediatr Hematol Oncol* 22:100-105.

- Poynter JN, Hooten AJ, Frazier AL, Ross JA. 2012. Associations between variants in *KITLG*, *SPRY4*, *BAK1*, and *DMRT1* and pediatric germ cell tumors. *Genes Chromosomes Cancer* 51:266-271.
- Rajpert-De Meyts E. 2006. Developmental model for the pathogenesis of testicular carcinoma in situ: genetic and environmental aspects. *Hum Reprod Update* 12:303-323.
- Rapley EA, Turnbull C, Olama AAA, Dermitzakis ET, Linger R, Huddart RA, Renwick A, Hughes D, Hines S, Seal S, Morrison J, Nsengimana J, Deloukas P, Rahman N, Bishop DT, Easton DF, Stratton MR. 2009. A genome-wide association study of testicular germ cell tumor. *Nat Genet* 41:807-810.
- Renedo DE, Trainer TD. 1994. Intratubular Germ Cell Neoplasia (ITGCN) with p53 and PCNA Expression and Adjacent Mature Teratoma in an Infant Testis An Immunohistochemical and Morphologic Study with a Review of the Literature. *Am J Surg Pathol* 18.
- Riopel MA, Spellerberg A, Griffin CA, Perlman EJ. 1998. Genetic analysis of ovarian germ cell tumors by comparative genomic hybridization. *Cancer Res* 58:3105-3110.
- Rodriguez E, Melamed J, Reuter V, Chaganti RS. 1995. Chromosome 12 abnormalities in malignant ovarian germ cell tumors. *Cancer Genet Cytogenet* 82:62-66.
- Rosenberg C, van Gurp R, Geelen E, Oosterhuis J, Looijenga L. 2000. Overrepresentation of the short arm of chromosome 12 is related to invasive growth of human testicular seminomas and nonseminomas. *Oncogene* 19:5858-5862.
- Ross JA, Schmidt PT, Perentesis JP, Davies SM. 1999. Genomic imprinting of H19 and insulin-like growth factor-2 in pediatric germ cell tumors. *Cancer Genet Cytogenet* 8:1389-1394.
- Schneider DT, Schuster AE, Fritsch MK, Calaminus G, Göbel U, Harms D, Lauer S, Olson T, Perlman EJ. 2002. Genetic analysis of mediastinal nonseminomatous germ cell tumors in children and adolescents. *Genes Chromosomes Cancer* 34:115-125.

- Schneider DT, Schuster AE, Fritsch MK, Hu J, Olson T, Lauer S, Gobel U, Perlman EJ. 2001. Multipoint imprinting analysis indicates a common precursor cell for gonadal and nongonadal pediatric germ cell tumors. *Cancer Res* 61:7268-7276.
- Shenouda S, Alahari S. 2009. MicroRNA function in cancer: oncogene or a tumor suppressor? *Cancer Metastasis Rev* 28:369-378.
- Skakkebaek N. 1978. Carcinomas in *situ* of the testis: Frequency and relationship to invasive germ cell tumours in infertile men. *Histopathology* 2:157-170.
- Skakkebaek NE. 1972. Possible carcinoma-in-situ of the testis. *Lancet* 2:516-517.
- Speleman F, De Potter C, Dal Cin P, Kathelijne. M, Herwig. I, Genevieve. L, Yves. B, Jules. L, Herman. VDB. 1990. i(12p) in a malignant ovarian tumor. *Cancer Genet Cytogenet* 45:49-53.
- Speleman F, Laureys G, Benoit Y, Cuvelier C. 1992. i(12p) in a near-diploid mature ovarian teratoma. *Cancer Genet Cytogenet* 60:216-218.
- Stamp IM, Barlebo H, Rix M, Jacobsen GK. 1993. Intratubular germ cell neoplasia in an infantile testis with immature teratoma. *Histopathology* 22:69-72.
- Stern JW, Bunin N. 2002. Prospective study of carboplatin-based chemotherapy for pediatric germ cell tumors. *Med Pediatr Oncol* 39:163-167.
- Summersgill B, Goker H, Weber-Hall S, Huddart R, Horwich A, Shipley J. 1998. Molecular cytogenetic analysis of adult testicular germ cell tumours and identification of regions of consensus copy number change. *Br J Cancer* 77:305-313.
- Summersgill B, Osin P, Lu Y, Huddart R, Shipley J. 2001. Chromosomal imbalances associated with carcinoma in situ and associated testicular germ cell tumours of adolescents and adults. *Br J Cancer* 85:213-220.
- Surti U, Hoffner L, Chakravarti A, Ferrell RE. 1990. Genetics and biology of human ovarian teratomas. Cytogenetic analysis and mechanism of origin. *Am J Hum Genet* 47:635-643.

- Teilum G. 1965. Classification of endodermal sinus tumour (mesoblastoma vitellinum) and so-called "embryonal carcinoma" of the ovary. *Acta Pathol Microbiol Scand* 64:407-429.
- Turnbull C, Rapley EA, Seal S, Pernet D, Renwick A, Hughes D, Ricketts M, Linger R, Nsengimana J, Deloukas P, Huddart RA, Bishop DT, Easton DF, Stratton MR, Rahman N. 2010. Variants near *DMRT1*, *TERT* and *ATF7IP* are associated with testicular germ cell cancer. *Nat Genetics* 42:604-607.
- van Echten J, Oosterhuis JW, Looijenga LHJ, van de Pol M, Wiersema J, Meerman GJT, Koops HS, Sleijfer DT, Jong BD. 1995. No recurrent structural abnormalities apart from i(12p) in primary germ cell tumors of the adult testis. *Genes Chromosomes Cancer* 14:133-144.
- van Gurp R, Oosterhuis J, Kalscheuer V, Mariman E, Looijenga L. 1994. Biallelic expression of the *H19* and *IGF2* genes in human testicular germ cell tumors. *J Natl Cancer Inst* 86:1070-1075.
- Veltman I, Schepens M, Looijenga L, Strong L, van Kessel A. 2003. Germ cell tumours in neonates and infants: a distinct subgroup? *APMIS* 111:152-160; discussion 160.
- Voorhoeve PM, le Sage C, Schrier M, Gillis AJM, Stoop H, Nagel R, Liu Y-P, van Duijse J, Drost J, Griekspoor A, Zlotorynski E, Yabuta N, De Vita G, Nojima H, Looijenga LHJ, Agami R. 2006. A Genetic Screen Implicates miRNA-372 and miRNA-373 As Oncogenes in Testicular Germ Cell Tumors. *Cell* 124:1169-1181.