

This is a repository copy of *Mental health pathways linking maltreatment to interpersonal revictimization during adolescence for girls in the child welfare system.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/94690/>

Version: Accepted Version

Article:

Auslander, Wendy, Myers Tlapek, Sarah, Threlfall, Jennifer Mary orcid.org/0000-0002-3266-1669 et al. (2 more authors) (2016) Mental health pathways linking maltreatment to interpersonal revictimization during adolescence for girls in the child welfare system. Journal of interpersonal violence. ISSN 1552-6518

<https://doi.org/10.1177/0886260515614561>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Mental Health Pathways Linking Childhood Maltreatment to Interpersonal
Revictimization during Adolescence for Girls in Child Welfare

Wendy Auslander^a, Sarah Myers Tlapek^b, Jennifer Threlfall^c, Tonya Edmond^a, and Jerry Dunn^d

This research was supported by grant No. 1R49CE001510 from the Centers for Disease Control and Prevention awarded to Washington University. The authors acknowledge the collaboration of Children's Division of Missouri of St. Louis City and County, and Jefferson County. Portions of this paper were presented at the American Public Health Association Annual Meeting (2012) in San Francisco, CA, the Society for Social Work and Research (2013) in San Diego, CA, and the 13th European Society for Traumatic Stress Studies (2013) in Bologna, Italy.

Correspondence should be addressed to: Wendy Auslander, Ph.D., Washington University, George Warren Brown School of Social Work, 1 Brookings Drive, Campus Box 1196, St. Louis, MO 63130. Phone: (314) 935-6634; E-mail: wendyaus@wustl.edu

^aWashington University in St. Louis, George Warren Brown School of Social Work;

^bUniversity of Missouri – Columbia, School of Social Work; ^cUniversity of York, Department of Social Policy and Social Work; ^dUniversity of Missouri-St. Louis, Department of Psychological Sciences.

Mental Health Pathways Linking Childhood Maltreatment to Interpersonal Revictimization
during Adolescence for Girls in Child Welfare

Abstract

This study compares the association of histories of childhood emotional, physical, sexual abuse, and physical neglect with revictimization among adolescent girls, and investigates the role of post-traumatic stress and symptoms of depression as mediators. Participants were 234 girls aged 12-19 years who have been involved with the child welfare system in a Midwestern urban area. Data were collected from baseline surveys of a trauma-focused group program to which the participants were referred. The majority of participants were youths of color (75%) who were primarily African American (70%), and the remaining participants were White, non-Hispanic (25%). Data were collected through surveys that assessed histories of child abuse and neglect, symptoms of posttraumatic stress and depression, and experiences of physical, verbal, and relational revictimization in the last three months. The mediating models were tested using steps established by Baron and Kenny (1986), followed by the bootstrapping technique to test the significance of the indirect effects (Preacher and Hayes, 2004). Control variables, such as placement instability, service use, age, and race were investigated in the analyses.

All types of abuse and neglect were significantly associated with higher frequencies of revictimization, and higher levels of posttraumatic stress and depressive symptoms. Parallel mediation analyses demonstrated that both posttraumatic stress and depression fully mediated the relationships between emotional abuse and revictimization, and sexual abuse and revictimization. Physical abuse was fully mediated by posttraumatic stress, but not by depression. Results also indicated that neither posttraumatic stress nor depression were mediators for the relationship

between neglect and revictimization. There were similar pathways to revictimization in adolescents from emotional and sexual abuse through posttraumatic stress and depression.

Evidence is mounting for the deleterious effects of emotional abuse. Based on the mediating analysis, there is evidence that treatment of both posttraumatic stress and depression in emotionally and sexually abused adolescents involved in child welfare is warranted to prevent future revictimization in those girls referred to services.

Keywords: child maltreatment, revictimization, post-traumatic stress, depression

Childhood abuse and neglect are common in the U.S. The most recent prevalence rates indicate that one in eight children under 18 years of age has experienced at least one form of child maltreatment (Wildeman et al., 2014). Childhood maltreatment has been associated with a number of negative outcomes. One serious consequence of child maltreatment is the increased risk for multiple types of revictimization during adolescence and adulthood (Desai, Arias, Thompson, & Basile, 2002; Irwin, 1999; Schaaf & McCanne, 1998; Widom, Czaja, & Dutton, 2008). Additionally, there are some serious mental health consequences to childhood abuse and neglect such as posttraumatic stress and depression. Adolescents involved in the child welfare system have high rates of serious mental health disorders compared with the general adolescent population, and females have higher rates of depression and other internalizing behavior disorders than males within the child welfare system (Auslander et al., 2002; McMillen, et al., 2005). Although there is research about the negative consequences of child maltreatment, there is limited knowledge of the mental health pathways linking childhood maltreatment to physical, verbal, or relational revictimization among adolescents in the child welfare system.

There is a growing literature of the association between childhood abuse and revictimization as reported by *adult women retrospectively* (West, Williams, & Siegel, 2000; for a review, see Classen et al., 2002). In some cases, specific types of maltreatment have been found to be more likely to lead to later revictimization than others; however, the research is inconclusive. There is also some evidence for a link between childhood maltreatment and revictimization in adolescence. Boney-McCoy and Finkelhor (1995) found that victims of childhood sexual abuse were almost twelve times as likely to be revictimized when compared to non-abused counterparts. Moreover, children who were maltreated were 20% more likely to experience peer or sibling victimization and 30% more likely to be physically assaulted

(Finkelhor, Ormrod, & Turner, 2007). Physical abuse, sexual abuse, and physical neglect have all been associated with peer victimization (Duncan, 1999). One of the only studies in a child welfare population found that emotional abuse was predictive of dating violence among female adolescents engaged with child protective services (Wekerle et al., 2009).

In addition to the evidence for the association between childhood abuse and revictimization during adolescence, research has demonstrated the linkages between posttraumatic stress and revictimization. A national longitudinal sample of adolescents found that PTSD symptoms at baseline significantly predicted interpersonal revictimization, controlling for the effects of prior victimization (McCart et al., 2012). Likewise, in a study of adolescent dating relationships over one year, trauma-related symptoms were a significant predictor of dating violence in a community sample (Wolfe, Wekerle, Scott, Straatman, & Grasley, 2004).

Information-processing models of memory provide one framework for how symptoms of mental health problems, particularly PTSD and depression resulting from child maltreatment, may contribute to revictimization (Beckham, Moore, & Reynolds, 2000). Research on cognitions associated with memory has recently focused on ways individuals with intrusive PTSD symptoms have difficulty ignoring trauma-related stimuli (Pineles, Shipherd, Mostoufi, Abramovitz, & Yovel, 2009) which hampers their ability to perceive new threats and may render them vulnerable to revictimization (Cougle, Resick, & Kilpatrick, 2009; Messman-Moore & Long, 2003). A diminished cognitive capacity to detect threat is also theorized to result from depression (Cougle et al., 2009), such that symptoms of depression hinder the ability of an adolescent with a history of child maltreatment to avoid revictimization (Wekerle et al., 2001). Maltreated children may also fail to develop secure attachment patterns with caregivers, interrupting affect regulation, or the ability to handle negative emotions in the face of danger or

distress (Schore & Schore, 2008; Mikulincer & Shaver, 2012). These children may be more vulnerable as adolescents and adults to anxiety, depression, and PTSD (Mikulincer & Shaver, 2012). Children with insecure attachment patterns may harbor feelings of unworthiness, be unable to correctly interpret traumatic experiences and, as a result, may be unable to prevent harmful interpersonal interactions such as revictimization (Tanaka, Wekerle, Schmuck, Paglia-Boak, & MAP Research Team, 2011).

Recent studies provide evidence for the mediating role of PTSD and depression in revictimization with adult samples, the majority of which have focused on the pathways between childhood sexual abuse and sexual revictimization among females (Fortier et al., 2009; Messman-Moore, Brown, & Koelsch, 2005; Risser, Hetzel-Riggin, Thomsen, & McCanne, 2006). Research on pathways between types of maltreatment and revictimization in adolescents is more limited, particularly with child welfare samples. For example, a study with a sample of students from 10 high schools demonstrated that trauma-related symptoms mediated the relationship between child maltreatment and of adolescent dating violence (Wolfe, et al., 2004). Another study that included a sample of juvenile offenders found that depression provided a pathway between childhood emotional abuse and peer victimization (Day et al., 2013). The only study that focused on adolescents involved in child welfare found that posttraumatic stress symptoms mediated the relationship between child maltreatment and dating violence victimization in girls (Wekerle et al., 2001).

The current study extends prior research by comparing multiple types of child maltreatment among girls in child welfare and their association with physical, verbal, and relational revictimization. Because pathways from childhood abuse and neglect to revictimization have been shown to be gender-based (Wekerle et al., 2009), and females are

more vulnerable to mental health problems and revictimization, the current study examined girls. Moreover, we operationalized *revictimization* more broadly than dating violence in order to capture the experiences of adolescent girls who may be experiencing interpersonal revictimization from a variety of perpetrators and in multiple contexts. Therefore, to further our knowledge of the mental health pathways by which histories of child maltreatment lead to revictimization (physical, verbal, and relational) among adolescent girls in child welfare, the following questions were addressed: 1) What is the association between different types of childhood abuse and neglect (emotional, physical, sexual abuse, and physical neglect) and revictimization among adolescent girls involved in the child welfare system? And 2) do posttraumatic stress and depression mediate this relationship among these adolescents? Findings will contribute to our understanding of the mechanisms by which childhood abuse and neglect increases the risk of revictimization during adolescence, and highlight potential modifiable factors to target for preventing future revictimization in female adolescents and young adults.

Methods

Subjects

Participants were 234 girls ages 12-19 ($M = 14.9$, $SD = 1.6$) who were involved with the child welfare system. Participants were recruited through referrals from the state's child protective agency (Children's Division), other agencies that provide services to adolescents in the child welfare system, and from caregivers. Inclusion criteria for the study included: 1) adolescent girls who were formally investigated or substantiated for child maltreatment by Children's Division and 2) girls 12-19 years old. Girls were excluded if they were unable to read or write, or if they had behaviors that would prohibit participation in either a structured interview

or group treatment focused on abuse and neglect. It was not necessary for girls to meet diagnostic thresholds for PTSD or depression to be included in the study.

Seventy-five percent of participants were youths of color, of whom the majority (70%) were African-American, and the other 5% self-identified as Native American, Hispanic, Asian/Asian American, or “other.” The remaining participants (25%) non-Hispanic, White. The living situations of the participants was as follows: 60% lived with biological parent(s), relatives, or an adoptive family, 27% lived with a foster family, and 13% living in a group or other congregate care facility. Of the participants, 72% were currently receiving 1 or more types of services such as outpatient, health, social, behavioral, mental health, and/or school-based services ($M = 1.35$, $SD = 1.14$).

Informed Consent and Interview Procedures

Baseline data collected for a trauma-focused cognitive behavioral intervention study were used for the current analysis. The study protocol was first approved by the Human Subjects Institutional Review Board of the two collaborating universities. Second, a Certificate of Confidentiality was issued by Centers for Disease Control and Prevention (CDC) (the funding agency) to protect the privacy of research subjects by withholding their identities from all persons not connected with the research. Additionally, the Research Committee of the state office of child protective services approved the research protocol.

The consent procedures included several steps. After a referral was made to the study team, and the adolescent expressed interest in participating, written consent was obtained from the legal custodian. Additionally, to the fullest extent possible, written consent was secured from all members of the youth’s Family Support Team (e.g., Guardian ad litem, Deputy Juvenile Officer, child’s current therapist). All adolescents under the age of 18 provided written assent to

participate in the study. Participants were given a \$20 gift card to compensate them for their time.

Data were collected by surveys verbally administered by interviewers who were masters-level or doctoral-level social work students who participated in eight hours of interview training. The training included background knowledge of the population, basic interviewing skills, confidentiality and ensuring privacy during the interview, procedures for reporting abuse, and procedures for participants who endorsed items related to suicide. The surveys were administered in participant homes or a community-based mental health agency and took approximately one hour to complete.

Variables and Measures

Child Maltreatment. The Child Trauma Questionnaire (CTQ; Bernstein & Fink, 1998) was used to obtain participant reports of the types and severity of child maltreatment experienced. Four subscales from the 28 item scale were used in the present study: emotional abuse, physical abuse, sexual abuse, and physical neglect. Items are rated from never true (1) to very often true (5). Previous research has demonstrated the validity and reliability of the CTQ (Bernstein & Fink, 1998). Additionally, guidelines for classification of CTQ scores are provided according to the following categories: None or minimal, low-moderate, moderate-severe, and severe-extreme (Bernstein & Fink, 1998). Internal consistency alpha coefficients for the current sample were as follows: emotional abuse ($\alpha = 0.86$), physical abuse ($\alpha = 0.87$), sexual abuse ($\alpha = 0.92$), and physical neglect ($\alpha = 0.81$).

PTSD Symptoms. Posttraumatic stress symptoms were assessed using the Child PTSD Symptom Scale (CPSS; Foa, Johnson, Feeny, & Treadwell, 2001). Seventeen items (e.g., Having bad dreams or nightmares; Trying not to think about, talk about, or have feelings about the

trauma) assessed domains of re-experiencing, avoidance, and arousal. Participants rated the frequency of posttraumatic stress symptoms over the past month using a 4 point scale from “not at all” (0) to “five or more times a week” (4). The CPSS has previously demonstrated convergent validity, correlating highly with a similar PTSD scale, as well as good internal consistency and test-retest reliability (Foa et al., 2001). For the present study, internal consistency reliability was $\alpha = 0.90$.

Depression. The Child Depression Inventory (CDI; Kovacs, 2003) was used to measure depressive symptoms over the previous two weeks. The 27 items were rated from 0 to 2 and summed, with higher scores indicating more severe symptoms. Concurrent, discriminative, and criterion validity have been demonstrated (Kovacs, 2003; Saylor, Finch, Spirito, & Bennett, 1984). Good internal consistency and test-retest reliability have been demonstrated for a child welfare population (Kolko et al., 2010). The internal consistency alpha coefficient for the current sample was $\alpha = 0.89$.

Interpersonal Revictimization. The Problem Behavior Frequency Scale (PBFS; Farrell, Kung, White, & Valois, 2000) consisting of 18 items was originally developed to assess peer aggression toward others among school-aged children and adolescents. For the current study of girls with prior histories of maltreatment, the items were modified to assess the girls’ experiences of being revictimized (i.e., aggressive behaviors targeting them in the past three months). Example items from the three subscales included: physical revictimization (e.g., *Another person hit or slapped you*); verbal revictimization (e.g., *Someone teased you to make you angry*); and relational revictimization (e.g., *Another person tried to keep others from liking you by saying mean things about you*). The language of each item did not specify who was the perpetrator (e.g. “another person” or “someone”) in order to capture the wide range of possible individuals who

may revictimize the girls. Items were rated to indicate the frequency with which they had happened over the past 3 months on a 6 point scale coded as follows: 1= '0 times,' 2= '1-2 times,' 3= '3-5 times,' 4= '6-9 times,' 5= '10-19 times,' and 6= '20+ times.' Items were summed to yield a *total revictimization score* with a possible range of 18 to 108. Previous studies have shown adequate to good internal consistency (Farrell et al., 2000; Miller-Johnson, Sullivan, & Simon, 2004), and the alpha coefficient for the current sample was $\alpha = 0.93$.

Potential Control Variables. Several control variables were investigated because they have been shown in previous research to predict poor mental health and behavioral outcomes among child welfare populations, such as placement instability, use of services, age, and race (0=youths of color; 1=white). The number of services used in the participant's lifetime was measured using the Service Assessment for Children and Adolescents (SACA; Stiffman et al., 2000). This structured interview asks about the number and types of inpatient, outpatient, and school-based social, behavioral, and mental health services the participant has ever received. Placement instability (ever) was measured using a count of the number of different types of placements that the adolescent had lived during their lifetime.

Data Analysis

The mediating roles of posttraumatic stress and depression were tested using the steps established by Baron and Kenny (1986), followed by the bootstrapping technique to test the significance of the indirect effects described by Preacher and Hayes (2004). First, a preliminary analysis was conducted to determine that there were significant correlations among the hypothesized predictors (types of abuse and neglect), mediators (posttraumatic stress and depression), and outcome variable (revictimization). Second, a mediation analysis was conducted using the Hayes (2013) PROCESS SAS macro. Because there is no theoretical causal

assumption between the two mediators, and they work relatively independently from each other, parallel mediation models were tested. In addition to calculating the standard a , b , c , and c' path coefficients, the Hayes macro uses a bootstrap re-sampling methodology to enable a significance test of the indirect effect (the impact of the predictor on the outcome variable through the mediator).

Results

Descriptive Analyses

Descriptive statistics (means and standard deviations) for the abuse and neglect (independent) variables, mediators (posttraumatic stress, depression), outcome variable (revictimization), and control variables are shown in Table 1. According to the Bernstein and Fink classification (1998), the mean scores of the childhood abuse and neglect subscales indicated that the girls experienced emotional abuse and physical neglect in the low to moderate range, physical abuse in the moderate to severe range, and sexual abuse in the severe range. Of the 234 adolescent girls in the sample, 83% had experienced childhood emotional abuse, 81% had experienced physical abuse, 64% sexual abuse, and 69% physical neglect (based on endorsing at least one of the items on the subscale).

Results indicated that 51% of the girls experienced PTSD symptoms in the clinical range (≥ 15), with a mean score of 16.90 (SD 10.95) for the entire sample. The participants' mean depression score was 12.05 (SD 8.57). Previous studies have reported CDI cut-off scores for clinically referred samples of ≥ 13 where higher scores are expected (Kovacs, 2003). In the current study, 35% of the girls had depression scores that were in the clinical range.

The frequencies of each type of revictimization that were experienced in the last 3 months are shown in Table 2. Notably, among the physical revictimization items, approximately

one in six of the girls ($n=37$) were threatened with a weapon, over one-half were shoved or pushed, and over one third were hit or slapped. Comparisons of different types of revictimization indicated that 84% endorsed at least one item in the verbal revictimization subscale; 74% endorsed at least one item in the relational revictimization subscale, and 68% endorsed at least one item in the physical revictimization subscale.

Bivariate and Mediation Analyses

Bivariate analyses were conducted between independent, mediating, and dependent variables to determine if they met the criteria to test a mediating hypothesis. Results shown in Table 3 indicated that significant relationships were found between all forms of abuse and neglect (emotional, physical, sexual abuse, and physical neglect) with revictimization, PTSD, and depression. Compared with other forms of abuse, emotional abuse was most strongly related to mental health symptoms and revictimization, and physical neglect showed the weakest relationships with these variables. Of the mediators, both PTSD and depression were strongly correlated with revictimization. Placement instability and service use were used as control variables because they were significantly associated with the dependent variable and the majority of independent variables, and conceptually may explain the relationship between histories of abuse and neglect and experiences of revictimization as adolescents. Because age and race were not associated with revictimization at the bivariate level, they were not included as control variables in the mediation models.

As shown in Figure 1, mediation analyses indicated that participants with greater emotional abuse were associated with more revictimization, and this relationship was fully mediated by both PTSD and depression. Results of the bootstrapping analysis described by Preacher and Hayes (2004) indicated significant indirect effects of emotional abuse on

revictimization via the hypothesized mediators of PTSD ($\beta = .3625$; 95% CI [0.1139, 0.6476]) and depression ($\beta = .2510$; 95% CI [0.0721, 0.5380]). Likewise, both PTSD and depression were significant mediators of the impact of sexual abuse on revictimization (as shown in Figure 2). Adolescent girls with histories of more severe sexual abuse experienced more revictimization, and this pathway was fully mediated by both PTSD and depression. In both pathways, the a and b paths were significant, and the c' path was not. Results of the bootstrapping analyses demonstrated a significant indirect effect of sexual abuse on revictimization for both PTSD and depression.

The parallel multiple mediation analyses testing the relationship between physical abuse and revictimization failed to support the pathway with depression as a mediator. As shown in Figure 3, results showed that the a_1 path from physical abuse to depression was not significant ($\beta = .1642$, $p = .12$) nor was the indirect effect of physical abuse on revictimization via depression ($\beta = .0805$, 95% CI [-0.0055, 0.2668]). However, results supported the mediating pathway of PTSD between physical abuse and revictimization.

Last, the parallel mediation model for PTSD and depression were not supported for the impact of physical neglect on revictimization. As shown in Figure 4, both a paths from physical neglect to PTSD and depression were not significant, nor were the indirect effects of physical neglect on revictimization (i.e., the confidence intervals contained zero).

Discussion

The results of the current study compared four types of child maltreatment and two different mental health pathways to revictimization among adolescent girls involved in child welfare. Perhaps the most compelling finding was that there are similar pathways to revictimization in adolescents from both emotional and sexual abuse through posttraumatic stress

and depression. The importance of this lies in the common assumption that emotional abuse is not as serious or damaging as sexual or physical abuse. Yet for the girls in this study, childhood histories of emotional abuse were significantly associated with posttraumatic stress and depression and increased their vulnerability to physical, verbal, and relational revictimization during adolescence.

In a recent study, Spinazzola and colleagues (2014) found that emotionally abused children experienced similar and in some cases higher rates of PTSD, depression, and other mental health problems than physically or sexual abused children. They found that when children experienced emotional abuse in conjunction with either physical or sexual abuse, they had significantly more severe negative psychological outcomes than those who had been physically and sexually abused without emotional abuse. In fact, both physical and sexual abuse had to occur simultaneously to induce the same negative effects as emotional abuse alone.

Clearly evidence is growing that demonstrates the deleterious effects of emotional abuse, including an increased vulnerability to revictimization as a consequence of PTSD and depression. However, there is still a tendency in the child welfare system and other systems of care to focus on what are perceived to be more serious forms of child maltreatment—physical and sexual abuse. It is often more difficult to detect, substantiate, and take action in cases involving emotional abuse, especially if that is the only type of abuse occurring. This is further complicated by the definitions of emotional abuse that determine grounds for intervention by state child protective agencies. Most of the states define emotional abuse as “injury to the psychological capacity or emotional stability of the child...as evidenced by anxiety, depression withdrawal, or aggressive behavior” (Child Welfare Information Gateway, 2014, p. 3). Consequently, there must already be evidence of the harmful effects of emotional abuse to the

child's behavioral and mental health before a report can be made to child protective services. Practitioners, therefore, outside of the child welfare system may need to be poised to detect this insidious form of abuse and to assess and treat the early symptoms PTSD and depression that these children and adolescents are likely to be experiencing. Assessment tools are available such as the Childhood Trauma Questionnaire (Bernstein & Fink, 1998) used in the current study, and a subscale of the Child Abuse and Trauma Scale (Kent & Waller, 1998).

For both emotionally abused and sexually abused adolescents, trauma-focused cognitive behavioral therapy in either the individual or group model has been shown to be effective in reducing PTSD and depression (Foa, Keane, Friedman, & Cohen, 2008; National Child Traumatic Stress Network, 2013). Moreover, among adult women, cognitive-behavioral therapy reduced PTSD and depressive symptoms and the likelihood of intimate partner violence (Iverson et al., 2011). Based on the mediating analysis of the current study, treatment of both PTSD and depression in emotionally and sexually abused girls involved in child welfare is warranted to prevent revictimization during adolescence.

It has been suggested that different types of maltreatment may lead to the same negative outcomes through different pathways or processes (Shaffer, Yates, & Egeland, 2009). Such appears to be the case with physical abuse. Posttraumatic stress was a significant mediator or pathway to revictimization, but depression was not. It has been theorized that depressed children can become targets for peer victimization because they are perceived as being less inclined to defend themselves or fight back (Fekkes, Pijpers, Fredriks, Vogels, & Verloove-Vanhorick, 2006). However, in the current study depression was not a significant pathway from physical abuse to revictimization, and in fact, the bivariate correlations indicate a weak significant relationship between physical abuse and depression. Another perspective is that children who

are physically abused are more easily angered, have a hyper vigilant interpersonal style, and are more likely to be reactively aggressive with peers (Dodge, Bates, & Pettit, 1990; Messman-Moore, et al., 2005). Further investigation comparing the role of depression as a pathway to revictimization and aggression towards others would be useful in designing prevention programs in this population.

The current study adds to the growing evidence that PTSD symptoms are associated with a heightened risk of revictimization and provides support for the mediating role of posttraumatic stress in the relationship between child maltreatment and revictimization (Messman-Moore et al., 2005; Messman-Moore, Ward, & Brown, 2009). Attentional interference to trauma stimuli in adolescent girls with intrusive PTSD symptoms may interfere with their inability to detect and avoid danger and may explain the heightened vulnerability to revictimization (Pineles, et al., 2009; Cogle et al., 2009). Additionally, in this population of girls who experienced childhood maltreatment, affect dysregulation is likely, with stressful situations leading to negative affective states and a reduced inability to avoid dangerous situations (Tanaka et al., 2011). To prevent revictimization, it is essential that adolescent girls involved in child welfare who have symptoms of posttraumatic stress as a result of physical, sexual, or emotional abuse have access to the most effective evidence based treatment available such as Trauma-Focused Cognitive-Behavior Therapy (California Evidence Based Clearinghouse, 2014).

Neither posttraumatic stress nor depression was a significant mediator of the relationship between physical neglect and revictimization. Although physical neglect is a form of reportable child maltreatment which can lead to removal from the home, the current study indicates that the pathways and inter-relationships between physical neglect, mental health outcomes, and revictimization are different from those of childhood emotional, physical, and sexual abuse. In

the current study, childhood neglect was measured by items such as, “*When I was growing up, I didn’t have enough to eat*” and “*When I was growing up, my parents were too drunk or high to take care of the family.*” Potentially, adolescents who experience physical neglect may hold different attributions than those who experience other forms of abuse, i.e., youths may believe that their experiences of neglect are neither their fault nor an intentional act by their caregivers (as compared with emotional, physical, or sexual abuse). This may explain the weaker relationships found between childhood neglect, posttraumatic stress, and depression in this population. The current findings indicate that childhood neglect is significantly associated with higher levels of revictimization, yet more research is needed to better understand the pathways involved. One possible modifiable pathway may be negative parenting behavior, which is frequently associated with parental substance and alcohol use.

There are several limitations to the present study. First, cross-sectional data were utilized to determine the relationships between childhood abuse and neglect, symptoms of PTSD and depression, and revictimization, and a longitudinal design would be more rigorous. Second, our measure of physical, verbal, and relational revictimization did not specify who the perpetrator of the aggression was; it could have been a peer, family member, or some other person, since the items use the word “someone” instead of a more specific term. Although many of these items are similar to those used to assess bullying or peer victimization in schools, we cannot assume that this is the case. Third, it is important to note that the adolescents who participated in the current study were not a representative sample of girls served by the child welfare system, but a convenience sample of those who were referred for a trauma-focused group treatment. However, because we did not use a clinical cut-off score for either PTSD or depression as inclusion criteria, the current sample may be more representative of a general child welfare sample of

female adolescents than a clinical sample. Furthermore, the majority of the sample was African American, and many of the girls may have experienced racially based revictimization that was not captured by our measures. Although white girls in the current study reported modestly higher rates of maltreatment, no racial differences were found in rates of revictimization, consistent with most of the existing literature (Jackson, O'Brien, & Pecora, 2011; Widom, et al., 2008). Future studies of minority populations should consider forms of racially based aggression (e.g., discrimination, stereotyping, microaggressions) as a type of revictimization that may be experienced by maltreated children. Last, because the purpose of the current study was to compare the mediating pathways of four types of abuse and neglect leading to revictimization, we did not examine the impact of co-occurring types of abuse and neglect. Some may argue that information is lost when maltreatment is classified into single types for analysis because it does not mirror reality. Moreover, we were not able to statistically control for the other types of abuse in each of the mediating models due to the high inter-correlations (i.e., multicollinearity) among the types of abuse. Examining the mediating pathways of polyvictimization and risk for revictimization (Hamby & Grych, 2013) in adolescent girls involved in child welfare would be beneficial in the future, particularly to create tailored treatment strategies for this population.

Despite these limitations, the current study has several strengths. Most importantly, many prior studies have included samples of adults who self-reported experiences of childhood abuse and/or neglect. This may be less accurate than the current study that included a sample that was formally investigated by workers from child protective services. Also, identifying mediating pathways during adolescence provides information that is useful for preventing dating and intimate partner violence in young adulthood. A second strength of the current study is the ability to compare pathways from four different types of maltreatment to revictimization. In particular,

the current study includes emotional abuse, which has often been overlooked despite some evidence that the effects of emotional abuse are considered disabling and enduring (Shaffer et al., 2009). The measure of abuse and neglect used a continuous scale instead of a single dichotomous item that allowed us to capture the severity of the abuse and neglect. Last, the current study examined parallel mediator models and was able to control for other significant covariates.

In addition to posttraumatic stress and depression, other researchers have proposed a number of other mediators to explain the relationship between maltreatment and revictimization. For example, emotional dysregulation, anger, and social skills deficits have been identified as potential pathways to negative outcomes (Hong, Espelage, Grogan-Kaylor, & Allen-Meares, 2012). Future research would benefit from including more comprehensive and complex mediation and path models to examine the pathways from childhood abuse and neglect to revictimization in adolescent females and to further investigate gender differences.

References

- Auslander, W. F., McMillen, J. C., Elze, D., Thompson, R., Jonson-Reid, M., & Stiffman, A. (2002). Mental health problems and sexual abuse among adolescents in foster care: Relationship to HIV risk behaviors and intentions. *AIDS and Behavior*, 6(4), 351-359.
- Baron, R. M., & Kenny, D. A. (1986). The moderator–mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*, 51(6), 1173-82.
- Beckham, J. C., Moore, S. D., & Reynolds, V. (2000). Interpersonal hostility and violence in Vietnam combat veterans with chronic posttraumatic stress disorder: A review of theoretical models and empirical evidence. *Aggression and Violent Behavior*, 5(5), 451 – 466.
- Bernstein, D. P., & Fink, L. (1998). *Childhood trauma questionnaire: A retrospective self-report: Manual*. San Antonio, TX: The Psychological Corporation.
- Boney-McCoy, S., & Finkelhor, D. (1995). Psychosocial sequelae of violent victimization in a national youth sample. *Journal of Consulting and Clinical Psychology*, 63(5), 726-36.
- California Evidence-Based Clearinghouse for Child Welfare. (2014). Trauma-Focused Cognitive-Behavioral Therapy (TF-CBT): Detailed report. Retrieved from <http://www.cebc4cw.org/program/trauma-focused-cognitive-behavioral-therapy/detailed>
- [Child Welfare Information Gateway. \(2014\). Definitions of child abuse and neglect. Washington, DC: U.S. Department of Health and Human Services, Children’s Bureau.](#)
- Classen, C., Nevo, R., Koopman, C., Nevill-Manning, K., Gore-Felton, C., Rose, D. S., & Spiegel, D. (2002). Recent stressful life events, sexual revictimization, and their relationship with traumatic stress symptoms among women sexually abused in childhood.

Journal of Interpersonal Violence, 17(12), 1274-1290.

Cougle, J.R., Resnick, H., & Kilpatrick, D.G. (2009). A prospective examination of PTSD symptoms as risk factors for subsequent exposure to potentially traumatic events among women. *Journal of Abnormal Psychology, 118*(2), 405-411.

Day, D. M., Hart, T. A., Wanklyn, S. G., McCay, E., Macpherson, A., & Burnier, N. (2013). Potential mediators between child abuse and both violence and victimization in juvenile offenders. *Psychological Services, 10*(1), 1-11.

Desai, S., Arias, I., Thompson, M. P., & Basile, K. C. (2002). Childhood victimization and subsequent adult revictimization assessed in a nationally representative sample of women and men. *Violence and Victims, 17*(6), 639-653.

Dodge, K. A., Bates, J. E., & Pettit G. S. (1990). Mechanisms in the cycle of violence. *Science, 250*(4988), 1678-1683.

Duncan, R. D. (1999). Maltreatment by parents and peers: The relationship between child abuse, bully victimization, and psychological distress. *Child Maltreatment, 4*(1), 45-55.

Farrell, A. D., Kung, E. M., White, K. S., & Valois, R. F. (2000). The structure of self-reported aggression, drug use, and delinquent behaviors during early adolescence. *Journal of Clinical Child Psychology, 29*(2), 282-292.

Fekkes, M., Pijpers, F. I., Fredriks, A. M., Vogels, T., & Verloove-Vanhorick, S. P. (2006). Do bullied children get ill, or do ill children get bullied? A prospective cohort study on the relationship between bullying and health-related symptoms. *Pediatrics, 117*, 1568–1574.

Finkelhor, D., Ormrod, R. K., & Turner, H. A. (2007). Poly-victimization: A neglected component in child victimization. *Child Abuse & Neglect, 31*(1), 7-26.

Foa, E. B., Johnson, K. M., Feeny, N. C., & Treadwell, K. R. (2001). The child PTSD symptom

- scale: A preliminary examination of its psychometric properties. *Journal of Clinical Child & Adolescent Psychology*, 30(3), 376-384.
- Foa, E. B., Keane, T. M., Friedman, M. J., & Cohen, J. A. (Eds.). (2008). *Effective treatments for PTSD: practice guidelines from the International Society for Traumatic Stress Studies* (2nd ed.). New York, NY: Guilford Press.
- Fortier, M. A., DiLillo, D., Messman-Moore, T. L., Peugh, J., DeNardi, K. A., & Gaffey, K. J. (2009). Severity of child sexual abuse and revictimization: The mediating role of coping and trauma symptoms. *Psychology of Women Quarterly*, 33(3), 308-320.
- Hamby, S., & Grych, J. (2013). *The web of violence: Exploring connections among forms of interpersonal violence and abuse*. Dordrecht, The Netherlands: Springer.
- Hayes, A. F. (2013). *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach*. Guilford Press.
- Hong, J. S., Espelage, D. L., Grogan-Kaylor, A., & Allen-Meares, P. (2012). Identifying potential mediators and moderators of the association between child maltreatment and bullying perpetration and victimization in school. *Educational Psychology Review*, 24(2), 167-186.
- Irwin, H. J. (1999). Violent and nonviolent revictimization of women abused in childhood. *Journal of Interpersonal Violence*, 14(10), 1095-1110.
- Iverson, K. M., Gradus, J. L., Resick, P. A., Suvak, M. K., Smith, K. F., & Monson, C. M. (2011). Cognitive-behavioral therapy for PTSD and depression symptoms reduces risk for future intimate partner violence among interpersonal trauma survivors. *Journal of Consulting and Clinical Psychology*, 79(2), 193-202.
- Jackson, L. J., O'Brien, K., & Pecora, P. J. (2011). Posttraumatic stress disorder among foster

- care alumni: The role of race, gender, and foster care context. *Child Welfare*, 90(5), 71-93.
- Kent, A. & Waller, G. (1998). The impact of childhood emotional abuse: An extension of the child abuse and trauma scale. *Child Abuse & Neglect*, 22(5), 393-399.
- Kolko, D. J., Hurlburt, M. S., Zhang, J., Barth, R. P., Leslie, L. K., & Burns, B. J. (2010). Posttraumatic stress symptoms in children and adolescents referred for child welfare investigation: A national sample of in-home and out-of-home care. *Child Maltreatment*, 15(1), 48-63.
- Kovacs, M. (2003). The children's depression inventory (CDI): Technical manual. North Tonawanda, New York: Multi-Health Systems, Inc.
- McCart, M. R., Zajac, K., Kofler, M. J., Smith, D. W., Saunders, B. E., & Kilpatrick, D. G. (2012). Longitudinal examination of PTSD symptoms and problematic alcohol use as risk factors for adolescent victimization. *Journal of Clinical Child & Adolescent Psychology*, 41(6), 822-836.
- McMillen, J. C., Zima, B. T., Scott, L. D., Auslander, W. F., Munson, M. R., Ollie, M. T., & Spitznagel, E. L. (2005). Prevalence of psychiatric disorders among older youths in the foster care system. *Journal of the American Academy of Child and Adolescent Psychiatry*, 44(1), 88-95.
- Messman-Moore, T. L., Brown, A. L., & Koelsch, L. E. (2005). Posttraumatic symptoms and self-dysfunction as consequences and predictors of sexual revictimization. *Journal of Traumatic Stress*, 18(3), 253-261.
- Messman-Moore, T. L., & Long, P. J. (2003). The role of childhood sexual abuse sequelae in the sexual revictimization of women: An empirical review and theoretical reformulation. *Clinical Psychology Review*, 23(4), 537-571.

- Messman-Moore, T. L., Ward, R. M., & Brown, A. L. (2009). Substance use and PTSD symptoms impact the likelihood of rape and revictimization in college women. *Journal of Interpersonal Violence, 24*(3), 499-521.
- Mikulincer, M., & Shaver, P. R. (2012). An attachment perspective on psychopathology. *World Psychiatry, 11*(1), 11-15.
- Miller-Johnson, S., Sullivan, T. N., & Simon, T. R. (2004). Evaluating the impact of interventions in the multisite violence prevention study: Samples, procedures, and measures. *American Journal of Preventive Medicine, 26*(1), 48-61.
- National Child Traumatic Stress Network. (2013). *National Child Traumatic Stress Network empirically supported treatments and promising practices*. Retrieved from <http://www.nctsnet.org/resources/topics/treatments-that-work/promising-practices>.
- Pineles, S. L., Shipherd, J. C., Mostoufi, S. M., Abramovitz, S. M., & Yovel, I. (2009). Attentional biases in PTSD: More evidence for interference. *Behaviour Research and Therapy, 47*(12), 1050 – 1057.
- Preacher, K. J., & Hayes, A. F. (2004). SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behavior Research Methods, Instruments, & Computers, 36*(4), 717-731.
- Risser, H. J., Hetzel-Riggin, M. D., Thomsen, C. J., & McCanne, T. R. (2006). PTSD as a mediator of sexual revictimization: The role of reexperiencing, avoidance, and arousal symptoms. *Journal of Traumatic Stress, 19*(5), 687-698.
- Saylor, C. F., Finch, A., Spirito, A., & Bennett, B. (1984). The children's depression inventory: A systematic evaluation of psychometric properties. *Journal of Consulting and Clinical Psychology, 52*(6), 955-967.

- Schaaf, K. K., & McCanne, T. R. (1998). Relationship of childhood sexual, physical, and combined sexual and physical abuse to adult victimization and posttraumatic stress disorder. *Child Abuse & Neglect*, 22(11), 1119-1133.
- Schore, J. R., & Schore, A. N. (2008). Modern attachment theory: The central role of affect regulation in development and treatment. *Clinical Social Work Journal*, 36(1), 9-20.
- Shaffer, A., Yates, T. M., & Egeland, B. R. (2009). The relation of emotional maltreatment to early adolescent competence: Developmental processes in a prospective study. *Child Abuse & Neglect*, 33, 36-44.
- Spinazzola, J., Hodgdon, H., Liang, L. J., Ford, J. D., Layne, C. M., Pynoos, R., ... & Kisiel, C. (2014). Unseen wounds: The contribution of psychological maltreatment to child and adolescent mental health and risk outcomes. *Psychological Trauma: Theory, Research, Practice, and Policy*, 6(S1), S18-S28.
- Stiffman, A. R., Horwitz, S. M., Hoagwood, K., Compton III, W., Cottler, L., Bean, D. L., . . . Weisz, J. R. (2000). The service assessment for children and adolescents (SACA): Adult and child reports. *Journal of the American Academy of Child & Adolescent Psychiatry*, 39(8), 1032-1039.
- Tanaka, M., Wekerle, C., Schmuck, M. L., Paglia-Boak, A., & MAP Research Team. (2011). The linkages among childhood maltreatment, adolescent mental health, and self-compassion in child welfare adolescents. *Child Abuse & Neglect*, 35(10), 887-898.
- Wekerle, C., Leung, E., Wall, A. M., MacMillan, H., Boyle, M., Trocme, N., & Waechter, R. (2009). The contribution of childhood emotional abuse to teen dating violence among child protective services-involved youth. *Child Abuse & Neglect*, 33(1), 45-58.
- Wekerle, C., Wolfe, D. A., Hawkins, D., Pittman, A. L., Glickman, A., & Loyal, B. E. (2001).

Childhood maltreatment, posttraumatic stress symptomatology, and adolescent dating violence: Considering the value of adolescent perceptions of abuse and a trauma mediational model. *Development and Psychopathology*, 13(04), 847-871.

West, C. M., Williams, L. M., & Siegel, J. A. (2000). Adult sexual revictimization among black women sexually abused in childhood: A prospective examination of serious consequences of abuse. *Child Maltreatment*, 5(1), 49-57.

Widom, C. S., Czaja, S. J., & Dutton, M. A. (2008). Childhood victimization and lifetime revictimization. *Child Abuse & Neglect*, 32(8), 785-796.

Wildeman, C., Emanuel, N., Leventhal, J.M., Putnam-Hornstein, E., Waldfogel, J., & Lee, H. (2014). The prevalence of confirmed maltreatment among US children, 2004 to 2011. *Journal of the American Medical Association Pediatrics*, 168(8), 706-713.

Wolfe, D., Wekerle, C., Scott, K., Straatman, A., Grasley, C. (2004). Predicting abuse in adolescent dating relationships over 1 year: The role of child maltreatment and trauma. *Journal of Abnormal Psychology*, 113(3), 406-415.

Table 1

Means (M) and standard deviations (SD) of variables (N = 234).

	<i>M</i>	<i>SD</i>
Emotional abuse	12.09	5.94
Physical abuse	11.05	5.78
Sexual abuse	12.65	7.38
Physical neglect	8.87	4.38
PTSD	16.90	10.95
Depression	12.05	8.57
Revictimization	32.67	15.25
Physical	10.46	4.84
Verbal	10.55	5.80
Relational	11.66	6.88
Home instability	3.79	2.08
Service use-ever	4.18	2.51

Table 2


Frequencies of types of physical, non-physical, and relational revictimization endorsed by adolescent girls (N=234).

	<i>n</i>	<i>%</i>
Physical Revictimization		
Another person shoved or pushed you?	120	51
Another person threatened to hit or physically harm you?	91	39
Another person hit or slapped you?	85	36
Been in a fight in which you were hit?	84	36
Someone threw something at you to hurt you?	70	30
Someone threatened you with a weapon (gun, knife, club, etc.)?	37	16
A teacher threatened to hurt you?	5	2
Verbal Revictimization		
Another person gave mean looks to you?	173	74
Someone teased you to make you angry?	142	61
Someone picked on you?	109	47
Someone put you down to your face?	80	34
Relational Revictimization		
Another person spread a false rumor about you?	142	61
Another person said things about you to make other people laugh?	129	55
Another person tried to keep others from liking you by saying mean things about you?	125	53
Another person left you out on purpose when it was time to do an activity?	78	33
Another person told you they wouldn't like you unless you did what they wanted you to do?	72	31
Another person didn't let you in the group anymore because they were mad at you?	66	28

Table 3*Bivariate correlations between child maltreatment types, mental health, revictimization, and demographics (N = 234).*


	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1. Emotional abuse	--													
2. Physical abuse	.70***	--												
3. Sexual abuse	.38***	.29***	--											
4. Physical neglect	.60***	.61***	.23***	--										
5. PTSD	.45***	.24***	.31***	.13*	--									
6. Depression	.39***	.15*	.24***	.13*	.74***	--								
7. Revictimization, Total	.37***	.25***	.23***	.16**	.56***	.54***	--							
8. Revictimization, Physical	.31***	.25***	.11	.14*	.44***	.44***	.76***	--						
9. Revictimization, Verbal	.35***	.18**	.20**	.13*	.49***	.47***	.92***	.5***	--					
10. Revictimization Relational	.31***	.22**	.27***	.15*	.53***	.50***	.91***	.49***	.79***	--				
11. Home instability	.26***	.37***	.08	.42***	.04	.06	.08	.08	.09	.04	--			
12. Service use	.38***	.35***	.20**	.33***	.19**	.17**	.22***	.16**	.23***	.19**	.44***	--		
13. Age	.09	.11	.05	.14*	-.08	-.01	.01	-.05	.05	.01	.26***	.19**	--	
14. Race ^a	.16*	.02	.14*	.14*	.06	.08	-.03	-.11	.02	-.01	-.03	.08	.04	--

* $p \leq .05$ ** $p \leq .01$ *** $p \leq .001$ ^a 0 = youths of color; 1 = white




Note. Covariates were non-significant.

Figure 1. Parallel mediation analysis predicting the impact of histories of childhood emotional abuse on revictimization in adolescent girls.




Note. Covariates were non-significant.

Figure 2. Parallel mediation analysis predicting the impact of histories of childhood sexual abuse on revictimization in adolescent girls.



Note. Covariates were non-significant.

Figure 3. Parallel mediation analysis predicting the impact of childhood physical abuse on revictimization in adolescent girls.



Note. Covariates were non-significant.

Figure 4. Parallel mediation analysis predicting the impact of childhood physical neglect on revictimization in adolescent girls.