



ELSEVIER

Contents lists available at ScienceDirect

Journal of Functional Analysis

[www.elsevier.com/locate/jfa](http://www.elsevier.com/locate/jfa)



## Nevanlinna representations in several variables <sup>☆</sup>



J. Agler <sup>a</sup>, R. Tully-Doyle <sup>b</sup>, N.J. Young <sup>c,d,\*</sup>

<sup>a</sup> Department of Mathematics, University of California at San Diego, CA 92103, USA

<sup>b</sup> Department of Mathematics, Hampton University, Hampton, VA 23668, USA

<sup>c</sup> School of Mathematics, Leeds University, Leeds LS2 9JT, England, United Kingdom

<sup>d</sup> School of Mathematics and Statistics, Newcastle University, Newcastle upon Tyne NE3 4LR, England, United Kingdom

### ARTICLE INFO

#### Article history:

Received 28 December 2014

Accepted 2 February 2016

Available online 23 February 2016

Communicated by G. Schechtman

#### MSC:

32A30

30E20

30E05

47B25

47A10

#### Keywords:

Pick class

Cauchy transform

Self-adjoint operator

Resolvent

### ABSTRACT

We generalize to several variables the classical theorem of Nevanlinna that characterizes the Cauchy transforms of positive measures on the real line. We show that for the Loewner class, a large class of analytic functions that have non-negative imaginary part on the upper polyhalf-plane, there are representation formulae in terms of densely-defined self-adjoint operators on a Hilbert space. We find four different representation formulae and we show that every function in the Loewner class has one of the four representations, corresponding precisely to four different growth conditions at infinity.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

<sup>☆</sup> The first author was partially supported by the National Science Foundation Grant on Extending Hilbert Space Operators DMS 1068830 and 1361720. The third author was partially supported by the UK Engineering and Physical Sciences Research Council grants EP/J004545/1 and EP/N03242X/1.

\* Corresponding author at: School of Mathematics and Statistics, Newcastle University, Newcastle upon Tyne NE3 4LR, England, United Kingdom.

E-mail addresses: [ryan.tullydoyle@hamptonu.edu](mailto:ryan.tullydoyle@hamptonu.edu) (R. Tully-Doyle), [Nicholas.Young@ncl.ac.uk](mailto:Nicholas.Young@ncl.ac.uk) (N.J. Young).

### 1. Introduction

In a classic paper [18] of 1922 R. Nevanlinna solved the problem of the determinacy of solutions of the Stieltjes moment problem. *En route* he proved several other theorems that have since been influential; in particular, the following theorem, which characterizes the Cauchy transforms of positive finite measures  $\mu$  on  $\mathbb{R}$ , has had a profound impact on the development of modern analysis. Let  $\mathcal{P}$  denote the Pick class, that is, the set of analytic functions on the upper half-plane,

$$\Pi \stackrel{\text{def}}{=} \{z \in \mathbb{C} : \text{Im } z > 0\},$$

that have non-negative imaginary part on  $\Pi$ .

**Theorem 1.1** (*Nevanlinna’s Representation*). *Let  $h$  be a function defined on  $\Pi$ . There exists a finite positive measure  $\mu$  on  $\mathbb{R}$  such that*

$$h(z) = \int \frac{d\mu}{t - z} \tag{1.1}$$

*if and only if  $h \in \mathcal{P}$  and*

$$\liminf_{y \rightarrow \infty} y |h(iy)| < \infty. \tag{1.2}$$

A closely related theorem, also referred to in the literature as Nevanlinna’s Representation, provides an integral representation for a general element of  $\mathcal{P}$ .

**Theorem 1.2.** *A function  $h : \Pi \rightarrow \mathbb{C}$  belongs to the Pick class  $\mathcal{P}$  if and only if there exist  $a \in \mathbb{R}$ ,  $b \geq 0$  and a finite positive Borel measure  $\mu$  on  $\mathbb{R}$  such that*

$$h(z) = a + bz + \int \frac{1 + tz}{t - z} d\mu(t) \tag{1.3}$$

*for all  $z \in \Pi$ . Moreover, for any  $h \in \mathcal{P}$ , the numbers  $a \in \mathbb{R}$ ,  $b \geq 0$  and the measure  $\mu \geq 0$  in the representation (1.3) are uniquely determined.*

What are the several-variable analogs of Nevanlinna’s theorems? In this paper we shall propose four types of Nevanlinna representation for various subclasses of the  $n$ -variable Pick class  $\mathcal{P}_n$ , where  $\mathcal{P}_n$  is defined to be the set of analytic functions  $h$  on the polyhalf-plane  $\Pi^n$  such that  $\text{Im } h \geq 0$ . In addition, we shall present necessary and sufficient conditions for a function defined on  $\Pi^n$  to possess a representation of a given type in terms of asymptotic growth conditions at  $\infty$ .

The integral representation (1.1) of those functions in the Pick class that satisfy condition (1.2) can be written in the form

$$h(z) = \langle (A - z)^{-1} \mathbf{1}, \mathbf{1} \rangle_{L^2(\mu)},$$

where  $A$  is the operation of multiplication by the independent variable on  $L^2(\mu)$  and  $\mathbf{1}$  is the constant function 1. We propose that an appropriate  $n$ -variable analog of the Cauchy transform is the formula

$$h(z_1, \dots, z_n) = \langle (A - z_1 Y_1 - \dots - z_n Y_n)^{-1} v, v \rangle_{\mathcal{H}} \quad \text{for } z_1, \dots, z_n \in \Pi, \quad (1.4)$$

where  $\mathcal{H}$  is a Hilbert space,  $A$  is a densely defined self-adjoint operator on  $\mathcal{H}$ ,  $Y_1, \dots, Y_n$  are positive contractions on  $\mathcal{H}$  summing to 1 and  $v$  is a vector in  $\mathcal{H}$ .

**Theorem 1.6** below characterizes those functions on  $\Pi^n$  that have a representation of the form (1.4). To state this theorem we require a notion based on the following classical result of Pick [20].

**Theorem 1.3.** *A function  $h$  defined on  $\Pi$  belongs to  $\mathcal{P}$  if and only if the function  $A$  defined on  $\Pi \times \Pi$  by*

$$A(z, w) = \frac{h(z) - \overline{h(w)}}{z - \bar{w}}$$

*is positive semidefinite, that is, for all  $n \geq 1$ ,  $z_1, \dots, z_n \in \Pi$ ,  $c_1, \dots, c_n \in \mathbb{C}$ ,*

$$\sum A(z_j, z_i) \bar{c}_i c_j \geq 0.$$

The following theorem, proved in [2], leads to a generalization of **Theorem 1.3** to two variables. The *Schur class of the polydisc*, denoted by  $\mathcal{S}_n$ , is the set of analytic functions on the polydisc  $\mathbb{D}^n$  that are bounded by 1 in modulus.

**Theorem 1.4.** *A function  $\varphi$  defined on  $\mathbb{D}^2$  belongs to  $\mathcal{S}_2$  if and only if there exist positive semidefinite functions  $A_1$  and  $A_2$  on  $\mathbb{D}^2 \times \mathbb{D}^2$  such that*

$$1 - \overline{\varphi(\mu)}\varphi(\lambda) = (1 - \bar{\mu}_1 \lambda_1)A_1(\lambda, \mu) + (1 - \bar{\mu}_2 \lambda_2)A_2(\lambda, \mu). \quad (1.5)$$

By way of the transformations

$$z = i \frac{1 + \lambda}{1 - \lambda}, \quad \lambda = \frac{z - i}{z + i}, \quad (1.6)$$

and

$$h(z) = i \frac{1 + \varphi(\lambda)}{1 - \varphi(\lambda)}, \quad \varphi(\lambda) = \frac{h(z) - i}{h(z) + i}, \quad (1.7)$$

there is a one-to-one correspondence between functions in the Schur and Pick classes. Under these transformations, **Theorem 1.4** becomes the following generalization of Pick’s theorem to two variables.

**Theorem 1.5.** *A function  $h$  defined on  $\Pi^2$  belongs to  $\mathcal{P}_2$  if and only if there exist positive semidefinite functions  $A_1$  and  $A_2$  on  $\Pi^2 \times \Pi^2$  such that*

$$h(z) - \overline{h(w)} = (z_1 - \overline{w_1})A_1(z, w) + (z_2 - \overline{w_2})A_2(z, w).$$

In the light of [Theorems 1.3 and 1.5](#) we define the *Loewner class*  $\mathcal{L}_n$  to be the set of analytic functions  $h$  on  $\Pi^n$  with the property that there exist  $n$  positive semidefinite functions  $A_1, \dots, A_n$  on  $\Pi^n$  such that

$$h(z) - \overline{h(w)} = \sum_{j=1}^n (z_j - \overline{w_j})A_j(z, w) \tag{1.8}$$

for all  $z, w \in \Pi^n$ . The Loewner class  $\mathcal{L}_n$  played a key role in [\[4\]](#), which gave a generalization to several variables of Loewner’s characterization of the one-variable operator-monotone functions [\[17\]](#). As the following theorem makes clear,  $\mathcal{L}_n$  also has a fundamental role to play in the understanding of Nevanlinna representations in several variables.

**Theorem 1.6.** *A function  $h$  defined on  $\Pi^n$  has a representation of the form [\(1.4\)](#) if and only if  $h \in \mathcal{L}_n$  and*

$$\liminf_{y \rightarrow \infty} y|h(iy, \dots, iy)| < \infty. \tag{1.9}$$

In the cases when  $n = 1$  and  $n = 2$ , [Theorems 1.3 and 1.5](#) assert that  $\mathcal{L}_n = \mathcal{P}_n$ , and so for  $n = 1$ , [Theorem 1.6](#) is Nevanlinna’s classical [Theorem 1.1](#), and when  $n = 2$ , [Theorem 1.6](#) is a straightforward generalization of that result to two variables. When there are more than two variables, it is known that the Loewner class is a proper subset of the Pick class,  $\mathcal{L}_n \neq \mathcal{P}_n$  [\[19,22\]](#). Nevertheless, Nevanlinna’s result survives as a theorem about the representation of elements of  $\mathcal{L}_n$ . Other than the work in [\[11\]](#) very little is known about the representation of functions in  $\mathcal{P}_n$  for three or more variables.

For a function  $h$  on  $\Pi^n$ , we call the formula [\(1.4\)](#) a *Nevanlinna representation of type 1*. Thus, [Theorem 1.6](#) can be rephrased as the assertion that  $h$  has a Nevanlinna representation of type 1 if and only if  $h \in \mathcal{L}_n$  and  $h$  satisfies condition [\(1.9\)](#). Somewhat more complicated representation formulae are needed to generalize [Theorem 1.2](#). We identify three further representation formulae, of increasing generality, and show that every function in  $\mathcal{L}_n$  has a representation of one or more of the four types.

For a function  $h$  defined on  $\Pi^n$ , we refer to a formula

$$h(z_1, \dots, z_n) = a + \langle (A - z_1 Y_1 - \dots - z_n Y_n)^{-1} v, v \rangle_{\mathcal{H}}$$

for  $z_1, \dots, z_n \in \Pi$ ,

(1.10)

where  $a$  is a constant,  $\mathcal{H}$  is a Hilbert space,  $A$  is a densely defined self-adjoint operator on  $\mathcal{H}$ ,  $Y_1, \dots, Y_n$  are positive contractions on  $\mathcal{H}$  summing to 1 and  $v$  is a vector in  $\mathcal{H}$ , as a *Nevanlinna representation of type 2*.

**Theorem 1.7.** *A function  $h$  defined on  $\Pi^n$  has a Nevanlinna representation of type 2 if and only if  $h \in \mathcal{L}_n$  and*

$$\liminf_{y \rightarrow \infty} y \operatorname{Im} h(iy, \dots, iy) < \infty. \tag{1.11}$$

A *Nevanlinna representation of type 3* of a function  $h$  defined on  $\Pi^n$  is of the form

$$h(z) = a + \langle (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 - iA)^{-1}v, v \rangle \quad \text{for all } z \in \Pi^n$$

for some real  $a$ , some self-adjoint operator  $A$  and some vector  $v$ , where  $Y_1, \dots, Y_n$  are operators as in equation (1.4) above and  $z_Y = z_1 Y_1 + \dots + z_n Y_n$ .

**Theorem 1.8.** *A function  $h$  defined on  $\Pi^n$  has a Nevanlinna representation of type 3 if and only if  $h \in \mathcal{L}_n$  and*

$$\liminf_{y \rightarrow \infty} \frac{1}{y} \operatorname{Im} h(iy, \dots, iy) = 0.$$

Finally, *Nevanlinna representations of type 4* are given by the formula

$$h(z) = a + \langle M(z)v, v \rangle, \tag{1.12}$$

where  $a \in \mathbb{R}$  and  $M(z)$  is an operator of the form

$$\begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \left( z_P \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix}^{-1}, \tag{1.13}$$

acting on an orthogonal direct sum of Hilbert spaces  $\mathcal{N} \oplus \mathcal{M}$ . In equation (1.12),  $v$  is a vector in  $\mathcal{N} \oplus \mathcal{M}$ . In equation (1.13),  $A$  is a densely-defined self-adjoint operator acting on  $\mathcal{M}$  and  $z_P$  is the operator acting on  $\mathcal{N} \oplus \mathcal{M}$  via the formula

$$z_P = \sum z_i P_i$$

where  $P_1, \dots, P_n$  are pairwise orthogonal projections acting on  $\mathcal{N} \oplus \mathcal{M}$  that sum to 1.

**Theorem 1.9.** *Let  $h$  be a function defined on  $\Pi^n$ . Then  $h$  has a Nevanlinna representation of type 4 if and only if  $h \in \mathcal{L}_n$ .*

A weaker, “generic” version of [Theorem 1.9](#) appeared in [\[4, Theorem 6.9\]](#), where it was used to show that elements in  $\mathcal{L}_n$  are locally operator-monotone.

It turns out that for  $1 \leq k \leq 4$ , if  $h$  is a function on  $\Pi^n$  and  $h$  has a Nevanlinna representation of type  $k$ , then for  $k \leq j \leq 4$ ,  $h$  also has a Nevanlinna representation of type  $j$ . Thus, it is natural to define the *type* of a function in  $\mathcal{L}_n$  to be the smallest  $k$  such that  $h$  has a Nevanlinna representation of type  $k$ .

For  $h \in \mathcal{L}_n$  the type of  $h$  can be characterized in function-theoretic terms through the use of a geometric idea due to Carathéodory. A *carapoint* for a function  $\varphi$  in the Schur class  $\mathcal{S}_n$  is a point  $\tau \in \mathbb{T}$  such that

$$\liminf_{\lambda \rightarrow \tau} \frac{1 - |\varphi(\lambda)|}{1 - \|\lambda\|_\infty} < \infty,$$

where

$$\|\lambda\|_\infty = \max_{1 \leq i \leq n} |\lambda_i|.$$

Carathéodory introduced this notion in one variable in [\[9\]](#), along the way to refining earlier results of Julia [\[14\]](#). The following was Carathéodory’s main result; the notation  $\lambda \xrightarrow{\text{nt}} \tau$  means that  $\lambda$  tends nontangentially to  $\tau$ .

**Theorem 1.10.** *Let  $\varphi \in \mathcal{S}_1$ ,  $\tau \in \mathbb{T}$ . If  $\tau$  is a carapoint for  $\varphi$ , then  $\varphi$  is nontangentially differentiable at  $\tau$ , that is, there exist values  $\varphi(\tau)$  and  $\varphi'(\tau)$  such that*

$$\lim_{\lambda \xrightarrow{\text{nt}} \tau} \frac{\varphi(\lambda) - \varphi(\tau) - \varphi'(\tau)(\lambda - \tau)}{\lambda - \tau} = 0.$$

*In particular, if  $\tau$  is a carapoint for  $\varphi$  then there exists a unique point  $\varphi(\tau) \in \mathbb{T}$  such that  $\varphi(\lambda) \rightarrow \varphi(\tau)$  as  $\lambda \xrightarrow{\text{nt}} \tau$ .*

In several variables, carapoints have been studied in [\[1,13,3\]](#). The strong conclusion of nontangential differentiability is lost in several variables; however, at a carapoint  $\tau$ , there still exists a unimodular nontangential limit  $\varphi(\tau)$ .

As the point  $\chi = (1, \dots, 1)$  is transformed to the point  $\infty = (\infty, \dots, \infty)$  by equation [\(1.6\)](#), it is natural to say that a function  $h \in \mathcal{L}_n$  has a carapoint at  $\infty$  if the associated Schur function  $\varphi$ , given by the transformation in equation [\(1.7\)](#), has a carapoint at  $\chi$ , and in that case to define  $h(\infty)$  by

$$h(\infty) = i \frac{1 + \varphi(\chi)}{1 - \varphi(\chi)}. \tag{1.14}$$

The connection between carapoints and function types is given in the following theorem.

**Theorem 1.11.** For a function  $h \in \mathcal{L}_n$ ,

- (1)  $h$  is of type 1 if and only if  $\infty$  is a carapoint of  $h$  and  $h(\infty) = 0$ ;
- (2)  $h$  is of type 2 if and only if  $\infty$  is a carapoint of  $h$  and  $h(\infty) \in \mathbb{R} \setminus \{0\}$ ;
- (3)  $h$  is of type 3 if and only if  $\infty$  is not a carapoint of  $h$ ;
- (4)  $h$  is of type 4 if and only if  $\infty$  is a carapoint of  $h$  and  $h(\infty) = \infty$ .

The paper is structured as follows. As is clear from the formulae used to define the various Nevanlinna representations, Nevanlinna representations are generalizations of the resolvent of a self-adjoint operator. These *structured resolvents*, studied in Sections 2 and 3, are analytic operator-valued functions on the polyhalf-plane  $\Pi^n$  with non-negative imaginary part, fully analogous to the familiar resolvent operator. There are also *structured resolvent identities* for them, studied in Section 10 of the paper.

In modern texts Nevanlinna's representation is derived from the Herglotz Representation with the aid of the Cayley transform [16,10]. In Section 4 we introduce the  $n$ -variable *strong Herglotz class* and then prove Theorem 1.9 by applying the Cayley transform to Theorem 1.8 of [2].

In Section 5 we derive the Nevanlinna representations of types 3, 2, and 1, we show how they arise naturally from the underlying Hilbert space geometry and we prove slight strengthenings of Theorems 1.6, 1.7 and 1.8. In Section 6 we give function-theoretic conditions for a function  $h \in \mathcal{L}_n$  to possess a representation of a given type.

In Section 7 we introduce the notion of carapoints for functions in the Pick class and in Section 8 we establish the criteria in Theorem 1.11 for the type of a function using the language of carapoints.

In Section 9 we give the growth estimates for functions in  $\mathcal{L}_n$  that flow from our analysis of structured resolvents, and in Section 10 we present resolvent identities for structured resolvents.

Results related to ours from a system-theoretic perspective have been obtained in recent works of J.A. Ball and D. Kalyuzhnyi-Verbovetzkyi [6,7]. See also [8], where Krein space methods are applied to similar problems.

## 2. Structured resolvents of operators

The resolvent operator  $(A - z)^{-1}$  of a densely defined self-adjoint operator  $A$  on a Hilbert space plays a prominent role in spectral theory. It has the following properties.

- (1) It is an analytic bounded operator-valued function of  $z$  in the upper half-plane  $\Pi$ ;
- (2) it satisfies the growth estimate  $\|(A - z)^{-1}\| \leq 1/\text{Im } z$  for  $z \in \Pi$ ;
- (3)  $(A - z)^{-1}$  has non-negative imaginary part for all  $z \in \Pi$ ;
- (4) it satisfies the "resolvent identity".

Here we are interested in several-variable analogs of the resolvent. These will again be operator-valued analytic functions with non-negative imaginary part, but now on

the polyhalf-plane  $\Pi^n$ . Because of the additional complexities in several variables we encounter three different types of resolvent; all of them have the four listed properties, with very slight modifications, and therefore deserve the name *structured resolvent*.

For any Hilbert space  $\mathcal{H}$ , a *positive decomposition* of  $\mathcal{H}$  will mean an  $n$ -tuple  $Y = (Y_1, \dots, Y_n)$  of positive contractions on  $\mathcal{H}$  that sum to the identity operator. For any  $z = (z_1, \dots, z_n) \in \mathbb{C}^n$  and any  $n$ -tuple  $T = (T_1, \dots, T_n)$  of bounded operators we denote by  $z_T$  the operator  $\sum_j z_j T_j$ . Here each  $T_j$  is a bounded operator from  $\mathcal{H}_1$  to  $\mathcal{H}_2$ , for some Hilbert spaces  $\mathcal{H}_1, \mathcal{H}_2$ , so that  $z_T$  is also a bounded operator from  $\mathcal{H}_1$  to  $\mathcal{H}_2$ .

**Definition 2.1.** Let  $A$  be a closed densely defined self-adjoint operator on a Hilbert space  $\mathcal{H}$  and let  $Y$  be a positive decomposition of  $\mathcal{H}$ . The *structured resolvent of  $A$  of type 2* corresponding to  $Y$  is the operator-valued function

$$z \mapsto (A - z_Y)^{-1} : \Pi^n \rightarrow \mathcal{L}(\mathcal{H}).$$

The following observation is essentially [4, Lemma 6.25].

**Proposition 2.2.** For  $A$  and  $Y$  as in Definition 2.1 the structured resolvent  $(A - z_Y)^{-1}$  is well defined on  $\Pi^n$  and satisfies, for all  $z \in \Pi^n$ ,

$$\|(A - z_Y)^{-1}\| \leq \frac{1}{\min_j \operatorname{Im} z_j}. \tag{2.1}$$

Moreover

$$\begin{aligned} \operatorname{Im} ((A - z_Y)^{-1}) &= (A - z_Y^*)^{-1} (\operatorname{Im} z_Y) (A - z_Y)^{-1} \\ &= (A - z_Y)^{-1} (\operatorname{Im} z_Y) (A - z_Y^*)^{-1} \\ &\geq 0. \end{aligned} \tag{2.2}$$

The range of the bounded operator  $(A - z_Y)^{-1}$  is of course  $\mathcal{D}(A)$ , the domain of  $A$ .

**Proof of Proposition 2.2.** For any vector  $\xi$  in the domain of  $A$ ,

$$\begin{aligned} \|(A - z_Y)\xi\| \|\xi\| &\geq |\langle (A - z_Y)\xi, \xi \rangle| \\ &\geq |\operatorname{Im} \langle (A - z_Y)\xi, \xi \rangle| \\ &= \langle (\operatorname{Im} z_Y)\xi, \xi \rangle \\ &= \sum_j (\operatorname{Im} z_j) \langle Y_j \xi, \xi \rangle \\ &\geq (\min_j \operatorname{Im} z_j) \left\langle \sum_j Y_j \xi, \xi \right\rangle \\ &= (\min_j \operatorname{Im} z_j) \|\xi\|^2. \end{aligned}$$

Thus  $A - z_Y$  has lower bound  $\min_j \operatorname{Im} z_j > 0$ , and so has a bounded left inverse. A similar argument with  $z$  replaced by  $\bar{z}$  shows that  $(A - z_Y)^*$  also has a bounded left inverse, and so  $A - z_Y$  has a bounded inverse and the inequality (2.1) holds.

The identities (2.2) are easy.  $\square$

Resolvents of type 2 are the simplest several-variable analogues of the familiar one-variable resolvent but they are not sufficient for the analysis of the several-variable Pick class. To this end we introduce two further generalizations. Let us first recall some basic facts about closed unbounded operators.

**Lemma 2.3.** *Let  $T$  be a closed densely defined operator on a Hilbert space  $\mathcal{H}$ , with domain  $\mathcal{D}(T)$ . The operator  $1 + T^*T$  is a bijection from  $\mathcal{D}(T^*T)$  to  $\mathcal{H}$ , and the operators*

$$B \stackrel{\text{def}}{=} (1 + T^*T)^{-1}, \quad C \stackrel{\text{def}}{=} T(1 + T^*T)^{-1}$$

*are everywhere defined and contractive on  $\mathcal{H}$ . Moreover  $B$  is self-adjoint and positive, and  $\operatorname{ran} C \subset \mathcal{D}(T^*)$ .*

**Proof.** All these statements are proved in [21, Sections 118, 119], although the final statement about  $\operatorname{ran} C$  is not explicitly stated. We must show that for all  $v \in \mathcal{H}$  there exists  $y \in \mathcal{H}$  such that, for all  $h \in \mathcal{H}$ ,

$$\langle Th, Cv \rangle = \langle h, y \rangle.$$

It is straightforward to check that this relation holds for  $y = v - Bv$ , and so  $\operatorname{ran} C \subset \mathcal{D}(T^*)$ .  $\square$

**Definition 2.4.** Let  $A$  be a closed densely defined self-adjoint operator on a Hilbert space  $\mathcal{H}$  and let  $Y$  be a positive decomposition of  $\mathcal{H}$ . The *structured resolvent of  $A$  of type 3* corresponding to  $Y$  is the operator-valued function  $M : \Pi^n \rightarrow \mathcal{L}(\mathcal{H})$  given by

$$M(z) = (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 - iA)^{-1}. \tag{2.3}$$

We denote the  $\ell_1$  norm on  $\mathbb{C}^n$  by  $\|\cdot\|_1$ . Note that  $\|z_Y\| \leq \|z\|_1$  for all  $z \in \mathbb{C}^n$  and all positive decompositions  $Y$ .

**Proposition 2.5.** *For  $A$  and  $Y$  as in Definition 2.4 the structured resolvent  $M(z)$  of type 3 given by equation (2.3) is well defined as a bounded operator on  $\mathcal{H}$  for all  $z \in \Pi^n$  and satisfies*

$$\|M(z)\| \leq (1 + 2\|z\|_1) \left( 1 + \frac{1 + \|z\|_1}{\min_j \operatorname{Im} z_j} \right). \tag{2.4}$$

**Proof.** Since

$$1 + z_Y A = 1 - iz_Y + iz_Y(1 - iA) : \mathcal{D}(A) \rightarrow \mathcal{H}$$

and  $(1 - iA)^{-1}$  is a contraction on all of  $\mathcal{H}$ , with range  $\mathcal{D}(A)$ , the operator  $(1 + z_Y A)(1 - iA)^{-1}$  is well defined as an operator on  $\mathcal{H}$  and

$$\begin{aligned} \|(1 + z_Y A)(1 - iA)^{-1}\| &= \|(1 - iz_Y)(1 - iA)^{-1} + iz_Y\| \\ &\leq \|1 - iz_Y\| + \|z_Y\| \\ &\leq 1 + 2\|z_Y\| \\ &\leq 1 + 2\|z\|_1. \end{aligned} \tag{2.5}$$

Similarly  $(1 - iA)(A - z_Y)^{-1}$  is well defined on  $\mathcal{H}$ , and since

$$i(A - z_Y) = -(1 - iA) + (1 - iz_Y) : \mathcal{D}(A) \rightarrow \mathcal{H}$$

we have

$$i = -(1 - iA)(A - z_Y)^{-1} + (1 - iz_Y)(A - z_Y)^{-1} : \mathcal{H} \rightarrow \mathcal{H}.$$

Thus, by virtue of the bound (2.1),

$$\begin{aligned} \|(1 - iA)(A - z_Y)^{-1}\| &= \|i - (1 - iz_Y)(A - z_Y)^{-1}\| \\ &\leq 1 + \|1 - iz_Y\| \|(A - z_Y)^{-1}\| \\ &\leq 1 + \frac{1 + \|z\|_1}{\min_j \operatorname{Im} z_j}. \end{aligned} \tag{2.6}$$

On combining the estimates (2.6) and (2.5) we obtain the bound (2.4).  $\square$

The following alternative formula for the structured resolvent of type 3, valid on the dense subspace  $\mathcal{D}(A)$  of  $\mathcal{H}$ , allows us to show that  $\operatorname{Im} M(z) \geq 0$ .

**Proposition 2.6.** *For  $A$  and  $Y$  as in Definition 2.4 and  $z \in \Pi^n$*

$$M(z)|_{\mathcal{D}(A)} = (1 - iA) \{ (A - z_Y)^{-1} - A(1 + A^2)^{-1} \} (1 + iA) \tag{2.7}$$

$$= (1 - iA)(A - z_Y)^{-1}(1 + iA) - A : \mathcal{D}(A) \rightarrow \mathcal{H}. \tag{2.8}$$

Moreover, for every  $v \in \mathcal{D}(A)$ ,

$$\operatorname{Im} \langle M(z)v, v \rangle = \langle (1 - iA)(A - z_Y^*)^{-1}(\operatorname{Im} z_Y)(A - z_Y)^{-1}(1 + iA)v, v \rangle \geq 0. \tag{2.9}$$

**Proof.** By Lemma 2.3 the operator  $A(1 + A^2)^{-1}$  is contractive on  $\mathcal{H}$  and has range contained in  $\mathcal{D}(A)$ . On  $\mathcal{D}(A^2)$  we have the identity

$$1 + z_Y A = 1 + A^2 - (A - z_Y)A.$$

Since  $(1 + A^2)^{-1}$  maps  $\mathcal{H}$  into  $\mathcal{D}(A^2)$  we have

$$(1 + z_Y A)(1 + A^2)^{-1} = 1 - (A - z_Y)A(1 + A^2)^{-1} : \mathcal{H} \rightarrow \mathcal{H},$$

and therefore

$$(A - z_Y)^{-1}(1 + z_Y A)(1 + A^2)^{-1} = (A - z_Y)^{-1} - A(1 + A^2)^{-1} : \mathcal{H} \rightarrow \mathcal{D}(A). \tag{2.10}$$

Clearly

$$(1 + A^2)^{-1}(1 + iA) = (1 - iA)^{-1} \quad \text{on } \mathcal{D}(A)$$

and so, on multiplying equation (2.10) fore-and-aft by  $1 \pm iA$ , we deduce that, as operators from  $\mathcal{D}(A)$  to  $\mathcal{H}$ ,

$$\begin{aligned} M(z)|_{\mathcal{D}(A)} &= (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 - iA)^{-1} \\ &= (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 + A^2)^{-1}(1 + iA) \\ &= (1 - iA) \{ (A - z_Y)^{-1} - A(1 + A^2)^{-1} \} (1 + iA). \end{aligned}$$

This establishes equation (2.7).

The expression (2.8) follows from equation (2.7) since

$$(1 - iA)A(1 + A^2)^{-1}(1 + iA) = A \quad \text{on } \mathcal{D}(A).$$

By equation (2.8) we have, for any  $z \in \Pi^n$  and  $v \in \mathcal{D}(A)$ ,

$$\begin{aligned} \operatorname{Im} \langle M(z)v, v \rangle &= \operatorname{Im} \langle (1 - iA)(A - z_Y)^{-1}(1 + iA)v, v \rangle - \operatorname{Im} \langle Av, v \rangle \\ &= \operatorname{Im} \langle (A - z_Y)^{-1}(1 + iA)v, (1 + iA)v \rangle \end{aligned}$$

and hence, by equation (2.2),

$$\operatorname{Im} \langle M(z)v, v \rangle = \langle (A - z_Y^*)^{-1}(\operatorname{Im} z_Y)(A - z_Y)^{-1}(1 + iA)v, (1 + iA)v \rangle,$$

and so equation (2.9) holds.  $\square$

**Corollary 2.7.** For  $A$  and  $Y$  as in Definition 2.4 the structured resolvent  $M(z)$  given by equation (2.3) satisfies  $\operatorname{Im} M(z) \geq 0$  for all  $z \in \Pi^n$ .

For, by [Propositions 2.5 and 2.6](#),  $M(z)$  is a bounded operator on  $\mathcal{H}$ , and  $\text{Im}\langle M(z)v, v \rangle \geq 0$  for  $v \in \mathcal{D}(A)$ . The conclusion follows by the density of  $\mathcal{D}(A)$  and continuity.

In the case of bounded  $A$  there is yet another expression for the structured resolvent of type 3.

**Proposition 2.8.** *If  $A$  is a bounded self-adjoint operator on  $\mathcal{H}$  and  $Y$  is a positive decomposition of  $\mathcal{H}$  then, for  $z \in \Pi^n$ ,*

$$M(z) = (1 + iA)^{-1}(1 + Az_Y)(A - z_Y)^{-1}(1 + iA). \tag{2.11}$$

**Proof.** Since  $A$  is bounded it is defined on all of  $\mathcal{H}$ . We have

$$1 + Az_Y = 1 + A^2 - A(A - z_Y)$$

and hence

$$(1 + Az_Y)(A - z_Y)^{-1} = (1 + A^2)(A - z_Y)^{-1} - A.$$

Thus

$$\begin{aligned} (1 + iA)^{-1}(1 + Az_Y)(A - z_Y)^{-1}(1 + iA) &= (1 - iA)(A - z_Y)^{-1}(1 + iA) - A \\ &= M(z) \end{aligned}$$

by equation [\(2.8\)](#).  $\square$

**Remark 2.9.** In the case of unbounded  $A$  the expression [\(2.11\)](#) for  $M(z)$  is valid wherever it is defined, but it is not to be expected that this will be a dense subspace of  $\mathcal{H}$  in general.

Here are two examples of structured resolvents of type 3, one on  $\mathbb{C}^2$  and one on an infinite-dimensional space.

**Example 2.10.** Let

$$\mathcal{H} = \mathbb{C}^2, \quad A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad Y_1 = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, \quad Y_2 = 1 - Y_1, \quad Y = (Y_1, Y_2).$$

Then

$$\begin{aligned} M(z) &= (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 - iA)^{-1} \\ &= \frac{1}{1 - z_1 z_2} \begin{bmatrix} (1 + z_1)(1 + z_2) & -i(z_1 - z_2) \\ i(z_1 - z_2) & -(1 - z_1)(1 - z_2) \end{bmatrix}. \end{aligned}$$

**Example 2.11.** Let  $\mathcal{H} = L^2(\mathbb{R})$ , let  $A$  be the operation of multiplication by the independent variable  $t$  and let  $Y = (P, Q)$  where  $P, Q$  are the orthogonal projection operators onto the subspaces of even and odd functions respectively in  $L^2$ . Thus

$$Pf(t) = \frac{1}{2} \{f(t) + f(-t)\}, \quad Qf(t) = \frac{1}{2} \{f(t) - f(-t)\}.$$

Let  $Y' = (Q, P)$ . Note that

$$PA = AQ, \quad QA = AP$$

and hence

$$z_Y A = A z_{Y'}, \quad z_{Y'} A = A z_Y, \quad z_Y z_{Y'} = z_1 z_2 = z_{Y'} z_Y.$$

It follows that  $z_Y$  and  $z_{Y'}$  commute with  $A^2$ , and it may be checked that

$$(A - z_Y)^{-1} = (A^2 - z_1 z_2)^{-1} (z_{Y'} + A) = (z_{Y'} + A) (A^2 - z_1 z_2)^{-1}$$

and hence

$$(A - z_Y)^{-1} (1 + z_Y A) = (A^2 - z_1 z_2)^{-1} ((1 + A^2) z_{Y'} + (1 + z_1 z_2) A).$$

A straightforward calculation now shows that the structured resolvent  $M(z)$  of  $A$  corresponding to  $Y$  is given by

$$(M(z)f)(t) = \frac{(\frac{1}{2}(z_1 + z_2)(1 + t^2) + (1 + z_1 z_2)t) f(t) + \frac{1}{2}(z_2 - z_1)(1 - it)^2 f(-t)}{t^2 - z_1 z_2}$$

for all  $z \in \Pi^2$ ,  $f \in L^2(\mathbb{R})$  and  $t \in \mathbb{R}$ . In particular, we note for future use that if  $f$  is an even function,

$$(M(z)f)(t) = \frac{t(1 + z_1 z_2) + (1 - it)(it z_1 + z_2)}{t^2 - z_1 z_2} f(t). \tag{2.12}$$

**3. The matricial resolvent**

The third and last form of structured resolvent that we consider has a  $2 \times 2$  matricial form. As will become clear, this extra complication is needed for the description of the most general type of function in the several-variable Loewner class.

By an *orthogonal decomposition* of a Hilbert space  $\mathcal{H}$  we shall mean an  $n$ -tuple  $P = (P_1, \dots, P_n)$  of orthogonal projection operators with pairwise orthogonal ranges such that  $\sum_{j=1}^n P_j$  is the identity operator.

**Proposition 3.1.** *Let  $\mathcal{H}$  be the orthogonal direct sum of Hilbert spaces  $\mathcal{N}, \mathcal{M}$ , let  $A$  be a densely defined self-adjoint operator on  $\mathcal{M}$  with domain  $\mathcal{D}(A)$  and let  $P$  be an orthogonal decomposition of  $\mathcal{H}$ . For every  $z \in \Pi^n$  the operator on  $\mathcal{H}$  given with respect to the decomposition  $\mathcal{N} \oplus \mathcal{M}$  by the matricial formula*

$$\begin{aligned}
 M(z) &= \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \\
 &\quad \times \left( z_P \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix}^{-1}
 \end{aligned} \tag{3.1}$$

is a bounded operator defined on all of  $\mathcal{H}$ , and

$$\|M(z)\| \leq (1 + \sqrt{10}\|z\|_1) \left( 1 + \frac{1 + \sqrt{2}\|z\|_1}{\min_j \operatorname{Im} z_j} \right). \tag{3.2}$$

**Proof.** Let  $z \in \Pi^n$ . Let the projection  $P_j$  have operator matrix

$$P_j = \begin{bmatrix} X_j & B_j \\ B_j^* & Y_j \end{bmatrix} \tag{3.3}$$

with respect to the decomposition  $\mathcal{H} = \mathcal{N} \oplus \mathcal{M}$ . Then

$$X = (X_1, \dots, X_n), \quad Y = (Y_1, \dots, Y_n)$$

are positive decompositions of  $\mathcal{N}, \mathcal{M}$  respectively, and

$$B = (B_1, \dots, B_n), \quad B^* = (B_1^*, \dots, B_n^*)$$

are  $n$ -tuples of contractions summing to 0, from  $\mathcal{M}$  to  $\mathcal{N}$  and from  $\mathcal{N}$  to  $\mathcal{M}$  respectively. Since the  $B_j$  are contractions we have

$$\|z_B\| \leq \|z\|_1.$$

For any  $z \in \mathbb{C}^n$ ,

$$z_P = \begin{bmatrix} z_X & z_B \\ z_{B^*} & z_Y \end{bmatrix}. \tag{3.4}$$

Consider the third and fourth factors in the product on the right hand side of equation (3.1); the product of these two factors is well defined as an operator on  $\mathcal{H}$  since  $(1 - iA)^{-1}$  maps  $\mathcal{M}$  to  $\mathcal{D}(A)$ . It is even a bounded operator, since, by virtue of equation (3.4),

$$\left( z_P \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix}^{-1} = \begin{bmatrix} iz_X & z_B A(1 - iA)^{-1} \\ iz_{B^*} & (1 + z_Y A)(1 - iA)^{-1} \end{bmatrix}. \tag{3.5}$$

Since

$$\|A(1 - iA)^{-1}\| = \|i(1 - (1 - iA)^{-1})\| \leq 2$$

we can immediately see that the operator (3.5) is bounded. We can get an estimate of the norm of the operator matrix (3.5) if we replace each of the four operator entries by an upper bound for its norm. We find that

$$\begin{aligned} \left\| \left( z_P \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix}^{-1} \right\| &\leq \left\| \begin{bmatrix} \|z\|_1 & 2\|z\|_1 \\ \|z\|_1 & 1 + 2\|z\|_1 \end{bmatrix} \right\| \\ &\leq 1 + \|z\|_1 \left\| \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix} \right\| \\ &= 1 + \sqrt{10}\|z\|_1. \end{aligned} \tag{3.6}$$

Now consider the second factor in the definition (3.1) of  $M(z)$ . We find that

$$\begin{aligned} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} &= \begin{bmatrix} 1 & -z_B \\ 0 & A - z_Y \end{bmatrix}^{-1} \\ &= \begin{bmatrix} 1 & z_B(A - z_Y)^{-1} \\ 0 & (A - z_Y)^{-1} \end{bmatrix}, \end{aligned} \tag{3.7}$$

which maps  $\mathcal{H}$  into  $\mathcal{N} \oplus \mathcal{D}(A)$ . Hence the product of the first two factors in the product on the right hand side of equation (3.1) is

$$\begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} = \begin{bmatrix} -i & -iz_B(A - z_Y)^{-1} \\ 0 & (1 - iA)(A - z_Y)^{-1} \end{bmatrix}. \tag{3.8}$$

Since

$$\begin{aligned} \|(1 - iA)(A - z_Y)^{-1}\| &= \|(1 - iz_Y)(A - z_Y)^{-1} - i\| \\ &\leq 1 + \|1 - iz_Y\| \|(A - z_Y)^{-1}\| \\ &\leq 1 + \frac{1 + \|z\|_1}{\min_j \operatorname{Im} z_j} \end{aligned}$$

we deduce from equation (3.8) that

$$\begin{aligned} &\left\| \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \right\| \\ &\leq \left\| \begin{bmatrix} 1 & \|z\|_1 \|(A - z_Y)^{-1}\| \\ 0 & 1 + (1 + \|z\|_1)\|(A - z_Y)^{-1}\| \end{bmatrix} \right\| \\ &\leq 1 + \left\| \begin{bmatrix} 0 & \|z\|_1 \\ 0 & 1 + \|z\|_1 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & \|(A - z_Y)^{-1}\| \end{bmatrix} \right\| \end{aligned}$$

$$\leq 1 + \frac{1 + \sqrt{2}\|z\|_1}{\min_j \operatorname{Im} z_j}. \tag{3.9}$$

On combining the estimates (3.9) and (3.6) we obtain the bound (3.2) for  $\|M(z)\|$ .  $\square$

**Remark 3.2.** On multiplying together the expressions (3.8) and (3.5) we obtain the formula

$$M(z) = \begin{bmatrix} z_X + z_B(A - z_Y)^{-1}z_{B^*} & -iz_B(A - z_Y)^{-1}(1 + iA) \\ i(1 - iA)(A - z_Y)^{-1}z_{B^*} & (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 - iA)^{-1} \end{bmatrix}.$$

Notice in particular that the (2, 2) entry (that is, the compression of  $M(z)$  to  $\mathcal{M}$ ) is the structured resolvent of  $A$  of type 3 corresponding to  $Y$ , the compression of  $P$  to  $\mathcal{M}$ , as in equation (2.3).

**Definition 3.3.** Let  $\mathcal{H}$  be the orthogonal direct sum of Hilbert spaces  $\mathcal{N}, \mathcal{M}$ , let  $A$  be a densely defined self-adjoint operator on  $\mathcal{M}$  with domain  $\mathcal{D}(A)$  and let  $P$  be an orthogonal decomposition of  $\mathcal{H}$ . The *structured resolvent of  $A$  of type 4* corresponding to  $P$  is the operator-valued function  $M : \Pi^n \rightarrow \mathcal{L}(\mathcal{H})$  given by equation (3.1).

We shall also refer to  $M(z)$  as the *matricial resolvent of  $A$  with respect to  $P$* . The important property that  $\operatorname{Im} M(z) \geq 0$  is not at once apparent from the formula (3.1); as with structured resolvents of type 3, there are alternative formulae from which this property is more easily shown. Once again the alternatives suffer the minor drawback that they give  $M(z)$  only on a dense subspace of  $\mathcal{H}$ .

**Proposition 3.4.** *With the notation of Definition 3.3, as operators on  $\mathcal{N} \oplus \mathcal{D}(A)$ ,*

$$M(z) = \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A(1 + A^2)^{-1} \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} \right) \times \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right)^{-1} \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix} \tag{3.10}$$

$$= \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right)^{-1} \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & A \end{bmatrix} \tag{3.11}$$

$$= \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \left( z_P \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & A \end{bmatrix} \tag{3.12}$$

for all  $z \in \Pi^n$ . Moreover, for all  $z, w \in \Pi^n$ ,

$$\begin{aligned}
 M(z) - M(w)^* &= \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - w_P^* \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \times \\
 &\quad (z_P - w_P^*) \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right)^{-1} \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix}
 \end{aligned}
 \tag{3.13}$$

on  $\mathcal{N} \oplus \mathcal{D}(A)$ .

**Proof.** By Lemma 2.3 the operators  $(1 + A^2)^{-1}$  and

$$C \stackrel{\text{def}}{=} \text{Im}(1 - iA)^{-1} = A(1 + A^2)^{-1}$$

are self-adjoint contractions defined on all of  $\mathcal{M}$ . Furthermore,

$$\text{ran}(1 + A^2)^{-1} = \mathcal{D}(A^2), \quad \text{ran } C \subset \mathcal{D}(A).$$

We claim that, as operators on  $\mathcal{N} \oplus \mathcal{D}(A)$ ,

$$\begin{aligned}
 \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \left( z_P \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) &= \\
 \left( \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} z_P \right)^{-1}.
 \end{aligned}
 \tag{3.14}$$

We have

$$\begin{aligned}
 &\left( z_P \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} z_P \right) \\
 &= \begin{bmatrix} 0 & 0 \\ 0 & C \end{bmatrix} + z_P \begin{bmatrix} 1 & 0 \\ 0 & AC \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} z_P - z_P \begin{bmatrix} 0 & 0 \\ 0 & C \end{bmatrix} z_P \\
 &= \begin{bmatrix} 0 & 0 \\ 0 & C \end{bmatrix} + z_P \left( \begin{bmatrix} 1 & 0 \\ 0 & AC \end{bmatrix} - 1 \right) + \left( 1 - \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} \right) z_P - z_P \begin{bmatrix} 0 & 0 \\ 0 & C \end{bmatrix} z_P \\
 &= \begin{bmatrix} 0 & 0 \\ 0 & C \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & AC \end{bmatrix} z_P - z_P \begin{bmatrix} 0 & 0 \\ 0 & C \end{bmatrix} z_P \\
 &= \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} \right).
 \end{aligned}$$

This is an identity between operators on  $\mathcal{H}$ , in both cases a composition  $\mathcal{H} \rightarrow \mathcal{N} \oplus \mathcal{D}(A) \rightarrow \mathcal{H}$ , and moreover the first factor on the left hand side and the second factor on the right hand side are invertible, from  $\mathcal{N} \oplus \mathcal{D}(A)$  to  $\mathcal{H}$  and from  $\mathcal{H}$  to  $\mathcal{N} \oplus \mathcal{D}(A)$  respectively. We may pre- and post-multiply appropriately to obtain equation (3.14), but note that the equation is then only valid as an identity between operators on  $\mathcal{N} \oplus \mathcal{D}(A)$ .

On combining equations (3.1) and (3.14) we deduce that

$$M(z) = \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} \right) \times \\ \left( \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} z_P \right)^{-1} \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix}^{-1}.$$

Since

$$\begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 + A^2 \end{bmatrix}^{-1} \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix}$$

and

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 + A^2 \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} z_P \right) = \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P,$$

we deduce further that

$$M(z) = \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} \right) \times \\ \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right)^{-1} \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix}, \tag{3.15}$$

which proves equation (3.10). It is straightforward to verify that

$$\left( \begin{bmatrix} 1 & 0 \\ 0 & C \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & (1 + A^2)^{-1} \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right)^{-1} \tag{3.16}$$

$$= \left( \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right)^{-1} - \begin{bmatrix} 0 & 0 \\ 0 & A(1 + A^2)^{-1} \end{bmatrix}. \tag{3.17}$$

Clearly

$$\begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & A(1 + A^2)^{-1} \end{bmatrix} \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & A \end{bmatrix},$$

and so on suitably pre- and post-multiplying equation (3.16), we obtain equation (3.11).

To prove equation (3.12), check first that

$$\left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) = \\ \left( z_P \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right)$$

as operators on  $\mathcal{N} \oplus \mathcal{D}(A)$ . It follows that

$$\left( \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right)^{-1} = \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \left( z_P \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)$$

as operators from  $\mathcal{H}$  to  $\mathcal{N} \oplus \mathcal{D}(A)$ . On combining this relation with equation (3.11) we derive the expression (3.12) for  $M(z)|_{\mathcal{N} \oplus \mathcal{D}(A)}$ .

We now derive the identity (3.13). Let

$$D = \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix}$$

and consider  $z, w \in \Pi^n$ . By equation (3.10)

$$M(z) = D^*W(z)D \tag{3.18}$$

on  $\mathcal{N} \oplus \mathcal{D}(A)$ , where

$$W(z) = R(z)S(z)^{-1} - \begin{bmatrix} 0 & 0 \\ 0 & A(1 + A^2)^{-1} \end{bmatrix} \tag{3.19}$$

and

$$R(z) = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \quad S(z) = \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P.$$

We have seen that  $S(z)$  is invertible for any  $z \in \Pi^n$ , so that  $W(z)$  is a bounded operator on  $\mathcal{H}$ . Clearly

$$\begin{aligned} M(z) - M(w)^* &= D^* (R(z)S(z)^{-1} - S(w)^{* -1}R(w)^*) D \\ &= D^* S(w)^{* -1} (S(w)^*R(z) - R(w)^*S(z)) S(z)^{-1} D. \end{aligned}$$

Here

$$\begin{aligned} S(w)^*R(z) - R(w)^*S(z) &= \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} z_P + \begin{bmatrix} 0 & 0 \\ 0 & A \end{bmatrix} - w_P^* \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} - \\ &\quad \left( w_P^* \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right) \\ &= z_P - w_P^*. \end{aligned}$$

Hence

$$M(z) - M(w)^* = D^*S(w)^{*^{-1}}(z_P - w_P^*)S(z)^{-1}D,$$

which is equation (3.13).  $\square$

The next result shows that the matricial resolvent belongs not just to the operator Pick class, but to the smaller *operator Loewner class*.

**Proposition 3.5.** *With the notation of Definition 3.3, there exists an analytic operator-valued function  $F : \Pi^n \rightarrow \mathcal{L}(\mathcal{H})$  such that for all  $z, w \in \Pi^n$ ,*

$$M(z) - M(w)^* = F(w)^*(z - \bar{w})_P F(z) \tag{3.20}$$

on  $\mathcal{H}$ .

**Proof.** The identity (3.13) shows that such a relation holds on  $\mathcal{N} \oplus \mathcal{D}(A)$ ; we must extend it to all of  $\mathcal{H}$ . Write  $P_j$  as an operator matrix with respect to the decomposition  $\mathcal{H} = \mathcal{N} \oplus \mathcal{M}$ , as in equation (3.3). Then  $z_P$  has the matricial expression (3.4). For  $z \in \Pi^n$  let

$$F^\sharp(z) = \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} z_P \right)^{-1} \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix}.$$

Then  $F^\sharp(z)$  is an operator from  $\mathcal{N} \oplus \mathcal{D}(A)$  to  $\mathcal{H}$ , and we find that

$$\begin{aligned} F^\sharp(z) &= \begin{bmatrix} 1 & 0 \\ -z_{B^*} & A - z_Y \end{bmatrix}^{-1} \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix} \\ &= \begin{bmatrix} i & 0 \\ i(A - z_Y)^{-1}z_{B^*} & (A - z_Y)^{-1}(1 + iA) \end{bmatrix} : \mathcal{N} \oplus \mathcal{D}(A) \rightarrow \mathcal{H}. \end{aligned}$$

Let

$$F(z) = \begin{bmatrix} i & 0 \\ i(A - z_Y)^{-1}z_{B^*} & i + (A - z_Y)^{-1}(1 + iz_Y) \end{bmatrix} : \mathcal{N} \oplus \mathcal{M} \rightarrow \mathcal{H}. \tag{3.21}$$

Since

$$(A - z_Y)^{-1}(1 + iA) = i + (A - z_Y)^{-1}(1 + iz_Y)$$

on  $\mathcal{N} \oplus \mathcal{D}(A)$  and the right hand side of the last equation is a bounded operator on all of  $\mathcal{H}$ , it is clear that, for every  $z \in \Pi^n$ ,  $F(z)$  is a continuous extension to  $\mathcal{H}$  of  $F^\sharp(z)$  and is a bounded operator. Furthermore  $F$  is analytic on  $\Pi^n$ .

By Proposition 3.4, equation (3.13), the relation (3.20) holds on the dense subspace  $\mathcal{N} \oplus \mathcal{D}(A)$  of  $\mathcal{H}$  for every  $z, w \in \Pi^n$ . Since the operators on both sides of equation (3.20) are continuous on  $\mathcal{H}$ , the equation holds throughout  $\mathcal{H}$ .  $\square$

**Corollary 3.6.** *A matricial resolvent has a non-negative imaginary part at every point of  $\Pi^n$ .*

**Proof.** In the notation of [Proposition 3.5](#), on choosing  $w = z$  in equation [\(3.20\)](#) and dividing by  $2i$  we obtain the relation

$$\operatorname{Im} M(z) = F(z)^*(\operatorname{Im} z_P)F(z)$$

on  $\mathcal{H}$ . We have

$$\operatorname{Im} z_P = \sum_j (\operatorname{Im} z_j)P_j \geq 0,$$

and so  $\operatorname{Im} M(z) \geq 0$  on  $\mathcal{H}$  for all  $z \in \Pi^n$ .  $\square$

Here is a concrete example of a matricial resolvent.

**Example 3.7.** The function

$$M(z) = \frac{1}{z_1 + z_2} \begin{bmatrix} 2z_1z_2 & i(z_1 - z_2) \\ -i(z_1 - z_2) & -2 \end{bmatrix} \tag{3.22}$$

is the matricial resolvent corresponding to

$$\mathcal{H} = \mathbb{C}^2, \quad \mathcal{N} = \mathcal{M} = \mathbb{C}, \quad A = 0 \text{ on } \mathbb{C}, \quad P_1 = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, \quad P_2 = 1 - P_1.$$

#### 4. Nevanlinna representations of type 4

In this section we derive a multivariable analog of the most general form of Nevanlinna representation for functions in the one-variable Pick class ([Theorem 1.2](#)). We start with a multivariable Herglotz theorem [[2](#), [Theorem 1.8](#)]. We shall say (following G. Herglotz [[12](#)]) that an analytic operator-valued function  $F$  on  $\mathbb{D}^n$  is a *Herglotz function* if  $\operatorname{Re} F(\lambda) \geq 0$  for all  $\lambda \in \mathbb{D}^n$ . For present purposes we need the following modification of the notion.

**Definition 4.1.** An analytic function  $F : \mathbb{D}^n \rightarrow \mathcal{L}(\mathcal{K})$ , where  $\mathcal{K}$  is a Hilbert space, is a *strong Herglotz function* if, for every commuting  $n$ -tuple  $T = (T_1, \dots, T_n)$  of operators on a Hilbert space and for  $0 \leq r < 1$ ,  $\operatorname{Re} F(rT) \geq 0$ .

In [[2](#)] these functions were called  $\mathcal{F}_n$ -Herglotz functions. The class of strong Herglotz functions has also been called the *Herglotz–Agler class* (for example [[15,7](#)]). It is clear that every strong Herglotz function is a Herglotz function, and in the cases  $n = 1$  and  $2$  the converse is also true [[2](#)].

**Theorem 4.2.** *Let  $\mathcal{K}$  be a Hilbert space and let  $F : \mathbb{D}^2 \rightarrow \mathcal{L}(\mathcal{K})$  be a strong Herglotz function such that  $F(0) = 1$ . There exist a Hilbert space  $\mathcal{H}$ , an orthogonal decomposition  $P$  of  $\mathcal{H}$ , an isometric linear operator  $V : \mathcal{K} \rightarrow \mathcal{H}$  and a unitary operator  $U$  on  $\mathcal{H}$  such that, for all  $\lambda \in \mathbb{D}^n$ ,*

$$F(\lambda) = V^* \frac{1 + U\lambda_P}{1 - U\lambda_P} V. \tag{4.1}$$

*Conversely, every function  $F : \mathbb{D}^n \rightarrow \mathcal{L}(\mathcal{K})$  expressible in the form (4.1) for some  $\mathcal{H}$ ,  $P$ ,  $V$  and  $U$  with the stated properties is a strong Herglotz function and satisfies  $F(0) = 1$ .*

Note that  $\lambda_P = \sum_j \lambda_j P_j$  has operator norm at most  $\|\lambda\|_\infty < 1$  for  $\lambda \in \mathbb{D}^n$ , and hence equation (4.1) does define  $F$  as an analytic operator-valued function on  $\mathbb{D}^n$ .

On specializing to scalar-valued functions in the  $n$ -variable Herglotz class we obtain the following consequence.

**Corollary 4.3.** *Let  $f$  be a scalar-valued strong Herglotz function on  $\mathbb{D}^n$ . There exists a Hilbert space  $\mathcal{H}$ , a unitary operator  $L$  on  $\mathcal{H}$ , an orthogonal decomposition  $P$  of  $\mathcal{H}$ , a real number  $a$  and a vector  $v \in \mathcal{H}$  such that, for all  $\lambda \in \mathbb{D}^n$ ,*

$$f(\lambda) = -ia + \langle (L - \lambda_P)^{-1} (L + \lambda_P)v, v \rangle. \tag{4.2}$$

*Conversely, for any  $\mathcal{H}$ ,  $L$ ,  $P$ ,  $a$  and  $v$  with the properties described, equation (4.2) defines  $f$  as an  $n$ -variable strong Herglotz function.*

Again, the right hand side of equation (4.2) is an analytic function of  $\lambda \in \mathbb{D}^n$  since

$$(L - \lambda_P)^{-1} = L^{-1} (1 - \lambda_P L^{-1})^{-1}$$

is a bounded operator and is analytic in  $\lambda$ .

**Definition 4.4.** A *Nevanlinna representation of type 4* of a function  $h : \Pi^n \rightarrow \mathbb{C}$  consists of an orthogonally decomposed Hilbert space  $\mathcal{H} = \mathcal{N} \oplus \mathcal{M}$ , a self-adjoint densely defined operator  $A$  on  $\mathcal{M}$ , an orthogonal decomposition  $P$  of  $\mathcal{H}$ , a real number  $a$  and a vector  $v \in \mathcal{H}$  such that

$$h(z) = a + \langle M(z)v, v \rangle \tag{4.3}$$

for all  $z \in \Pi^n$ , where  $M(z)$  is the structured resolvent of  $A$  of type 4 corresponding to  $P$  (given by the formula (3.1)).

We wish to convert Corollary 4.3 to a representation theorem for suitable analytic functions on  $\Pi^n$ . The fact that the corollary only applies to *strong* Herglotz functions

results in representation theorems for a subclass of the Pick class  $\mathcal{P}_n$ . Recall from the introduction:

**Definition 4.5.** The *Loewner class*  $\mathcal{L}_n$  is the set of analytic functions  $h$  on  $\Pi^n$  with the property that there exist  $n$  positive semi-definite functions  $A_1, \dots, A_n$  on  $\Pi^n \times \Pi^n$ , analytic in the first argument, such that

$$h(z) - \overline{h(w)} = \sum_{j=1}^n (z_j - \overline{w_j}) A_j(z, w)$$

for all  $z, w \in \Pi^n$ .

A function  $h$  on  $\Pi^n$  belongs to  $\mathcal{L}_n$  if and only if it corresponds under conjugation by the Cayley transform to a function in the Schur–Agler class of the polydisc [4, Lemma 2.13]. Another characterization:  $h \in \mathcal{L}_n$  if and only if, for every commuting  $n$ -tuple  $T$  of bounded operators with strictly positive imaginary parts,  $h(T)$  has positive imaginary part.

We can now prove Theorem 1.9 from the introduction: a function  $h$  defined on  $\Pi^n$  has a Nevanlinna representation of type 4 if and only if  $h \in \mathcal{L}_n$ .

**Proof of Theorem 1.9.** Let  $h \in \mathcal{L}_n$ . Define an  $n$ -variable Herglotz function  $f : \mathbb{D}^n \rightarrow \mathbb{C}$  by

$$f(\lambda) = -ih(\lambda) \tag{4.4}$$

where

$$z_j = i \frac{1 + \lambda_j}{1 - \lambda_j} \quad \text{for } j = 1, \dots, n. \tag{4.5}$$

When  $\lambda \in \mathbb{D}^n$  the point  $z$  belongs to  $\Pi^n$ , and so  $f(\lambda)$  is well defined, and since  $\text{Im } h(z) \geq 0$  we have  $\text{Re } f(\lambda) \geq 0$ , so that  $f$  is indeed a Herglotz function. In fact  $f$  is even a strong Herglotz function: since  $h \in \mathcal{L}_n$ , the function  $\varphi \in \mathcal{S}_n$  corresponding to  $h$  lies in the Schur–Agler class of the polydisc, and so  $f = (1 + \varphi)/(1 - \varphi)$  is a strong Herglotz function.

By Corollary 4.3 there exist a real number  $a$ , a Hilbert space  $\mathcal{H}$ , a vector  $v \in \mathcal{H}$ , a unitary operator  $L$  on  $\mathcal{H}$  and an orthogonal decomposition  $P$  on  $\mathcal{H}$  such that, for all  $z \in \Pi^n$ ,

$$\begin{aligned} h(z) &= if(\lambda) = a + \langle i(L - \lambda)^{-1}(L + \lambda)v, v \rangle \\ &= a + \langle i[L - (z - i)(z + i)^{-1}]^{-1}[L + (z - i)(z + i)^{-1}]v, v \rangle. \end{aligned} \tag{4.6}$$

Here and in the rest of this section  $z, \lambda$  are identified with the operators  $z_P, \lambda_P$  on  $\mathcal{H}$ , and in consequence the relation

$$\lambda = \frac{z - i}{z + i}$$

is meaningful and valid.

For  $z \in \Pi^n$  let

$$M(z) = i(L - \lambda)^{-1}(L + \lambda) = i \left( L - \frac{z - i}{z + i} \right)^{-1} \left( L + \frac{z - i}{z + i} \right). \tag{4.7}$$

Since  $L$  is unitary on  $\mathcal{H}$  and  $\lambda \in \mathbb{D}^n$ , the operator  $M(z)$  is bounded on  $\mathcal{H}$  for every  $z \in \Pi^n$  and, by equation (4.6), we have

$$h(z) = a + \langle M(z)v, v \rangle \tag{4.8}$$

for all  $z \in \Pi^2$ . Theorem 1.9 will follow provided we can show that  $M(z)$  is given by equation (3.1) for a suitable self-adjoint operator  $A$ .

Observe that

$$\begin{aligned} M(z) &= i((z + i)L - (z - i))^{-1}((z + i)L + (z - i)) \\ &= i(z(L - 1) + i(L + 1))^{-1}(z(L + 1) + i(L - 1)). \end{aligned} \tag{4.9}$$

We wish to take out a factor  $1 - L$  from both factors in equation (4.9), but this may be impossible since  $1 - L$  can have a nonzero kernel. Accordingly we decompose  $\mathcal{H}$  into  $\mathcal{N} \oplus \mathcal{M}$  where  $\mathcal{N} = \ker(1 - L)$ ,  $\mathcal{M} = \mathcal{N}^\perp$ . With respect to this decomposition we can write  $L$  as an operator matrix

$$L = \begin{bmatrix} 1 & 0 \\ 0 & L_0 \end{bmatrix},$$

where  $L_0$  is unitary and  $\ker(1 - L_0) = \{0\}$ . Substituting into equation (4.9) we have

$$\begin{aligned} M(z) &= i \left( z \begin{bmatrix} 0 & 0 \\ 0 & L_0 - 1 \end{bmatrix} + i \begin{bmatrix} 2 & 0 \\ 0 & L_0 + 1 \end{bmatrix} \right)^{-1} \left( z \begin{bmatrix} 2 & 0 \\ 0 & L_0 + 1 \end{bmatrix} + i \begin{bmatrix} 0 & 0 \\ 0 & L_0 - 1 \end{bmatrix} z \right) \\ &= \left( -z \begin{bmatrix} 0 & 0 \\ 0 & 1 - L_0 \end{bmatrix} + \begin{bmatrix} 2i & 0 \\ 0 & i(1 + L_0) \end{bmatrix} \right)^{-1} \left( z \begin{bmatrix} 2i & 0 \\ 0 & i(1 + L_0) \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 - L_0 \end{bmatrix} \right) \end{aligned} \tag{4.10}$$

Formally we may now write

$$\begin{aligned} M(z) &= \begin{bmatrix} -\frac{1}{2}i & \\ 0 & (1 - L_0)^{-1} \end{bmatrix} \left( -z \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & i\frac{1+L_0}{1-L_0} \end{bmatrix} \right)^{-1} \times \\ &\quad \left( z \begin{bmatrix} 1 & 0 \\ 0 & i\frac{1+L_0}{1-L_0} \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} 2i & 0 \\ 0 & 1 - L_0 \end{bmatrix}, \end{aligned} \tag{4.11}$$

but whereas equation (4.10) is a relation between bounded operators defined on all of  $\mathcal{H}$ , equation (4.11) involves unbounded, partially defined operators and we must verify that the product of operators on the right hand side is meaningful.

Let

$$A = i \frac{1 + L_0}{1 - L_0}.$$

Since  $L_0$  is unitary on  $\mathcal{M}$  and  $\ker(1 - L_0) = \{0\}$ , the operator  $A$  is self-adjoint and densely defined on  $\mathcal{M}$  [21, Section 121]. The domain  $\mathcal{D}(A)$  of  $A$  is the dense subspace  $\text{ran}(1 - L_0)$  of  $\mathcal{M}$ . It follows from the definition of  $A$  that

$$(1 - L_0)^{-1} = \frac{1}{2}(1 - iA), \tag{4.12}$$

which is an equation between bijective operators from  $\mathcal{D}(A)$  to  $\mathcal{M}$ . Likewise

$$1 + L_0 = -2iA(1 - iA)^{-1} : \mathcal{M} \rightarrow \mathcal{D}(A) \tag{4.13}$$

are bounded operators.

Let us continue the calculation from the first factor on the right hand side of equation (4.10). Since  $\ker(1 - L_0) = \{0\}$ , the right hand side of the relation

$$-z \begin{bmatrix} 0 & 0 \\ 0 & 1 - L_0 \end{bmatrix} + \begin{bmatrix} 2i & 0 \\ 0 & i(1 + L_0) \end{bmatrix} = \left( -z \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} \right) \begin{bmatrix} 2i & 0 \\ 0 & 1 - L_0 \end{bmatrix}$$

comprises a bijective map from  $\mathcal{H}$  to  $\mathcal{N} \oplus \mathcal{D}(A)$  followed by a bijection from  $\mathcal{N} \oplus \mathcal{D}(A)$  to  $\mathcal{H}$  (recall the equation (3.7)). We may therefore take inverses in the equation to obtain

$$\begin{aligned} & \left( -z \begin{bmatrix} 0 & 0 \\ 0 & 1 - L_0 \end{bmatrix} + \begin{bmatrix} 2i & 0 \\ 0 & i(1 + L_0) \end{bmatrix} \right)^{-1} \\ &= \begin{bmatrix} -\frac{1}{2}i & 0 \\ 0 & (1 - L_0)^{-1} \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \\ &= \begin{bmatrix} -\frac{1}{2}i & 0 \\ 0 & \frac{1}{2}(1 - iA) \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \end{aligned} \tag{4.14}$$

as operators on  $\mathcal{N} \oplus \mathcal{D}(A)$ .

Similar reasoning applies to the equation

$$\begin{aligned} & z \begin{bmatrix} 2i & 0 \\ 0 & i(1 + L_0) \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 - L_0 \end{bmatrix} \\ &= \left( z \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} 2i & 0 \\ 0 & 1 - L_0 \end{bmatrix} \\ &= \left( z \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} -\frac{1}{2}i & 0 \\ 0 & \frac{1}{2}(1 - iA) \end{bmatrix}^{-1}; \end{aligned} \tag{4.15}$$

it is valid as an equation between operators on  $\mathcal{H}$ . The right hand side comprises an operator from  $\mathcal{H}$  to  $\mathcal{N} \oplus \mathcal{D}(A)$  followed by an operator from  $\mathcal{N} \oplus \mathcal{D}(A)$  to  $\mathcal{H}$ , and so both sides of the equation denote an operator on  $\mathcal{H}$ .

On combining equations (4.10), (4.14) and (4.15) we obtain

$$M(z) = \begin{bmatrix} -\frac{1}{2}i & 0 \\ 0 & \frac{1}{2}(1 - iA) \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \\ \times \left( z \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} -\frac{1}{2}i & 0 \\ 0 & \frac{1}{2}(1 - iA) \end{bmatrix}^{-1}.$$

Pre-multiply this equation by 2 and post-multiply by  $\frac{1}{2}$  to deduce that  $M(z)$  is indeed the structured resolvent of  $A$  of type 4 corresponding to  $P$ , as defined in equation (3.1). Thus the formula (4.8) is a Nevanlinna representation of  $h$  of type 4.

Conversely, let  $h \in \mathcal{L}_n$  have a type 4 representation (4.3). By Proposition 3.5 there exists an analytic operator-valued function  $F : \Pi^n \rightarrow \mathcal{L}(\mathcal{H})$  such that, for all  $z, w \in \Pi^n$ ,

$$M(z) - M(w)^* = F(w)^*(z - \bar{w})_P F(z) \tag{4.16}$$

on  $\mathcal{H}$ . Hence

$$h(z) - \overline{h(w)} = \langle (M(z) - M(w)^*)v, v \rangle \\ = \langle F(w)^*(z - \bar{w})_P F(z)v, v \rangle \\ = \sum_{j=1}^n (z_j - \bar{w}_j) A_j(z, w)$$

for all  $z, w \in \Pi^n$ , where

$$A_j(z, w) = \langle P_j F(z)v, F(w)v \rangle.$$

The  $A_j$  are clearly positive semidefinite on  $\Pi^n$ , and hence  $h$  belongs to the Loewner class  $\mathcal{L}_n$ .  $\square$

### 5. Nevanlinna representations of types 3, 2 and 1

Nevanlinna representations of type 4 have the virtue of being general for functions in  $\mathcal{L}_n$ , but they are undeniably cumbersome. In this section we shall show that there are three simpler representation formulae, corresponding to increasingly stringent growth conditions on  $h \in \mathcal{L}_n$ .

In Nevanlinna’s one-variable representation formula of Theorem 1.2,

$$h(z) = a + bz + \int \frac{1 + tz}{t - z} d\mu(t), \tag{5.1}$$

it may be the case for a particular  $h \in \mathcal{P}$  that the  $bz$  term is absent. The analogous situation in two variables is that the space  $\mathcal{N}$  in a type 4 representation may be zero. Equivalently, in the corresponding Herglotz representation, the unitary operator  $L$  does not have 1 as an eigenvalue. This suggests the following notion.

**Definition 5.1.** A *Nevanlinna representation of type 3* of a function  $h$  on  $\Pi^n$  consists of a Hilbert space  $\mathcal{H}$ , a self-adjoint densely defined operator  $A$  on  $\mathcal{H}$ , a positive decomposition  $Y$  of  $\mathcal{H}$ , a real number  $a$  and a vector  $v \in \mathcal{H}$  such that, for all  $z \in \Pi^n$ ,

$$h(z) = a + \langle (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 - iA)^{-1}v, v \rangle. \quad (5.2)$$

Thus  $h$  has a type 3 representation if  $h(z) = a + \langle M(z)v, v \rangle$  where  $M(z)$  is the structured resolvent of  $A$  of type 3 corresponding to  $Y$ , as given by equation (2.3).

In [5] the authors derived a somewhat simpler representation which can also be regarded as an analog of the case  $b = 0$  of Nevanlinna's one-variable formula (5.1).

**Definition 5.2.** A *Nevanlinna representation of type 2* of a function  $h$  on  $\Pi^n$  consists of a Hilbert space  $\mathcal{H}$ , a self-adjoint densely defined operator  $A$  on  $\mathcal{H}$ , a positive decomposition  $Y$  of  $\mathcal{H}$ , a real number  $a$  and a vector  $\alpha \in \mathcal{H}$  such that, for all  $z \in \Pi^n$

$$h(z) = a + \langle (A - z_Y)^{-1}\alpha, \alpha \rangle. \quad (5.3)$$

This means of course that, for all  $z \in \Pi^n$ ,

$$h(z) = a + \langle M(z)\alpha, \alpha \rangle$$

where  $M(z)$  is the structured resolvent of  $A$  of type 2 corresponding to  $Y$  (compare equation (2.1)).

We wish to understand the relationship between type 3 and type 2 representations.

**Proposition 5.3.** *If  $h \in \mathcal{P}_n$  has a type 2 representation then  $h$  has a type 3 representation. Conversely, if  $h \in \mathcal{P}_n$  has a type 3 representation as in equation (5.2) with the additional property that  $v \in \mathcal{D}(A)$  then  $h$  has a type 2 representation.*

**Proof.** Suppose that  $h \in \mathcal{P}_n$  has the type 2 representation

$$h(z) = a_0 + \langle (A - z_Y)^{-1}\alpha, \alpha \rangle$$

for some  $a_0 \in \mathbb{R}$ , positive decomposition  $Y$  and  $\alpha \in \mathcal{H}$ . We must show that  $h$  has a representation of the form (5.2) for some  $a \in \mathbb{R}$  and  $v \in \mathcal{H}$ . By Proposition 2.6, it suffices to find  $a \in \mathbb{R}$  and  $v \in \mathcal{D}(A)$  such that

$$h(z) = a + \langle (1 - iA) \{ (A - z_Y)^{-1} - A(1 + A^2)^{-1} \} (1 + iA)v, v \rangle$$

for all  $z \in \Pi^n$ .

To this end, let  $C = A(1 + A^2)^{-1}$  and let

$$a = a_0 + \langle C\alpha, \alpha \rangle. \tag{5.4}$$

Since  $1 + iA$  is invertible on  $\mathcal{H}$  and  $\text{ran}(1 + iA)^{-1} \subset \mathcal{D}(A)$  we may define

$$v = (1 + iA)^{-1}\alpha \in \mathcal{D}(A). \tag{5.5}$$

Then

$$\begin{aligned} h(z) &= a_0 + \langle (A - z_Y)^{-1}\alpha, \alpha \rangle \\ &= a - \langle C\alpha, \alpha \rangle + \langle (A - z_Y)^{-1}\alpha, \alpha \rangle \\ &= a + \langle \{(A - z_Y)^{-1} - C\} (1 + iA)v, (1 + iA)v \rangle \\ &= a + \langle (1 - iA) \{(A - z_Y)^{-1} - C\} (1 + iA)v, v \rangle \end{aligned}$$

as required. Thus  $h$  has a type 3 representation.

Conversely, let  $h$  have a type 3 representation (5.2) such that  $v \in \mathcal{D}(A)$ , that is

$$h(z) = a + \langle M(z)v, v \rangle$$

where  $a \in \mathbb{R}$  and  $M$  is the structured resolvent of  $A$  of type 3 corresponding to  $Y$ , as in equation (2.3). Since  $v \in \mathcal{D}(A)$  we may define the vector  $\alpha \stackrel{\text{def}}{=} (1 + iA)v \in \mathcal{H}$ , and furthermore, by Proposition 2.6,

$$\begin{aligned} h(z) &= a + \langle (1 - iA) \{(A - z_Y)^{-1} - C\} (1 + iA)v, v \rangle \\ &= a + \langle \{(A - z_Y)^{-1} - C\} \alpha, \alpha \rangle \\ &= a - \langle C\alpha, \alpha \rangle + \langle (A - z_Y)^{-1}\alpha, \alpha \rangle \\ &= a_0 + \langle (A - z_Y)^{-1}\alpha, \alpha \rangle, \end{aligned}$$

where  $a_0 \in \mathbb{R}$  is given by equation (5.4). Thus  $h$  has a representation of type 2.  $\square$

A special case of a type 2 representation occurs when the constant term  $a$  in equation (5.3) is 0. In one variable, this corresponds to Nevanlinna’s characterization of the Cauchy transforms of positive finite measures on  $\mathbb{R}$ . Accordingly we define a *type 1 representation* of  $h \in \mathcal{L}_n$  to be the special case of a type 2 representation of  $h$  in which  $a = 0$  in equation (5.3).

**Definition 5.4.** An analytic function  $h$  on  $\Pi^n$  has a *Nevanlinna representation of type 1* if there exist a Hilbert space  $\mathcal{H}$ , a densely defined self-adjoint operator  $A$  on  $\mathcal{H}$ , a positive decomposition  $Y$  of  $\mathcal{H}$  and a vector  $\alpha \in \mathcal{H}$  such that, for all  $z \in \Pi^n$ ,

$$h(z) = \langle (A - z_Y)^{-1}\alpha, \alpha \rangle. \tag{5.6}$$

A representation of type 1 is obviously a representation of type 2. The following proposition is an immediate corollary of [Proposition 5.3](#).

**Proposition 5.5.** *A function  $h \in \mathcal{L}_n$  has a type 1 representation if and only if  $h$  has a type 3 representation as in equation (5.2) with the additional properties that  $v \in \mathcal{D}(A)$  and*

$$a - \langle A(1 + A^2)^{-1}\alpha, \alpha \rangle = 0.$$

For consistency with our earlier terminology for structured resolvents and representations we should have to define a structured resolvent of type 1 to be the same as a structured resolvent of type 2. We refrain from making such a confusing definition.

We conclude this section by giving examples of the four types of Nevanlinna representation in two variables.

**Example 5.6.** (1) The formula

$$h(z) = -\frac{1}{z_1 + z_2} = \langle (0 - z_Y)^{-1}v, v \rangle_{\mathbb{C}},$$

where  $Y = (\frac{1}{2}, \frac{1}{2})$  and  $v = 1/\sqrt{2}$ , exhibits a representation of type 1, with  $A = 0$ .

(2) Likewise

$$h(z) = 1 - \frac{1}{z_1 + z_2} = 1 + \langle (0 - z_Y)^{-1}v, v \rangle_{\mathbb{C}}$$

is a representation of type 2.

(3) Let

$$h(z) = \begin{cases} \frac{1}{1 + z_1 z_2} \left( z_1 - z_2 + \frac{iz_2(1 + z_1^2)}{\sqrt{z_1 z_2}} \right) & \text{if } z_1 z_2 \neq -1 \\ \frac{1}{2}(z_1 + z_2) & \text{if } z_1 z_2 = -1 \end{cases} \tag{5.7}$$

where we take the branch of the square root that is analytic in  $\mathbb{C} \setminus [0, \infty)$  with range  $\Pi$ . We claim that  $h \in \mathcal{P}_2$  and that  $h$  has the type 3 representation

$$h(z) = \langle M(z)v, v \rangle_{L^2(\mathbb{R})}, \tag{5.8}$$

where  $M(z)$  is the structured resolvent of type 3 given in [Example 2.11](#) and  $v(t) = 1/\sqrt{\pi(1 + t^2)}$ . To see this, let  $h$  be temporarily defined by equation (5.8). Since  $v$  is an even function in  $L^2(\mathbb{R})$ , equation (2.12) tells us that

$$h(z) = \int_{-\infty}^{\infty} \frac{t(1 + z_1 z_2) + (1 - it)(it z_1 + z_2)}{\pi(t^2 - z_1 z_2)(1 + t^2)} dt.$$

Since the denominator is an even function of  $t$ , the integrals of all the odd powers of  $t$  in the numerator vanish, and we have, provided  $z_1 z_2 \neq -1$ ,

$$\begin{aligned} h(z) &= \frac{2}{\pi} \int_0^{\infty} \frac{z_2 + t^2 z_1}{(t^2 - z_1 z_2)(1 + t^2)} dt \\ &= \frac{2}{\pi} \int_0^{\infty} \frac{z_2(1 + z_1^2)}{1 + z_1 z_2} \frac{1}{t^2 - z_1 z_2} + \frac{z_1 - z_2}{1 + z_1 z_2} \frac{1}{1 + t^2} dt. \end{aligned}$$

Now, for  $w \in \Pi$ ,

$$\int_0^{\infty} \frac{dt}{t^2 - w^2} = \frac{i\pi}{2w},$$

and so we find that  $h$  is indeed given by equation (5.7) in the case that  $z_1 z_2 \neq -1$ . When  $z_1 z_2 = -1$  we have

$$\begin{aligned} h(z) &= \frac{2}{\pi} \int_0^{\infty} \frac{z_2 + z_1 t^2}{(1 + t^2)^2} dt \\ &= \frac{2}{\pi} \int_0^{\infty} \frac{z_1}{1 + t^2} + \frac{z_2 - z_1}{(1 + t^2)^2} dt \\ &= \frac{1}{2}(z_1 + z_2). \end{aligned}$$

Thus equation (5.8) is a type 3 representation of the function  $h$  given by equation (5.7). This function is *constant* and equal to  $i$  on the diagonal  $z_1 = z_2$ .

(4) The function

$$h(z) = \frac{z_1 z_2}{z_1 + z_2} = - \left( -\frac{1}{z_1} - \frac{1}{z_2} \right)^{-1}$$

clearly belongs to  $\mathcal{P}_2$ . It has the representation of type 4

$$h(z) = \langle M(z)v, v \rangle_{\mathbb{C}^2}$$

where  $M(z)$  is the matricial resolvent given in Example 3.7 and

$$v = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

We claim that each of the above representations is of the simplest available type for the function in question; for example, the function  $h$  in part (4) does not have a Nevanlinna representation of type 3. To prove this claim (which we shall do in [Example 8.2](#) below) we need characterizations of the types of functions – the subject of the next two sections.

## 6. Asymptotic behavior and types of representations

In this section we shall give function-theoretic conditions for a function in  $\mathcal{L}_n$  to have a representation of a given type. These conditions will be in terms of the asymptotic behavior of the function at  $\infty$ .

Every function in  $\mathcal{L}_n$  has a type 4 representation, by [Theorem 1.9](#). Let us characterize the functions that possess a type 3 representation. We denote by  $\chi$  the vector  $(1, \dots, 1)$  of ones in  $\mathbb{C}^n$ . The following statement contains [Theorem 1.8](#).

**Theorem 6.1.** *The following three conditions are equivalent for a function  $h \in \mathcal{L}_n$ .*

- (1) *The function  $h$  has a Nevanlinna representation of type 3;*
- (2)

$$\liminf_{s \rightarrow \infty} \frac{1}{s} \operatorname{Im} h(is\chi) = 0; \tag{6.1}$$

- (3)

$$\lim_{s \rightarrow \infty} \frac{1}{s} \operatorname{Im} h(is\chi) = 0. \tag{6.2}$$

**Proof.** (1) $\Rightarrow$ (3) Suppose that  $h$  has a Nevanlinna representation of type 3:

$$h(z) = a + \langle (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 - iA)^{-1}v, v \rangle \tag{6.3}$$

for suitable  $a \in \mathbb{R}$ ,  $\mathcal{H}$ ,  $A$ ,  $Y$  and  $v \in \mathcal{H}$ . Since

$$(is\chi)_Y = \sum_j isY_j = is$$

we have

$$h(is\chi) = a + \langle (1 - iA)(A - is)^{-1}(1 + isA)(1 - iA)^{-1}v, v \rangle.$$

Let  $\nu$  be the scalar spectral measure for  $A$  corresponding to the vector  $v \in \mathcal{H}$ . By the Spectral Theorem

$$\begin{aligned}
 h(is\chi) &= a + \int (1 - it)(t - is)^{-1}(1 + ist)(1 - it)^{-1} \, d\nu(t) \\
 &= a + \int \frac{1 + ist}{t - is} \, d\nu(t).
 \end{aligned}$$

Since

$$\operatorname{Im} \frac{1 + ist}{t - is} = \frac{s(1 + t^2)}{s^2 + t^2},$$

we have

$$\frac{1}{s} \operatorname{Im} h(is\chi) = \int \frac{1 + t^2}{s^2 + t^2} \, d\nu(t).$$

The integrand decreases monotonically to 0 as  $s \rightarrow \infty$  and so, by the Monotone Convergence Theorem, equation (6.2) holds.

(3) $\Rightarrow$ (2) is trivial.

(2) $\Rightarrow$ (1) Now suppose that  $h \in \mathcal{L}_n$  and

$$\liminf_{s \rightarrow \infty} \frac{1}{s} \operatorname{Im} h(is\chi) = 0.$$

By Theorem 1.9,  $h$  has a Nevanlinna representation of type 4: that is, there exist  $a, \mathcal{H}, \mathcal{N} \subset \mathcal{H}$ , operators  $A, Y$  on  $\mathcal{N}^\perp$  and a vector  $v \in \mathcal{H}$  with the properties described in Definition 5.1 such that

$$h(z) = a + \langle M(z)v, v \rangle$$

for all  $z \in \Pi^n$ , where

$$\begin{aligned}
 M(z) &= \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)^{-1} \\
 &\quad \times \left( z_P \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix}^{-1}.
 \end{aligned} \tag{6.4}$$

Thus, for  $s > 0$ , since once again  $(is\chi)_P = is$ ,

$$\begin{aligned}
 M(is\chi) &= \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & (A - is)^{-1} \end{bmatrix} \begin{bmatrix} is & 0 \\ 0 & 1 + isA \end{bmatrix} \begin{bmatrix} i & 0 \\ 0 & (1 - iA)^{-1} \end{bmatrix} \\
 &= \begin{bmatrix} is & 0 \\ 0 & (1 - iA)(A - is)^{-1}(1 + isA)(1 - iA)^{-1} \end{bmatrix}.
 \end{aligned}$$

Let the projections of  $v$  onto  $\mathcal{N}, \mathcal{N}^\perp$  be  $v_1, v_2$  respectively. Then

$$\begin{aligned} h(is\chi) &= a + \langle M(is\chi)v, v \rangle \\ &= a + is \|v_1\|^2 + \langle (1 - iA)(A - is)^{-1}(1 + isA)(1 - iA)^{-1}v_2, v_2 \rangle \end{aligned}$$

and therefore

$$\begin{aligned} \frac{1}{s} \operatorname{Im} h(is\chi) &= \|v_1\|^2 + \frac{1}{s} \operatorname{Im} \langle (1 - iA)(A - is)^{-1}(1 + isA)(1 - iA)^{-1}v_2, v_2 \rangle \\ &\geq \|v_1\|^2 \end{aligned}$$

by [Corollary 2.7](#). Hence

$$\begin{aligned} 0 &= \liminf_{s \rightarrow \infty} \frac{1}{s} \operatorname{Im} h(is\chi) \\ &\geq \|v_1\|^2. \end{aligned}$$

It follows that  $v_1 = 0$ .

Let the compression of the projection  $P_j$  to  $\mathcal{N}^\perp$  be  $Y_j$ : then  $Y = (Y_1, \dots, Y_n)$  is a positive decomposition of  $\mathcal{N}^\perp$ , and the compression of  $z_P$  to  $\mathcal{N}^\perp$  is  $z_Y$ . By [Remark 3.2](#) the  $(2, 2)$  block  $M_{22}(z)$  in  $M(z)$  is

$$M_{22}(z) = (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 - iA)^{-1}.$$

Since  $v_1 = 0$  it follows that

$$\begin{aligned} h(z) &= a + \langle M(z)v, v \rangle \\ &= a + \langle M_{22}(z)v_2, v_2 \rangle \\ &= a + \langle (1 - iA)(A - z_Y)^{-1}(1 + z_Y A)(1 - iA)^{-1}v_2, v_2 \rangle, \end{aligned}$$

which is the desired type 3 representation of  $h$ . Hence (2) $\Rightarrow$ (1).  $\square$

In [\[7\]](#) it is shown that condition (3) in the above theorem is also a necessary and sufficient condition that  $-ih$  have a  $\Pi^n$ -impedance-conservative realization.

Type 2 representations were characterized by the following theorem in [\[5\]](#) in the case of two variables. The following result, which contains [Theorem 1.7](#), shows that the result holds generally.

**Theorem 6.2.** *The following three conditions are equivalent for a function  $h \in \mathcal{L}_n$ .*

- (1) *The function  $h$  has a Nevanlinna representation of type 2;*
- (2)

$$\liminf_{s \rightarrow \infty} s \operatorname{Im} h(is\chi) < \infty; \tag{6.5}$$

(3)

$$\lim_{s \rightarrow \infty} s \operatorname{Im} h(is\chi) < \infty. \tag{6.6}$$

**Proof.** (1) $\Rightarrow$ (3) Suppose that  $h$  has the type 2 representation  $h(z) = a + \langle (A - z_Y)^{-1}v, v \rangle$  for a suitable real  $a$ , self-adjoint  $A$ , positive decomposition  $Y$  and vector  $v$ . Let  $\nu$  be the scalar spectral measure for  $A$  corresponding to the vector  $v$ . Then, for  $s > 0$ ,  $A - (is\chi)_Y = A - is$  and so

$$\begin{aligned} s \operatorname{Im} h(is\chi) &= s \operatorname{Im} \int \frac{d\nu(t)}{t - is} \\ &= \int \frac{s^2 d\nu(t)}{t^2 + s^2}. \end{aligned}$$

The integrand is positive and increases monotonically to 1 as  $s \rightarrow \infty$ . Hence, by the Dominated Convergence Theorem

$$\lim_{s \rightarrow \infty} s \operatorname{Im} h(is\chi) = \nu(\mathbb{R}) = \|v\|^2 < \infty.$$

Hence (1) $\Rightarrow$ (3).

(3) $\Rightarrow$ (2) is trivial.

(2) $\Rightarrow$ (1) Suppose (2) holds. *A fortiori*,

$$\liminf_{s \rightarrow \infty} \frac{1}{s} \operatorname{Im} h(is\chi) = 0.$$

By [Theorem 6.1](#)  $h$  has a type 3 representation (6.3) for suitable  $a \in \mathbb{R}$ ,  $\mathcal{H}$ ,  $A$ ,  $Y$  and  $v \in \mathcal{H}$ . Let  $\nu$  be the scalar spectral measure for  $A$  corresponding to the vector  $v$ . Then for  $s > 0$

$$\begin{aligned} s \operatorname{Im} h(is\chi) &= s \operatorname{Im} \int \frac{1 + ist}{t - is} d\nu(t) \\ &= \int \frac{s^2(1 + t^2)}{t^2 + s^2} d\nu(t). \end{aligned}$$

As  $s \rightarrow \infty$  the integrand increases monotonically to  $1 + t^2$ . Condition (2) now implies that

$$\int 1 + t^2 d\nu(t) < \infty.$$

It follows that  $v \in \mathcal{D}(A)$ . Hence, by [Proposition 5.3](#),  $h$  has a representation of type 2.  $\square$

In [5] we proved [Theorem 6.2](#) for  $n = 2$  using a different approach from the present one.

From this theorem the characterization of type 1 representations follows just as in the one-variable case. We obtain a strengthening of [Theorem 1.6](#).

**Theorem 6.3.** *The following three conditions are equivalent for a function  $h \in \mathcal{L}_n$ .*

- (1) *The function  $h$  has a Nevanlinna representation of type 1;*
- (2)

$$\liminf_{s \rightarrow \infty} s |h(is\chi)| < \infty;$$

- (3)

$$\lim_{s \rightarrow \infty} s |h(is\chi)| < \infty. \tag{6.7}$$

**Proof.** We follow Lax’s treatment [\[16\]](#) of the one-variable Nevanlinna theorem.

(1) $\Rightarrow$ (3) Suppose that  $h$  has a type 1 representation as in equation [\(5.6\)](#) for some  $\mathcal{H}$ ,  $A$ ,  $Y$  and  $v$ . Then

$$\begin{aligned} h(is\chi) &= \langle (A - is)^{-1}\alpha, \alpha \rangle \\ &= \langle (A + is)(A^2 + s^2)^{-1}\alpha, \alpha \rangle, \end{aligned}$$

and so

$$\operatorname{Re} sh(is\chi) = \langle sA(A^2 + s^2)^{-1}\alpha, \alpha \rangle, \quad \operatorname{Im} sh(is\chi) = \langle s^2(A^2 + s^2)^{-1}\alpha, \alpha \rangle.$$

Let  $\nu$  be the scalar spectral measure for  $A$  corresponding to the vector  $\alpha \in \mathcal{H}$ . Then

$$\operatorname{Re} sh(is\chi) = \int \frac{st}{t^2 + s^2} d\nu(t), \quad \operatorname{Im} sh(is\chi) = \int \frac{s^2}{t^2 + s^2} d\nu(t).$$

The integrand in the first integral tends pointwise in  $t$  to 0 as  $s \rightarrow \infty$ , and by the inequality of the means it is no greater than  $\frac{1}{2}$ ; thus the integral tends to 0 as  $s \rightarrow \infty$  by the Dominated Convergence Theorem. The integrand in the second integral increases monotonically to 1 as  $s \rightarrow \infty$ . Thus

$$\operatorname{Re} sh(is\chi) \rightarrow 0, \quad \operatorname{Im} sh(is\chi) \rightarrow \|\alpha\|^2 \quad \text{as } s \rightarrow \infty.$$

Hence the inequality [\(6.7\)](#) holds. Thus (1) $\Rightarrow$ (3).

(3) $\Rightarrow$ (2) is trivial.

(2) $\Rightarrow$ (1) Suppose that

$$\liminf_{s \rightarrow \infty} s |h(is\chi)| < \infty. \tag{6.8}$$

As

$$\liminf_{s \rightarrow \infty} s \operatorname{Im} h(is\chi) \leq \liminf_{s \rightarrow \infty} s |h(is\chi)| < \infty,$$

$h$  satisfies condition (6.5) of Theorem 6.2. Therefore  $h$  has a representation of type 2, say

$$h(z) = a + \langle (A - z_Y)^{-1} \alpha, \alpha \rangle.$$

It remains to show that  $a = 0$ . The inequality (6.8) implies that there exists a sequence  $(s_j)$  tending to  $\infty$  such that  $h(is_j\chi) \rightarrow 0$ . But

$$\operatorname{Re} h(is_j\chi) = a + \langle A(A^2 + s_n^2)^{-1} \alpha, \alpha \rangle \rightarrow a.$$

Hence  $a = 0$  and  $h$  has a type 1 representation. This establishes (2) $\Rightarrow$ (1).  $\square$

### 7. Carapoints at infinity

How can we recognize from function-theoretic properties whether a given function in the  $n$ -variable Loewner class admits a Nevanlinna representation of a given type? In the preceding section it was shown that it depends on growth along a single ray through the origin. In this section we describe the notion of carapoints at infinity for a function in the Pick class, and in the next section we shall give succinct criteria for the four types in the language of carapoints.

Carapoints (though not with this nomenclature) were first introduced by Carathéodory in 1929 [9] for a function  $\varphi$  on the unit disc, as a hypothesis in the “Julia–Carathéodory Lemma”. For any  $\tau \in \mathbb{T}$ , a function  $\varphi$  in the Schur class satisfies the Carathéodory condition at  $\tau$  if

$$\liminf_{\lambda \rightarrow \tau} \frac{1 - |\varphi(\lambda)|}{1 - |\lambda|} < \infty. \tag{7.1}$$

The notion has been generalized to other domains by many authors. Consider domains  $U \subset \mathbb{C}^n$  and  $V \subset \mathbb{C}^m$  and an analytic function  $\varphi$  from  $U$  to the closure of  $V$ . The function  $\varphi$  is said to satisfy Carathéodory’s condition at  $\tau \in \partial U$  if

$$\liminf_{\lambda \rightarrow \tau} \frac{\operatorname{dist}(\varphi(\lambda), \partial V)}{\operatorname{dist}(\lambda, \partial U)} < \infty.$$

Thus, for example, when  $U = \Pi^n$ ,  $V = \Pi$ , a function  $h \in \mathcal{P}_n$  satisfies Carathéodory’s condition at the point  $x \in \mathbb{R}^n$  if

$$\liminf_{z \rightarrow x} \frac{\operatorname{Im} h(z)}{\min_j \operatorname{Im} z_j} < \infty. \tag{7.2}$$

This definition works well for finite points in  $\partial U$ , but for our present purpose we need to consider points at infinity in the boundaries of  $\Pi^n$  and  $\Pi$ . We shall introduce a variant of Carathéodory’s condition for the class  $\mathcal{P}_n$  with the aid of the Cayley transform

$$z = i \frac{1 + \lambda}{1 - \lambda}, \quad \lambda = \frac{z - i}{z + i}, \tag{7.3}$$

which furnishes a conformal map between  $\mathbb{D}$  and  $\Pi$ , and hence a biholomorphic map between  $\mathbb{D}^n$  and  $\Pi^n$  by coordinatewise action. We obtain a one-to-one correspondence between  $\mathcal{S}_n \setminus \{\mathbf{1}\}$  and  $\mathcal{P}_n$  via the formulae

$$h(z) = i \frac{1 + \varphi(\lambda)}{1 - \varphi(\lambda)}, \quad \varphi(\lambda) = \frac{h(z) - i}{h(z) + i} \tag{7.4}$$

where  $\mathbf{1}$  is the constant function equal to 1 and  $\lambda, z$  are related by equations (7.3). For  $\varphi \in \mathcal{S}_n$  we define  $\tau \in \mathbb{T}^n$  to be a *carapoint* of  $\varphi$  if

$$\liminf_{\lambda \rightarrow \tau} \frac{1 - |\varphi(\lambda)|}{1 - \|\lambda\|_\infty} < \infty. \tag{7.5}$$

We can now extend the notion of carapoints to points at infinity. The point  $(\infty, \dots, \infty)$  in the boundary of  $\Pi^n$  corresponds to the point  $\chi$  in the closed unit polydisc; as in the last section,  $\chi$  denotes the point  $(1, \dots, 1) \in \mathbb{C}^n$ .

**Definition 7.1.** Let  $h$  be a function in the Pick class  $\mathcal{P}_n$  with associated function  $\varphi$  in the Schur class  $\mathcal{S}_n$  given by equation (7.4). Let  $\tau \in \mathbb{T}^n, x \in (\mathbb{R} \cup \infty)^n$  be related by

$$x_j = i \frac{1 + \tau_j}{1 - \tau_j} \quad \text{for } j = 1, \dots, n. \tag{7.6}$$

We say that  $x$  is a *carapoint* for  $h$  if  $\tau$  is a carapoint for  $\varphi$ . We say that  $h$  has a *carapoint at  $\infty$*  if  $h$  has a carapoint at  $(\infty, \dots, \infty)$ , that is, if  $\varphi$  has a carapoint at  $\chi$ .

Note that, for a point  $x \in \mathbb{R}^n$ , to say that  $x$  is a carapoint of  $h$  is *not* the same as saying that  $h$  satisfies the Carathéodory condition (7.2) at  $x$ . Consider the function  $h(z) = -1/z_1$  in  $\mathcal{P}_n$ . Clearly  $h$  does not satisfy Carathéodory’s condition at  $0 \in \mathbb{R}^n$ . However, the function  $\varphi$  in  $\mathcal{S}_n$  corresponding to  $h$  is  $\varphi(\lambda) = -\lambda_1$ , which does have a carapoint at  $-\chi$ , the point in  $\mathbb{T}^n$  corresponding to  $0 \in \mathbb{R}^n$ . Hence  $h$  has a carapoint at 0.

We shall be mainly concerned with carapoints at 0 and  $\infty$ . The following observation will help us identify them. For any  $h \in \mathcal{P}_n$  we define  $h^b \in \mathcal{P}_n$  by

$$h^b(z) = h \left( -\frac{1}{z_1}, \dots, -\frac{1}{z_n} \right) \quad \text{for } z \in \Pi^n.$$

For  $\varphi \in \mathcal{S}_n$  we define

$$\varphi^b(\lambda) = \varphi(-\lambda).$$

If  $h$  and  $\varphi$  are corresponding functions, as in equations (7.4), then so are  $h^b$  and  $\varphi^b$ .

**Proposition 7.2.** *The following conditions are equivalent for a function  $h \in \mathcal{P}_n$ .*

- (1)  $\infty$  is a carapoint for  $h$ ;
- (2)  $0$  is a carapoint for  $h^b$ ;
- (3)

$$\liminf_{y \rightarrow 0^+} \frac{\text{Im } h^b(iy\chi)}{y|h^b(iy\chi) + i|^2} < \infty;$$

(4)

$$\liminf_{y \rightarrow \infty} \frac{y \text{Im } h(iy\chi)}{|h(iy\chi) + i|^2} < \infty.$$

**Proof.** (1) $\Leftrightarrow$ (2) Since  $-\chi \in \mathbb{T}^n$  corresponds under the Cayley transform to  $0 \in \mathbb{R}^n$ , we have

$$\begin{aligned} \infty \text{ is a carapoint of } h &\Leftrightarrow \chi \text{ is a carapoint of } \varphi \\ &\Leftrightarrow -\chi \text{ is a carapoint of } \varphi^b \\ &\Leftrightarrow 0 \text{ is a carapoint of } h^b. \end{aligned}$$

(2) $\Leftrightarrow$ (3) A consequence of the  $n$ -variable Julia–Carathéodory Theorem [13,1], is that  $\tau \in \mathbb{T}^n$  is a carapoint of  $\varphi \in \mathcal{S}_n$  if and only if

$$\liminf_{r \rightarrow 1^-} \frac{1 - |\varphi(r\tau)|}{1 - r} < \infty.$$

It follows that

$$\begin{aligned} 0 \text{ is a carapoint for } h^b &\Leftrightarrow -\chi \text{ is a carapoint for } \varphi^b \\ &\Leftrightarrow \liminf_{r \rightarrow 1^-} \frac{1 - |\varphi^b(-r\chi)|}{1 - r} < \infty \\ &\Leftrightarrow \liminf_{r \rightarrow 1^-} \frac{1 - |\varphi^b(-r, -r)|^2}{1 - r^2} < \infty. \end{aligned}$$

Let  $iy \in \Pi$  be the Cayley transform of  $-r \in (-1, 0)$ , so that  $y \rightarrow 0^+$  as  $r \rightarrow 1^-$ . In view of the identity

$$\frac{1 - |\varphi(\lambda)|^2}{1 - \|\lambda\|_\infty^2} = \left( \max_j \frac{|z_j + i|^2}{\text{Im } z_j} \right) \frac{\text{Im } h(z)}{|h(z) + i|^2} \tag{7.7}$$

we have

$$\begin{aligned}
 0 \text{ is a carapoint for } h^b &\Leftrightarrow \liminf_{y \rightarrow 0^+} \frac{|iy + i|^2}{y} \frac{\operatorname{Im} h^b(iy\chi)}{|h^b(iy\chi) + i|^2} < \infty \\
 &\Leftrightarrow \liminf_{y \rightarrow 0^+} \frac{\operatorname{Im} h^b(iy\chi)}{y|h^b(iy\chi) + i|^2} < \infty.
 \end{aligned}$$

(3) $\Leftrightarrow$ (4) Replace  $y$  by  $1/y$ .  $\square$

**Corollary 7.3.** *If  $f \in \mathcal{P}_n$  satisfies Carathéodory’s condition*

$$\liminf_{z \rightarrow x} \frac{\operatorname{Im} f(z)}{\operatorname{Im} z} < \infty \tag{7.8}$$

at  $x \in \mathbb{R}^n$  then  $x$  is a carapoint for  $f$ . If

$$\liminf_{y \rightarrow \infty} y \operatorname{Im} f(iy\chi) < \infty$$

then  $\infty$  is a carapoint for  $f$ .

**Proof.** Let  $h = f^b \in \mathcal{P}_n$ . Clearly  $|h^b(z) + i| \geq 1$  for all  $z \in \Pi^n$ . If the condition (7.8) holds for  $x = 0$  then

$$\liminf_{z \rightarrow 0} \frac{\operatorname{Im} h^b(z)}{|h^b(z) + i|^2 \min_j \operatorname{Im} z_j} \leq \liminf_{z \rightarrow 0} \frac{\operatorname{Im} h^b(z)}{\min_j \operatorname{Im} z_j} < \infty$$

and hence, by (2) $\Leftrightarrow$ (3) of Proposition 7.2, 0 is a carapoint for  $h^b = f$ . The case of a general  $x \in \mathbb{R}^n$  follows by translation.  $\square$

If  $h \in \mathcal{P}_n$  has a carapoint at  $x \in (\mathbb{R} \cup \infty)^n$  then it has a value at  $x$  in a natural sense. If  $\varphi \in \mathcal{S}_n$  has a carapoint at  $\tau \in \mathbb{T}^n$ , then by [13] there exists a unimodular constant  $\varphi(\tau)$  such that

$$\lim_{\lambda \overset{\text{nt}}{\rightarrow} \tau} \varphi(\lambda) = \varphi(\tau). \tag{7.9}$$

Here  $\lambda \overset{\text{nt}}{\rightarrow} \tau$  means that  $\lambda$  tends nontangentially to  $\tau$  in  $\mathbb{D}^n$ .

**Definition 7.4.** If  $h \in \mathcal{P}_n$  has a carapoint at  $x \in (\mathbb{R} \cup \infty)^n$  then we define

$$h(x) = \begin{cases} \infty & \text{if } \varphi(\tau) = 1 \\ i \frac{1 + \varphi(\tau)}{1 - \varphi(\tau)} & \text{if } \varphi(\tau) \neq 1 \end{cases}$$

where  $\tau \in \mathbb{T}^n$  corresponds to  $x$  as in equation (7.6).

Thus  $h(\infty) \in \mathbb{R} \cup \{\infty\}$  when  $\infty$  is a carapoint of  $h$ .

In the example  $h(z) = -1/z_1$ , since the value of  $\varphi(-\lambda)$  at  $-\chi$  is 1, we have  $h(0) = \infty$ .

Although the value of  $h(\infty)$  is defined in terms of the Schur class function  $\varphi$ , it can be expressed more directly in terms of  $h$ .

**Proposition 7.5.** *If  $\infty$  is a carapoint of  $h$  then*

$$h(\infty) = h^b(0) = \lim_{z \xrightarrow{nt} \infty} h(z). \tag{7.10}$$

Here we say that  $z \xrightarrow{nt} \infty$  if  $z \rightarrow (\infty, \dots, \infty)$  in the set  $\{z \in \Pi^n : (-1/z_1, \dots, -1/z_n) \in S\}$  for some set  $S \subset \Pi^n$  that approaches 0 nontangentially, or equivalently, if  $z \rightarrow (\infty, \dots, \infty)$  in a set on which  $\|z\|_\infty / \min_j \text{Im } z_j$  is bounded.

**Proof of Proposition 7.5.** Clearly

$$h(\infty) = \infty \iff \varphi(\chi) = 1 \iff \varphi^b(-\chi) = 1 \iff h^b(0) = \infty.$$

Similarly, for  $\xi \in \mathbb{R}$ ,

$$h(\infty) = \xi \iff \varphi(\chi) = \frac{\xi - i}{\xi + i} \iff \varphi^b(-\chi) = \frac{\xi - i}{\xi + i} \iff h^b(0) = \xi.$$

Thus, whether  $h(\infty)$  is finite or infinite,  $h(\infty) = h^b(0)$ . Equation (7.10) follows from the relation (7.9).  $\square$

### 8. Types of functions in the Loewner class

In this section we shall show that the type of a function  $h \in \mathcal{L}_n$  is entirely determined by whether or not  $\infty$  is a carapoint of  $h$  and by the value of  $h(\infty)$ . Let us make precise the notion of the *type* of a function in  $\mathcal{L}_n$ .

**Definition 8.1.** A function  $h \in \mathcal{L}_n$  is of *type 1* if it has a Nevanlinna representation of type 1. For  $n = 2, 3$  or 4 we say that  $h$  is of *type  $n$*  if  $h$  has a Nevanlinna representation of type  $n$  but has no representation of type  $n - 1$ .

Clearly every function in  $\mathcal{L}_n$  is of exactly one of the types 1 to 4. We shall now prove [Theorem 1.11](#). Recall that it states the following, for any function  $h \in \mathcal{L}_n$ .

- (1)  $h$  is of type 1 if and only if  $\infty$  is a carapoint of  $h$  and  $h(\infty) = 0$ ;
- (2)  $h$  is of type 2 if and only if  $\infty$  is a carapoint of  $h$  and  $h(\infty) \in \mathbb{R} \setminus \{0\}$ ;
- (3)  $h$  is of type 3 if and only if  $\infty$  is not a carapoint of  $h$ ;
- (4)  $h$  is of type 4 if and only if  $\infty$  is a carapoint of  $h$  and  $h(\infty) = \infty$ .

**Proof of Theorem 1.11.** (2) Let  $h \in \mathcal{L}_n$  have a type 2 representation  $h(z) = a + \langle (A - z_Y)^{-1}v, v \rangle$  with  $a \neq 0$ . By [Theorem 6.2](#),

$$\liminf_{y \rightarrow \infty} y \operatorname{Im} h(iy\chi) < \infty.$$

By [Corollary 7.3](#),  $\infty$  is a carapoint for  $h$ . Furthermore, by [Proposition 7.5](#)

$$h(\infty) = \lim_{y \rightarrow \infty} h(iy\chi) = a \in \mathbb{R} \setminus \{0\}.$$

Conversely, suppose that  $\infty$  is a carapoint for  $h$  and  $h(\infty) \in \mathbb{R} \setminus \{0\}$ . By [Proposition 7.2](#)

$$\liminf_{y \rightarrow \infty} \frac{y \operatorname{Im} h(iy\chi)}{|h(iy\chi) + i|^2} < \infty$$

while by [Proposition 7.5](#)

$$\lim_{y \rightarrow \infty} |h(iy\chi) + i|^2 = h(\infty)^2 + 1 \in (1, \infty).$$

On combining these two limits we find that

$$\liminf_{y \rightarrow \infty} y \operatorname{Im} h(iy\chi) < \infty,$$

and so, by [Theorem 6.2](#),  $h$  has a representation of type 2. Since  $h(\infty) \neq 0$  it is clear that  $h$  does not have a representation of type 1. Thus (2) holds.

A trivial modification of the above argument proves that (1) is also true.

(4) Let  $h$  be of type 4. Then  $h$  has no type 3 representation, and so, by [Theorem 6.1](#), there exists  $\delta > 0$  and a sequence  $(s_n)$  of positive numbers tending to  $\infty$  such that

$$\frac{1}{s_n} \operatorname{Im} h(is_n\chi) \geq \delta > 0.$$

Let  $y_n = 1/s_n$ ; then  $-1/(is_n) = iy_n$ , and we have

$$y_n \operatorname{Im} h^b(iy_n\chi) \geq \delta \quad \text{for all } n \geq 1. \tag{8.1}$$

Since  $|h^b(z) + i| > \operatorname{Im} h^b(z)$  for all  $z$ , we have

$$\begin{aligned} \liminf_{z \rightarrow 0} \frac{\operatorname{Im} h^b(z)}{|h^b(z) + i|^2 \min_j \operatorname{Im} z_j} &\leq \liminf_{z \rightarrow 0} \frac{1}{\operatorname{Im} h^b(z) \min_j \operatorname{Im} z_j} \\ &\leq \liminf_{n \rightarrow \infty} \frac{1}{y_n \operatorname{Im} h^b(iy_n\chi)} \\ &\leq 1/\delta. \end{aligned}$$

Hence  $(0, 0)$  is a carapoint of  $h^b$ , and so  $\infty$  is a carapoint of  $h$ .

Since  $y_n \rightarrow 0$  it follows from the inequality (8.1) that  $\text{Im } h^b(iy_n\chi) \rightarrow \infty$ , hence that  $h^b(0) = \infty$ , and therefore that  $h(\infty) = \infty$ .

Conversely, suppose that  $\infty$  is a carapoint of  $h$  and that  $h(\infty) = \infty$ . We shall show that

$$\lim_{s \rightarrow \infty} \frac{1}{s} \text{Im } h(is\chi) \neq 0, \tag{8.2}$$

and it will follow from Theorem 6.1 that  $h$  does not have a representation of type 3, that is,  $h$  is of type 4.

Let  $\varphi \in \mathcal{S}_n$  correspond to  $h$  and let  $r \in (0, 1)$  correspond to  $is \in \Pi$ . Then

$$\begin{aligned} \frac{1}{s} \text{Im } h(is\chi) &= \frac{1-r}{1+r} \frac{1 - |\varphi(r\chi)|^2}{|1 - \varphi(r\chi)|^2} \\ &= \frac{1 - |\varphi(r\chi)|^2}{1 - r^2} \frac{(1-r)^2}{|1 - \varphi(r\chi)|^2}. \end{aligned} \tag{8.3}$$

By hypothesis,  $\chi$  is a carapoint for  $\varphi$  and  $\varphi(\chi) = 1$ . By definition of carapoint,

$$\liminf_{z \rightarrow \chi} \frac{1 - |\varphi(z)|^2}{1 - \|z\|_\infty^2} = \alpha < \infty \quad \text{for all } s > 0.$$

The  $n$ -variable Julia–Carathéodory Lemma (see [13,1]) now tells us that  $\alpha > 0$  and

$$\frac{|1 - \varphi(r\chi)|^2}{|1 - r|^2} \leq \alpha \frac{1 - |\varphi(r\chi)|^2}{1 - r^2} \quad \text{for all } r \in (0, 1). \tag{8.4}$$

On combining equations (8.3) and (8.4) we obtain

$$\frac{1}{s} \text{Im } h(is\chi) \geq \frac{1}{\alpha} > 0 \quad \text{for all } s > 0.$$

Thus the relation (8.2) is true, and so, by Theorem 6.1,  $h$  is of type 4.

Statement (3) now follows easily. The function  $h \in \mathcal{L}_n$  is of type 3 if and only if it is not of types 1, 2 or 4, hence if and only if it is not the case that  $\infty$  is a carapoint for  $h$  and  $h(\infty) \in \mathbb{R} \cup \{\infty\}$ , hence if and only if  $\infty$  is not a carapoint of  $h$ .  $\square$

We now show that there are functions in the Pick class  $\mathcal{P}_2$  of all four types. We return to Example 5.6 and show that the functions in  $\mathcal{P}_2$  which we presented there are indeed of the stated types.

**Example 8.2.** (1) The function

$$h(z) = -\frac{1}{z_1 + z_2} = \langle (0 - z_Y)^{-1}v, v \rangle_{\mathbb{C}},$$

where  $Y = \frac{1}{2}$  and  $v = 1/\sqrt{2}$ , is obviously of type 1. Let us nevertheless check that  $\infty$  is a carapoint of  $h$  and  $h(\infty) = 0$ , in accordance with [Theorem 1.11](#). We have  $h(iy, iy) = \frac{1}{2}i/y$  and hence

$$\liminf_{y \rightarrow 0^+} y \operatorname{Im} h(iy, iy) = \frac{1}{2}.$$

Thus  $\infty$  is a carapoint for  $h$  by [Proposition 7.2](#). Moreover  $h(iy, iy) \rightarrow 0$  as  $y \rightarrow \infty$ , and therefore  $h(\infty) = 0$ .

(2) It is immediate that the function  $1 + h$ , with  $h$  as in (1), is of type 2, and that  $\infty$  is a carapoint of  $1 + h$  with value 1.

(3) We have seen that the function

$$h(z) = \begin{cases} \frac{1}{1 + z_1 z_2} \left( z_1 - z_2 + \frac{i z_2 (1 + z_1^2)}{\sqrt{z_1 z_2}} \right) & \text{if } z_1 z_2 \neq -1 \\ \frac{1}{2}(z_1 + z_2) & \text{if } z_1 z_2 = -1 \end{cases} \tag{8.5}$$

has a representation of type 3. To show that  $h$  is indeed of type 3 we must prove that  $\infty$  is not a carapoint of  $h$ .

For all  $y > 0$  we have  $h(iy, iy) = i$ . Hence

$$\liminf_{y \rightarrow \infty} \frac{y \operatorname{Im} h(iy, iy)}{|h(iy, iy) + i|^2} = \liminf_{y \rightarrow \infty} \frac{y}{4} = \infty.$$

By [Proposition 7.2](#),  $\infty$  is not a carapoint for  $h$ . Thus  $h$  is of type 3.

(4) The function

$$h(z) = \frac{z_1 z_2}{z_1 + z_2} = -1 \Big/ \left( -\frac{1}{z_1} - \frac{1}{z_2} \right)$$

is clearly in  $\mathcal{P}_2$ . We gave a type 4 representation of  $h$  in [Example 5.6](#). We claim that  $\infty$  is a carapoint of  $h$ . We have  $h(iy, iy) = \frac{1}{2}iy$ , and thus

$$\liminf_{y \rightarrow \infty} \frac{y \operatorname{Im} h(iy, iy)}{|h(iy, iy) + i|^2} = \liminf_{y \rightarrow \infty} \frac{\frac{1}{2}y^2}{|\frac{1}{2}iy + i|^2} = 2.$$

Hence  $\infty$  is a carapoint for  $h$ . Furthermore  $h(iy, iy) = \frac{1}{2}iy \rightarrow \infty$  as  $y \rightarrow \infty$ , and so  $h(\infty) = \infty$ . Thus  $h$  is of type 4.

Another example of a function of type 4 is  $h(z) = \sqrt{z_1 z_2}$ .

### 9. Rates of growth in the Loewner class

The Nevanlinna representation formulae give rise to growth estimates for functions in the  $n$ -variable Loewner class. It turns out that growth is mild, both at infinity and close to the real axis. Even though the type of a function is determined by its growth on the single ray  $\{iy\chi : y > 0\}$ , in turn the growth of the function on the entire polyhalf-plane is constrained by its type.

Consider first the one-variable case. If  $h$  is the Cauchy transform of a finite positive measure  $\mu$  then

$$|h(z)| \leq \int \frac{d\mu(t)}{|t - z|} \leq \int \frac{d\mu(t)}{\text{Im } z} = \frac{C}{\text{Im } z}$$

for some  $C > 0$  and for all  $z \in \Pi$ . For a general function  $h$  in the Pick class, by Nevanlinna’s representation ([Theorem 1.2](#)) there exist  $a \in \mathbb{R}$ ,  $b \geq 0$  and a finite positive measure  $\mu$  on  $\mathbb{R}$  such that, for all  $z \in \Pi$ ,

$$\begin{aligned} h(z) &= a + bz + \int \frac{1 + tz}{t - z} d\mu(t) \\ &= a + bz + \int \frac{1 + z^2}{t - z} + z d\mu(t) \end{aligned}$$

and therefore

$$\begin{aligned} |h(z)| &\leq |a| + b|z| + \left( \frac{1 + |z|^2}{\text{Im } z} + |z| \right) \mu(\mathbb{R}) \\ &\leq C \left( 1 + |z| + \frac{1 + |z|^2}{\text{Im } z} \right) \end{aligned}$$

for some  $C > 0$ .

Similar estimates hold for the Loewner class.

**Proposition 9.1.** *For any function  $h \in \mathcal{L}_n$  there exists a non-negative number  $C$  such that, for all  $z \in \Pi^n$ ,*

$$|h(z)| \leq C \left( 1 + \|z\|_1 + \frac{1 + \|z\|_1^2}{\min_j \text{Im } z_j} \right). \tag{9.1}$$

*For any function  $h \in \mathcal{L}_n$  of type 2 there exists a non-negative number  $C$  such that, for all  $z \in \Pi^n$ ,*

$$|h(z)| \leq C \left( 1 + \frac{1}{\min_j \text{Im } z_j} \right). \tag{9.2}$$

For any function  $h \in \mathcal{L}_n$  of type 1 there exists a non-negative number  $C$  such that, for all  $z \in \Pi^n$ ,

$$|h(z)| \leq \frac{C}{\min_j \operatorname{Im} z_j}. \tag{9.3}$$

**Proof.** Let  $h \in \mathcal{L}_n$ . Let  $\mathcal{N}, \mathcal{M}, A, P, a$  and  $v$  be as in [Theorem 1.9](#), so that

$$h(z) = a + \langle M(z)v, v \rangle$$

for all  $z \in \Pi^n$ , where  $M(z)$  is the matricial resolvent given by equation [\(3.1\)](#). By [Proposition 3.1](#) we have, for all  $z \in \Pi^n$ ,

$$\begin{aligned} \|M(z)\| &\leq (1 + \sqrt{10}\|z\|_1) \left( 1 + \frac{1 + \sqrt{2}\|z\|_1}{\min_j \operatorname{Im} z_j} \right) \\ &\leq 1 + \sqrt{10}\|z\|_1 + B \frac{1 + \|z\|_1 + \|z\|_1^2}{\min_j \operatorname{Im} z_j} \end{aligned}$$

for a suitable choice of  $B \geq 0$ . Hence

$$\begin{aligned} |h(z)| &\leq |a| + \|M(z)\| \|v\|^2 \\ &\leq |a| + \left( 1 + \sqrt{10}\|z\|_1 + B \frac{1 + \|z\|_1 + \|z\|_1^2}{\min_j \operatorname{Im} z_j} \right) \|v\|^2. \end{aligned}$$

Since

$$1 + \|z\|_1 + \|z\|_1^2 \leq \frac{3}{2}(1 + \|z\|_1^2),$$

we have

$$|h(z)| \leq C \left( 1 + \|z\|_1 + \frac{1 + \|z\|_1^2}{\min_j \operatorname{Im} z_j} \right)$$

for some choice of  $C > 0$  and for all  $z \in \Pi^n$ . Thus the estimate [\(9.1\)](#) holds.

Similarly, the estimates [\(9.2\)](#) and [\(9.3\)](#) follow easily from the simple resolvent estimate [\(2.1\)](#).  $\square$

### 10. Structured resolvent identities

To conclude the paper we point out that there are structured analogs of the classical resolvent identity

$$(A - z)^{-1} - (A - w)^{-1} = (z - w)(A - z)^{-1}(A - w)^{-1}$$

for any  $z, w$  in the resolvent set of an operator  $A$ .

**Proposition 10.1.** *Let  $A$  be a densely defined self-adjoint operator on a Hilbert space  $\mathcal{H}$  and let  $Y$  be a positive decomposition of  $\mathcal{H}$ . For all  $z, w \in \Pi^n$*

$$(A - z_Y)^{-1} - (A - w_Y)^{-1} = (A - z_Y)^{-1}(z - w)_Y(A - w_Y)^{-1}. \tag{10.1}$$

*If  $M(z)$  is the structured resolvent of type 3 corresponding to  $A$  and  $Y$  then*

$$M(z) - M(w) |_{\mathcal{D}(A)} = (1 - iA)(A - z_Y)^{-1}(z - w)_Y(A - w_Y)^{-1}(1 + iA). \tag{10.2}$$

**Proof.** The first of these identities is immediate. For the second, by equation (2.8),

$$M(z) - M(w) |_{\mathcal{D}(A)} = (1 - iA) \left( (A - z_Y)^{-1} - (A - w_Y)^{-1} \right) (1 + iA),$$

and the identity (10.2) follows from equation (10.1).  $\square$

**Proposition 10.2.** *Let  $\mathcal{H}$  be the orthogonal direct sum of Hilbert spaces  $\mathcal{N}, \mathcal{M}$ , let  $A$  be a densely defined self-adjoint operator on  $\mathcal{M}$  with domain  $\mathcal{D}(A)$  and let  $P$  be an orthogonal decomposition of  $\mathcal{H}$ . For every  $z, w \in \Pi^n$ , as operators on  $\mathcal{N} \oplus \mathcal{D}(A)$ ,*

$$\begin{aligned} M(z) - M(w) &= \begin{bmatrix} -i & 0 \\ 0 & 1 - iA \end{bmatrix} \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \right)^{-1} (z - w)_P \\ &\quad \times \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} w_P \right)^{-1} \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix}. \end{aligned} \tag{10.3}$$

**Proof.** Let

$$D = \begin{bmatrix} i & 0 \\ 0 & 1 + iA \end{bmatrix} : \mathcal{N} \oplus \mathcal{D}(A) \rightarrow \mathcal{H}.$$

By equations (3.11) and (3.12) we have

$$\begin{aligned} &M(z) - M(w) |_{\mathcal{N} \oplus \mathcal{D}(A)} \\ &= D^* \left\{ \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \right)^{-1} \left( z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \right. \\ &\quad \left. - \left( \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} w_P + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} w_P \right)^{-1} \right\} D \\ &= D^* \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \right)^{-1} \left\{ \left( z_P \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} w_P \right) \right. \\ &\quad \left. - \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - z_P \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \right) \left( \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} w_P + \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right) \right\} \\ &\quad \times \left( \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix} - \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} w_P \right)^{-1} D. \end{aligned}$$

The term in braces in the last expression reduces to  $(z - w)_P$ , and the identity (10.3) follows.  $\square$

**Corollary 10.3.** *With the assumptions of Proposition 10.2, there exists an analytic function  $F : \Pi^n \rightarrow \mathcal{L}(\mathcal{H})$  such that, for all  $z, w \in \Pi^n$ ,*

$$M(z) - M(w) = F(\bar{z})^*(z - w)_P F(w). \quad (10.4)$$

The statement follows from Proposition 10.2 just as Proposition 3.5 follows from Proposition 3.4. If  $F$  is defined by equation (3.21) then  $F(z)$  is a bounded operator on  $\mathcal{H}$ ,  $F$  is analytic on  $\Pi^n$  and Proposition 10.2 states that equation (10.4) holds on  $\mathcal{N} \oplus \mathcal{D}(A)$ . It follows by continuity that equation (10.4) holds on  $\mathcal{H}$ .

## References

- [1] M. Abate, The Julia–Wolff–Carathéodory Theorem in polydisks, *J. Anal. Math.* 74 (1998) 275–306.
- [2] J. Agler, On the representation of certain holomorphic functions defined on a polydisc, in: *Operator Theory: Advances and Applications*, vol. 48, Birkhäuser, Basel, 1990, pp. 47–66.
- [3] J. Agler, J.E. McCarthy, N.J. Young, A Carathéodory theorem for the bidisk via Hilbert space methods, *Math. Ann.* 352 (2012) 581–624.
- [4] J. Agler, J.E. McCarthy, N.J. Young, Operator monotone functions and Löwner functions of two variables, *Ann. of Math.* 176 (2012) 1783–1826.
- [5] J. Agler, R. Tully-Doyle, N.J. Young, Boundary behavior of analytic functions of two variables via generalized models, *Indag. Math.* 23 (2012) 995–1027.
- [6] J.A. Ball, D. Kalyuzhnyi-Verbovetzkyi, Conservative dilations of dissipative multidimensional systems: the commutative and non-commutative settings, *Multidimens. Syst. Signal Process.* 19 (2008) 79–122.
- [7] J.A. Ball, D. Kalyuzhnyi-Verbovetzkyi, Schur–Agler and Herglotz–Agler classes of functions: positive-kernel decompositions and transfer-function realizations, *Adv. Math.* 280 (2015) 121–187.
- [8] J.A. Ball, O.J. Staffans, Conservative state-space realizations of dissipative system behaviors, *Integral Equations Operator Theory* 54 (2006) 151–213.
- [9] C. Carathéodory, Über die Winkelderivierten von beschränkten analytischen Funktionen, *Sitzunber. Preuss. Akad. Wiss.*, 1929.
- [10] W.F. Donoghue, *Monotone Matrix Functions and Analytic Continuation*, Springer, Berlin, 1974.
- [11] A. Grinshpan, D.S. Kalyuzhnyi-Verbovetzkyi, V. Vinnikov, H.J. Woerdeman, Classes of tuples of commuting contractions satisfying the multivariable von Neumann inequality, *J. Funct. Anal.* 256 (2009) 3035–3054.
- [12] G. Herglotz, Über Potenzreihen mit positivem reellen Teil im Einheitskreis, *Leipz. Ber.* 63 (1911) 501–511.
- [13] F. Jafari, Angular derivatives in polydisks, *Indian J. Math.* 35 (1993) 197–212.
- [14] G. Julia, Extension nouvelle d’un lemme de Schwarz, *Acta Math.* 42 (1920) 349–355.
- [15] D.S. Kalyuzhnyi-Verbovetzkyi, Carathéodory interpolation on the non-commutative polydisk, *J. Funct. Anal.* 229 (2005) 241–276.
- [16] P. Lax, *Functional Analysis*, Wiley, 2002.
- [17] K. Löwner, Über monotone Matrixfunktionen, *Math. Z.* 38 (1934) 177–216.
- [18] R. Nevanlinna, Asymptotische Entwicklungen beschränkter Funktionen und das Stieltjesche Momentenproblem, *Ann. Acad. Sci. Fenn. Ser. A* 18 (1922).
- [19] S. Parrott, Unitary dilations for commuting contractions, *Pacific Math. J.* 34 (1970) 481–490.
- [20] G. Pick, Über die Beschränkungen analytischer Funktionen, welche durch vorgegebene Funktionswerte bewirkt werden, *Math. Ann.* 77 (1916) 7–23.
- [21] F. Riesz, B. Sz. Nagy, *Functional Analysis*, Dover, New York, 1990.
- [22] N.Th. Varopoulos, On an inequality of von Neumann and an application of the metric theory of tensor products to operators theory, *J. Funct. Anal.* 16 (1974) 83–100.