



This is a repository copy of *On the Integral Cohomology of Bianchi groups*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/97992/>

Version: Accepted Version

Article:

Sengun, M.H. (2011) *On the Integral Cohomology of Bianchi groups*. *Experimental Mathematics*, 20 (4). pp. 487-505. ISSN 1058-6458

<https://doi.org/10.1080/10586458.2011.594671>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

On the Integral Cohomology of Bianchi groups

Mehmet Haluk Şengün

Abstract

Extensive and systematic machine computations are carried out to investigate the integral cohomology of the Euclidean Bianchi groups and their congruence subgroups. The collected data give insight on several aspects, including the asymptotic behaviour of the torsion in the first homology. Along with the experimental work, some basic properties of the integral cohomology are recorded with an eye towards the liftability issue of Hecke eigenvalue systems.

1. Introduction

Bianchi groups are groups of the form $\mathrm{PSL}_2(\mathcal{O})$ where \mathcal{O} is the ring of integers of an imaginary quadratic field. First studied by L. Bianchi in 1892 [Bi], these groups form an important class of arithmetic Kleinian groups. In fact, it is well known that any non-cocompact arithmetic Kleinian group, after conjugation, is commensurable with a Bianchi group.

In this paper, I will be interested in the cohomology of Bianchi groups with certain \mathcal{O} -module coefficients. These cohomology groups are fundamental to the study of Bianchi modular forms, that is, modular forms (for \mathbf{GL}_2) over an imaginary quadratic field. Unlike their analogs over totally real fields, i.e. Hilbert modular forms, the arithmetic of Bianchi modular forms is little understood. One of the features that obstruct the application of standard methods is the torsion in the cohomology of Bianchi groups.

The first computations of torsion were performed in 1981 by Elstrodt, Grunewald and Mennicke [EGM] where they computed the abelianizations $\Gamma_0(\mathfrak{p})^{ab} (\simeq H_1(\Gamma_0(\mathfrak{p}), \mathbb{Z}))$ of congruence subgroups $\Gamma_0(\mathfrak{p})$ for prime ideals \mathfrak{p} of $\mathbb{Z}[i]$ of residue degree 1 and norm ≤ 400 . In the same paper, numerical evidence suggesting a connection between some of the 2-torsion classes and certain S_3 -extension of $\mathbb{Q}(i)$ was exhibited. Later in 1994, Figueiredo [Fi] provided several more examples which suggested the same kind of connection between certain 3-torsion classes and certain $\mathrm{SL}_2(\mathbb{F}_3)$ -extensions of the fields $\mathbb{Q}(\sqrt{-d})$ with $d = 1, 2, 3$. Recently further examples have been found by myself in [Se2] and by Torrey in [To] where she formulates and tests an analogue of the, now proven, strong modularity conjecture of Serre [S2].

In 1983, Grunewald and Schwermer [GS] determined the conjugacy classes of small index subgroups of several Bianchi groups. In particular, they computed the abelianizations of a vast amount of small index (≤ 12) subgroups. Depending on the data they collected and the data in [EGM], they speculated that for any finite index subgroup Γ

¹2010 *Mathematics Subject Classification*. Primary 11F41, 11F75; Secondary 22E40

²*Key words and phrases*. Bianchi groups, torsion in the homology, cuspidal cohomology

of a Bianchi group G , if a prime p appears as an exponent for an element of Γ^{ab} , then p should be smaller than half of the index of Γ in G .

In 1988, Taylor in his unpublished thesis [T] proved the existence of p^m -torsion classes in $H_1(\Gamma_1(Np^r), E_{k,\ell}(\mathcal{O}))$ (notation will be explained later) with $k \neq \ell$ under special circumstances. In her unpublished 2000 thesis [P], Priplata numerically investigated the torsion for some Bianchi groups, mostly for coefficient modules $E_{k,\ell}$ where $k \neq \ell$. It is worth remarking that the (co)homology of Bianchi groups is related to cuspidal Bianchi modular forms only in the parallel weight case, that is, in the case $k = \ell$.

In this paper I report on my extensive systematic computations of the integral (co)homology of Bianchi groups. More specifically, I worked with the groups $\mathrm{PSL}(\mathcal{O}_d)$ and $\mathrm{PGL}_2(\mathcal{O}_d)$ with $-d = 1, 2, 3, 7, 11$ and also with their congruence subgroups. Motivated by the possibility of congruences between cuspidal Bianchi modular forms and torsion classes, and also the arithmetical connections mentioned above, I limited myself to the (co)homology with parallel weights. The data I collected

- show that H_{cusp}^2 can have sporadically large torsion part with very little torsion-free part, in particular the speculations of Grunewald and Schwermer mentioned above are false,
- support a recent conjecture of Long, Maclachlan and Reid [LMR] on existence of certain families of rational homology spheres,
- strongly suggest that an analogue of the very recent result of Bergeron-Venkatesh [BV] on the asymptotic behaviour of the torsion in the homology of cocompact arithmetic congruence lattices in $\mathrm{SL}_2(\mathbb{C})$ holds for Bianchi groups.

Along with the computational work, I record some basic properties of the integral cohomology with an eye towards the liftability issue of Hecke eigenvalue systems.

Acknowledgments I started this work towards the end of my postdoctoral fellowship within the Arithmetic Geometry Research Group of the University of Duisburg-Essen under SFB/TR 45 of the Deutsche Forschungsgemeinschaft. I am grateful for the wonderful ambiance and hospitality provided by the Mathematics Department. I gratefully acknowledge that all the computations were carried out on the computers of the Institute for Experimental Mathematics, Essen. I thank the excellent Hausdorff Institute for Mathematics in Bonn where I completed the rest of this work as a visitor. I would like to thank Gebhard Böckle and Gabor Wiese for enlightening discussions and many comments/corrections on this work. My correspondence with Nicolas Bergeron and Akshay Venkatesh was very encouraging and gave me more directions for experimentation. I am grateful to Nansen Petrosyan for his help with the spectral sequences. Finally I thank J-P. Serre and the referees whose numerous comments improved the paper.

Dedication After the first draft of this work was completed, *Fritz Grunewald*, whom I consider my second PhD adviser, tragically passed away. Over the years, I benefited greatly from his invaluable guidance and generous support. I dedicate this paper to his memory with admiration, gratitude and love.

2. The modules

Given a commutative a ring R , let $E_k(R) \simeq R[x, y]_k$ where the latter is the space of homogeneous degree k polynomials in two variables over R . Note that $\{x^{k-i}y^i : 0 \leq i \leq k\}$ is an R -basis of $E_k(R)$.

For a polynomial $P(X, Y)$ in $E_k(R)$ and a matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in $M_2(R)$, then we have the right action

$$(P \cdot \begin{pmatrix} a & b \\ c & d \end{pmatrix})(X, Y) = P\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix}\right) = P(aX + bY, cX + dY).$$

Let \mathcal{O} be the ring of integers of an imaginary quadratic field. Consider the $M_2(\mathcal{O})$ -module

$$E_{k,\ell}(\mathcal{O}) := E_k(\mathcal{O}) \otimes_{\mathcal{O}} \overline{E_{\ell}(\mathcal{O})}.$$

Here the overline on the second factor is to indicate that action on the second factor is twisted with complex conjugation. Note that we should insist that $k + \ell$ is even so that $-Id$ acts trivially and thus $\mathrm{PSL}(2, \mathcal{O})$ acts on it as well.

It is useful to remark that $E_{k,\ell}(\mathcal{O}) \simeq \mathrm{Sym}^k(\mathcal{O}^2) \otimes_{\mathcal{O}} \overline{\mathrm{Sym}^{\ell}(\mathcal{O}^2)}$ as $M_2(\mathcal{O})$ -modules where $\mathrm{Sym}^i(\mathcal{O}^2)$ is the i th symmetric power of the standard representation of $M_2(\mathcal{O})$ on \mathcal{O}^2 . Here the overline on the second factor means that the action is twisted with complex conjugation.

Let π be a prime element of \mathcal{O} over a rational prime p . Put κ_{π} for its residue field. We put

$$E_{k,\ell}(\kappa_{\pi}) := E_{k,\ell}(\mathcal{O}) \otimes_{\mathcal{O}} \kappa_{\pi}.$$

If p splits in \mathcal{O} , then

$$E_{k,\ell}(\kappa_{\pi}) \simeq E_k(\kappa_{\pi}) \otimes E_{\ell}(\kappa_{\bar{\pi}}).$$

Thus $\mathrm{PSL}_2(\mathcal{O})$ acts on it by reduction mod π on the first factor and by reduction mod $\bar{\pi}$ on the second. If p is inert in \mathcal{O} , then

$$E_{k,\ell}(\kappa_{\pi}) \simeq E_k(\kappa_{\pi}) \otimes E_{\ell}(\kappa_{\pi})^{\sigma}.$$

Here $\mathrm{PSL}_2(\mathcal{O})$ acts by reduction mod π . The action on the second factor is twisted by the nontrivial automorphism σ of κ_{π} .

Finally, when p is ramified in \mathcal{O} , we have

$$E_{k,\ell}(\kappa_{\pi}) \simeq E_k(\kappa_{\pi}) \otimes E_{\ell}(\kappa_{\pi}).$$

Here the action of $\mathrm{PSL}_2(\mathcal{O})$ is via reduction mod π and is the same on both factors.

A result of Brauer and Nesbitt [BN] tells us that in the inert and split cases, the $\mathrm{PSL}_2(\mathcal{O})$ -modules $E_{k,\ell}(\kappa_{\pi})$ are irreducible only when $0 \leq k, \ell \leq p - 1$. In the ramified case, $E_{k,\ell}(\kappa_{\pi})$ is never an irreducible $\mathrm{PSL}_2(\mathcal{O})$ -module unless $k = 0 \leq \ell \leq p - 1$ or $\ell = 0 \leq k \leq p - 1$. For more on the structure of these modules, we refer to reader to [ST].

The following will be used later.

Proposition 2.1. *Let \mathcal{O} be the ring of integers of an imaginary quadratic field. Let $k \geq \ell$ and put $R = \mathcal{O}[\frac{1}{k\ell}]$. Then there is a $\mathrm{PSL}_2(\mathcal{O})$ -equivariant perfect pairing*

$$E_{k,\ell}(R) \times E_{k,\ell}(R) \rightarrow R.$$

It is well known (see e.g. [Wi] Lemma 2.4) that there is a perfect pairing on $\text{Sym}^n(R^2)$ coming from the determinant pairing on R^2 whenever $n!$ is invertible in the ring R . The proposition follows by taking the product of the two pairings associated to the two factors of $E_{k,\ell}$. For an explicit description of this pairing, see Section 2.4. of [Be]. As a corollary, we see that the modules $E_{k,\ell}(R)$ are self-dual.

3. the cohomology

In this section I will investigate the integral cohomology of Bianchi groups. My treatment is heavily influenced by work of Hida [Hi], Wang [Wa] and Wiese [Wi].

Let K be an imaginary quadratic field. Let \mathcal{O} be its ring of integers. Let G be the associated Bianchi group. Let Γ be a finite index subgroup of G .

In this paper, we will focus on the \mathcal{O} -modules

$$H^i(\Gamma, E_{k,\ell}(\mathcal{O})), \quad i = 1, 2.$$

It is well known that these are finitely generated \mathcal{O} -modules.

Definition 3.1. Let $\pi \in \mathcal{O}$ be a prime element over the rational prime p . Assume that $H^i(\Gamma, E_{k,\ell}(\mathcal{O}))$ has π -torsion, i.e. it contains a non-zero class c such that $\pi \cdot c = 0$. We say that π is a *large torsion* if $k, \ell < p$. Otherwise, we say that π is a *small torsion*.

Proposition 3.2. *Let π be prime element of \mathcal{O} over the rational prime p . Put κ_π for its residue field. Let Γ be a torsion-free finite index subgroup of the Bianchi group G .*

- (a) *If Γ surjects onto $\text{PSL}_2(\kappa_\pi)$ and π is unramified, then $H^1(\Gamma, E_{k,\ell}(\mathcal{O}))$ has no large π -torsion.*
- (b) *If Γ surjects onto $\text{PSL}_2(\kappa_\pi)$ and π is ramified, then $H^1(\Gamma, E_{k,\ell}(\mathcal{O}))$ has no π -torsion if and only if $k = 0$ and $\leq \ell \leq p - 1$ or vice versa.*
- (c) *The obstruction to the lifting of a class in $H^1(\Gamma, E_{k,\ell}(\kappa_\pi))$ to $H^1(\Gamma, E_{k,\ell}(\mathcal{O}))$ is the π -torsion in $H^2(\Gamma, E_{k,\ell}(\mathcal{O}))$.*
- (d) *$H^2(\Gamma, E_{k,\ell}(\mathcal{O})) \otimes \kappa_\pi \simeq H^2(\Gamma, E_{k,\ell}(\kappa_\pi))$ for every k, ℓ .*

Proof. In the following, let us put $E = E_{k,\ell}$. Consider the following short exact sequence

$$0 \rightarrow E(\mathcal{O}) \xrightarrow{\cdot\pi} E(\mathcal{O}) \rightarrow E(\kappa_\pi) \rightarrow 0.$$

where $\cdot\pi$ is the multiplication by π map.

The associated long exact sequence gives the following short exact sequence

$$0 \rightarrow H^i(\Gamma, E(\mathcal{O})) \otimes \kappa_\pi \rightarrow H^i(\Gamma, E(\kappa_\pi)) \rightarrow H^{i+1}(\Gamma, E(\mathcal{O}))[\pi] \rightarrow 0$$

for $i \geq 0$. Here $H^j(\Gamma, E(\mathcal{O}))[\pi]$ denotes the kernel of the map induced by $\cdot\pi$.

Putting $i = 0$, we get

$$E(\kappa_\pi)^\Gamma \simeq H^1(\Gamma, E(\mathcal{O}))[\pi]$$

Now (a) and (b) follow via the irreducibility discussions of the previous section. For $i = 1$ we get

$$0 \rightarrow H^1(\Gamma, E(\mathcal{O})) \otimes \kappa_\pi \rightarrow H^1(\Gamma, E(\kappa_\pi)) \rightarrow H^2(\Gamma, E(\mathcal{O}))[\pi] \rightarrow 0$$

which explains the claim (c). It is known that the virtual cohomological dimension of a Bianchi group is 2. Setting $i = 2$, we get

$$H^2(\Gamma, E(\mathcal{O})) \otimes \kappa_\pi \simeq H^2(\Gamma, E(\kappa_\pi)),$$

finishing the proof. \square

Each cohomology space comes equipped with a commuting family \mathbb{T} of Hecke operators acting on it, see [ST]. A (*Hecke*) *eigenvalue system* with values in a ring R is a ring homomorphism $\Phi : \mathbb{T} \rightarrow R$. We say that an eigenvalue system Φ occurs in an $R\mathbb{T}$ -module A if there is a nonzero element $a \in A$ such that $Ta = \Phi(T)a$ for all T in \mathbb{T} . Using a lifting theorem of Ash and Stevens ([AS], Prop.1.2.2), we can lift an eigenvalue systems occurring in $H^2(\Gamma, E(\kappa_\pi))$ to one occurring in $H^2(\Gamma, E(\mathcal{R}))$ where \mathcal{R} is some finite extension of the completion of \mathcal{O} at π . The possible p -torsion in $H^2(\Gamma, E(\mathcal{O}))$ obstructs us from applying the lifting theorem to lift eigenvalue systems occurring in $H^1(\Gamma, E(\kappa_\pi))$.

3.1. Cuspidal cohomology

There is a subspace of the cohomology that is of special interest due to the fact that it can be identified with cuspidal Bianchi modular forms.

Let $K = \mathbb{Q}(\sqrt{-d})$ be an imaginary quadratic field of class number h_K with ring of integers $\mathcal{O} = \mathcal{O}_d$. Let \mathbb{P} denote the projective line over K and $G = \mathrm{PSL}_2(\mathcal{O})$. The group $\mathrm{PSL}_2(K)$ acts naturally on K^2 and thus on \mathbb{P} . It is well-known that the cardinality $|\mathbb{P}/G|$ of the set \mathbb{P}/G of G -orbits of \mathbb{P} is equal to h_K . Hence $|\mathbb{P}/\Gamma|$ is finite for any finite index subgroup Γ of G . We will call the elements of \mathbb{P}/Γ the *cusps* of Γ .

For every $D \in \mathbb{P}$, let B_D be the Borel subgroup of G defined by the (setwise) stabilizer of D in G . Then the pointwise stabilizer of D in G is the unipotent radical U_D of the Borel subgroup B_D . Let Γ be a finite index subgroup of G and D_c be a representative for a cusp c of Γ . Define

$$\Gamma_c := B_{D_c} \cap \Gamma.$$

If Γ_c is torsion-free (this is automatic if Γ is itself torsion-free or $-d \neq 1, 3$), then $\Gamma_c = U_{D_c} \cap \Gamma$ and Γ_c is free abelian of rank two (see [S1] p.507). The group

$$U(\Gamma) := \bigoplus_{c \in \mathbb{P}/\Gamma} \Gamma_c$$

is independent of the choice of representatives taken for the cusps of Γ .

Let E be a Γ -module. Consider the long exact sequence of relative group cohomology for the pair $(\Gamma, U(\Gamma))$

$$\dots \rightarrow H_c^{i-1}(\Gamma, E) \rightarrow H^i(\Gamma, E) \rightarrow H^i(U(\Gamma), E) \rightarrow \dots$$

where $H_c^n(\Gamma, E) := H^n(\Gamma; U(\Gamma), E)$ and the third arrow is given by the restriction maps.

Definition 3.3. The *cuspidal cohomology* $H_{cusp}^i(\Gamma, E)$ is defined as the image of the cohomology with compact support in $H^i(\Gamma, E)$, or equivalently as the kernel of the restriction map $H^i(\Gamma, E) \rightarrow H^i(U(\Gamma), E)$.

Remark 3.4. Let $S_{k,\ell}(\Gamma)$ denote the space of cuspidal Bianchi modular forms with level Γ and weight (k, ℓ) . Harder proved in [Ha] the so called Eichler-Shimura-Harder isomorphism

$$S_{k,\ell}(\Gamma) \simeq H_{cusp}^1(\Gamma, E_{k,\ell}(\mathbb{C})) \simeq H_{cusp}^2(\Gamma, E_{k,\ell}(\mathbb{C}))$$

of Hecke modules. Note that the second isomorphism is an instance of a duality result which says that if F is a field in which 6 is invertible, then

$$H_{cusp}^1(\Gamma, E(F))^\vee \simeq H_{cusp}^2(\Gamma, E(F)^\vee)$$

as Hecke modules where $-\vee$ denotes the dual, see [AS] Lemma 1.4.3.

Deep results of Borel and Wallach (see Section II of their book [BW]) imply that whenever $k \neq \ell$, the cuspidal cohomology $H_{cusp}^i(\Gamma, E_{k,\ell}(\mathbb{C}))$ vanishes. It is important to remark that this is not true anymore when the module $E_{k,\ell}$ is not over a field of characteristic 0. In particular, $H_{cusp}^i(\Gamma, E_{k,\ell}(\mathcal{O}))$ is completely torsion when $k \neq \ell$.

Proposition 3.5. *Let Γ be a finite index subgroup of the Bianchi group $\mathrm{PSL}_2(\mathcal{O})$. Assume either that Γ is torsion-free or that $-d \neq 1, 3$. Then $H^2(U(\Gamma), E_{k,\ell}(\mathcal{O}))$ has no large torsion.*

Proof. It is enough to prove the claim for a single cusp c of Γ , that is, for $H^2(\Gamma_c, E)$. So fix a cusp c and Γ_c . Let $E = E_{k,\ell}$ and $t = \max\{k, \ell\}$. Put $R = \mathcal{O}[\frac{1}{t}]$. Composition of the cup product and the perfect pairing of Proposition 2.1 gives us a pairing

$$\begin{array}{ccc} H^0(\Gamma_c, E(R)) \times H^2(\Gamma_c, E(R)) & \xrightarrow{\cup} & H^2(\Gamma_c, E(R)) \otimes_R E(R) \\ & & \downarrow (\cdot, \cdot) \\ & & H^2(\Gamma_c, R) \simeq R. \end{array}$$

That $H^2(\Gamma_c, R) \simeq R$ can be shown as follows. Recall that Γ_c is free abelian with two generators, say a, u . It is known, see [M] p.188, that the tensor product of the two resolutions

$$0 \longrightarrow R[\langle a \rangle] \xrightarrow{1-a} R[\langle a \rangle] \xrightarrow{\varepsilon} R \longrightarrow 0,$$

$$0 \longrightarrow R[\langle u \rangle] \xrightarrow{1-u} R[\langle u \rangle] \xrightarrow{\varepsilon} R \longrightarrow 0,$$

where ε is the usual augmentation map, gives a resolution of Γ_c . One sees from this resolution that the second cohomology of Γ_c with any (right) R -module M can be described as

$$H^2(\Gamma_c, M) \simeq M / (M(1-a) + M(1-u)).$$

In the case of trivial module R , it follows immediately that $H^2(\Gamma_c, R) \simeq R$.

The above pairing gives that $H^2(\Gamma_c, E(R)) \simeq H^0(\Gamma_c, E(R))^\vee$. Clearly $H^0(\Gamma_c, E(R)) \simeq E(R)^{\Gamma_c}$ is torsion-free. This implies that its dual and hence $H^2(\Gamma_c, E(R))$ is torsion-free. The claim that there can only be small torsion in $H^2(\Gamma_c, E(\mathcal{O}))$ now follows as $R = \mathcal{O}[\frac{1}{t}]$. \square

As a corollary we see that the cuspidal part of H^2 is responsible for the possible large torsion. The referee brought to my attention that the analogue of the above result for $H^1(U(\Gamma), E_{k,k}(\mathcal{O}))$ was proven by Urban in [U] Prop.2.4.1. This result of Urban similarly implies that the possible large torsion in $H_c^2(\Gamma, E_{k,k}(\mathcal{O}))$ comes from $H_{cusp}^2(\Gamma, E_{k,k}(\mathcal{O}))$ as well.

Proposition 3.6. *Let G be the Bianchi group $\mathrm{PSL}_2(\mathcal{O})$. Let π be prime element of \mathcal{O} over the rational prime p and put κ_π for its residue field. Let Γ be a torsion-free finite index subgroup of G . Then*

$$H_{cusp}^2(\Gamma, E_{k,\ell}(\mathcal{O})) \otimes \kappa_\pi \simeq H_{cusp}^2(\Gamma, E_{k,\ell}(\kappa_\pi))$$

for every $k, \ell < p$.

Proof. Put $E = E_{k,\ell}(\mathcal{O})$. Now consider the commutative diagram

$$\begin{array}{ccccccc} H^2(\Gamma, E) & \xrightarrow{\cdot\pi} & H^2(\Gamma, E) & \longrightarrow & H^2(\Gamma, E(\kappa_\pi)) & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ 0 \longrightarrow & H^2(U(\Gamma), E) & \xrightarrow{\cdot\pi} & H^2(U(\Gamma), E) & \longrightarrow & H^2(U(\Gamma), E(\kappa_\pi)) & \end{array}$$

Here the vertical maps are given by the usual restriction maps.

The horizontal lines are exact. The exactness of the first line comes from Proposition 3.2 part (d). The exactness of the second line amounts to Proposition 3.5.

Observe that the cokernel of the restriction map $H^2(\Gamma, E) \rightarrow H^2(U(\Gamma), E)$ is isomorphic to $H_{cusp}^3(\Gamma, E) \subset H^3(\Gamma, E)$. Since the virtual cohomological dimension of a Bianchi group is two and Γ is torsion-free, we have $H^3(\Gamma, E) = 0$. Now the claim follows by the Snake Lemma. \square

Let us end this section with the following observation on lifting eigenvalue systems.

Proposition 3.7. *Let $\pi \in \mathcal{O}$ be a prime element over the rational prime $p > 3$. Let Γ be a torsion-free finite index subgroup of $\mathrm{PSL}_2(\mathcal{O})$. Let Φ be an eigenvalue system occurring in $H_{cusp}^1(\Gamma, E_{k,\ell}(\kappa_\pi))$ with $k, \ell < p$ and $\kappa_\pi = \mathcal{O}/(\pi)$. If Φ does not lift to $H_{cusp}^1(\Gamma, E_{k,\ell}(\mathcal{R}))$ for any finite extension \mathcal{R} of the completion \mathcal{O}_π of \mathcal{O} at π , then there is a π -torsion eigenclass $c \in H_{cusp}^2(\Gamma, E_{k,\ell}(\mathcal{O}))$ realizing a lift of Φ .*

Proof. As $p > 3$, by the duality result I mentioned in Remark 3.4, we deduce that Φ^\vee lives in $H_{cusp}^2(\Gamma, E_{k,\ell}(\kappa_\pi))$. Note that our coefficient modules are self-dual. Since $k, \ell < p$, using Proposition 3.6 and the lifting theorem of Ash and Stevens mentioned after Prop.3.2, we infer that there is an eigenvalue system Ψ living in $H_{cusp}^2(\Gamma, E_{k,\ell}(\mathcal{R}))$ lifting Φ^\vee for some finite extension \mathcal{R} of \mathcal{O}_π . If Ψ is not realized by a torsion eigenclass $c \in H_{cusp}^2(\Gamma, E_{k,\ell}(\mathcal{R}))$, then we can realize Ψ in $H_{cusp}^2(\Gamma, E_{k,\ell}(L))$ where L is the field of fractions of \mathcal{R} . By duality again, Ψ^\vee occurs in $H_{cusp}^1(\Gamma, E_{k,\ell}(L))$. As Ψ^\vee has integral values, it can be realized in $H_{cusp}^1(\Gamma, E_{k,\ell}(\mathcal{R}))$. Clearly Ψ^\vee is a lift of Φ and this contradicts our starting assumption of non-liftability. To finish, observe that $H_{cusp}^i(\Gamma, E_{k,\ell}(\mathcal{R})) \simeq H_{cusp}^i(\Gamma, E_{k,\ell}(\mathcal{O})) \otimes_{\mathcal{O}} \mathcal{R}$. \square

4. First cohomology

I will now describe a method, first observed by Fox [F], that allows us to compute H^1 of any finitely presented group with coefficients in a finite dimensional module. It is well known that Bianchi groups are finitely presented. Presentations for many Bianchi groups are in the literature, see, for example, [FGT].

Let me illustrate the method through an example. A formal exposition is contained in [FGT]. Let $w = \sqrt{-2}$ and $G = \mathrm{PSL}_2(\mathbb{Z}[w])$. It is known that

$$G = \langle A, B, U \mid B^2 = (AB)^3 = [A, U] = (BU^2BU^{-1})^2 = 1 \rangle$$

where A, B, U can be realized as $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix}$ respectively.

Let E be any G -module. Given any cocycle $f : G \rightarrow E$, any value $f(X)$ can be expressed linearly in terms of the images $f(A), f(B), f(U)$ of the generators of G , e.g.

$$f(ABU) = f(A) \cdot BU + f(B) \cdot U + f(U).$$

Moreover, $f(A), f(B), f(U)$ satisfy the linear equations coming from the relations of the presentation. For example,

$$B^2 = 1 \implies f(B)(B + 1) = 0.$$

Conversely, any pair $(x, y, z) \in E^3$ satisfying the linear equations coming from the presentation gives uniquely a cocycle. Thus the space of cocycles can be seen as the kernel of the matrix corresponding to this linear system. One gets the coboundaries similarly and hence computes $H^1(G, E)$ as the quotient of the two spaces.

Note that to compute with a finite index subgroup Γ of G , it is not practical to apply the method to a presentation of Γ (which can be derived from that of G once the coset representatives are known). It is best to use Shapiro's lemma and compute $H^1(G, \mathrm{Coind}_\Gamma^G(E))$.

4.1. Data on the integral first cohomology

I have implemented the above algorithm in MAGMA [BCP] for the five Euclidean imaginary quadratic fields $K = \mathbb{Q}(\sqrt{-d})$ with $d = 1, 2, 3, 7, 11$. In the following, let \mathcal{O}_d denote the corresponding ring of integers.

By the theory of modules over principal ideal domains, we know that our \mathcal{O} -module $H^1(\Gamma, E(\mathcal{O}))$ has a decomposition

$$H^1(\Gamma, E(\mathcal{O})) \simeq \mathcal{O}/(a_1) \oplus \dots \oplus \mathcal{O}/(a_m) \oplus \mathcal{O}^r.$$

with $a_i \neq 0, 1$ and $a_i | a_{i+1}$. The a_i are called elementary divisors and are unique up to multiplication by units. The exponent r is called the rank.

Below I report on some of my computations. Observe that the torsion is always "small" as proved in Proposition 3.2. The only exception to this is the ramifying prime which always appears in the torsion. We show the rank in a separate column as it provides a means to check our work against the dimension computations of [FGT].

data for $H^1(\mathrm{PSL}_2(\mathcal{O}_1), E_{n,n}(\mathcal{O}_1))$

n	norms of elt. divisors	primes	rank
0			0
1	[4]	(2)	1
2	[2, 16]	(2)	0
3	[2, 2, 4]	(2)	1
4	[2, 2, 2, 8, 1152]	(2,3)	0
5	[2, 2, 2, 2, 4, 4]	(2)	2
6	[2, 2, 2, 2, 2, 2, 8, 8, 800]	(2,5)	0
7	[2, 2, 2, 2, 2, 2, 2, 2, 4, 4, 4]	(2)	3
8	[2, 2, 2, 2, 2, 2, 2, 2, 2, 4, 8, 8, 32, 225792]	(2,3,7)	0
9	[2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 4, 4, 4, 4, 4]	(2)	3
10	[2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 8, 8, 8, 8, 16, 288]	(2,3)	1

data for $H^1(\mathrm{PSL}_2(\mathcal{O}_2), E_{n,n}(\mathcal{O}_2))$

n	norms of elt. divisors	primes	rank
1	[8]	(2)	1
2	[2, 32]	(2)	1
3	[2, 2, 8]	(2)	2
4	[2, 2, 2, 8, 1152]	(2,3)	1
5	[2, 2, 2, 2, 8, 8]	(2)	3
6	[2, 2, 2, 2, 2, 2, 8, 8, 7200]	(2,3,5)	2
7	[2, 2, 2, 2, 2, 2, 2, 2, 8, 8, 8]	(2)	4
8	[2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 4, 8, 8, 32, 225792]	(2,3,7)	2
9	[2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 8, 8, 8, 8, 8, 8]	(2)	5
10	[2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 8, 8, 8, 8, 8, 32, 288]	(2,3)	3

data for $H^1(\mathrm{PSL}_2(\mathcal{O}_3), E_{n,n}(\mathcal{O}_3))$

n	norms of elt. divisors	primes	rank
0			0
1	[3]	(3)	0
2	[3]	(3)	1
3	[3, 108]	(2,3)	0
4	[3, 3, 12]	(2,3)	0
5	[3, 3, 12]	(2,3)	1
6	[3, 3, 3, 3, 10800]	(2,3,5)	1
7	[3, 3, 3, 3, 3, 12]	(2,3)	1
8	[3, 3, 3, 3, 3, 12, 2352]	(2,3,7)	1
9	[3, 3, 3, 3, 3, 3, 3, 108, 972]	(2,3)	1

5. Second cohomology

The main method I employ for computing the second cohomology is based on reduction theory as used in [SV]. The cohomological dimension of Bianchi groups is 2 and the symmetric space they act on, namely the hyperbolic 3-space $\mathbb{H} \simeq \mathbb{C} \times \mathbb{R}^+$, is 3 dimensional. Reduction theory gives us a contractible 2 dimensional CW-complex inside \mathbb{H} which is a deformation retract for the action of the Bianchi group. Moreover, the cellular action of the Bianchi group on the CW-complex is cocompact. This makes the CW-complex a suitable tool for cohomological computations.

I will continue to focus on the Euclidean imaginary quadratic fields. The reduction theory for Bianchi groups has been worked out for these fields by Mendoza [Me] and Flöge [Fl]. See also [BLW, RF].


For an overview of Mendoza's construction, I refer readers to [SV]. I will exhibit the method for the case of the Bianchi group $\Gamma = \mathrm{PSL}_2(\mathbb{Z}[w])$ with $w = \sqrt{-2}$.

Let \mathcal{C} be the 2-dimensional CW-complex constructed by Mendoza for Γ . Then a fundamental cellular domain \mathcal{F} for the action of Γ on \mathcal{C} is given by the area on the unit hemisphere centered at the origin of \mathbb{H} above the rectangle in $\mathbb{C} \times \{0\}$ with vertices $(\pm \frac{w}{2}, 0)$ and $(\frac{1}{2} \pm \frac{w}{2}, 0)$.

Let

$$a := \begin{pmatrix} 1 & w \\ w & -1 \end{pmatrix}, \quad b := \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}, \quad c := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

The stabilizers of the edges (1-cells) and the vertices (0-cells) of \mathcal{F} are shown in the following picture



The horizontal edges are identified by the element $g = \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix}$, that is $gP_1P_2 = P_4P_3$. Thus the quotient by Γ is a cylinder. Moreover, the stabilizer of the whole rectangle (2-cell) is trivial.

From this combinatorial data, one can compute the (co)homology. One way to do this is to feed the data into the equivariant cohomology spectral sequence

$$E_1^{p,q}(M) = \bigoplus_{\sigma \in \Sigma_p} H^q(\Gamma_\sigma, M) \implies H^{p+q}(\Gamma, M).$$

where M is any $\mathbb{Z}\Gamma$ -module and Σ_p is a set of representatives of all the Γ -orbits of the p -cells of \mathcal{C} . See page 164 of [Br] for a description. The homological version of

this spectral sequence has been used in [SV], [RF]. See Section 10 of [Ya] for another method to extract the same information.

Let Γ_i, Γ_{ij} stand for the stabilizers of the vertex P_i and the edge between P_i and P_j respectively. Let M be a right Γ -module over $\mathbb{Z}[w][\frac{1}{6}]$. As primes above 2 and 3 are inverted, the cohomology of the (finite) stabilizers vanish in degree greater than 0. Hence, we have $E_1^{p,q}(M) = 0$ for all $q > 0$. Therefore, the spectral sequence is concentrated on the horizontal axis $q = 0$ and the cohomology of the cochain complex

$$E_1^{0,0} \xrightarrow{d_1^{0,0}} E_1^{1,0}(M) \xrightarrow{d_1^{1,0}} E_1^{2,0}(M)$$

gives $H^*(\Gamma, M)$, that is

$$H^0(\Gamma, M) = \text{Ker}(d_1^{0,0}) \quad H^1(\Gamma, M) = \text{Ker}(d_1^{1,0})/\text{Im}(d_1^{0,0}), \quad H^2(\Gamma, M) = M/\text{Im}(d_1^{1,0}).$$

Now with the appropriate substitutions, the cochain complex reads

$$\bigoplus_{\text{vertex } i} H^0(\Gamma_i, M) \xrightarrow{d_1^{0,0}} \bigoplus_{\text{edge } ij} H^0(\Gamma_{ij}, M) \xrightarrow{d_1^{1,0}} H^0(\langle Id \rangle, M).$$

Here $\langle Id \rangle$ is the trivial stabilizer of the 2-cell \mathcal{F} .

To explicitly compute H^2 , it remains to describe the differential $d_1^{1,0}$. One can choose the orientation on \mathcal{F} so that the differential map becomes as follows

$$M^{\Gamma_1} \oplus M^{\Gamma_2} \xrightarrow{d_1^{0,0}} M^{\Gamma_{12}} \oplus M^{\Gamma_{23}} \oplus M^{\Gamma_{41}} \xrightarrow{d_1^{1,0}} M.$$

$$d_1^{1,0}(m_{12}, m_{23}, m_{41}) = m_{12} + m_{23} + m_{41} - m_{12} \cdot g^{-1}$$


The information on the fundamental 2-cell for the groups $\text{PSL}_2(\mathcal{O}_d)$ with $-d = 1, 2, 3, 7, 11$ is included in the article [SV], so I do not repeat it here. The same information for the groups $\text{PGL}_2(\mathcal{O}_d)$ with $-d = 1, 2, 3, 7, 11$ is not included in that article and one needs to go to the above mentioned thesis of Mendoza (although a few of them are contained in [BLW] as well) which is very hard to access from outside of Germany. So I will now describe the information on these groups in the pictorial form as above.

- $\text{PGL}_2(\mathcal{O}_1)$

Put $i = \sqrt{-1}$. Let

$$a := \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad b := \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}, \quad c := \begin{pmatrix} 0 & i \\ 1 & 0 \end{pmatrix}.$$

The stabilizers of the edges and the vertices of \mathcal{F} are shown in the following picture




There are no identifications and the stabilizer of the triangle (2-cell) is trivial.

- $\mathrm{PGL}_2(\mathcal{O}_2)$

Put $w = \sqrt{-2}$. Let

$$a := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad b := \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}, \quad c := \begin{pmatrix} w & 1 \\ 1 & 0 \end{pmatrix}, \quad d := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

The stabilizers of the edges and the vertices of \mathcal{F} are shown in the following picture




There are no identifications and the stabilizer of the rectangle (2-cell) is trivial.

- $\mathrm{PGL}_2(\mathcal{O}_3)$

Put $w = \frac{1+\sqrt{-3}}{2}$. Let

$$a := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad b := \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}, \quad c := \begin{pmatrix} 0 & w \\ 1 & 0 \end{pmatrix}.$$

The stabilizers of the edges and the vertices of \mathcal{F} are shown in the following picture




There are no identifications and the stabilizer of the triangle (2-cell) is trivial.

- $\mathrm{PGL}_2(\mathcal{O}_7)$

Put $w = \frac{1+\sqrt{-7}}{2}$. Let

$$a := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad b := \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}, \quad c := \begin{pmatrix} 1 & -w \\ w & -1 \end{pmatrix}, \quad d := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

The stabilizers of the edges and the vertices of \mathcal{F} are shown in the following picture




The two adjacent short edges on the top are identified via $g = \begin{pmatrix} 1 & -w \\ 0 & -1 \end{pmatrix}$ which fixes the vertex between them. Thus these two edges are oppositely oriented and the stabilizer of the vertex between them is $\mathbf{D}_2 \simeq \langle c, g \rangle$. Again the stabilizer of the whole 2-cell is trivial.

- $\mathrm{PGL}_2(\mathcal{O}_{11})$

Put $w = \frac{1+\sqrt{-11}}{2}$. Let

$$a := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad b := \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}, \quad c := \begin{pmatrix} 1 & -w \\ w & -2 \end{pmatrix}, \quad d := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

The stabilizers of the edges and the vertices of \mathcal{F} are shown in the following picture



The two adjacent short edges on the top are identified via $g = \begin{pmatrix} 1 & -w \\ 0 & -1 \end{pmatrix}$ which fixes the vertex between them. Thus these two edges are oppositely oriented and the stabilizer of the vertex between them is $\mathbf{S}_3 \simeq \langle c, g \rangle$. Again the stabilizer of the whole 2-cell is trivial.

5.1. data on the integral second cohomology: level 1

I have implemented the above algorithm in MAGMA. I have not inverted the primes above 2,3 and thus the computations may not give correct data on 2,3 torsion. Below I give a complete list of the primes that appear in the torsion part of the second cohomology of both PSL and PGL. The large primes are highlighted in bold face.

The data imply that if (p) ramifies in \mathcal{O} then there is p -torsion in the integral second cohomology (except the case $k = \ell = 0$) but I have not been able to prove that this is always the case.

$$H^2(\mathrm{PSL}_2(\mathcal{O}_1), E_{n,n}(\mathcal{O}_1))$$

n	primes	rank
1	[]	1
2	[2]	1
3	[2, 3]	1
4	[2, 3]	1
5	[2]	2
6	[2, 3, 5]	1
7	[2, 3, 7]	3
8	[2, 3, 5, 7]	1
9	[2, 3]	3
10	[2, 3, 5, 7]	2
11	[2, 3, 5, 11]	4
12	[2, 3, 5, 7, 11]	1
13	[2, 3, 5]	5
14	[2, 3, 5, 7, 11, 13]	2
15	[2, 3, 5, 7]	5
16	[2, 3, 5, 7, 11, 13]	2
17	[2, 3, 5, 7]	6
18	[2, 3, 5, 7, 11, 13, 17, 19 , 23]	2
19	[2, 3, 5, 7, 13, 19]	7
20	[2, 3, 5, 7, 11, 13, 17, 19, 409 , 6997]	2
21	[2, 3, 5, 7, 59]	7
22	[2, 3, 5, 7, 11, 13, 17, 19, 13707791]	3
23	[2, 3, 5, 7, 11, 23 , 113]	8
24	[2, 3, 5, 7, 11, 13, 17, 19, 23, 1033 , 4457 , 18743]	2
25	[2, 3, 5, 7, 11, 13, 17, 1523]	9

$$H^2(\mathrm{PGL}_2(\mathcal{O}_1), E_{n,n}(\mathcal{O}_1))$$

n	primes	rank
1	[]	1
2	[2]	1
3	[2, 3]	1
4	[2]	1
5	[2]	2
6	[2, 3, 5]	1
7	[2, 3, 7]	2
8	[2, 3, 5, 7]	1
9	[2, 3]	3
10	[2, 3, 5, 7]	2
11	[2, 3, 11]	3
12	[2, 3, 5, 11]	1
13	[2, 3, 5]	4
14	[2, 3, 5, 7, 11, 13]	2
15	[2, 3, 5, 7]	4
16	[2, 3, 5, 7, 13]	2
17	[2, 3, 5, 7]	5
18	[2, 3, 5, 7, 11, 17]	2
19	[2, 3, 5, 19]	5
20	[2, 3, 5, 7, 13, 17, 19, 409]	2
21	[2, 3, 5, 7]	6
22	[2, 3, 5, 7, 11, 19]	3
23	[2, 3, 5, 7, 11, 23]	6
24	[2, 3, 5, 7, 11, 13, 17, 23, 1033]	2
25	[2, 3, 5, 7, 11, 17]	7
26	[2, 3, 5, 7, 11, 13, 19, 23, 157, 683]	3
27	[2, 3, 5, 7]	7
28	[2, 3, 5, 7, 11, 13, 17, 664197637]	3
29	[2, 3, 5, 7, 11, 13, 89]	8
30	[2, 3, 5, 7, 11, 13, 19, 23, 29, 211, 36312691]	3

$$H^2(\mathrm{PSL}_2(\mathcal{O}_2), E_{n,n}(\mathcal{O}_2))$$

n	primes	rank
1	[]	1
2	[2]	1
3	[2, 3]	2
4	[2, 3]	1
5	[2, 3, 5]	3
6	[2, 3, 5]	2
7	[2, 3, 5, 7]	4
8	[2, 3, 5, 7]	2
9	[2, 3, 5, 7, 31]	5
10	[2, 3, 5, 7]	3
11	[2, 3, 5, 7, 11]	6
12	[2, 3, 5, 7, 11, 37]	3
13	[2, 3, 5, 7, 11, 13, 547]	7
14	[2, 3, 5, 7, 11, 13, 439, 110281]	4
15	[2, 3, 5, 7, 11, 13, 61, 163]	8

$$H^2(\mathrm{PGL}_2(\mathcal{O}_2), E_{n,n}(\mathcal{O}_2))$$

n	primes	rank
1	[]	1
2	[2]	1
3	[2]	2
4	[2, 3]	1
5	[2, 5]	3
6	[2, 3, 5]	1
7	[2, 3, 7]	4
8	[2, 3, 5, 7]	1
9	[2, 3]	5
10	[2, 3, 5, 7]	2
11	[2, 3, 5]	6
12	[2, 3, 5, 7, 11, 37]	1
13	[2, 3, 5, 13]	7
14	[2, 3, 5, 7, 11, 13, 110281]	2
15	[2, 3, 5, 7]	8
16	[2, 3, 5, 7, 11, 13, 1671337]	2
17	[2, 3, 5, 7, 103]	9
18	[2, 3, 5, 7, 11, 13, 17, 3812807473]	2
19	[2, 3, 5, 7, 907]	10
20	[2, 3, 5, 7, 11, 13, 17, 19, 3511, 879556698451244053]	2

$$H^2(\mathrm{PSL}_2(\mathcal{O}_3), E_{n,n}(\mathcal{O}_3))$$

n	primes	rank
1	[]	1
2	[3]	1
3	[3]	1
4	[2, 3]	1
5	[2, 3, 5]	1
6	[2, 3, 5]	2
7	[2, 3, 5]	2
8	[2, 3, 7]	1
9	[2, 3, 7]	2
10	[2, 3, 5]	3
11	[2, 3, 5, 11]	2
12	[2, 3, 5, 7, 11]	2
13	[2, 3, 5, 11]	3
14	[2, 3, 5, 7, 13]	3
15	[2, 3, 5, 7, 13]	3
16	[2, 3, 5, 7, 11]	3
17	[2, 3, 5, 7, 17]	3
18	[2, 3, 5, 7, 13, 17]	4
19	[2, 3, 5, 7, 11, 17, 61]	4
20	[2, 3, 5, 7, 19]	3
21	[2, 3, 5, 7, 13, 19, 151]	4
22	[2, 3, 5, 7, 11, 17]	5
23	[2, 3, 5, 7, 11, 23, 103]	4
24	[2, 3, 5, 7, 11, 13, 17, 19, 23, 53]	4
25	[2, 3, 5, 7, 11, 17, 23, 29, 947]	5
26	[2, 3, 5, 7, 11, 13]	5

$$H^2(\mathrm{PGL}_2(\mathcal{O}_3), E_{n,n}(\mathcal{O}_3))$$

n	primes	rank
1	[]	1
2	[]	1
3	[2, 3]	1
4	[2, 3]	1
5	[2, 3, 5]	1
6	[2, 3, 5]	1
7	[2, 3]	2
8	[2, 3, 7]	1
9	[2, 3]	2
10	[2, 3, 5]	2
11	[2, 3, 5, 11]	2
12	[2, 3, 5, 7, 11]	1
13	[2, 3]	3
14	[2, 3, 5, 7, 13]	2
15	[2, 3, 5]	3
16	[2, 3, 5, 7, 11]	2
17	[2, 3, 5, 17]	3
18	[2, 3, 5, 7, 13, 17]	2
19	[2, 3, 5, 7]	4
20	[2, 3, 5, 7, 19]	2
21	[2, 3, 5, 7]	4
22	[2, 3, 5, 7, 11, 17]	3
23	[2, 3, 5, 11, 23]	4
24	[2, 3, 5, 7, 11, 13, 19, 23, 53]	2
25	[2, 3, 5, 7]	5
26	[2, 3, 5, 7, 11, 13]	3
27	[2, 3, 5, 7, 11]	5
28	[2, 3, 5, 7, 11, 13, 17, 23]	3
29	[2, 3, 5, 7, 29]	5

$$H^2(\mathrm{PSL}_2(\mathcal{O}_7), E_{n,n}(\mathcal{O}_7))$$

n	primes	rank
1	[]	1
2	[2, 7]	1
3	[2, 3, 7]	1
4	[2, 3, 7]	2
5	[2, 3, 5, 7]	2
6	[2, 3, 5, 7]	2
7	[2, 3, 5, 7]	3
8	[2, 3, 5, 7]	3
9	[2, 3, 5, 7]	3
10	[2, 3, 5, 7]	4
11	[2, 3, 5, 7, 11]	4
12	[2, 3, 5, 7, 11, 127]	6
13	[2, 3, 5, 7, 11, 13, 31]	5
14	[2, 3, 5, 7, 11, 13, 73]	5
15	[2, 3, 5, 7, 11, 13, 271, 431]	5
16	[2, 3, 5, 7, 11, 13]	6
17	[2, 3, 5, 7, 11, 13, 17, 37, 67, 89, 101, 277]	6
18	[2, 3, 5, 7, 11, 13, 17, 43, 457, 2069, 3323]	6

$$H^2(\mathrm{PGL}_2(\mathcal{O}_7), E_{n,n}(\mathcal{O}_7))$$

n	primes	rank
1	[]	1
2	[2]	1
3	[2, 3, 7]	1
4	[2, 3, 7]	1
5	[2, 5, 7]	2
6	[2, 3, 5, 7]	1
7	[2, 3, 7]	3
8	[2, 3, 5, 7]	1
9	[2, 3, 7]	3
10	[2, 3, 5, 7]	2
11	[2, 3, 5, 7]	4
12	[2, 3, 5, 7, 11, 127]	1
13	[2, 3, 5, 7, 13]	5
14	[2, 3, 5, 7, 11, 13, 73]	2
15	[2, 3, 5, 7, 431]	5
16	[2, 3, 5, 7, 11, 13]	2
17	[2, 3, 5, 7, 17, 37]	6
18	[2, 3, 5, 7, 11, 13, 17, 43, 457, 2069, 3323]	2
19	[2, 3, 5, 7, 13, 19, 311]	7
20	[2, 3, 5, 7, 11, 13, 17, 19, 42197, 12272815271]	2

$$H^2(\mathrm{PSL}_2(\mathcal{O}_{11}), E_{n,n}(\mathcal{O}_{11}))$$

n	primes	rank
1	[]	1
2	[2]	2
3	[2, 3, 11]	2
4	[2, 3, 11]	2
5	[2, 3, 5, 11]	3
6	[2, 3, 5, 11]	4
7	[2, 3, 5, 7, 11]	4
8	[2, 3, 5, 7, 11]	4
9	[2, 3, 5, 7, 11, 23]	5
10	[2, 3, 5, 7, 11]	8
11	[2, 3, 5, 7, 11, 37]	6
12	[2, 3, 5, 7, 11]	6
13	[2, 3, 5, 7, 11, 13, 43, 19973]	7
14	[2, 3, 5, 7, 11, 13]	8
15	[2, 3, 5, 7, 11, 13, 31, 47, 1409, 30817]	8
16	[2, 3, 5, 7, 11, 13, 17, 19, 41, 281]	8

$$H^2(\mathrm{PGL}_2(\mathcal{O}_{11}), E_{n,n}(\mathcal{O}_{11}))$$

n	primes	rank
1	[]	1
2	[2]	1
3	[2]	2
4	[2, 3, 11]	1
5	[2, 11]	3
6	[2, 3, 5, 11]	1
7	[2, 3, 7, 11]	4
8	[2, 3, 5, 7, 11]	1
9	[2, 3, 11]	5
10	[2, 3, 5, 7, 11]	4
11	[2, 3, 5, 11]	6
12	[2, 3, 5, 7, 11]	1
13	[2, 3, 5, 11, 13]	7
14	[2, 3, 5, 7, 11, 13]	2
15	[2, 3, 5, 7, 11, 47]	8
16	[2, 3, 5, 7, 11, 13, 41, 281]	2
17	[2, 3, 5, 7, 11, 17, 67]	9
18	[2, 3, 5, 7, 11, 13, 17, 449, 20147, 201797]	2

5.2. Congruence subgroups

Now let us focus on the second cohomology of congruence subgroups. For computational considerations, we will focus on the subgroups of the type

$$\Gamma_0(\mathfrak{a}) : \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G : c \equiv 0 \pmod{\mathfrak{a}} \right\}.$$

Here G is the Bianchi group $\mathrm{PSL}_2(\mathcal{O}_d)$ and \mathfrak{a} is an ideal of \mathcal{O}_d , which is called the level.

5.2.1. Trivial weight: torsion

To compute the second cohomology with trivial weight, the approach employed in [EGM] is more efficient than the reduction theory approach used above. The idea is to compute the abelianization of the congruence subgroup Γ using a (finite) presentation for the Bianchi group G and the knowledge of the permutation action of the generators of G on a set of coset representatives of Γ in G .

The relationship between first homology and the second cohomology is given by the Lefschetz duality. Let R be any module in which 6 is invertible. Then for any $R[\Gamma]$ -module E , we have

$$H_1(\Gamma, E) \simeq H_c^2(\Gamma, E)$$

where the right hand side is the cohomology with compact support, see Section 2.

We are only interested in the case of the trivial coefficients $R = \mathbb{Z}[1/6]$. We need to study the exact sequence

$$H^1(\Gamma, R) \rightarrow H^1(U(\Gamma), R) \rightarrow H_c^2(\Gamma, R) \rightarrow H_{cusp}^2(\Gamma, R) \rightarrow 0.$$

Assume that K has class number one and that $\Gamma = \Gamma_0(\mathfrak{p})$ where \mathfrak{p} is a prime ideal of residue degree one. It is easy to see that Γ has only two cusps; $\{0, \infty\}$. That is $|\Gamma \backslash \mathbb{P}| = 2$ and the two classes are represented by the elements $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. If $-d \neq 1, 3$, then the stabilizers of the cusps are free abelian of rank 2 and by Corollaire 3 of [S1] p.517, the image of $H^1(\Gamma, R)$ has rank 2 in $H^1(U(\Gamma), R)$ which has rank 4. For $-d = 1, 3$, the situation is complicated by the existence of torsion in the stabilizers of cusps. In this case, let $\Gamma_c^+ := \Gamma \cap U_{D_c}$ for each cusp c of Γ in the terminology of Section 2.1. Then $\Gamma_c/\Gamma_c^+ \simeq \mu := \langle \begin{pmatrix} \varepsilon & 0 \\ 0 & \varepsilon^{-1} \end{pmatrix} \rangle$ where ε is a generator of the roots of unity in \mathcal{O}_d . The inflation-restriction sequence then gives

$$H^1(\Gamma_c, R) \simeq H^0(\mu, H^1(\Gamma_c^+, R)).$$

Now let us directly show that the latter is trivial. Without loss of generality, assume that $c = \infty$ and thus that $\Gamma_c^+ := \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} \cap \Gamma$. The action of μ on $H^1(\Gamma_c^+, R)$ is given as $(\tau f)(x) = f(\tau x \tau^{-1})\tau = f(\tau x \tau^{-1})$ for every $\tau \in \mu$ and every 1-cocycle $f : \Gamma_c^+ \rightarrow R$. Take $\varepsilon \in \mathcal{O}_d^*$ such that $\varepsilon^2 = -1$ and put $\tau = \begin{pmatrix} \varepsilon & 0 \\ 0 & \varepsilon^{-1} \end{pmatrix}$. Then for any element $x \in \Gamma_c^+$, $\tau x \tau^{-1} = x^{-1}$. The condition $f(x) = f(x^{-1})$ for every x forces the 1-cocycle f to be the trivial cocycle $f = 0$. We are done. An alternative way is to use the geometric approach which I mostly avoided in this paper. Then our discussion here follows from the observation that the cross section of a cusp is an orbifold with underlying manifold a sphere in the cases $-d = 1, 3$ and is a torus in the remaining cases.

I wrote programs to compute $\Gamma_0(\mathfrak{p})^{ab}$ in MAGMA for the Euclidean \mathcal{O}_d , that is, for $-d = 1, 2, 3, 7, 11$. It is easily seen that the torsion get very large very quickly compared

to the norm of the level of the congruence subgroup. Recall from the introduction that Grunewald and Schwermer speculated in [GS] that no p -torsion appearing in abelianization of a finite index subgroup Γ is greater than half of the index of Γ inside the Bianchi group. In the case of $\Gamma_0(\mathfrak{p})$, their speculation says that any p -torsion that appear in $\Gamma_0(\mathfrak{p})^{ab}$ should satisfy

$$p \leq \frac{N\mathfrak{p} + 1}{2}$$

where $N\mathfrak{p}$ is the norm of the prime ideal \mathfrak{p} .

The smallest primes that witness the falseness of this speculation for our five Bianchi groups $\mathrm{PSL}_2(\mathcal{O}_d)$ are listed below in Table 1.

d	Norm of \mathfrak{p}	rank of $\Gamma_0(\mathfrak{p})^{ab}$	prime torsion of $\Gamma_0(\mathfrak{p})^{ab}$
1	401	0	[2, 5, 41, 271]
2	193	2	[2, 3, 23, 251]
3	937	0	[2, 3, 13, 599]
7	137	2	[2, 17, 83]
11	103	2	[2, 3, 17, 19, 71]

Table 1: smallest counter examples to Grunewald-Schwermer

My computations of $\Gamma_0(\mathfrak{p})^{ab}$ agree perfectly with that of [EGM] mentioned in the introduction in the part they overlap. It is easily observed that the primes in the torsion get to astronomical sizes even within the range $N\mathfrak{p} \leq 5000$. Below in Table 2 is a sample of the primes that appear in the torsion of $\Gamma_0(\mathfrak{p})^{ab}$ with $4900 \leq N\mathfrak{p} \leq 5000$ for the five Euclidean imaginary quadratic fields. The complete list is available on my website. It is interesting to observe that in the cases $-d = 1, 3$, where we have other units besides ± 1 , the torsion grows much slower than the other three cases.

5.2.2. Trivial weight: rank

In this section I will report on the rank of $H_{cusp}^2(\Gamma_0(\mathfrak{a}), \mathcal{O})$. This rank is clearly equal to the dimension of $H_{cusp}^2(\Gamma_0(\mathfrak{a}), \mathbb{C})$ and thus its nonvanishing is conjecturally connected to abelian varieties of GL_2 -type over imaginary quadratic fields (see [Cr, EGM, Se1]). Moreover, the vanishing of this rank in certain cases is equivalent to the existence of rational homology spheres (see [LMR]). In this section I will report on my computations related to these two aspects.

As explained in the previous subsection, the rank of $\Gamma_0(\mathfrak{a})^{ab}$ is related to the rank r of $H_{cusp}^2(\Gamma_0(\mathfrak{a}), \mathcal{O})$. More precisely, when \mathfrak{p} is a prime ideal of residue degree one, our discussion above shows that

$$r = \mathrm{rank}(\Gamma_0(\mathfrak{p})^{ab}), \quad \text{for } -d = 1, 3$$

and

$$r + 2 = \mathrm{rank}(\Gamma_0(\mathfrak{p})^{ab}), \quad \text{for } -d \neq 1, 3$$

I have computed the rank of $\Gamma_0(\mathfrak{p})^{ab}$ for prime ideals \mathfrak{p} of residue degree one and norm up to 45000 for $-d = 1, 3$, 30000 for $-d = 2$, and 21000 for $-d = 7, 11$. I report on the distribution of prime levels according to the ranks in Table 3 where I use $N_r(x)$ to denote the number of primes of residue degree one with norm $< x$ and such that

Np	some of the primes that appear in the torsion of $\Gamma_0(\mathfrak{p})^{ab}$
PSL(\mathcal{O}_1)	
4909	[2, 3, 7, 13, 409, 10691, 22871, 29423, 56980673, 71143433]
4933	[2, 3, 37, 101, 137, 577, 947, 21169, 194981]
4937	[2, 7, 37, 617, 10859, 108893, 4408530403, 157824962047]
4957	[2, 3, 7, 13, 31, 59, 14347, 3051863, 9405667, 23132267]
4969	[2, 3, 23, 71, 373, 191299, 39861006443, 8672729371087]
4973	[2, 11, 13, 47, 71, 113, 127, 331, 6317, 7949, 39023, 628801, 2995319]
4993	[2, 3, 5, 7, 11, 13, 101, 173, 798569, 5995036891, 18513420749]
PSL(\mathcal{O}_2)	
4931	[... , 3772418780827, 67462419379713541, 442541106225737082232052179]
4937	[... , 1889149903, 7397090738497, 880941232181841675673769]
4969	[... , 2728733329370698225919458399, 114525595847400940348788195788260381871]
4987	[... , 1354882997352809, 167973141926075800477, 109210638303577813415629]
4993	[... , 15997185593, 14633678967206157243930187, 4844017554743814674462620193]
PSL(\mathcal{O}_3)	
4903	[3, 7, 19, 29, 37, 43, 61, 137, 191, 733]
4909	[2, 3, 7, 13, 19, 47, 67, 409, 1409]
4933	[2, 3, 5, 137, 173, 383, 719, 1451, 100057]
4951	[3, 5, 7, 11, 271, 3797, 6696049]
4957	[2, 3, 5, 7, 23, 43, 59, 233, 823, 62207]
4969	[2, 3, 5, 7, 23, 181, 2591, 516336433]
4987	[2, 3, 11, 71, 277, 619, 21977, 1971691]
4993	[2, 3, 11, 13, 29, 727, 4153, 27127]
4999	[2, 3, 7, 17, 29, 41, 83, 38593, 179623]
PSL(\mathcal{O}_7)	
4909	[... , 3354447021713, 666100957349057134013, 13363557375430202095093]
4937	[... , 836083247742263, 60001748772648369971, 1344885261548364695671]
4943	[... , 94861335404089, 157213239530981, 345644733766517, 714087340201211]
4951	[... , 42137202713, 11756096619570265637, 47745831545933513537]
4957	[... , 6803766726937001299, 21088956680308937473, 34130091188757085391]
4967	[... , 42061245937, 3414861551033731, 385786872173747641]
4993	[... , 16112554517, 22230923149, 47405513059, 17179435084786759]
4999	[... , 47183940647, 47747826462797, 176725513764138170761817312541116531]
PSL(\mathcal{O}_{11})	
4909	[... , 491602700153184794115037, 3160753948740219890398523741106925031]
4931	[... , 59242366654994144915737, 397153057377536493107457514082773]
4933	[... , 471591580131222099301009, 753357254439534230416253]
4937	[... , 774606120056702384410790118960699805738139]
4943	[... , 49685906201385872741, 7533150099701393721041, 1806172579157695730540919793]
4951	[... , 32561299447966536475490232836221, 575858582707156517384453334853901]
4973	[... , 668079334182971453623, 2223356120717452698676440064717]
4987	[... , 26685596532560442049106969671, 121708009502005164710374726093]
4999	[... , 35270997998154652004835942597708494620078410433635847]

Table 2: a modest sample of large torsion occurring in $\Gamma_0(\mathfrak{p})^{ab}$

	$d = -1$		$d = -2$		$d = -3$		$d = -7$		$d = -11$	
r	$N_r(45000)$	%	$N_r(30000)$	%	$N_r(45000)$	%	$N_r(21000)$	%	$N_r(21000)$	%
0	2061	88.8	1480	91.8	2033	87.4	1054	89.5	1056	89.7
1	177	7.6	94	5.83	184	7.91	89	7.6	96	8.15
2	66	2.8	31	1.92	82	3.52	29	2.5	22	1.90
3	10	0.4	5	0.31	21	0.91	4	0.3	2	0.17
4	4	0.2	1	0.07	5	0.22	0	0	0	0
5	1	0.04	0	0	0	0	1	0	0	0
6	1	0.04	1	0.07	0	0	0	0.1	1	0.08
7	1	0.04	0	0	1	0.04	0	0	0	0
≥ 8	0	0	0	0	0	0	0	0	0	0
> 0	260	11.2	132	8.2	293	12.6	123	10.5	121	10.3

Table 3: distribution of dimension of $\mathbb{H}_{cusp}^2(\Gamma_0(\mathfrak{p}), \mathbb{C})$

$H_{cusp}^2(\Gamma_0(\mathfrak{a}), \mathcal{O})$ has rank r . It is curious that for all five \mathcal{O}_d , approximately 90% of the time the rank was 0. Note that in [FGT], Finis, Grunewald and Tirao used an efficient method which works with finite fields and approximated the ranks up to norm 60000 for $-d = 1$. My computations for $-d = 1$ agree with theirs in the part that they overlap except that the values 12113, 12373 are missing from their Table 10 and 12941 is missing from their Table 11.

It is believed that there are infinitely many prime ideals \mathfrak{p} of residue degree one such that $H_{cusp}^2(\Gamma_0(\mathfrak{p}), \mathbb{C}) = 0$. On the other hand, in analogy with the conjecture that there are infinitely many elliptic curves over \mathbb{Q} with prime conductor, see [BS] p.97, it is reasonable to expect that there are infinitely many prime ideals \mathfrak{p} of residue degree one and $H_{cusp}^2(\Gamma_0(\mathfrak{p}), \mathbb{C}) \neq 0$. In [FGT], in analogy with the distribution questions for elliptic curves (see [BM]), the following question was posed (stated here in a slightly more general form).

Question 5.1. Let \mathcal{O}_d be the ring of integers of an imaginary quadratic number field. Is there constant C_d such that the asymptotic relation

$$\sum_{\mathfrak{p}, \mathbf{N}\mathfrak{p} \leq x} \dim H_{cusp}^1(\Gamma_0(\mathfrak{p}), \mathbb{C}) \sim C_d \frac{x^{\frac{5}{6}}}{\log x}$$

holds as x goes to infinity, where the sum ranges over residue degree one prime ideals $\mathfrak{p} \triangleleft \mathcal{O}_d$.

Let us put $L_d(x) := \sum_{\mathfrak{p}, \mathbf{N}\mathfrak{p} \leq x} \dim H_{cusp}^1(\Gamma_0(\mathfrak{p}), \mathbb{C})$ for the ring \mathcal{O}_d and $R(x) := x^{\frac{5}{6}}/\log x$. Table 4 compares the two functions $L(x)$ and $R(x)$ within the range of my computations.

In [CD], Calegari and Dunfield constructed a family of commensurable arithmetic rational homology 3-spheres, that is, commensurable arithmetic Kleinian groups Γ such that $H_1(\Gamma \backslash \mathbb{H}, \mathbb{Q}) \simeq H_1(\Gamma, \mathbb{Q}) = 0$. In [LMR], Long, Maclahlan and Reid asked the question whether there are infinitely many commensurability classes of arithmetic rational homology 3-spheres. In the same paper, they posed the following two con-

x	$R(x)/L_1(x)$	$R(x)/L_2(x)$	$R(x)/L_3(x)$	$R(x)/L_7(x)$	$R(x)/L_{11}(x)$
3000	1.793	3.654	1.827	2.294	2.099
6000	1.650	3.172	1.540	2.489	2.101
9000	1.720	3.334	1.435	2.462	2.281
12000	1.828	3.608	1.534	2.617	2.495
15000	1.927	3.610	1.524	2.662	2.731
18000	1.950	3.292	1.482	2.457	2.678
21000	1.912	3.114	1.575	2.464	2.642
24000	1.801	2.993	1.543	-	-
27000	1.782	3.000	1.594	-	-
30000	1.781	2.884	1.591	-	-
33000	1.830	-	1.632	-	-
36000	1.831	-	1.627	-	-
39000	1.825	-	1.612	-	-
42000	1.885	-	1.628	-	-
45000	1.887	-	1.607	-	-

Table 4: data related to the asymptotics of nonvanishing of cuspidal cohomology

jectures (they are slightly rephrased in an equivalent form that fits better with this paper).

Conjecture 5.1.

1. *There exist infinitely many pairs of prime ideals $\{\mathfrak{p}_1, \mathfrak{p}_2\} \subset \mathbb{Z}[i]$ such that*

$$H_{cusp}^2(\Gamma_0(\mathfrak{p}_1\mathfrak{p}_2), \mathbb{Q}) = 0.$$

2. *Let $\mathfrak{p} = (1 + i)$. There are infinitely many prime ideals $\mathfrak{q} \subset \mathbb{Z}[i]$ with $\mathbf{N}\mathfrak{q} = 1 \pmod{12}$ such that*

$$H_{cusp}^2(\Gamma_0(\mathfrak{p}\mathfrak{q}), \mathbb{Q}) = 0.$$

If the second conjecture holds, then using Jacquet-Langlands correspondence one gets (see [LMR] p.29) a positive answer to the question of Long-MacLachlan-Reid stated above.

I computed the ranks of $\Gamma_0(\mathfrak{p}\mathfrak{q})^{ab}$ where $\mathfrak{p} = (1 + i)$ and $\mathfrak{q} \subset \mathbb{Z}[i]$ prime with $\mathbf{N}\mathfrak{q} = 1 \pmod{12}$ of norm ≤ 14850 . There are 423 such prime ideals \mathfrak{q} , 245 of them satisfied the desired property that

$$H_{cusp}^2(\Gamma_0(\mathfrak{p}\mathfrak{q}), \mathbb{Q}) = \Gamma_0(\mathfrak{p}\mathfrak{q})^{ab} \otimes \mathbb{Q} = 0.$$

The uniform distribution of the primes with vanishing rank supports the second conjecture.

5.3. Asymptotics of torsion

Very recently Müller [Mü] and Bergeron-Venkatesh [BV] obtained significant results that relate the asymptotic behaviour of the size of the torsion in the homology of

certain cocompact lattices in $SL_2(\mathbb{C})$ to that of the volume of the associated 3-folds. The following is a special case of the main result of Bergeron and Venkatesh.

Theorem 5.2. *Let $\{\Gamma_n\}$ be a decreasing tower of cocompact arithmetic congruence subgroups of $SL_2(\mathbb{C})$ such that $\bigcap_n \Gamma_n = \{1\}$. Put $X = \Gamma_1 \backslash \mathbb{H}$ where \mathbb{H} is the hyperbolic 3-space. Then*

$$\lim_{n \rightarrow \infty} \frac{\log |H_1(\Gamma_n, E_{k,\ell})_{tor}|}{[\Gamma_1 : \Gamma_n]} = \frac{1}{6\pi} \cdot c_{k,\ell} \cdot \text{vol}(X), \quad k \neq \ell$$

where $E_{k,\ell}$ is standard module $Sym^k(\mathbb{Z}^2) \otimes \overline{Sym^\ell(\mathbb{Z}^2)}$. Here $c_{k,\ell}$ is a positive integer depending only on the non-equal parameters k, ℓ .

Bianchi groups and their congruence subgroups are outside of the scope of these results as they are not cocompact. It is very interesting to numerically investigate whether similar asymptotic relations hold for them as well.

I will compare the size of the torsion in the first homology and the volume of the associated 3-folds. As it is computationally costly to increase the weight, I will concentrate on increasing the level.

Let $\Gamma_0(\mathfrak{p})$ where \mathfrak{p} is a prime ideal of \mathcal{O} with residue degree one. Let $H_1(\Gamma_0(\mathfrak{p}), \mathbb{Z})_{tor}$ denote the torsion part of the first homology of $\Gamma_0(\mathfrak{p})$ with coefficients in \mathbb{Z} . Let $\text{vol}(\Gamma_0(\mathfrak{p}) \backslash \mathbb{H})$ denote the volume of the 3-fold $\Gamma_0(\mathfrak{p}) \backslash \mathbb{H}$ where \mathbb{H} is the hyperbolic 3-space. In light of the result of Bergeron-Venkatesh, the following is an interesting question.

Question 5.3. With the notation of the above paragraph, is there a constant C , independent of d , such that the asymptotic relation

$$\log |H_1(\Gamma_0(\mathfrak{p}), \mathbb{Z})_{tor}| \sim C \cdot \text{vol}(\Gamma_0(\mathfrak{p}) \backslash \mathbb{H})$$

holds as norm of the ideals $\mathfrak{p} \triangleleft \mathcal{O}_d$, which are prime with residue degree one, tends to infinity?

To computationally investigate the question, we need to approximate the volumes first. Using the well-known formula, see [GK],

$$\mathbb{V}_d := \text{vol}(\text{PSL}_2(\mathcal{O}_d) \backslash \mathbb{H}) = \frac{|\Delta_d|^{3/2}}{4\pi^2} \zeta_{K_d}(2),$$

where Δ_d is the discriminant of the field K_d and ζ_{K_d} is the Dedekind zeta function of K_d , we get

$$\begin{aligned} \mathbb{V}_1 &\simeq 0.305321864725739671684867838311 \\ \mathbb{V}_2 &\simeq 1.00384100334119813727236488577 \\ \mathbb{V}_3 &\simeq 0.169156934401608937503533759046 \\ \mathbb{V}_7 &\simeq 0.888914927816353263598904154202 \\ \mathbb{V}_{11} &\simeq 1.38260830790264587367165334450 \end{aligned}$$

Now for $\mathfrak{p} \triangleleft \mathcal{O}_d$ prime of residue degree one over the rational prime p , we have

$$\text{vol}(\Gamma_0(\mathfrak{p}) \backslash \mathbb{H}) = (p+1) \cdot \mathbb{V}_d.$$

I have collected data on the ratio of $\log |H_1(\Gamma_0(\mathfrak{p}), \mathbb{Z})_{tor}|$ to $\text{vol}(\Gamma_0(\mathfrak{p}) \backslash \mathbb{H})$ in the case of the five Euclidean \mathcal{O}_d . Ignoring the first 500 entries in each case, the average ratios

\mathcal{O}_1		\mathcal{O}_2		\mathcal{O}_3		\mathcal{O}_7		\mathcal{O}_{11}	
$N_{\mathfrak{p}}$	$T_{\mathfrak{p}}/V_{\mathfrak{p}}$	$N_{\mathfrak{p}}$	$T_{\mathfrak{p}}/V_{\mathfrak{p}}$	$N_{\mathfrak{p}}$	$T_{\mathfrak{p}}/V_{\mathfrak{p}}$	$N_{\mathfrak{p}}$	$T_{\mathfrak{p}}/V_{\mathfrak{p}}$	$N_{\mathfrak{p}}$	$T_{\mathfrak{p}}/V_{\mathfrak{p}}$
44533	0.05342	27817	0.05338	44533	0.05288	20549	0.05337	19583	0.05368
44537	0.05391	27827	0.05247	44563	0.05288	20563	0.05414	19603	0.05324
44549	0.05250	27851	0.05300	44587	0.05559	20693	0.05411	19661	0.05212
44617	0.05467	27883	0.05463	44617	0.05279	20707	0.05410	19699	0.05244
44621	0.05509	27947	0.05282	44623	0.05352	20717	0.05297	19717	0.05331
44633	0.05390	27953	0.05221	44641	0.05558	20731	0.05269	19727	0.05327
44641	0.05317	27961	0.05439	44647	0.05581	20743	0.05353	19739	0.05346
44657	0.05203	28001	0.05342	44683	0.05509	20749	0.05172	19759	0.05385
44701	0.05520	28019	0.05161	44701	0.05791	20759	0.05121	19793	0.05410
44729	0.05351	28027	0.05359	44773	0.05280	20771	0.05204	19801	0.05296
44741	0.05355	28051	0.05231	44797	0.05357	20773	0.05411	19853	0.05157
44753	0.05533	28057	0.05238	44809	0.05239	20857	0.05258	19867	0.05442
44773	0.05604	28081	0.05214	44839	0.05606	20897	0.05290	19889	0.05311
44777	0.05573	28097	0.05198	44851	0.05300	20899	0.05470	19891	0.05352
44789	0.05172	28099	0.05353	44887	0.05332	20903	0.05326	19913	0.05324
44797	0.05480	28123	0.05271	44893	0.05427	20939	0.05348	19919	0.05191
44809	0.05220	28163	0.05233	44917	0.05308	20959	0.05395	19937	0.05389
44893	0.05476	28201	0.05140	44953	0.05433	20981	0.05243	19963	0.05383
44909	0.05227	28211	0.05325	44959	0.05292	20983	0.05425	19979	0.05266
44917	0.05281	28219	0.05185	44971	0.05547	21001	0.04985	19991	0.05346
44953	0.05441	28283	0.05312	44983	0.05481	21011	0.05473	20021	0.05318

Table 5: the ratio the size of the torsion to the volume as level grows

read 0.054291, 0.053140, 0.055386, 0.053206, 0.053131 respectively for $-d = 1, 2, 3, 7, 11$. The range of my computations were up to norm 45000, 30000, 45000, 21000, 21000 respectively. It is very significant that the ratio is very close to

$$\frac{1}{6\pi} \simeq 0.0530516476972984452562945877908$$

which is the constant for the Lie group $SL_2(\mathbb{C})$ that appears in the above result of Bergeron-Venkatesh. A small sample is given in Table 5 where I use the convention $T_{\mathfrak{p}} := \log|H_1(\Gamma_0(\mathfrak{p}), \mathbb{Z})_{tor}|$ and $V_{\mathfrak{p}} := \text{vol}(\Gamma_0(\mathfrak{p}) \backslash \mathbb{H})$. The complete data is presented in my website.

References

- [AS] A.Ash, G.Stevens. Cohomology of arithmetic groups and congruences between systems of Hecke eigenvalues. *J. Reine Angew. Math.* **365** (1986), 192–220.
- [Be] T.Berger. Denominators of Eisenstein cohomology classes for GL_2 over imaginary quadratic fields. *Manuscripta Math.* **125.4** (2008), 427–470.
- [BV] N.Bergeron, A.Venkatesh. The asymptotic growth of torsion homology for arithmetic groups. preprint.

- [Bi] L.Bianchi. Sui gruppi di sostituzioni lineari con coefficienti appartenenti a corpi quadratici immaginari. *Math. Ann.* **40.3** (1892), 332–412.
- [BW] A.Borel, N.Wallach. Continuous Cohomology, Discrete Subgroups, and Representations of Reductive Groups. Vol. 94. Princeton, N.J.: Princeton University Press, 1980.
- [BCP] W.Bosma, J.Cannon, and C.Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, **24(3-4)** (1997), 235–265.
- [BN] R.Brauer, C.Nesbitt. On the modular characters of groups. *Ann. of Math. (2)* **42** (1941), 556–590.
- [Br] K.S.Brown. Cohomology of groups. New York, Springer-Verlag, 1994.
- [BM] A.Brumer, O.McGuinness. The behavior of the Mordell-Weil group of elliptic curves. *Bulletin of the A.M.S.* **23** (1990), 375–382
- [BS] A.Brumer, J. Silverman. The number of elliptic curves over \mathbb{Q} with conductor N , *Manuscripta Math.* **91** (1996), 95–102.
- [BLW] A.M.Brunner, Y.Lee, and N.J.Wielenberg. Polyhedral groups and graph amalgamation products. *Topology Appl.* **20.3** (1985), 289–304.
- [CD] F.Calegari, N.Dunfield. Automorphic forms and rational homology 3-spheres. *Geom. Topol.* **10** (2006), 295–329
- [Cr] J.E.Cremona. Hyperbolic tessellations, modular symbols, and elliptic curves over complex quadratic fields. *Compositio Math.* **51.3** (1984), 275–324.
- [EGM] J.Elstrodt, F.Grunewald, J.Mennicke. PSL(2) over imaginary quadratic integers. Arithmetic Conference (Metz, 1981), 43–60, Astérisque, 94, Soc. Math. France, Paris, 1982.
- [F] R.Fox. Free Differential Calculus. II. Derivation in the free group ring. *Ann. of Math. (2)* **57** (1953), 547–560.
- [FGT] T.Finis, F.Grunewald, and P.Tirao. Cohomology of lattices in $SL_2(\mathbb{C})$. *Exp. Math.* (2010) in press.
- [Fi] L.M.Figueiredo. Serre’s conjecture for imaginary quadratic fields. *Compositio Math.* **118.1** (1999), 103–122
- [Fl] D.Flöge. Zur Struktur der PSL_2 über einigen imaginär-quadratischen Zahlringen. *Math. Z.* **183.2** (1983), 255–279.
- [GK] F.Grunewald, S.Kühnlein. On the proof of Humbert’s volume formula. *Manuscripta Math.* **95.4** (1998), 431–436.
- [GS] F.Grunewald, J.Schwermer. Subgroups of Bianchi groups and arithmetic quotients of hyperbolic 3-Space. *Trans. Amer. Math. Soc.* **335.1** (1993), 47–78.
- [Ha] G.Harder. Eisenstein cohomology of arithmetic groups. The case GL_2 . *Invent. Math.* **89.1** (1987), 37–118.
- [Hi] H.Hida. Congruence of cusp forms and special values of their zeta functions. *Invent. Math.* **63.2** (1981), 225–261.
- [LMR] D.Long, C.Maclachlan, A.Reid. Arithmetic Fuchsian groups of genus zero. *Pure Appl. Math. Q.* **2.2**, part 2 (2006), 569–599.

- [M] S. Mac Lane. Homology. Berlin: Springer-Verlag, 1963.
- [Me] E.R.Mendoza. Cohomology of PGL_2 over imaginary quadratic integers. PhD Thesis, Universität Bonn Mathematisches Institut, 1979.
- [Mü] W.Müller. The asymptotics of the Ray-Singer analytic torsion of hyperbolic 3-manifolds. preprint.
- [P] C.Priplata. Cohomology and homology of PSL_2 over imaginary quadratic integers with general coefficients $M_{n,m}(\mathcal{O}_K)$ and Hecke eigenvalues. PhD. Thesis, Heinrich Heine Universität, Düsseldorf, 2000.
- [RF] A.Rahm, M.Fuchs. The integral homology of PSL_2 over imaginary quadratic integers of class number two. to appear in *J. of Pure and Applied Algebra*
- [S1] J-P.Serre. Le problème des groupes de congruence pour SL_2 . *Ann. of Math.* (2) **92** (1970), 489–527.
- [S2] J-P.Serre. Sur les représentations modulaires de degré 2 de $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. *Duke Math. J.*, **54**, (1987)
- [Se1] M.H.Şengün. Hyperelliptic curves of genus 2 and Bianchi modular forms. In preparation.
- [Se2] M.H.Şengün. A numerical study of mod p Bianchi modular forms and Galois representations. preprint.
- [ST] M.H.Şengün, S.Türkelli. Weight reduction for mod ℓ Bianchi modular forms. *J. Number Theory***129** (2009), 2010–2019.
- [SV] J.Schwermer, K.Vogtmann. The integral homology of SL_2 and PSL_2 of Euclidean imaginary quadratic integers. *Comment. Math. Helv.* **58.4** (1983), 573–598.
- [T] R.Taylor. On congruences between modular forms. PhD. Thesis, Princeton University 1988.
- [To] R.Torrey. On Serre’s conjecture over imaginary quadratic fields, preprint.
- [U] E.Urban. Formes automorphes cuspidales pour GL_2 sur un corps quadratique imaginaire. Valeurs spéciales de fonctions L et congruences. *Compositio Math.* **99.3** (1995), 283–324.
- [V] K.Vogtmann. Rational homology of Bianchi groups. *Math. Ann.* **272.3** (1985), 399–419.
- [Wa] X.D. Wang. The Hecke operators on the cusp forms of $\Gamma_0(N)$ with Nebentype. On Artin’s Conjecture for Odd 2-Dimensional Representations. LNM Vol. 1585. Berlin: Springer, (1994), 95–108.
- [Wi] G.Wiese. On the faithfulness of parabolic cohomology as a Hecke module over a finite field. *J. Reine Angew. Math.* **606** (2007), 79–103
- [Ya] D.Yasaki. An explicit spine for the Picard modular group over the Gaussian integers. *J. Number Theory* **128.1** (2008), 207–234.

DEPARTAMENT D’ALGEBRA I GEOMETRIA, FACULTAT DE MATEMÀTIQUES,
 UNIVERSITAT DE BARCELONA, GRAN VIA DE LES CORTS CATALANES, 585,
 BARCELONA, SPAIN
E-mail address: mehmet.sengun@uni-due.de