



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/97994/>

Other:

Sengun, M.H. and Tsaknias, P. (2013) Dimension Formulae for Spaces of Lifted Bianchi Modular Forms. UNSPECIFIED. (Unpublished)

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

DIMENSION FORMULAE FOR SPACES OF LIFTED BIANCHI MODULAR FORMS

MEHMET HALUK ŞENGÜN AND PANAGIOTIS TSAKNIAS

ABSTRACT. In this note, we work out the dimension of the subspace of base-change forms and their twists inside the space of Bianchi modular forms with fixed (Galois stable) level and weight. The formulae obtained here have been used for the numerical experiments of [1]. Extended versions of the formulae obtained in this note, treating CM-forms and spaces with non-trivial nebentypus, will be employed in the upcoming paper [3].

1. SETTING

Let $K = \mathbb{Q}(\sqrt{-D})$ be an imaginary quadratic field. Let $f \in S_k(\Gamma_0(N))$ be classical newform. To such an object one attaches an irreducible automorphic representation $\pi := \pi_f$ of $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$. One then has the so-called *Base Change* operator:

$$\mathrm{BC}_{\mathbb{Q}}^K : \mathcal{A}_n(\mathbb{Q}) \rightarrow \mathcal{A}_n(K)$$

We would like to understand how the base change operator affects the level of an automorphic representation. We will in fact restrict our attention to the 2-dimensional case (which will in turn force us to consider the 1-dimensional case as well).

Let $\pi \in \mathcal{A}_n(\mathbb{Q})$. One can write it as a restricted tensor product of local ones $\pi_{\mathfrak{p}} \in \mathcal{A}_n(\mathbb{Q}_{\mathfrak{p}})$, where p runs through all finite primes and infinity. It is well-known that Base-Change is compatible with taking local components, i.e for every rational prime p and every prime \mathfrak{p} of K above it we have:

$$\mathrm{BC}_{\mathbb{Q}}^K(\pi)_{\mathfrak{p}} = \mathrm{BC}_{\mathbb{Q}_{\mathfrak{p}}}^{K_{\mathfrak{p}}}(\pi_{\mathfrak{p}})$$

In light of this it is enough to examine what happens at each prime individually. This allows us to freely exploit the local Langlands correspondence throughout this note. In fact the base change operator is the one that corresponds on the restriction operation (locally and globally)

$$\mathrm{Res}_{W_K}^{W_{\mathbb{Q}}} : \mathcal{G}_n(\mathbb{Q}) \rightarrow \mathcal{G}_n(K),$$

$$\mathrm{Res}_{W_{K_{\mathfrak{p}}}}^{W_{\mathbb{Q}_p}} : \mathcal{G}_n(\mathbb{Q}_p) \rightarrow \mathcal{G}_n(K_{\mathfrak{p}}),$$

on n -dimensional Weil-Deligne representations. We will denote by $\mathrm{Art}_F : W_F \rightarrow F^{\times}$ the local Artin map of local class field theory. Then the local Langlands correspondence in the 1-dimensional case is composition with Art_F^{-1} .

Going back to the automorphic side, if p is finite then one has the following classification of the types of the $\pi_{\mathfrak{p}}$ that can occur:

- Principal series $\mathrm{PS}(\chi_1, \chi_2)$, where $\chi_i \in \mathcal{A}_1(\mathbb{Q}_p)$ and $\chi_1/\chi_2 \neq \|\cdot\|^{\pm 1}$. The corresponding Weil-Deligne representation $WD(\pi_p) = (\rho_p, N_p)$ is in this case split (decomposable), i.e. $\rho_p \sim \tilde{\chi}_1 \oplus \tilde{\chi}_2$ and $N_p = 0$, where $\tilde{\chi}_i$ is the corresponding character of the Weil Group, i.e. $\tilde{\chi} = \chi \circ \mathrm{Art}_p$. We will study this type in Section 3.1.

- Special $S(\chi||\cdot||, \chi)$, where $\chi \in \mathcal{A}_1(\mathbb{Q}_p)$. The corresponding local Galois representation is in this case reducible but not decomposable: $\rho_{\pi_p} \sim \tilde{\chi}||\cdot|| \oplus \tilde{\chi}$ and $N = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. This will be the subject of Section 3.2.
- Supercuspidal, which is all the rest. We will describe this type in more detail in section Section 3.3.

Let now p be a rational prime and \mathfrak{p} a prime in K above it. We have the following:

- $D_p = D_{\mathfrak{p}}$ iff p splits, i.e. $p = \mathfrak{p}\mathfrak{p}'$.
- $I_p = I_{\mathfrak{p}}$ iff p is unramified, i.e. $p \neq \mathfrak{p}^2$.

This trivial observation tells us that $\alpha(\rho|_{I_p}) = \alpha(\rho|_{I_{\mathfrak{p}}})$ for all the unramified primes p in K and therefore the conductor stays the same at these primes. Lets examine what happens for the ramified primes in K whose (finite) set we will denote by R_K .

Remark 1. Since we have established the finiteness of the set of primes where any change in the conductor can possibly occur one can wlog consider a suitable ℓ -adic representation of $G_{\mathbb{Q}}$ instead of the Weil-Deligne representation of $W_{\mathbb{Q}}$: This follows from the Deligne-Grothendieck classification of ℓ -adic representations if we pick ℓ big enough so that ℓ not in R_K and π is unramified at it. Same thing goes locally.

In what follows, we will assume for simplicity that 2 is unramified in K . We treat first the case $n = 1$, and then in the following three sections we treat the case $n = 2$, one type at a time. We then finish with some dimension formulae for spaces of modular forms of a given type. In the future, we will also consider the Automorphic Induction operator and spaces with non-trivial Nebentypus.

2. THE CASE $n = 1$

We will work out this case not just for quadratic extensions of \mathbb{Q}_p but for any quadratic extension E/F of local fields over \mathbb{Q}_p . Let $\chi \in \mathcal{A}_1(F)$. Let \mathfrak{p}_F and \mathfrak{p}_E be the primes in \mathcal{O}_F and \mathcal{O}_E respectively such that $\mathfrak{p}_E|\mathfrak{p}_F|p$. We then have an explicit description of the Base Change operator thanks to local class field theory:

$$\mathrm{BC}_F^E(\chi) = \chi \circ \mathbf{N}_{E/F}.$$

Recall that the conductor of a character η of F^* is the smallest integer such that $\eta|_{U_F^i}$ is trivial, with U_F^i defined as:

$$U_F^i = \begin{cases} \mathcal{O}_F^\times & i = 0 \\ 1 + \mathfrak{p}_F^i & i > 0 \end{cases}$$

Clearly if χ is unramified, then $\mathrm{BC}_F^E(\chi)$ is unramified too. Assume that χ is tamely ramified (and therefore $p > 2$). It then factors through the quotient $\mathcal{O}_F^*/U_F^1 \cong k_F^*$. By [2, Lemma 18.1] we have that $\mathrm{BC}_{\mathbb{Q}_p}^{K_p}(\chi_p)$ can be at most tame. It is unramified if and only if $\chi_p|_{\mathbb{Z}_p^*}$ has order 2.

The following Proposition explains what happens when the conductor of χ_p is at least 2 in the case where K_p/\mathbb{Q}_p is tamely ramified (which is the only possible case since $p > 2$):

Proposition 2. [2, Proposition 18.1] *Assume that K_p/\mathbb{Q}_p is tamely ramified. Then:*

$$\alpha(\mathrm{BC}_{\mathbb{Q}_p}^{K_p}(\chi_p)) = e_{K_p/\mathbb{Q}_p}(\alpha(\chi_p) - 1) + 1 \quad \forall m \geq 2$$

Remark 3. Bushnell and Henniart use the term *level*, which is equal to the conductor minus 1.

Putting everything together:

Proposition 4.

$$\alpha(BC_{\mathbb{Q}_p}^{K_p}(\chi_p)) = \begin{cases} 0 & \alpha(\chi_p) = 0 \\ 0 & \alpha(\chi_p) = 1 \text{ and } \chi_p^2|_{Z_p^*} = 1 \\ 1 & \alpha(\chi_p) = 1 \text{ and } \chi_p^2|_{Z_p^*} \neq 1 \\ e(K_p/\mathbb{Q}_p)(m-1) + 1 & \alpha(\chi_p) = m \geq 2 \end{cases}$$

Remark 5. Notice that this proposition covers the conductor at all primes of K , not just the ramified ones. As we expected, the conductor does not change at the unramified ones though.

3. THE CASE $n = 2$

3.1. The split case. In this case π_p is a principal series $I(\chi_1, \chi_2)$ parametrized by two Hecke characters $\chi_1, \chi_2 \in \mathcal{A}_1(\mathbb{Q}_p)$. Obviously one has (by looking at the Galois side for example) that

$$BC_{\mathbb{Q}_p}^{K_p}(I(\chi_1, \chi_2)) = I(BC_{\mathbb{Q}_p}^{K_p}(\chi_1), BC_{\mathbb{Q}_p}^{K_p}(\chi_2)).$$

Combining this with the conductor formula

$$\alpha(I(\chi_1, \chi_2)) = \alpha(\chi_1) + \alpha(\chi_2)$$

for principal series one then reduces the problem to the 1-dimensional case that we just treated in the previous Section.

3.2. The Special case. In this case π_p is a (twisted) Steinberg representation $S(\chi||\cdot||, \chi) = \chi \otimes \text{St}$ parametrized by one Hecke character $\chi \in \mathcal{A}_1(\mathbb{Q}_p)$. We have the following conductor formula:

$$\alpha(\chi||\cdot||, \chi) = \begin{cases} 1 & \alpha(\chi) = 0 \\ 2\alpha(\chi) & \alpha(\chi) > 0. \end{cases}$$

In order to understand base change in this case we will exploit the Galois side. The base change corresponds to restriction there and therefore it is easy to see that:

$$BC_{\mathbb{Q}_p}^{K_p}(S(\chi||\cdot||, \chi)) = S(BC_{\mathbb{Q}_p}^{K_p}(\chi)||\cdot||, BC_{\mathbb{Q}_p}^{K_p}(\chi))$$

Since the same dimension formula applies for the RHS, we have again reduced the problem to one in the 1-dimensional case.

3.3. The Supercuspidal case. In this case π_p is none of the previous types. Since we have the ongoing assumption that $p > 2$, all these local automorphic representations are parametrized by pairs (E, θ) where E/\mathbb{Q}_p is quadratic and $\theta \in \mathcal{A}_1(E) \setminus BC_{\mathbb{Q}_p}^E(\mathcal{A}_1(\mathbb{Q}))$. We also get that the corresponding Galois representation looks like

$$\rho_{\pi_p} \sim \text{Ind}_{G_E}^{G_p}(\theta \circ \text{Art}_p).$$

We will write $\tilde{\theta}$ for $\theta \circ \text{Art}_p$. Base change corresponds to restriction on the Galois side so we want to examine the conductor of $\text{Res}_{G_p}^{G_p} \text{Ind}_{G_E}^{G_p} \tilde{\theta}$. Using [6, Proposition 22] we get:

$$\text{Res}_{G_p}^{G_p} \text{Ind}_{G_E}^{G_p} \tilde{\theta} = \begin{cases} \tilde{\theta} \oplus \tilde{\theta}^\sigma & G_p = G_E, \\ \text{Ind}_{G_E \cap G_p}^{G_p} \tilde{\theta}|_{G_E \cap G_p} & G_p \neq G_E \end{cases}.$$

Here σ is a generator of $\text{Gal}(K_p/\mathbb{Q}_p)$ and $\tilde{\theta}^\sigma(x) = \tilde{\theta}(g^{-1}xg)$, for all $x \in G_p$ and $g \in G_p$ a lift of σ . Notice that since K_p/\mathbb{Q}_p is ramified we have that $I_p \not\subseteq G_p$ so we can pick g to be in I_p .

Let's treat the first case. The conductor of the representation in the Galois side is equal to $\alpha(\tilde{\theta}) + \alpha(\tilde{\theta}^\sigma)$. We also have that $\alpha(\tilde{\theta}) = \alpha(\tilde{\theta}^\sigma)$ so we get that:

$$\alpha(\mathrm{BC}_{\mathbb{Q}_p}^{K_p}(\pi_p)) = 2\alpha(\theta).$$

For the second case we employ [5, Theorem 8.2] which gives the following formula:

$$\alpha(\mathrm{Ind}_{G_{EK_p}}^{G_p} \tilde{\theta}|_{G_{EK_p}}) = f_{EK_p/K_p} \left(d_{EK_p/K_p} + \alpha(\tilde{\theta}|_{G_{EK_p}}) \right)$$

Translating it into a statement on the automorphic side gives:

$$\alpha(\mathrm{BC}_{\mathbb{Q}_p}^{K_p}(\pi_p)) = f_{EK_p/K_p} \left(d_{EK_p/K_p} + \alpha(\mathrm{BC}_{K_p}^{EK_p}(\theta)) \right)$$

The conductor on the RHS can then be computed using the formula obtained for the 1-dimensional case.

4. DIMENSION FORMULAE FOR SPACES OF FIXED TYPE

Let N be a positive integer. We will denote the automorphic forms of level N by $\mathcal{A}_2^N(\mathbb{Q})$, i.e. $\mathcal{A}_2^N(\mathbb{Q}) := \mathcal{A}_2(\mathbb{Q})^{U_1(N)}$. In the same fashion, $\mathcal{A}_2^{\mathfrak{n}}(K) := \mathcal{A}_2(K)^{U_1(\mathfrak{n})}$ will denote the automorphic forms over K of level \mathfrak{n} , where \mathfrak{n} is an ideal of \mathcal{O}_K .

Assume for simplicity that K be an imaginary quadratic field with **odd** discriminant $d = \mathrm{disc}(K/\mathbb{Q})$ and that N is a **square-free** positive integer that is coprime to d . Let $\Pi \in \mathcal{A}_{2,bc}^{(N)}(K)$ and $\pi \in \mathcal{A}_2(\mathbb{Q})$ such that Π is a twist of $BC_{\mathbb{Q}}^K(\pi)$.

Definition 6. For $k \geq 0$, let $\mathcal{B}_2^N(\mathbb{Q}, k)$ denote the set of all cuspidal automorphic representations π of $\mathrm{GL}_2(\mathbb{A}_{\mathbb{Q}})$ such that

- π_∞ is the holomorphic discrete series representation of weight $k+2$,
- for every prime $p|N$, π_p is of Steinberg type,
- for every prime $\ell|d$, there are three possibilities for π_ℓ :
 - (1) unramified principal series,
 - (2) $PS(\alpha, \omega_{K,\ell}\beta)$ with α, β unramified characters of \mathbb{Q}_ℓ^* and ω_K is the quadratic character associated to K ,
 - (3) $AI_{K_\ell/\mathbb{Q}_\ell}(\theta_\ell) \otimes \gamma$ with an unramified character γ of \mathbb{Q}_ℓ^* .

Assume that Π is of weight k , that is, Π_∞ is the principal representation $\rho_\infty^k = PS(\chi_\infty^{k+1}, \chi_\infty^{-k-1})$ of $\mathrm{GL}_2(\mathbb{C})$ with $\chi_\infty(z) = z/|z|$. Then one can find $\pi \in \mathcal{B}_2^N(\mathbb{Q}, k)$ such that $\Pi \simeq BC(\pi) \otimes \chi$ for some finite order idele class character χ of K . Conversely, for any non-CM π in $\mathcal{B}_2^N(\mathbb{Q}, k)$, there exists a finite order idele class character χ of K such that $\Pi \simeq BC(\pi) \otimes \chi$. Moreover, it is easy to see that the base change map $\mathcal{B}_2^N(\mathbb{Q}, k) \rightarrow \mathcal{A}_{2,bc}^{(N)}(K, k)$ is at most two-to-one.

Remark 7. Assume that $\ell = \mathrm{disc}(K/\mathbb{Q})$ is prime. In this special case, the elliptic modular form associated to $\pi \in \mathcal{B}_2^N(\mathbb{Q}, k)$ lives in $S_k(\Gamma_0(N))^{new}$, $S_k(\Gamma_0(N\ell), \omega_K)^{new}$ and $S_K(\Gamma_0(N\ell^2), \epsilon)^{N-new}$ depending on whether π_ℓ has type 1, 2 or 3 as above. The base change map is one-to-one on the subset of π 's with local type 1 at ℓ , and is two-to-one on the subset of π 's with local type 2 or 3 at ℓ .

To compute the dimensions of spaces of elliptic modular forms corresponding to π with the local conditions as above, we use a formula (see [4, Proposition 4.18.]) that computes the multiplicity of any given representation of $\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ in the space $S_k(\Gamma(N))$ of elliptic modular forms of weight k for the *principal* congruence subgroup of level N .

Proposition 8. *Let $N \geq 1$ and $k \geq 2$ be integers and σ a representation of $G_N = \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ such that $\sigma(-I_2)$ is $(-1)^k$. Let $U_N \leq G_N$ be the subgroup of all upper triangular unipotent elements and S_3 and S_4 be the images of elements of $\mathrm{SL}_2(\mathbb{Z})$ of order 3 and 4, respectively. Then*

$$\dim \mathrm{Hom}_{G_N}(\sigma, S_k(\Gamma(N))) = \frac{k-1}{12} \dim \sigma - \frac{1}{2} \dim \sigma^{U_N} + \varepsilon_k \mathrm{tr} \sigma(S_4) + \rho_k \mathrm{tr} \sigma(S_3) + \delta_{k,2} \dim \sigma^{G_N}$$

where

$$\varepsilon_k = \begin{cases} \frac{(-1)^{k/2}}{4}, & \text{if } k \equiv 0 \pmod{2}, \\ 0, & \text{otherwise.} \end{cases}$$

and

$$\rho_k = \begin{cases} 0, & \text{if } k \equiv 1 \pmod{3}, \\ -1/3, & \text{if } k \equiv 2 \pmod{3}, \\ 1/3, & \text{if } k \equiv 0 \pmod{3}. \end{cases}$$

and $\delta_{k,2}$ is the Kronecker delta.

Remark 9. Let us assume the situation of the previous remark. Then to compute the dimension of $S_k(\Gamma_0(N))^{new}$, we need to consider (the restriction to $\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ of) the representation σ of $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})$ which is the tensor product of the representations σ_p of $\mathrm{GL}_2(\mathbb{F}_p)$ where σ_p is the (virtual) representation $[St] - [1]$ where St is the Steinberg representation. Using the character table of $\mathrm{GL}_2(\mathbb{F}_p)$, we conclude that for **even** $k \geq 2$,

$$\dim S_k(\Gamma_0(N))^{new} = \frac{k-1}{12} \prod_{p|N} (p-1) + \varepsilon_k \prod_{p|N} \left(\left(\frac{-4}{p} \right) - 1 \right) + \rho_k \prod_{p|N} \left(\left(\frac{-3}{p} \right) - 1 \right) + \delta_{k,2} (-1)^{div(N)}$$

where $div(N)$ is the number of prime divisors of N .

Using Finis-Grunewald-Tirao [4] and Bushnell-Henniart [2], we also compute that for **odd** $k \geq 3$,

$$\begin{aligned} \dim S_k(\Gamma_0(N\ell), \omega_K)^{new} &= \frac{k-1}{12} (\ell+1) \prod_{p|N} (p-1) \\ &+ \varepsilon_k \left(\left(\frac{-4}{\ell} \right) + 1 \right) \cdot \prod_{p|N} \left(\left(\frac{-4}{p} \right) - 1 \right) \\ &+ \rho_k \left(\left(\frac{-3}{\ell} \right) + 1 \right) \cdot \prod_{p|N} \left(\left(\frac{-3}{p} \right) - 1 \right) \end{aligned}$$

Finally, to compute the subspace of $S_K(\Gamma_0(N\ell^2), \epsilon)^{corr}$ of $S_K(\Gamma_0(N\ell^2), \epsilon)^{N-new}$ that have the local type 3 at ℓ , we proceed as above and find that for **even** $k \geq 2$,

$$\begin{aligned} \dim S_k(\Gamma_0(N\ell^2), \epsilon)^{corr} &= \frac{k-1}{12} (\ell-1) \prod_{p|N} (p-1) \\ &+ \varepsilon_k \mathrm{tr} \sigma_\ell(S_4) \cdot \prod_{p|N} \left(\left(\frac{-4}{p} \right) - 1 \right) \\ &+ \rho_k \left(\left(\frac{-3}{\ell} \right) - 1 \right) \cdot \prod_{p|N} \left(\left(\frac{-3}{p} \right) - 1 \right) \end{aligned}$$

where

$$\mathrm{tr} \sigma_\ell(S_4) = \begin{cases} 2 \cdot (-1)^{(\ell-3)/4}, & \text{if } \ell \equiv 3 \pmod{4}, \\ 0, & \text{otherwise.} \end{cases}$$

Let $BS_k^{bc}(\Gamma_0(N))^{new}$ denote the base-change subspace of space of *new* cuspidal *Bianchi modular forms* over K of weight k for $\Gamma_0(N)$, that is, the subspace of $BS_k(\Gamma_0(N))^{new}$ that is accounted for by Π 's in $\mathcal{A}_{bc}^N(K, k)$. Then for **even** $k \geq 2$, we have

$$\dim BS_k^{bc}(\Gamma_0(N))^{new} = \dim S_k(\Gamma_0(N))^{new} + \frac{1}{2} \dim S_k(\Gamma_0(N\ell^2), \epsilon)^{corr}$$

and for **odd** $k \geq 3$, we have

$$\dim BS_k^{bc}(\Gamma_0(N))^{new} = \frac{1}{2} \dim S_k(\Gamma_0(N\ell), \omega_K)^{new}.$$

REFERENCES

- [1] N. Bergeron, M. H. Şengün, and A. Venkatesh, *Torsion homology growth and cycle complexity of arithmetic manifolds*, in progress.
- [2] C. J. Bushnell and G. Henniart, *The local Langlands conjecture for $GL(2)$* , Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 335, Springer-Verlag, Berlin, 2006. MR 2234120 (2007m:22013)
- [3] M. H. Şengün and P. Tsaknias, *Even Galois representations and the torsion homology of Bianchi groups*, in progress.
- [4] T. Finis, F. Grunewald, and P. Tirao, *The cohomology of lattices in $SL(2, \mathbb{C})$* , Experiment. Math. **19** (2010), no. 1, 29–63.
- [5] G. Henniart, *Représentations du groupe de Weil d'un corps local*, Publications Mathématiques d'Orsay 79 [Mathematical Publications of Orsay 79], vol. 2, Université de Paris-Sud Département de Mathématique, Orsay, 1979, With an English summary. MR 551495 (80m:12016)
- [6] J.-P. Serre, *Linear representations of finite groups*, Springer-Verlag, New York, 1977, Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42. MR 0450380 (56 #8675)

MATHEMATICS INSTITUTE, ZEEMAN BUILDING, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL, UK
E-mail address: M.H.Sengun@warwick.ac.uk

FACULT DES SCIENCES, DE LA TECHNOLOGIE ET DE LA COMMUNICATION, UNIVERSITÉ DU LUXEMBOURG
E-mail address: panagiotis.tsaknias@uni.lu
E-mail address: p.tsaknias@gmail.com