



This is a repository copy of *Speaker sex effects on temporal and spectro-temporal measures of speech*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/98009/>

Version: Accepted Version

---

**Article:**

Herrmann, F. [orcid.org/0000-0002-6672-9369](http://orcid.org/0000-0002-6672-9369), Cunningham, S.P. and Whiteside, S.P. (2014) Speaker sex effects on temporal and spectro-temporal measures of speech. *Journal of the International Phonetic Association* , 44 (01). pp. 59-74. ISSN 0025-1003

<https://doi.org/10.1017/S0025100313000315>

---

This article has been published in a revised form in *Journal of the International Phonetic Association* [<https://doi.org/10.1017/S0025100313000315>]. This version is free to view and download for private research and study only. Not for re-distribution, re-sale or use in derivative works. © International Phonetic Association 2014.

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

**Speaker sex effects on temporal and spectro-temporal measures of speech****Frank Herrmann**

Department of English

University of Chester

Parkgate Road

Chester, CH1 4BJ

Phone: +44(0)1244 511392

*f.herrmann@chester.ac.uk***Stuart P. Cunningham****Sandra P. Whiteside**

Department of Human Communication Sciences

University of Sheffield

*s.cunningham@sheffield.ac.uk**s.whiteside@sheffield.ac.uk*

Published in: Journal of the International Phonetic Association / Volume 44 / Issue 01 / April 2014, pp 59-74. DOI: 10.1017/S0025100313000315, Published online: 21 March 2014.

Link to this article: [http://journals.cambridge.org/abstract\\_S0025100313000315](http://journals.cambridge.org/abstract_S0025100313000315)

## **Abstract**

This study investigated speaker sex differences in the temporal and spectro-temporal parameters of English monosyllabic words spoken by thirteen women and eleven men. Vowel and utterance duration were investigated. A number of formant frequency parameters were also analysed to assess the spectro-temporal dynamic structures of the monosyllabic words as a function of speaker sex. Absolute frequency changes were measured for the first (F1), second (F2) and third (F3) formant frequencies ( $\Delta F1$ ,  $\Delta F2$ , and  $\Delta F3$ , respectively). Rates of these absolute formant frequency changes were also measured and calculated to yield measurements for  $rF1$ ,  $rF2$  and  $rF3$ . Normalised frequency changes ( $\text{norm}\Delta F1$ ,  $\text{norm}\Delta F2$ , and  $\text{norm}\Delta F3$ ), and normalised rates of change ( $\text{norm}rF1$ ,  $\text{norm}rF2$ , and  $\text{norm}rF3$ ) were also calculated. F2 locus equations were then derived from the F2 measurements taken at the onset and temporal mid points of the vowels. Results indicated that there were significant sex differences in the spectro-temporal parameters associated with F2:  $\Delta F2$ ,  $\text{norm}\Delta F2$ ,  $rF2$ , and F2 locus equation slopes; women displayed significantly higher values for  $\Delta F2$ ,  $\text{norm}\Delta F2$  and  $rF2$ , and significantly shallower F2 locus equation slopes. Collectively, these results suggested lower levels of coarticulation in the speech samples of the women speakers, and corroborates evidence reported in earlier studies.

## **1 Introduction**

### **1.1 Sex differences in speech production**

Speaker sex differences in adults have been reported in the temporal domain of speech. For example, faster speaking rates (Jacewicz, Fox & Wei 2010) and shorter acoustic vowel durations (e.g. Simpson 1998, Whiteside 1996) have previously been attributed to male speakers. However, there is also evidence to suggest that sex differences in vowel durations are complex, and these differences are context dependent (Simpson & Ericsson 2003). In addition, there are reports of speaker sex differences in the spectral parameters of speech. For example, a more distinct and larger vowel space has been found for women's formant data (e.g. Lee, Potamianos, & Narayanan 1999, Simpson 2009, Traunmüller 1988, Whiteside 2001). These speaker sex differences in vowel spaces could be attributed to the sexual dimorphism in supralaryngeal dimensions (i.e. the oral cavity to pharynx ratio). However, evidence suggests that some sex differences in speech production might also be a function of sociocultural factors (e.g. Byrd 1992, 1994; Henton & Bladon 1985; Whiteside 2001). The sex differences in speaking style in more formal settings observed by Byrd (1992, 1994), the fuller phonetic forms displayed by women (Whiteside 1996), as well as the greater phonetic distinctiveness of vowels (e.g. a larger vowel space for women) found in Lee et al.'s 1999 database (Whiteside 2001), suggest that lower degrees of coarticulation might also be a feature of women's speech. This suggestion is supported by McLeod, Baillargeon, Metz, Schiavetti & Whitehead (2001), who found significantly steeper second formant (F2) locus equation slopes for male than for female speakers, thereby suggesting higher levels of coarticulation for male speakers. Considering these previously reported speaker sex differences, the following hypotheses were investigated in this study. Firstly, female speakers were expected to exhibit longer utterance and vowel durations

than male speakers. Secondly, it was predicted that male speakers would show greater degrees of coarticulation than female speakers. The faster speaking rates attributed to male speakers may suggest faster mean rates of formant change. However, differences in vocal tract dimensions (Fitch & Giedd 1999) and the more peripheral and larger vowel space of female speakers (Whiteside 2001) may contribute to larger changes in formant frequencies over time. On this basis, an additional hypothesis was that male speakers would exhibit slower mean rates of formant change in the acoustic signal, despite articulatory movements which cover a greater distance.

## 1.2 Temporal and spectro-temporal measures of speech

The focus of this study was on spectro-temporal measures which may hypothetically vary as a function of speaker sex. In addition to measuring utterance and vowel durations, this study quantified the degree of coarticulation by calculating the absolute formant frequency changes in the first three formants between vowel onset and temporal midpoint for each participant (see Equation 1).

$$\text{(Eq. 1) } \Delta F_n = ||F_{n_{\text{midpoint}}} - F_{n_{\text{onset}}}|$$

Smaller absolute formant changes (e.g. smaller  $\Delta F_2$  values) have been associated with greater degrees of coarticulation, because smaller differences between vowel onset and midpoint formant values may indicate an approximation of both vowel and consonant targets (e.g. Brancazio & Fowler 1998, Fowler 1994). In order to control for sex differences in vocal tract dimensions, the absolute formant frequency changes were also normalised for each participant (see Equation 2).

$$\text{(Eq. 2) } \text{norm}\Delta F_n = [2\Delta F_n / (F_{n_{\text{onset}}} + F_{n_{\text{midpoint}}})]$$

As a measure of dynamic changes in the formant frequencies, mean rates of formant frequency change were calculated based on the absolute formant changes and vowel durations for each participant (see Equation 3) where  $t_{\text{vowel}}$  is the duration of the vowel.

$$\text{(Eq. 3) } rFn = \Delta Fn / (0.5 \times t_{\text{vowel}})$$

During the production of CV sequences, these have been shown to indirectly correlate with the mean velocity of the active articulators while moving from the release of the preceding consonant to the vowel target (e.g. Chang, Ohde & Conture 2002, Simpson 2001). Greater mean rates of formant change may be interpreted as being indicative of faster movements of the active articulators. To allow for direct comparisons between the male and female data, mean rates of formant frequency change were also normalised for each participant (see Equation 4).

$$\text{(Eq. 4) } \text{normrFn} = [2 rFn / (Fn_{\text{onset}} + Fn_{\text{midpoint}})]$$

In addition to absolute formant changes, F2 locus equations were derived to gauge the degree of coarticulation (see Equation 5). These are linear regression functions based on the F2 values at vowel midpoint (independent variable), and F2 values at vowel onset (dependent variable).

$$\text{(Eq. 5) } F2_{\text{onset}} = \text{slope} \times F2_{\text{midpoint}} + \text{y-intercept}$$

The linear relationship of the second formant in vowel onset and target position was first observed by Lindblom (1963). Later studies, such as Krull (1987), applied F2 locus equations to quantify coarticulation by analysing the slope of the linear regression function; steeper slopes (and corresponding lower y-intercept values) were associated with greater degrees of coarticulation and shallower slopes (and corresponding higher y-intercept values), with the converse. Although the relationship between articulatory data and F2 locus equations is not linear, it has been shown to be lawful (Löfqvist 1999; Tabain 2000, 2002). In addition, more recently, Iskarous, Fowler & Whalen (2010) and Lindblom & Sussman (2012) have demonstrated the link between articulatory data and F2 locus equations.

Studies on F2 locus equations for CV sequences have consistently reported place of articulation effects. For example, slopes for the bilabial plosives such as /b/ are consistently steeper and corresponding y-intercepts, lower, compared to those observed for the alveolar cognate /d/ (e.g. Fowler 1994, McLeod et al. 2001, Sussman, Bessell, Dalston & Majors 1997, Sussman, Dalston & Gumbert 1998). These observations suggest that /b/ in CV sequences has the highest levels of coarticulation with the ensuing vowel, and therefore the lowest levels of articulatory resistance due to the independence between the main articulators involved in the production of /bV/; the lips and the lingual system. However, while bilabials display the steepest slopes across different places of articulation, the slope values for /bilabial plosive V/ sequences vary across a range of studies (see Table 1 for examples).

In addition to place of articulation effects on F2 locus equations (e.g. Sussman et al. 1997, 1998), studies have also found evidence for allophonic variation and vowel context effects (Sussman et al. 1991, 1997, 1998; Fowler 1994). While numerous studies report vowel context effects in locus equations between front (shallower slopes and higher y-intercept values) and back vowels (steeper slopes and lower y-intercept values) for velar plosives (Sussman et al. 1991, 1997, 1998), others report on similar, if less marked, effects for both bilabial and alveolar plosives (Fowler 1994). Taken together, the place of articulation and vowel context effects on F2 locus equations provide evidence for phonetic context effects on coarticulation effects indexed by F2 locus equations.

This additional measure of coarticulation was used in the current study because F2 locus equations preserve the phonetic context (F2 values at onset and midpoint), whereas the absolute formant changes merely encode the differences. F2 locus equation slopes can be compared using a large sample Z-test for parallelism (Kleinbaum & Kupper 1978) based on all individual data points (multiple repetitions/contexts), while comparisons of absolute formant changes are based on mean values per speaker group and subtle effects may not be detected. Lastly, as the phonetic context is largely preserved in F2 locus equations, and onset values are plotted as a function of vowel midpoint/target values, they have the capacity to encode the degree of anticipatory (right-to-left) coarticulation, which has been attributed to speech planning processes (e.g. Fowler & Saltzman 1993), while absolute formant changes encode both anticipatory and perseverative (left-to-right) coarticulation simultaneously.

## **2 Methods**

## **2.1 Stimuli**

The stimuli used in the current investigation formed part of a larger study on syllable frequency and speaker sex effects on speech production (Herrmann 2011). The aim of the current study was to focus specifically on the effects of speaker sex on the temporal and spectro-temporal parameters of speech in the production of the following monosyllabic words by twenty-four native speakers of British English: /bɪz, bɪs, bɪt, bɪb, bɛd, bɛk, bɛst, bu:st/. The original primary selection criteria for these stimuli were their frequency of occurrence (per million words, pmw) in spoken language (Baayen, Piepenbrock, & van Rijn 1993) and a matched phonetic makeup across two narrow frequency bands (325–56 pmw and 1–7 pmw). Other places of articulation for the word-initial plosive were not considered, as no further stimuli could be phonetically matched across frequency categories (Herrmann 2011). However, frequency effects are not the focus of this analysis.

## **2.2 Speakers**

Eleven male and thirteen female native speakers of English (mean age: 20;1 years; SD: 1;8; range: 18 – 24) participated in the study. None of the participants had a known speech or language impairment and all passed a hearing test prior to the experiment using a Kamplex Screening Audiometer (AS7) with an upper hearing level of 20 dB.

## **2.3 Data Collection**

Speech accommodation in interactional settings is a well-established phenomenon; speakers have been found to adjust their speech and vocal patterns to accommodate to those of their interlocutors (Giles, Coupland & Coupland 1991). Furthermore, behavioural studies have found

strong links between perception and behaviour in face-to-face social interactions with evidence for ‘chameleon effects’ (the unintentional, non-conscious mimicry of others) between individuals (Chartrand & Bargh 1999). Moreover, there is also evidence for individual differences in these effects where those who are profiled as displaying higher levels of empathy also appear to exhibit higher levels of mimicry (Chartrand & Bargh 1999). Therefore, in order to limit the effects of speech accommodation and mimicry which might occur in a social interactional face-to-face setting, the list of speech stimuli were presented to individual participants auditorily as a series of recorded prompts. A single-walled sound attenuating booth was used to secure high quality recordings. Furthermore, no experimenter was present within the sound booth during the recording sessions. While it is acknowledged that some speech accommodation and mimicry might occur when only auditory information is presented, the same fixed protocol for speech data collection was adopted for each of the 24 participants to control for these effects across speakers. The speech stimuli were presented to each participant via loudspeakers to avoid the Lombard effect (Lombard 1911, Junqua 1993). All the prompt speech stimuli were produced by a single adult male speaker and were presented in a randomised order. Participants were required to listen to a stimulus, repeat it during a constant 2.5 second inter-stimulus interval, listen again, repeat again, and so forth for ten repetitions of the same stimulus; a tone after the tenth repetition signalled the presentation of a new stimulus. For example: the stimulus /best/ would be presented to be repeated by the participant during a 2.5 s inter-stimulus interval. This would be repeated until ten instances of /best/ were recorded. A tone would then alert the participant that the next stimulus would be a new word, e.g. /bis/. The recorder used to collect the speech samples was a Marantz PMD670, and its settings were: mono at a sampling frequency of 22.05 kHz and a 16-

bit amplitude resolution. The microphone was a Sennheiser MD425, and was placed on a microphone stand at a distance of approximately 20 cm from each participant's mouth. For each participant 80 sound files were recorded (8 monosyllables x 10 repetitions), yielding a potential total of 1,920 sound files (80 x 24 participants) for analysis. Incorrect or incomplete productions (truncated by the inter-stimulus interval) were discarded, as well as a number of productions, which did not warrant an acoustic analysis due to voice quality (whisper/breathy). These amounted to 119 sound files, leaving a total of 1,801 sound files for acoustic analysis.

## 2.4 Data Analysis

The acoustic analyses were carried out with Praat (Boersma & Weenink 2008). TextGrid files were used to mark the acoustic onset and offset of the speech productions in order to calculate the *utterance duration*. Utterances with unreleased plosives in word-final position were measured to the acoustic offset of the speech signal; e.g. to the end of the final glottal pulse or to the breathy vowel offset. The *vowel duration* was measured between the first glottal pulse and the final glottal pulse. This interval was also used to calculate the temporal midpoint of the vowel.

Figure 1 shows an overview of these points for the stimulus /best/. The start point of the vowel interval was marked at the first glottal pulse as seen in the spectrogram; the peak of the first excitation within the sound pressure waveform was selected. The end point was similarly marked based on the visual examination of both spectrogram and sound pressure waveform: after the last glottal pulse as seen in the spectrogram, the end of the last periodic excitation of the sound pressure waveform was selected. In cases where the final consonant was voiced, visual

examination was extended to the acoustic energy distribution of higher frequencies, in particular around the second formant and its end.

<Figure 1 about here>

The frequencies of the first three formants were measured at the vowel onset and the calculated temporal midpoint. Based on these measures, the *absolute formant frequency change* in F1, F2, and F3 (see Eq. 1) as well as the *mean rate of formant frequency change* in F1, F2, and F3 (see Eq. 3) were calculated. Absolute formant frequency changes and the mean rates of change for F1, F2 and F3 were also normalised (see Eq. 2 and Eq. 4, respectively). *F2 locus equations* (e.g. Sussman 1994; Sussman et al. 1991, 1997, 1998; Iskarous et al. 2010; Lindblom & Sussman 2012) were derived from the F2 measurements obtained at the vowel onset (first glottal pulse) and the temporal midpoint of the vowel (see Eq. 5). This method of measuring the F2 midpoint at the temporal midpoint varies from those adopted in other investigations. For example, some studies have adopted visual inspection methods to determine steady state vowel midpoints or the maxima/minima points within a vowel's formant structure (e.g. Fowler 1994; Lindblom et al. 2007; Sussman et al. 1991; Sussman 1994), while others have measured vowel targets at a specified time point from the first glottal pulse/plosive release of a CVC syllable (e.g. Nearey & Shammass 1987; McLeod et al. 2001). It also varies from a hybrid approach which is based on both visual inspection and the use of temporal vowel midpoints (e.g. Sussman et al. 1991; Fowler 1994). One advantage of the adopted method is the increased consistency in conducting measurements across different CV contexts. Linear regression analyses based on the vowel midpoints and vowel onsets yielded slope, y-intercept, and  $R^2$  values for all speech tokens.

## 2.5 Intra-rater reliability

All measures were subjected to an intra-rater reliability test. A data sample (ca. 10 %) of the recordings was analysed by the first author for a second time approximately one year after the first measurements were taken. One year was considered to be a significant time lapse to ensure robust intra-rater reliability measurements. Table 2 summarises the mean and standard deviation values of utterance and vowel duration, as well as the first three formants at both vowel onset and temporal midpoint. The Pearson correlation values (between 0.945 and 0.998) indicate high levels of agreement between the two sets of measurements.

<Table 2 about here>

## 2.6 Statistical Analysis

The temporal and spectro-temporal measures investigated in the study were analysed statistically using analysis of variance with speaker sex as the between subjects factor. The linear regression analyses representing the F2 locus equations yielded slope, y-intercept, and  $R^2$  values for all speech tokens. The slope values for the F2 locus equations were tested for statistical sex differences using a ‘large-sample Z test for parallelism’ (Kleinbaum & Kupper 1978: 101-102), and the y-intercept values were tested for statistical sex differences using a ‘large-sample Z test for common intercept’ (Kleinbaum & Kupper 1978: 103-105). The former of these two statistical methods tests whether the slopes of two linear regression lines are the same or different, and the latter, whether their y-intercepts are the same or different. If the slopes are the same (i.e. not statistically significantly different), they are described as parallel if they do not share the same

intercept, or as coincident lines if they share the same slope and intercept. Two linear regression lines can share the same intercept but have different slopes, and intersecting regression lines have different slopes and different intercepts (see Kleinbaum & Kupper 1978: 97-98). The difference in the correlation coefficients underlying the  $R^2$  values of the F2 locus equations was tested using Fisher's  $z_r$  transformation test (e.g. Ferguson 1959: 195-6; Fisher 1950).

### **3 Results**

#### **3.1 Utterance and Vowel Duration**

Tables 3 and 4 summarise the mean and standard deviation values for utterance and vowel duration by stimulus item and speaker sex, as well as by speaker sex across all stimuli. The differences between the male and female data were not significant for either utterance duration (mean difference: 9.48 msec; [ $F(1,22) = 0.309, p > .05$ ]) or vowel duration (mean difference: 10.57 msec; [ $F(1,22) = 0.951, p > .05$ ]).

<Tables 3 & 4 about here>

#### **3.2 Absolute formant frequency and normalised frequency changes**

Table 5 summarises the mean values of the absolute formant frequency changes  $\Delta F1$ ,  $\Delta F2$ , and  $\Delta F3$ , and corresponding normalised values by speaker sex. There was no significant speaker sex effect for either  $\Delta F1$  (mean difference: 16.77 Hz; [ $F(1,22) = 1.901, p > .05$ ]) or  $\Delta F3$  values (mean difference: 21.97 Hz; [ $F(1,22) = 1.040, p > .05$ ]), which were lower for male speakers. However, male speakers displayed significantly lower  $\Delta F2$  values than female speakers (mean

difference: 84.96 Hz; [ $F(1,22) = 10.609, p < .005$ ]). The pattern of speaker sex effects was replicated across the normalised values for the three formant frequencies (norm $\Delta F1$  – mean difference: 0.007 [ $F(1,22) = .148, p > .05$ ]; norm $\Delta F2$  – mean difference: 0.032 [ $F(1,22) = 6.069, p < .05$ ]; norm $\Delta F3$  – mean difference: 0.000 [ $F(1,22) = .001, p > .05$ ]).

<Table 5 about here>

### 3.3 Mean rate of formant frequency change and normalised rates of frequency change

Table 6 summarises the mean values of the mean rate of formant frequency changes for rF1, rF2, and rF3, and corresponding normalised values by speaker sex. There was a significant effect of speaker sex on rF2, where female speakers showed faster mean rates of formant change than male speakers (mean difference: 0.68 Hz/msec; [ $F(1,22) = 8.657, p < .01$ ]). There were no significant sex differences for either rF1 (mean difference: 0.10 Hz/msec; [ $F(1,22) = 1.321, p > .05$ ]) or rF3 (mean difference: 0.12 Hz/msec; [ $F(1,22) = 0.326, p > .05$ ]). The pattern of speaker sex effects was broadly similar across the normalised values for the three formant frequencies. However, the sex effects for F2 were diminished, and only approached significance (normrF1 – [ $F(1,22) = 0.000, p > .05$ ]; normrF2 – [ $F(1,22) = 4.003, p = .058$ ]; normrF3 – [ $F(1,22) = 0.209, p > .05$ ]).

<Table 6 about here>

### 3.4 F2 locus equation data

Figure 2 displays the F2 locus equations of the monosyllables grouped by speaker sex. The males showed a significantly steeper slope (0.661) than the females (0.598) [ $Z = 2.84, p < .005$ ]. The y-intercept for the men's data samples (503 Hz) was found to be significantly lower compared to the women's (645 Hz) [ $Z = -3.27, p < .001$ ]. The correlation coefficients underlying the  $R^2$  values for the male ( $r = 0.819$ ) and female ( $r = 0.785$ ) data were also significantly different [ $z_r = 2.02, p < .05$ ].

<Figure 2 about here>

### 3.5 Post-hoc analysis of vowel effects

There is evidence to suggest that vowel quality plays some role in speaker sex differences (Simpson & Ericsson 2003). Therefore, a post-hoc analysis examining vowel effects was also conducted. Vowel duration,  $\Delta F2$  and  $\text{norm}\Delta F2$  values were examined as a function of vowel quality for men and women. Firstly,  $\Delta F2$  and vowel duration values for all stimuli except /bust/ were examined using Pearson's product moment correlation test. This was done separately for the men and women. This was repeated for the  $\text{norm}\Delta F2$  and vowel duration values. Secondly, the whole set of the aforementioned analyses was repeated for the data solely representing /bust/. This yielded eight sets of correlation coefficients, which are provided in Table 7.

<Table 7 about here>

The results summarised in Table 7 suggest a *vowel quality x speaker sex* interaction. The correlation between vowel duration and  $\Delta F2$  values was significant for the female speakers for

all stimuli combined except /bu:st/ ( $r = 0.646, p < .05$ ), but not for male speakers ( $r = -0.010, p > .05$ ). This sex difference was replicated for the correlation between vowel duration and norm $\Delta$ F2 values. However, when considering solely the data obtained for /bu:st/, the correlations between vowel duration and  $\Delta$ F2 values or vowel duration and norm $\Delta$ F2 values were not significant for either women or men.

#### **4 Discussion**

Female speakers exhibited trends for longer utterance and vowel durations than male speakers, but these differences were not significant. This provides only weak support for the first hypothesis and some of the temporal sex differences reported in the literature, where female speakers have been found to exhibit significantly longer vowel and utterance durations (Simpson 1998, 2001; Whiteside 1996). As mentioned in section 2.3, the data elicitation method may have been a contributing factor to this result. Participants listened to the audio recordings of a male speaker and may have accommodated their speech productions to some degree to his productions. Nevertheless, when Simpson & Ericsson (2003) analysed sex-specific durational differences in American English and Swedish their results revealed more complex interdependencies. While the mean durations of a series of sentences did not vary as a function of speaker sex for either English or Swedish, a plot of the percentage differences in segment durations revealed quite systematic sex differences. In those syllables which carried ‘a degree of stress’ (Simpson & Ericsson 2003: 1114), female vowel durations trended for longer values. On the other hand, many of the back vowels trended for longer durations when produced by male speakers. Male speakers also trended for longer consonant productions than female speakers. The stimuli in the current study contained predominantly intrinsically short front vowels and one long

back vowel (/u:/) as opposed to Simpson (2001), who analysed diphthongs, and Whiteside (1996) and Jacewicz et al. (2010), who analysed a range of vowels. For one long back vowel (/u:/) male speakers may have exhibited longer vowel durations, which may have diminished the overall sex difference. Further investigation was conducted as a function of the different stimulus items to investigate this, and an analysis of /bust/ found this to be the case; males exhibited the trend of a longer mean vowel duration value than the females (see Table 4). In contrast, the remaining set of short front vowels displayed the reverse pattern; females displayed a trend for longer vowel duration values for all stimuli and a longer mean vowel duration value (219.72 msec) than the males (207.91 msec). These results corroborate those of Simpson & Ericsson (2003) suggesting that sex differences in temporal parameters such as vowel duration are determined by an intricate interaction between factors such as linguistic stress, vowel quality and phonetic context, and might therefore explain the results reported here.

There were no significant sex differences in the spectral and spectro-temporal parameters associated with F1 and F3 (see Tables 5 and 6). However, significant sex differences were found for the spectro-temporal parameters associated with F2:  $\Delta F2$  and corresponding normalised values (see Table 5), the rate in the absolute change in F2 (see Table 6), and the slope of the F2 locus equations representing the speech samples of the women and men (Figure 2). Here women displayed higher values for  $\Delta F2$ , normalised  $\Delta F2$  and  $rF2$ , and shallower slopes for the F2 locus equations. The higher values for  $\Delta F2$  and normalised  $\Delta F2$ , and the shallower slopes of the F2 locus equations are both indicative of lower degrees of coarticulation in the women's speech samples.

Concerning the sex differences found in the distribution of F2 related parameters, the stimuli used in this study may provide some further insights, as they contained mainly high front and mid front vowels /i, ε/, and one back vowel /u:/. Hillenbrand, Getty, Clark, & Wheeler (1995) and Traunmüller (1988) found that sex differences in high front vowels are particularly prominent in F2 (see also Lee et al. 1999; Whiteside 2001). In comparison, the differences in F2 are much smaller for high back vowels, for which female and male values are relatively close to each other (e.g. Simpson 2009). Whiteside (2001), who further investigated the data collected by Lee et al. (1999), reported that data for the back vowel (/u:/) was indicative of a more fronted (palatalized) quality for the female speakers when compared to their male peers. These sex differences in the second formant, as well as the predominantly high front vowels in the data set investigated here, may explain the consistent speaker sex effects found for the F2 related data (absolute and normalised formant frequency changes, mean rate of formant change, and F2 locus equations). Female speakers showed patterns of higher mean values of the absolute formant change in F1, F2, and F3 than male speakers, which was significant for  $\Delta F2$ . The higher absolute formant changes for female speakers provide some support to the hypothesis of lower degrees of coarticulation. However, Simpson (2001) compared acoustic and articulatory data and found that although the articulatory distances covered by the female speakers during the production of the diphthong /aɪ/ in 'light' are smaller for female than for male speakers, the absolute formant changes were larger. A similar discrepancy was observed for the articulatory speeds seen in the mean rates of the formant changes. This suggests that the lower degrees of coarticulation, as

indicated by the higher absolute formant change values, may be in part attributed to the different vocal tract dimensions of male and female speakers.

Female speakers displayed faster mean rates of formant change in F1 (rF1), F2 (rF2), and F3 (rF3). This difference was significant for rF2, marginally significant for normalised rF2, and mirrors the results of the absolute formant changes in F2 ( $\Delta F2$ ). However, these faster mean rates of formant change do not necessarily imply faster articulatory movements for female speakers, but may be an acoustic dynamic consequence of different vocal tract dimensions (see Simpson 2001). On the basis of differences in vocal tract dimensions and the different articulatory postures involved in the production of different vowels, the links between absolute formant changes and mean rates of formant change need to be interpreted in the context of vowel duration and vowel quality. There were no significant sex differences in vowel duration. However, when the data were probed further, there was some indication that the women and men exhibited different patterns of vowel duration as a function of vowel quality; women trended for longer durations for the front vowels, whereas men displayed a pattern of longer vowel durations for the back vowel /u:/. When the  $\Delta F2$  data were analysed further in a post-hoc analysis as a function of vowel quality, a significant sex difference emerged which might explain why the  $\Delta F2$  values were higher for the women (see Table 7). When  $\Delta F2$  and vowel duration data were examined for all stimuli except /bu:st/ using Pearson's product moment correlation test for each sex group, the women's data displayed a significant correlation ( $n = 13$ ,  $r = 0.646$ ,  $p < .05$ ). This was in contrast to the men's data which displayed no significant correlation ( $n = 11$ ,  $r = -0.010$ , ns). When the correlations between  $\Delta F2$  and vowel duration data were examined only for /bu:st/ neither the women ( $n = 13$ ,  $r = 0.239$ , ns) nor the men ( $n = 11$ ,  $r = -0.560$ , ns) displayed

significant correlations. These results were replicated when normalised  $\Delta F2$  and vowel duration data were examined (see Table 7). These data provide further support for the complex interaction between vowel quality and the spectro-temporal dynamics observed for men and women speakers; the data set with the high front vowels displayed higher  $\Delta F2$  and higher normalised  $\Delta F2$  values for the women which appeared to be a function of vowel duration and vowel quality. The fact that the stimuli in the current study were dominated by high/front vowels might go some way to explain the sex differences reported here.

For the F2 locus equations, women displayed a significantly higher y-intercept value which can be explained by vocal tract differences (e.g. Fitch & Giedd 1999, Lee et al. 1999, Whiteside 2001), and will not be discussed further. The correlation coefficients underlying the  $R^2$  values for the men were significantly higher than those for the women, but will not be discussed further as they are not the focus of this study. The F2 locus equations data in the current study centred on the slope values as a measure of coarticulation, and were used to assess the pattern and extent of sex differences, which was the focus of this study. Results indicated significantly steeper slope values for men (0.661) compared to the women (0.598) (see Figure 2). The mean slope values reported here are lower than those reported in other studies (see Section 1.2 above). There are several possible explanations for this difference. One reason could be the method which was used in the current study to measure the F2 vowel midpoint; this varied from those of earlier reports (see Section 2.3 above). The temporal midpoint of the vowel was used to measure the F2 vowel midpoint in the current study. It is therefore possible that temporal midpoint measurements did not always coincide with the vowel nucleus (steady state, or maxima/minima points) for all data samples. In cases where the vowel nucleus was temporally surpassed, it is

therefore possible that the word final velar and alveolar consonants, and consonant clusters (e.g. /k/, /s/, /z/, /st/, /t/, /d/) might have had some influence on the temporal midpoint values which might have resulted in shallower slopes. Another possible source of variation is individual speaker differences; this suggestion is supported by the range and variation in slope values reported across different speakers in different studies (see Table 1). However, another possible source for the difference could be vowel context. The majority of vowels in the current study were high/front vowels. Some studies have systematically investigated the F2 locus equations of velar plosives as a function of vowel context (e.g. [g] palatal (front) versus [g] velar (back) vowel contexts) where shallower slopes for /g/ have been reported for front/palatal vowel contexts compared to back/velar contexts (Fowler 1994; Sussman et al. 1997, 1998). However, there is also evidence to suggest that vowel context also affects the F2 locus equation slopes of /bV/ sequences, and that /bV/ sequences in front vowel contexts have shallower slopes (e.g. approximately 0.65 – estimated from Figure 3, Fowler 1994: 602) compared to those produced in back vowel contexts (approximately 0.8 – estimated from Figure 3, Fowler 1994: 602). What is worth noting here is that the aforementioned slope value for the front vowel context (0.65) is closer to the values reported in this study (0.661 and 0.598 for males and females respectively).

The steeper slope for the male speakers found in this study is in line with McLeod et al. (2001), who reported significantly steeper F2 locus equation slopes for male than female speakers (see also the trends reported by Löfqvist (1999)). This suggests greater degrees of anticipatory coarticulation for male than for female speakers in the current study, as the F2 values at consonantal release were closer to and varied more systematically with the F2 values at vowel

midpoint. The greater degree of anticipatory coarticulation for male speakers may be indicative of a different pattern of movement synchronisation; the data suggest that male speakers may have initiated the vowel gesture earlier than female speakers (Simpson 2003), which would also contribute to smaller absolute formant change values in speech stimuli of the current data set which predominantly consisted of high front vowels.

The current study examined /bV/ sequences which, in theory, allow for maximum degrees of coarticulation between the voiced bilabial plosive and the ensuing vowel due to the independence of the labial and lingual articulators. Therefore, in light of the predominantly high front vowels in the current data set, it might be argued that the results are not truly sex-specific differences in coarticulation, but merely consequences of differences in vowel space sizes between women and men. While there may be some degree of influence on the absolute formant change in F2 given the overall higher F2 values and their wider distribution for the female data, the latter of which may be caused by greater interharmonic spacing (e.g. Diehl, Lindblom, Hoemeke & Fahey 1996, Simpson 2011), these effects are attenuated when analysing F2 locus equations. The higher F2 values at both vowel onset and midpoint are largely contextually preserved in F2 locus equations, but are removed when calculating the absolute formant change in F2. Higher F2 onset and F2 midpoint pairs for front vowels are represented as data points further along the linear regression line of the F2 locus equations, but do not affect the steepness of the slope (see Figure 2: the data points for the female speakers in the left panel are distributed further along the regression line towards the higher F2 values). As the F2 locus equations of this data set exhibited lower degrees of coarticulation for the women (shallower slope values), which complements the effect found in the absolute formant change in F2, this suggests that the effects

reported for  $\Delta F_2$  may be similarly attributed to sex-specific differences in coarticulation and are not exclusively a consequence of differences in vowel space sizes.

## 5 Conclusions

The sex differences previously reported in the literature were corroborated in the current study: although there was only weak support for the first hypothesis, men trended for shorter acoustic vowel durations (e.g. Simpson 2003, Whiteside 1996), and this appeared to be interacting with vowel quality differences (e.g. Simpson & Ericsson 2003). In addition, women displayed significantly faster mean rates of formant change for F2 (e.g. Simpson 2001), and significantly lower degrees of coarticulation than men as indexed by  $\Delta F_2$ , normalised  $\Delta F_2$  and shallower F2 locus equation slopes (e.g. McLeod et al. 2001, and see trends in Löfqvist 1999), thereby supporting both the second and third hypotheses. The sex differences in the formant frequency changes, and the higher levels of systematic variation between vowel onsets and vowel midpoints in the data were interpreted as in part, being indicative of differences in the synchronisation of articulatory gestures: it was suggested that male speakers may initiate the vowel gesture earlier relative to the release of the consonant (see Simpson 2002, 2003). The predominantly high front vowels used in the current data set may also explain the significant sex differences found in this investigation. In order to further elucidate the role of factors such as sex-determined biological differences in the spectro-temporal dynamics of speech and in coarticulation, further systematic acoustic and articulatory investigations with a range of stimuli and phonetic contexts are warranted.

## References

- Baayen, R. Harald, Richard Piepenbrock & Hedderik van Rijn. 1993. The CELEX database (Release 1) [CD Rom]. Philadelphia PA.
- Boersma, Paul & David Weenink. 2008. Praat: Doing phonetics by computer (version 5.0.35). <http://www.praat.org> (accessed 7 October 2008).
- Brancazio, Lawrence & Carol A. Fowler. 1998. On the relevance of locus equations for production and perception of stop consonants. *Perception & Psychophysics* 60(1), 24–50.
- Byrd, Dani. 1992. Preliminary results on speaker-dependent variation in the TIMIT database. *Journal of the Acoustical Society of America* 92(1), 593–596.
- Byrd, Dani. 1994. Relations of sex and dialect to reduction. *Speech Communication* 15, 39–54.
- Chang, Soo-Eyn, Ralph N. Ohde & Edward G. Conture. 2002. Coarticulation and formant transition rate in young children who stutter. *Journal of Speech, Language, and Hearing Research* 45(4), 676–688.
- Chartrand, Tanya L & John A. Bargh. 1999. The chameleon effect: The perception-behavior link and social interaction. *Journal of Personality and Social Psychology* 76(6), 839–910.
- Diehl, Randy L., Björn Lindblom, Kathryn A. Hoemeke & Richard P. Fahey. 1996. On explaining certain male-female differences in the phonetic realization of vowel categories. *Journal of Phonetics*, 24(2), 187–208.
- Ferguson, George A. 1959. *Statistical analysis in psychology and education*, 5th edn. London: McGraw-Hill Book Company.
- Fisher, Ronald A. 1950. *Statistical methods for research workers*, 11th edn. Edinburgh: Oliver and Boyd.
- Fitch, W. Tecumseh & Jay Giedd. 1999. Morphology and development of the human vocal tract: A study using magnetic resonance imaging. *Journal of the Acoustical Society of America* 106(3), 1511–1522.
- Fowler, Carol A. 1994. Invariants, specifiers, cues: An investigation of locus equations as information for place of articulation. *Perception & Psychophysics* 55(6), 597–610.
- Fowler, Carol A. & Elliot L. Saltzman. 1993. Coordination and coarticulation in speech production. *Language and Speech* 36, 171–195.
- Giles, Howard, Nikolas Coupland & Justine Coupland. 1991. Accommodation theory: Communication, context and consequence. In H. Giles, N. Coupland & J. Coupland (eds.),

*Contexts of Accommodation: Developments in Applied Sociolinguistics*, 1–68, Cambridge: Cambridge University Press.

- Henton, Caroline G. & R. Anthony W. Bladon. 1985. Breathiness in normal female speech: Inefficiency versus desirability. *Language and Communication* 5(3), 221–227.
- Herrmann, Frank. 2011. *Syllable frequency and speaker sex effects on coarticulation: An acoustic investigation*. Ph.D. thesis. Department of Human Communication Sciences, University of Sheffield.
- Hillenbrand, James M., Laura A. Getty, Michael J. Clark & Kimberlee Wheeler. 1995. Acoustic characteristics of American English vowels. *Journal of the Acoustical Society of America* 97(5), 3099–3111.
- Iskarous, Khalil, Carol A. Fowler & Douglas H. Whalen. 2010. Locus equations are an acoustic expression of articulator synergy. *Journal of the Acoustical Society of America* 128(4), 2021–2032.
- Jacewicz, Ewa, Robert A. Fox & Lai Wei. 2010. Between-speaker and within-speaker variation in speech tempo of American English. *Journal of the Acoustical Society of America* 128(2), 839–850.
- Junqua, Jean-Claude. 1993. The Lombard reflex and its role on human listeners and automatic speech recognizers. *Journal of the Acoustical Society of America* 93(1), 510–524.
- Kleinbaum, David G. & Lawrence L. Kupper. 1978. *Applied regression analysis and other multivariate methods*. Boston, MA: Duxbury Press.
- Krull, Diana. 1987. Second formant locus patterns as a measure of consonant-vowel coarticulation. *Phonetic Experimental Research at the Institute of Linguistics, University of Stockholm* V, 43–61.
- Lee, Sungbok, Alexandros Potamianos & Shrikanth Narayanan. 1999. Acoustics of children's speech: Developmental changes of temporal and spectral parameters. *Journal of the Acoustical Society of America* 105(3), 1455–1468.
- Lindblom, Björn. 1963. Spectrographic study of vowel reduction. *Journal of the Acoustical Society of America* 35, 1773–1781.
- Lindblom, Björn, Augustine Agwuele, Harvey M. Sussman, & Elisabet E. Cortes. 2007. The effect of emphatic stress on consonant vowel articulation. *Journal of the Acoustical Society of America* 121(6), 3802–3813.
- Lindblom, Björn & Harvey M. Sussman. 2012. Dissecting coarticulation: How locus equations happen. *Journal of Phonetics* 40(1), 1–19.

- Löfqvist, Anders. 1999. Interarticulator phasing, locus equations, and degree of coarticulation. *Journal of the Acoustical Society of America* 106(4), 2022–2030.
- Lombard, Étienne. 1911. Le signe de l'élevation de la voix. *Ann Maladies Oreille Larynx, Nez, Pharynx*(37), 101–119.
- McLeod, Amy, Megan Baillargeon, Dale E. Metz, Nicholas Schiavetti, & Robert L Whitehead. 2001. Locus Equations as a source of relational invariance for stop place categorization: A direct replication. *Contemporary Issues in Communication Science and Disorders* 28, 98–103.
- Nearey, Terrence M. & S. E. Shammass. 1987. Formant transitions as partly distinctive invariant properties in the identification of voiced stops. *Canadian Acoustics* 15(4), 17–24.
- Simpson, Adrian P. 1998. *Phonetische Datenbanken des Deutschen in der empirischen Sprachforschung und der phonologischen Theoriebildung*. In K. J. Kohler (Ed.), *Arbeitsberichte des Instituts für Phonetik und digitale Sprachverarbeitung (AIPUK)* 33. Kiel.
- Simpson, Adrian P. 2001. Dynamic consequences of differences in male and female vocal tract dimensions. *Journal of the Acoustical Society of America* 109(5), 2153–2164.
- Simpson, Adrian P. 2002. Gender-specific articulatory-acoustic relations in vowel sequences. *Journal of Phonetics* 30, 417–435.
- Simpson, Adrian P. 2003. Possible articulatory reasons for sex-specific differences in vowel duration. *Proceedings of the 6th International Seminar on Speech Production* (pp. 261–266). Sydney.
- Simpson, Adrian P. 2009. Phonetic differences between male and female speech. *Language and Linguistics Compass* 3(2), 621–640.
- Simpson, Adrian P. 2011. Is there a relationship between acoustic vowel space size and fundamental frequency? In *17th International Congress of Phonetic Sciences, Hong-Kong, City University of Hong-Kong (ICPhS 17)*, vol. 1, 1854–1857.
- Simpson, Adrian P. & Christine Ericsdotter. 2003. Sex-specific durational differences in English and Swedish. In *15th International Congress of Phonetic Sciences, Barcelona, Universitat Autònoma de Barcelona (ICPhS 15)*, vol. 1, 1113–1116.
- Sussman, Harvey M. 1994. The phonological reality of locus equations across manner class distinctions: Preliminary observations. *Phonetica* 51, 119–131.
- Sussman, Harvey M., Nicola Bessell, Eileen Dalston & Tivoli Majors. 1997. An investigation of stop place articulation as a function of syllable position: A locus equation perspective. *Journal of the Acoustical Society of America* 101(5), 2826–2838.

- Sussman, Harvey M., Eileen Dalston & S. Gumbert. 1998. The effect of speaking style on a locus equation characterization of stop place articulation. *Phonetica* 55, 204–225.
- Sussman, Harvey M., Helen A. McCaffrey & Sandra A. Matthews. 1991. An investigation of locus equations as a source of relational invariance for stop place categorization, *Journal of the Acoustical Society of America* 90(3), 1309–1325.
- Tabain, Marija. 2000. Coarticulation in CV syllables: A comparison of locus equation and EPG data. *Journal of Phonetics* 28, 137–159.
- Tabain, Marija. 2002. Voiceless consonants and locus equations: A comparison with electropalatographic data on coarticulation. *Phonetica* 59, 20–37.
- Traunmüller, Hartmut. 1988. Paralinguistic variation and invariance in the characteristic frequencies of vowels. *Phonetica* 45, 1–29.
- Whiteside, Sandra P. 1996. Temporal-based acoustic-phonetic patterns in read speech: Some evidence for speaker sex differences. *Journal of the International Phonetic Association* 26, 23–40.
- Whiteside, Sandra P. 2001. Sex-specific fundamental and formant frequency patterns in a cross-sectional study. *Journal of the Acoustical Society of America* 110(1), 464–478.