



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/98411/>

Version: Accepted Version

Article:

Flanagan, Lindsey A, Wright, John J, Roessler, Maxie M et al. (2016) Re-engineering a NiFe hydrogenase to increase the H₂ production bias while maintaining native levels of O₂ tolerance. *Chemical Communications*. pp. 9133-6. ISSN: 1364-548X

<https://doi.org/10.1039/C6CC00515B>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Re-engineering a NiFe hydrogenase to increase the H₂ production bias while maintaining native levels of O₂ tolerance

Received 00th January 20xx,
Accepted 00th January 20xx

Lindsey A. Flanagan,^a John J. Wright,^b Maxie M. Roessler,^b James W. B. Moir^c and Alison Parkin^{*a}

DOI: 10.1039/x0xx00000x

www.rsc.org/

Naturally occurring oxygen tolerant NiFe membrane bound hydrogenases have a conserved catalytic bias towards hydrogen oxidation which limits their technological value. We present an *Escherichia coli* Hyd-1 amino acid exchange that apparently causes the catalytic rate of H₂ production to double but does not impact the O₂ tolerance.

Biology offers a source of inspiration for the discovery of highly active, precious-metal-free molecular H₂ production (proton reduction) catalysts, because many microbes have evolved to produce hydrogen from protons and electrons (2H⁺ + 2e⁻ → H₂) using enzymes known as hydrogenases.^{1, 2} Most nickel-iron hydrogenases, enzymes containing a NiFe bimetallic active site, have limited application as H₂ catalysts in “one-pot” light-driven water-splitting devices with the O₂-generating anode and H₂-producing cathode in same compartment. This is because O₂ binds at the active site to produce the catalytically inactive Ni-A species (such inhibition is described as “O₂ sensitivity”). There is a well-studied subclass of NiFe hydrogenases which retain catalytic activity in O₂, these are the membrane-bound, O₂ tolerant NiFe hydrogenases (O₂-tolerant MBH).³ Unfortunately, the evolution of O₂ tolerance was concomitant with the evolution of a bias towards H₂ oxidation.³ In this study we have rationally redesigned an O₂-tolerant MBH to yield a catalyst with an increased H₂ production : H₂ oxidation bias but native levels of O₂ tolerance.

NiFe MBHs contain a NiFe active site which is buried within a “large” protein subunit and three iron sulfur (FeS) clusters ligated by a “small” protein subunit.^{1, 2} The FeS clusters act as an electron transfer conduit, mediating the flow of electrons between the protein surface of the small subunit and the buried active site. While the active site binding pocket is remarkably well conserved in all NiFe hydrogenases, it is

plasticity in the FeS relay which appears to control both the catalytic bias (the ratio of H₂ oxidation to production activity) and O₂ sensitivity in NiFe MBH.²




Fig. 1 (A) Sequence alignment showing the conservation of large subunit residue 73 amongst oxygen tolerant and oxygen sensitive NiFe hydrogenases. Ec= *E. coli*; Se= *Salmonella enterica*; Re= *Ralstonia eutropha*; Ae= *Aquifex aeolicus*; Av= *Allochroamium vinosum*; Dg= *Desulfovibrio gigas*; Dv= *Desulfovibrio vulgaris*; Db= *Desulfomicrobium baculatum*. (B) Crystal structures of (i) an oxygen tolerant MBH (PDB: 3RGW) and (ii) and oxygen sensitive NiFe hydrogenase (PDB: 4UD2), showing the location of residue 73 relative to the active site and the proximal cluster. The H-bonding capacity of residue 73 is denoted by the dashed black lines. *E. coli* numbering is used throughout.

All O₂ tolerant MBH contain an unusual 4Fe3S “proximal” (closest to the active site) cluster, while the O₂ sensitive MBH have a standard 4Fe4S centre in the same position.^{2, 4-6} The “medial” (middle of the relay) FeS centre is always a 3Fe4S centre but despite the structural invariance this centre still plays a vital role in enabling O₂ tolerant MBH to catalyse H₂ oxidation in the presence of O₂.⁷ Armstrong and co-workers have hypothesised that the “distal” (furthest from the active site) FeS cluster controls the thermodynamic driving force which is needed to induce catalysis in a MBH.⁸⁻¹⁰ One of the reasons that the proximal cluster was disregarded as an important tuning point for catalysis is because previous mutations within both *Ralstonia* MBH and *Escherichia coli*

^a Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK, E-mail: alison.parkin@york.ac.uk; Tel: +44 1904 322561

^b School of Biological and Chemical Sciences, Queen Mary University of London, Mile End Road, London E1 4NS, UK

^c Department of Biology, University of York, York YO10 5DD, UK.

† Electronic Supplementary Information (ESI) available: Experimental protocol details and Fig. S1 to S7. See DOI: 10.1039/x0xx00000x

hydrogenase-1 were not noted to increase the H₂ production of these O₂ tolerant hydrogenases.^{5, 11} However, other researchers have shown that the rate of catalysis in a [NiFe] hydrogenase can be altered by changes to the proximal cluster. An increase in the cellular level of H₂ production by the cyanobacteria *Nostoc punctiforme* accompanies the replacement of the 4Fe4S proximal cluster of the O₂-sensitive HupSL uptake hydrogenase with a 3Fe4S cluster.¹² Conversely, a drop in the H₂ production activity of *Desulfovibrio fructosovorans* variants¹³ has been attributed to single site amino acid changes having an impact on the proximal cluster.⁸




Fig. 2 Cyclic voltammetry (CV) measurements at 5 mV s⁻¹ of (A) Native *Escherichia coli* hydrogenase-1, and (B) the E73Q variant under gas atmospheres of both 3% H₂ in N₂ (dotted lines) and 0% H₂ (100% N₂, solid lines). Other experimental conditions: pH 4.5; 37 °C, electrode rotation rate 3500 rpm; carrier gas N₂ and total gas flow rate 100 scc min⁻¹. (C) The difference in H₂ production activity is emphasised through an overlay of the averaged forward and back scan sweeps of the CVs in the top two panels, the selected region is highlighted by the dotted grey boxes in A and B.

Escherichia coli (*E. coli*) produce two different membrane-bound hydrogenases, the O₂ tolerant hydrogenase-1 (Hyd-1) and the O₂ sensitive hydrogenase-2 (Hyd-2).¹⁴ Sequence comparisons between *E. coli* Hyd-1 and Hyd-2 can help suggest which amino acid residues play a vital role in controlling how a hydrogenase reacts with substrate and/or inhibitors. The conserved presence of a glutamine (Q) in a position between the active site and proximal cluster of all O₂ sensitive hydrogenases (position 73 using *E. coli* Hyd-1 numbering), contrasts with the occurrence of a glutamic acid (E) in the

same position in most O₂ tolerant enzymes (Fig. 1). In previous work¹⁵ on *Salmonella* Hyd-5 we found that an E73A variant sustained lower levels of H₂ oxidation activity in the presence of O₂ when compared to Native enzyme, although the O₂ inhibition remained substantially reversible. Since the mechanism of O₂ tolerance relies on electron transfer between the proximal cluster and the active site,⁴⁻⁶ we hypothesised that E73 might influence the proximal cluster redox potential. As a result of this, we have explored how an E73Q amino acid exchange in *E. coli* Hyd-1 impacts the enzyme's reactivity with both H⁺ and H₂ substrates and the inhibitor O₂.

Methylene blue (MB) assays in H₂-saturated buffer were used to compare the enzyme turnover rate of purified E73Q and Native enzyme at pH 4.5, 25 °C.¹⁶ Very similar H₂ oxidation rates were measured for both hydrogenases (Native: 21 ± 4 s⁻¹; E73Q: 22 ± 3 s⁻¹), suggesting that the amino acid exchange does not impact on the enzyme's ability to catalyse H₂-uptake at the MB redox potential (voltammetry measurements (data not shown) determined E_{mid}(MB) = +0.113 V vs SHE at pH 4.5, 25 °C). Protein film electrochemistry experiments (Fig. 2A and 2B) were then performed to explore the enzyme activity over a wider potential range and under different levels of H₂. To further facilitate the comparison of H₂ production activity, Fig. 2C shows overlay plots of the data in Fig. 2A and Fig. 2B in the region -0.45 to -0.15 V vs SHE after background electrode charging currents have been removed by averaging the current measured in forward and back scan sweeps.

From Fig. 2A and 2B, comparing the ratio of the maximum positive (H₂ oxidation) current measured under 3% H₂ to the maximum negative (H⁺ reduction/H₂ production) current measured under 0% H₂ immediately reveals that the catalytic bias of the E73Q variant is significantly different to that of Native enzyme. A value of approx. 16 is generated from calculating $i_{\max}(\text{H}_2 \text{ oxidation, 3\% H}_2) \div i_{\max}(\text{H}_2 \text{ reduction, 0\% H}_2)$ for Native enzyme, whereas the equivalent ratio for E73Q is approx. 8. If we assume that the rate of H₂ oxidation is unaffected by the amino acid exchange, then this means that the rate of catalytic H₂ production by E73Q is approximately double that of Native enzyme.




Fig. 3 Inhibition and recovery from 3% O₂ measured at +0.113 V vs SHE with 3% H₂ throughout. The black line shows data for Native *Escherichia coli* Hyd-1 while the red line depicts the measurement for the E73Q variant. Other experimental conditions: pH 4.5; 37 °C, electrode rotation rate 3500 rpm; carrier gas N₂ and total gas flow rate 100 scc min⁻¹.

Although the E to Q amino acid exchange has a dramatic impact on the catalytic activity, the oxygen tolerance remains

unchanged. Fig. 3 shows that for both Native and E73Q hydrogenase the same proportion of H₂ oxidation current is sustained at +0.113 V vs SHE when the experimental headgas is changed from 3% H₂ in N₂ to 3% H₂, 3% O₂ and 94% N₂. When the inhibitory O₂ is removed from the gas flow, both enzymes rapidly recover almost the full extent of their pre-O₂ activity, demonstrating characteristic “O₂ tolerant” behaviour. We therefore suggest that we have artificially created an *E. coli* Hyd-1 variant with twice the rate of H₂ production of the Native enzyme, but equal levels of O₂ tolerance.

In order to probe the origin of the novel reactivity of E73Q, molecular biology was used to create *E. coli* strains to encode the Hyd-1 variants E73K, E73N and E73A (see Fig. S1 ESI). Unfortunately, only very low levels of the proteins E73K and E73N could be purified (see Fig. S2 ESI). Electrochemical analysis of E73A showed the same phenotype as was previously observed for the *Salmonella* Hyd-5 variant,¹⁵ namely Native-like catalytic activity and increased sensitivity to O₂ (see Fig. S3, ESI). Dance¹⁷ has rationalised that the loss of O₂ tolerance in a *Salmonella* Hyd-5 E73A variant is attributable to the decoupling of the proton transfer network between the active site and proximal cluster. This decoupling was proposed to arise because the non-polar alanine is incapable of proton donation/acceptance. However, our E73Q variant is O₂ tolerant despite the fact that glutamine does not have an acidic or basic side chain. Instead, a glutamine in this position would be expected to participate in extensive H-bonding, based on the structure of O₂ sensitive hydrogenases (Fig. 1). We therefore hypothesise that the critical function of residue 73 in the O₂ tolerance mechanism is to participate in stabilising the H-bonding interaction between the active site and the proximal cluster (Fig. 1), and a direct role in proton exchange is not required.

Modifications to a NiFe hydrogenase gas channel have previously been demonstrated to impact on the reactivity of the enzyme,¹ but there is no evidence that the catalytic bias of E73Q can be ascribed to a gas channel effect. Firstly, Fig. 3 suggests that O₂ access to the NiFe active site is unaffected by the E to Q amino acid exchange. Further experiments also suggest that H₂ movement through the enzyme is not altered in E73Q relative to Native enzyme. Using data extracted from cyclic voltammograms measured under different percentages of H₂ (Fig. S4, ESI) permits a Hanes Woolf analysis to be conducted (Fig. S5A, ESI). The Michaelis constant for H₂, $K_M(\text{H}_2)$, at +0.113 V vs SHE is thus calculated for Native and E73Q, yielding very similar values of $4.2 \pm 1.0 \mu\text{M}$ and $3.8 \pm 1.7 \mu\text{M}$, respectively (average value \pm standard error, $n=4$). The product inhibition constant (K_i) for H₂ production at -0.285 V vs SHE has also been analysed (Fig. S5B, ESI), for Native enzyme and E73Q, with $9.5 \pm 1.7 \mu\text{M}$ for the former and $4.6 \pm 0.8 \mu\text{M}$ for the latter. This difference is to be expected, since the variants have similar H₂ production activity under 3% H₂ despite the greater H₂ production activity of E73Q under 0% H₂ (Fig. 2C).

Since the H-bonding network around E73 in an O₂ tolerant NiFe MBH is different to that around the equivalent Q in an O₂ sensitive enzyme (Fig. 1), we also explored whether changing

the pH revealed any characteristic differences between Native and E73Q hydrogenase. As shown in Fig. S6, ESI, apart from the enhanced H₂ production activity of E73Q enzyme, no substantial change in the pH profile is detected, including in the high potential region where reversible formation of the “Ni-B” OH⁻ bound Ni(III) oxidised inactive state, is observed. In order to further confirm this, the experiments shown in Fig. S7, ESI, were conducted, permitting extraction of the potential of maximum rate of reactivation, E_{switch} , which is the same for both enzymes, as shown in Fig. S8, ESI. It is therefore difficult to rationalise the changes in catalytic bias in light of proton availability.




Fig. 4 Potentiometric titration of Hyd-1 E73Q. The $g = 1.97$ peak of the $[\text{Fe}_4\text{S}_3\text{Cys}_2]^{3+}$ EPR signal (inset) was monitored as a function of potential and fitted to the one-electron Nernst equation (solid line, $R=0.9923$).

Following the careful elimination of substantial gas or proton transfer pathway changes as the possible origin of changes in reactivity, it is logical to speculate that changes to the iron-sulfur relay underlie the differences between the catalytic bias of the E73Q and Native hydrogenases. We have explored this by carrying out an electron paramagnetic resonance (EPR) redox titration¹⁸ on E73Q (Fig. 4 and Fig. S9), monitoring the signal corresponding to EPR-visible “super-oxidised” state of the proximal cluster. This has yielded a midpoint potential of $+211 \pm 10 \text{ mV}$ at pH 7 for the transition of the proximal cluster between the super-oxidised state at very high potentials and the EPR-silent “oxidised” state at intermediate potentials ($[\text{Fe}_4\text{S}_3\text{Cys}_2]^{3+} \rightarrow [\text{Fe}_4\text{S}_3\text{Cys}_2]^{2+} + 1e^-$). The potential for this proximal-cluster process is very similar to that previously reported for Native hydrogenase, $+230 \pm 15 \text{ mV}$ at pH 6.¹⁹ We consider that it is appropriate to compare these midpoint potential values despite the difference in pH because it has been shown for the *Aquifex aeolicus* MBH that the high-potential redox transition of the proximal cluster is not affected by pH over the range 6.4 to 7.4, and pH-related changes to the lower potential $[\text{Fe}_4\text{S}_3\text{Cys}_2]^{2+/1+}$ redox couple only occur at $\text{pH} > 7$.²⁰

The development of the medial cluster EPR signal as a function of potential (Fig. S9) also correlates with previous measurements of Native *E. coli* Hyd-1, suggesting that the midpoint potential of this cluster is unaffected by the E73Q amino acid exchange. (Unfortunately a quantitative analysis of

the redox potential of spectroscopic transitions at lower potential than the proximal cluster was not possible due to the low protein concentration). Given the large distance between E73 and the distal cluster,²¹ we predict that this FeS centre (invisible in EPR)¹⁹ will also have a redox potential which is the same in both the E73Q variant and Native enzyme. The changes in catalytic bias of E73Q therefore cannot be interpreted in light of changes to the energetics of the electron transfer relay.

Based on the evidence above, we speculate that the E73Q amino acid exchange impacts on the active site chemistry of Hyd-1. Fourier transform InfraRed spectroelectrochemical experiments, beyond the scope of this study, would be the logical suggestion for testing this hypothesis.

In conclusion, we have confirmed that it is possible to significantly perturb the *in vitro* catalytic bias of an O₂ tolerant NiFe MBH while maintaining the native levels of resistance to aerobic inhibition. We have engineered an apparent doubling of H₂ production rates through altering an amino acid residue which is located close to the proximal cluster. We have followed a reverse engineering strategy suggested by comparing *E. coli* Hyd-1 and Hyd-2, emphasising the utility of working with bacteria which express multiple hydrogenases.

The research leading to these results has received funding from the Wellcome Trust through the Combating Infectious Disease: Computational Approaches in Translational Science (CIDCATS) CDT program, from a Royal Society small research grant and from EP/M024393/1 (to MMR) and EP/M506394/1 (supporting JJW).

Notes and references

1. A. Parkin, in *The Metal-Driven Biogeochemistry of Gaseous Compounds in the Environment*, eds. P. M. H. Kroneck and M. E. S. Torres, Springer Netherlands, 2014, vol. 14, p. 99.
2. W. Lubitz, H. Ogata, O. Rüdiger and E. Reijerse, *Chem. Rev.*, 2014, **114**, 4081.
3. A. Parkin and F. Sargent, *Current Opinion in Chemical Biology*, 2012, **16**, 26.
4. Y. Shomura, K.-S. Yoon, H. Nishihara and Y. Higuchi, *Nature*, 2011, **479**, 253.
5. M. J. Lukey, M. M. Roessler, A. Parkin, R. M. Evans, R. A. Davies, O. Lenz, B. Friedrich, F. Sargent and F. A. Armstrong, *Journal of the American Chemical Society*, 2011, **133**, 16881.
6. T. Goris, A. F. Wait, M. Saggu, J. Fritsch, N. Heidary, M. Stein, I. Zebger, F. Lenzian, F. A. Armstrong, B. Friedrich and O. Lenz, *Nat Chem Biol*, 2011, **7**, 310.
7. R. M. Evans, A. Parkin, M. M. Roessler, B. J. Murphy, H. Adamson, M. J. Lukey, F. Sargent, A. Volbeda, J. C. Fontecilla-Camps and F. A. Armstrong, *J. Am. Chem. Soc.*, 2013, **135**, 2694.
8. S. V. Hexter, T. F. Esterle and F. A. Armstrong, *Phys. Chem. Chem. Phys.*, 2014, **16**, 11822.
9. S. V. Hexter, F. Grey, T. Happe, V. Climent and F. A. Armstrong, *PNAS*, 2012, **109**, 11516.
10. B. J. Murphy, F. Sargent and F. A. Armstrong, *Energ. Environ. Sci.*, 2014, **7**, 1426.
11. T. Goris, A. F. Wait, M. Saggu, J. Fritsch, N. Heidary, M. Stein, I. Zebger, F. Lenzian, F. A. Armstrong, B. Friedrich and O. Lenz, *Nat Chem Biol*, 2011, **7**, 310-318.
12. P. Raleiras, N. Khanna, H. Miranda, L. S. Meszaros, H. Krassen, F. Ho, N. Battchikova, E.-M. Aro, A. Magnuson, P. Lindblad and S. Styring, *Energ. Environ. Sci.*, Advance article, DOI:10.1039/c1035ee02694f.
13. A. A. Hamdan, S. Dementin, P. P. Liebgott, O. Gutierrez-Sanz, P. Richaud, A. L. D. Lacey, M. Roussett, P. Bertrand, L. Cournac and C. Léger, *J. Am. Chem. Soc.*, 2012, **134**, 8368.
14. M. J. Lukey, A. Parkin, M. M. Roessler, B. J. Murphy, J. Harmer, T. Palmer, F. Sargent and F. A. Armstrong, *J. Biol. Chem.*, 2010, **285**, 3928.
15. L. Bowman, L. Flanagan, P. K. Fyfe, A. Parkin, W. N. Hunter and F. Sargent, *Biochem. J.*, 2014, **458**, 449.
16. R. Cammack, in *Hydrogen as a Fuel: Learning from Nature*, eds. R. Cammack, M. Frey and R. Robson, CRC Press, 2001.
17. I. Dance, *Chemical Science*, 2015, **6**, 1433-1443.
18. J. J. Wright, E. Salvadori, H. Bridges, J. Hirst and M. M. Roessler, Invited Paper, Under review in *Journal of Inorganic Biochemistry*.
19. M. M. Roessler, R. M. Evans, R. A. Davies, J. Harmer and F. A. Armstrong, *J. Am. Chem. Soc.*, 2012, **134**, 15581.
20. M.-E. Pandelia, W. Nitschke, P. Infossi, M.-T. Giudici-Orticoni, E. Bill and W. Lubitz, *Proceedings of the National Academy of Sciences*, 2011, **108**, 6097-6102.
21. A. Volbeda, P. Amara, C. Darnault, J.-M. Mouesca, A. Parkin, M. M. Roessler, F. A. Armstrong and J. C. Fontecilla-Camps, *Proceedings of the National Academy of Sciences*, 2012, **109**, 5305-5310.