



UNIVERSITY OF LEEDS

This is a repository copy of *An extremal eigenvalue problem in Kähler geometry*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/98566/>

Version: Accepted Version

Article:

Apostolov, V, Jakobson, D and Kokarev, G (2015) An extremal eigenvalue problem in Kähler geometry. *Journal of Geometry and Physics*, 91. pp. 108-116. ISSN 0393-0440

<https://doi.org/10.1016/j.geomphys.2015.01.008>

© 2015, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

AN EXTREMAL EIGENVALUE PROBLEM IN KÄHLER GEOMETRY

VESTISLAV APOSTOLOV, DMITRY JAKOBSON, AND GERASIM KOKAREV

Dedicated to Professor Paul Gauduchon on the occasion of his 70'th birthday.

ABSTRACT. We study Laplace eigenvalues λ_k on Kähler manifolds as functionals on the space of Kähler metrics with cohomologous Kähler forms. We introduce a natural notion of a λ_k -extremal Kähler metric and obtain necessary and sufficient conditions for it. A particular attention is paid to the λ_1 -extremal properties of Kähler-Einstein metrics of positive scalar curvature on manifolds with non-trivial holomorphic vector fields.

1. INTRODUCTION

1.1. **Motivation.** Let M be a closed manifold. For a Riemannian metric g on M we denote by

$$0 = \lambda_0(g) < \lambda_1(g) \leq \lambda_2(g) \leq \dots \leq \lambda_k(g) \leq \dots$$

the eigenvalues of the Laplace-Beltrami operator Δ_g repeated according to their multiplicity. In real dimension 2, by classical results of Hersch [23], Yang and Yau [43], and Li and Yau [32], the first eigenvalue $\lambda_1(g)$ is bounded when the Riemannian metric g ranges over metrics of fixed volume. A basic question is: for a given conformal class c on M , is there a metric that maximizes $\lambda_1(g)$ among metrics $g \in c$ with $\text{vol}(M, g) = 1$? What are its properties? When M is a sphere or a projective plane, the answers go back to the classical results of Hersch [23] and Li and Yau [32]. For higher genus surfaces this circle of questions have been studied extensively in the last decades, see [14, 15, 16, 24, 25, 34] and the most recent papers [31, 35, 36].

In particular, Nadirashvili and Sire [35], developing earlier ideas by Nadirashvili [34], have stated an existence theorem for λ_1 -maximizers in conformal classes along with an outlined proof. This statement has been improved by Petrides [36], who has also given a rigorous argument for it, using previous work [19, 31]. Mention that the λ_1 -maximizers given by the above existence theorems may have conical singularities, and can be described as metrics that admit harmonic maps into round spheres by their first eigenfunctions. The latter statement actually holds for arbitrary λ_1 -maximizers (and even λ_k -extremals for any $k \geq 1$) with conical singularities, see [31], where many more general statements in this direction have been proven. Besides, the combination of [36, Theorem 1] and [31, Theorem E_1] shows that the set of all conformal C^∞ -metrics with conical singularities that maximize λ_1 is compact, see also [30].

1.2. **Eigenvalue problems on Kähler manifolds.** The purpose of this paper is to describe an extremal eigenvalue problem in higher dimensions on a Kähler manifold (M, J, g, ω) , which generalizes the above extremal problem on Riemannian surfaces. Here we view the eigenvalue $\lambda_k(g)$ as a functional on the space $\mathcal{K}_\Omega(M, J)$ that is formed by Kähler metrics g whose Kähler forms ω represent a given de Rham cohomology class Ω . When (M, J, g) is a Riemannian surface with a volume form ω , the space $\mathcal{K}_{[\omega]}(M, J)$ is precisely the set of metrics of a fixed area

2010 *Mathematics Subject Classification.* 49R05, 35P99, 53C55, 32Q20.

Key words and phrases. Laplacian eigenvalues, extremal metric, Kähler manifold, Kähler-Einstein manifold.

that are conformal to g , see the discussion in Sect. 2. By classical work due to Bourguignon, Li, and Yau [9], the first eigenvalue $\lambda_1(g)$ is bounded on $\mathcal{K}_\Omega(M, J)$ when (M, J) is projective and the de Rham class Ω belongs to $H^2(M, \mathbb{Q})$. The eigenvalue bound in [9] shows that the Fubini-Study metric on $\mathbb{C}P^m$ is a λ_1 -maximizer in its Kähler class. Recently, these results have been generalized by Arezzo, Ghigi, and Loi [6] to more general Kähler manifolds that admit holomorphic stable vector bundles over M with sufficiently many sections. In particular, they show that the symmetric Kähler-Einstein metrics on the Grassmannian spaces are also λ_1 -maximizers in their Kähler classes. Moreover, as is shown in [8], so are symmetric Kähler-Einstein metrics on Hermitian symmetric spaces of compact type. These results motivate a further investigation of spectral geometry of Kähler-Einstein metrics of positive scalar curvature; see [10, 41] for the general existence theory of such metrics.

Following the ideas in [34, 17] for other extremal eigenvalue problems, we introduce the notion of a λ_k -extremal Kähler metric under the deformations in its Kähler class. The class of such extremal metrics contains λ_k -maximizers in $\mathcal{K}_\Omega(M, J)$. We assume that the metrics under consideration are always smooth, and show that for a λ_k -extremal Kähler metric there exists a collection of non-trivial λ_k -eigenfunctions f_1, \dots, f_ℓ such that

$$\lambda_k^2 \left(\sum_{i=1}^{\ell} f_i^2 \right) - 2\lambda_k \left(\sum_{i=1}^{\ell} |\nabla f_i|^2 \right) + \sum_{i=1}^{\ell} |dd^c f_i|^2 = 0.$$

For the first eigenvalue the latter hypothesis is also sufficient for a metric to be λ_1 -extremal. This statement shows that for a λ_k -extremal Kähler metric g the eigenvalue $\lambda_k(g)$ is always multiple (Corollary 2.2), and allows to produce examples of λ_1 -extremal metrics as products.

We proceed with considering in more detail the case when (M, J) is a complex Fano manifold with non-trivial holomorphic vector fields. Recall that, by a classical result of Matsushima [33], for a Kähler-Einstein metric g on (M, J) the λ_1 -eigenfunctions are potentials of Killing vector fields. Using this fact, we show that a Kähler-Einstein metric g is λ_1 -extremal if and only if there exist non-trivial Killing potentials f_1, \dots, f_ℓ such that the function $\sum f_i^2$ is also a Killing potential. As an application, we conclude that a toric Fano Kähler-Einstein manifold whose connected group of automorphisms is a torus is not λ_1 -extremal. For example, so is a Kähler-Einstein metric on $\mathbb{C}P^2$ blown up at three points in a generic position, see [39, 40]. As another example, we show that Kähler-Einstein manifolds different from $\mathbb{C}P^m$ that admit hamiltonian 2-forms of order ≥ 1 are also never λ_1 -extremal. This conclusion applies to the non-homogeneous Kähler-Einstein metrics from [27, 28, 29].

2. AN EXTREMAL EIGENVALUE PROBLEM

2.1. Statement of the problem. Let (M, g, J, ω) be a compact Kähler manifold of real dimension $n = 2m$. Recall that due to the $\partial\bar{\partial}$ -lemma any Kähler metric \tilde{g} whose Kähler form $\tilde{\omega}(\cdot, \cdot) = \tilde{g}(J\cdot, \cdot)$ is co-homologous to ω has the form $\omega + dd^c\varphi$, where $d^c = JdJ^{-1} = i(\partial - \bar{\partial})$ and the action of J on the cotangent bundle is defined via the duality with respect to g . The smooth function φ above is determined uniquely by the condition

$$\int_M \varphi v_g = 0,$$

where $v_g = \omega^m/m!$ is the Riemannian volume form on M . By $\mathcal{K}_\Omega(M, J)$ we denote the space of Kähler metrics on (M, J) whose Kähler forms represent a given de Rham cohomology class Ω .

For a representative Kähler form ω it can be identified with the space of functions

$$\{\varphi \in C^\infty(M) : \omega + dd^c \varphi > 0, \int_M \varphi v_g = 0\}.$$

Besides, for any function φ with a zero mean-value there exists $\varepsilon > 0$ such that $\omega + tdd^c \varphi \in \mathcal{K}_\Omega(M, J)$ for all $|t| < \varepsilon$, and the space $C_0^\infty(M)$, formed by such functions φ , can be thought as the tangent space at $g \in \mathcal{K}_\Omega(M, J)$.

For a Kähler metric g on (M, J) we denote by

$$0 = \lambda_0(g) < \lambda_1(g) \leq \lambda_2(g) \leq \dots \leq \lambda_k(g) \leq \dots$$

the eigenvalues of the Laplace–Beltrami operator $\Delta_g = \delta d$ acting on functions, where δ is the L_2 -adjoint of the exterior derivative d with respect to g . Recall that due to the standard Kato–Rellich perturbation theory the functions $t \mapsto \lambda_k(g_t)$ have left and right derivatives for analytic deformations g_t . We view the eigenvalues $\lambda_k(g)$ as functionals on the space $\mathcal{K}_\Omega(M, J)$ and introduce the following definition.

Definition 2.1. A Kähler metric $g \in \mathcal{K}_\Omega(M, J)$ is called λ_k -extremal, if for any analytical deformation $g_t \in \mathcal{K}_\Omega(M, J)$ with $g_0 = g$ the following relation holds

$$(2.1) \quad \left. \frac{d}{dt} \right|_{t=0-} \lambda_k(g_t) \cdot \left. \frac{d}{dt} \right|_{t=0+} \lambda_k(g_t) \leq 0.$$

It is straightforward to see that a Kähler metric is λ_k -extremal if and only if either the inequality

$$\lambda_k(g_t) \leq \lambda_k(g) + o(t) \quad \text{as } t \rightarrow 0,$$

or the inequality

$$\lambda_k(g_t) \geq \lambda_k(g) + o(t) \quad \text{as } t \rightarrow 0$$

occurs. In particular, we see that any λ_k -maximizer in $\mathcal{K}_\Omega(M, J)$ is λ_k -extremal. The following two remarks are consequences of the results in [17]. First, for the first eigenvalue only the first of the above inequalities may occur. Second, for a deformation $\omega_t = \omega + dd^c \varphi_t$ the validity or the failure of relation (2.1) depends only on the function $\dot{\varphi} = (d/dt)|_{t=0} \varphi_t$, and hence, in Definition 2.1 we may consider only deformations with $\varphi_t = t\varphi$, where $\varphi \in C_0^\infty(M)$.

In the sequel we use the fourth order differential operator $L(f)$ defined as $\delta^c \delta(fdd^c f)$, where δ and δ^c stand for the L_2 -adjoints of d and d^c respectively. Recall that they satisfy the relations

$$\delta \psi = - \sum_{i=1}^{2m} \iota_{e_i} (D_{e_i} \psi) \quad \text{and} \quad \delta^c \psi = - \sum_{i=1}^{2m} \iota_{J e_i} (D_{e_i} \psi),$$

where D is the Levi-Civita connection of g , and $\{e_1, J e_1, \dots, e_m, J e_m\}$ is a J -adapted orthonormal frame. We then calculate

$$\begin{aligned} (2.2) \quad L(f) &= \delta^c \left(- (dd^c f)(df^\sharp, \cdot) + f d^c \delta df \right) \\ &= \sum_{i=1}^{2m} \iota_{J e_i} (D_{e_i} dd^c f)(df^\sharp, \cdot) - \sum_{i=1}^{2m} (\iota_{J e_i} (dd^c f)(D_{e_i} df^\sharp), \cdot) + f \delta^c d^c \delta df - (df, d\Delta f) \\ &= (\delta^c dd^c f, df) + (dd^c f, dd^c f) + f \Delta_g^2 f - (\Delta df, df) \\ &= (dd^c f, dd^c f) + f \Delta_g^2 f - 2(\Delta df, df), \end{aligned}$$

where we used the facts that $dd^c f$ is J -invariant and that J commutes with D and Δ_g (when acting on 1-forms), as well as the standard Kähler identities between the operators d , d^c , δ , δ^c , and Δ_g .

The proof of the following statement is close in the spirit to the arguments in [34, 17] and is given at the end of the section.

Theorem 2.1. *Let $g \in \mathcal{K}_\Omega(M, J)$ be a λ_k -extremal Kähler metric. Then there exists a finite collection f_1, \dots, f_ℓ of non-trivial eigenfunctions corresponding to $\lambda_k(g)$ such that*

$$(2.3) \quad \sum_{i=1}^{\ell} L(f_i) = \lambda_k^2 \left(\sum_{i=1}^{\ell} f_i^2 \right) - 2\lambda_k \left(\sum_{i=1}^{\ell} |\nabla f_i|^2 \right) + \sum_{i=1}^{\ell} |dd^c f_i|^2 = 0.$$

For $k = 1$ the existence of such a collection of eigenfunctions is sufficient for a Kähler metric g to be λ_1 -extremal.

Denote by $E_k = E_k(g)$ the eigenspace formed by eigenfunctions corresponding to the eigenvalue $\lambda_k(g)$. The operator L restricted to E_k takes the form

$$L(f) = \lambda_k^2 f^2 - 2\lambda_k |\nabla f|^2 + |dd^c f|^2.$$

Considering maximal and minimal values of a function $f \in E_k$, it is straightforward to see that it has a trivial kernel. Thus, we obtain the following corollary.

Corollary 2.2. *Let $g \in \mathcal{K}_\Omega(M, J)$ be a λ_k -extremal Kähler metric. Then the eigenvalue $\lambda_k(g)$ is multiple.*

As another consequence of Theorem 2.1, we see that the notion of λ_1 -extremality behaves well under products.

Corollary 2.3. *Let $g \in \mathcal{K}_\Omega(M)$ be a λ_1 -extremal Kähler metric on (M, J) . Then for any Kähler metric g' on (M', J') such that $\lambda_1(g') \geq \lambda_1(g)$ the product metric $g \times g'$ is λ_1 -extremal along deformations in its Kähler class on $(M, J) \times (M', J')$.*

Mention that the hypothesis $\lambda_1(g') \geq \lambda_1(g)$ in the corollary above always holds after an appropriate scaling of either of the metrics.

2.2. Discussion and basic examples. We proceed with considering the case when $m = 1$, that is when M is an oriented Riemann surface. Let g be a Riemannian metric and $\omega = v_g$ be its volume form. It is straightforward to see that for any smooth function $\varphi \in C^\infty(M)$ the hypothesis $\omega + dd^c \varphi > 0$ holds if and only if $1 - \Delta_g \varphi > 0$. Thus, the space $\mathcal{K}_{[\omega]}(M, J)$ can be defined as

$$\mathcal{K}_{[\omega]}(M, J) = \left\{ \varphi \in C^\infty(M) : 1 - \Delta_g \varphi > 0, \int_M \varphi v_g = 0 \right\}.$$

The following lemma is elementary; we state it for the convenience of references.

Lemma 2.4. *Let (M, g) be a Riemannian surface, and ω be its volume form. Then the space of Kähler metrics in $\mathcal{K}_{[\omega]}(M, J)$ coincides with the space of Riemannian metrics \tilde{g} that are conformal to g and such that $\text{vol}(M, \tilde{g}) = \text{vol}(M, g)$.*

Proof. Let φ be a function from the above space $\mathcal{K}_{[\omega]}(M, J)$. Then the metric $\tilde{g} = (1 - \Delta_g \varphi)g$ is clearly has the same volume as the metric g . Conversely, for a given conformal metric $\tilde{g} = e^\sigma g$ of the same volume as g the equation

$$e^\sigma = 1 - \Delta_g \varphi$$

has a unique solution $\varphi \in C^\infty(M)$ with zero mean-value, see [21] for standard existence results for solutions of elliptic equations. \square

As a consequence of the lemma above, we see that a Riemannian metric g on M is λ_k -extremal in the sense of Definition 2.1 if and only if it is λ_k -extremal under volume preserving conformal deformations. There have been constructed a number of examples of various λ_1 -extremal and λ_1 -maximal metrics in the literature, and as is known [35, 36], every conformal class on a closed surface contains a λ_1 -maximizer, which may have conical singularities. Thus, using Corollary 2.3, we obtain a variety of examples of λ_1 -extremal Kähler metrics by taking the products of λ_1 -extremal Riemannian surfaces and Kähler manifolds.

Recall that by [15, 17] for a metric g that is λ_k -extremal under the volume preserving conformal deformations there exists a collection f_1, \dots, f_ℓ of non-trivial λ_k -eigenfunctions such that $\sum f_i^2 = 1$. When $m = 1$, we see that the latter condition coincides with the necessary condition given by Theorem 2.1. Indeed, in this case, for a λ_k -eigenfunction f the operator $L(f)$ takes the form

$$L(f) = 2(\lambda_k^2 f^2 - \lambda_k |\nabla f|^2),$$

where we used the identities $(\Delta_g f)\omega = -dd^c f$ and $|\omega|^2 = 1$. Now the relation

$$\Delta_g(\sum f_i^2) = 2(\lambda_k \sum f_i^2 - \sum |\nabla f_i|^2) = \lambda_k^{-1} \sum L(f_i)$$

implies the claim. More generally, in higher dimensions $m \geq 2$ the hypothesis $\sum L(f_i) = 0$ in Theorem 2.1 is equivalent to the relation

$$\Delta_g(\sum f_i^2) = \lambda_k^{-1} m(m-1) (\sum dd^c f_i \wedge dd^c f_i) \wedge \omega^{m-2} / \omega^m.$$

Here we used the fact that $(\Delta_g f)^2 - |dd^c f|^2 = m(m-1) dd^c f \wedge dd^c f \wedge \omega^{m-2} / \omega^m$, see identity (2.8) in the proof of Theorem 2.1 below.

2.3. Proof of Theorem 2.1. Below we use the conventions and basic identities from [7, Ch. 2]; in particular, $|\omega|^2 = m$ and $m \cdot dd^c f \wedge \omega^{m-1} = (-\Delta_g f)\omega^m$. We start with the following lemma.

Lemma 2.5. *Let $g \in \mathcal{H}_\Omega(M, J)$ be a λ_k -extremal Kähler metric, and E_k be an eigenspace for $\lambda_k(g)$. Then for any function $\varphi \in C_0^\infty(M)$ the quadratic form*

$$Q_\varphi(f) = \int_M \varphi L(f) v_g$$

is indefinite on E_k . For $k = 1$ the hypothesis that the form Q_φ is indefinite for any $\varphi \in C_0^\infty(M)$ is also sufficient for a Kähler metric g to be λ_k -extremal.

Proof. For any $\varphi \in C_0^\infty(M)$ consider the Kähler deformation $\omega_t = \omega + t dd^c \varphi$, defined for a sufficiently small $|t|$. By [17, Theorem 2.1] for a proof of the lemma it is sufficient to show that the form Q_φ satisfies the relation

$$(2.4) \quad Q_\varphi(f) = \int_M f(\dot{\Delta}_\varphi f) v_g,$$

where $\dot{\Delta}_\varphi f$ stands for the value $(d/dt)|_{t=0}(\Delta_{g_t} f)$. First, we claim that the operator $\dot{\Delta}_\varphi$ satisfies the identity

$$(2.5) \quad \dot{\Delta}_\varphi f = (dd^c f, dd^c \varphi).$$

Indeed, differentiating the relation

$$m \cdot dd^c f \wedge \omega_t^{m-1} = (-\Delta_{g_t} f)\omega_t^m,$$

we obtain

$$(2.6) \quad dd^c f \wedge dd^c \varphi \wedge \omega^{m-2} / (m-2)! = \left((\Delta_g f)(\Delta_g \varphi) - \dot{\Delta}_\varphi f \right) \omega^m / m!.$$

Denote by $\wedge^{1,1}(M)$ the bundle of real $(1,1)$ -forms on the manifold (M, J) , that is 2-forms ψ such that $\psi(J\cdot, J\cdot) = \psi(\cdot, \cdot)$. For a given Kähler metric (g, ω) on (M, J) , it decomposes as a direct g -orthogonal sum

$$(2.7) \quad \wedge^{1,1}(M) = \mathbb{R}\omega \oplus \wedge_0^{1,1}(M)$$

of the irreducible $U(m)$ -invariant subspaces of $(1,1)$ -forms proportional to ω and primitive (trace-free) $(1,1)$ -forms, respectively. Let us consider the symmetric $U(m)$ -invariant bilinear form on $\wedge^{1,1}(M)$ defined as

$$q(\phi, \psi) = (\mathrm{tr}_\omega \phi)(\mathrm{tr}_\omega \psi) - \frac{\phi \wedge \psi \wedge (\omega^{m-2}/(m-2)!)}{\omega^m/m!},$$

where

$$\mathrm{tr}_\omega \psi = (\psi, \omega)_g = \frac{\psi \wedge (\omega^{m-1}/(m-1)!)}{\omega^m/m!}.$$

Since the form $q(\cdot, \cdot)$ leaves the two irreducible factors in the decomposition (2.7) orthogonal, by the Schür lemma we conclude that it is proportional to the induced Euclidean product $(\cdot, \cdot)_g$ on each factor in (2.7). Evaluating $q(\omega, \omega)$ and $q(\psi_0, \psi_0)$ with $\psi = \alpha \wedge J\alpha$ for a unitary 1-form α , we see that in fact the form $q(\cdot, \cdot)$ coincides with $(\cdot, \cdot)_g$, that is

$$(2.8) \quad (\mathrm{tr}_\omega \phi)(\mathrm{tr}_\omega \psi) - \frac{\phi \wedge \psi \wedge (\omega^{m-2}/(m-2)!)}{\omega^m/m!} = (\phi, \psi)_g.$$

Now combining the last relation with (2.6), we obtain identity (2.5). Using the latter we have

$$Q_\varphi(f) = \int_M \varphi L(f) v_g = \int_M f(dd^c f, dd^c \varphi) v_g = \int_M f(\Delta_\varphi f) v_g,$$

and thus, obtain relation (2.4). \square

By Lemma 2.5, in order to prove the theorem it is sufficient to show that the quadratic form $Q_\varphi(f)$ is indefinite on E_k if and only if there exists a collection of eigenfunctions $f_1, \dots, f_\ell \in E_k$ such that

$$(2.9) \quad \sum_{i=1}^{\ell} L(f_i) = 0 \quad \text{and} \quad \sum_{i=1}^{\ell} \int_M f_i^2 v_g = 1.$$

Consider the convex subset

$$K = \left\{ \sum_i L(f_i) : f_i \in E_k, \sum_i \int_M f_i^2 v_g = 1 \right\}$$

in the space $L^2(M)$. We are going to show that the form $Q_\varphi(f)$ is indefinite if and only if $0 \in K$. Suppose that $Q_\varphi(f)$ is indefinite for any $\varphi \in C_0^\infty(M)$ and $0 \notin K$. Then by the Hahn–Banach separation theorem there exists a function $\psi \in L^2(M)$ and $\varepsilon > 0$ such that

$$\int_M \psi uv_g \geq \varepsilon > 0 \quad \text{for any } u \in K.$$

Since the set K lies in a finite-dimensional subspace, then choosing $\varepsilon > 0$ smaller, if necessary, by approximation we may assume that the function ψ belongs to $C^\infty(M)$. Define ψ_0 as the zero mean-value part of ψ , that is

$$\psi_0 = \psi - \frac{1}{\mathrm{vol}(M, g)} \int_M \psi v_g.$$

Since the operator $L(f)$ takes values among zero mean-value functions, we obtain

$$Q_{\psi_0}(f) = \int_M \psi_0 L(f) v_g = \int_M \psi L(f) v_g > 0$$

for any non-trivial $f \in E_k$. Thus, we arrive at a contradiction with the assumption that the form Q_φ is indefinite for any $\varphi \in C_0^\infty(M)$.

Conversely, given a collection $f_1, \dots, f_\ell \in E_k$ that satisfy relationships (2.9), we have

$$\sum_{i=1}^{\ell} Q_\varphi(f_i) = \int_M \varphi \left(\sum_{i=1}^{\ell} L(f_i) \right) v_g = 0.$$

Thus, the quadratic form $Q_\varphi(f)$ is indeed indefinite for any $\varphi \in C_0^\infty(M)$. \square

3. KÄHLER–EINSTEIN MANIFOLDS WITH A NON-TRIVIAL AUTOMORPHISM GROUP

We now specialize the considerations to the case when (g, J, ω) is a Kähler–Einstein manifold with positive scalar curvature, that is the Ricci form ρ is a positive constant multiple of the Kähler form ω . Rescaling the metric, we may assume that $\rho = \omega$, or equivalently, the Kähler class is $\Omega = 2\pi c_1(M, J)$. Under this assumption, the scalar curvature Scal_g equals $2m$.

Since the manifold (M, J) is Fano, by [26, 44] it is simply-connected. In particular, the first de Rham cohomology group vanishes, and hence, any real holomorphic vector field X can be uniquely written in the form

$$X = \text{grad}_g h_X + J \text{grad}_g f_X,$$

where h_X and f_X are smooth functions with zero mean-values, see [7, Cor. 2.126] for details. The complex-valued function $h_X + if_X$ is called the *holomorphy potential* of X . For a Killing vector field Y , the above decomposition reduces to

$$Y = J \text{grad}_g f_Y,$$

and the corresponding function $f_Y \in C_0^\infty(M)$ is called the *Killing potential* of Y . The following statement is due to Matsushima [33], see also [20, Ch. 3].

Proposition 3.1. *Let (M, g, J, ω) be a compact Kähler–Einstein manifold with scalar curvature $\text{Scal}_g = 2m$. Then the Lie algebra $\mathfrak{h}(M)$ of real holomorphic vector fields on (M, J) decomposes as the direct sum*

$$\mathfrak{h}(M) = \mathfrak{k}(M, g) \oplus J\mathfrak{k}(M, g),$$

where $\mathfrak{k}(M, g)$ is the sub-algebra of Killing vector fields for g . Moreover, the algebra $\mathfrak{k}(M, g)$ is Lie algebra isomorphic to the space

$$E_1(g) = \{f \in C_0^\infty(M) : \Delta_g(f) = 2f\}$$

equipped with the Poisson bracket of functions with respect to ω , via the map $f \rightarrow J \text{grad}_g f$. Furthermore, the first eigenvalue satisfies the inequality $\lambda_1(g) \geq 2$.

As a consequence of the proposition above, we see that $\lambda_1(g) = 2$ if and only if the connected component of the identity $\text{Aut}_0(M, J)$ of the group of biholomorphic automorphisms of (M, J) is non-trivial. In this case, we have the following statement.

Theorem 3.2. *Let (M, g, J, ω) be a compact Kähler–Einstein manifold with scalar curvature $\text{Scal}_g = 2m$, and suppose that the connected component $\text{Aut}_0(M, J)$ of the automorphism group is non-trivial. Then the metric g is λ_1 -extremal in $\Omega = 2\pi c_1(M, J)$ if and only if there exist*

non-trivial λ_1 -eigenfunctions f_1, \dots, f_ℓ such that the zero mean-value part of the sum $\sum_{i=1}^\ell f_i^2$ is a (possibly trivial) λ_1 -eigenfunction, that is

$$(3.1) \quad \sum_{i=1}^\ell f_i^2 - \frac{1}{\text{vol}(M, g)} \int_M \left(\sum_{i=1}^\ell f_i^2 \right) \nu_g \in E_1(g).$$

Proof. We show that the necessary and sufficient condition in Theorem 2.1 is equivalent to relation (3.1). First, note that by Proposition 3.1, we have $\lambda_1(g) = 2$, and any eigenfunction $f \in E_1(g)$ is a Killing potential. In particular, the form $d^c f$ is dual to a Killing vector field, and therefore $Dd^c f = (1/2)dd^c f$. Thus, using the standard identity $D^*D|T|^2 = 2(D^*DT, T) - 2(DT, DT)$ for a tensor field T , we obtain

$$\begin{aligned} \Delta_g |df|^2 &= D^*D(d^c f, d^c f) = 2\left((D^*D(d^c f), d^c f) - (D(d^c f), D(d^c f)) \right) \\ &= (\delta d^c f, d^c f) - (dd^c f, dd^c f) = (\Delta_g d^c f, d^c f) - (dd^c f, dd^c f) \\ &= 2|df|^2 - |dd^c f|^2. \end{aligned}$$

Above we also used the relation $\Delta_g d^c f = 2df^c$ and the fact that the tensor norm of $dd^c f$ is twice its norm as a differential 2-form. On the other hand, we clearly have

$$(3.2) \quad \Delta_g f^2 = 2f\Delta_g f - 2|df|^2 = 4f^2 - 2|df|^2.$$

Combining the last two relations, for any eigenfunction $f \in E_1(g)$ we obtain

$$(3.3) \quad \Delta_g (f^2 - |df|^2) = 4f^2 - 4|df|^2 + |dd^c f|^2.$$

Now comparing (3.3) with the relation $\sum L(f_i) = 0$ in Theorem 2.1, we conclude that the metric g is λ_1 -extremal if and only if there exist $\{f_1, \dots, f_\ell\} \in E_1(g)$ such that

$$(3.4) \quad \sum_{i=1}^\ell (f_i^2 - |df_i|^2) = c$$

for some constant c . Integrating the last relation, we see that the constant c is minus the mean-value of the sum $\sum_{i=1}^\ell f_i^2$. Setting $f_0 := \sum_{i=1}^\ell f_i^2 + c$, and using (3.2), we further obtain

$$\begin{aligned} \Delta_g f_0 &= \sum_{i=1}^\ell (4f_i^2 - 2|df_i|^2) \\ (3.5) \quad &= 2f_0 + 2\left(\sum_{i=1}^\ell (f_i^2 - |df_i|^2) - c \right) = 2f_0. \end{aligned}$$

Thus, relation (3.4) is in turn equivalent to the hypothesis that there exist non-trivial $f_1, \dots, f_\ell \in E_1(g)$ such that the zero mean-value part f_0 of the sum $\sum_{i=1}^\ell f_i^2$ is a λ_1 -eigenfunction itself. \square

Theorem 3.2 is a useful criterion for verifying whether the Kähler–Einstein metric on (M, J) is λ_1 -extremal in $2\pi c_1(M, J)$. We demonstrate this in the corollaries below.

Corollary 3.3. *Let (M, g, J, ω) be a compact homogeneous Kähler–Einstein manifold. Then the metric g is a λ_1 -extremal metric within its Kähler class.*

Proof. Let $\{f_1, \dots, f_\ell\}$ be an orthonormal basis of $E_1(g)$ with respect to the L_2 -global product $\langle \cdot, \cdot \rangle_g$ induced by g . The group of Kähler isometries G of (M, g, J, ω) acts isometrically on the

space $(E_1(g), \langle \cdot, \cdot \rangle_g)$. It follows that the function

$$f = \sum_{i=1}^{\ell} f_i^2$$

is G -invariant, and since G acts transitively on M , is constant. \square

As another application, we consider *toric* Kähler–Einstein manifolds which have been studied in many places, see [1, 13, 22, 42], and for which the existence theory takes a fairly concrete shape.

Corollary 3.4. *Let (M, g, J, ω) be a compact Kähler–Einstein manifold of real dimension $2m$ whose connected identity component $\text{Aut}_0(M, J)$ of the automorphism group is the complexification of an m -dimensional real torus. Then the metric g is not λ_1 -extremal.*

Proof. The assumption on $\text{Aut}_0(M, J)$ implies that (M, g, J, ω) is a *toric* Kähler–Einstein metric in the sense of [1, 13, 22, 42]. Indeed, by Proposition 3.1, the connected component of the isometry group of the Kähler–Einstein metric is a maximal connected compact subgroup of $\text{Aut}_0(M, J)$. In our case, by assumption, it must be a real m -dimensional torus T . The latter acts in a hamiltonian way (as any induced vector field is individually hamiltonian because M is simply-connected, and T is abelian), and by Delzant theorem [12], the momentum map $\mu : M \rightarrow \mathfrak{t}^*$ sends M onto a compact convex polytope in the dual vector space \mathfrak{t}^* of the Lie algebra $\mathfrak{t} = \text{Lie}(T)$. By Proposition 3.1 the pullback $f = (u, \mu) + \lambda$ to M of an affine function $(u, x) + \lambda$ on \mathfrak{t}^* , where $u \in \mathfrak{t}$ and $\lambda \in \mathbb{R}$, defines an element in $E_1(g) \oplus \mathbb{R}$. Conversely, all elements of $E_1(g) \oplus \mathbb{R}$ are of this form, again by Proposition 3.1 and our assumption for $\text{Aut}_0(M, J)$.

Suppose that the metric g is λ_1 -extremal. Then, by Theorem 3.2, there exist non-trivial eigenfunctions $f_i = (u_i, \mu) + \lambda_i$ such that the sum $\sum_{i=1}^{\ell} f_i^2$ is the pull-back $f = (u, \mu) + \lambda$ of an affine function $(u, x) + \lambda$ on \mathfrak{t}^* . It follows that

$$\sum_{i=1}^{\ell} \left((u_i, x) + \lambda_i \right)^2 = (u, x) + \lambda,$$

which implies $u_i = 0$ for any i , and hence, $f_i = \lambda_i$ is constant for any i . Thus, we arrive at a contradiction with the hypothesis that the f_i 's are non-trivial. \square

The above corollary, for instance, shows that the Kähler–Einstein metric on $\mathbb{C}P^2$ blown-up at three points in general position (see [39, 40]) is not λ_1 -extremal, and in particular, can not be a maximizer for λ_1 in its Kähler class.

As a final example, we consider non-homogeneous Kähler–Einstein metrics on projective bundles over the product of compact Kähler–Einstein manifolds with positive scalar curvatures that have been found and studied by Koiso and Sakane [27, 28, 29, 38], see also [4, 11, 37] for alternative treatments. It is convenient to use the characterization from [4] saying that these are Kähler–Einstein metrics admitting a *Hamiltonian 2-form* of order 1, in the sense of the theory in [2, 3]. We can then prove the following statement.

Corollary 3.5. *Let (M, g, J, ω) be a compact Kähler–Einstein manifold that is different from $\mathbb{C}P^m$ and which admits a Hamiltonian 2-form of order ≥ 1 . Then the metric g is not λ_1 -extremal.*

Proof. As follows from the theory in [2, 3], a Hamiltonian 2-form ϕ on (M, g, J, ω) gives rise to an ℓ -dimensional (real) torus T in the connected component $I_0(M, g)$ of the isometry group of (g, J, ω) , where $\ell \geq 1$ is the order of ϕ . By the general classification [3, Thm. 5] and [2, Prop. 16], the corresponding moment map $\mu : M \rightarrow \mathfrak{t}^*$ sends M to a Delzant simplex in the dual

vector space of $\mathfrak{t} = \text{Lie}(T)$. Since M is simply connected by the Kähler–Einstein assumption (see [4, Sec. 2.1] for the refinement in this case) and $(M, J) \not\cong \mathbb{C}P^m$, we conclude that (M, J) is the total space of a holomorphic projective bundle $P(\mathcal{E}_0 \oplus \mathcal{E}_1 \oplus \cdots \oplus \mathcal{E}_\ell) \rightarrow S$ over the product $S = \prod_j S_j$ of Kähler–Einstein manifolds $(S_j, g_j, J_j, \omega_j)$, where \mathcal{E}_i are projectively-flat holomorphic vector bundles over S , satisfying certain topological conditions. Moreover, the metric g on M is obtained by the *generalized Calabi construction* associated to this bundle, see [5] for a detailed treatment of this class of metrics, but the existence of a Hamiltonian 2-form of order $\ell \geq 2$ is a more restrictive condition for the metric (and the bundles).

To describe the eigenspace E_1 corresponding to $\lambda_1(g)$, we first show that the torus T lies in the centre of $I_0(M, g)$. By the general theory in [2, 3], the torus T is generated by the Hamiltonian vector fields corresponding to the elementary symmetric functions of the m -eigenvalues of the Hamiltonian form ϕ viewed as a Hermitian operator via the Kähler form ω . Thus, it is sufficient to show that ϕ is invariant under any isometry $\Phi \in I_0(M, g)$. As shown in [18], when $(M, J) \not\cong \mathbb{C}P^m$ any other Hamiltonian 2-form $\tilde{\phi}$ on (M, g, J, ω) must be a linear combination of ϕ and ω . Since the property of being Hamiltonian is preserved by Kähler isometries, we see that $\tilde{\phi} = \Phi^*(\phi) = a\omega + b\phi$ for some constants a, b . In addition, we also have

$$\text{tr}_\omega(\phi) = \text{tr}_\omega \tilde{\phi} = am + b \text{tr}_\omega \phi.$$

As a Hamiltonian 2-form of order $\ell \geq 1$, ϕ cannot have a constant trace (otherwise it must be parallel by its very definition, and thus of order $\ell = 0$, see [2]), and we obtain that $a = 0$ and $b = 1$; in other words, $\Phi^*(\phi) = \phi$.

Denote by $\mathfrak{i}(M, g)$ the Lie algebra of $I_0(M, g)$. Since M is Fano, it is simply connected, and we may identify $\mathfrak{i}(M, g)$ with the space of zero mean-value Killing potentials. It follows from our previous argument that the centralizer of \mathfrak{t} in $\mathfrak{i}(M, g)$ equals $\mathfrak{i}(M, g)$, and we may use the description of $\mathfrak{i}(M, g)$ in [5, Lemma 5] in terms of the pullback metric on the T -equivariant blow-up

$$\hat{M} = P\left(\bigoplus_{k=0}^{\ell} \mathcal{O}(-1)_{\mathcal{E}_k}\right) \rightarrow \hat{S}$$

of (M, J) along the sub-manifolds $P(\mathcal{E}_i) \cong \mathbb{C}P^{r_i-1} \times S$, where r_i is the rank of \mathcal{E}_i and

$$\hat{S} = P(\mathcal{E}_0) \times_S \cdots \times_S P(\mathcal{E}_\ell) \cong \left(\prod_{i=0}^{\ell} \mathbb{C}P^{r_i-1}\right) \times \left(\prod_j S_j\right)$$

is equipped with the product of the Kähler–Einstein metrics on its factors. (Above we use the facts that S is simply connected and the bundles \mathcal{E}_i are projectively flat.) Note also that the induced action of T on \hat{M} arises from the diagonal action on each fibre of $\left(\bigoplus_{k=0}^{\ell} \mathcal{O}(-1)_{\mathcal{E}_k}\right) \rightarrow \hat{S}$, so that \hat{M} has a structure of a toric $\mathbb{C}P^\ell$ -bundle over \hat{S} , such that the pull-back of ω to \hat{M} restricts to each fibre to define a T -invariant symplectic form on $\mathbb{C}P^\ell$ whose momentum map is the pull-back of μ to \hat{M} .

Denote by $(S_\alpha, g_\alpha, J_\alpha, \omega_\alpha)$ a Kähler–Einstein factor in the definition of \hat{S} , where α ranges over the values of the i 's and j 's. As is shown in the proof of [5, Lemma 5], any Killing potential f on (M, g, J, ω) when pulled-back to \hat{M} has the form

$$f = \sum_{\alpha} \left((u_{\alpha}, \mu) + \lambda_{\alpha} \right) f_{\alpha} + \left((u, \mu) + \lambda \right),$$

where f_α is the pull-back to \hat{M} of a Killing potential on S_α , $u, u_\alpha \in \mathfrak{t}$, and $\lambda, \lambda_\alpha \in \mathbb{R}$. Here the values u_α, λ_α are determined by (M, J) . More precisely, the u_α 's can be expressed in terms of the degrees of \mathcal{E}_i over S_j while the λ_α 's are determined by the Kähler class of ω .

By Theorem 3.2, the hypothesis that the metric g is λ_1 -extremal reduces to the relation

$$\sum_{n=1}^r \left(\sum_{\alpha} \left((u_\alpha, \mu) + \lambda_\alpha \right) f_{\alpha,n} + (u_n, \mu) + \lambda_n \right)^2 = \sum_{\alpha} \left((u_\alpha, \mu) + \lambda_\alpha \right) f_\alpha + (u, \mu) + \lambda,$$

where $r \geq 2$, and the functions

$$f_n := \left(\sum_{\alpha} \left((u_\alpha, \mu) + \lambda_\alpha \right) f_{\alpha,n} + (u_n, \mu) + \lambda_n \right)$$

are non-constant. Restricting to a $\mathbb{C}P^\ell$ fibre over \hat{S} , the argument from the proof of Corollary 3.4 shows that the following relations hold:

$$(3.6) \quad \sum_{\alpha} u_\alpha f_{\alpha,n} + u_n = 0, \quad \sum_{\alpha} u_\alpha f_\alpha + u = 0,$$

where $n = 1, \dots, r$ is arbitrary. Here $f_{\alpha,n}$ and $f_{\beta,n}$ are the pullbacks of Killing potentials on S_α and S_β respectively, and hence, are functionally independent when $\alpha \neq \beta$. Thus, we see that $u_\alpha f_{\alpha,n}$ and $u_\alpha f_\alpha$ are constant vectors in the Lie algebra \mathfrak{t} for any α . By the classification in [3, Thm. 5], we get that $u_\alpha \neq 0$; relation (40) in [3] cannot vanish for each j . Hence, the potentials $f_{\alpha,n}$ and f_α are constant, and by the relations in (3.6), so are the functions

$$f_n = \left(\sum_{\alpha} \left((u_\alpha, \mu) + \lambda_\alpha \right) f_{\alpha,n} + (u_n, \mu) + \lambda_n \right) = \sum_{\alpha} \left(\lambda_\alpha f_{\alpha,n} \right) + \lambda_n.$$

Thus, we arrive at a contradiction. □

Acknowledgements. The project has originated out of a number of discussions the authors had while GK was visiting the Centre de recherches mathématiques in Montréal. Its hospitality is gratefully acknowledged. During the work on the paper VA and DJ were supported by NSERC Discovery Grants; DJ was also partially supported by FQRNT. The authors are grateful to the referee for carefully reading the manuscript and suggesting a number of improvements.

REFERENCES

- [1] M. Abreu, *Kähler geometry of toric varieties and extremal metrics*, Internat. J. Math **9** (1998), 641–651.
- [2] V. Apostolov, D. M. J. Calderbank and P. Gauduchon, *Hamiltonian 2-forms in Kähler Geometry I. General Theory*, J. Differential Geom. **73** (2006), 359–412.
- [3] V. Apostolov, D. M. J. Calderbank, P. Gauduchon and C. Tønnesen-Friedman, *Hamiltonian 2-forms in Kähler Geometry II. Global Classification*, J. Differential Geom. **68** (2004), 277–345.
- [4] V. Apostolov, D. M. J. Calderbank, P. Gauduchon and C. Tønnesen-Friedman, *Hamiltonian 2-forms in Kähler geometry. IV. Weakly Bochner-flat Kähler manifolds*, Comm. Anal. Geom. **16** (2008), 91–126.
- [5] V. Apostolov, D. M. J. Calderbank, P. Gauduchon and C. Tønnesen-Friedman, *Extremal Kähler metrics on projective bundles over a curve*, Adv. Math. **227** (2011), 2385–2424.
- [6] C. Arezzo, A. Ghigi, and A. Loi, *Stable bundles and the first eigenvalue of the Laplacian*, J. Geom. Anal. **17** (2007), 375–386.
- [7] A. Besse, *Einstein Manifolds*. Reprint of the 1987 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2008. xii+516 pp.
- [8] L. Biliotti, A. Ghigi, *Satake-Furstenberg compactifications, the moment map and λ_1* , Amer. J. Math. **135** (2013), 237–274.
- [9] J.-P. Bourguignon, P. Li, S.-T. Yau, *Upper bound for the first eigenvalue of algebraic sub manifolds*, Comment. Math. Helv. **69** (1994), 199–207.
- [10] X. X. Chen, S. K. Donaldson, and S. Sun, *Kähler-Einstein metrics and stability*, arXiv:1210.7494.
- [11] A. Dancer, M. Wang, *Kähler-Einstein metrics of cohomogeneity one*, Math. Ann. **312** (1998), 503–526.

- [12] T. Delzant, *Hamiltoniens périodiques et image convexe de l'application moment*, Bull. Soc. Math. France **116** (1988), 315–339.
- [13] S. K. Donaldson, *Scalar curvature and stability of toric varieties*, J. Differential Geom. **62** (2002), 289–349.
- [14] El Soufi, A., Ilias, S. *Riemannian manifolds admitting isometric immersions by their first eigenfunctions*. Pacific J. Math. **195** (2000), 91–99.
- [15] El Soufi, A., Ilias, S. *Extremal metrics for the first eigenvalue of the Laplacian in a conformal class*. Proc. Amer. Math. Soc. **131** (2003), 1611–1618.
- [16] El Soufi, A., Giacomini, H., Jazar, M. *A unique extremal metric for the least eigenvalue of the Laplacian on the Klein bottle*. Duke Math. J. **135** (2006), 181–202.
- [17] A. El Soufi and S. Ilias, *Laplacian eigenvalue functionals and metric deformations on compact manifolds*, J. Geom. Physics **58** (2008), 89–104.
- [18] A. Fedorova, V. Kiosak, V. Matveev, S. Rosemann, *The only Kähler manifold with degree of mobility at least 3 is $(\mathbb{C}P(n), g_{\text{Fubini-Study}})$* , Proc. Lond. Math. Soc. (3) **105** (2012), 153–188.
- [19] Fraser, A., Schoen, R. *Sharp eigenvalue bounds and minimal surfaces in the ball*. arXiv: 1209.3789.
- [20] P. Gauduchon, *Calabi's extremal metrics: An elementary introduction*, in preparation.
- [21] Gilbarg, D., Trudinger, N. S. *Elliptic partial differential equations of second order*. Classics in Mathematics. Springer-Verlag, Berlin, 2001. xiv+517 pp.
- [22] V. Guillemin, *Kähler structures on toric varieties*, J. Differential Geom. **40** (1994), 285–309.
- [23] J. Hersch, *Quatre propriétés isopérimétriques de membranes sphériques homogènes*, C.R. Acad. Sci. Paris Sér. A-B **270** (1970), A1645–A1648.
- [24] Jakobson, D., Levitin, M., Nadirashvili, N., Nigam, N., Polterovich, I. *How large can the first eigenvalue be on a surface of genus two?* Int. Math. Res. Not. (2005), 3967–3985.
- [25] Jakobson, D., Nadirashvili, N., Polterovich, I. *Extremal metric for the first eigenvalue on the Klein bottle*. Canad. J. Math. **58** (2006), 381–400.
- [26] S. Kobayashi, *On Kähler manifolds with positive definite Ricci tensor*, Ann. of Math. **74** (1961), 570–574.
- [27] N. Koiso, *On rationally symmetric Hamilton's equations for Kähler–Einstein metrics*, in ‘Recent topics in differential and analytic geometry’, Adv. Studies in Pure Math. **18-I**, Academic Press, Boston, 1990, 327–337.
- [28] N. Koiso and Y. Sakane, *Nonhomogeneous Kähler–Einstein metrics on compact complex manifolds*, in Curvature and Topology of Riemannian Manifolds (Kataka, 1985), Lecture Notes in Math. 1201, Springer-Verlag, Berlin, 1986, 165–179.
- [29] N. Koiso and Y. Sakane, *Nonhomogeneous Kähler–Einstein metrics on compact complex manifolds II*, Osaka J. Math. **25** (1988), 933–959.
- [30] Kokarev, G. *On the concentration-compactness phenomenon for the first Schrodinger eigenvalue*. Calc. Var. Partial Differential Equations **38** (2010), 29–43.
- [31] Kokarev, G. *Variational aspects of Laplace eigenvalues on Riemannian surfaces*. Adv. Math. **258** (2014), 191–239.
- [32] P. Li, Y.-S. Yau, *A new conformal invariant and its application to the Willmore conjecture and the first eigenvalue of compact surfaces*, Invent. Math. **69** (1982), 269–291.
- [33] Y. Matsushima, *Sur la structure du groupe d'homéomorphismes analytiques d'une certaine variété kählérienne*, Nagoya Math. J. **11** (1957), 145–150.
- [34] Nadirashvili, N. *Berger's isoperimetric problem and minimal immersions of surfaces*. Geom. Funct. Anal. **6** (1996), 877–897.
- [35] Nadirashvili, N., Sire, Y. *Conformal spectrum and harmonic maps*. arXiv:1007.3104.
- [36] Petrides, R. *Existence and regularity of maximal metrics for the first Laplace eigenvalue on surfaces*. Geom. Funct. Anal., to appear.
- [37] F. Podestà and A. Spiro, *Kähler manifolds with large isometry group*, Osaka J. Math. **36** (1999), 805–833.
- [38] Y. Sakane, *Examples of compact Einstein Kähler manifolds with positive Ricci tensor*. Osaka J. Math. **23** (1986), 585–616.
- [39] Y.-T. Siu, *The Existence of Kähler–Einstein Metrics on Manifolds with Positive Anticanonical Line Bundle and a Suitable Finite Symmetry Group*, Ann. Math. **127** (1988), 585–627.
- [40] G. Tian, *On Kähler–Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$* . Invent. Math. **89** (1987), 225–246.
- [41] G. Tian, *K-stability and Kähler–Einstein metrics*, arXiv:1211.4669.
- [42] X.-J. Wang, X. Zhu, *Kähler–Ricci solitons on toric manifolds with positive first Chern class*, Adv. Math. **188** (2004), 87–103.

- [43] P. Yang, S.-T. Yau, *Eigenvalues of the Laplacian of compact Riemann surfaces and minimal sub-manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **7** (1980), 55–63.
- [44] S.-Y. Yau, *On Calabi's conjecture and some new results in algebraic geometry*, Proc. Nat. acad. U. S. A. **74** (1977), 1789–1799.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, CASE POSTALE 8888, SUC-CURSALE CENTRE-VILLE MONTRÉAL (QUÉBEC) H3C 3P8, CANADA
E-mail address: apostolov.vestislav@uqam.ca

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY, 805 SHERBROOKE STREET WEST, MONTRÉAL (QUÉBEC) H3A0B9, CANADA
E-mail address: jakobson@math.mcgill.ca

MATHEMATISCHES INSTITUT DER UNIVERSITÄT MÜNCHEN, THERESIENSTR. 39, D-80333 MÜNCHEN, GERMANY
E-mail address: Gerasim.Kokarev@math.lmu.de