



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/99241/>

Version: Accepted Version

---

**Article:**

Segura-Noguera, M., Cruzado, A. and Blasco, D. (2016) The biogeochemistry of nutrients, dissolved oxygen and chlorophyll a in the Catalan Sea (NW Mediterranean Sea). *Scientia Marina*, 80 (S1). pp. 39-56. ISSN: 0214-8358

<https://doi.org/10.3989/scimar.04309.20A>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

## **The biogeochemistry of nutrients, dissolved oxygen and chlorophyll *a* in the Catalan Sea (NW Mediterranean Sea)**

Mariona Segura-Noguera<sup>1,2\*</sup>, Antoni Cruzado<sup>3,4</sup>, and Dolors Blasco<sup>1</sup>

5 <sup>1</sup> Department of Marine Biology and Oceanography, Institut de Ciències del Mar, CSIC, Barcelona, Catalonia.

<sup>2</sup> Present address: ChELSI Institute, Department of Chemical and Biological Engineering, The University of Sheffield, Sheffield, UK

10 <sup>3</sup> Department of Aquatic Biogeochemistry, Centre d'Estudis Avançats de Blanes, CSIC, Blanes, Catalonia

<sup>4</sup> Present address: Oceans.cat, Anselm Clavé 8, Blanes 17300, Catalonia

\* corresponding author: ChELSI, Sir Robert Hadfield Building, Portobello Street, Sheffield, S1 3JD. Phone: +44 114 222 7594. Email: [mariona@marionasegura.cat](mailto:mariona@marionasegura.cat).

15

### **Summary**

Reference depth profiles of dissolved inorganic nutrients, dissolved oxygen and chlorophyll *a* are described for the Catalan Sea using quality controlled data. Phosphate, nitrate and silicate show typical nutrient profiles, with nutriclines at different depths.

20 Maximums of nitrite, dissolved oxygen, and occasionally ammonium, are found within the photic zone, close to the deep chlorophyll maximum. In intermediate waters we found a minimum of dissolved oxygen coincident with maximum concentrations of phosphate and nitrate. Ammonium concentration is unexpectedly high in the mesopelagic zone, where there are still measurable nitrite concentration, and their origin remains unclear. We also identify and describe anomalous data and profiles resulting from eutrophication, Western Mediterranean Deep Water formation, or dense shelf-water cascading. N:P ratio in deep waters is 22.4, which indicates P-limitation relative to the Redfield ratio, however, the N:P ratio above the deep chlorophyll maximum in stratified surface waters is <4 (<8 with ammonium). The depth profiles of key biogeochemical variables described in this study will be a useful reference for future studies in the Catalan Sea (NW Mediterranean Sea): to validate data sampled in this area, to identify

25

30

anomalous processes, and to study the evolution of the system following the undergoing global change.

**Keywords**

5 Dissolved inorganic nutrients; dissolved oxygen; chlorophyll *a*; stoichiometry; P-limitation; NW Mediterranean Sea; Catalan Sea

**Running title:** Biogeochemistry of the Catalan Sea

10

## INTRODUCTION

The Catalan Sea (Fig. 1) is an area of the North Western Mediterranean Sea that has been thoroughly studied, starting early 1900, with almost yearly oceanographic cruises from 1960 (e.g. Table A1). As a result of these studies, the dynamics of circulation, phytoplankton composition and primary production of the Catalan Sea are well known. In this study we use quality controlled data from cruises in the Catalan Sea to describe reference concentration depth profiles of key biogeochemical variables, including dissolved inorganic nutrients, dissolved oxygen and chlorophyll *a*.

The Mediterranean Sea is a concentration basin where evaporation greatly exceeds precipitation and river runoff. The water deficit is compensated for by the entrance of surface waters through the Strait of Gibraltar, while the outgoing flow of deep waters (about 4.7% nutrient enriched after spending between 75 – 100 years at the Mediterranean Sea, Cruzado 1985) prevents the accumulation of nutrients in deep Mediterranean Sea waters. As a consequence of this water regime deep water nutrient concentrations are lower than those at the neighboring Atlantic Ocean (Cruzado 1985). In addition, the Mediterranean Sea is characterized by having almost permanently stratified water columns, even though their intensity throughout the year is variable. In stratified water columns the nutrients regenerated in deep waters are prevented to reenter into the euphotic zone, where nutrient concentrations are low due to their consumption by osmotrophic organisms (Riley 1971). Consequently, the Mediterranean sea is an oligotrophic sea, with low levels of chlorophyll *a* in surface waters and hence, low levels of primary production are expected.

However, in the Western Mediterranean Sea moderate levels of primary production have been recorded, an observation that cannot be explained by the low nutrient content of surface waters. This observation has been named by Sournia (1973) as the “Mediterranean paradox”. In the Catalan Sea, mesoscale structures like permanent fronts, eddies and filaments, act favoring the upwelling of nutrients to surface waters, allowing the existence of the moderate levels of primary production (e.g. Estrada and Margalef 1988, Salat 1995, Granata et al. 2004). Moreover, the general cyclonic circulation of the Catalan sea creates a central divergence, where intermediate waters, richer in nutrients than surface waters, are closer to the photic zone, enhancing primary production (Estrada and Margalef 1988, Estrada 1999). However, it is during the late

winter months that the most important fertilizing event occurs during the formation of Western Mediterranean Deep Water (WMDW) at the so-called MEDOC area, located between the northern part of the Catalan Sea and south of the Gulf of Lion (Estrada 1996). The WMDW formation is a complex process, which involves strong, cold and dry northerly winds blowing on an area with cyclonic circulation and maximum surface salinity. As a result, a deep convection occurs and the whole water column (from 0 to 2800 m) may be homogeneous, leading to the formation of dense water at the surface that rapidly sinks and becomes a new water mass (MEDOC Group 1970, Salat 1983, 1996).

Maximum surface chlorophyll *a* concentrations are found during winter months (Estrada 1999, Morales-Blake et al. submitted), related to mixing of surface waters with deeper nutrient-rich waters, as well as to continental discharges from the rivers Ebre and Rhône. The Ebre River collects the runoff from Pyrenees, wastewater from large cities and intensive agricultural and industrial activities. Its outflow may account for 10–25% of the total nutrient content in the water column on the adjacent continental shelf (Cruzado et al. 2002, Salat et al. 2002). During the rest of the year, in stratified water column conditions, maximum chlorophyll *a* concentrations are found at the deep chlorophyll maximum (DCM), a typical structure found in oligotrophic systems that results from an accumulation of cells actively growing, and an increased pigment content per cell due to photo acclimation (Estrada 1985a). In the Catalan sea, the DCM is located close to the nitracline, between 40 – 60 m depth, where there is still light enough (1% surface radiation) to carry on photosynthesis (Margalef 1985, Estrada 1985b, Bahamón and Cruzado 2003). Coincident with the DCM is found the primary nitrite maximum (PNM) (Estrada et al. 1993, Bahamón and Cruzado 2003). The coincidence has been explained by the phytoplankton excretion of nitrite as a result of the incomplete nitrate assimilation for cellular requirements (Riley 1971), for example because of the low light intensities (e.g. Blasco 1971, Olson 1981). Finally, a maximum of dissolved oxygen is formed above the DCM as a result of photosynthetic activity (Estrada 1999), as found in other parts of the Mediterranean sea (e.g. Minas and Bonin 1988).

In this study we build up and describe depth concentration profiles of dissolved inorganic nutrients, dissolved oxygen and chlorophyll *a*. To do so, biogeochemical data

from cruises in the Catalan Sea were collected. Previously, we used part of this data to build a quality control for dissolved inorganic nutrients, dissolved oxygen and chlorophyll *a* (Segura-Noguera et al. 2011). In this study we present a complete and exhaustive description of the biogeochemistry of the Catalan Sea made with quality controlled data, including the description of the concentration depth profiles of these basic oceanographic variables, its seasonal variation in surface waters, and the description of the stoichiometry between phosphorus, nitrogen, silicon and apparent oxygen utilization. This basic description of the biogeochemistry of the Catalan Sea will be a useful reference for future studies in the NW Mediterranean Sea, as well as to validate data sampled in the same area, to identify anomalous processes, and to study the evolution of the system following the undergoing global change.

## **MATERIAL AND METHODS**

### **Circulation and water masses of the Catalan Sea**

The Catalan Sea is the part of the Balearic Sea situated between the northeastern part of the Iberian Peninsula and the Balearic Archipelago (Fig. 1). It limits with the Gulf of Lion and the Ligurian Sea at the North and with the Balearic Sea at the South. The Catalan Sea is a 400 km long subbasin that reaches a maximum depth of 2500 m. Its general circulation is cyclonic, with a central dense water mass surrounded by lighter continental and Atlantic surface waters (Font et al. 1988). Between the central and peripheral waters there are two permanent fronts, the Catalan and the Balearic fronts, that follow the continental shelf-break of the Iberian Peninsula and the Balearic islands, respectively. The Catalan front is caused by differences in salinity (S) between continental and central waters, while the Balearic front is caused by differences in temperature (T) and S between Atlantic and central waters (Font et al. 1988). The Catalan current, associated to the Catalan front, is an extension of the Liguro-Provençal current, also known as Northern current. It occupies the top 300 – 400 m of the water column, flowing towards the southeast following the 1000 isobath. In front of the Gulf of Valencia, part of this current creates a cyclonic gyre, incorporates Atlantic waters, and continues flowing towards the northwest, following the Balearic shelf-break, which generates the Balearic current (Millot 1987, Font et al. 1988, Salat 1995). The Catalan current, with average velocities between 20 – 30 cm s<sup>-1</sup>, has a marked seasonality, its

intensity being maximum during autumn and decreasing towards a minimum during the summer months (Font et al. 1988, Send et al. 1999).

The northernmost continental shelf of the Catalan Sea is narrow (less than 25 km width) and irregular due to the presence of submarine canyons, most of which finish in submarine canals (Canals et al. 1982). Submarine canyons and other topographic features, like the expansion of the continental shelf in front of the Ebre Delta (Fig. 1), lead to deviations of the Catalan current, creating eddies and intrusions (filaments) of denser and saltier water to the continental shelf (e.g. Font et al. 1988, Masó et al. 1990, Granata et al. 2004).

In the Catalan Sea four water masses can be found: Surface Water (SW), Winter Intermediate Water (WIW), Levantine Intermediate Water (LIW) and Western Mediterranean Deep Water (WMDW). Three different surface water masses can be distinguished by its S and location (Salat and Cruzado 1981, Font et al. 1988; Salat 1995): (1) surface water with continental influence is located close to the peninsular coast, containing water from the rivers Rhône and Ebre, which results in a S between 35.5 and 37; (2) close to the Balearic Islands there is surface water with Atlantic influence, which S increases from 36.5 to around 37 – 37.3 as a result of progressive evaporation from its entrance to the Mediterranean Sea; and (3) at the center of the basin are found resident surface waters (also known as “Mediterranean Waters”, MW, or “Modified Atlantic Waters”, MAW), with the highest S (37.8 – 38) as a result of evaporation during the time spent in the Mediterranean basin.

Below surface waters two types of intermediate waters can be found. WIW is formed during winter by convection of surface waters at the continental shelf, both from the Gulf of Lion and the Catalan Sea, without mixing with deeper waters. It is characterized by a minimum of potential temperature around 12 – 12.5°C, and salinity between 38.1 and 38.3. The LIW is located between 200 and 600 m, and it is the result of the mixing of Levantine water formed in winter at the Aegean Sea and Cyprus with winter water formed at the Adriatic Sea. It is the saltiest water, with a S maximum of 38.55 and a maximum of potential T of 13.5°C. In case of coexistence, WIW is found above LIW (Salat and Font 1987). Finally, the Western Mediterranean Deep Water (WMDW) (MEDOC Group 1970) is found below the LIW down to the bottom of the

basin. It has a potential temperature between 12.65 and 12.9°C, a S of 38.45 – 38.48, and a density between 29.08 and 29.11 kg m<sup>-3</sup>.

### **Biogeochemical datasets**

5           The biogeochemical data used in this study included published historical data, biogeochemical datasets of cruises where the authors have participated, and data extracted from MEDAR and MEDATLAS databases (0 – 4.5 E, 38.7 – 42.5 N) (Table A1). Apart from T, S and depth, each cruise included some or all of the following biogeochemical variables: dissolved inorganic nutrients (phosphate, nitrate+nitrite  
10 –from now on, nitrate–, nitrite, ammonium and silicate), dissolved oxygen and chlorophyll *a* (from now on, chl<sub>a</sub>) concentration. The list of biogeochemical variables available per each cruise gathered, as well as the source of the data, is shown in Additional Information Table A1. The final dataset was carefully checked to avoid any duplicated data. The whole pool of data was quality controlled using flags, where data  
15 within the ranges of concentration described in Segura-Noguera et al. (2011) were flagged “0”, data above were flagged “1”, and below were flagged “2”. In the present study, only data with flag “0” (Table 1) have been used to describe depth profiles and to characterize water masses. Sampling stations for each oceanographic variable are shown in Additional Information, Fig. A1. The main features of the analyses of dissolved  
20 inorganic nutrient, dissolved oxygen and chlorophyll *a* (chl<sub>a</sub>) of the dataset, when available, are exhaustively described in Segura-Noguera et al. (2011) and summarized below.

          Four different autoanalyzers were used for nutrient analysis (Technicon, Skalar, Evolution-II of Alliance and AA3 of Bran+Luebbe), using similar methodologies  
25 (Hansen and Koroleff 1999). A salycilate-hypochlorite method was used for ammonium analysis between 2001 and 2004 (Bower and Holm-Hansen 1980). Until 1999, samples were usually analyzed on board, and after that year they were preserved by freezing and analyzed on land. Nutrient standards were always prepared at room temperature (20-25°C). The analysis precision of each biogeochemical variable was estimated for  
30 several cruises using different methods (Segura-Noguera et al. 2011). Following, the detection limit for the whole dataset was estimated as 3 times the average of the analysis precision calculated after removing the outliers with the interquartile method. The

5 resulting detection limits were: 0.02  $\mu\text{M}$  for phosphate, 0.01  $\mu\text{M}$  for nitrite, 0.08  $\mu\text{M}$  for  
nitrate, 0.07  $\mu\text{M}$  for ammonium, and 0.12  $\mu\text{M}$  for silicate. Detection limits of the  
2001-2004 cruises “Canyons” I to IV and “Caco” 1 to 4 were calculated during the  
present study, and were 0.006  $\mu\text{M}$  for phosphate, 0.004  $\mu\text{M}$  for nitrite, 0.015  $\mu\text{M}$  for  
10 nitrate, 0.020  $\mu\text{M}$  for ammonium, and 0.017  $\mu\text{M}$  for silicate. Nutrient concentrations are  
given in  $\mu\text{M}$  instead of  $\mu\text{mol kg}^{-1}$  because data of salinity and temperature were not  
available for all the correspondent nutrient concentrations. Consequently, the conversion  
to  $\mu\text{mol kg}^{-1}$  would represent a loss of around 6% of inorganic nutrients data and up to  
11.2% of ammonium data, a very scarce nutrient in our dataset. Except for very few  
15 stations close to the Ebre Delta with freshwater signature, the  $\sigma\text{-T}$  range in our dataset is  
13.7 to 29.4  $\text{kg m}^{-3}$ . This difference means that the values in  $\mu\text{M}$  are 2.9% larger, on  
average, than those in  $\mu\text{mol kg}^{-1}$ , and that the maximum difference between both units  
(corresponding to the largest concentrations of these variables in the Catalan Sea) are  
around 0.27 of nitrate, 0.30 of silicate, 0.07 of ammonium, 0.02 of nitrite, 0.01 of  
15 phosphate and 8.9 of dissolved oxygen.

Throughout the years, dissolved oxygen was consistently measured following  
Winkler (1988). The difference between cruises was the method for the determination of  
the final titration point, which was visual or automatic. Dissolved oxygen data were  
converted to  $\mu\text{M}$  using the molar volume of oxygen gas (22.3916  $\text{L mol}^{-1}$ ) and the  
20 density of water at zero pressure calculated with SEABIRD software from CTD  
temperature ( $^{\circ}\text{C}$ , converted to potential temperature with the same software) and salinity  
(psu). The detection limit of dissolved oxygen of the whole dataset was 7.7  $\mu\text{M}$ . All our  
dissolved oxygen values were above detection limit.

The Apparent Oxygen Utilization (AOU,  $\mu\text{M}$ ) was calculated as the difference  
25 between dissolved oxygen in saturation ( $[\text{O}_2]^*$ ) calculated from *in situ* T and S following  
Owens and Millard (1985), and the measured dissolved oxygen in the water ( $[\text{O}_2]_{\text{meas}}$ ).

Chla data were determined fluorometrically after filtering with glass fiber filters  
and extracting with acetone (Yentsch and Menzel 1963). The average precision of chla  
analysis of all cruises was 0.01  $\mu\text{g l}^{-1}$  (Segura-Noguera et al. 2011).

30

## Depth profiles and stoichiometry

Depth profiles have been built up for dissolved nutrients, chlorophyll *a* and dissolved oxygen by calculating the median value at each standard level depth (SLD). SLDs are the depths that have been historically most sampled, and include data  
5 comprised between 25% of the distance with the upper SLD and 75% distance with the lower SLD. We have used the SLDs proposed for the Mediterranean Sea by Fichaut et al. (1997): 0, 5, 10, 20, 30, 40, 50, 60, 80, 100, 120, 160, 200, 250, 300, 400, 500, 600, 800, 1000, 1200, 1400, 1600, 1800, 2000, and 2500 m. Continental shelf and open sea profiles were differentiated because the area closest to the coast (bottom depth up to 200  
10 m, as in Fichaut et al. 1997) is theoretically influenced by coastal and anthropogenic processes that could presumably increase the concentration of the different oceanographic variables. The data were grouped according to the stratification index estimated for each station and profile depth as described in Segura-Noguera et al. (2011). The most representative seasons for each stratification index are winter for “1”,  
15 spring for “2”, summer for “3” and autumn for “4”.

Depth median concentration profiles of each variable, as well as stoichiometry profiles, were created for each stratification index. Depth-integrated (0 – 100 m) T, S, density, as well as nutrients, chl*a* and dissolved oxygen concentrations were calculated for surface waters. Seasonal differences in depth-integrated concentrations in surface  
20 waters were studied with the software PAST v 3.06 (Hammer et al. 2001). Univariate non-parametric Kruskal-Wallis test was used to find differences, followed by a post-hoc Mann-Whitney pairwise comparison with Bonferroni corrected *p* values. Statistical significance was accepted when  $p \leq 0.05$ .

The molar stoichiometry between nutrients as well as dissolved oxygen was  
25 calculated for each available pair of data. Data below the estimated detection limits were not used to calculate the stoichiometry. Element:element correlations were studied using Standard Major Axis (SMA), a regression model II that takes into account differences in the scale of the axes (e.g. N:P relationship). The software SMATR Ver. 2.0 (Falster et al. 2006) was used to calculate SMA slopes and intercepts, along with *p* values and  
30 confidence intervals, as well as seasonal differences in slopes and intercepts.

## RESULTS

### Concentration depth profiles

Figs. 2, 3 and 4 show the seasonal depth profiles of T, S, density anomaly and of biogeochemical variables. The concentration of each variable at each SLD and season can be obtained from Additional Information Table A2. A clear seasonality in temperature is found in surface waters (Fig. 2, Table 2). Warming of surface waters in spring outsets the stratification of the water column and the development of a mixed layer. During summer, the mixed layer depth is minimum and a strong thermocline can be observed. The stratification weakens during autumn and the thermocline disappears in winter (Fig. 2). Surface nutrients and chl<sub>a</sub> concentration follow the general pattern of thermal stratification, with larger concentrations in winter and minimum surface concentrations during maximal stratification of the water column (Figs. 3 and 4).

Phosphate, nitrate and silicate show the typical nutrient depth profile, with low surface concentrations that increase with depth (Fig. 3). In open sea stations, the largest surface (0 – 50 m) phosphate, nitrate and silicate concentration is found in winter (0.11, 1.07 and 1.98  $\mu\text{M}$ , respectively), and decrease throughout spring to reach minimum values in summer (0.04, 0.11 and 0.63  $\mu\text{M}$ , respectively). In autumn, nitrate and phosphate concentrations increase and are similar to those in spring and winter, respectively, while silicate concentration remains low. Depth profiles from continental shelf stations show a similar pattern, except that summer phosphate concentration is similar to spring (0.06  $\mu\text{M}$ ) and that nitrate concentration in winter is higher than in the open sea (1.70  $\mu\text{M}$ ). In winter mixed water columns, nutrients concentration increase linearly from surface to 400 m depth at different rates (phosphate: 0.7  $\text{nM m}^{-1}$ ,  $r^2=0.92$ ; nitrate: 19.4  $\text{nM m}^{-1}$ ,  $r^2=0.98$ ; silicate: 12.7  $\text{nM m}^{-1}$ ,  $r^2=0.95$ ). In spring, a nitracline and a phosphocline starts to develop in open ocean at 60 and 100 m depth, respectively. The nitracline is deeper in summer and autumn (80 m, Fig. 3). In the continental shelf, the phosphocline is smoother than in open sea stations. Phosphate and nitrate concentrations continue to increase with depth until 500 m. In spring, a maximum of phosphate concentration (0.44  $\mu\text{M}$ ) can be observed at this depth, but not of nitrate, whose concentration is around 8.6  $\mu\text{M}$  from 500 m to the bottom of the basin. Silicate concentration continues to increase below 500 m depth, and concentrations around 8.8 – 9.2  $\mu\text{M}$  are found below 1200 – 1500 m depth. The average ( $\pm$  standard deviation)

nutrients concentration in deep waters (WMDW with depth  $\geq 800$  m) are  $0.40 \pm 0.05$   $\mu\text{M}$  of phosphate,  $8.68 \pm 0.04$   $\mu\text{M}$  of nitrate and  $8.72 \pm 0.91$   $\mu\text{M}$  of silicate.

Ammonium concentration in open sea stations (0 – 200 m depth) is higher in winter, decreasing throughout spring to a minimum in summer (around 0.6, 0.2 and 0.1  $\mu\text{M}$ , respectively) (Fig. 4). In autumn, ammonium concentration increases again and is higher than that in spring (around 0.4  $\mu\text{M}$ ). In continental shelf stations this seasonality can not be clearly seen, and the largest concentrations are found in winter below 30 m depth. It is difficult to detect clear features below 200 m depth because of the scarcity of data. The average concentration below 250 m depth is around  $0.64 \pm 0.34$   $\mu\text{M}$ .

Surface chla, nitrite and dissolved oxygen concentrations show a similar pattern, with highest concentrations in winter and lowest in summer, and with a presence of a maximum (Fig. 4). In winter, high chla concentration is found between 0 – 60 m (around  $0.75$   $\mu\text{g l}^{-1}$  in open ocean and  $0.61$   $\mu\text{g l}^{-1}$  in continental shelf stations), and two maximums can be observed, one above 20 m ( $0.86$  and  $0.75$   $\mu\text{g l}^{-1}$  in open sea and continental shelf stations, respectively) and another at 50 m ( $0.83$  and  $0.68$   $\mu\text{g l}^{-1}$ , respectively). The maximum at 50 m is coincident with a maximum of ammonium and of nitrite, although only in open sea stations ( $1.21$  and  $0.23$   $\mu\text{M}$ , respectively). Surface chla, nitrite and dissolved oxygen concentrations decrease during spring. Chla and nitrite accumulate between 40 – 100 m depth, forming the DCM around 50 – 60 m depth ( $0.58$  and  $0.49$   $\mu\text{g l}^{-1}$  in open sea and continental shelf, respectively) and PNM around 60 – 80 m depth ( $0.11$  and  $0.16$   $\mu\text{M}$ , in open ocean and continental shelf, respectively), while dissolved oxygen forms a shallower maximum around 30 – 40 m depth ( $261.7$  and  $262.6$   $\mu\text{M}$  in open ocean and continental shelf, respectively). In summer, the DCM is larger and deeper than in spring ( $0.61$   $\mu\text{g l}^{-1}$  at 80 m in open sea, and  $0.62$   $\mu\text{g l}^{-1}$  at 60 m in continental shelf stations), as well as it is the dissolved oxygen maximum ( $272.9$   $\mu\text{M}$  at 40 m in open sea, and  $268.4$   $\mu\text{M}$  at 50 m in continental shelf stations). Contrarily, the PNM is lower ( $0.11$  and  $0.08$   $\mu\text{M}$  in open sea and continental shelf stations, respectively) and only deeper in open sea stations (80 m). In autumn, surface chla and nitrite increase and are similar to spring values, while dissolved oxygen is lower than in spring. The DCM is shallower (around 50 m) and lower than the rest of the year ( $0.38$  and around  $0.3$   $\mu\text{g l}^{-1}$  in open ocean and continental shelf stations, respectively), while the PNM is higher in continental shelf than in open ocean ( $0.21$  and  $0.07$  –  $0.09$   $\mu\text{M}$ ,

respectively). The dissolved oxygen maximum is at the same depth than in summer, but its intensity is much lower (246.5  $\mu\text{M}$  in open sea, and 242.9  $\mu\text{M}$  in continental shelf stations).

Below the DCM, chl $a$  concentration decreases until it falls below the detection limit around 200 m, except in winter, when concentrations of 0.03 – 0.06  $\mu\text{g l}^{-1}$  have been measured between 300 and 600 m depth. Below the PNM, the nitrite concentration also decreases, but its concentration remains detectable throughout the whole water column (up to 0.08  $\mu\text{M}$ ). Dissolved oxygen concentration decreases below the surface maximum to reach minimum values at 500 – 800 m depth, although its intensity is variable (189.8 – 201.0  $\mu\text{M}$ ), and below 1400 m to the bottom of the basin, it increases again up to around 205.9 $\pm$ 3.64  $\mu\text{M}$ .

Surface water (0 – 50 m) dissolved oxygen concentrations, as well as saturation concentrations, plotted vs T are shown in Figure 5. Open sea stratified surface waters (0–50 m) of the Catalan Sea are oversaturated in dissolved oxygen (104.9% in spring, 108.5% in summer and 101.8% in autumn). In open sea mixed water columns in winter, are undersaturated (mean value 98.0%). This winter value is highly influenced by the low values (<230  $\mu\text{M}$ ) at 13°C, that corresponds to stations sampled during WMDW formation. In the continental shelf, the oversaturation is found throughout the year in surface waters between 0 – 40 m in spring (106.8%) and between 0 – 50 m in summer and autumn (107.0% and 102.3%, respectively), but the oversaturation in continental shelf mixed water columns is restricted to the upper 20 m (102.5% in winter). Both in stratified open sea waters and continental shelf waters, the depth where the consumption of oxygen overcomes its production, i.e. where median AOU values turn positive, is just above the DCM. The absolute minimum dissolved oxygen concentration found at 500-600 m depth (189.8 – 201.0  $\mu\text{M}$ ), the same depth of maximum phosphate and nitrate concentrations are found, is corresponded by an absolute maximum of AOU (58.4 – 69.5  $\mu\text{M}$ ), which supposes a consumption of around 23.7 – 25.2% of the saturating oxygen.

## Stoichiometry

Surface water ratios are shown in Table 3. In open sea, the N:P ratio under stratified conditions (spring to autumn) from 0 to 50 m depth is lower than the Redfield ratio (median±median absolute deviation,  $2.6\pm 1.3$ ,  $n=717$ ) (Fig. 6). In the continental shelf, the N:P ratio is significantly larger in autumn than in spring and summer. The N:P ratio significantly increases when ammonium is added to the calculations, but it is still below the Redfield ratio. In open sea, DIN:P in summer is higher than in spring and autumn; while in the continental shelf the DIN:P ratio in summer and autumn is larger than in spring. In mixed water winter conditions, the N:P ratio is higher and more variable. Below the photic layer (around 200 m depth to the bottom of the basin) the average±standard deviation N:P ratio is  $22.6\pm 5.7$  ( $n=1203$ ), and DIN:P is  $21.9\pm 5.8$  ( $n=194$ ). These values are similar to the slopes obtained with regression lines for all N:P ( $22.6$ ,  $r^2=0.646$ ,  $n=6560$ ) and DIN:P data ( $22.2$ ,  $r^2=0.578$ ,  $n=1605$ ), which are significantly different from 16 ( $p\leq 0.001$ ) (Table 4). Lower DIN:P compared to N:P ratios are a consequence of using different datasets, since not all the variables are analyzed in all the cruises (See Table A1).

Median Si:N ratio shows a clear seasonality in surface open sea waters (0 – 50 m), with significantly lower ratio in winter and higher ratio in summer, both in terms of average and standard deviation (Fig. 6, Table 3). The median Si:N is not different from spring to autumn (pooled Si:N= $4.4\pm 2.7$ ,  $n=1825$ ). As in open ocean, the lowest Si:N ratio in the continental shelf is found in winter, and there are no significant differences between the ratio from spring to autumn (pooled ratio:  $5.3\pm 2.6$ ,  $n=1108$ ). Below 1200 m depth average Si:N concentration is  $1.0\pm 0.1$  ( $n=144$ ). In surface waters (0 – 50 m) median Si:P ratios are highly variable (Fig. 6, Table 3) and, like the Si:N ratio, the largest values are in winter and the lowest are in autumn in open sea, and in summer in the continental shelf (Table 3). In open sea, Si:P ratios in spring and summer are the same (pooled ratio:  $16.3\pm 9.1$ ,  $n=854$ ). Below 1200 m depth, the average Si:P ratio is  $23.2\pm 3.4$  ( $n=69$ ).

The standard major axis regression slopes of AOU:P and AOU:N are 150.2 ( $r^2=0.353$ ,  $n=1420$ ) and 8.9 ( $r^2=0.603$ ,  $n=1864$ ) (Table 4). The scatter plots in Figure 7 show that Si:N, Si:P and AOU:Si relationships are not linear, since silicate concentration still increases below the depth where phosphate and nitrate have been completely

remineralized, corresponding to the WMDW and deepest LIW (from 500 – 600 m depth). Therefore, we have not calculated the linear relationship for these ratios. The average AOU:Si calculated from pairs of data in deep waters (>1200 m depth) is  $6.06 \pm 0.7$  (n=58).

5

## DISCUSSION

In this study we have used quality controlled data from the Catalan Sea to build seasonal depth profiles of dissolved inorganic nutrients, oxygen and chlorophyll *a*, and to study nutrients and dissolved oxygen stoichiometry in the water column. These general profiles are useful as a reference values for future biogeochemical studies in the Catalan Sea. Furthermore, they allow the identification of anomalous profiles resulting from eutrophication, WMDW formation, and dense shelf-water cascading (DSWC). Below we describe depth profiles of key biogeochemical variables, comparing them with other areas of the Mediterranean Sea and of the global oceans, we show and discuss anomalous profiles and describe surface seasonality and stoichiometry.

10  
15

### Concentration depth profiles

Depth distribution of phosphate, nitrate and silicate described in this study (Fig. 3) show the typical nutrient profile, with low surface concentrations due to their assimilation by phytoplankton and other osmotrophic organisms, followed by an increase in concentration with depth, as described in other parts of the NW Mediterranean Sea (e.g. Marty et al. 2002 at the neighboring DYFAMED site). Deep water phosphate, nitrate and silicate concentrations are in agreement with published concentrations from the Western Mediterranean Sea (e.g. Ballester et al. 1967, Cruzado 1985, Béthoux et al. 1998).

20  
25

Maximum concentrations of phosphate and nitrate in intermediate waters have been observed in other areas of the Mediterranean Sea (around 400–700 m, Béthoux et al. 1998, Moutin and Raimbault 2002). Although a clear maximum can not always be distinguished in our seasonal median profiles, the observation of individual profiles confirms the presence of higher nitrate and phosphate concentrations between 500 to 800 m depth, that follow the pattern of the depth profiles of salinity. In addition, we have also observed in individual profiles the occasional presence of a nutrients peak

30

between 150 – 300 m depth, associated again to a peak in salinity. Although the intensity of this peak is variable, it still can be observed in the median profiles of nitrate and phosphate (Fig 3).

A maximum of nutrients and AOU associated to a minimum of dissolved oxygen in the mesopelagic zone (100 – 1000 m) is a feature present in most of the oceans (Sarmiento and Gruber 2006), as well as between 400 and 800 m depth in other areas of the NW Mediterranean Sea (e.g. Marty et al. 2002, Moutin and Raimbault 2002). It has been associated to both animal respiration and active microbial remineralization of organic material, mostly dead phytoplankton, faecal pellets and other detrital particles, that sink from surface waters (e.g. Sarmiento and Gruber 2006). Moreover, large zooplankton organisms and micronekton also play an important role in transporting dissolved organic carbon and nutrients from surface waters to the mesopelagic zone through diel vertical migrations (e.g. Longhurst and Harrison 1988, in the NW Mediterranean Sea: Isla et al. 2015). A higher abundance of animals and microbes in certain depth of the mesopelagic zone, for example because of the accumulation of sinking organic matter in the interface of two water masses of different density (like WIW and LIW or LIW and WMDW), could lead to the accumulation of nutrients, and hence to the formation of a maximum. Also, the association of the nutrients maximum and dissolved oxygen minimum with salinity indicates that their origin is the remineralization of surface organic matter that has been trapped within the LIW (or WIW) during its formation and sinking. It is unlikely that the decrease in nitrate from this maximum to the bottom of the basin (Fig. 3) is due to denitrification, since this process requires lower oxygen concentrations than those measured at deep Catalan Sea waters (Riley 1971, Millero 1996).

Phosphate and nitrate concentrations remain approximately constant below the maximum in intermediate waters to the bottom of the basin, indicating that all the labile phosphate and nitrate contained in sinking organic matter have already been remineralized. On the other hand, silicate concentration steadily increases from surface to 1400 m depth, from where it remains constant down to the bottom of the basin. This difference in the nutrient profiles is a consequence of the slow chemical dissolution of siliceous diatom frustules throughout the water column, as opposed to the active remineralization of phosphate and nitrate, carried out mainly by bacteria and other

protozoa (Millero 1996). The different remineralization rates between these nutrients results in the non-linearity of the Si:N, Si:P and AOU:Si ratios, as opposed to AOU:P, AOU:N and N:P, as can be observed in Figure 7.

5 A statistically significant relationship between the depths of the DCM, the PNM and the nitracline has been described in the Catalan Sea (Estrada 1999) using a small set of stations included in the present study (Fronts 90, 91, 92 and Varimed 93). Estrada (1999) compared the position of the maximums and nitracline in 58 stations, showing that the PNM was almost coincident with the nitracline, the DCM was slightly above them, and finally the maximum of dissolved oxygen was the shallower. This pattern is almost the same that the resulting of comparing our median profiles, except that in our profiles the DCM is found at the same depth than both the PNM and the nitracline in winter and summer, and immediately below in spring and autumn (Fig. 4). However, the observation of individual profiles show that these maximums are not always coincident, even in consecutive days of the same cruise. This differences could be related to changes in the nitrite release by phytoplankton at the DCM driven by light or nutrient availability, that could also affect the abundance and/or activity of nitrifiers. Unfortunately, there are no data about light or organisms abundances in our dataset that could help us to understand the relative changes in the peaks position.

20 In other oligotrophic areas (North Pacific, Bering Sea, Greenland Sea, Sargasso Sea and Gulf Stream) a maximum in ammonium concentration has been detected close to the thermocline (i.e. under stratified conditions) and to the DCM (e.g. Saino et al. 1983, Brzezinski 1988), an indication of the importance of recycled production in oligotrophic seas. The examination of individual profiles also confirms the accumulation of ammonium around the DCM and PNM in stratified surface waters. However, this pattern is not reflected in the median profiles because a clear peak is not always observed (Fig. 4). On the other hand, under mixed water conditions in the Catalan Sea, an absolute maximum of the median ammonium concentration ( $1.21 \mu\text{M}$ ) is commonly observed at 50 m overlapping with a relative maximum of chl<sub>a</sub> and an absolute maximum of nitrite.

30 The analysis of ammonium is very sensitive to environmental contaminants and processes both *in situ* and on land (e.g. cleaning products, smoke, unclean glassware, or freezing as preservation method, see Holmes et al. 1999). Thus, large ammonium

concentrations are usually discarded assuming possible sampling or analysis contamination. Taking into account the ease of contamination, the lack of sampling details and the scarcity of the ammonium dataset it is striking the clear seasonality found. Ammonium concentrations in open sea surface waters follow the same seasonal pattern than phosphate, nitrate and silicate, with highest concentrations in winter and lowest in summer. The accumulation of ammonium in surface waters demonstrates an imbalance in the production-consumption processes, especially in winter, when the highest primary production takes place. In the North Atlantic Ocean has been described an increase in ammonium concentration after a silicate-limited diatom bloom, which reversed the direction of the air-sea flux of this nutrient (Johnson et al 2007). The molecules of atmospheric ammonium suffer a series of chemical reactions and can act as cloud condensation nuclei (Johnson et al. 2007). In the Catalan Sea, diatoms dominate the winter blooms (Estrada 1999). Although diatom late winter-spring bloom in the Catalan Sea are most probably not silicate limited (Si:N and Si:P ratios in winter surface waters are larger than in deep waters, Fig. 6), the accumulation of ammonium (median ~ 0.6  $\mu\text{M}$ , similar to that in the study of Johnson et al. 2007) indicates that a release of ammonium to the atmosphere could also be happening in the NW Mediterranean Sea during winter, something that remains to be quantified.

Ammonium and nitrite concentrations are still measurable below the euphotic layer, throughout the water column (below 400 m depth:  $0.59\pm 0.36 \mu\text{M}$  ammonium and  $0.04\pm 0.02 \mu\text{M}$  nitrite, Fig. 4). This result is unexpected in an oxic environment like the Catalan Sea, and the origin of such high values remains unclear. Relatively high ammonium concentrations ( $0.4 \mu\text{M}$  at 800 m depth) in deep waters have also been reported in other oligotrophic areas, like the North Pacific Subtropical Gyre (Karl and Knauer 1984). The authors suggested that its origin was the chemolithotrophic activity associated to sinking particles, and their accumulation was the result of the inefficiency of the aphotic microbial community to use them (Karl and Knauer 1984). Assuming that the amount of sinking particles is proportional to surface water primary production, our average ammonium concentration is larger than that measured in the North Pacific Subtropical Gyre, both systems having similar primary production estimates:  $300 - 400 \text{ mg C m}^{-2} \text{ day}^{-1}$  in the Western Mediterranean (Estrada 1996) and  $463 \text{ mg C m}^{-2} \text{ day}^{-1}$  in the North Pacific Subtropical Gyre (Karl et al. 1996). However, in the Catalan Sea,

additional particles from the shelf and slope can also reach the water column, e.g. through nepheloid layers, trawling (e.g. Palanques et al. 2006), or through dense shelf water cascading (DSWC, see below). Also, several studies point out that surface particulate organic matter in the NW Mediterranean Sea is N-rich (e.g. Copin-Montégut and Copin-Montégut 1983, Mostajir et al. 1998). Then, chemolithotrophy activity in sinking particles appears to be a realistic pathway for the accumulation of ammonium in the mesopelagic zone of the Catalan Sea.

### **Deviations from reference depth profiles**

In this section we inspect data that have failed the quality control, i.e. they have been excluded from the establishment of the reference depth profiles described above. We have identified three processes that affect the concentration of biogeochemical variables in the Catalan Sea: coastal eutrophication, dense shelf water cascading and deep water formation.

High nutrients concentration in surface coastal waters due to river discharges or anthropogenic influence can be easily identified in our dataset, even though coastal conditions are poorly represented in this study: only 5.3% of the stations have a bottom depth lower than 50 m, the lowest bottom depth being 20 m. The Catalan coast shows a high variability in nutrient concentrations in response to the characteristics of the coast (i.e., beaches, rocky coast, harbors), and the existence of continental or anthropogenic outflows (Flo and Camp 2005). This variability is also influenced by seasonality. For example, before the installation of a wastewater treatment plant, nutrient and chl<sub>a</sub> concentration in surface waters of the Blanes Bay reached maximum values during summer months related to the increase in population (Duarte et al. 1999, Guadayol et al. 2009). However, the present study shows that in the continental shelf the lowest nutrient concentrations are found in summer, suggesting that occasional coastal fertilization processes have local effects, and can not be distinguished with the dataset used. Still, data outside the QC ranges of concentration in surface waters occur mainly near the Ebre River, and large cities like Barcelona, Tarragona and Palma (see Fig. 1), indicating that these high values are related to anthropogenic activity and river discharges.

In the open sea, anomalous depth profiles are found in the northern part of the Catalan Sea in winter, and are related to Western Mediterranean Deep Water (WMDW)

formation and to dense shelf water cascading (DSWC). WMDW is formed in winter in the Gulf of Lion and the northern part of the Catalan Sea (MEDOC Group 1970, Salat 1983, Font et al. 1988). During WMDW formation, T, S, and density characteristics are homogeneous throughout the whole water column (0 – 2800 m) (Salat 1996). Nutrients, density anomaly and dissolved oxygen profiles of two stations of the cruise Hivern 2000, sampled during the process of WMDW formation, are shown in Figure 8. It can be observed that the biogeochemical depth profiles resulting of the mixing during the WMDW formation are also homogeneous throughout the water column, with concentrations equivalent to deep waters. For comparison, the median winter concentration profiles of the same variables are also plotted in Fig. 8, as well as the limits of the winter quality control ranges. These profiles “fail” the quality control, which is constructed based on depth. However, the concentrations are in agreement with those measured at deep waters, which can be easily identified by its S,  $\theta$ , and  $\sigma$ -T characteristics.

DSWC also occurs in winter, when cold and dry northerly winds cause heat loss and evaporation of surface Gulf of Lion shelf waters. The waters, then, become cooler and mix with off-shelf waters, becoming denser than the surrounding waters and rapidly sinking, overflowing the shelf edge and cascading downslope through submarine canyons. Depending on the resulting density, the waters can reach the bottom of the basin, or detach from the slope when they reach its equilibrium with surrounding waters (Canals et al. 2006). In the Cap de Creus Canyon (Fig. 1) there have been identified four major cascading events on 1980, 1988, 1999 and 2005 (Canals et al. 2009). In our dataset, we have identified relatively large ( $\sim 12 \mu\text{M}$ ) nitrate and silicate concentrations in La Fonera Canyon, and very large concentrations below 500 m depth in winter of 1999 (cruise Hivern 1999), up to  $41.5 \mu\text{M}$  nitrate and  $24.5 \mu\text{M}$  silicate (Fig. 9), which are more than 3 times the corresponding concentrations in deep waters. The corresponding phosphate values are also high (between  $0.37$  and  $0.71 \mu\text{M}$ ), but because the range of concentration of phosphate in the Catalan Sea is narrower, they can not be as easily identified. In shelf surface waters (0-5 m depth) of the Gulf of Lion in winter, high nutrients concentrations have been measured in several cruises (up to  $41.0 \mu\text{M}$  nitrate,  $42.7 \mu\text{M}$  silicate and  $2.00 \mu\text{M}$  phosphate), including years of intense cascading (Pelagolion 2 in 1986, Discovery88 and Pelagolion 3 in 1988, and Tyro in 1991). These

high concentrations are related to nutrient discharges from the Rhône River (Cruzado and Velásquez 1990). We suggest that deep shelf cascading of winter surface waters, enriched in nutrients through the discharge of the Rhône River, could be the process that causes the occasional presence of high nutrients in intermediate waters in the Catalan sea.

### **Biogeochemical seasonality in surface waters**

Seasonal differences in the biogeochemistry of surface waters have been studied by the statistical comparison of integrated concentrations (0 – 100 m). As expected, a clear seasonality in T is found in surface waters (Salat 1996, Duarte et al. 1999), which drives the stratification of the water column (Segura-Noguera et al. 2011). The stratification is broken by convective processes (vertical mixing) related to an increase in the wind velocity and storms during autumn and winter (Salat 1996, Send et al. 1999, Guadayol and Peters, 2006), resulting in an enrichment of surface waters. Surface S does not show such strong seasonality, but even so, higher values are found in autumn and winter due to wind-driven evaporation and because of winter mixing with deeper and saltier waters (Salat 1996), which results in larger surface density anomaly in this season. Seasonal biogeochemical differences are not always statistically significant (Table 2). For example, phosphate average (0 – 100 m) concentration in winter is not statistically different from autumn, which could be the result of their fast uptake by phytoplankton, as soon as nutrients become available.

Chla average concentration (0 – 100 m) is larger in mixing conditions (winter) than during the rest of the year, in contrast to the most usual spring bloom (Longhurst 1988). This is possible because of the higher irradiance (around 20%) as well as warmer temperatures (3 to 5°C higher) in the Mediterranean Sea compared to the Atlantic Ocean at the same latitude (Duarte et al. 1999), as well as because deep mixing occurs during winter (e.g. Salat 1996), which result in nutrient enriched surface waters. Chla concentration decreases from winter through summer following a decrease in autotrophic organisms, both because the nutrients are being consumed (and hence the DCM moves towards deeper waters), and because grazing pressure is increased (Bahamón and Cruzado 2003).

In autumn, after the stratification is broken and nutrient concentration increase in surface waters, the DCM is found at shallower depths, as already described (Estrada 1985a,b, Bahamón and Cruzado 2003 ). However its intensity, as well as that of the PNM, is smaller than in summer and spring. As a consequence, and unlike nutrients concentration seasonal dynamics, integrated chla concentration in autumn is smaller than in summer (Table 2). Assuming that there are no changes in irradiance compared to the summer months, the smaller DCM in autumn could be in part resulting from a decrease in pigment concentration per cell due to photoacclimation. In addition, the lower %O<sub>2</sub> saturation in autumn suggests that the decrease in chla in this season could also be related to a decrease in the abundance of autotrophic organisms. Finally, the sudden increase of the Catalan current intensity in autumn driven by wind and storms (Astraldi and Gasparini 1992, Font et al. 1995), could result in an increase in chla exported from the Catalan Sea. Nevertheless, additional data, like particulate organic carbon or plankton composition, are needed to explain if the observed decrease in average chla concentration corresponds to a decrease in pigment cell content, in total biomass, or to a change of populations with different pigment composition.

#### **Nutrient limitation in the Catalan sea**

The nitrate to phosphate ratio described by Redfield in deep waters (N:P=16:1, Redfield 1963) is typically used to determine if a system is N- or P-limited. Our N:P ratio in deep waters , calculated with values above detection limit, is larger than the Redfield ratio (N:P>22), as usually found in the western Mediterranean Sea (Copin-Montégut and Copin-Montégut 1983, Béthoux et al. 1998, Marty et al. 2002, Ribera d'Alcalà et al. 2003, Pujo-Pay et al. 2011). Similarly, our deep waters (>1200 m) Si:N and Si:P are similar to other deep values in the western Mediterranean Sea (Si:N=0.9 – 1.1, Ribera d'Alcalà et al. 2003; N:Si:P=22:19.5:1, Béthoux et al. 2002; N:Si:P=31.5:26.5:1, at Gibraltar Strait: Dafner et al. 2003), and the differences can be explained by the ratios being calculated at different depths. Moreover, as observed in the Eastern Mediterranean Sea (Krom et al. 1991, Kress and Herut 2001), the N:P and DIN:P scatterplots show that the relationship is non-lineal at low and high N and P concentrations (Fig. 7). This results suggest that P is the element that generally limits the production in the Catalan Sea, as found in the rest of the Mediterranean Sea (e.g.

Margalef 1963, Marty et al. 2002, Krom et al. 2010). There are several hypotheses to explain the high N:P in the Mediterranean Sea: atmospheric deposition, unbalanced river discharges, phosphate adsorption onto Saharan dust, and nitrogen fixation by cyanophyceae and marine phanerogams (Béthoux and Copin-Montégut 1986, Krom et al. 1991, Ribera d'Alcalà et al. 2003). Indeed, in the Mediterranean Sea, P has long been known to limit phytoplankton production in surface waters (Margalef 1963), as it has been confirmed on numerous experimental and descriptive studies (e.g. Thingstad et al. 1998, Moutin and Raimbault 2002).

However, our data also show that in stratified surface waters above the DCM (0–50 m), N:P and DIN:P ratios are below the Redfield ratio, while in mixed surface waters they are above it, both at open sea and continental shelf waters. This suggests a shift from N-limitation when the water column is stratified in some degree, to P-limitation when the water column is mixed (Fig. 6). On one side, using a model II regression method (our data are not normally distributed), the intercepts of both N:P and DIN:P regressions are negative and significantly different from 0 (Table 4), indicating an excess of P when N is 0. However, using Ordinary Least Squares model I (e.g. like in Kress and Herut 2001), the intercepts are positive and not significantly different from 0 ( $p>0.05$ ). An examination of paired data from 0-50 m depth shows that it is more frequent to find phosphate data below detection limit when there are still measurable concentrations (i.e. above detection limit) of nitrate (59% of the cases) or DIN (67%), than vice versa. Low N:P ratios in surface waters, as well as median phosphate concentrations above detection limits, are unexpected results compared to the observed deep waters N:P ratio and to experimental studies about nutrient limitation in the Mediterranean Sea. Low N:P ratios could be the result of an overestimation of phosphate, as found when comparing chemical and radiochemical methods (Hudson et al. 2000), and would point out the necessity of using more sensitive methods to measure phosphate in surface waters of the Catalan Sea, that provide equivalent detection limits for both nutrients.

In addition, as observed in other areas of the Mediterranean Sea (Western Mediterranean Sea: Raimbault and Coste 1990, Estrada 1999, Moutin and Raimbault 2002, Levantine Basin: e.g. Krom et al. 1991), we have found a shift from N-limitation to P-limitation within the DCM, where the nitracline is found above the phosphocline

(Pujo-Pay et al. 2011). The uncoupling of nutriclines causes N:P ratio to reach maximum values at the depth of the DCM (Fig. 10). It is suggested that the lack of phosphorus prevents the utilization of nitrogen and hence, nitrate (and silicate) is accumulated in the water column at a shallower depth than phosphate (Raimbault and Coste 1990).

5           The N:P ratio below the DCM (160 – 400 m depth) is around 23, which corresponds to the remineralization of particulate matter, mostly dead phytoplankton, faecal pellets and other detrital particles (Sarmiento and Gruber 2006), from waters immediately above. Consequently, in the Catalan Sea this sinking organic material would contain more N than P relative to the Redfield ratio. The rapid turnover rate of  
10 phosphorus in the euphotic zone (Benitez-Nelson and Buesseler 1999) could result in a larger N:P ratio of non-living sinking particles. Indeed, Mostajir et al. (1998) found higher than Redfield N:P ratios in detritic matter from the Ligurian Sea (N:P=32). In addition, some analyses of particulate matter from the NW Mediterranean Sea indicate that the N:P ratio of the whole phytoplankton community is also larger than Redfield.  
15 Copin-Montégut and Copin-Montégut (1983) found a particulate N:P ratio between 20 and 23 in the NW Mediterranean Sea, and suggested that the high N:P ratio in deep waters was the result of remineralization of microphytoplankton, which has a N:P~30 and higher than smaller organisms. Later studies in living cells found that the N:P ratios of cultured picoplanktonic cells (*Synechococcus* and *Prochlorococcus*, including one  
20 strain from the NW Mediterranean Sea) were higher than Redfield in cells grown without nutrient limitation, and up to N:P=109 under P-limitation (Bertilsson et al. 2003). Single-cell analysis of diatoms and dinoflagellates sampled directly from the Catalan Sea also indicate that the cells are N-rich (slope N:P quota=24.6 in dinoflagellates and 38.7 in diatoms, Segura-Noguera et al. submitted).

25           Redfield noted that the N:P obtained from nutrients in deep oceanic waters should be an average of plankton stoichiometry, which is able to adapt to oceanic nutrients availability (Redfield 1963). Several studies confirm the plasticity of plankton stoichiometry as a result of changes in the metabolic cellular machinery in response to environmental nutrient availability (e.g. Margalef 1998, Geider and LaRoche 2002),  
30 with a theoretical optimal N:P stoichiometry between 8.2 and 45.0 (Klausmeier et al. 2004). For example, diatoms, which dominate the phytoplankton community in the NW Mediterranean Sea in winter (Estrada 1999, Arin et al. 2013), are able to store nitrate, as

well as phosphate, inside intracellular vacuoles, allowing them to continue growing for some generations after the nutrients are depleted (Rosen and Lowe 1984, Margalef 1998). Also, diatoms could store nitrate as pigments (Rosen and Lowe 1984), and since each chlorophyll molecule contain four atoms of N, the resulting N:P ratio would be higher. A larger chlorophyll concentration in diatoms compared to cultured dinoflagellates and other autotrophic organisms in natural samples, has actually been suggested (e.g. Ríos et al. 1998). Another adaptation that would lead to higher cellular N:P ratios is the preferential synthesis of N-rich substitute lipids in response to P limitation, as found in the Adriatic Sea (Ivančić et al. 2012). Since the cellular stoichiometry is the result of the species-specific genetic stoichiometry and the adaptation to the nutrients availability, the coincidence in the Catalan Sea of the N:P ratio in particulate matter and in dissolved nutrients below the euphotic zone, both of them higher than the canonical Redfield ratio, suggests an adaptation of the phytoplankton to the low P availability of the Mediterranean Sea (Klausmeier et al. 2004), and/or an evolutive selection of high N:P stoichiometry plankton cells (Kay et al. 2007). To confirm this, more measurements of the elemental composition of plankton are needed in the NW Mediterranean Sea, including other abundant groups (e.g. Primmeciophyceae), as well as of the community composition and participation in the downward flux of particulate organic matter.

## ACKNOWLEDGEMENTS

This work was supported by project CANYONS (MAR99-1060-C03, CYCIT), CACO (REN2002-01339/MAR, MEC), and PUDEM (REN2003-06637-C02) and by an I3P predoctoral fellowship (CSIC, MEC) to M.S.-N. We thank Marta Estrada, Jordi Salat, Miquel Alcaraz and Laura Arin for providing biogeochemical data from cruises, as well as Ana Sabatés and Albert Palanques for providing cruise opportunities between 2001 and 2004 (“Canyons” I to IV in 2001, and “Caco” 1 to 4 in 2003 and 2004). We thank Mikhail Emelianov, Jordi Salat and Jordi Font (Institut de Ciències del Mar, CSIC) for useful guidance about the physical oceanography key aspects of the Catalan Sea. We also thank the UTM staff (CSIC) for their assistance during the above mentioned cruises, as well as two anonymous reviewers for helpful comments that greatly improved the original manuscript submitted.

## REFERENCES

- Arin L., Guillén J., Segura-Noguera M., Estrada M. 2013. Open sea hydrographic forcing of nutrient and phytoplankton dynamics in a Mediterranean coastal ecosystem. *Est. Coast. Shelf Sci.* 133: 116-128.
- 5 Astraldi M., Gasparini G.P. 1992. The seasonal characteristics of the circulation in the North Mediterranean basin and their relationship with the atmospheric-climatic conditions. *J. Geophys. Res.* 97: 9531-9540.
- Bahamón N., Cruzado A. 2003. Modelling nitrogen fluxes in oligotrophic environments : NW Mediterranean and NE Atlantic. *Ecol. Model.* 163: 223-244.
- 10 Ballester A., Arias E., Cruzado A., Blasco D., Camps J.M. 1967. Estudio hidrográfico de la costa catalana, de junio de 1965 a mayo de 1967. *Investigación Pesquera* 31: 621-662.
- Benitez-Nelson C.R., Buesseler K.O. 1999. Variability of inorganic and organic phosphorus turnover rates in the coastal ocean. *Nature* 398: 502-505.
- Bertilsson S., Berglund O., Karl D.M., Chisholm S.W. 2003. Elemental composition of marine *Prochlorococcus* and *Synechococcus*: Implications for the ecological stoichiometry of the sea. *Limnol. Oceanogr.* 48: 1721-1731.
- 15 Béthoux J.P., Copin-Montégut G. 1986. Biological fixation of atmospheric dinitrogen in the Mediterranean Sea. *Limnol. Oceanogr.* 31: 1353-1358.
- Béthoux J.P., Morin P., Chaumery C., Connan O., Gentili B., Ruiz-Pino D. 1998. Nutrients in the Mediterranean Sea, mass balance and statistical analysis of concentrations with respect to environmental change. *Mar. Chem.* 63: 155-169.
- 20 Béthoux J.P., Morin P., Ruiz-Pino D. 2002. Temporal trends in nutrient ratios: chemical evidence of Mediterranean ecosystem changes driven by human activity. *Deep-Sea Res. II* 49(11): 2007-2016.
- 25 Blasco D. 1971. Acumulación de nitritos en determinados niveles marinos por acción del fitopláncton. Ph.D. Thesis, Univ. Barcelona.
- Bower C., Holm-Hansen T. 1980. A salycilate-hypochlorite method for determining ammonia in seawater. *Can. J. Aquat. Sci.* 37: 794-798.
- Brzezinski M.A., 1988. Vertical distribution of ammonium in stratified oligotrophic waters. *Limnol. Oceanogr.* 33(5): 1176-1182.
- 30 Canals M., Serra J., Riba O. 1982. Toponímia de la Mar Catalano-Balear (amb un glossari de termes genèrics). *Bull. Soc. Hist. Nat. Balears* 26: 169-194.

- Canals M., Puig P., Durrieu de Madron X., Heussner S., Palanques A., Fabres J. 2006. Flushing submarine canyons. *Nature* 444: 354-357.
- Canals M., Danovaro R., Heussner S., Lykousis V., Puig P., Trincadi F., Calafat A.M., Durrieu de Madron X., Palanques A., Sànchez-Vidal A. 2009. Cascades in Mediterranean submarine grand canyons. *Oceanography* 22(1): 26-43.
- 5 Copin-Montégut C., Copin-Montégut G. 1983. Stoichiometry of carbon, nitrogen, and phosphorus in marine particulate matter. *Deep-Sea Res.* 30: 31-46.
- Cruzado A., 1985. Chemistry of Mediterranean Waters. In: Margalef R. (ed.), *Western Mediterranean*. Pergamon Press, Oxford, pp. 126-147.
- 10 Cruzado A., Velásquez Z. 1990. Nutrients and phytoplankton in the Gulf of Lions, north-western Mediterranean. *Cont. Shelf Res.* 10(9-11): 931-942.
- Cruzado A., Velásquez Z., Perez M.D., Bahamón N., Grimaldo N.S., Ridolfi F. 2002. Nutrient fluxes from the Ebre river and subsequent across-shelf dispersion. *Cont. Shelf Res.* 22: 349-360.
- 15 Dafner E.V., Boscolo R., Bryden H.L. 2003. The N:Si:P molar ratio in the Strait of Gibraltar. *Geophysical Research Letters* 30(10): 1506.
- Duarte C.M., Agustí S., Kennedy H., Vaqué D. 1999. The Mediterranean climate as a template for Mediterranean marine ecosystems: the example of the northeast Spanish littoral. *Prog. Oceanogr.* 44: 245-270.
- 20 Estrada M. 1985a. Deep phytoplankton and chlorophyll maxima in the Western Mediterranean. In: Moraitou-Apostolopoulou, M., Kiorstis, V. (eds.), *Mediterranean Marine Ecosystems*. Plenum Press, New York, pp. 247-277.
- Estrada M. 1985b. Primary production at the deep chlorophyll maximum in the Western Mediterranean. In: P.E. Gibbs (ed.), *Proc. 19th European Marine Biology Symp*, Cambridge University Press, pp. 109-121.
- 25 Estrada M. 1996. Primary production in the Northwestern Mediterranean. *Sci. Mar.* 60 (Supl. 2): 55-64.
- Estrada M. 1999. Hidrodinàmica i fitoplàncton en el Mar Català. *Memorias de la Real Academia de Ciencias y Artes de Barcelona* 961, vol. LVIII (6), 189-247.
- 30 Estrada M., Margalef R. 1988. Supply of nutrients to the Mediterranean photic zone along a persistent front. *Oceanol. Acta* 9 (Spec. Issue): 133-142.

- Estrada M., Marrasé C., Latasa M., Berdalet E., Delgado M., Riera T. 1993. Variability of deep chlorophyll maximum characteristics in the northwestern Mediterranean. *Mar. Ecol. Prog. Ser.* 92: 289-300.
- 5 Estrada M., Varela R.A., Salat J., Cruzado A., Arias E. 1999. Spatio-temporal variability of the winter phytoplankton distribution across the Catalan and north Balearic fronts (NW Mediterranean). *J. Plankton Res.* 21: 1-20.
- Falster D S, Warton D I, Wright I J. SMATR: Standardised major axis tests and routines. 2006. Available: <http://www.bio.mq.edu.au/ecology/SMATR/>
- 10 Fichaut M., Balopoulos E., Baudet L., Dooley H., García-Fernández M.-J., Iona A., Jourdan D., Maillard C. 1997. A common protocol to assemble a coherent database from distributed heterogeneous data sets: the MEDATLAS database experience. MAST Workshop on Project Data Management, 11–13 June, Ispra, Italy.
- Flo E., Camp J. 2005. Programa de Vigilància i Control de la Qualitat Ambiental de les Aigües Litorals a Catalunya 8, Anual any 2.
- 15 Font J., Salat J., Tintoré J. 1988. Permanent features of the circulation in the Catalan Sea. *Oceanol Acta* 9 (Spec. Issue): 51-57.
- Font J., García-Ladona E., Gorriz E.G. 1995. The seasonality of mesoscale motion in the Northern Current of the western Mediterranean: several years of evidence. *Oceanol. Acta* 18(2): 207-219.
- 20 Geider R., La Roche J. 2002. Redfield revisited: variability of C : N : P in marine macroalgae and its biochemical basis. *Eur. J. Phycol.* 37: 1-17.
- Granata T.C., Estrada M., Zika U., Merry C. 2004. Evidence for enhanced primary production resulting from relative vorticity induced upwelling in the Catalan current. *Sci. Mar.* 68 (Suppl.1): 113-119.
- 25 Guadayol Ò., Peters F. 2006. Analysis of wind events in a coastal area: a tool for assessing turbulence variability for studies on plankton. *Sci. Mar.* 70: 9-20.
- Guadayol Ò., Marrasé C., Peters F., Berdalet E., Roldán C., Sabata A. 2009. Responses of coastal osmotrophic planktonic communities to simulated events of turbulence and nutrient load throughout a year. *J. Plankton Res.* 31(6): 583-600.
- 30 Hammer Ø., Harper D.A.T., Ryan P.D. 2001. Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontol. Electron.* 4: 9.

- Hansen H.P., Koroleff F. 1999. Determination of nutrients. In: Grasshoff K., Kremling K., Ehrhardt M. (eds.), *Methods of Seawater Analysis*. Wiley-VCH, Weinheim, New York, pp. 191-228.
- Holmes R.M., Aminot A., K  rouel R., Hooker B.A., Peterson B.J. 1999. A simple and precise method for measuring ammonium in marine and freshwater ecosystems. *Can. J. Fish. Aquat. Sci.* 56: 1801-1808.
- Hudson J.J., Taylor W.D., Schindler D.W. 2000. Phosphate concentration in lakes. *Nature* 406: 54-56.
- Isla E., Scharek R., Latasa M. 2015. Zooplankton diel vertical migration and contribution to deep active carbon flux in the NW Mediterranean. *J. Mar. Sys.* 143: 86-97.
- Ivan  i   I., Godrijan J., Pfannkuchen M., Mari   D., Gas  parovi   B., Djakivac T, Najdeka M. 2012. Survival mechanisms of phytoplankton in conditions of stratification-induced deprivation of orthophosphate: Northern Adriatic case study. *Limnol. Oceanogr.* 57(6): 1721-1731.
- Johnson M., Sanders R., Avgoustidi V., Lucas M., Brown L., Hansell D., Moore M., Gibb S., Liss P., Jickells T. 2007. Ammonium accumulation during a silicate-limited diatom bloom indicates the potential for ammonia emission events. *Mar. Chem.* 106: 63-75.
- Karl D.M., Knauer G.A. 1984. Vertical distribution, transport, and exchange of carbon in the Northeast Pacific Ocean: evidence for multiple zones of biological activity. *Deep-Sea Res.* 31: 221-243.
- Karl D.M., Christian J.R., Dore J.E., Hebel D.V., Letelier R.M., Tupas L.M., Winn C.D. 1996. Seasonal and interannual variability in primary production and particle flux at Station ALOHA. *Deep-Sea Res. II* 43(2-3): 539-568.
- Kay A.D., Ashton I.W., Gorokhova E., Kerkhoff A.J., Liess A., Litchman E. 2005. Toward a stoichiometric framework for evolutionary biology. *Oikos* 109: 6-17.
- Klausmeier C.A., Litchman E., Daufresne T., Levin, S.A. 2004. Optimal nitrogen-to-phosphorus stoichiometry of phytoplankton. *Nature* 429: 171-174.
- Kress N., Herut B. 2001. Spatial and seasonal evolution of dissolved oxygen and nutrients in the Southern Levantine Basin (Eastern Mediterranean Sea): chemical characteri-

zation of the water masses and inferences on the N : P ratios. *Deep-Sea Res. I* 48: 2347-2372.

Krom M.D., Kress N., Brenner S., Gordon L.I., 1991. Phosphorus limitation of primary productivity in the Eastern Mediterranean Sea. *Limnol. Oceanogr.* 36(6): 424-432.

5 Krom M.D., Emeis K.-C., Van Cappellen P., 2010. Why is the Eastern Mediterranean phosphorus limited? *Prog. Oceanogr.* 85: 236-244.

Longhurst A., 1998. *Ecological Geography of the Sea*. Academic Press, San Diego.

Longhurst A. R., Harrison W.G. 1988. Vertical nitrogen flux from the oceanic photic zone by diel migrant zooplankton and nekton. *Deep Sea Res. A.* 35 (6): 881-889.

10 Margalef R., 1985. *Western Mediterranean*. Pergamon Press, Oxford.

Margalef R., 1963. El ecosistema pelagico de un area costera del Mediterraneo Occidental. *Memorias de la Real Academia de Ciencias y Artes de Barcelona* 699, vol. XXXV (1): 1-48.

15 Margalef R. 1998. Elements limitants, explotabilitat i diversitat. Homenatge a Bolós i al fósfor. *Acta Bot. Barc.* 45 (Homenatge a Oriol de Bolós): 633-643.

Marty J.-C., Chiavérini J., Pizay M.-D., Avril, B. 2002. Seasonal and interannual dynamics of nutrients and phytoplankton pigments in the western Mediterranean Sea at the DYFAMED time-series station (1991–1999). *Deep-Sea Res. II* 49: 1965-1985.

20 Masó M., La Violette P.E., Tintoré J. 1990. Coastal flow modification by submarine canyons along the NE Spanish coast. *Sci. Mar.* 54(4): 343-348.

MEDOC Group. 1970. Observation of formation of deep water in the Mediterranean Sea. *Nature* 277 : 1037-1040.

Millero, F.J. 1996. *Chemical Oceanography*. CRC Press, Boca Raton, Florida.

Millot C. 1987. Circulation in the western Mediterranean sea. *Oceanol. Acta* 10: 149.

25 Minas H.J., Bonin M.C. 1988. Oxygénation physique et biologique de la Méditerranée nord-occidentale en hiver et au printemps. *Oceanol. Acta* 9 (Spec. Issue): 123-132.

Mostajir B., Fagerbakke K.M., Heldal M., Thingstad T.F., Rassoulzadegan F. 1998. Elemental composition of individual pico- and nano-sized marine detrital particles in the northwestern Mediterranean Sea. *Oceanol. Acta* 21(4):589-596.

30 Moutin T., Raimbault P., 2002. Primary production, carbon export and nutrients availability in western and Eastern Mediterranean Sea in early summer 1996 (MINOS cruise). *J. Mar. Sys.* 33: 273-288.

- Olson R.J., 1981. Differential photoinhibition of marine nitrifying bacteria-A possible mechanism for the formation of the primary nitrite maximum. *J. Mar. Res.* 39: 227-238.
- Owens W.B., Millard Jr.R.C., 1985. A new algorithm for CTD oxygen calibration. *J. Phys. Oceanogr.* 15: 621-631.
- 5 Palanques A., Martín J., Puig P., Guillén J., Company J.B., Sardà F. 2006. Evidence of sediment gravity flows induced by trawling in the Palamós (Fonera) submarine canyon (northwestern Mediterranean). *Deep-Sea Res. I* 53: 201-214.
- Pascual A., Buongiorno Nardelli B., Larnicol G., Emelianov M., Gomis D. 2002. A case of an intense anticyclonic eddy in the Balearic Sea (western Mediterranean). *J. Geophys. Res.* 107(C11): 3183.
- 10 Pujó-Pay M., Conan P., Oriol L., Cornet-Barthaux V., Falco C., Ghiglione J.-F., Goyet C., Moutin T., Prieur L. 2011. Integrated survey of elemental stoichiometry (C, N, P) from the western to eastern Mediterranean Sea. *Biogeosciences* 8: 883-899.
- Raimbault P., Coste B. 1990. Very high values of the nitrate/phosphate ratio (>30) in the subsurface layers of the Western Mediterranean Sea. *Rapp. CIESMM* 32 (1): 44.
- 15 Redfield A.C., Ketchum B.H., Richards F.A., 1963. The influence of organisms on the composition of sea-water. In: Hill, M.N. (ed.), *The Sea*, Vol. 2. The Composition of Sea-water. Wiley, New York.
- Ribera d'Alcalà M., Civitarese G., Conversano F., Lavezza R. 2003. Nutrient ratios and fluxes hint at overlooked processes in the mediterranean sea. *J. Geophys. Res.* 108: 8106.
- 20 Riley J.P. 1971. Micronutrient elements. In: Riley, J.P., Chester, R. (eds.), *Introduction to Marine Chemistry*. Academic Press, London.
- Ríos A.G., Fraga F. Pérez F.F., Figueiras F.G. 1998. Chemical composition of phytoplankton and particulate organic matter in the Ría de Vigo (NW Spain). *Sci. Mar.* 62(3): 257-271.
- 25 Rosen, B., Lowe, R. 1984. Physiological and ultrastructural responses of *Cyclotella meneghiniana* (*Bacillariophyta*) to light-intensity and nutrient limitation. *J. Phycol.* 20: 173-183.
- 30 Saino T., Otake H., Wada E., Hattori A. 1983. Subsurface ammonium maximum in the northern Pacific and the Bering Sea in summer. *Deep-Sea Res.* 30: 1157-1171.

- Salat J. 1983. Formation d'eau profonde près de la côte catalane au sud du 42°N. Croisière "Caron 82": premiers resultats. Rapp. Comm. Int. Mer Médit. 28(2): 73-77.
- Salat J. 1995. The interaction between the Catalan and Balearic currents in the southern Catalan Sea. Oceanol. Acta 18(2): 227-234.
- 5 Salat J. 1996. Review of hydrographic environmental factors that may influence anchovy habitats in northwestern Mediterranean. Sci. Mar. 60(Suppl. 2): 21-32.
- Salat J., Cruzado A. 1981. Masses d'eau dans la Méditerranée Occidentale: Mer Catalane et eaux adjacentes. Rapp. Comm Int. Mer Médit. 27(6): 201-209.
- Salat J., Font J. 1987. Water mass structure near and offshore the Catalan coast during the winters of 1982 and 1983. Ann. Geophys. B- Terr. Planet. Phys. 5: 49-54.
- 10 Salat J., Garcia M., Cruzado A., Palanques A., Arin L., Gomis D., Guillen J., de Leon A., Puigdefabregas J., Sospedra J., Velásquez Z. 2002. Seasonal changes of water mass structure and shelf slope exchanges at the Ebre shelf (NW Mediterranean). Cont. Shelf Res. 22: 327-348.
- 15 Sarmiento J.L., Gruber N. 2006. Ocean Biogeochemical Dynamics. Princeton Univ. Press.
- Send U., Font J., Krahnemann G., Millot C., Rhein M., Tintoré J. 1999. Recent advances in observing the physical oceanography of the western Mediterranean Sea. Prog. Oceanogr. 44: 37-64.
- 20 Segura-Noguera M., Cruzado A., Blasco D. 2011. Nutrient preservation, analysis precision and quality control of an oceanographic database (inorganic nutrients, dissolved oxygen and chlorophyll *a*) from the NW Mediterranean Sea. Sci. Mar. 75(2): 321-339.
- Sournia A. 1973. La production primaire planctonique en Méditerranée : Essai de mise à jour. Bulletin de l'Étude en Commun de la Méditerranée 5: 1-28.
- 25 Thingstad T.F., Zweifel U.L., Rassoulzadegan F. 1998. P limitation of heterotrophic bacteria and phytoplankton in the northwest Mediterranean. Limnol. Oceanogr. 43: 88-94.
- Winkler L.W., 1888. Die Bestimmung des in Wasser gelosten Sauerstoffes. Berichte der Deutschen Chemischen Gesellschaft 21: 2842-2855.
- Yentsch C.S., Menzel D.W. 1963. A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence. Deep-Sea Research 10: 221-231.
- 30

**TABLE 1** Some characteristics of the oceanographic variables of the dataset used in this study: range of years, number of cruises, profiles and data from continental shelf (stations with bottom depth  $\leq 200$  m, CS) and from the open sea (stations with a bottom depth  $> 200$  m, OS). Only data flagged “0” during the quality control are included in the Table. In addition, only data above detection limits are included in the case of the dissolved inorganic nitrogen (DIN), apparent oxygen utilization (AOU), oxygen saturation (%O<sub>2</sub> sat) and all the ratios.

Variable	Range of years	Num of cruises	Num of profiles			Num of data		
			CS	OS	Total	CS	OS	Total
Phosphate	1957 – 2004	88	490	731	1221	2994	7478	10472
Nitrate	1979 – 2004	38	622	1029	1652	4117	11262	15379
Nitrite	1976 – 2004	67	538	891	1430	3612	9841	13453
Ammonium	1982 – 2004	26	247	470	717	1196	3180	4376
DIN	1982 – 2004	21	139	369	508	416	1645	2061
Silicate	1970 – 2004	82	620	1027	1648	4235	11794	16029
Oxygen	1910 – 2000	89	309	679	989	2119	8174	10293
AOU, % O <sub>2</sub> sat	1910 – 2000	74	1996	649	948	298	7543	9539
Chlorophyll <i>a</i>	1976 – 2004	64	468	725	1194	2153	6035	8188
N:P	1982 – 2003	23	294	591	885	1609	4515	6124
DIN:P	1982 – 2003	20	121	294	414	329	1022	1351
Si:P	1982 – 2003	22	290	581	871	1684	4824	6508
Si:N	1979 – 2003	30	407	877	1285	2561	7921	10482
AOU:P	1970 – 2000	17	154	391	545	762	3191	3953
AOU:N	1979 – 2000	24	246	583	830	1029	5201	6515
AOU:Si	1982 – 2000	21	238	571	810	1440	5885	7325

**TABLE 2** Physico-chemical seasonal properties of surface water (average 100 m  $\pm$  standard deviation, number of profiles between parenthesis) in the Catalan Sea (NW Mediterranean Sea). Units are  $\mu\text{M}$  for nutrients, and dissolved oxygen ( $\text{O}_2$ ), and  $\mu\text{g L}^{-1}$  for chlorophyll *a*. Significant differences ( $p \leq 0.05$ ) are indicated with “<” or “>”, while the absence of statistically significant differences is indicated with “=”.

5

Variable	Winter	Spring	Summer	Autumn
Temperature	14.0 $\pm$ 0.6 (41)	< 15.7 $\pm$ 0.9 (183)	< 16.9 $\pm$ 0.6 (52)	> 16.6 $\pm$ 1.2 (66) >
Salinity	38.4 $\pm$ 0.1 (42)	< 38.3 $\pm$ 0.2 (183)	= 38.3 $\pm$ 0.2 (53)	< 38.5 $\pm$ 0.1 (66) =
Density ( $\sigma\text{-T}$ )	28.9 $\pm$ 0.1 (41)	> 28.4 $\pm$ 0.2 (183)	> 28.0 $\pm$ 0.3 (52)	> 28.2 $\pm$ 0.3 (66) <
Phosphate	0.09 $\pm$ 0.05 (37)	> 0.06 $\pm$ 0.04 (140)	= 0.05 $\pm$ 0.02 (42)	< 0.11 $\pm$ 0.08 (72) =
Nitrate	1.33 $\pm$ 0.48 (40)	> 1.03 $\pm$ 0.48 (210)	> 0.74 $\pm$ 0.33 (48)	< 0.99 $\pm$ 0.56 (75) <
Nitrite	0.15 $\pm$ 0.07 (35)	> 0.10 $\pm$ 0.05 (194)	> 0.05 $\pm$ 0.02 (52)	< 0.09 $\pm$ 0.04 (58) <
Ammonium	0.75 (1)	0.31 $\pm$ 0.20 (14)	= 0.15 $\pm$ 0.13 (22)	< 0.33 $\pm$ 0.17 (25)
Silicate	1.60 $\pm$ 0.72 (41)	= 1.51 $\pm$ 0.59 (212)	> 0.99 $\pm$ 0.50 (51)	< 1.28 $\pm$ 0.37 (78) <
Chlorophyll <i>a</i>	0.54 $\pm$ 0.17 (42)	> 0.37 $\pm$ 0.12 (112)	= 0.36 $\pm$ 0.13 (46)	> 0.26 $\pm$ 0.08 (64) <
$\text{O}_2$	258.6 $\pm$ 9.0 (16)	> 249.5 $\pm$ 10.8 (134)	= 247.9 $\pm$ 11.4 (23)	> 240.0 $\pm$ 6.7 (40) <
% $\text{O}_2$ sat	100.8 $\pm$ 3.4 (16)	= 100.9 $\pm$ 5.2 (91)	= 102.9 $\pm$ 5.1 (21)	> 98.4 $\pm$ 3.6 (38) <

**TABLE 3** Seasonal changes in the nitrate (N), dissolved inorganic nitrogen (DIN) and silicate (Si) vs. phosphate (P), and silicate vs. nitrate ratios in surface water (0 to 50 m depth). Ratios are shown as median±median absolute deviation, number of data between parenthesis. Significant differences ( $p \leq 0.05$ ) are indicated with “<” or “>”, while the absence of statistically significant differences is indicated with “=”. “n.d.”: no data.

5


Ratio	Winter		Spring		Summer		Autumn	
<b>Open Sea</b>								
N:P	17.3±12.6 (515)	>	2.5±1.3 (453)	=	3.0±1.4 (66)	=	2.8±1.3 (198)	<
DIN:P	n.d.		5.9±2.8 (107)	<	7.5±3.4 (34)	>	4.1±1.9 (82)	
Si:P	23.5±15.2 (561)	>	15.8±9.3 (746)	=	17.2±8.1 (109)	>	8.1±7.1 (189)	<
Si:N	1.5±0.6 (644)	<	4.7±2.8 (1400)	=	6.0±2.8 (146)	=	2.6±1.4 (279)	>
<b>Continental Shelf</b>								
N:P	25.8±13.4 (388)	>	2.5±1.3 (248)	=	2.5±1.0 (108)	<	3.7±1.4 (191)	<
DIN:P	22.1±10.4 (68)	>	6.1±3.09 (40)	<	10.7±5.5 (34)	=	11.0±6.6 (47)	<
Si:P	40.3±19.9 (406)	>	17.7±10.1 (302)	=	9.9±4.4 (118)	<	16.3±4.6 (216)	<
Si:N	1.4±0.4 (447)	<	5.4±2.6 (758)	=	4.5±2.2 (132)	=	5.0±2.5 (218)	>

**TABLE 4** Results of the standard major axis regressions of nitrate (N), dissolved inorganic nitrogen (DIN) and apparent oxygen utilization (AOU) vs. phosphate (P), and AOU vs. nitrate in the Catalan Sea. Shown are the y-intercept and the slope of the regression equations, the interval of confidence (Low CI and Upp CI), the coefficient of determination ( $r^2$ ), and the number of data points included (n). All intercepts, except AOU:P, and slopes are significantly different from 0 ( $p \leq 0.001$ ).


Ratio	intercept	Low CI	Upp CI	slope	Low CI	Upp CI	$r^2$	n
N:P	-0.71	-0.78	-0.64	22.62	22.29	22.94	0.646	6560
DIN:P	-0.77	-0.92	-0.62	22.23	21.53	22.94	0.578	1605
AOU:P	-0.18	-2.19	-1.82	150.2	144.0	156.6	0.353	1420
AOU:N	-12.5	-14.4	-10.6	8.89	8.64	9.15	0.603	1864

## FIGURES


**Figure 1** Bathymetric map of the Catalan Sea (2005 Catalano-Balearic Sea. Bathymetric Chart and Toponyms, [www.icm.csic.es/geo/gma/MCB/MCBexplora\\_cat.htm](http://www.icm.csic.es/geo/gma/MCB/MCBexplora_cat.htm)). The pathway of the Liguro-Provençal (or Northern Current) and Balearic currents is shown (arrows), as well as an approximate position of the Catalan and Balearic fronts (dashed and dotted lines, respectively), as in Pascual et al. (2002). The inlet map shows the location of the Catalan Sea in the Mediterranean Sea (created using [www.aquarius.geomar.de/omc](http://www.aquarius.geomar.de/omc)).




**Figure 2** Seasonal depth concentration profiles of (A) density anomaly, (B) Salinity and (C) Temperature in open sea stations (bottom depth > 200 m) of the Catalan Sea. The first 200 m are enlarged in the small plots.




**Figure 3** Seasonal depth concentration profiles of (A) phosphate, (B) nitrate, and (C) silicate in open sea stations (bottom depth > 200 m) of the Catalan Sea. The first 200 m are enlarged in the small plots.




**Figure 4** Seasonal depth concentration profiles of (A) ammonium, (B) dissolved oxygen, (C) nitrite, and (D) chlorophyll *a* in open sea stations (bottom depth > 200 m) of the Catalan Sea. The first 200 m are enlarged in the small plots.




**Figure 5** Surface water (0 – 50 m) dissolved oxygen concentration (green dots: open sea, orange dots: continental shelf (orange) , and dissolved oxygen saturation (red, calculated from *in situ* S and T), as a function of temperature.



**Figure 6** Seasonal stoichiometry depth profiles of N:P (A), dissolved inorganic nitrogen (DIN= $\text{NO}_{2+3} + \text{NH}_4$ ):P (B), Si:P (C), and Si:N (D) in open sea stations (bottom depth > 200 m) of the Catalan Sea. The first 200 m are enlarged in the small plots.




**Figure 7** Scatter plots of nitrate vs. phosphate (N:P), dissolved inorganic nitrogen vs. phosphate (DIN:P), silicate vs. phosphate (Si:P), silicate vs. nitrate (Si:N), apparent oxygen utilization vs. phosphate (AOU:P) and AOU vs. nitrate (AOU:N). Plotted are quality controlled data flagged with “0”.




5


**Figure 8** Depth profiles of density anomaly, phosphate, nitrate, silicate, dissolved oxygen and chlorophyll *a* of two stations of Hivern2000 cruise, at the WMDW formation area. The arrows in the inlet map point towards the position of stations 25/26 (41.80 N 3.92 E) and 27/28 (41.73 N 4.11 E). As a reference, the median winter depth profile (Median) of the same variables, and the lower (LW) and upper (UR) ranges of concentration where data are flagged “0” in winter, are also plotted.



**Figure 9** Scatter plots of nitrate and silicate vs. depth, including all the data collected. Blue dots are from cruises in winter, and orange dots are from cruises between spring and autumn. High values are observed in mesopelagic and deep waters, corresponding to Hivern 1999 cruise, a year of intense deep shelf water cascading in the “Cap de Creus” Canyon, as well as inside of the “La Fonera” Canyon (south of “Cap the Creus” Canyon) in 2001.



**Figure 10** Depth profile of N:P for the Catalan Sea. Data for winter mixing are shown with blue circles, and data for stratified water columns (spring-autumn) are shown with orange circles.



### Additional Information

**Figure A1** Position of oceanographic stations in the Catalan Sea with phosphate, nitrate, nitrite, ammonium, silicate, dissolved oxygen and chlorophyll *a* data. Isobaths correspond to 100, 200, 400, 1000, 2000 and 2500 m depth.




TABLE A1. Detailed list of the cruises used in this study, including the year, maximum depth sampled (z max), number of profiles (n), biogeochemical variables available (shaded), and the source of the data.

Cruise	year	z max	n	PO <sub>4</sub>	NO <sub>2+3</sub>	NO <sub>2</sub>	NH <sub>4</sub>	Si(OH) <sub>4</sub>	O <sub>2</sub>	Chla	source
ARCHIVES	1910	1000	4								1
MUSEUM	1930	1500	1								1
MUSEUM	1951	796	1								1
NODC-1468-1	1951	1775	1								1
57051411	1957	1000	1								1
NODC-322-3-ATLANT	1961	2297	22								1
s.h 11-61	1961	1936	36								1
NODC-0210-3-ATLANT	1962	2540	2								1
S.H 05-62	1962	2212	35								1
S.H 10-63	1963	2240	3								1
S.H 03-64	1964	289	3								1
S.H 10-64	1964	2141	17								1
S.H 01-65	1965	591	15								1
S.H 04-65	1965	1850	14								1
S.H 03-66	1966	1940	4								1
S.H 04-67	1967	1182	4								1
MEDOC 1969	1969	2480	18								1
NODC-1355-5	1969	2429	13								1
MEDOC 1970	1970	2504	60								1
3131	1971	814	5								1
3132	1971	75	3								1
S.H 02-71	1971	495	1								1
S.H 04-71	1971	485	1								1
S.H 07-71	1971	489	1								1
S.H 09-71	1971	535	1								1
3767	1972	514	9								1
S.H	1972	590	2								1
7224	1973	500	14								1
CONTA 75	1975	1800	8								1
MAHON-III	1975	24	4								1
OBM-II	1975	1000	1								1
OBM-III	1975	1000	9								1
13 CRUISE	1976	2451	10								1
CAMBRILS-1	1976	70	13								1
CAMBRILS-2	1976	70	16								1
CAMBRILS-3	1976	70	17								1
CAMBRILS-4	1976	70	17								1
MAHON-IV	1976	35	4								1
NDCA-00039	1976	1400	18								1
OBM-IV	1976	600	11								1
OBM-VI	1976	1000	11								1
ROMB-1	1978	40	4								1
ROMB-2	1978	40	4								1
ROMB-3	1978	40	4								1
ROMB-4	1979	40	4								1
TANIT	1979	2100	16								2
MAHON-80	1980	27	49								1
MAHON-81	1981	27	51								1

Cruise	year	z max	n	PO <sub>4</sub>	NO <sub>2+3</sub>	NO <sub>2</sub>	NH <sub>4</sub>	Si(OH) <sub>4</sub>	O <sub>2</sub>	Chla	source
PEP 82	1982	1400	54								3
ROMM-82	1982	26	5								1
PEP 83	1983	2236	94								3
ROMM-83	1983	25	13								1
PEP 84	1984	2318	52								3
PEMTA-1	1984	70	22								1
PEMTA-2	1984	51	4								1
BALEAR-I	1985	767	15								1
BALEAR-II	1985	715	7								1
BALEAR-III	1985	628	20								1
CARON85	1985	1606	20								2
FRONTS-3-85	1985	1966	17								4
FRONTS-6-85	1985	2200	23								4
CARBAL-III	1986	48	15								1
PELAGOLION-I	1986	1500	4								2
PELAGOLION-II	1986	2235	5								2
PEP 86	1986	2331	39								4
FRONTS-11-86	1986	2150	15								4
PEP 87	1987	1983	20								4
BALEAR-IV	1987	875	23								1
BALEAR-V	1987	960	29								1
BALEAR-VI	1987	965	30								1
BAHIAS-88	1988	30	22								1
BALEAR-VII	1988	960	23								1
BALEAR-VIII	1988	941	27								1
DISCOVERY 88	1988	1354	2								2
PELAGOLION-III	1988	1500	3								2
FRONTS 89	1989	2145	28								5
BAHIAS 89	1989	30	22								1
BAHIAS-90	1990	30	21								1
CYBELE	1990	1700	33								2
RHODIBER	1990	1764	6								2
FRONTS 90	1990	2332	30								5
BAHIAS-91	1991	30	49								1
IBIZA-791	1991	500	5								1
TYRO	1991	2284	7								2
FRONTS 91	1991	2110	37								5
FRONTS 92	1992	2174	27								6
BAHIAS-92	1992	30	7								1
FORMENTERA	1993	3	2								1
VARIMED 93 I	1993	2214	60								6
VARIMED 93 II	1993	2048	21								6
VARIMED 93 III	1993	1860	41								6
RABDAL-94	1994	200	31								1
EUROMODEL-JUL95	1995	2028	5								1
RABDAL-95	1995	200	36								1
MESO 95	1995	2208	181								2
VARIMED 95	1995	2120	32								7
CNL APR96	1996	200	15								1
CNL JUL96	1996	200	15								1
CNL MAR96	1996	200	15								1
CNL MAY96	1996	200	15								1
CNL1 JUN96	1996	200	15								1

Cruise	year	z max	n	PO <sub>4</sub>	NO <sub>2+3</sub>	NO <sub>2</sub>	NH <sub>4</sub>	Si(OH) <sub>4</sub>	O <sub>2</sub>	Chla	source
FANS 1	1996	1152	54								2
FANS 1	1996	1152	54								2
FANS 2	1997	1303	107								2
FANS 3	1997	1200	110								2
CNL APR97	1997	200	15								1
CNL AUG97	1997	200	15								1
CNL JAN97	1997	200	14								1
CNL JUN97	1997	200	8								1
CNL MAY97	1997	200	11								1
CNL OCT97	1997	200	15								1
CNL JAN98	1998	200	12								1
CNL JUN98	1998	200	36								1
HIVERN 99	1999	2389	47								7
HIVERN 2000	2000	2367	50								7
CANYONS I	2001	1954	32								8
CANYONS II	2001	1922	44								8
CANYONS III	2001	1955	24								8
CANYONS IV	2001	1081	14								8
CACO 1	2003	1331	67								8
CACO 2	2003	1328	65								8
CACO 3	2004	1326	66								8
CACO 4	2004	1322	66								8

Source of the data:

1. data extracted from MEDAR and MEDATLAS databases between 0 – 4.5 E, 38.7 – 42.5 N (Mater Group. 2001. MTPII-MATER 1996–1999 (Mass transfer and ecosystem response) database. IFREMER Edition, Medar Group. 2002. MEDATLAS/2002 database. Mediterranean and Black Sea database of temperature salinity and bio-chemical parameters. Climatological Atlas. IFREMER Edition).
2. Dr. A. Cruzado, Center for Advances Studies of Blanes (CEAB-CSIC), Catalonia.
3. Grup PEPS. 1986. Datos oceanográficos básicos de las campañas PEP-82, PEP-83 y PEP-84 en el mar Catalán. Datos Informativos, Institute of Marine Ciencias (ICM-CSIC), 19: 1-100.
4. Masó M. and Grup PEPS. 1988. Datos oceanográficos básicos de las campañas Fronts-3-85, Fronts-6-85, PEP-86, Fronts-11-86 y PEP-87 en el mar Catalán. Datos informativos, Institute of Marine Ciencias (ICM-CSIC), 24: 1-83.
5. Varela R. and Group FRONTS. 1991. Datos Oceanográficos Básicos de las Campañas FRONTS 1989, FRONTS 1990 y FRONTS 1991 en el Mar Catalán. Datos informativos, Institute of Marine Ciencias (ICM-CSIC), 26: 1-71.

6. Masó M. and Grup Varimed. 1995. Datos Oceanográficos Básicos de las Campañas «FRONTS 1992» (octubre – noviembre 1992) y «VARIABILIDAD DE MESOESCALA EN EL MEDITERRANEO OCCIDENTAL» (junio 1993). Datos informativos, Institute of Marine Ciencias (ICM-CSIC), 27: 1-117.
- 5 7. Dr. M. Estrada, Institute of Marine Sciences (ICM-CSIC), Barcelona, Catalonia.
8. This study.