

This is a repository copy of *Integral polynomials with small discriminants and resultants*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/99522/>

Version: Published Version

---

**Article:**

Beresnevich, Victor [orcid.org/0000-0002-1811-9697](https://orcid.org/0000-0002-1811-9697), Bernik, Vasili and Goetze, Friedrich (2016) Integral polynomials with small discriminants and resultants. *Advances in Mathematics*. pp. 393-412. ISSN: 0001-8708

<https://doi.org/10.1016/j.aim.2016.04.022>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**


If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim



# Integral polynomials with small discriminants and resultants <sup>☆</sup>

Victor Beresnevich <sup>a,\*</sup>, Vasili Bernik <sup>b</sup>, Friedrich Götze <sup>c</sup><sup>a</sup> University of York, Heslington, York, YO10 5DD, England, United Kingdom<sup>b</sup> Institute of Mathematics, Surganova 11, Minsk, 220072, Belarus<sup>c</sup> University of Bielefeld, 33501, Bielefeld, Germany

## ARTICLE INFO

*Article history:*

Received 8 February 2015

Received in revised form 18

February 2016

Accepted 21 April 2016

Available online xxxx

Communicated by Kenneth Falconer

*MSC:*

11J83

11J13

11K60

11K55

*Keywords:*Counting discriminants and  
resultants of polynomials

Algebraic numbers

Metric theory of Diophantine  
approximation

Polynomial root separation

## ABSTRACT

Let  $n \in \mathbb{N}$  be fixed,  $Q > 1$  be a real parameter and  $\mathcal{P}_n(Q)$  denote the set of polynomials over  $\mathbb{Z}$  of degree  $n$  and height at most  $Q$ . In this paper we investigate the following counting problems regarding polynomials with small discriminant  $D(P)$  and pairs of polynomials with small resultant  $R(P_1, P_2)$ :

- (i) given  $0 \leq v \leq n-1$  and a sufficiently large  $Q$ , estimate the number of polynomials  $P \in \mathcal{P}_n(Q)$  such that

$$0 < |D(P)| \leq Q^{2n-2-2v};$$

- (ii) given  $0 \leq w \leq n$  and a sufficiently large  $Q$ , estimate the number of pairs of polynomials  $P_1, P_2 \in \mathcal{P}_n(Q)$  such that

$$0 < |R(P_1, P_2)| \leq Q^{2n-2w}.$$

Our main results provide lower bounds within the context of the above problems. We believe that these bounds are best possible as they correspond to the solutions of naturally arising linear optimisation problems. Using a counting result for the number of rational points near planar curves due to

<sup>☆</sup> This work was supported by the Engineering and Physical Sciences Research Council (EPSRC) [Programme Grant EP/J018260/1]; and Collaborative Research Centre at Bielefeld University [SFB 701].

\* Corresponding author.

E-mail addresses: victor.beresnevich@york.ac.uk, vb8@york.ac.uk (V. Beresnevich), bernik@im.bas-net.by, bernik.vasili@mail.ru (V. Bernik), goetze@math.uni-bielefeld.de (F. Götze).

R. C. Vaughan and S. Velani we also obtain the complementary optimal upper bound regarding the discriminants of quadratic polynomials.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Throughout this paper  $n$  will denote a positive integer. In what follows, given a polynomial  $P = a_n x^n + \cdots + a_0 \in \mathbb{Z}[x]$  of degree  $n$ , let

$$H(P) := \max_{0 \leq i \leq n} |a_i|$$

denote the standard (naive) height of  $P$  and, given a real parameter  $Q > 1$ , let

$$\mathcal{P}_n(Q) = \{P \in \mathbb{Z}[x] : \deg P = n, H(P) \leq Q\} \quad (1)$$

denote the set of integral polynomials  $P$  of degree  $n$  and height  $H(P) \leq Q$ . Throughout,  $D(P)$  will stand for the discriminant of a polynomial  $P$  and  $R(P_1, P_2)$  will stand for the resultant of polynomials  $P_1, P_2$ . The formal definitions and basic properties of these important number theoretic characteristics will be recalled below.

In this paper we investigate the following counting problems regarding the discriminant and resultant of polynomials in  $\mathcal{P}_n(Q)$ .

**Problem 1.** Let  $n \geq 2$  be an integer. Given  $0 \leq v \leq n - 1$  and a sufficiently large  $Q$ , estimate the number of polynomials  $P \in \mathcal{P}_n(Q)$  such that

$$0 < |D(P)| \leq Q^{2n-2-2v}. \quad (2)$$

**Problem 2.** Let  $n \geq 1$  be an integer. Given  $0 \leq w \leq n$  and a sufficiently large  $Q$ , estimate the number of pairs of polynomials  $P_1, P_2 \in \mathcal{P}_n(Q)$  such that

$$0 < |R(P_1, P_2)| \leq Q^{2n-2w}.$$

These natural problems of intrinsic interest originate from transcendental number theory. For instance, Davenport's estimate [21] for the sums of reciprocals to square roots of  $|D(P)|$  was crucial to Volkmann's proof [35] of the cubic case of a long standing conjecture of Mahler on  $S$ -numbers [28]. On the other hand, Volkmann [36] used resultants for counting pairs of irreducible polynomials with close roots to attack the higher degree case of Mahler's conjecture. The conjecture was eventually settled by Sprindžuk [31]. It is worth mentioning that properties of discriminants and resultants form the backbone of Sprindžuk's techniques as well as various generalisations, see for instance [1, 8–12, 17, 26,

32]. In particular, upper bounds for the resultant of irreducible polynomials play a crucial role in the proof of the far-reaching generalisation of Mahler’s conjecture for Hausdorff dimension [9].

There are also  $p$ -adic and ‘mixed’ analogues of the above problems which alongside the size of the discriminant and resultant address their arithmetic structure. More precisely, their formulation requires that the discriminant/resultant is divisible by large powers of given prime numbers, see [7] for an overview. In recent years there has been substantial activity in attempting to resolve Problems 1 and 2, see [4,5,13,24,23,27], as well as their  $p$ -adic versions, see [14,15] and also [7].

The discriminant and resultant naturally encode the information regarding the distance between different algebraic numbers including conjugate algebraic numbers – see definitions (3) and (4) below. Thus Problems 1 and 2 complement various questions regarding close algebraic numbers. Studying the latter dates back to a work of Mahler [29] who proved a general lower bound on the distance between two algebraic numbers. Establishing ‘correct’ upper bounds as well as quantitative results have been the subject of numerous papers including [3,16,19,20,17,18,22,30], which thereby further motivate understanding the above problems.

We now proceed by recalling some basic facts regarding discriminants and resultants. Given a polynomial  $P = a_n x^n + \cdots + a_0 \in \mathbb{Z}[x]$  of degree  $n$ , the discriminant of  $P$  is defined by

$$D(P) := a_n^{2n-2} \prod_{1 \leq i < j \leq n} (\alpha_i - \alpha_j)^2, \quad (3)$$

where  $\alpha_1, \dots, \alpha_n \in \mathbb{C}$  are the roots of  $P$ . Given two polynomials

$$P_1 = a_n x^n + \cdots + a_0 \quad \text{and} \quad P_2 = b_m x^m + \cdots + b_0$$

of degrees  $n$  and  $m$  respectively, the resultant of  $P_1$  and  $P_2$  is defined to be

$$R(P_1, P_2) := a_n^m b_m^n \prod_{\substack{1 \leq i \leq n \\ 1 \leq j \leq m}} (\alpha_i - \beta_j) = a_n^m \prod_{1 \leq i \leq n} P_2(\alpha_i) = (-1)^{mn} R(P_2, P_1), \quad (4)$$

where  $\alpha_1, \dots, \alpha_n \in \mathbb{C}$  are the roots of  $P_1$  and  $\beta_1, \dots, \beta_m \in \mathbb{C}$  are the roots of  $P_2$ . It is clear that  $D(P) = 0$  if and only if  $P$  has a repeated root, while  $R(P_1, P_2) = 0$  if and only if  $P_1$  and  $P_2$  have a common root.

It is well known and easily verified that

$$D(P) = \frac{(-1)^{\frac{n(n-1)}{2}} R(P, P')}{a_n}, \quad (5)$$

where  $P'$  is the derivative of  $P$ . Furthermore, there is a classical explicit formula for  $D(P)$  and  $R(P_1, P_2)$  via the determinant of a Sylvester matrix composed from the coefficients

of the polynomials, see [33]. Namely,  $R(P_1, P_2)$  can be computed as the following  $(n + m) \times (n + m)$  determinant

$$R(P_1, P_2) = \begin{vmatrix} a_n & a_{n-1} & \dots & a_0 & 0 & \dots & 0 \\ 0 & a_n & a_{n-1} & \dots & a_0 & \dots & 0 \\ \vdots & & \ddots & \ddots & & \ddots & \\ 0 & \dots & 0 & a_n & a_{n-1} & \dots & a_0 \\ b_m & b_{m-1} & \dots & b_0 & 0 & \dots & 0 \\ 0 & b_m & b_{m-1} & \dots & b_0 & \dots & 0 \\ \vdots & & \ddots & \ddots & & \ddots & \\ 0 & \dots & 0 & b_m & b_{m-1} & \dots & b_0 \end{vmatrix}. \quad (6)$$

The corresponding formula for  $D(P)$  can be found from the above expression and (5). Two obvious consequences of (5) and (6) are that  $D(P)$  is an integral polynomial of the coefficients of  $P$ , while  $R(P_1, P_2)$  is an integral polynomial of the coefficients of  $P_1$  and  $P_2$ . In particular,  $D(P)$  and  $R(P_1, P_2)$  return integer numbers for any choice of  $P, P_1, P_2 \in \mathbb{Z}[x] \setminus \{0\}$ . Hence, whenever  $D(P) \neq 0$  we have that  $|D(P)| \geq 1$  and whenever  $R(P_1, P_2) \neq 0$  we have that  $|R(P_1, P_2)| \geq 1$  for all choices of non-zero integral polynomials  $P, P_1, P_2$ .

Another straightforward consequences of (6) is that  $D(P)$  and  $R(P_1, P_2)$  are bounded in terms of the heights and degrees of the polynomials. In particular, it is readily verified that for every  $n \geq 2$  there exists a constant  $\gamma > 0$  which depends on  $n$  only such that for any  $P \in \mathcal{P}_n(Q)$  we have that

$$|D(P)| \leq \gamma Q^{2n-2}. \quad (7)$$

This together with the inequality  $|D(P)| \geq 1$  clarifies why the range of  $v$  is  $[0, n - 1]$  within the context of Problem 1.

Similarly, for any  $n \in \mathbb{N}$  there exists a constant  $\rho > 0$  which depends on  $n$  only such that for any two polynomials  $P_1, P_2 \in \mathcal{P}_n(Q)$  we have that

$$|R(P_1, P_2)| \leq \rho Q^{2n}.$$

In turn, this together with the inequality  $|R(P_1, P_2)| \geq 1$  clarifies why the range of  $w$  is  $[0, n]$  within the context of Problem 2.

Regarding Problem 1, in order to deal with the ‘extreme’ case  $v = n - 1$  efficiently we will weaken inequality (2) by introducing the constant  $\gamma$  the same way it appears in (7). Thus, when addressing Problem 1 we will be rather estimating the size of the following subset of  $\mathcal{P}_n(Q)$ :

$$\mathcal{D}_{n,\gamma}(Q, v) := \{P \in \mathcal{P}_n(Q) : 1 \leq |D(P)| \leq \gamma Q^{2n-2-2v}\}. \quad (8)$$

In the case  $\gamma = 1$ , we will write  $\mathcal{D}_n(Q, v)$  for  $\mathcal{D}_{n,\gamma}(Q, v)$ . Similarly, in order to deal with the extreme case  $w = n$  within [Problem 2](#) we will be estimating the size of

$$\mathcal{R}_{n,\rho}(Q, w) = \{(P_1, P_2) \in \mathcal{P}_n(Q)^2 : 0 < |R(P_1, P_2)| \leq \rho Q^{2n-2w}\}, \quad (9)$$

where  $\rho$  is a fixed constant,  $0 \leq w \leq n$  and  $Q$  is sufficiently large. In the case  $\rho = 1$ , we will write  $\mathcal{R}_n(Q, w)$  for  $\mathcal{R}_{n,\rho}(Q, w)$ .

In what follows we will use the Vinogradov symbol  $\ll$ . By definition,  $X \ll Y$  means that  $X \leq CY$  for some constant  $C$  which will depend on  $n$  only. Also we will write  $X \asymp Y$  if  $X \ll Y$  and  $Y \ll X$  simultaneously.

We now briefly recall the results that have been obtained to date. The first general estimate regarding [Problem 1](#) was established in [\[13\]](#) by showing that

$$\#\mathcal{D}_n(Q, v) \gg Q^{n+1-2v} \quad (10)$$

for  $0 < v < 1/2$ . In the case of quadratic polynomials it was shown in [\[23\]](#) that

$$\#\mathcal{D}_{2,\gamma}(Q, v) = 20(1 + \ln 2)Q^{3-2v} + O(Q^{3-3v} + Q^2) \quad (11)$$

if  $0 < v < 1/2$ . This was obtained by calculating the volume of the body in  $\mathbb{R}^3$  defined by the inequality  $|a_1^2 - 4a_2a_0| < Q^{4-2v}$ , which clearly contains all the integer vectors  $(a_0, a_1, a_2)$  that define the polynomials of interest.

Once again, by calculating the volume of relevant bodies, this time in  $\mathbb{R}^4$ , it was shown in [\[24\]](#) that

$$\#\mathcal{D}_{3,\gamma}(Q, v) \asymp Q^{4-\frac{5}{3}v} \quad (12)$$

for  $0 < v < 3/5$ . The latter estimate was in some way surprising as it led to the realisation of the fact that [\(10\)](#) is not in general sharp. One of the main goals of this paper is to make a further step in determining the ‘right size’ of  $\mathcal{D}_n(Q, v)$ . The following main result of this paper extends the lower bound given by [\(12\)](#) to the full range of  $v$  and to arbitrary degrees  $n$ .

**Theorem 1.** *Let  $n \in \mathbb{N}$ ,  $n \geq 2$  be given. Then there exists  $\gamma > 0$  depending on  $n$  only such that for any sufficiently large  $Q$  and  $0 \leq v \leq n-1$  one has that*

$$\#\mathcal{D}_{n,\gamma}(Q, v) \gg Q^{n+1-\frac{n+2}{n}v}, \quad (13)$$

where the constant implied by the Vinogradov symbol depends on  $n$  only.

**Remark 1.** If we require that  $v < n-1$ , then  $\gamma$  within the above result can be taken to be any constant, in particular 1. This readily follows from the following trivial equality

$$\mathcal{D}_{n,\gamma}(Q, v) = \mathcal{D}_{n,\gamma'}(Q, v')$$

which holds whenever  $\gamma'Q^{2v} = \gamma Q^{2v'}$ . Since  $\gamma$  and  $\gamma'$  are fixed, when  $v < n - 1$  we will also have that  $v' < n - 1$  for sufficiently large  $Q$ . Hence (13) will imply the corresponding bound for  $\#\mathcal{D}_{n,\gamma'}(Q, v')$ .

**Remark 2.** It will be immediate from the proof of Theorem 1 that (13) remains true if we additionally require that the polynomials are irreducible and of height  $H(P) \gg Q$  for some suitably chosen constant in the Vinogradov symbol. Formally, let  $\mathcal{D}_{n,\gamma,\eta}^*(Q, v)$  be the set of irreducible primitive polynomials  $P \in \mathcal{P}_n(Q)$  such that  $1 \leq |D(P)| \leq \gamma Q^{2n-2-2v}$  and  $H(P) \geq \eta Q$ . We establish that there exist  $\gamma, \eta > 0$  depending on  $n$  only such that for any sufficiently large  $Q$  and  $0 \leq v \leq n - 1$

$$\mathcal{D}_{n,\gamma,\eta}^*(Q, v) \gg Q^{n+1-\frac{n+2}{n}v}. \quad (14)$$

Much less is known about Problem 2, although some counting estimates are implicit in various papers, e.g. upper bounds for pairs of irreducible polynomials are a vital ingredient in [9]. Explicitly, it was shown in [5] that for any integers  $n \geq 2$  and  $m \in [0, n - 1]$  there exists a constant  $\rho$  depending on  $n$  only such that

$$\#\mathcal{R}_{n,\rho}\left(Q, \frac{(n+1)(m+1)}{m+2}\right) \gg Q^{\frac{2(n+1)}{(m+1)(m+2)}}. \quad (15)$$

This only gives a bound on  $\#\mathcal{R}_{n,\rho}(n, w)$  for a specific finite number of values of  $w$ , namely  $w = \frac{(n+1)(m+1)}{m+2}$ . For example, taking  $m = n - 1$  in the above estimate gives the non-trivial lower bound of  $\text{CONST} \times Q^{\frac{2}{n}}$ . Our second main result extends (15) to the whole range of  $w$  and indeed improves (15) for  $w$  of the form  $\frac{(n+1)(m+1)}{m+2}$  as above.

**Theorem 2.** Let  $n \in \mathbb{N}$  be given. Then there exists  $\rho > 0$  depending on  $n$  only such that for any sufficiently large  $Q$  we have that

$$\#\mathcal{R}_{n,\rho}(Q, w) \gg \begin{cases} Q^{2n+2-2w} & \text{if } 0 \leq w \leq \frac{n+1}{2}, \\ Q^{2n+2-2w-\frac{2}{n}(2w-n-1)} & \text{if } \frac{n+1}{2} \leq w \leq n, \end{cases} \quad (16)$$

where the constant implied by the Vinogradov symbol depends on  $n$  only.

**Remark 3.** If we require that  $w < n$ , then  $\rho$  within the above result can be taken to be any constant, in particular 1. The proof is similar to that given within the remark to Theorem 1. Similarly to Theorem 1, the proof of Theorem 2 will imply that the polynomials within  $\mathcal{R}_{n,\rho}(Q, w)$  can be required to be irreducible and of height  $H(P) \gg Q$  (see Remark 2 above).

**Remark 4.** To compare Theorem 2 to (15) take  $w = \frac{(n+1)(m+1)}{m+2}$  with integer  $m \in [0, n - 1]$ . When  $m = 0$  we are within the first case of (16), which implies

$$\#\mathcal{R}_{n,\rho}(Q, w) \gg Q^{n+1},$$

the same as (15). When  $m > 0$  we are within the second case of (16), which implies

$$\#\mathcal{R}_{n,\rho}(Q, w) \gg Q^{\frac{2(n+1)(n-m)}{(m+2)n}}. \quad (17)$$

When  $m = n - 1$  this again is the same as (15). However, as is easily seen, for  $0 < m < n - 1$  estimate (17) is significantly stronger than (15).

Theorems 1 and 2 are obtained by constructing polynomials (*resp.* pairs of polynomials) in  $\mathcal{P}_n(Q)$  with a prescribed configuration of roots. This method is not new and, in view of (3) and (4), is not surprising. Indeed, the distribution of roots instantly gives an estimate for the discriminant/resultant of polynomials. However, previously the configuration of roots was designed by using ‘hands-on techniques’. As a result the bounds for  $\#\mathcal{D}_n(Q, v)$  and  $\#\mathcal{R}_n(Q, w)$  were far from being sharp. In this paper we develop a general approach which enables us to determine the optimal configuration that maximises the number of choices for polynomials while keeping their discriminants/resultants under a given bound. Indeed, the approach boils down to a linear optimisation problem and the estimates for  $\#\mathcal{D}_n(Q, v)$  and  $\#\mathcal{R}_n(Q, w)$  we obtain correspond to optimal solutions to these problems. As a result, we believe that the estimates obtained in the above theorems are best possible, perhaps up to an arbitrarily small additive constant  $\delta > 0$  in the exponents within (13) and (16).

It should be noted that the case  $v = 0$  within Problem 1 corresponds to essentially imposing no restriction on the discriminant. Naturally, our lower bound for  $\#\mathcal{D}_n(Q, v)$  in this case gives  $\gg Q^{n+1}$ , which is however quite trivial. Nevertheless, a much more detailed insight into the distribution of such (typical) values of the discriminant is provided in [25]. The main result of [25] proves an asymptotic formula for the number of polynomials  $P \in \mathcal{P}_n(Q)$  such that

$$a \leq \frac{|D(P)|}{Q^{2n-2}} \leq b$$

with a logarithmically small error term.

## 2. Rational points near planar curves and the quadratic case

Considering the case  $n = 2$  in this section we obtain the complementary upper bound for  $\mathcal{D}_2(Q, v)$  which holds for all  $v \in (0, 1)$  and thus extends (11) to the full range of  $v$ .

**Theorem 3.** *For any  $v \in (0, 1)$  and any sufficiently large  $Q$  we have that*

$$\#\mathcal{D}_2(Q, v) \asymp Q^{3-2v}. \quad (18)$$



By [Theorem 1](#), we only have to prove the upper bound, that is  $\mathcal{D}_2(Q, v) \ll Q^{3-2v}$ . The proof will be based on the following counting result for rational points near planar curves due to Vaughan and Velani which is a straightforward consequence of [Theorem 3](#) from [\[34\]](#). Note that the case  $f(x) = x^2$  that will be of interest for us is also treated within [\[6, Appendix 2 and Case \(c\) of § 2.1\]](#).

**Theorem 4** (Vaughan & Velani [\[34\]](#)). Let  $f : [\alpha, \beta] \rightarrow \mathbb{R}$  be a  $C^{(3)}$  function. Suppose that

$$\inf_{\alpha \leq x \leq \beta} |f''(x)| > 0.$$

Let

$$N(T, \varepsilon) := \#\{(a, q) \in \mathbb{Z} \times \mathbb{N} : q \leq T, \alpha q < a \leq \beta q, \|qf(a/q)\| < \varepsilon\},$$

where  $\|\cdot\|$  denotes the distance to the nearest integer. Then for any  $T \geq 1$ ,  $\varepsilon \in (0, \frac{1}{2})$  and any  $\delta > 0$  one has that

$$N(T, \varepsilon) \ll \varepsilon T^2 + \varepsilon^{-1/2} T^{1/2+\delta} + \varepsilon^{-\delta} T^{1+\delta},$$

where the implied constant in the Vinogradov symbol depends on  $f$  and  $\delta$  only.

**Proof of Theorem 3.** First of all note that there is no loss of generality in assuming that  $v > 0$  as otherwise the statement of [Theorem 3](#) is trivial. Note that any polynomial  $P \in \mathcal{P}_2(Q)$  is of the form

$$P = ax^2 + bx + c, \quad a, b, c \in \mathbb{Z}, \quad a \neq 0, \quad \max\{|a|, |b|, |c|\} \leq Q.$$

Thus the cardinality of  $\mathcal{D}_2(Q, v)$  is bounded by that of the set

$$\{(a, b, c) \in \mathbb{Z}^3 : \max\{|a|, |b|, |c|\} \leq Q, \quad 0 < |b^2 - 4ac| \leq Q^{2-2v}\}. \quad (19)$$

We shall consider two cases depending on the size of the coefficients  $a, b, c$ .

**Case 1:**  $\max\{|a|, |c|\} \leq \frac{1}{3}|b|$ . Then,  $|4ac| < \frac{1}{2}|b|^2$  and therefore  $|b^2 - 4ac| \geq \frac{1}{2}|b|^2$ . Since we are interested in the triples  $(a, b, c)$  which lie in [\(19\)](#), we get that  $|b| \ll Q^{1-v}$ . Since  $\max\{|a|, |b|\} \leq \frac{1}{3}|b|$ , we also have that  $\max\{|a|, |b|\} \ll Q^{1-v}$ . Thus, the number of triples in question is  $\ll (Q^{1-v})^3 = Q^{3-3v} \leq Q^{3-2v}$  and we obtain the required bound.

**Case 2:**  $\max\{|a|, |c|\} > \frac{1}{3}|b|$ . Since the discriminant  $b^2 - 4ac$  does not change if we swap  $a$  and  $c$ , without loss of generality we can assume that  $|a| = \max\{|a|, |c|\}$ . In particular, we have that  $\frac{1}{3}|b| \leq |a| \leq Q$ . Next, for each  $a$  as above there exists an integer  $t \geq 0$  such that  $2^t \leq Q$  and  $2^t \leq |a| < 2^{t+1}$ . Then dividing the inequality in [\(19\)](#) through by  $a$  we get that

$$|a(b/a)^2 - 4c| \ll \varepsilon_t := 2^{-t} Q^{2-2v}. \quad (20)$$

If  $\varepsilon_t < \frac{1}{2}$ , then using Theorem 4 with

$$\begin{aligned} f(x) &= x^2, & T &= 2^{t+1} & \varepsilon &= \varepsilon_t \\ [\alpha, \beta] &= [0, 3], & \delta &= \min\{3 - 3v, v^{-1} - 1\}, \end{aligned}$$

where the choice of  $\delta$  is justified by the conditions  $0 < v < 1$ , we conclude that the number of triples  $(a, b, c)$  in (19) such that  $|a| = \max\{|a|, |c|\} > \frac{1}{3}|b|$  and  $2^t \leq |a| < 2^{t+1}$  is

$$\ll \varepsilon_t 2^{2t} + \varepsilon_t^{-1/2} 2^{t(1/2+\delta)} + \varepsilon_t^{-\delta} 2^{t(1+\delta)} \ll 2^t Q^{2-2v} + 2^t Q^{-1+v} 2^{t\delta} + Q^{2v\delta} 2^t. \quad (21)$$

This estimate also holds when  $\varepsilon_t \geq \frac{1}{2}$ . Indeed, for  $\varepsilon_t \geq \frac{1}{2}$ , assuming  $a$  and  $b$  are fixed, the number of different integers  $c$  satisfying (20) is  $\ll \varepsilon_t$ . Also the number of different integer pairs  $(a, b)$  such that  $2^t \leq |a| < 2^{t+1}$  and  $\frac{1}{3}|b| \leq |a|$  is  $\ll 2^{2t}$ . Hence, when  $\varepsilon_t \geq \frac{1}{2}$  the number of triples  $(a, b, c)$  in question is  $\ll 2^t Q^{2-2v}$ .

Summing (21) over non-negative integers  $t$  such that  $2^t \leq Q$  gives the following estimate for the number of triples  $(a, b, c)$  in question:

$$\ll Q^{3-2v} + Q^{v+\delta} + Q^{1+2v\delta}$$

which is  $\ll Q^{3-2v}$  since  $\delta = \min\{3 - 3v, v^{-1} - 1\}$ . The proof is thus complete.  $\square$

### 3. Auxiliary lemmas

We begin with the following statement which is a version of Lemma 4 from [3] and based on [2, Theorem 5.8]. In what follows, given a Lebesgue measurable set  $X \subset \mathbb{R}^d$ ,  $|X|$  will stand for its (ambient) Lebesgue measure.

**Lemma 1.** *Let  $n \geq 2$  and  $v_0, \dots, v_n$  be a collection of real numbers such that*

$$v_0 + \dots + v_n = 0 \quad (22)$$

*and that*

$$v_0 \geq v_1 \geq \dots \geq v_n = -1. \quad (23)$$

*Then there are positive constants  $\delta_0$  and  $c_0$  depending on  $n$  only with the following property. For any interval  $J \subset [-\frac{1}{2}, \frac{1}{2}]$  there is a sufficiently large  $Q_0$  such that for all  $Q > Q_0$  there is a measurable set  $G_J \subset J$  satisfying*

$$|G_J| \geq \frac{3}{4}|J| \quad (24)$$

*such that for every  $x \in G_J$  there are  $n+1$  linearly independent primitive irreducible polynomials  $P \in \mathbb{Z}[x]$  of degree exactly  $n$  such that*

$$\begin{cases} \delta_0 Q^{-v_0} \leq |P(x)| \leq c_0 Q^{-v_0}, \\ \delta_0 Q^{-v_j} \leq |P^{(j)}(x)| \leq c_0 Q^{-v_j} \quad (1 \leq j \leq n). \end{cases} \quad (25)$$

**Remark 5.** Note the constant  $\frac{3}{4}$  is not crucial for the above lemma and can be replaced with any other positive constant strictly less than 1.

We now establish a general statement that relates the derivatives of  $P$  and its roots.

**Lemma 2.** Let  $x \in \mathbb{C}$  be a fixed point,  $P$  be a polynomial with complex coefficients of degree  $\deg P = n > 0$  and let  $\alpha_1, \dots, \alpha_n$  be the roots of  $P$  ordered so that

$$|x - \alpha_1| \leq |x - \alpha_2| \leq \dots \leq |x - \alpha_n|. \quad (26)$$

Let  $a_n$  be the leading coefficient of  $P$ . Then for  $0 \leq j < n$  we have that

$$|P^{(j)}(x)| \leq \binom{n}{j} |a_n| |x - \alpha_{j+1}| \cdots |x - \alpha_n|, \quad (27)$$

where as usually  $\binom{n}{j} = \frac{n!}{j!(n-j)!}$  is a binomial coefficient. If we further assume that for some  $j > 0$

$$|x - \alpha_j| < \frac{1}{2} \binom{n}{j}^{-1} |x - \alpha_{j+1}|, \quad (28)$$

then we also have that

$$|P^{(j)}(x)| \geq \frac{1}{2} |a_n| |x - \alpha_{j+1}| \cdots |x - \alpha_n|. \quad (29)$$

**Proof.** First observe that

$$P^{(j)}(x) = a_n \sum_{1 \leq i_1 < \dots < i_{n-j} \leq n} (x - \alpha_{i_1}) \cdots (x - \alpha_{i_{n-j}}). \quad (30)$$

Note that the right hand side contains exactly  $\binom{n}{j}$  terms. By (26), the largest of these terms in absolute value is  $T = (x - \alpha_{j+1}) \cdots (x - \alpha_n)$  and so (27) readily follows from (30).

Further, (29) is trivially true when  $j = 0$  (it actually becomes an equality when the factor  $\frac{1}{2}$  is removed). When (28) is satisfied for some  $j > 0$ , then any term of the sum in the right hand side of (30) different from  $T$  is  $\leq \frac{1}{2} \binom{n}{j}^{-1} |T|$ . Hence,

$$\left| \sum_{1 \leq i_1 < \dots < i_{n-j} \leq n} (x - \alpha_{i_1}) \cdots (x - \alpha_{i_{n-j}}) - T \right| \leq \frac{1}{2} |T|$$

whence (29) readily follows upon using the triangle inequality.  $\square$

Now we apply [Lemma 2](#) to inequalities [\(25\)](#) to obtain the following

**Lemma 3.** *Let  $n$ ,  $m$  and  $v_j$  be the same as in [Lemma 1](#). Let*

$$d_j = v_{j-1} - v_j \quad (1 \leq j \leq n). \quad (31)$$

Suppose that

$$d_1 \geq d_2 \geq \dots \geq d_n \geq 0 \quad (32)$$

and that for some  $x \in \mathbb{C}$  and  $Q > 1$  inequalities [\(25\)](#) are satisfied by some polynomial  $P$  over  $\mathbb{C}$  of degree  $\deg P = n$ . Then the roots  $\alpha_1, \dots, \alpha_n \in \mathbb{C}$  of  $P$ , ordered according to their distance to  $x$  as in [\(26\)](#), satisfy the following inequalities

$$|x - \alpha_j| \leq c_j Q^{-d_j} \quad (1 \leq j \leq n), \quad (33)$$

where

$$c_1 = nc_0 \delta_0^{-1} \quad \text{and} \quad c_{j+1} = \max \left\{ \frac{2c_0}{\delta_0} \binom{n}{j+1}, 2c_j \binom{n}{j} \right\} \quad (1 \leq j \leq n-1).$$

**Proof.** Since  $P$  is of degree  $n$  it has  $n$  complex roots, say  $\alpha_1, \dots, \alpha_n$ . Order the roots with respect to their distance to  $x$ , thus ensuring the validity of [\(26\)](#).

First take  $j = 1$ . Then, by [Lemma 2](#), we have that

$$|P'(x)| \leq n|a_n||x - \alpha_2| \cdots |x - \alpha_n| = n \frac{|P(x)|}{|x - \alpha_1|}.$$

Then, by [\(25\)](#), we obtain that

$$|x - \alpha_1| \leq nc_0 \delta_0^{-1} Q^{-v_0+v_1} = nc_0 \delta_0^{-1} Q^{-d_1}$$

as required. We proceed by induction. Suppose [\(33\)](#) holds for some  $j < n$  and we now want to prove it for  $j+1$ . If [\(28\)](#) is fulfilled, then we are in a position similar to the case  $j = 1$ . Namely, by [Lemma 2](#), we have [\(29\)](#) while, by [\(27\)](#) written for  $P^{(j+1)}(x)$ , we have

$$|P^{(j+1)}(x)| \cdot |x - \alpha_{j+1}| \leq \binom{n}{j+1} |a_n| |x - \alpha_{j+1}| \cdots |x - \alpha_n|. \quad (34)$$

This together with [\(29\)](#) now gives

$$|P^{(j+1)}(x)| \cdot |x - \alpha_{j+1}| \leq 2 \binom{n}{j+1} |P^{(j)}(x)|.$$

By [\(25\)](#), we get

$$|x - \alpha_{j+1}| \leq 2c_0 \binom{n}{j+1} \delta_0^{-1} Q^{-v_j+v_{j+1}} = 2c_0 \binom{n}{j+1} \delta_0^{-1} Q^{-d_{j+1}} \leq c_{j+1} Q^{-d_{j+1}}.$$

On the other hand, if (28) is not fulfilled, then,  $|x - \alpha_{j+1}| \leq 2\binom{n}{j}|x - \alpha_j|$ . Therefore, using induction and (32) we obtain that

$$|x - \alpha_{j+1}| \leq 2\binom{n}{j}c_j Q^{-d_j} \leq 2\binom{n}{j}c_j Q^{-d_{j+1}} \leq c_{j+1} Q^{-d_{j+1}}.$$

This completes the proof.  $\square$

The last auxiliary statement, which will only be used in the proof of Theorem 2, concerns measurable sets in the plane which lie near the line  $y = x$ .

**Lemma 4.** *Let  $I \subset \mathbb{R}$  be a finite interval,  $0 < \Delta < |I|$  and*

$$X_\Delta = \{(x, y) \in \mathbb{R}^2 : |x - y| \leq \Delta\}.$$

*Let  $A$  be a Lebesgue measurable subset of  $I$  such that  $|A| \geq \lambda|I|$  for some  $\lambda \in (0, 1)$ . Let  $A^2 := \{(x, y) \in \mathbb{R}^2 : x, y \in A\}$ . Then*

$$|A^2 \cap X_\Delta| \geq 2^{-6}\lambda^3\Delta|I|.$$

**Proof.** Let  $T$  be an integer such that

$$\frac{|I|}{\Delta} < T < \frac{2|I|}{\Delta}. \quad (35)$$

The existence of  $T$  follows from the inequality  $\Delta < |I|$ .

Divide  $I$  into  $T$  equal subintervals  $I_1, \dots, I_T$ . By (35),  $\frac{1}{2}\Delta \leq |I_j| \leq \Delta$ . Let  $N$  be the number of intervals  $I_j$  such that  $|A \cap I_j| \geq \frac{1}{4}\lambda\Delta$ . Then

$$|A| \leq \frac{1}{4}\lambda\Delta T + N\Delta.$$

On the other hand, using (35) we get that

$$|A| \geq \lambda|I| \geq \frac{1}{2}\lambda\Delta T.$$

Hence  $\frac{1}{2}\lambda\Delta T \leq \frac{1}{4}\lambda\Delta T + N\Delta$  whence we obtain that  $N \geq \frac{1}{4}\lambda T$ .

Since  $|I_j| \leq \Delta$  we trivially have that  $I_j^2 \subset X_\Delta$ . Hence, for each  $j$  we have the inclusion

$$(A \cap I_j)^2 \subset A^2 \cap X_\Delta.$$

In particular,  $|(A \cap I_j)^2| \geq \frac{1}{16}\lambda^2\Delta^2$  whenever  $|A \cap I_j| \geq \frac{1}{4}\lambda\Delta$ . Recall that we have  $N \geq \frac{1}{4}\lambda T$  intervals  $I_j$  satisfying this condition. Hence,

$$|A^2 \cap X_\Delta| \geq N \cdot \frac{1}{16}\lambda^2\Delta^2 \geq 2^{-6}\lambda^3 T \Delta^2 \stackrel{(35)}{\geq} 2^{-6}\lambda^3 \frac{|I|}{\Delta} \Delta^2 = 2^{-6}\lambda^3 |I| \Delta$$

as required.  $\square$

#### 4. Proof of Theorem 1

Let  $v_0, \dots, v_n$  be given and satisfy (22), (23) and let the parameters  $d_j$  be given by (31) and satisfy (32). First of all let us show that

$$\sum_{j=1}^n j d_j = n + 1. \quad (36)$$

Indeed, by (31), we have that  $v_{j-1} = d_j + v_j$ . Hence  $v_{j-1} = d_j + \dots + d_n + v_n$ . Since  $v_n = -1$  we have that  $v_{j-1} + 1 = d_j + \dots + d_n$ . Also  $v_n + 1 = 0$ . Summing these equations over  $j = 0, \dots, n$ , by (22), we get

$$n + 1 = \sum_{j=0}^n (v_j + 1) = \sum_{j=0}^{n-1} (v_j + 1) = \sum_{j=0}^{n-1} (d_{j+1} + \dots + d_n) = \sum_{j=1}^n j d_j$$

as stated in (36).

Now, let  $J = [-\frac{1}{2}, \frac{1}{2}]$ ,  $Q$  be sufficiently large and  $x \in G_J$ , where  $G_J$  is the same as in Lemma 1. By Lemma 1, inequalities (25) are satisfied for some irreducible polynomial  $P \in \mathbb{Z}[x]$  of degree  $n$ . Then, by Lemma 3, we have (33). Hence, for any pair of integers  $(i, j)$  satisfying  $1 \leq i < j \leq n$  we have that

$$|\alpha_i - \alpha_j| \leq |x - \alpha_i| + |x - \alpha_j| \ll Q^{-d_j}.$$

By (25), we have that

$$H(P) \leq h_0 Q \quad (37)$$

for some constant  $h_0$  which depends on  $n$  only. Therefore, using (3) we conclude that

$$1 \leq |D(P)| \ll Q^{2n-2} \prod_{1 \leq i < j \leq n} Q^{-2d_j} = Q^{2n-2-2 \sum_{j=2}^n (j-1)d_j}. \quad (38)$$

Note that the left hand side inequality is due to the irreducibility of  $P$ . The goal is to construct polynomials with

$$1 \leq |D(P)| \leq \gamma Q^{2n-2-2v} \quad (39)$$

with some suitably chosen constant  $\gamma$ . By (38), inequalities (39) are fulfilled if we impose the condition  $\sum_{j=2}^n (j-1)d_j = v$ . Subtracting this from (36) we obtain the equivalent equation

$$\sum_{j=1}^n d_j = n + 1 - v. \quad (40)$$

Now, we estimate the number of polynomials that we can obtain this way. By (33) and Lemma 1, for every  $x \in G_J$  we have that  $|x - \alpha_1(P)| \ll Q^{-d_1}$ , where  $P$  arises from Lemma 1. Therefore, since (39) holds whenever (40) is satisfied, we have that

$$G_J \subset \bigcup_{P \in \mathcal{D}_{n,\gamma}(h_0 Q, v)} \bigcup_{j=1}^n \{|x - \alpha_j(P)| \leq c_1 Q^{-d_1}\},$$

where  $\gamma$  and  $c_1$  depend on  $n$  only. Here we have used bound (37) on the height. Hence

$$\frac{3}{4} = \frac{3}{4}|J| \leq 2nc_1 Q^{-d_1} \#\mathcal{D}_{n,\gamma}(h_0 Q, v)$$

and we get that

$$\#\mathcal{D}_{n,\gamma}(h_0 Q, v) \gg Q^{d_1}.$$

To obtain the best lower bound we should maximise  $d_1$  subject to conditions (32), (36) and (40). This is a linear optimisation problem. By (36),  $d_1$  would be maximal if we could ensure that  $d_2 = \dots = d_n$ . Then (36) and (40) give a system of two linear equations with two variables, namely  $d_1$  and  $d_2$ , from which we easily find that

$$d_2 = \dots = d_n = \frac{2v}{n(n-1)} \quad \text{and} \quad d_1 = n+1 - \frac{n+2}{n}v.$$

A quick check shows that  $d_1 \geq d_2$  because  $v \leq n-1$ . Thus (32) is also fulfilled. One can also verify that

$$v_0 = n-v \quad \text{and} \quad v_j = -1 + (n-j)d_2 \quad (1 \leq j \leq n-1).$$

Thus,

$$\#\mathcal{D}_{n,\gamma}(h_0 Q, v) \gg Q^{d_1} \geq Q^{n+1 - \frac{n+2}{n}v}.$$

To complete the proof it remains to rescale the bound on the height by making the following change of variables  $\tilde{Q} = h_0 Q$  in the above expression. This results in the required lower bound

$$\#\mathcal{D}_{n,\gamma}(\tilde{Q}, v) \gg \tilde{Q}^{n+1 - \frac{n+2}{n}v}$$

and completes the proof.

## 5. Proof of Theorem 2

The start is the same as in the proof of Theorem 1. Let  $v_0, \dots, v_n$  be given and satisfy (22) and (23) and let the parameters  $d_j$  be given by (31) and satisfy (32). Equation (36) is then again satisfied.

Let  $J = [-\frac{1}{2}, \frac{1}{2}]$ ,  $Q$  be sufficiently large,  $G_J$  be the same as in Lemma 1 and  $(x_1, x_2) \in G_J^2 \cap X_{Q^{-t}}$ , where  $X_{Q^{-t}}$  is  $X_\Delta$  with  $\Delta = Q^{-t}$  as defined in Lemma 4. By Lemma 1, for each  $i = 1, 2$  there is an irreducible polynomial  $P_i$  over  $\mathbb{Z}$  of degree  $n$  such that inequalities (25) with  $x = x_i$  are satisfied. Furthermore, for each  $i = 1, 2$  Lemma 1 offers  $n + 1$  linearly independent choices of irreducible (over  $\mathbb{Q}$ ) polynomials  $P_i$ . Therefore, we can assume that  $P_1$  and  $P_2$  are linearly independent. If  $P_1$  and  $P_2$  had a common root, by the fact they are irreducible, each of them would have to divide the other, meaning  $P_1 = \pm P_2$  and contradicting their linear independence. Therefore,  $P_1$  and  $P_2$  have no common roots and we have that  $|R(P_1, P_2)| \geq 1$ . We will impose the condition

$$d_1 \geq t \geq d_2. \quad (41)$$

By Lemma 3, the roots  $\alpha_1, \dots, \alpha_n \in \mathbb{C}$  of  $P_1$ , ordered according to their distance to  $x_1$ , satisfy the inequalities

$$|x_1 - \alpha_i| \leq c_i Q^{-d_i} \quad (1 \leq i \leq n) \quad (42)$$

and the roots  $\beta_1, \dots, \beta_n \in \mathbb{C}$  of  $P_2$ , ordered according to their distance to  $x_2$ , satisfy the inequalities

$$|x_2 - \beta_j| \leq c_j Q^{-d_j} \quad (1 \leq j \leq n). \quad (43)$$

Hence, for any pair of integers  $(i, j)$  such that  $1 \leq i, j \leq n$  using the triangle inequality we obtain that

$$|\alpha_i - \beta_j| \leq |x_1 - x_2| + |x_1 - \alpha_i| + |x_2 - \beta_j| \ll Q^{-\min\{t, d_i, d_j\}}.$$

By (25), we have that

$$H(P_i) \leq h_0 Q \quad (i = 1, 2) \quad (44)$$

for some constant  $h_0$  which depends on  $n$  only. Therefore, by (4), we get that

$$1 \leq |R(P_1, P_2)| \ll Q^{2n} \prod_{1 \leq i, j \leq n} Q^{-\min\{t, d_i, d_j\}}.$$

In view of (32) and (41) we then have that

$$1 \leq |R(P_1, P_2)| \ll Q^{2n-t-d_2-\dots-d_n-2\sum_{1 \leq i < j \leq n} d_j}. \quad (45)$$

The goal is to construct pairs of polynomials  $P_1, P_2$  with



$$1 \leq |R(P_1, P_2)| \leq \rho Q^{2n-2w} \quad (46)$$

with some suitably chosen constant  $\rho$ . By (45), inequalities (46) are fulfilled if we impose the condition

$$t + d_2 + \cdots + d_n + 2 \sum_{1 \leq i < j \leq n} d_j = 2w. \quad (47)$$

Using (36) we get that

$$\sum_{1 \leq i < j \leq n} d_j = \sum_{j=1}^n (j-1)d_j = n+1 - \sum_{j=1}^n d_j.$$

Hence (47) transforms into

$$2d_1 - t + \sum_{j=2}^n d_j = 2n + 2 - 2w. \quad (48)$$

Now, we estimate the number of pairs  $(P_1, P_2)$  that we can obtain this way. By Lemma 4, we have that  $|G_J^2 \cap X_{Q^{-t}}| \gg Q^{-t}|J| = Q^{-t}$ . Next, by Lemma 1, (44) and (46), we have that

$$G_J^2 \cap X_{Q^{-t}} \subset \bigcup_{(P_1, P_2) \in \mathcal{R}_{n, \rho}(h_0 Q, w)} \bigcup_{i, j=1}^n \left\{ (x_1, x_2) : \begin{array}{l} |x_1 - \alpha_i(P_1)| \leq c_1 Q^{-d_1} \\ |x_2 - \beta_j(P_2)| \leq c_1 Q^{-d_1} \end{array} \right\},$$

where  $\rho$  and  $c_1$  depend on  $n$  only. Hence

$$Q^{-t} \ll |G_J^2 \cap X_{Q^{-t}}| \ll Q^{-2d_1} \# \mathcal{R}_{n, \rho}(h_0 Q, w)$$

and we get that

$$\# \mathcal{R}_{n, \rho}(h_0 Q, w) \gg Q^{2d_1-t}.$$

To obtain the best lower bound we should maximise  $2d_1 - t$  subject to conditions (32), (36), (41) and (48). This is again a linear optimisation problem. By (36),  $2d_1 - t$  would be maximal if we could ensure that  $d_2 = \cdots = d_n$ . Assuming these equations, (32), (36), (41) and (48) give

$$\begin{aligned} 2d_1 + (n-1)(n+2)d_2 &= 2n+2, \\ (n^2-1)d_2 + t &= 2w, \\ d_1 &\geq t \geq d_2. \end{aligned} \quad (49)$$

In particular, (48) with  $d_2 = \cdots = d_n$  implies that

$$2d_1 - t + (n - 1)d_2 = 2n + 2 - 2w. \quad (50)$$

Thus, we have to maximise  $2d_1 - t$ , bearing in mind that  $0 < w \leq n$  together with the above constraints. Clearly, the best we can get is  $2d_1 - t = 2n + 2 - 2w$  in the case  $d_2 = 0$ . However, this does not a priori mean that the last condition of (49) is fulfilled. For this reason we are forced to consider the following two cases.

**Case (i):**  $w \leq \frac{n+1}{2}$ . Then we can indeed take  $d_2 = 0$ ,  $d_1 = n + 1$  and  $t = 2w$ . Clearly, the last condition of (49) holds. In this case

$$v_0 = n, \quad v_j = -1 \quad (1 \leq j \leq n - 1)$$

and we get that

$$\#\mathcal{R}_{n,\rho}(h_0Q, w) \gg Q^{2d_1-t} \geq Q^{2n+2-2w}.$$

**Case (ii):**  $\frac{n+1}{2} < w \leq n$ . In this case  $d_2 = 0$  would imply via (49) that  $t > n + 1 > d_1$ , contrary to the requirement  $d_1 \geq t$ . It is easily calculated from (49) that the smallest value of  $d_2$  which enables the condition  $t \leq d_1$  is

$$d_2 = \frac{4w - 2n - 2}{n(n-1)}.$$

In view of (50) this maximises  $2d_1 - t$ . Then, from (49) we obtain that

$$t = d_1 \quad \text{and} \quad d_1 = 2n + 2 - 2w - \frac{2}{n}(2w - n - 1).$$

A quick check shows that  $0 \leq d_2 \leq d_1$  since  $\frac{n+1}{2} < w \leq n$ . Thus, all the required conditions are met and we have that

$$\#\mathcal{R}_{n,\rho}(h_0Q, w) \gg Q^{2d_1-t} = Q^{d_1} \geq Q^{2n+2-2w-\frac{2}{n}(2w-n-1)}.$$

Finally, to complete the proof it remain to re-scale the bound on the height the same way as we did in the proof of Theorem 1, that is by setting  $\tilde{Q} = h_0Q$ .

## 6. Comparing the estimates for different degrees

Given  $n \in \mathbb{N}$ ,  $n \geq 2$ ,  $Q > 1$  and  $\xi \geq 0$ , consider the set

$$\mathcal{D}_{\leq n}(Q, \xi) := \{P \in \mathbb{Z}[x] : 2 \leq \deg P \leq n, H(P) \leq Q, 1 \leq |D(P)| \leq \gamma Q^\xi\}.$$

This set is composed of the polynomials of degree up to  $n$  with a given restriction on the discriminant. By Theorem 1, there is a  $\gamma > 0$  which depends on  $n$  only such that the

number of polynomials  $P \in \mathcal{P}_k(Q)$  lying in this set is

$$\gg Q^{k+1-\frac{k+2}{k}v},$$


where  $v$  is determined from the equation  $\xi = 2k - 2 - 2v$  when  $\xi \leq 2k - 2$  and  $v = 0$  when  $\xi > 2k - 2$ . Let

$$f_k(\xi) = k + 1 - \frac{k+2}{k}\xi = \begin{cases} \frac{2}{k} + \frac{k+2}{2k}\xi & \text{if } 0 < \xi \leq 2k - 2, \\ k + 1 & \text{if } \xi > 2k - 2. \end{cases}$$

Then the number of polynomials  $P \in \mathcal{P}_k(Q)$  lying in  $\mathcal{D}_{\leq n}(Q, \xi)$  is  $\gg Q^{f_k(\xi)}$ . Consequently,

$$\mathcal{D}_{\leq n}(Q, \xi) \gg Q^{d_n(\xi)}, \quad \text{where} \quad d_n(\xi) := \max_{2 \leq k \leq n} f_k(\xi).$$

Since the slopes of the lines  $y_k = \frac{2}{k} + \frac{k+2}{2k}\xi$  and the points of their intersection with the  $y$ -axis get smaller as  $k$  gets bigger, the graph of  $y_k$  lies under the graph of any other line  $y_m$  with  $m < k$ . Hence, the graph of  $f_k(\xi)$  will always intersect that of  $d_{k-1}(\xi)$  at some point  $\xi > 2k - 4$ . Hence the contribution to  $\mathcal{D}_{\leq n}(Q, \xi)$  by polynomials of degree  $k$  will be the maximum only when the contribution by polynomials of smaller degrees is no longer growing. This may seem rather counterintuitive as there are generally many more polynomials of higher degree. Below we sketch the graph of  $y = d_n(\xi)$  for the case  $n = 4$ , which is enough to exhibit the ‘staircase’ nature of this function.



The following explicit formula for  $d_n(\xi)$  is also readily computed:

$$d_n(\xi) = \begin{cases} \xi + 1 & \text{if } 0 \leq \xi \leq 2, \\ k + 1 & \text{if } 2k - 4 \leq \xi \leq 2k - 4 + \frac{4}{k+2} \text{ for } 3 \leq k < n, \\ \frac{2}{k} + \frac{k+2}{2k}\xi & \text{if } 2k - 4 + \frac{4}{k+2} \leq \xi \leq 2k - 2 \text{ for } 3 \leq k \leq n, \\ n + 1 & \text{if } \xi \geq 2n - 2. \end{cases}$$

*Resultants.* Given  $n \in \mathbb{N}$ ,  $Q > 1$  and  $\xi \geq 0$ , consider the set

$$\mathcal{R}_{\leq n}(Q, \xi) := \left\{ (P_1, P_2) \in \mathbb{Z}[x] \times \mathbb{Z}[x] : \begin{array}{l} 2 \leq \deg P_i \leq n, \ H(P_i) \leq Q, \\ 1 \leq |R(P_1, P_2)| \leq \rho Q^\xi \end{array} \right\}.$$

By [Theorem 2](#), there is a constant  $\rho > 0$  such that the number of pair of polynomials  $P_1, P_2 \in \mathcal{P}_k(Q)$  lying in this set is

$$\gg Q^{2k+2-2w} = Q^{\xi+2},$$

where  $w$  is determined from the equation  $\xi = 2k - 2w$  when  $0 \leq 2w \leq k + 1$ . This gives the following restriction on  $\xi$ :  $k - 1 \leq \xi \leq 2k$ . Consequently,

$$\mathcal{R}_{\leq n}(Q, \xi) \gg Q^{\xi+2} \quad \text{for} \quad 0 \leq \xi \leq 2n + 2.$$

## Acknowledgments

The first and second authors are grateful to EPSRC for the support of their exchange visits through grant EP/J018260/1 and to SFB701 for supporting their visits to the University of Bielefeld. The authors are also grateful to the anonymous reviewer for many useful comments.

## References

- [1] V. Beresnevich, On approximation of real numbers by real algebraic numbers, *Acta Arith.* 90 (2) (1999) 97–112.
- [2] V. Beresnevich, Rational points near manifolds and metric Diophantine approximation, *Ann. of Math.* (2) 175 (1) (2012) 187–235.
- [3] V. Beresnevich, V. Bernik, F. Götze, The distribution of close conjugate algebraic numbers, *Compos. Math.* 146 (5) (2010) 1165–1179.
- [4] V. Beresnevich, V. Bernik, F. Götze, Simultaneous approximations of zero by an integral polynomial, its derivative, and small values of discriminants, *Dokl. Nats. Akad. Nauk Belarusi* 54 (2) (2010) 26–28, 125 (in Russian).
- [5] V. Beresnevich, V. Bernik, F. Götze, On the distribution of the values of the resultants of integral polynomials, *Dokl. Nats. Akad. Nauk Belarusi* 54 (5) (2010) 21–23, 125 (in Russian).

- [6] V. Beresnevich, D. Dickinson, S. Velani, Diophantine approximation on planar curves and the distribution of rational points, with an Appendix II by R.C. Vaughan, *Ann. of Math.* (2) 166 (2) (2007) 367–426.
- [7] V. Beresnevich, V. Bernik, F. Götze, O. Kukso, Distribution of algebraic numbers and metric theory of Diophantine approximation, in: *Limit Theorems in Probability, Statistics and Number Theory*, in: Springer Proc. Math. Stat., vol. 42, Springer, Heidelberg, 2013, pp. 23–48.
- [8] V. Bernik, A metric theorem on the simultaneous approximation of zero by values of integer polynomials, *Izv. Akad. Nauk SSSR Ser. Mat.* 44 (1) (1980) 24–45, 238 (in Russian).
- [9] V. Bernik, An application of Hausdorff dimension in the theory of Diophantine approximation, *Acta Arith.* 42 (1983) 219–253 (in Russian). English transl. in *Amer. Math. Soc. Transl.* 140 (1988) 15–44.
- [10] V. Bernik, The exact order of approximating zero by values of integral polynomials, *Acta Arith.* 53 (1) (1989) 17–28 (in Russian).
- [11] V. Bernik, N. Budarina, D. Dickinson, A divergent Khintchine theorem in the real, complex, and  $p$ -adic fields, *Lith. Math. J.* 48 (2) (2008) 158–173.
- [12] V. Bernik, N. Budarina, D. Dickinson, Simultaneous Diophantine approximation in the real, complex and  $p$ -adic fields, *Math. Proc. Cambridge Philos. Soc.* 149 (2) (2010) 193–216.
- [13] V. Bernik, F. Götze, O. Kukso, Lower bounds for the number of integral polynomials with given order of discriminants, *Acta Arith.* 133 (2008) 375–390.
- [14] V. Bernik, F. Götze, O. Kukso, On the divisibility of the discriminant of an integral polynomial by prime powers, *Lith. Math. J.* 48 (2008) 380–396.
- [15] N. Budarina, D. Dickinson, Y. Jin, On the number of polynomials with small discriminants in the euclidean and  $p$ -adic metrics, *Acta. Math. Sin.* 28 (3) (2012) 469–476.
- [16] N. Budarina, F. Götze, Distance between conjugate algebraic numbers in clusters, *Transl. Mat. Zametki* 94 (5) (2013) 780–783, *Math. Notes* 94 (5–6) (2013) 816–819.
- [17] Y. Bugeaud, *Approximation by Algebraic Numbers*, Cambridge Tracts in Mathematics, vol. 160, Cambridge University Press, Cambridge, 2004, 274 pp.
- [18] Y. Bugeaud, A. Dujella, Root separation for reducible integer polynomials, *Acta Arith.* 162 (4) (2014) 393–403.
- [19] Y. Bugeaud, M. Mignotte, On the distance between roots of integer polynomials, *Proc. Edinb. Math. Soc.* 47 (2) (2004) 553–556.
- [20] Y. Bugeaud, M. Mignotte, Polynomial root separation, *Int. J. Number Theory* 6 (2010) 587–602.
- [21] H. Davenport, A note on binary cubic forms, *Mathematika* 8 (1961) 58–62.
- [22] J.-H. Evertse, Distances between the conjugates of an algebraic number, *Publ. Math. Debrecen* 65 (2004) 323–340.
- [23] F. Götze, D. Kaliada, M. Korolev, On the number of quadratic polynomials with bounded discriminants, *Mat. Zametki* (2016), in press (in Russian), arXiv:1308.2091.
- [24] F. Götze, D. Kaliada, O. Kukso, The asymptotic number of integral cubic polynomials with bounded heights and discriminants, *Lith. Math. J.* 54 (2) (2014) 150–165.
- [25] F. Götze, D. Zaporozhets, Discriminant and root separation of integral polynomials, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 441 (2015) 144–153.
- [26] J.F. Koksma, Über die Mahlersche Klasseneinteilung der transzendenten Zahlen und die Approximation komplexer Zahlen durch algebraische Zahlen, *Monatshefte Math. Phys.* 48 (1939) 176–189.
- [27] O. Kukso, Optimal regular systems consisting of the roots of polynomials with small discriminants and their applications, *Vestsī Nats. Akad. Navuk Belarusī Ser. Fiz.-Mat. Navuk* (2) (2007) 41–47, 124.
- [28] K. Mahler, Über das Mass der Menge aller  $S$ -Zahlen, *Math. Ann.* 106 (1932) 131–139.
- [29] K. Mahler, An inequality for the discriminant of a polynomial, *Michigan Math. J.* 11 (1964) 257–262.
- [30] M. Mignotte, Some useful bounds, in: *Computer Algebra*, Springer, Vienna, 1983, pp. 259–263.
- [31] V. Sprindžuk, The proof of Mahler’s conjecture on the measure of the set of  $S$ -numbers, *Izv. Akad. Nauk SSSR, Math. Ser.* 19 (1965) 191–194 (in Russian).
- [32] V. Sprindžuk, Mahler’s Problem in the Metric Theory of Numbers, vol. 25, Amer. Math. Soc., Providence, RI, 1969.
- [33] B.L. Van Der Waerden, *Algebra*, Springer-Verlag, Berlin, Heidelberg, 1971.
- [34] R.C. Vaughan, S. Velani, Diophantine approximation on planar curves: the convergence theory, *Invent. Math.* 166 (1) (2006) 103–124.
- [35] B. Volkmann, The real cubic case of Mahler’s conjecture, *Mathematika* 8 (1961) 55–57.
- [36] B. Volkmann, Zur metrischen Theorie der  $S$ -Zahlen. II (German), *J. Reine Angew. Math.* 213 (1963/1964) 58–65.