



UNIVERSITY OF LEEDS

This is a repository copy of *Characterization of gold mineralization in the northern Cariboo Gold District, British Columbia, Canada, through integration of compositional studies of lode and detrital gold with historical placer production: a template for evaluation of orogenic gold districts.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/99588/>

Version: Accepted Version

Article:

Chapman, RJ and Mortensen, JK (2016) Characterization of gold mineralization in the northern Cariboo Gold District, British Columbia, Canada, through integration of compositional studies of lode and detrital gold with historical placer production: a template for evaluation of orogenic gold districts. *Economic Geology*, 111 (6). pp. 1321-1345. ISSN 0361-0128

<https://doi.org/10.2113/econgeo.111.6.1321>

(c) 2016, Society of Economic Geologists, Inc. This is an author produced version of a paper published in *Economic Geology*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

TABLE 4. Analysis of placer-lode relationships in the northern CGD

Placer locality	Contributing populations	Comments
Lower Lowhee Ck	Cow Mtn veins (ma), B.C. vein (mi), Lowhee showing	
Upper Lowhee Ck	B.C. Vein (ma), local type 2 gold; e.g Wells adit and Myrtle	
Chisholm Ck	Regional type 2; e.g. Fosters Ledge	Source of type 1 gold unknown
Mosquito Ck	Mosquito mine	Potential for minor contribution from larger particles of type 4 gold
Burns Ck		Source of type 1 gold unknown
Williams Ck	BC vein for low Ag population	Source of high- Ag type 2 gold unknown
Slough Ck drainage	Medium Ag type 2	Source of type 2 gold unknown
Beggs Gulch	High Ag type 2 gold (ma)	Source of gold unknown
Antler Ck	Type 2 gold	Source unknown, but represents two Ag ranges
Cunningham Ck	Type 2 gold ± arsenopyrite	Many possible sources but not specifically linked to local placers
Little Snowshoe Ck	Midas – Snowshoe Creek	
Keithley Ck	Sources on Yanks Peak	Many possible sources but not specifically linked to local placers

ma= major component, mi= minor component