



UNIVERSITY OF LEEDS

This is a repository copy of *Characterization of gold mineralization in the northern Cariboo Gold District, British Columbia, Canada, through integration of compositional studies of lode and detrital gold with historical placer production: a template for evaluation of orogenic gold districts.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/99588/>

Version: Accepted Version

Article:

Chapman, RJ and Mortensen, JK (2016) Characterization of gold mineralization in the northern Cariboo Gold District, British Columbia, Canada, through integration of compositional studies of lode and detrital gold with historical placer production: a template for evaluation of orogenic gold districts. *Economic Geology*, 111 (6). pp. 1321-1345. ISSN 0361-0128

<https://doi.org/10.2113/econgeo.111.6.1321>

(c) 2016, Society of Economic Geologists, Inc. This is an author produced version of a paper published in *Economic Geology*. Uploaded in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

TABLE 5. Information Relating to Economically Important Placer Creeks in the Study Area.

Locality	Figure	Fineness range ¹	Mean Au % ²	Gold type(s)	Production (oz) ¹	Estimation of Au mass by type (oz)		
						Type 1	Type 2a	Type 2b
Lowhee Ck.	7A	816-914	91	1, 2a	74,023	59218	14805	0
Mosquito Ck.	7A	896-912	89	1, 4?	18,295	18295	0	0
Williams Ck.	7A	811-869	85	2a, 2b	85,530		42765	42765
Chisholm Ck.	7A	909-911	90	1,2a	U			
Burns Ck.	7A	904-919	92	1	5,655	4524	1131	0
Dragon Ck.	7B	888-926	82	2b, 3	2,498	0	2498	0
Nelson Ck.	7B	898-919	14	2b	13,620	0	0	13620
Coulter Ck.	7B	894-905	90	2a,2b	1,015	0	203	812
Slough Ck.	7B	900-920	90	2a, 2b	29,977	0	11991	17986
Antler Ck.	7C	819-861	84	2a	33,652	0	33652	0
Beggs Gulch	7C	789-801	80	2a	U			
Grouse Ck.	n/a	813-833	n.a.	2a	14,435	0	14435	0
Amador Gulch	7D	850	86	2a, 2b	178	0	89	89
Perkins Gulch	7D	889-892	87	2a, 2b	1,466	0	733	733
Lightening Ck.	7D	868-908	88	2a, 2b	98,602	0	49301	49301
Cunningham Ck.	7E	853-870	85	2a	12,857	0	12857	0
Peter Ck.	7E	U	83	2a	1,717	0	1717	0
Keithley Ck.	7E	872-895	89	2a	35,395	0	35395	0
Snowshoe Ck.	7E	U	86	2a	13,940	0	13940	0
TOTAL						82037	235511	125306
% by mass						19	53	30

. ¹ Data from Holland, 1950. ² Data from the present study. U= unrecorded.